

THE COINVARIANT ALGEBRA OF THE SYMMETRIC GROUP AS A DIRECT SUM OF INDUCED MODULES

HIDEAKI MORITA and TATSUHIRO NAKAJIMA*

(Received September 18, 2003)

Abstract

Let R_n be the coinvariant algebra of the symmetric group S_n . The algebra has a natural gradation. For a fixed l ($1 \leq l \leq n$), let $R_n(k; l)$ ($0 \leq k \leq l-1$) be the direct sum of all the homogeneous components of R_n whose degrees are congruent to k modulo l . In this article, we will show that for each l there exists a subgroup H_l of S_n and a representation $\Psi(k; l)$ of H_l such that each $R_n(k; l)$ is induced by $\Psi(k; l)$.

1. Introduction

Throughout this article, we follow [5] for fundamental terminology on partitions, Young tableaux and symmetric functions.

A *partition* of a positive integer n is a weakly decreasing sequence $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_k)$ of nonnegative integers with $\lambda_1 + \lambda_2 + \dots + \lambda_k = n$. We also denote the partition λ by $(1^{m_1} 2^{m_2} \dots n^{m_n})$, where m_i is the multiplicity of i in λ for $1 \leq i \leq n$. If λ is a partition of n , we simply write $\lambda \vdash n$. The *Young diagram* of a partition λ is a set of points

$$Y_\lambda = \{(i, j) \in \mathbb{Z}^2 \mid 1 \leq j \leq \lambda_i\},$$

in which we regard the coordinates increase from left to right, and from top to bottom. Let $[n]$ denote the set of integers $\{1, 2, \dots, n\}$. A *standard tableau* T of shape λ is a bijection $T: Y_\lambda \rightarrow [n]$ with the condition that the assigned numbers strictly increase along both the rows and the columns in Y_λ . We illustrate the Young diagram Y_λ and a standard tableau T for $\lambda = (3, 2, 2) \vdash 7$ in the following:

$$Y_\lambda = \begin{array}{ccc} \bullet & \bullet & \bullet \\ \bullet & \bullet & \\ \bullet & \bullet & \end{array}, \quad T = \begin{array}{ccc} 1 & 3 & 4 \\ 2 & 5 & \\ 6 & 7 & \end{array}.$$

We denote by $\text{STab}(\lambda)$ the set of all the standard tableaux of shape λ .

For a standard tableau T of shape $\lambda \vdash n$, define the *descent set* $\text{Des}(T)$ by

$$\text{Des}(T) := \{i \in [n-1] \mid i+1 \text{ is located in a lower row than } i \text{ in } T\}.$$

*The second author was partially supported by the grants from the Miyata Foundation

We call the sum of the elements of $\text{Des}(T)$ the *major index* of T , and denote it by $\text{maj}(T)$. In the preceding example, $\text{Des}(T) = \{1, 4, 5\}$ and $\text{maj}(T) = 1 + 4 + 5 = 10$.

Let S_n be the symmetric group of degree n , and

$$P_n = \mathbb{C}[x_1, x_2, \dots, x_n]$$

denote the polynomial ring with n variables over \mathbb{C} . As customary, S_n acts on P_n from the left as permutations of variables by setting

$$(wf)(x_1, x_2, \dots, x_n) = f(x_{w(1)}, x_{w(2)}, \dots, x_{w(n)}),$$

where $w \in S_n$ and $f(x_1, x_2, \dots, x_n) \in P_n$. Let $I_n = \bigoplus_{d \geq 0} I^d$ denote the graded S_n -stable ideal of P_n generated by the elementary symmetric functions. Hence the quotient algebra $R_n = P_n/I_n$ is also a graded S_n -module. We write its homogeneous decomposition as

$$R_n = \bigoplus_{d \geq 0} R_n^d,$$

and call R_n the *coinvariant algebra* of S_n . It is well known that the coinvariant algebra R_n affords the left regular representation of S_n .

Let us consider, for each integer $k = 0, \dots, n - 1$, the direct sum $R_n(k; n)$ of homogeneous components of R whose degrees are congruent to k modulo n , i.e.,

$$R_n(k; n) = \bigoplus_{d \equiv k \pmod n} R_n^d.$$

Since each homogeneous component R_n^d is S_n -invariant, these subspaces also afford representations of S_n , and the dimensions of these representations do not depend on k , i.e.,

$$\dim R_n(k; n) = (n - 1)!$$

for all $k = 0, \dots, n - 1$.

In [4], W. Kraśkiewicz and J. Weyman consider these S_n -modules, and prove that each $R_n(k; n)$ is induced from a corresponding irreducible representation of a cyclic subgroup of S_n (see also [2, Proposition 8.2] [6, Theorem 8.9]). Precisely, let γ be the cyclic permutation $(12 \cdots n)$, and C_n the subgroup of S_n generated by γ . The cyclic subgroup C_n of degree n has n inequivalent irreducible representations

$$\psi^{(k)}: C_n \longrightarrow \mathbb{C}^\times, \quad \gamma \longmapsto \zeta_n^k,$$

where ζ_n is the primitive root of unity, and the following equivalence of S_n -modules holds for each $k = 0, \dots, n - 1$:

$$R_n(k; n) \cong_{S_n} \text{ind}_{C_n}^{S_n}(\psi^{(k)}).$$

REMARK. In fact, the number n by which we take modulo is the *Coxeter number* of S_n , i.e., the order of the Coxeter elements of the Coxeter group of type A_{n-1} . They also obtain similar results for Coxeter groups of type B_n and D_n . Stembridge obtains more general results [8]. He treats the Complex reflection groups G and shows that the coinvariant algebra of G has the similar properties for the irreducible representation of the cyclic subgroup of G generated by a *Springer's regular element* [7]. We can easily see that the Coxeter elements are regular.

They also prove that the multiplicity of a irreducible representation of S_n in R_n^d ($d \geq 0$) is described by the major index of standard tableaux. It is well known that the irreducible representations of S_n are in one to one correspondence with the partitions of n . For $\lambda \vdash n$ let V^λ denote the corresponding irreducible representation of S_n . They showed that the multiplicity $[R_n^d : V^\lambda]$ of V^λ in R_n^d equals the number of standard tableaux whose major indices are d :

$$[R_n^d : V^\lambda] = \#\{T \in \text{STab}(\lambda) \mid \text{maj}(T) = d\}.$$

(see also [2, Theorem 8.6] [6, Theorem 8.8].) Combining these results, the multiplicities of the irreducible representation V^λ in the induced representations $\psi^{(k)} \uparrow_{C_n}^{S_n} \cong_{S_n} R_n(k; n)$ are easily obtained:

$$[R_n(k; n) : V^\lambda] = \#\{T \in \text{STab}(\lambda) \mid \text{maj}(T) \equiv k \pmod n\}.$$

It should be mentioned here that a more refined result is obtained by R. Adin, F. Brenti and Y. Roichman [1] recently. For each subset $S \subseteq [n - 1]$, they construct an S_n -module R_S satisfying

$$R_n^d = \bigoplus_S R_n^S,$$

where the direct sum is taken over the subsets $S \subseteq [n - 1]$ such that $\sum_{i \in S} i = d$, and describe the multiplicities of irreducible constituents on R_n^S as follows:

$$[R_n^S : V^\lambda] = \#\{T \in \text{STab}(\lambda) \mid \text{Des}(T) = S\}.$$

They also consider an analogue of the theorem of Krařkiewicz and Weyman for the Weyl groups of type B , and obtain a result on the irreducible decompositions of the coinvariant algebras of type B finer than one already obtained by Stembridge in [8].

The aim of the present article is to achieve a generalization of these results in the following sense. Fix an integer $l \in [n]$ and consider subspaces of R_n obtained by gathering homogeneous components whose degrees are congruent modulo l . Precisely,

for each $k = 0, \dots, l - 1$ we will consider

$$R_n(k; l) = \bigoplus_{d \equiv k \pmod l} R_n^d.$$

We can see that the dimension of the space $R_n(k; l)$ is independent of k , i.e.,

$$\dim R_n(k; l) = \frac{n!}{l}$$

for all $k = 0, \dots, l - 1$ (Proposition 4). In this article we will seek out a systematic realization of each submodule $R_n(k; l)$ as a S_n -module induced from a subgroup of S_n that is determined by l . First we settle a subgroup H_l of S_n for each $l \in [n]$, then construct a representation $\Psi(k; l)$ of H_l for each $k = 0, \dots, l - 1$. When we write $n = dl + r$ with $0 \leq r \leq l - 1$, the subgroup H_l turns out to be isomorphic to a direct product of the cyclic group of order l and the symmetric group of degree r , i.e.,

$$H_l \cong C_l \times S_r.$$

The representation $\Psi(k; l)$ of H_l is not necessarily irreducible in contrast to the case $l = n$ (Section 4). Finally, we verify that

$$R_n(k; l) \cong_{S_n} \text{ind}_{H_l}^{S_n} (\Psi(k; l))$$

for each l and k by comparing the graded characters of R_n and $\bigoplus_{k=0}^{l-1} \text{ind}_{H_l}^{S_n} (\Psi(k; l))$ as polynomials in q modulo $q^l - 1$ (Theorem 8).

2. Coinvariant algebra and its graded character

Let $R_n = \bigoplus_{d \geq 0} R_n^d$ be the coinvariant algebra of S_n and its homogeneous decomposition. Let q be an indeterminate over \mathbb{C} . Define the graded character of R_n by

$$X_n(q) = \sum_{d \geq 0} q^d \chi^{n,d},$$

where $\chi^{n,d}$ is the character of the representation R_n^d of S_n . We denote by $X_{n,\rho}(q)$ and $\chi_\rho^{n,d}$ the value of $X_n(q)$ and $\chi^{n,d}$ at elements of cycle-type $\rho \vdash n$, respectively. Precisely, $X_{n,\rho}(q)$ is a polynomial in q whose coefficient in q^d is $\chi_\rho^{n,d}$. This polynomial $X_{n,\rho}(q)$ is also known as a *Green polynomial* $Q_\rho^{(1^n)}(q)$ of type A [3] [5, III.7].

The graded character of R_n has a well-known product formula ([3, Appendix]. see also [2, Proposition 8.1]), that plays an essential role in the present article.

Proposition 1. *For any partition $\rho = (1^{m_1} 2^{m_2} \dots n^{m_n})$ of n , we have*

$$X_{n,\rho}(q) = \frac{(1 - q)(1 - q^2) \dots (1 - q^n)}{(1 - q)^{m_1} (1 - q^2)^{m_2} \dots (1 - q^n)^{m_n}}.$$

From the Proposition above, we can prove the following auxiliary result.

Proposition 2. *Fix a integer $l \in [n]$. Let p be a divisor of l , $n = ep + s$ ($0 \leq s \leq p - 1$), and θ a primitive p -th root of unity. If $\rho \vdash n$ satisfies*

$$X_{n,\rho}(\theta) \neq 0,$$

then $\rho = (1^{m_1} \dots s^{m_s} p^e)$, where $m_1 + 2m_2 + \dots + sm_s = s$.

Proof. We apply Stembridge's argument for the case $l = n$ (see [2, Section 8]) to our situation. By Proposition 1, we have

$$X_{n,\rho}(\theta) = \frac{(1-q)(1-q^2)\dots(1-q^n)}{(1-q)^{m_1}(1-q^2)^{m_2}\dots(1-q^n)^{m_n}} \Big|_{q=\theta},$$

for $\rho = (1^{m_1} 2^{m_2} \dots n^{m_n}) \vdash n$. Thus $X_{n,\rho}(\theta) \neq 0$ implies that all the vanishing factors in the numerator are canceled by corresponding factors in the denominator. There are e vanishing factors: $1-q^p, 1-q^{2p}, \dots, 1-q^{ep}$ in the numerator, and $m_p + m_{2p} + \dots + m_{ep}$ vanishing factors: $(1-q^p)^{m_p}, (1-q^{2p})^{m_{2p}}, \dots, (1-q^{ep})^{m_{ep}}$ in the denominator. Since

$$pm_p + 2pm_{2p} + \dots + epm_{ep} \leq m_1 + 2m_2 + \dots + nm_n = n (= ep + s),$$

we have

$$m_p + 2m_{2p} + \dots + em_{ep} \leq e.$$

Therefore,

$$e = m_p + m_{2p} + \dots + m_{ep} \leq m_p + 2m_{2p} + \dots + em_{ep} \leq e.$$

Hence, we have $m_p = e$. We also obtain $m_i = 0$ for $s + 1 \leq i \leq n$ ($i \neq p$) since $n - pm_p = n - pe = s$. Thus, we have

$$m_1 + 2m_2 + \dots + sm_s = s. \quad \square$$

Let $l \in [n]$ be a fixed integer. For each $k = 0, 1, \dots, l - 1$, we define

$$R_n(k; l) := \bigoplus_{d \equiv k \pmod{l}} R_n^d,$$

i.e.,

$$R_n = \bigoplus_{k=0}^{l-1} R_n(k; l).$$

We prove that the dimensions of the spaces $R_n(k; l)$ are independent of the choice of k . We first show the following lemma.

Lemma 3. *Let q be an indeterminate and $f(q) = \sum_{i \geq 0} a_i q^i \in \mathbb{C}[q]$ a polynomial in q . Let $l \geq 2$ be an integer and ζ_l a primitive l -th root of unity. Then the following conditions are equivalent:*

- (1) $f(\zeta_l^k) = 0$ for each $k = 1, \dots, l-1$,
- (2) The partial sums $c_k = \sum_{i \equiv k \pmod{l}} a_i$ ($k = 0, 1, \dots, l-1$) of coefficients of the polynomial $f(q)$ are independent of the choice of k .

Proof. If the condition (b) holds, then $f(q)$ is divisible by

$$1 + q + q^2 + \dots + q^{l-1} = \frac{1 - q^l}{1 - q},$$

and hence we have (a).

We shall prove the converse. From (a) we have

$$f(\zeta_l^k) = a_0 + a_1 \zeta_l^k + a_2 (\zeta_l^k)^2 + \dots = 0 \quad (k = 0, 1, \dots, l-1).$$

By the definition of c_k , it reduces to the linear equation system in c_0, \dots, c_{l-1} :

$$\begin{cases} c_0 + c_1 \zeta_l + c_2 \zeta_l^2 + \dots + c_{l-1} \zeta_l^{l-1} = 0, \\ c_0 + c_1 \zeta_l^2 + c_2 (\zeta_l^2)^2 + \dots + c_{l-1} (\zeta_l^2)^{l-1} = 0, \\ \vdots \\ c_0 + c_1 \zeta_l^{l-1} + c_2 (\zeta_l^{l-1})^2 + \dots + c_{l-1} (\zeta_l^{l-1})^{l-1} = 0. \end{cases}$$

Since the rank of the coefficient matrix of the equation system is $l-1$, it has an one dimensional solution space. It is clear that $(c_0, c_1, \dots, c_{l-1}) = (1, 1, \dots, 1)$ satisfies the equation system, hence we have $c_0 = c_1 = \dots = c_{l-1}$. \square

By using the above lemma, we easily reach our aim.

Proposition 4. *Let $l \in [n]$ be a fixed integer. Then the dimension of $R_n(k; l)$ is independent of the choice of $k = 0, 1, \dots, l-1$, i.e., we have*

$$\dim R_n(k; l) = \frac{n!}{l}$$

for all $k = 0, 1, \dots, l$.

Proof. If $l = 1$, then the assertion is trivial. Suppose that $l \geq 2$. Let ζ_l be a primitive l -th root of unity. If we evaluate the formula in Proposition 1 at the identity ele-

ment $e \in S_n$, then we have

$$\begin{aligned} [X_n(q)](e) &= X_{n,(1^n)}(q) \\ &= \frac{(1-q)(1-q^2)\cdots(1-q^n)}{(1-q)^n} \\ &= (1+q)(1+q+q^2)\cdots(1+q+\cdots+q^{n-1}) \end{aligned}$$

It follows immediately that, for each $k = 0, \dots, l-1$,

$$X_{n,(1^n)}(\zeta_l^k) = \sum_{d \geq 0} (\dim R_n^d) q^d |_{q=\zeta_l^k} = 0.$$

By Lemma 3, we obtain that $\dim R_n(k; l) = \sum_{d \equiv k \pmod{l}} \dim R_n^d$ is independent of $0 \leq k \leq l-1$ and is equal to $n!/l$. □

If $w \in S_n$, the cycle type $\rho(w)$ of w is the partition $\rho(w) = (1^{m_1} 2^{m_2} \cdots n^{m_n})$. For a partition ρ of n , let C_ρ be the conjugacy class in S_n containing $w \in S_n$ such that $\rho(w) = \rho$. For any partition $\rho = (1^{m_1} 2^{m_2} \cdots n^{m_n})$, define

$$z_\rho = \frac{n!}{|C_\rho|} = 1^{m_1} m_1! 2^{m_2} m_2! \cdots n^{m_n} m_n!.$$

Let f and g be arbitrary class functions on S_n . There is a natural scalar product of f and g defined by

$$\langle f, g \rangle_{S_n} := \frac{1}{n!} \sum_{w \in S_n} f(w)g(w).$$

(For a general finite group G , the scalar product is defined by $\langle f, g \rangle := (1/|G|) \times \sum_{w \in G} f(w)\overline{g(w)}$, where $\overline{g(w)}$ denotes the complex conjugate of $g(w)$. However, we can use $g(w)$ instead of $\overline{g(w)}$ here since all characters of S_n are rational.) Note that if δ_λ ($\lambda \vdash n$) is the class function defined by

$$\delta_\lambda(w) = \begin{cases} 1 & \text{if } \rho(w) = \lambda \\ 0 & \text{otherwise,} \end{cases}$$

then $\langle \delta_\lambda, f \rangle_{S_n} = z_\lambda^{-1} f(\lambda)$.

If $n = dl + r$ ($0 \leq r \leq l-1$), then we can embed $S_{dl} \times S_r$ in S_n by

$$(2.1) \quad \begin{aligned} S_{dl} &= \{w \in S_n \mid w(i) = i \text{ for all } i = dl + 1, \dots, n\}, \\ S_r &= \{w \in S_n \mid w(i) = i \text{ for all } i = 1, \dots, dl\}. \end{aligned}$$

We see that, if $u \in S_{dl}$ and $v \in S_r$, the element $u \times v \in S_n$ has cycle-type $\rho(u \times v) = \rho(u) \cup \rho(v)$.

Let f and g be characters of the representations ϕ of S_{dl} and ψ of S_r , respectively. Then $f \times g$ defined by

$$(f \times g)(u, v) = f(u)g(v) \quad (u \in S_{dl}, v \in S_r)$$

is the character of the tensor product representation $\phi \otimes \psi$ of $S_{dl} \times S_r$. We define

$$f \cdot g = \text{ind}_{S_{dl} \times S_r}^{S_n}(f \times g),$$

which is a character of the induced representation $\text{ind}_{S_{dl} \times S_r}^{S_n}(\phi \otimes \psi)$ of S_n .

The following is a key proposition to the main result.

Proposition 5. *Let n be a positive integer, and choose an integer l ($1 \leq l \leq n$). If $n = dl + r$ ($0 \leq r < l$), then we have*

$$X_n(q) \equiv (X_{dl}(q) \cdot X_r(q)) \pmod{q^l - 1}.$$

Proof. We show that

$$(2.2) \quad X_{n,\rho}(q) \equiv (X_{dl}(q) \cdot X_r(q))_\rho \pmod{q^l - 1}$$

for each $\rho \vdash n$, where $(X_{dl}(q) \cdot X_r(q))_\rho$ is the value of $(X_{dl}(q) \cdot X_r(q))$ at elements of cycle-type ρ . By the Lagrange interpolation and Proposition 2, in order to verify (2.2), it is sufficient to show that

$$(X_{dl}(\theta) \cdot X_r(\theta))_\rho = \begin{cases} X_{n,\rho}(\theta) & \text{if } \rho = (1^{m_1} \dots s^{m_s} p^e) \\ 0 & \text{otherwise.} \end{cases}$$

for each $\theta = \zeta_l^k$ ($k = 0, \dots, l-1$), where p is the multiplicative order of θ . Note that p divides l . Using the property of the class function δ_ρ , we then have

$$\begin{aligned} & z_\rho^{-1} (X_{dl}(\theta) \cdot X_r(\theta))_\rho \\ &= \langle (X_{dl}(\theta) \cdot X_r(\theta)), \delta_\rho \rangle_{S_n} \\ &= \left\langle (X_{dl}(\theta) \times X_r(\theta)), \text{res}_{S_{dl} \times S_r}^{S_n}(\delta_\rho) \right\rangle_{S_{dl} \times S_r} \quad (\text{by Frobenius reciprocity}) \\ &= \frac{1}{(dl)! r!} \sum_{u \in S_{dl}} \sum_{v \in S_r} (X_{dl}(\theta) \times X_r(\theta))(u, v) \delta_\rho(u \times v) \\ &= \frac{1}{(dl)! r!} \sum_{u \in S_{dl}} \sum_{v \in S_r} \sum_{\rho^1, \rho^2} X_{dl,\rho(u)}(\theta) X_{r,\rho(v)}(\theta) \delta_{\rho^1}(u) \delta_{\rho^2}(v) \\ &= \sum_{\rho^1, \rho^2} z_{\rho^1}^{-1} z_{\rho^2}^{-1} X_{dl,\rho^1}(\theta) X_{r,\rho^2}(\theta), \end{aligned}$$

where $\rho^1 \vdash dl$ and $\rho^2 \vdash r$ are partitions such that $\rho^1 \cup \rho^2 = \rho$. Now let $n = ep + s$ and $r = fp + s$ ($0 \leq s < p$). Then $dl/p = e - f$. By Proposition 2, $X_{dl, \rho^1} X_{r, \rho^2} = 0$ unless $\rho^1 = (p^{e-f})$ and $\rho^2 = (1^{m_1} \dots s^{m_s} p^f)$. Hence, if ρ is not of the form $(1^{m_1} \dots s^{m_s} p^e)$ for some $(1^{m_1} \dots s^{m_s}) \vdash s$, we have $(X_{dl}(\theta) \cdot X_r(\theta)) = 0$. On the other hand, we pick $\rho^1 = (p^{e-f})$ and $\rho^2 = (1^{m_1} \dots s^{m_s} p^f)$ so that $\rho = (1^{m_1} \dots s^{m_s} p^e)$, and finally we have

$$\begin{aligned} & z_\rho^{-1} (X_{dl}(\theta) \cdot X_r(\theta))_\rho \\ &= z_{(p^{e-f})}^{-1} z_{(1^{m_1} \dots s^{m_s} p^f)}^{-1} X_{dl, (p^{e-f})}(\theta) X_{r, (1^{m_1} \dots s^{m_s} p^f)}(\theta) \\ &= z_{(p^{e-f})}^{-1} z_{(1^{m_1} \dots s^{m_s} p^f)}^{-1} \frac{(1-q) \dots (1-q^{dl})}{(1-q^p)^{e-f}} \frac{(1-q) \dots (1-q^r)}{(1-q)^{m_1} \dots (1-q^s)^{m_s} (1-q^p)^f} \Big|_{q=\theta} \\ &= z_{(p^{e-f})}^{-1} z_{(1^{m_1} \dots s^{m_s} p^f)}^{-1} \binom{e}{f}^{-1} \frac{(1-q) \dots (1-q^{dl})(1-q^{dl+1}) \dots (1-q^{dl+r})}{(1-q)^{m_1} \dots (1-q^s)^{m_s} (1-q^p)^e} \Big|_{q=\theta} \\ &= z_\rho^{-1} \frac{(1-q)(1-q^2) \dots (1-q^n)}{(1-q)^{m_1} \dots (1-q^s)^{m_s} (1-q^p)^e} \Big|_{q=\theta} \\ &= z_\rho^{-1} X_{n, \rho}(\theta) \end{aligned} \quad \square$$

Translating Proposition 2 and Proposition 5 into the language of the Green polynomials, we obtain the following formula.

Corollary 6. *Let $n > l$ be positive integers, p a divisor of l , and θ a primitive p -th root of unity. If we write $n = dl + r = ep + s$ ($0 \leq r \leq l - 1$, $0 \leq s \leq p - 1$), then*

- (1) $Q_\rho^{(1^n)}(\theta) = 0$ unless $\rho = (1^{m_1} \dots s^{m_s} p^e)$ and $m_1 + 2m_2 + \dots + sm_s = s$.
- (2) If $\rho = (1^{m_1} \dots s^{m_s} p^e)$,

$$Q_\rho^{(1^n)}(q) \equiv Q_{\rho^1}^{(1^{dl})}(q) Q_{\rho^2}^{(1^r)}(q) \pmod{q^l - 1},$$

where $\rho^1 = (p^{e-f}) \vdash dl$ and $\rho^2 = (1^{m_1} \dots s^{m_s} p^f) \vdash r$.

3. $l|n$ case

In this section, we consider the case where l divides n , and show that each $R_n(k; l)$ is induced from a representation of a cyclic subgroup of S_n .

Suppose that l divides n , and say $d = n/l$. Let C_l be the cyclic group of order l , and we embed C_l into S_n as follows:

$$C_l \cong \langle \gamma_1 \gamma_2 \dots \gamma_d \rangle \subset S_n,$$

where $\gamma_1 = (1, 2, \dots, l)$, $\gamma_2 = (l + 1, l + 1, \dots, 2l)$, \dots , $\gamma_d = ((d - 1)l + 1, \dots, dl)$. The cyclic group C_l has inequivalent l irreducible representations $\psi^{(0)}, \dots, \psi^{(l-1)}$, i.e.,

$$\psi^{(k)}: C_l \longrightarrow \mathbb{C}^\times, \quad \gamma_1 \gamma_2 \dots \gamma_d \longmapsto \zeta_l^k,$$

where ζ_l denotes a primitive l -th root of unity. Let

$$\tau^{(k)} := \frac{1}{l} \sum_{i=0}^{l-1} \zeta_l^{-ik} (\gamma_1 \cdots \gamma_d)^i \quad (k = 0, 1, \dots, l-1).$$

We can easily check that each $\tau^{(k)}$ is an idempotent by a direct calculation.

Let $\mathbb{C}[S_n]$ be the group algebra of S_n . Consider the representation of S_n afforded by the left ideal $\mathbb{C}[S_n]\tau^{(k)}$, which is equivalent to the induced representation $\text{ind}_{C_l}^{S_n}(\psi^{(k)})$. Its character $\chi[\mathbb{C}[S_n]\tau^{(k)}]$ is given by $\Gamma_n\tau^{(k)}$, where Γ_n is an operator defined by

$$\Gamma_n: \mathbb{C}[S_n] \longrightarrow \mathbb{C}[S_n], \quad \rho \longmapsto \sum_{w \in S_n} w^{-1} \rho w$$

(see e.g., [2, Proposition 5.2] [6, Lemma 8.4]). Here we regard an element $\rho = \sum_{w \in S_n} \rho_w w \in \mathbb{C}[S_n]$ as the function on S_n that maps $w \in S_n$ to the coefficient ρ_w :

$$\text{ind}_{C_l}^{S_n}(\chi[\psi^{(k)}]) = \Gamma_n\tau^{(k)},$$

where $\chi[\psi^{(k)}]$ stands for the C_l -character of $\psi^{(k)}$.

We have shown in Proposition 4 that the dimension of the space

$$R_n(k; l) = \bigoplus_{d \equiv k \pmod{l}} R_n^d$$

is constant with respect to $k = 0, \dots, l-1$. This fact suggests that every $R_n(k; l)$ ($k = 0, \dots, l-1$) are induced from the same dimensional representations of a certain subgroup of S_n . In fact, we can verify that, for each $k = 0, \dots, l-1$, there exists an irreducible representation of C_l that yields $R_n(k; l)$.

Proposition 7. *Let n be a positive integer and l a divisor of n . Write $d = n/l$. For $i = 1, 2, \dots, d$, let γ_i be the cyclic permutation $((i-1)l+1, (i-1)l+2, \dots, il)$. Let C_l be the cyclic subgroup of S_n generated by $\gamma_1 \cdots \gamma_d$ and $\{\psi^{(k)} \mid k = 0, 1, \dots, l-1\}$ the set of its inequivalent irreducible representations. Then, we have an isomorphism of S_n -modules*

$$R_n(k; l) \cong_{S_n} \text{ind}_{C_l}^{S_n}(\psi^{(k)}) \quad (k = 0, 1, \dots, l-1).$$

Proof. We prove that

$$(3.1) \quad X_n(q) \equiv \sum_{k=0}^{l-1} q^k \text{ind}_{C_l}^{S_n}(\chi[\psi^{(k)}]) \pmod{q^l - 1}.$$

Using the Lagrange interpolation again, we only have to show that the both sides of (3.1) coincide when $q = \zeta_l^s$ ($s = 0, 1, \dots, l-1$).

Recall that

$$\text{ind}_{C_l}^{S_n}(\chi[\psi^{(k)}]) = \Gamma_n \tau^{(k)}$$

for each $k = 0, \dots, l-1$. Substituting $q = \zeta_l^s$ in the right hand side of (3.1), we obtain

$$\begin{aligned} \sum_{k=0}^{l-1} (\zeta_l^s)^k \text{ind}_{C_l}^{S_n}(\chi[\psi^{(k)}]) &= \sum_{k=0}^{l-1} \zeta_l^{ks} \Gamma_n \tau^{(k)} = \Gamma_n(\gamma_1 \cdots \gamma_d)^s \sum_{k=0}^{l-1} \tau^{(k)} \\ &= \Gamma_n(\gamma_1 \cdots \gamma_d)^s \sum_{k=0}^{l-1} \frac{1}{l} \sum_{i=0}^{l-1} \zeta_l^{-ik} (\gamma_1 \cdots \gamma_d)^i \\ &= \Gamma_n(\gamma_1 \cdots \gamma_d)^s \frac{1}{l} \sum_{i=0}^{l-1} (1 + \zeta_l^{-i} + \zeta_l^{-2i} + \cdots + \zeta_l^{-(l-1)i}) (\gamma_1 \cdots \gamma_d)^i \\ &= \Gamma_n(\gamma_1 \cdots \gamma_d)^s \end{aligned}$$

for each $s = 0, 1, \dots, l-1$. Since the cycle-type of $(\gamma_1 \cdots \gamma_d)^s$ can be written as (p^e) ($e = n/p$), where p is the multiplicative order of $(\zeta_l^s)^p = 1$, we have

$$\sum_{k=0}^{l-1} (\zeta_l^s)^k \text{ind}_{C_l}^{S_n}(\chi[\psi^{(k)}])_\rho = \begin{cases} z_{(p^e)}, & \text{if } \rho = (p^e) \\ 0, & \text{otherwise} \end{cases}$$

for a partition ρ . Hence the congruence (3.1) immediately follows from Proposition 1 and Proposition 2. □

4. Main result

Let n be a positive integer, and choose an integer $l = 1, 2, \dots, n$. Suppose that $n = dl + r$, where $0 \leq r \leq l-1$. Let R_n be the coinvariant algebra of S_n , and $R_n = \bigoplus_{d \geq 0} R_n^d$ its homogeneous decomposition. For each $k = 0, 1, \dots, l-1$, define

$$R_n(k; l) := \bigoplus_{d \equiv k \pmod{l}} R_n^d.$$

Now, for each $l = 1, 2, \dots, n$, we define a subgroup H_l of S_n by

$$\begin{aligned} H_l &= \langle \gamma_1 \gamma_2 \cdots \gamma_d \rangle \times S_r \\ &\cong C_l \times S_r, \end{aligned}$$

where γ_i is the cyclic permutation $((i-1)l+1, (i-1)l+2, \dots, il)$, and the symmetric group S_r of degree r is identified as the subgroup $\{w \in S_n \mid w(i) = i \text{ for all } i = 1, 2, \dots, n-r\}$ of S_n .

For each $k = 0, 1, \dots, l-1$, we construct a representation $\Psi(k; l)$ of H_l as follows:

$$\Psi(k; l) := \bigoplus_{\lambda \vdash r} \bigoplus_{T \in \text{STab}(\lambda)} \psi^{(k - \text{maj}(T))} \otimes V^\lambda,$$

where $k - \text{maj}(T) = k - \text{maj}(T) \pmod l$, $\{\psi^{(i)} \mid i = 0, \dots, l-1\}$ is the set of inequivalent irreducible representation of C_l , and V^λ ($\lambda \vdash r$) is the irreducible representation of S_r corresponding to the partition λ of r . Then it can be seen that the dimension of $\Psi(k; l)$ does not depend on k and hence so does $\text{deg ind}_{H_l}^{S_n}(\Psi(k; l))$. Actually, since $\text{deg } V^\lambda = \# \text{STab}(\lambda)$ and $\sum_{\lambda \vdash r} \# \text{STab}(\lambda)^2 = r!$, we have

$$\begin{aligned} \text{deg } \Psi(k; l) &= \sum_{\lambda \vdash r} \sum_{T \in \text{STab}(\lambda)} \text{deg } \psi^{(k - \text{maj}(T))} \otimes V^\lambda \\ &= \sum_{\lambda \vdash r} \sum_{T \in \text{STab}(\lambda)} \# \text{STab}(\lambda) \\ &= \sum_{\lambda \vdash r} \# \text{STab}(\lambda)^2 \\ &= r!, \end{aligned}$$

and $\text{deg ind}_{H_l}^{S_n}(\Psi(k; l)) = r!n!/r!l = n!/l$, which coincides with the dimension of $R_n(k; l)$. Moreover, we prove that these two representations are equivalent.

Theorem 8 (Main result). *Let n be a positive integer. Fix an integer $l \in [n]$ and write $n = dl + r$ ($0 \leq r \leq l-1$). Let $H_l \cong C_l \times S_r$ be the subgroup of S_n defined above and $\Psi(k; l)$ ($k = 0, 1, \dots, l-1$) representations of it defined by*

$$\Psi(k; l) := \bigoplus_{\lambda \vdash r} \bigoplus_{T \in \text{STab}(\lambda)} \psi^{(k - \text{maj}(T))} \otimes V^\lambda,$$

where $\psi^{(i)}$ and V^λ stand for the irreducible representations of C_l and S_r , respectively. Then, for each $k = 0, 1, \dots, l-1$, there is an isomorphism

$$R_n(k; l) \cong_{S_n} \text{ind}_{H_l}^{S_n}(\Psi(k; l)).$$

as an S_n -module.

Proof. By the definition of $\Psi(k; l)$, it suffices to show

$$(4.1) \quad X_n(q) \equiv \sum_{k=0}^{l-1} q^k \sum_{\lambda \vdash r} \sum_{T \in \text{STab}(\lambda)} \text{ind}_{H_l}^{S_n} (\chi [\psi^{(k - \text{maj}(T))} \otimes V^\lambda]) \pmod{q^l - 1}.$$

Let S_{dl} and S_r be the subgroup of S_n defined in (2.1). Since H_l is a subgroup

of $S_{dl} \times S_r$, we have

$$\text{ind}_{H_l}^{S_n} (\psi^{\overline{(k-\text{maj}(T))}} \otimes V^\lambda) \cong_{S_n} \text{ind}_{S_{dl} \times S_r}^{S_n} \left(\text{ind}_{H_l}^{S_{dl} \times S_r} (\psi^{\overline{(k-\text{maj}(T))}} \otimes V^\lambda) \right)$$

for any $\lambda \vdash r$. Therefore, the right hand side of (4.1) equals

$$\begin{aligned} & \sum_{k=0}^{l-1} \sum_{\lambda \vdash r} \sum_{T \in \text{STab}(\lambda)} q^k \text{ind}_{S_{dl} \times S_r}^{S_n} \left(\text{ind}_{C_l \times S_r}^{S_{dl} \times S_r} (\chi [\psi^{\overline{(k-\text{maj}(T))}} \otimes V^\lambda]) \right) \\ &= \sum_k \sum_\lambda \sum_T q^k \text{ind}_{S_{dl} \times S_r}^{S_n} \left(\text{ind}_{C_l}^{S_{dl}} (\chi [\psi^{\overline{(k-\text{maj}(T))}}]) \times \chi[V^\lambda] \right) \\ &= \text{ind}_{S_{dl} \times S_r}^{S_n} \left(\sum_k \sum_\lambda \sum_T q^{k-\text{maj}(T)} \text{ind}_{C_l}^{S_{dl}} (\chi [\psi^{\overline{(k-\text{maj}(T))}}]) \times q^{\text{maj}(T)} \chi[V^\lambda] \right) \\ (4.2) \quad & \equiv \text{ind}_{S_{dl} \times S_r}^{S_n} X_{dl}(q) \left(\sum_\lambda \sum_T q^{\text{maj}(T)} \chi[V^\lambda] \right) \pmod{q^l - 1} \text{ by (3.1)}. \end{aligned}$$

By the theorem of Kraškiewicz-Weyman, the multiplicity $[R_n^d : V^\lambda]$ of irreducible components isomorphic to V^λ ($\lambda \vdash n$) is the number of standard Young tableaux of shape λ whose major index equals d , that is,

$$[R_n^d : V^\lambda] = \#\{T \in \text{STab}(\lambda) : \text{maj}(T) = d\}.$$

Hence we have

$$(4.3) \quad X_r(q) = \sum_{\lambda \vdash r} \sum_{T \in \text{STab}(\lambda)} q^{\text{maj}(T)} \chi[V^\lambda].$$

Applying (4.3) and Proposition 5, we see that (4.2) equals

$$\begin{aligned} & \text{ind}_{S_{dl} \times S_r}^{S_n} X_{dl}(q) \left(\sum_{\lambda \vdash r} \sum_{T \in \text{STab}(\lambda)} q^{\text{maj}(T)} \chi[V^\lambda] \right) \\ &= \text{ind}_{S_{dl} \times S_r}^{S_n} X_{dl}(q) \times X_r(q) \\ &= (X_{dl}(q) \cdot X_r(q)) \\ &\equiv X_n(q) \pmod{q^l - 1}, \end{aligned}$$

and complete the proof. □

When $r = 0$ or 1 , H_l is a cyclic group and $\Psi(k; l)$ is irreducible. In this case, the generator of H_l coincides with a regular element of S_n defined by Springer [7].

It is obvious that the multiplicity of V^λ in $R_n(k; l)$ is obtained by counting the number of standard Young tableaux of shape λ with the major index congruent

to k modulo l , that is,

$$[R_n(k; l) : V^\lambda] = \#\{T \in \text{STab}(\lambda) \mid \text{maj}(T) \equiv k \pmod{l}\}.$$

EXAMPLE. In the case of $n = 5$ and $l = 3$, the subgroup H_3 is $\langle(123)\rangle \times \langle(45)\rangle$, which is isomorphic to $C_3 \times S_2$. Then we have

$$R_5(k; 3) \cong_{S_5} \text{ind}_{H_3}^{S_5} \left((\psi^{(k)} \otimes V^{(2)}) \oplus (\psi^{(k-1)} \otimes V^{(1,1)}) \right)$$

for each $k = 0, 1, 2$.

If we consider the case $n = 11$ and $l = 4$ (thus $r = 3$), then the subgroup H_4 is $\langle(1234)(5678)\rangle \times \langle(9, 10), (10, 11)\rangle$ isomorphic to $C_4 \times S_3$. Hence, for each $R_{11}(k; 4)$ ($k = 0, 1, 2, 3$) is isomorphic to the representation induced by

$$\begin{aligned} \Psi(0; 4) &= (\psi^{(0)} \otimes V^{(3)}) \oplus (\psi^{(3)} \otimes V^{(2,1)}) \oplus (\psi^{(2)} \otimes V^{(2,1)}) \oplus (\psi^{(1)} \otimes V^{(1,1,1)}), \\ \Psi(1; 4) &= (\psi^{(1)} \otimes V^{(3)}) \oplus (\psi^{(0)} \otimes V^{(2,1)}) \oplus (\psi^{(3)} \otimes V^{(2,1)}) \oplus (\psi^{(2)} \otimes V^{(1,1,1)}), \\ \Psi(2; 4) &= (\psi^{(2)} \otimes V^{(3)}) \oplus (\psi^{(1)} \otimes V^{(2,1)}) \oplus (\psi^{(0)} \otimes V^{(2,1)}) \oplus (\psi^{(3)} \otimes V^{(1,1,1)}), \\ \Psi(3; 4) &= (\psi^{(3)} \otimes V^{(3)}) \oplus (\psi^{(2)} \otimes V^{(2,1)}) \oplus (\psi^{(1)} \otimes V^{(2,1)}) \oplus (\psi^{(0)} \otimes V^{(1,1,1)}). \end{aligned}$$

ACKNOWLEDGEMENT. The authors wish to thank Professor Toshio Oshima for his valuable comments, in particular, on the proof of Lemma 3. The authors also thank Professor Toshiaki Shoji for helpful comments on regular elements of Coxeter groups. Professor Ken-ichi Shinoda, Professor Itaru Terada and Professor Tohru Umeda gave useful comments that improve the article. The authors wish to thank Professor Hiro-Fumi Yamada for useful discussions. Finally, we thank for the referee's suggestions for improving our manuscript.

References

- [1] R. Adin, F. Brenti and Y. Roichman: *Descent representations and multivariate statistics*, preprint, 2001.
- [2] A.M. Garsia: *Combinatorics of the free Lie algebra and the symmetric group*; in *Analysis, et cetera...*, Jürgen Moser Festschrift, Academic Press, New York, 1990, 309–382.
- [3] J.A. Green: *The character of the finite general linear groups*, *Trans. Amer. Math. Soc.* **80** (1955), 402–447.
- [4] W. Kraśkiewicz and J. Weyman: *Algebra of coinvariants and the action of Coxeter elements*, *Bayreuth. Math. Schr.* **63** (2001), 265–284.
- [5] I.G. Macdonald: *Symmetric Functions and Hall Polynomials*, 2nd ed., Oxford University Press, 1995.
- [6] C. Reutenauer: *Free Lie Algebras*, Oxford University Press, Oxford, 1993.
- [7] T.A. Springer: *Regular elements of finite reflection groups*, *Invent. Math.* **25** (1974), 159–198.
- [8] J.R. Stembridge: *On the eigenvalues of representations of reflection groups and wreath products*, *Pacific J. Math.* **140** (1989), 359–396.

Hideaki Morita
School of Science
Tokai University
Hiratsuka 259-1292, Japan
e-mail: moritah@msg.biglobe.ne.jp

Tatsuhiro Nakajima
Faculty of Economics
Meikai University
Urayasu 279-8550, Japan
e-mail: tatsu.nkjm@mac.com