

## COHOMOLOGY OPERATIONS IN THE LOOP SPACE OF THE COMPACT EXCEPTIONAL GROUP $F_4$

Dedicated to Professor A. Komatu on his 70-th birthday

TAKASHI WATANABE

(Received April 14, 1978)

### 1. Introduction

Let  $F_4$  be the compact, simply connected, exceptional Lie group of rank 4. In [6] we have described the Hopf algebra structure of  $H_*(\Omega F_4; Z)$ . Using this thoroughly, we can compute the action of the mod  $p$  Steenrod algebra  $\mathcal{A}_p$  on  $H^*(\Omega F_4; Z_p)$  for every prime  $p$ . But here we deal with the cases  $p=2$  (Theorem 4) and  $p=3$  (Theorem 5) only, because in the other cases the result follows immediately from a spectral sequence argument for the path fibration  $\Omega F_4 \rightarrow PF_4 \rightarrow F_4$ .

Let  $C(=C_s) = T^1 \cdot Sp(3)$  in the notation of [3], which is a closed connected subgroup of  $F_4$ . Then in [6] the homogeneous space  $F_4/C$  has been found to be a *generating variety* for  $F_4$ . That is, there exists a map  $f_s: F_4/C \rightarrow \Omega F_4$  such that the image of  $f_{s*}: H_*(F_4/C; Z) \rightarrow H_*(\Omega F_4; Z)$  generates the Pontrjagin ring  $H_*(\Omega F_4; Z)$ . In this situation Bott [1, §6] asserted that the Steenrod operations in  $H^*(\Omega F_4; Z_p)$  can be deduced from their effect on  $H^*(F_4/C; Z_p)$ . This is the motive of our work.

Throughout the paper  $X$  will always denote any connected space such that  $H_*(X; Z)$  is of finite type.

### 2. The generating variety

In this section we shall compute the  $\mathcal{A}_p$ -module structure of  $H^*(F_4/C; Z_p)$  for  $p=2$  and 3.

First since  $C$  contains a maximal torus  $T$  of  $F_4$ , we have a commutative diagram

$$(2.1) \quad \begin{array}{ccc} F_4/T & \xrightarrow{q} & F_4/C \\ \downarrow \iota & & \downarrow j \\ BT & \xrightarrow{\rho} & BC. \end{array}$$

We require the following notations and results (2.2)-(2.6), whose details can be found in [3, §4]:

$$(2.2) \quad H^*(BT; Z) = Z[t, y_1, y_2, y_3] \quad \text{where } \deg t = \deg y_i = 2 \quad (i=1, 2, 3).$$

Put  $z_i = y_i(t - y_i) \in H^4(BT; Z)$  and let  $q_i = \sigma_i(z_1, z_2, z_3) \in H^{4i}(BT; Z)$  for  $i=1, 2, 3$  where  $\sigma_i$  denotes the  $i$ -th elementary symmetric function. Then we have

$$(2.3) \quad H^*(BC; Z) = Z[t, q_1, q_2, q_3] \quad \text{where } \deg t = 2 \text{ and } \deg q_i = 4i \quad (i=1, 2, 3).$$

$$(2.4) \quad \rho^*(t) = t \text{ and } \rho^*(q_i) = q_i \quad (i=1, 2, 3).$$

On the other hand we have

$$(2.5) \quad H^*(F_4/C; Z) = Z[t, u, v, w]/(t^3 - 2u, u^2 - 3t^2v + 2w, 3v^2 - t^2w, v^3 - w^2) \quad \text{where} \\ \deg t = 2, \deg u = 6, \deg v = 8 \text{ and } \deg w = 12.$$

$$(2.6) \quad j^*(t) = t, j^*(q_1) = t^2, j^*(q_2) = 3v \text{ and } j^*(q_3) = w.$$

We shall say that an  $\mathcal{A}_p$ -action on  $H^*(X; Z_p)$  is *non-trivial* if it does not follow directly from the axioms 1), 4) or 5) of [4, p.1 and p.76]. With these preliminaries we have

**Proposition 1.** *The non-trivial  $\mathcal{A}_2$ -action on*

$$H^*(F_4/C; Z_2) = Z_2[t, u, v, w]/(t^3, u^2 - t^2v, v^2 - t^2w, v^3 - w^2)$$

is given by:

- (1)  $Sq^2(t) = t^2.$
- (2)  $Sq^2(u) = v, Sq^4(u) = tv \text{ and } Sq^6(u) = t^2v.$
- (3)  $Sq^2(v) = 0, Sq^4(v) = w, Sq^6(v) = tw \text{ and } Sq^8(v) = v^2.$
- (4)  $Sq^2(w) = tw, Sq^4(w) = 0, Sq^6(w) = 0, Sq^8(w) = vzw, Sq^{10}(w) = tvw$   
and  $Sq^{12}(w) = w^2.$

Proof. (1) and the last equalities in (2), (3) and (4) are immediate from the axiom 3) of [4, p.1].

First we consider (3) and (4). Since  $v = j^*(q_2)$  and  $w = j^*(q_3)$  in  $H^*(F_4/C; Z_2)$  by (2.6), it suffices to determine  $Sq^i(q_2)$  and  $Sq^i(q_3)$  in  $H^*(BC; Z_2)$ . To do so, by (2.4), it suffices to compute  $Sq^i(q_2)$  and  $Sq^i(q_3)$  in  $H^*(BT; Z_2)$ . But this is a direct calculation, for  $H^*(BT; Z_2)$  is multiplicatively generated by the elements of degree 2 (see (2.2)).

Finally we show the remaining part of (2). By [3, Corollary 4.5] we may set

$$Sq^2(u) = k \cdot tu + l \cdot v \text{ and} \\ Sq^4(u) = m \cdot t^2u + n \cdot tv$$

for some  $k, l, m, n \in Z_2$ . Then from the Adem relations  $Sq^2Sq^2 = Sq^3Sq^1$ ,  $Sq^2Sq^4 = Sq^6 + Sq^5Sq^1$  and  $Sq^4Sq^4 = Sq^7Sq^1$ , it follows that  $kl=0$ ,  $lm+n=1$  and  $l=n$  respectively. Hence  $k=m=0$  and  $l=n=1$ , which proves (2).

Next we turn to the case  $p=3$ . To begin with we need some preparations.

As in [6, §3], put  $x=t/2$  and  $x_i=x-y_i$  for  $i=1, 2, 3$ . Thus  $x, x_i \in H^2(BT; Z[1/2])$  ( $i=1, 2, 3$ ). Furthermore put

$$(2.7) \quad t_1 = -x_1 + x_2, t_2 = x_1 + x_2, t_3 = -x_3 - x \text{ and } t_4 = -x_3 + x.$$

Note that  $t_i \in H^2(BT; Z)$  ( $i=1, 2, 3, 4$ ). For later convenience we introduce the notation:

$$c_i = \sigma_i(t_1, t_2, t_3, t_4) \in H^{2i}(BT; Z);$$

$$p_j = \sigma_j(x_1^2, x_2^2, x_3^2) \in H^{4j}(BT; Q),$$

where  $1 \leq i \leq 4$  and  $1 \leq j \leq 3$ . A straightforward calculation using (2.7) yields:

$$(2.8) \quad c_1^2 - 2c_2 = 2(p_1 + x^2);$$

$$c_2^2 - 2c_3c_1 + 2c_4 = (p_1 + x^2)^2 + 2(p_2 + p_1x^2 - 3x_1^2x_2^2 - 3x_3^2x^2).$$

We also need:

$$(2.9) \quad p_1 = -q_1 + 3x^2; \quad p_2 = q_2 - 2q_1x^2 + 3x^4.$$

This follows from [6, (3.6)].

Now put  $\gamma_1 = c_1/2$ . From the discussion in [3, §4.2] we observe that

$$H^*(BT; Z) = Z[t_1, t_2, t_3, t_4, \gamma_1]/(c_1 - 2\gamma_1)$$

on which the Weyl group  $\Phi(F_4)$  acts as follows:

	$\tilde{R}$	$R_1$	$R_2$	$R_3$	$R_4$
$t_1$			$t_2$	$-t_1$	$t_1 - \gamma_1$
$t_2$		$t_3$	$t_1$		$t_2 - \gamma_1$
$t_3$	$t_4$	$t_2$			$t_3 - \gamma_1$
$t_4$	$t_3$				$t_4 - \gamma_1$

This allows us to identify the  $t_i$  with that given in [5, §4(A)]. Then by [5, Theorem A] we have

$$(2.10) \quad H^*(F_4/T; Z) = Z[t_1, t_2, t_3, t_4, \gamma_1, \gamma_3, w_4]/(c_1 - 2\gamma_1, c_2 - 2\gamma_1^2, c_3 - 2\gamma_3, c_4 - 2c_3\gamma_1 + 2\gamma_1^4 - 3w_4, -c_4\gamma_1^2 + \gamma_3^2, 3c_4\gamma_1^4 - \gamma_1^8 + 3c_3\gamma_1w_4 + 3w_4^2, w_4^3)$$

where  $\deg t_i = \deg \gamma_1 = 2$  ( $i=1, 2, 3, 4$ ),  $\deg \gamma_3 = 6$  and  $\deg w_4 = 8$ .

(By abuse of notation we have written  $t_i, c_i$ , etc. for their images under  $\iota^*$ .)

It is well known that  $\text{Spin}(9)$  is a closed connected subgroup of  $F_4$ , and

the homogeneous space  $F_4/\text{Spin}(9)$  can be identified with the Cayley projective plane  $\Pi$ , whose integral cohomology is given by:

$$(2.11) \quad H^*(\Pi; Z) = Z[w]/(w^3) \text{ where } \deg w = 8.$$

Since  $T \subset \text{Spin}(9) \subset F_4$ , we have a natural map

$$F_4/T \xrightarrow{p} F_4/\text{Spin}(9) = \Pi.$$

Then it follows from [5, (6.9)] that

$$(2.12) \quad p^*(w) = w_4.$$

The following result may be of independent interest.

**Lemma 2.**  $q^*(v) = w_4 + x_1^2 x_2^2 + x_3^2 x^2 + 2x^4$  in  $H^8(F_4/T; Z)$ .

Proof. From (2.6), the commutativity of (2.1), and (2.4) we see that  $q^*(v) = q_2/3 \in H^8(F_4/T; Z)$ . (2.10), together with (2.8) and (2.9), gives:

$$\begin{aligned} w_4 &= \frac{1}{3} (c_4 - c_3 c_1 + \frac{1}{2} c_2^2) \\ &= \frac{1}{3} (p_2 + p_1 x^2 - 3x_1^2 x_2^2 - 3x_3^2 x^2) \\ &= \frac{1}{3} q_2 - x_1^2 x_2^2 - x_3^2 x^2 - 2x^4. \end{aligned}$$

Combining these we get the result. (It is easy to verify that  $x_1^2 x_2^2 + x_3^2 x^2 + 2x^4$  is in fact an integral class.)

**Proposition 3.** *The non-trivial  $\mathcal{A}_3$ -action on*

$$H^*(F_4/C; Z_3) = Z[t, v]/(t^8, v^3)$$

is given by:

- (1)  $\mathcal{P}^1(t) = t^3.$
- (2)  $\mathcal{P}^1(v) = -t^6, \mathcal{P}^2(v) = 0, \mathcal{P}^3(v) = 0$  and  $\mathcal{P}^4(v) = 0.$

Proof. (1) and the last equality in (2) are immediate from the axiom 3) of [4, p.76].

To show (2), we must compute  $\mathcal{P}^i(v)$  for  $i=1, 2, 3$ . Since  $q^*: H^*(F_4/C; Z) \rightarrow H^*(F_4/T; Z)$  is a split monomorphism (see [3, §3]) and its image is known with  $Z_3$ -coefficients (see Lemma 2), it suffices to determine  $q^*(\mathcal{P}^i(v))$  in  $H^*(F_4/T; Z_3)$ . (2.10) and the same calculation as in (2.8) yield:

$$\begin{aligned} H^*(F_4/T; Z_3) &= Z_3[t_1, t_2, t_3, t_4, w_4]/(c_2+c_1^2, c_4-c_3c_1-c_4^1, \\ &\quad c_3^2-c_4c_1^2, c_1^8, w_4^3) \\ &= Z_3[x_1, x_2, x_3, x, w_4]/(x_1^2+x_2^2+x_3^2+x^2, x_2^4+ \\ &\quad x_2^2x_3^2+x_3^4+x_2^2x^2+x_3^2x^2+x^4, x_3^6+x_3^4x^2+x_3^2x^4 \\ &\quad +x^6, x^8, w_4^3). \end{aligned}$$

Moreover (2.12) and (2.11) imply that  $\mathcal{P}^i(w_4)=0$  for all  $i \geq 1$ . Therefore we have

$$\begin{aligned} q^*(\mathcal{P}^1(v)) &= \mathcal{P}^1(q^*(v)) \\ &= \mathcal{P}^1(x_1^2x_2^2+x_3^2x^2-x^4) \\ &= -x_1^4x_2^2-x_1^2x_2^4-x_3^4x^2-x_3^2x^4-x^6 \\ &= -x^6. \end{aligned}$$

So since  $q^*(t)=-x$  it follows that  $\mathcal{P}^1(v)=-t^6$ . Using the Adem relation  $\mathcal{P}^2=-\mathcal{P}^1\mathcal{P}^1$ , we also get  $\mathcal{P}^2(v)=0$ . Finally we consider  $\mathcal{P}^3(v)$ . By [3, Corollary 4.5] we may set

$$\mathcal{P}^3(v) = k \cdot t^6v + l \cdot t^2v^2$$

for some  $k, l \in Z_3$ . Then  $q^*(\mathcal{P}^3(v))=k \cdot x^6w_4 + l \cdot x^2w_4^2 + \dots$ . On the other hand notice that  $\mathcal{P}^3(q^*(v))$  does not involve  $x^6w_4$  or  $x^2w_4^2$ , and they are linearly independent in  $H^{20}(F_4/T; Z_3)$ . This implies that  $k=l=0$  as required.

### 3. Main results

As seen in [6], the algebraic description of  $H_*(\Omega F_4; Z)$  is much easier than that of  $H^*(\Omega F_4; Z)$ . For this reason we shall treat the right  $\mathcal{A}_p$ -action on  $H_*(X; Z_p)$  which dualizes to the usual left  $\mathcal{A}_p$ -action on  $H^*(X; Z_p)$ .

We first consider the case  $p=2$  and follow the notation of [2]. For  $i \geq 0$  let  $( )Sq^i$  be the dual to  $Sq^i( )$ . Then these operations have the following properties (cf. [4, p. 1]):

(3.1)  $( )Sq^i: H_n(X; Z_2) \rightarrow H_{n-i}(X; Z_2)$ .

(3.2) If  $\deg \alpha < 2i$ ,  $(\alpha)Sq^i=0$ .

(3.3) If  $\deg \alpha = 2i$ ,  $(\alpha)Sq^i = \sqrt{\alpha}$  where  $\sqrt{\quad}$  is the dual of the squaring map for  $Z_2$ -algebras.

(3.4) (diagonal Cartan formula) Let  $\psi: H_*(X; Z_2) \rightarrow H_*(X; Z_2) \otimes H_*(X; Z_2)$  be the coproduct (induced from the diagonal map  $\Delta: X \rightarrow X \times X$ ). If  $\psi(\alpha) = \sum \alpha' \otimes \alpha''$ , then

$$\psi((\alpha)Sq^k) = \sum_{i+j=k} (\alpha')Sq^i \otimes (\alpha'')Sq^j.$$

Suppose now that  $X$  is an  $H$ -space, and  $\alpha \cdot \beta$  denotes the Pontrjagin product of  $\alpha$  and  $\beta$  in  $H_*(X; Z_2)$ . Then one can readily check:

(3.5) (*internal Cartan formula*)  

$$(\alpha \cdot \beta)Sq^k = \sum_{i+j=k} (\alpha)Sq^i \cdot (\beta)Sq^j .$$

We shall say that an  $\mathcal{A}_2$ -action on  $H_*(X; Z_2)$  is *non-trivial* if it does not follow from (3.1), (3.2) or (3.5).

Let us now consider the case  $X = \Omega F_4$ . Hereafter we shall use the notations and results of [6] without specific reference.

First we have

(3.6)  $H_*(\Omega F_4; Z_2) = Z_2[\sigma_1, \sigma_2, \sigma_5, \sigma_7, \sigma_{11}] / (\sigma_1^2)$  where  $\deg \sigma_i = 2i$  ( $i = 1, 2, 5, 7, 11$ ).  
 Moreover  $\sigma_1, \sigma'_5 = \sigma_5 + \sigma_2^2 \sigma_1, \sigma_7$  and  $\sigma'_{11} = \sigma_{11} + \sigma_5^2 \sigma_1 + \sigma_7 \sigma_2^2$  are primitive, and  $\tilde{\psi}(\sigma_2) = \sigma_1 \otimes \sigma_1$ .

Therefore (by (3.5)) we have only to determine the  $( )Sq^i$  on the elements  $\sigma_1, \sigma_2, \sigma'_5, \sigma_7$  and  $\sigma'_{11}$ . On the other hand, (3.4) implies that for  $i \geq 1$   $( )Sq^i$  sends a primitive element to another primitive element. In view of (3.6), the primitive elements of  $H_*(\Omega F_4; Z_2)$  which appear in degrees  $\leq 22$  are:

deg	2	8	10	14	16	20	22
	$\sigma_1$	$\sigma_2^2$	$\sigma'_5$	$\sigma_7$	$\sigma_2^4$	$\sigma_5'^2$	$\sigma'_{11}$

These, together with (3.1) and (3.2), show that possible non-zero operations (among non-trivial operations) are:

deg	2	8	10	14	16	20
	$(\sigma_2)Sq^2$	$(\sigma'_5)Sq^2$	$(\sigma_7)Sq^4$	$(\sigma'_{11})Sq^8$	$(\sigma'_{11})Sq^6$	$(\sigma'_{11})Sq^2$
		$(\sigma_7)Sq^6$				

Let us compute these operations. First by (3.3) we have  $(\sigma_2)Sq^2 = \sigma_1$ . Next we want to determine the coefficient  $k \in Z_2$  in the equation  $(\sigma'_5)Sq^2 = k \cdot \sigma_2^2$ . By use of (3.5) we have  $(\sigma'_5)Sq^2 = (\sigma_5)Sq^2 + (\sigma_2^2 \sigma_1)Sq^2 = (\sigma_5)Sq^2$  and so  $(\sigma_5)Sq^2 = k \cdot \sigma_2^2$ . Dualizing this gives  $Sq^2(a_4) = k \cdot b_5 + l \cdot a_5$  for some  $l \in Z_2$ . Since  $f_s^*(Sq^2(a_4)) = Sq^2(f_s^*(a_4)) = Sq^2(tu) = t^2u + tv$  by use of (1) and (2) of Proposition 1, and since  $f_s^*(b_5) = t^2u + tv$  and  $f_s^*(a_5) = tv$ , it follows that  $k = 1$  (and also  $l = 0$ ). Thus we obtain  $(\sigma'_5)Sq^2 = \sigma_2^2$ .

Instead of proceeding further, we state here a pattern of computation: The problem is to determine the coefficient  $k' \in Z_2$  in the equation

$$(\alpha')Sq^i = k' \cdot \beta$$

where  $\alpha'$  and  $\beta$  are primitive. In particular  $\alpha' = \alpha + \text{decomposables}$  and  $\alpha$  is

the image under the mod 2 reduction of an integral class which is indecomposable. Using (3.5) we get

$$(\alpha)Sq^i = k \cdot \beta + \dots$$

where  $k(\in Z_2)$  and  $k'$  determine each other. Dualizing this gives

$$(*) \quad Sq^i(b) = k \cdot a + \dots$$

where  $a$  and  $b$  are dual to  $\alpha$  and  $\beta$  respectively. In particular  $a$  is the image under the mod 2 reduction of an integral class which is primitive. Since the composite

$$PH^*(\Omega F_4; Z) \xrightarrow{\subset} H^*(\Omega F_4; Z) \xrightarrow{f_s^*} H^*(F_4/C; Z)$$

is a split monomorphism, it is sufficient to consider (\*) in  $H^*(F_4/C; Z_2)$  via  $f_s^*$ . But in [6, §4] the cohomology ring  $H^*(\Omega F_4; Z)$  and its image under  $f_s^*$  have been described, and by Proposition 1 we already know the  $\mathcal{A}_2$ -action on  $H^*(F_4/C; Z_2)$ . Thus  $k$  and hence  $k'$  are computable.

In this way routine computations yield

**Theorem 4.** *The non-trivial  $\mathcal{A}_2$ -action on*

$$H_*(\Omega F_4; Z_2) = Z_2[\sigma_1, \sigma_2, \sigma'_5, \sigma_7, \sigma'_{11}]/(\sigma_1^2)$$

is given by:

- (1)  $(\sigma_2)Sq^2 = \sigma_1$ .
- (2)  $(\sigma'_5)Sq^2 = \sigma_2^2$  and  $(\sigma'_5)Sq^4 = 0$ .
- (3)  $(\sigma_7)Sq^2 = 0, (\sigma_7)Sq^4 = \sigma'_5$  and  $(\sigma_7)Sq^6 = 0$ .
- (4)  $(\sigma'_{11})Sq^2 = \sigma_5'^2, (\sigma'_{11})Sq^4 = 0, (\sigma'_{11})Sq^6 = \sigma_2^4, (\sigma'_{11})Sq^8 = \sigma_7$  and  $(\sigma'_{11})Sq^{10} = 0$ .

The argument for the case  $p=3$  is similar (we have prepared Proposition 3 in place of Proposition 1) and so we only present the result.

**Theorem 5.** *The non-trivial  $\mathcal{A}_3$ -action on*

$$H_*(\Omega F_4; Z_3) = Z_3[\sigma_1, \sigma_3, \sigma'_5, \sigma'_7, \sigma'_{11}]/(\sigma_1^3)$$

is given by:

- (1)  $(\sigma_3)\mathcal{O}^1 = \sigma_1$ .
- (2)  $(\sigma'_5)\mathcal{O}^1 = 0$ .

$$(3) (\sigma'_7)\mathcal{P}^1 = \sigma'_5 \text{ and } (\sigma'_7)\mathcal{P}^2 = 0.$$

$$(4) (\sigma'_{11})\mathcal{P}^1 = \sigma^3_{3,(\sigma'_{11})}\mathcal{P}^2 = 0 \text{ and } (\sigma'_{11})\mathcal{P}^3 = 0.$$

OSAKA CITY UNIVERSITY

---

### References

- [1] R. Bott: *The space of loops on a Lie group*, Michigan Math. J. **5** (1958), 35–61.
- [2] A.K. Bousfield and E.B. Curtis: *A spectral sequence for the homotopy of nice spaces*, Trans. Amer. Math. Soc. **151** (1970), 457–479.
- [3] K. Ishitoya and H. Toda: *On the cohomology of irreducible symmetric spaces of exceptional type*, J. Math. Kyoto Univ. **17** (1977), 225–243.
- [4] N.E. Steenrod and D.B.A. Epstein: *Cohomology operations*, Princeton University Press, Princeton, N.J., 1962.
- [5] H. Toda and T. Watanabe: *The integral cohomology rings of  $F_4/T$  and  $E_6/T$* , J. Math. Kyoto Univ. **14** (1974), 257–286.
- [6] T. Watanabe: *The homology of the loop space of the exceptional group  $F_4$* , Osaka J. Math. **15** (1978), 463–474.