

ON THE NON-TRIVIALITY OF THE GREEK LETTER ELEMENTS IN THE ADAMS-NOVIKOV E_2 -TERM

MICHIMASA TANABE

(Received July 6, 1989)

1. Introduction

For a prime p , there is a spectral sequence, called the Adams-Novikov spectral sequence, converging to the stable homotopy ring of spheres localized at p , $\pi_{*}^S(p)$, whose E_2 -term is isomorphic to $\text{Ext}_{BP_*(BP)}(BP_*, BP_*)$, where BP denotes the Brown-Peterson spectrum at p ([2]).

In [1], the elements $\alpha_i^{(n)}$ were defined in $\text{Ext}_{BP_*(BP)}^n(BP_*, BP_*)$ for every positive integer n, t . (In [1], these elements were denoted by $\eta(v_n^t)$ instead of $\alpha_i^{(n)}$.) Here $\alpha^{(n)}$ stands for the n -th letter of Greek alphabet and we call them Greek letter elements.

For $n \leq 3$, it has already been proved that these elements are represented by non-trivial elements in $\pi_{*}^S(p)$ if $p \geq 2n$ ([3], [4], [1], [2]) but in the case of $n \geq 4$, we have had few information on them yet.

The purpose of this paper is to prove the non-triviality of $\alpha_i^{(n)}$ in $\text{Ext}_{BP_*(BP)}^n(BP_*, BP_*)$ for $n \geq 4$ under suitable restrictions on p, t and we succeed for $p \geq n$ and $1 \leq t \leq p-1$. Moreover we also prove p does not divide them.

In the next section, we recall the necessary information on BP and state our results proved in §3.

I would like to thank Professor Akira Kono for useful conversations, kind encouragement and reading this manuscript.

2. Recollections on BP and Statement of results

Let BP denote the Brown-Peterson spectrum at a prime p ([2]). Then $BP_* = \mathbf{Z}_{(p)}[v_1, v_2, \dots]$ and $BP_*(BP) = BP_*[t_1, t_2, \dots]$ where the generators v_m and t_m are defined as follows.

$BP_* \otimes \mathbf{Q} = \mathbf{Q}[\lambda_1, \lambda_2, \dots]$ for canonical generators $\lambda_m, |\lambda_m| = 2(p^m - 1)$. Then v_m are determined inductively by

$$p\lambda_m = \sum_{0 \leq i \leq m} \lambda_i v_{m-i}^{p^i} \quad (\lambda_0 = 1, v_0 = p) \quad ([2] \text{ A 2.2.2})$$

and t_m by

$$\eta_R(\lambda_m) = \sum_{0 \leq i \leq m} \lambda_i t_{m-i}^{\beta^i} \quad (t_0 = 1) \quad ([2] \text{ A } 2.1.27)$$

where $\eta_R: BP_* \otimes Q \rightarrow BP_*(BP) \otimes Q$ is the right unit of the Hopf algebroid $(BP_*, BP_*(BP))$ tensored with Q .

Under the above choice of generators we have

Theorem 2.1. ([2] A. 2.2.5).

$$\sum_{i,j \geq 0}^F t_i \eta_R(v_j)^{\beta^i} = \sum_{i,j \geq 0}^F v_i t_j^{\beta^i},$$

Theorem 2.2 ([2] A 2.1.27).

$$\sum_{i > 0}^F \Delta(t_i) = \sum_{\substack{i,j \geq 0 \\ i+j > 0}}^F t_i \otimes t_j^{\beta^i},$$

where \sum^F denotes the formal group sum associated with BP and $\Delta: BP_*(BP) \rightarrow BP_*(BP) \otimes_{BP_*} BP_*(BP)$ is the coproduct of $BP_*(BP)$.

Let I_m be the ideal in BP_* generated by p, v_1, \dots, v_{m-1} . Using 2.1 we see easily that I_m is an invariant ideal in BP_* . In fact we have the following theorem.

Theorem 2.3. ([2] 4.3.2). *Let $I_m = (p, v_1, \dots, v_{m-1})$*

- (a) I_m is invariant.
- (b) For $m > 0$.

$$\text{Ext}^0(BP_*/I_m) = F_p[v_m],$$

and

$$\text{Ext}^0(BP_*) = Z_{(p)}.$$

- (c) $0 \rightarrow \sum^{2(p^m-1)} BP_*/I_m \xrightarrow{v_m} BP_*/I_m \rightarrow BP_*/I_{m+1} \rightarrow 0$
is a short exact sequence of comodules.

- (d) *The only invariant prime ideals in BP_* are the I_m for $0 \leq m \leq \infty$.*

(From here we abbreviate $\text{Ext}(M)$ for $\text{Ext}_{BP_*(BP)}(BP_*, M)$.)

This result allows us to define Greek letter elements.

We consider the short exact sequence given by (c) which leads to a long exact sequence of Ext and let

$$\delta_m: \text{Ext}^s(BP_*/I_{m+1}) \rightarrow \text{Ext}^{s+1}(BP_*/I_m)$$

denote the connecting homomorphism of the resulting long exact sequence.

DEFINITION. For $t, n > 0$, let

$$\alpha_t^{(n)} = \delta_0 \delta_1 \cdots \delta_{n-1}(v_n^t) \in \text{Ext}^n(BP_*).$$

We now state our results. Let

$$\varphi_n: \text{Ext}^s(BP_*) \rightarrow \text{Ext}^s(BP_*/(I_{n-1} + I_n^{p-n+1}))$$

be the homomorphism induced by the natural projection $BP_* \rightarrow BP_*/(I_{n-1} + I_n^{p-n+1})$. (From now on we always assume $p \geq n \geq 3$.) Then we have

Theorem 2.4. $\varphi_n(\alpha_i^{(n)}) \neq 0$ if $3 \leq n \leq p$ and $1 \leq t \leq p-1$.

As an immediate consequence of 2.4 we have

Corollary 2.5. $\alpha_i^{(n)} \neq 0$ if $3 \leq n \leq p$ and $1 \leq t \leq p-1$. Moreover p does not divide them.

REMARK. For $n \leq 3$ we have much more general results than 2.4. (See [1], [2].)

The rest of this section is devoted to describing the cobar construction which we need in the next section.

Let (A, Γ) be a Hopf algebroid such that Γ is flat over A . Then the category of (left) Γ -comodules becomes an abelian category with enough injectives, so we can define $\text{Ext}_\Gamma^s(L, M)$ for (left) Γ -comodules L, M as the s -th right derived functor of $\text{Hom}_\Gamma(L, M)$.

In the case of $L=A$, these Ext groups can be computed as the homology of the cobar complex $C_\Gamma(M)$ defined below.

DEFINITION. Let $\varepsilon: \Gamma \rightarrow A$ be the counit and $\bar{\Gamma} = \ker \varepsilon$. The cobar complex $C_\Gamma(M)$ is defined by $C_\Gamma^s(M) = \bar{\Gamma}^{\otimes s} \otimes_A M$ with the differential $d: C_\Gamma^s(M) \rightarrow C_\Gamma^{s+1}(M)$ given by

$$\begin{aligned} d(\gamma_1 \otimes \cdots \otimes \gamma_s \otimes m) &= \sum_{1 \leq i \leq s} (-1)^i \gamma_1 \otimes \cdots \otimes \gamma_{i-1} \otimes \gamma_i' \otimes \gamma_i'' \otimes \gamma_{i+1} \otimes \cdots \otimes \gamma_s \otimes m \\ &\quad + (-1)^{s+1} \gamma_1 \otimes \cdots \otimes \gamma_s \otimes m' \otimes m'' \end{aligned}$$

for $\gamma_1, \dots, \gamma_s \in \bar{\Gamma}$ and $m \in M$ where $\bar{\Gamma}^{\otimes s}$ denotes the s -fold tensor product of $\bar{\Gamma}$ over A , $\Delta(\gamma_i) = 1 \otimes \gamma_i + \sum \gamma_i' \otimes \gamma_i'' + \gamma_i \otimes 1$ and $\psi(m) = 1 \otimes m + \sum m' \otimes m''$. (Δ denotes the coproduct of Γ and ψ denotes the coaction map of M). The element $\gamma_1 \otimes \cdots \otimes \gamma_s \otimes m$ will be denoted by $\gamma_1 | \cdots | \gamma_s | m$.

Then the following isomorphism holds.

Theorem 2.6 ([2] A 1.2.12).

$$\text{Ext}_\Gamma^s(A, \Gamma) \cong H^s(C_\Gamma(M)).$$

Finally we define a certain quotient complex of $C_\Gamma(A)$ associated with a sequence of non negative integers (a_1, \dots, a_i) if $\Gamma = A[\gamma_1, \dots, \gamma_m]$ (m may be

infinity.) and γ_i is primitive modulo $(\gamma_1, \dots, \gamma_{i-1})$ for all i .

Let $E=(e_1, e_2, \dots)$ be a sequence of non negative integers such that $e_i=0$ for all but a finite number of i . We introduce an order between such sequences by saying that $E < F (= (f_1, f_2, \dots))$ iff there is a positive integer i such that $e_j=f_j$ for $j < i$ and $e_i < f_i$. Let $E+F$ denote a sequence $(e_1+f_1, e_2+f_2, \dots)$ and $\gamma^E = \gamma_1^{e_1} \dots \gamma_m^{e_m} \in \Gamma$.

DEFINITION.

$$C_\Gamma((a_1, \dots, a_i)) = \frac{C_\Gamma(A)}{\bigoplus_{s \geq 1} A\{\gamma^{E_1} | \dots | \gamma^{E_s}; E_1 + \dots + E_s > (a_1, \dots, a_i)\}}$$

where $A\{\cdot\}$ denotes the submodule of $C_\Gamma(A)$ generated by the indicated generators.

(Clearly $C_\Gamma((a_1, \dots, a_i))$ depends on the choice of γ_i but we do not indicate the generators in our notation because our choice is always evident in this paper.)

Now we show that $C_\Gamma((a_1, \dots, a_i))$ is a quotient complex of $C_\Gamma(A)$. By our assumption

$$\Delta(\gamma_i) = 1 \otimes \gamma_i + \sum \gamma'_i \otimes \gamma''_i + \gamma_i \otimes 1$$

where $\gamma'_i \in (\gamma_1, \dots, \gamma_{i-1})$ or $\gamma''_i \in (\gamma_1, \dots, \gamma_{i-1})$. Thus $\Delta(\gamma_i) \in A\{\gamma^F \otimes \gamma^G; F+G \geq (0, \dots, 0, 1)\}$ and more generally we have $\Delta(\gamma^E) \in A\{\gamma^F \otimes \gamma^G; F+G \geq E\}$ since Δ is an algebra homomorphism and $(\gamma^F \otimes \gamma^G)(\gamma^{F'} \otimes \gamma^{G'}) = \gamma^{F+F'} \otimes \gamma^{G+G'}$. Therefore $\bigoplus_{s \geq 1} A\{\gamma^{E_1} | \dots | \gamma^{E_s}; E_1 + \dots + E_s > (a_1, \dots, a_i)\}$ is a subcomplex of $C_\Gamma(A)$ as desired.

3. Proof of Theorem 2.4

Let $C(n, m)$ (resp. $D(n, m)$) denote

$$C_{BP_*(BP)/J_{n,m}}((p^{n-2}+1, p^{n-3}, \dots, p, 1))$$

(resp. $C_{BP_*(BP)/K_{n,m}}((p^{n-2}+1, p^{n-3}, \dots, p, 1))$)

where $J_{n,m} = I_m + I_{n-1}^{m+1} + I_n^{p-n+m+1}$ and $K_{n,m} = I_m + I_{n-1}^{m+2} + I_n^{p-n+m+2}$.

(Note that $BP_*(BP) = BP_*[t_1, t_2, \dots]$ and $\Delta(t_i)$ has the form

$$(3.1) \quad \Delta(t_i) = 1 \otimes t_i + t_i \otimes 1 \quad \text{in} \quad BP_*(BP) \otimes_{BP_*} BP_*(BP) / (t_1, \dots, t_{i-1})$$

for degree reasons.) It is obvious that the sequence

$$0 \rightarrow C(n, m) \xrightarrow{v_m} D(n, m) \rightarrow C(n, m+1) \rightarrow 0$$

is a short exact sequence of complexes and letting

$$\tilde{\delta}_m: H^s(C(n, m+1)) \rightarrow H^{s+1}(C(n, m))$$

denote the corresponding connecting homomorphism we have a commutative diagram

$$\begin{CD} \text{Ext}^s(BP_*/I_{m+1}) @>\delta_m>> \text{Ext}^{s+1}(BP_*/I_m) \\ @V\psi_{m+1}VV @VV\psi_mV \\ H^s(C(n, m+1)) @>\delta_m>> H^{s+1}(C(n, m)) \end{CD}$$

where ψ_m is the homomorphism induced by the natural projection $C_{BP_*(BP)}(BP_*/I_m) \rightarrow C(n, m)$.

Thus it is sufficient to show 3.2 below for the proof of 2.4 since ψ_0 factors through

$$\varphi_n: \text{Ext}^s(BP_*) \rightarrow \text{Ext}^s(BP_*/(I_{n-1} + I_n^{p-n+1})).$$

Proposition 3.2. $\tilde{\alpha}_i^{(n)} \neq 0$ in $H^n(C(n, 0))$ if $1 \leq i \leq p-1$ where $\tilde{\alpha}_i^{(n)}$ denotes the element $\psi_0(\alpha_i^{(n)}) = \tilde{\delta}_0 \cdots \tilde{\delta}_{n-1} \psi_n(v_n^i) \in H^n(C(n, 0))$.

In order to prove 3.2 we begin with giving an explicit representative for $\tilde{\alpha}_i^{(n)}$ and this requires some formulas on η_R of BP .

Lemma 3.3 ([2] 4.3.21).

$$\eta_R(v_m) \equiv v_m + v_{m-1} t_1^{p^{m-1}} - v_{m-1}^p t_1 \pmod{I_{m-1}}.$$

Lemma 3.4.

$$\eta_R(v_m) \equiv \sum_{0 \leq i \leq m} v_i t_{m-i}^{p^i} \pmod{I_m^p}.$$

For the proof of 3.4 we first prove the following simple fact about the formal group law associated with BP .

Lemma 3.5.

$$X +_F Y = X + Y \text{ in } BP_*[[X, Y]]/(X, Y)^p.$$

Proof. Note that $X +_F Y$ has the form

$$X + Y + \sum_{i, j \geq 1} a_{i, j} X^i Y^j \text{ in } BP_*[[X, Y]]$$

where $a_{i, j} = a_{j, i} \in BP_{2(i+j-1)}$.

Considering the degree of $a_{i, j}$, it is clear that $a_{i, j} = 0$ if $i + j < p$ so we get the desired result. \square

Proof of 3.4. In the degree of $\eta_R(v_m)$, the left hand side of 2.1 is congruent to $\eta_R(v_m)$ modulo I_m^p by 2.3 (a) and 3.5.

The right hand side of 2.1 is congruent to $\sum_{0 \leq i \leq m} v_i t_{m-i}^{p^i}$ modulo I_m^p and the

result follows. \square

We now describe a representative for $\tilde{\alpha}_i^{(n)}$.

Lemma 3.6. $\tilde{\alpha}_i^{(n)} \in H^n(C(n, 0))$ is represented by a cocycle

$$-t \frac{(p-1)!}{(p-n)!} v_n^{t-1} v_{n-1}^{p-n} t_{n-1} | t_{n-2}^p | \cdots | t_1^{p^{n-2}} | t_1 \in C^n(n, 0).$$

Proof. In $D(n, n-1)$,

$$\begin{aligned} d(v_n^t) &= \eta_R(v_n)^t - v_n^t \\ &= (v_n + v_{n-1} t_1^{p^{n-1}} - v_{n-1}^p t_1)^t - v_n^t \quad (\text{by 3.3}) \\ &= -t v_n^{t-1} v_{n-1}^p t_1. \end{aligned}$$

So we have

$$\tilde{\delta}_{n-1}(v_n^t) = -t v_n^{t-1} v_{n-1}^{p-1} t_1 \in H^1(C(n, n-1)).$$

(We often abuse the same notation for a cocycle and its representing element in the cohomology.)

In $D(n, n-2)$,

$$\begin{aligned} d(v_n^{t-1} v_{n-1}^{p-1} t_1) &= d(v_n^{t-1} v_{n-1}^{p-1}) | t_1 \quad (\text{by 3.1}) \\ &= \{ \eta_R(v_n)^{t-1} \eta_R(v_{n-1})^{p-1} - v_n^{t-1} v_{n-1}^{p-1} \} | t_1 \\ &= \{ v_n^{t-1} (v_{n-1} + v_{n-2} t_1^{p^{n-2}})^{p-1} - v_n^{t-1} v_{n-1}^{p-1} \} | t_1 \quad (\text{by 3.4}) \\ &= (p-1) v_n^{t-1} v_{n-1}^{p-2} v_{n-2} t_1^{p^{n-2}} | t_1 \end{aligned}$$

and thus

$$\tilde{\delta}_{n-2} \tilde{\delta}_{n-1}(v_n^t) = -t(p-1) v_n^{t-1} v_{n-1}^{p-2} t_1^{p^{n-2}} | t_1 \in H^2(C(n, n-2)).$$

More generally, by induction on k , we can easily show

$$(3.7) \quad \begin{aligned} \tilde{\delta}_{n-k} \cdots \tilde{\delta}_{n-1}(v_n^t) &= -\frac{t(p-1)!}{(p-k)!} v_n^{t-1} v_{n-1}^{p-k} t_{k-1}^{p^{n-k}} | \cdots | t_1^{p^{n-2}} | t_1 \\ &\in H^k(C(n, n-k)) \quad \text{for all } k, 2 \leq k \leq n. \end{aligned}$$

Let $k=n$ in 3.7 then we obtain the lemma. \square

Next we define a subcomplex of $C(n, 0)$ which will be denoted by $\overline{C(n, 0)}$.

Let $(P(v_{n-1}, v_n)/(v_{n-1}^{p^{n+1}}), P(v_{n-1}, v_n, t_1, \dots, t_n)/(v_{n-1}^{p^{n+1}}))$ be the sub-Hopf algebroid of $(BP_*/J_{n,0}, BP_*(BP)/J_{n,0})$ where $P(\cdot)$ denotes the polynomial algebra which has the indicated generators over \mathbf{F}_p . We define

$$\overline{C(n, 0)} = C_{p(v_{n-1}, v_n, t_1, \dots, t_n)/(v_{n-1}^{p^{n+1}})}((p^{n-2} + 1, p^{n-3}, \dots, p, 1))$$

and let

$$B(n, m) = C_{p(t_1, \dots, t_m)}((p^{n-2} + 1, p^{n-3}, \dots, p, 1))$$

where $P(t_1, \dots, t_m)$ is considered as a Hopf algebra over F_p whose coproduct is given by $\Delta(t_i) = \sum_{0 \leq j \leq i} t_j \otimes t_i^{p^j}$ ($1 \leq i \leq m$). Then the following isomorphism of differential graded algebras holds.

$$(3.8) \quad \overline{C(n, 0)} \cong P(v_{n-1}, v_n) / (v_{n-1}^{p-n+1}) \otimes_{F_p} B(n, n).$$

This follows from 3.4 and the formulas on the coproduct of $BP_*(BP)$ given by the next lemma.

Lemma 3.9 ([2] 4.3.15). *For $m \geq 1$*

$$\Delta(t_m) = \sum_{0 \leq i \leq m} t_i \otimes t_{m-i}^{p^i} \text{ in } BP_*(BP) \otimes_{BP_*} BP_*(BP) / I_m,$$

and

$$\begin{aligned} \Delta(t_{m+1}) &= \sum_{0 \leq i \leq m+1} t_i \otimes t_{m+1-i}^{p^i} \\ &\text{in } BP_*(BP) \otimes_{BP_*} BP_*(BP) / (I_m + BP_* \{t_1^{p^i} \otimes t_1^{p^i}; e_1 + e_2 \geq p^m\}). \end{aligned}$$

Now note that $|\alpha_i^{(n)}| < |v_{n+1}|$ for $t \leq p-1$ and $C(n, 0)$ is equal to the sub-complex $\overline{C(n, 0)}$ defined above in the internal degree less than $|v_{n+1}|$ and therefore 3.2 is equivalent to

Proposition 3.10. $t_{n-1} |t_{n-2}^p| \cdots |t_1^{p^{n-2}}| t_1 \neq 0$ in $H^n(B(n, n-1))$

by 3.6 and 3.8 since $B(n, n-1) = B(n, n)$ in the internal degree less than $|t_n|$ ($> |t_{n-1}| |t_{n-2}^p| \cdots |t_1^{p^{n-2}}| t_1$ for $p \geq n$).

In order to show 3.10 we need the following lemma proved at the end of this section.

Lemma 3.11. *There is a spectral sequence converging to $H^*(B(n, m))$ with*

$$E_2^{a,b} = H^b(C_{p(t_m)}(F_p)) \otimes H^a(B(n, m-1)) / R_{a,b}$$

and

$$d_r: E_r^{a,b} \rightarrow E_r^{a+r, b-r+1}$$

where $P(t_m)$ is considered as a Hopf algebra over F_p with t_m primitive and

$$R_{a,b} = F_p \{x \otimes y \in H^b(C_{p(t_m)}(F_p)) \otimes H^a(B(n, m-1)); \text{ Both } x \text{ and } y \text{ have representative cocycles } \tilde{x} \text{ and } \tilde{y} \text{ such that } \tilde{x}\tilde{y} = 0 \text{ in } B^{a+b}(n, m)\}.$$

Moreover this spectral sequence has the third grading induced by the internal degree in the cohomology which is preserved by all differentials.

Proof of 3.10. First note that

$$t_1^{p^n-2} | t_1 \neq 0 \text{ in } H^2(B(n, 1))$$

by 3.12 below since $B(n, 1)$ is a direct summand of $C_{p(t_1)}(\mathbf{F}_p)$ as a complex. (Recall our assumption $n \geq 3$ which assures $p^{n-2} > 1$.)

Lemma 3.12.

$$H^*(C_{p(t_m)}(\mathbf{F}_p)) = E(h_{m,0}, h_{m,1}, \dots) \otimes P(b_{m,0}, b_{m,1}, \dots)$$

where $h_{m,i}$ (resp. $b_{m,i}$) is represented by $t_m^{p^i}$ (resp. $\frac{1}{p} \sum_{0 < j < p} \binom{p}{j} t_m^{p^i j} | t_m^{p^i(p-j)}$) and $E(\cdot)$ denotes the exterior algebra which has the indicated generators over \mathbf{F}_p .

Proof. This result is obtained by a routine calculation. \square

Now suppose

$$(3.13) \quad t_{m-1}^{p^n-m} | \dots | t_1^{p^n-2} | t_1 \neq 0 \text{ in } H^m(B(n, m-1))$$

holds for some m , $1 < m \leq n-1$. Then the element $t_m^{p^n-m-1} \otimes t_{m-1}^{p^n-m} | \dots | t_1^{p^n-2} | t_1$ ($\in H^1(C_{p(t_m)}(\mathbf{F}_p)) \otimes H^m(B(n, m-1))$) defines a non-trivial element in the E_2 -term of the spectral sequence given by 3.11 which is clearly a permanent cycle and moreover there is no differential killing this element as observed below.

Let c_m denote the internal degree of the above element then

$$(3.14) \quad c_m = 2(p-1) \{ m p^{n-2} + (m-1) p^{n-3} + \dots + 2 p^{n-m} + p^{n-m-1} + 1 \}$$

and it is enough to prove $E_r^{m-r, r, c_m} = 0$ for all $r \geq 2$.

Using 3.11 and 3.12 we can identify the E_r with an appropriate subquotient of $\bigotimes_{\substack{1 \leq i \leq m \\ j \geq 0}} (E(h_{i,j}) \otimes P(b_{i,j}))$ and let $c_{i_1, j_1, \dots, i_l, j_l, i'_1, j'_1, \dots, i'_s, j'_s}$ denote the internal degree of

$h_{i_1, j_1} \dots h_{i_l, j_l} b_{i'_1, j'_1} \dots b_{i'_s, j'_s}$ ($1 \leq i_1 \leq \dots \leq i_l \leq m$, $1 \leq i'_1 \leq \dots \leq i'_s \leq m$) then

$$(3.15) \quad c_{i_1, j_1, \dots, i_l, j_l, i'_1, j'_1, \dots, i'_s, j'_s} = 2(p-1) \left\{ \sum_{1 \leq k \leq l} p^j (p^{i_k-1} + \dots + p + 1) + \sum_{1 \leq k \leq s} p^{j'_k+1} (p^{i'_k-1} + \dots + p + 1) \right\} .$$

Comparing 3.14 with 3.15 we see easily that $c_m = c_{i_1, j_1, \dots, i_l, j_l, i'_1, j'_1, \dots, i'_s, j'_s}$ does not hold for $l+2s \leq m$ under our assumption $m < n \leq p$ (≥ 3) and consequently $E_r^{m-r, r, c_m} = 0$ for all $r \geq 2$. Therefore

$$t_m^{p^n-m-1} | \dots | t_1^{p^n-2} | t_1 \neq 0 \text{ in } H^{m+1}(B(n, m))$$

and by induction on m we have shown 3.13 for all m , $1 < m \leq n$.

Letting $m=n$ in 3.13 we get 3.10 and thus complete the proof of 2.4 assuming 3.11. \square

Proof of 3.11. We begin with recalling the construction of the Cartan-Eilenberg spectral sequence for the following cocentral Hopf algebra extension

$$P(t_1, \dots, t_{m-1}) \xrightarrow{f} P(t_1, \dots, t_m) \xrightarrow{g} P(t_m). \quad (\text{cf. [2] A 1.3.17})$$

We first define a decreasing filtration on $C_{p(t_1, \dots, t_m)}(\mathbf{F}_p)$ by

$$(3.16) \quad \begin{aligned} \tilde{F}^{a,b} &= \mathbf{F}_p \{t^{E_1} | \dots | t^{E_{a+b}}; \text{ at least } a \text{ of the } t^{E_i} \text{ lie in } \ker g\} \\ &\subset C_{p(t_1, \dots, t_m)}^{a+b}(\mathbf{F}_p) \end{aligned}$$

and let \tilde{E}_r denote the E_r -term of the spectral sequence associated with this filtration.

Next define a homomorphism

$$\tilde{h}_{a,b}: C_{p(t_m)}^b(\mathbf{F}_p) \otimes C_{p(t_1, \dots, t_{m-1})}^a(\mathbf{F}_p) \rightarrow \tilde{E}_0^{a,b}$$

which is given by

$$\begin{aligned} \tilde{h}_{a,b}(t_m^{e_1} | \dots | t_m^{e_b} \otimes t^{E_1} | \dots | t^{E_a}) &= t_m^{e_1} | \dots | t_m^{e_b} | t^{E_1} | \dots | t^{E_a} \\ &\in \tilde{F}^{a,b} / \tilde{F}^{a+1,b-1} = \tilde{E}_0^{a,b} \end{aligned}$$

for $t_m^{e_1} | \dots | t_m^{e_b} \in C_{p(t_m)}^b(\mathbf{F}_p)$ and $t^{E_1} | \dots | t^{E_a} \in C_{p(t_1, \dots, t_{m-1})}^a(\mathbf{F}_p)$. If we consider $C_{p(t_m)}^b(\mathbf{F}_p) \otimes C_{p(t_1, \dots, t_{m-1})}^a(\mathbf{F}_p)$ as a complex with its differential $d \otimes 1$ then $\tilde{h}_{a,b}$ becomes a chain map and induces

$$\tilde{h}'_{a,b}: H^b(C_{p(t_m)}(\mathbf{F}_p)) \otimes H^a(C_{p(t_1, \dots, t_{m-1})}(\mathbf{F}_p)) \rightarrow \tilde{E}_1^{a,b}.$$

Moreover we can prove \tilde{h}' is an isomorphism and if we consider $H^b(C_{p(t_m)}(\mathbf{F}_p)) \otimes C_{p(t_1, \dots, t_{m-1})}^a(\mathbf{F}_p)$ as a complex with its differential $(-1)^b 1 \otimes d$ then $\tilde{h}'_{a,b}$ is also a chain map.

Hence we obtain an isomorphism

$$\tilde{h}''_{a,b}: H^b(C_{p(t_m)}(\mathbf{F}_p)) \otimes H^a(C_{p(t_1, \dots, t_{m-1})}(\mathbf{F}_p)) \xrightarrow{\cong} \tilde{E}_2^{a,b}$$

induced by \tilde{h}' .

Therefore we have a spectral sequence converging to $H^*(C_{p(t_1, \dots, t_m)}(\mathbf{F}_p))$ whose E_2 -term is isomorphic to $H^*(C_{p(t_m)}(\mathbf{F}_p)) \otimes H^*(C_{p(t_1, \dots, t_{m-1})}(\mathbf{F}_p))$. This spectral sequence is called the Cartan-Eilenberg spectral sequence.

We now turn to our case. It is trivial that $\tilde{F}^{a,b}$ given by 3.16 also defines a decreasing filtration on $B(n, m)$ naturally. Thus we obtain a spectral sequence E_r converging to $H^*(B(n, m))$ and a homomorphism

$$h'_{a,b}: H^b(C_{p(t_m)}(\mathbf{F}_p)) \otimes H^a(B(n, m-1)) / R_{a,b} \rightarrow E_2^{a,b}$$

induced by a chain map

$$h''_{a,b}: H^b(C_{p(t_m)}(\mathbf{F}_p)) \otimes B(n, m-1) / R'_{a,b} \rightarrow E_1^{a,b}$$

where

$$R'_{a,b} = \mathbf{F}_p \{x \otimes y \in H^b(C_{p(t_m)}(\mathbf{F}_p)) \otimes B^a(n, m-1); x \text{ has a representative cocycle } \tilde{x} \text{ such that } \tilde{x}y = 0 \text{ in } B^{a+b}(n, m)\}$$

and h' (resp. h'') is the map induced by \tilde{h}' (resp. \tilde{h}'') naturally. So we will show that h' is an isomorphism.

Let

$$R'' = \mathbf{F}_p \{x \otimes y \in C_{p(t_m)}(\mathbf{F}_p) \otimes B(n, m-1); xy = 0 \text{ in } B(n, m)\}.$$

It is easy to see that $C_{p(t_m)}(\mathbf{F}_p) \otimes B(n, m-1)/R''$ (resp. E_0) is a direct summand of $C_{p(t_m)}(\mathbf{F}_p) \otimes C_{p(t_1, \dots, t_{m-1})}(\mathbf{F}_p)$ (resp. \tilde{E}_0) as a complex where $C_{p(t_m)}(\mathbf{F}_p) \otimes C_{p(t_1, \dots, t_{m-1})}(\mathbf{F}_p)$ is endowed with the differential $d \otimes 1$ and $C_{p(t_m)}(\mathbf{F}_p) \otimes B(n, m-1)/R''$ with the induced one, and moreover through \tilde{h} , $C_{p(t_m)}(\mathbf{F}_p) \otimes B(n, m-1)/R''$ corresponds to E_0 and another summand of $C_{p(t_m)}(\mathbf{F}_p) \otimes C_{p(t_1, \dots, t_{m-1})}(\mathbf{F}_p)$ corresponds to another one of \tilde{E}_0 .

Hence the fact \tilde{h}' is an isomorphism implies h' is also an isomorphism and we complete the proof of 3.11. \square

References

- [1] H.R. Miller, D.C. Ravenel, and W.S. Wilson: *Periodic phenomena in the Adams-Novikov spectral sequence*, Ann. of Math. **106** (1977), 469–516.
- [2] D.C. Ravenel: *Complex Cobordism and Stable Homotopy Groups of Spheres*, Academic Press, 1986.
- [3] L. Smith: *On realizing complex bordism modules*, Amer. J. Math. **92** (1970), 793–856.
- [4] H. Toda: *On spectra realizing exterior parts of the Steenrod algebra*, Topology **10** (1971), 53–65.

Department of Mathematics
Faculty of Science
Kyoto University
Kyoto 606,
Japan

Current address
Department of Mathematics
The Johns Hopkins University
Baltimore, MD 21218
U.S.A.