

TWISTED COHOMOLOGY FOR HYPERBOLIC THREE MANIFOLDS

PERE MENAL-FERRER and JOAN PORTI

(Received February 24, 2010, revised February 3, 2011)

Abstract

For a complete hyperbolic three manifold M , we consider the representations of $\pi_1(M)$ obtained by composing a lift of the holonomy with complex finite dimensional representations of $\mathrm{SL}(2, \mathbf{C})$. We prove a vanishing result for the cohomology of M with coefficients twisted by these representations, using techniques of Matsushima–Murakami. We give some applications to local rigidity.

Let M be an orientable complete hyperbolic three manifold. The holonomy representation of the complete hyperbolic structure

$$\mathrm{Hol}: \pi_1(M) \rightarrow \mathrm{Isom}^+ \mathbf{H}^3 \cong \mathrm{PSL}(2, \mathbf{C}),$$

can be lifted to a representation $\widetilde{\mathrm{Hol}}: \pi_1(M) \rightarrow \mathrm{SL}(2, \mathbf{C})$ (see for instance [10]), and there is a one-to-one correspondence between these lifts and spin structures on M . Composing one of these lifts with a finite representation V of $\mathrm{SL}(2, \mathbf{C})$, we obtain a representation $\rho: \pi_1(M) \rightarrow \mathrm{SL}(V)$. Then we can consider the associated flat vector bundle E_ρ .

We will consider only *complex* and *finite dimensional* representations of $\mathrm{SL}(2, \mathbf{C})$. It is well known that for every positive integer n there exists only one complex irreducible representation V_n of $\mathrm{SL}(2, \mathbf{C})$ of dimension n . Moreover, V_n is $(n - 1)$ -th symmetric power of the standard representation $V_2 = \mathbf{C}^2$. Let

$$\rho_n: \pi_1(M) \rightarrow \mathrm{SL}(n, \mathbf{C}).$$

denote the representation ρ defined above for V_n .

A hyperbolic 3-manifold M is said to be *topologically finite* if it is the interior of a compact manifold \overline{M} . This is equivalent to say that $\pi_1(M)$ is finitely generated, by the proof of Marden’s conjecture [1, 9].

Along the paper we shall assume that M is *nonelementary*, which means, in the context of three manifolds, that its holonomy is an irreducible representation in $\mathrm{PSL}(2, \mathbf{C})$,

2010 Mathematics Subject Classification. Primary 57M50; Secondary 20C15.

Both authors partially supported by the Spanish Micinn through grant MTM2009-0759 and by the Catalan AGAUR through grant SGR2009-1207. The second author received the prize ICREA Acadèmia for excellence in research, funded by the Generalitat de Catalunya.

namely that there is no proper invariant subspace of \mathbf{C}^2 . Elementary manifolds have a simple geometry and topology (cf. Lemma 3.3) and the following results still hold and have a straightforward proof.

Theorem 0.1. *Let M be a complete, nonelementary, hyperbolic 3-manifold that is topologically finite, and $n \geq 2$. Then the inclusion $\partial\overline{M} \subset \overline{M}$ induces an injection,*

$$H^1(M; E_{\rho_n}) \hookrightarrow H^1(\partial\overline{M}; E_{\rho_n}),$$

with $\dim H^1(M; E_{\rho_n}) = (1/2) \dim H^1(\partial\overline{M}; E_{\rho_n})$, and an isomorphism

$$H^2(M; E_{\rho_n}) \cong H^2(\partial\overline{M}; E_{\rho_n}).$$

If M is a complete hyperbolic 3-manifold of finite volume with a single cusp, then $\partial\overline{M}$ is a torus. An analysis of the cohomology groups $H^*(\partial\overline{M}; E_{\rho})$ shows that all these groups vanish for the representations ρ_{2k} , with $k > 0$ (see Section 3.1). Hence, using Theorem 0.1 we get the following result.

Theorem 0.2. *Let M be a complete hyperbolic 3-manifold of finite volume with a single cusp. Then for $k \geq 1$ we have*

$$H^*(M; E_{\rho_{2k}}) = 0.$$

Notice that this theorem applies to hyperbolic knot exteriors in S^3 . For instance, it allows to compute Reidemeister torsions for hyperbolic knot exteriors.

Theorem 0.1 has applications to infinitesimal rigidity. The space of infinitesimal deformations of ρ_n is isomorphic to $H^1(M; E_{\text{Ad}\circ\rho_n})$, where

$$\text{Ad}: \text{SL}(n, \mathbf{C}) \rightarrow \text{Aut}(\mathfrak{sl}(n, \mathbf{C}))$$

is the adjoint representation.

The following theorem is an infinitesimal rigidity result for ρ_n in $\text{SL}(n, \mathbf{C})$ relative to the boundary. Its proof uses the decomposition of representation $\mathfrak{sl}(n, \mathbf{C})$ into irreducible factors, and will be given in Section 4.

Theorem 0.3. *Let M be a complete, hyperbolic, nonelementary and orientable 3-manifold that is topologically finite. If $\partial\overline{M}$ is the union of k tori and l surfaces of genus $g_1, \dots, g_l \geq 2$, and $n \geq 2$, then*

$$\dim_{\mathbf{C}} H^1(M; E_{\text{Ad}\circ\rho_n}) = k(n - 1) + \sum (g_i - 1)(n^2 - 1).$$

In particular, if M is closed then $H^1(M; E_{\text{Ad}\circ\rho_n}) = 0$. In addition, all nontrivial elements in $H^1(M; E_{\text{Ad}\circ\rho_n})$ are nontrivial in $H^1(\partial\overline{M}; E_{\text{Ad}\circ\rho_n})$ and have no L^2 representative.

When $n = 2$, this is Weil’s infinitesimal rigidity in the compact case, and Garland’s L^2 -infinitesimal rigidity in the noncompact case. This has been generalized to cone three manifolds by Hodgson–Kerckhoff [16], Weiss [28] and Bromberg [7].

Let $X(M, \text{SL}(n, \mathbf{C}))$ be the variety of characters of $\pi_1(M)$ in $\text{SL}(n, \mathbf{C})$. The character of ρ_n is denoted by χ_{ρ_n} . From the previous theorem and standard results on the variety of characters, we deduce:

Theorem 0.4. *Let M be a topologically finite, hyperbolic, nonelementary and orientable 3-manifold as in Theorem 0.3. If $n \geq 2$, then the character χ_{ρ_n} is a smooth point of $X(M, \text{SL}(n, \mathbf{C}))$ with tangent space $H^1(M; E_{\text{Ad} \circ \rho_n})$.*

For $n = 2$, this is Theorem 8.44 of Kapovich [18].

This paper is organized as follows. In Section 1 we recall some results about finite dimensional complex representations of $\text{SL}(2, \mathbf{C})$. Section 2 is devoted to Raghunathan’s vanishing theorem, from which Theorem 0.1 will follow. Theorem 0.2 is proved in Section 3, where we compute the cohomology of the ends and discuss some properties of lifts of representations. Section 4 deals with applications to infinitesimal and local rigidity, in particular we prove Theorems 0.3 and 0.4.

Appendix A reviews some results about principal bundles that are required in Section 2.

1. Finite dimensional complex representations of $\text{SL}(2, \mathbf{C})$

Irreducible complex finite dimensional representation of $\text{SL}(2, \mathbf{C})$ are well known to be the symmetric powers of the standard representation \mathbf{C}^2 . Therefore, there is exactly one irreducible representation in each dimension. Let V_n denote the irreducible complex n -th dimensional representation of \mathbf{C}^2 . We have $V_n = \text{Sym}^{n-1} V_2$, with the convention that Sym^0 is the base field.

The decomposition into irreducible factors of the tensor product of two given complex irreducible representation is given by the Clebsch–Gordan formula (cf. [11, §11.2]).

Theorem 1.1 (Clebsch–Gordan theorem). *For non-negative integer numbers n, k we have*

$$V_n \otimes V_{n+k} = \bigoplus_{i=0}^{n-1} V_{2(n-i)+k-1}.$$

Lemma 1.2. *Let V a finite dimensional complex representation of $\text{SL}(2, \mathbf{C})$. Then there exists a nondegenerate \mathbf{C} -bilinear invariant pairing*

$$\phi: V \times V \rightarrow \mathbf{C}.$$

Moreover, if V is irreducible this pairing is unique up to multiplication by nonzero scalars.

Proof. From the classification of the irreducible representations of $\mathrm{SL}(2, \mathbf{C})$, we deduce that V^* is isomorphic to V . Thus we get an invariant bilinear pairing by composing the isomorphism $V \times V \cong V^* \times V$ with the natural pairing between V^* and V . If V is irreducible, $V = V_n$, then the Clebsch–Gordan formula implies that $(V_n \otimes V_n)^* \cong V_n \otimes V_n$ has only one irreducible factor of dimension 1, so the bilinear pairing is unique in this case. \square

From this lemma we get (cf. [13, Section 2.2]):

Corollary 1.3. *Poincaré duality with coefficients in E_ρ holds true.*

Let $\mathrm{Ad}: \mathrm{SL}(n, \mathbf{C}) \rightarrow \mathrm{Aut}(\mathfrak{sl}(n, \mathbf{C}))$ denote the adjoint representation of $\mathrm{SL}(n, \mathbf{C})$. Composing it with the representation V_n we get a representation $\mathrm{SL}(2, \mathbf{C}) \rightarrow \mathrm{Aut}(\mathfrak{sl}(n, \mathbf{C}))$, which makes $\mathfrak{sl}(n, \mathbf{C})$ a $\mathrm{SL}(2, \mathbf{C})$ -module. Next we want to decompose this module into irreducible ones.

Lemma 1.4. *As $\mathrm{SL}(2, \mathbf{C})$ -modules, we have*

$$\mathfrak{sl}(n, \mathbf{C}) \cong V_{2n-1} \oplus V_{2n-3} \oplus \cdots \oplus V_3.$$

Proof. Consider the action of $\mathrm{SL}(2, \mathbf{C})$ on $\mathfrak{gl}(n, \mathbf{C})$ obtained by composing the n -dimensional representation V_n with the adjoint. We have the following isomorphisms of $\mathrm{SL}(2, \mathbf{C})$ -modules:

$$V_n \otimes V_n^* \cong \mathfrak{gl}(n, \mathbf{C}) \cong \mathfrak{sl}(n, \mathbf{C}) \oplus \mathbf{C},$$

where the factor \mathbf{C} corresponds to diagonal matrices. The result now follows from the Clebsch–Gordan formula applied to $V_n \otimes V_n^* \cong V_n \otimes V_n$. \square

2. Raghunathan’s cohomology vanishing theorem

The aim of this section is to prove Theorem 2.1 stated below. This theorem is a particular case of a theorem due to Raghunathan [24]. Before stating it, let us recall some facts.

The homogeneous manifold $\mathrm{SL}(2, \mathbf{C})/\mathrm{SU}(2)$ is endowed with a Riemannian structure using the Killing form on $\mathrm{SL}(2, \mathbf{C})$ (see Section 2.1 for details), which makes this space isometric to hyperbolic 3-dimensional space \mathbf{H}^3 .

Let Γ be a discrete torsion-free subgroup of $\mathrm{SL}(2, \mathbf{C})$, and $M = \Gamma \backslash \mathbf{H}^3$ the corresponding complete hyperbolic manifold. Let V be a finite dimensional representation of $\mathrm{SL}(2, \mathbf{C})$, and $\rho: \Gamma \rightarrow \mathrm{SL}(V)$ the induced representation. We can consider the associated flat vector bundle over M ,

$$E_\rho = \tilde{M} \times_\Gamma V.$$

The space of E_ρ -valued differential forms on M will be denoted by $\Omega^*(M; E_\rho)$. A $SU(2)$ -invariant hermitian product on V yields a well defined hermitian metric on E_ρ , and hence on $\Omega^*(M; E_\rho)$. In particular, it makes sense to talk about L^2 -forms of $\Omega^*(M; E_\rho)$ as those which are square summable.

Theorem 2.1 ([24]). *Let Γ be a discrete torsion-free subgroup of $SL(2, \mathbf{C})$. Let V be an irreducible finite dimensional complex representation of $SL(2, \mathbf{C})$, and $\rho: \Gamma \rightarrow SL(V)$ the induced representation. Then, for $p = 1, 2$, every closed L^2 -form in $\Omega^p(\Gamma \backslash \mathbf{H}^3; E_\rho)$ is exact.*

As an immediate corollary of Theorem 2.1 we get a particular case of Raghunathan’s cohomology vanishing theorem.

Corollary 2.2 ([24]). *Let M be a closed hyperbolic three-manifold. If V is an irreducible finite dimensional complex representation of $SL(2, \mathbf{C})$, then*

$$H^1(M; E_\rho) = 0.$$

REMARK. Raghunathan’s theorem applies to lattices of a semisimple Lie group G , and a broader family of representations, see [24].

From Theorem 2.1 we can easily deduce Theorem 0.1.

Proof of Theorem 0.1. We have $M = \Gamma \backslash \mathbf{H}^3$ for some discrete torsion-free subgroup Γ of $SL(2, \mathbf{C})$. If M is compact then the result is clear from Theorem 2.1, so we can assume M is noncompact. The space $H^p(\overline{M}, \partial \overline{M}; E_\rho)$ can be identified with the cohomology group of compactly supported E_ρ -valued p -forms on M ; hence, an element $[\alpha] \in H^p(\overline{M}, \partial \overline{M}; E_\rho)$ is represented by a closed compactly supported form α on M . Therefore, Theorem 2.1 implies that for $p = 1, 2$ the image of $[\alpha]$ under the map $H^p(\overline{M}, \partial \overline{M}; E_\rho) \rightarrow H^p(M; E_\rho)$ induced by the inclusion is zero.

The theorem now follows from the long exact sequence of the pair, and Poincaré duality. Indeed the long exact sequence of the pair $(\overline{M}, \partial \overline{M})$ gives short exact sequences

$$\begin{aligned} 0 \rightarrow H^1(M; E_{\rho_n}) &\rightarrow H^1(\partial \overline{M}; E_{\rho_n}) \rightarrow H^2(\overline{M}, \partial \overline{M}; E_{\rho_n}) \rightarrow 0, \\ 0 \rightarrow H^2(M; E_{\rho_n}) &\rightarrow H^2(\partial \overline{M}; E_{\rho_n}) \rightarrow H^3(\overline{M}, \partial \overline{M}; E_{\rho_n}). \end{aligned}$$

By Poincaré duality we have $\dim H^1(M; E_{\rho_n}) = \dim H^2(\overline{M}, \partial \overline{M}; E_{\rho_n})$, and $\dim H^3(\overline{M}, \partial \overline{M}; E_{\rho_n}) = \dim H^0(M; E_{\rho_n}) = 0$, by Lemma 3.5. □

Raghunathan’s original proof of the theorem, a particular case of which is Theorem 2.1, uses two results as starting points. The first one the following theorem due to Andreotti and Vesentini [2]. Although the original theorem is for complex manifolds, there is an adaptation of Garland [12, Theorem 3.22] to the real case.

Theorem 2.3 (Andreotti–Vesentini [2], Garland [12]). *Suppose that M is complete. Assume that there exists $c > 0$ such that for every $\alpha \in \Omega^p(M; E)$ with compact support $(\Delta\alpha, \alpha) \geq c(\alpha, \alpha)$, where (\cdot, \cdot) denotes the hermitian (or inner) product on the space of E -valued forms. Then every square-integrable closed p -form is exact.*

The second point is the work of Matsushima–Murakami concerning the theory of harmonic forms in a locally symmetric manifold [21]. One of the goals of that work consists in proving a Weitzenböck formula for the Laplacian. Using that formula, the strong-positivity hypothesis of the Laplacian required in Theorem 2.3 can be proved by establishing the positivity of a certain linear operator defined on a finite dimensional space, see Subsection 2.1. Although this is an important conceptual reduction, it remains to prove the positivity of that operator. Raghunathan was able to prove it for a large family of locally symmetric manifolds and representations, see [24].

The rest of this section is divided into two parts. The first one is a review of the work of Matsushima and Murakami concerning the Laplacian of a locally symmetric manifold. The material presented here is almost entirely based on Matsushima–Murakami [21], and Raghunathan’s book [25]. Although it does not bring in a new conceptual approach, seeking completeness, we hope the exposition given here will be more accessible to the non-expert. Using this material, we give a simple proof of Theorem 2.1 in Subsection 2.2.

2.1. Review of harmonic forms on a locally symmetric manifold. Let G be a connected semisimple Lie group and $K < G$ a maximal compact subgroup of G . The respective Lie algebras are denoted by \mathfrak{g} and \mathfrak{k} , with the convention that they are the Lie algebras of left invariant vector fields on G and K , respectively.

Let B denote the Killing form of \mathfrak{g} . We recall that it is defined by

$$B(V, W) = \text{tr}(\text{ad}_V \circ \text{ad}_W),$$

for $V, W \in \mathfrak{g}$. Cartan’s criterion implies that B is nondegenerate if, and only if, \mathfrak{g} is semisimple. In that case, we have a canonical decomposition $\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{k}$, where \mathfrak{m} is the orthogonal complement to \mathfrak{k} respect to B . This decomposition satisfies the following properties: B is negatively defined on \mathfrak{k} ; B positively defined on \mathfrak{m} ; $[\mathfrak{k}, \mathfrak{m}] \subset \mathfrak{m}$; and $[\mathfrak{m}, \mathfrak{m}] \subset \mathfrak{k}$.

The Killing form defines a pseudo-Riemannian metric on G , which is invariant by the action of G by right translations, and is positively (resp. negatively) defined on \mathfrak{m} (resp. \mathfrak{k}). Therefore, the Killing form defines a Riemannian metric on the homogeneous space $X = G/K$. Note that G acts on the left on X by orientation preserving isometries.

Let Γ be a discrete subgroup of G that acts freely on X . Since Γ acts by isometries, the quotient $M = \Gamma \backslash X$ is a Riemannian manifold. It is said that M is a *locally symmetric manifold*.

For our purposes, it will be convenient to regard the universal covering $X \rightarrow M$ as a principal bundle over M with structure group Γ . We follow the convention that the action of the structure group of a principal bundle is on the right. Hence we only need to convert the action of Γ into a right action (if $g \in \Gamma$, then $x \cdot g = g^{-1} \cdot x$, for $x \in X$). We will also regard X as a flat bundle.

Consider the G -principal bundle $P = X \times_{\Gamma} G$ over M (see Appendix A for notation) endowed with the flat connection induced from the trivial connection of the product $X \times G$. We can embed X on P using the section $X \rightarrow X \times G$ whose second coordinate is constant and equal to the identity element. We can think of X as a reduction of the structure group. Obviously, the horizontal leaves of X are also horizontal leaves of P , so the connection on P is reducible to X .

On the other hand, the principal bundle P has a canonical reduction of its structure group from G to K . In order to get such a reduction, consider the embedding $i: G \hookrightarrow X \times G$ given by $i(g) = (gK, g)$. The image of G by this embedding is invariant by the bundle action of K , so it defines an embedding $\Gamma \backslash G \hookrightarrow X \times_{\Gamma} G$, which will be also denoted by i . Therefore, $Q = i(\Gamma \backslash G)$ is a reduction of the structure group.

The connection defined on P is not reducible to Q , because its horizontal distribution is not tangent to Q (a curve on $X \times G$ whose second component is constant, gives an horizontal curve on P ; hence, if the horizontal distribution were tangent to Q , this curve would be contained in Q , and this does not happen). Nevertheless, since the action of K on \mathfrak{g} respects the decomposition $\mathfrak{g} = \mathfrak{m} \oplus \mathfrak{k}$, we can state the following.

Proposition 2.4. *Let $\eta \in \Omega^1(P; \mathfrak{g})$ be the connection form of the connection defined on P above. Put $\eta = \eta_{\mathfrak{m}} + \eta_{\mathfrak{k}}$, where $\eta_{\mathfrak{m}}$ and $\eta_{\mathfrak{k}}$ are the \mathfrak{m} and \mathfrak{k} components of η respectively. Then, the restriction of $\eta_{\mathfrak{k}}$ to Q is a connection form on Q .*

OBSERVATION. We can identify \mathfrak{g} with the space of vector fields on $\Gamma \backslash G$ that are projection of left invariant vector fields on G . In what follows, we will tacitly do this identification.

Let $\omega \in \Omega^1(\Gamma \backslash G; \mathfrak{g})$ be the left Maurer–Cartan form of G . It is easily checked that $i^*(\eta) = \omega$. Hence, if we decompose $\omega = \omega_{\mathfrak{m}} + \omega_{\mathfrak{k}}$ into the \mathfrak{m} -component and the \mathfrak{k} -component, $\omega_{\mathfrak{k}}$ is the connection form of the connection defined on $\Gamma \backslash G$, and the horizontal distribution is given by \mathfrak{m} .

Consider a finite linear representation $\rho: G \rightarrow \text{Aut}(V)$, and the associated vector bundle $E = X \times_{\Gamma} V$ (note that E is canonically identified with $P \times_G V$ and $Q \times_K V$).

The flat connection on P defines an exterior covariant differential d_{ρ} on the space $\Omega^*(M; E)$. Via the canonical isomorphism between $\Omega^*_{\text{Hor}}(\Gamma \backslash G; V)^K$ and $\Omega^*(M; E)$, we can transfer the operator d_{ρ} to an operator D_{ρ} , in such a way that this isomorphism is a chain complex isomorphism. If we denote by D the exterior covariant differential defined by the connection $\omega_{\mathfrak{k}}$ on Q , then the relation between D and D_{ρ} is given by the following proposition.

Proposition 2.5. *Let α be a form in $\Omega_{\text{Hor}}^r(\Gamma \backslash G; V)^K$. We have the following decomposition*

$$D_\rho \alpha = D\alpha + T\alpha,$$

where $T\alpha = \rho(\omega_m) \wedge \alpha$.

Proof. On P the differential covariant is given by $d\alpha + \rho(\eta) \wedge \alpha$ (see Proposition A.2). Hence, if we transfer it to Q via i , we get $D_\rho \alpha = d\alpha + \rho(i^* \eta) \wedge \alpha$, and the proposition follows from the fact that $i^* \eta = \omega$. □

Let's fix an orientation on \mathfrak{k} and \mathfrak{m} , and take an orthonormal basis for \mathfrak{g} , $(X_1, \dots, X_n, Y_1, \dots, Y_m)$, such that (X_1, \dots, X_n) and (Y_1, \dots, Y_m) are positively oriented orthonormal bases for \mathfrak{k} and \mathfrak{m} , respectively. Here, orthonormality means that

$$B(X_i, X_j) = -\delta_{ij}, \quad B(Y_i, Y_j) = \delta_{ij}, \quad B(X_i, Y_j) = 0.$$

NOTATION. We will follow the following conventions. Let V be a finite dimensional vector space. If e_1, \dots, e_n is a basis for V , then its dual basis will be denoted by $e^1, \dots, e^n \in V^*$, with $e^i(e_j) = \delta_{ij}$. If $A \in \otimes^r V^*$ is an r -times covariant tensor, then its components relative to the basis defined by e^1, \dots, e^n will be denoted by A_{i_1, \dots, i_r} . Concerning the exterior product on $\wedge^* V^*$, we will follow the convention such that $e^1 \wedge \dots \wedge e^n$ is the determinant. We will also use Einstein notation. Hence, given $\alpha \in \wedge^r V^*$, we have $\alpha = \alpha_{i_1, \dots, i_r} e^{i_1} \otimes \dots \otimes e^{i_r}$, where α_{i_1, \dots, i_r} are scalars satisfying $\alpha_{i_{\sigma(1)}, \dots, i_{\sigma(r)}} = \text{sgn}(\sigma) \alpha_{i_1, \dots, i_r}$, for any permutation $\sigma \in \Sigma_r$. Then we also have

$$\alpha = \sum_{1 \leq i_1 < \dots < i_r \leq r} \alpha_{i_1, \dots, i_r} e^{i_1} \wedge \dots \wedge e^{i_r} = \frac{1}{r!} \alpha_{i_1, \dots, i_r} e^{i_1} \wedge \dots \wedge e^{i_r}.$$

From now on, all the tensors will be written in the basis of \mathfrak{g} given by $\{X_1, \dots, X_n, Y_1, \dots, Y_m\}$.

Proposition 2.6. *For $\alpha \in \Omega_{\text{Hor}}^r(\Gamma \backslash G; V)^K$, the operators D and T are given by the following equations.*

$$(1) \quad (D\alpha)_{i_1, \dots, i_{r+1}} = \sum_{k=1}^{r+1} (-1)^{k+1} Y_{i_k} \alpha_{i_1, \dots, \hat{i}_k, \dots, i_{r+1}},$$

$$(2) \quad (T\alpha)_{i_1, \dots, i_{r+1}} = \sum_{k=1}^{r+1} (-1)^{k+1} \rho(Y_{i_k}) \alpha_{i_1, \dots, \hat{i}_k, \dots, i_{r+1}}.$$

Proof. Put $\alpha = (1/r!) \alpha_{i_1, \dots, i_r} Y^{i_1} \wedge \dots \wedge Y^{i_r}$. By definition, $D\alpha$ is the horizontal component of $d\alpha$. It is immediate that dY^k has no horizontal component: $dY^k(Y_i, Y_j) =$

$Y^k([Y_i, Y_j]) = 0$. Hence, $D\alpha = (1/r!)Y_j\alpha_{i_1, \dots, i_r} \otimes Y^j \wedge Y^{i_1} \wedge \dots \wedge Y^{i_r}$. Rearranging the indices we get equation (1). The other equation follows immediately from the definition of T . \square

Let us define the forms $\Omega_K = X^1 \wedge \dots \wedge X^n$ and $\Omega_M = Y^1 \wedge \dots \wedge Y^m$. It is clear that these forms are independent of the orthonormal bases chosen. Hence, Ω_K and Ω_M are well defined forms on $\Gamma \backslash G$. Note that Ω_K is vertical and Ω_M is horizontal, and both are right K -invariant (it is a consequence of the fact the right action of K on \mathfrak{g} leaves both the Killing form and the decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$ invariant). Observe that Ω_M defines a volume form on M , which is compatible with the metric structure of M .

Next we want to define an inner product on the fibers of E . In order to do that, fix a K -invariant inner product $\langle \cdot, \cdot \rangle_V$ on V , and use it to define a metric on the fibers of $E = Q \times_K V$. Then define an inner product on $\Omega^*(M; E)$ as usual: if $\alpha, \beta \in \Omega^*(M; E)$ then

$$(\alpha, \beta) = \int_M \langle \alpha(x), \beta(x) \rangle_x \Omega_M,$$

where $\langle \cdot, \cdot \rangle_x$ is the inner product defined on the fiber E_x , and Ω_M is interpreted as a form on M . On the other hand, we can define the inner product of two forms $\tilde{\alpha}, \tilde{\beta} \in \Omega^r_{\text{Hor}}(\Gamma \backslash G; V)^K$ by

$$(\tilde{\alpha}, \tilde{\beta}) = \frac{1}{\mu(K)} \int_{\Gamma \backslash G} \langle \tilde{\alpha}(u), \tilde{\beta}(u) \rangle_u \Omega_K \wedge \Omega_M,$$

where $\langle \cdot, \cdot \rangle_u$ is the inner product on $\bigwedge^r H^* \otimes V$ induced by the Killing form, and the inner product on V , and $\mu(K) = \int_K \Omega_K$ the volume of K . Proposition A.4, gives the relation between these two products.

Proposition 2.7. *The canonical isomorphism between $\Omega^*_{\text{Hor}}(\Gamma \backslash G; V)^K$ and $\Omega^*(M; E)$ is an isometry.*

Using the Hodge dual operator on the horizontal bundle

$$* : \Omega^r_{\text{Hor}}(\Gamma \backslash G; V)^K \rightarrow \Omega^{n-r}_{\text{Hor}}(\Gamma \backslash G; V)^K,$$

we can give a characterization of the formal adjoint of the operators D and T .

Proposition 2.8. *Let $\alpha \in \Omega^r_{\text{Hor}}(\Gamma \backslash G; V)^K$ with compact support. Then,*

(3)
$$D^* \alpha = (-1)^r *^{-1} D * \alpha,$$

(4)
$$T^* \alpha = (-1)^{r-1} *^{-1} \overline{\rho(\omega)}^f \wedge (*\alpha).$$

Proof. We want to use Proposition A.5. We claim that

$$\int_P D\alpha \wedge \beta \wedge \Omega_K = (-1)^r \int_P \alpha \wedge D\beta \wedge \Omega_K,$$

for α and β forms of $\Omega_{\text{Hor}}^*(\Gamma \backslash G; V)$ with compact support of degree $r - 1$ and $m - r$ respectively. Indeed, since $D\alpha$ is the horizontal component of $d\alpha$, we have $D\alpha \wedge \Omega_K = d\alpha \wedge \Omega_K$. Then,

$$d(\alpha \wedge \beta \wedge \Omega_K) = d\alpha \wedge \beta \wedge \Omega_K + (-1)^{r-1} \alpha \wedge d\beta \wedge \Omega_K,$$

for Ω_K being closed. Therefore, by Stokes' theorem we get the equality we wanted to prove. Now, Proposition A.5 gives Formula (3).

Now, let us prove (4). By Proposition A.5, it suffices to prove that

$$(\rho(\omega) \wedge \alpha) \wedge \beta = (-1)^{r-1} \alpha \wedge (\rho(\omega)^* \wedge \beta).$$

If we take an orthonormal basis for V , then α and β are column vectors of forms of degree $r - 1$ and $m - r$ respectively, and $\rho(\omega)$ a matrix of one forms. Hence, in this basis $(\rho(\omega) \wedge \alpha) \wedge \beta$ is $(\rho(\omega)\alpha)^t \bar{\beta}$, but $(\rho(\omega)\alpha)^t \beta = (-1)^{r-1} \alpha^t \rho(\omega)^t \bar{\beta}$, as we wanted to prove. □

A similar proof of Proposition 2.6, using the formulae found in the previous proposition, gives the following.

Proposition 2.9. *For $\alpha \in \Omega_{\text{Hor}}^r(\Gamma \backslash G; V)^K$ with compact support, the operators D^* and T^* are given by the following equations.*

$$(5) \quad (D^*\alpha)_{i_1, \dots, i_{r-1}} = \sum_{k=1}^m -Y_k \alpha_{k, i_1, \dots, i_{r-1}},$$

$$(6) \quad (T^*\alpha)_{i_1, \dots, i_{r-1}} = \sum_{k=1}^m \rho(Y_k) \alpha_{k, i_1, \dots, i_{r-1}}.$$

Lemma 2.10. *If the inner product on V is symmetric respect to the action of \mathfrak{m} , then the operator $S = TD^* + T^*D + DT^* + D^*T$ is zero for every form with compact support.*

Before proving the lemma, we need the following result.

Lemma 2.11. *For every function f with compact support, and $Y \in \mathfrak{g}$,*

$$\int_{\Gamma \backslash G} (Yf) \Omega_M \wedge \Omega_K = 0.$$

Proof. Since Y is an infinitesimal isometry we have $L_Y(f\Omega_M \wedge \Omega_K) = (Yf)\Omega_M \wedge \Omega_K$. On the other hand, the formula $L_Y = i_Y \circ d + d \circ i_Y$ gives $L_Y(f\Omega_M \wedge \Omega_K) = d(i_Y f \Omega_M \wedge \Omega_K)$, and Stokes' theorem implies

$$0 = \int_{\Gamma \backslash G} L_Y(f\Omega_M \wedge \Omega_K) = \int_{\Gamma \backslash G} (Yf)\Omega_M \wedge \Omega_K,$$

as we wanted to prove. □

Proof of Lemma 2.10. Since S is a self-adjoint operator, $S = 0$ if, and only if, $(S\alpha, \alpha) = 0$ for every α with compact support. Let's take $\alpha \in \Omega_{\text{Hor}}^*(\Gamma \backslash G; V)^K$ with compact support. We must show that

$$(S\alpha, \alpha) = (D\alpha, T\alpha) + (T\alpha, D\alpha) + (D^*\alpha, T^*\alpha) + (T^*\alpha, D^*\alpha) = 0.$$

Observe that it suffices to prove that $(D\alpha, T\alpha) + (D^*\alpha, T^*\alpha) = 0$. Moreover, using the m -symmetry of the inner product and the fact that the Hodge $*$ operator is an isometry, we must prove $(D\alpha, T\alpha) + (D(*\alpha), T(*\alpha)) = 0$. Let's compute $(D\alpha, T\alpha)$. Put $\alpha = \alpha_{i_1, \dots, i_r} \otimes Y^{i_1} \wedge \dots \wedge Y^{i_r}$. If we use the expression of D and T given in Proposition 2.6, we see that $(D\alpha, T\alpha)$ is the sum of terms of the form

$$(-1)^{i+j} \int_{\Gamma \backslash G} \langle Y_{i_j} \alpha_{i_1, \dots, \hat{i}_j, \dots, i_{r+1}}, \rho(Y_{i_k}) \alpha_{i_1, \dots, \hat{i}_k, \dots, i_{r+1}} \rangle_V d\mu_G.$$

It is convenient to group the summands according to whether the avoided sub-indices \hat{i}_j and \hat{i}_k are equal or not. Therefore, one term is a sum of factors of the form

$$\int_{\Gamma \backslash G} \langle Y_j \alpha_{i_1, \dots, i_r}, \rho(Y_j) \alpha_{i_1, \dots, i_r} \rangle_V d\mu_G, \quad j \notin \{i_1, \dots, i_r\},$$

and the rest is a sum of terms of the form

$$(7) \quad (-1)^{j+k} \int_{\Gamma \backslash G} \langle Y_{i_j} \alpha_{i_1, \dots, \hat{i}_j, \dots, i_k, \dots, i_r}, \rho(Y_{i_k}) \alpha_{i_1, \dots, i_j, \dots, \hat{i}_k, \dots, i_r} \rangle d\mu_G,$$

with $i_j \neq i_k$. We can apply this formula to $*\alpha$ to compute $(D(*\alpha), T(*\alpha))$. The formula we get is just the above formula with the range of the indices changed by their complementary; that is, on one hand we get terms of the form

$$\int_{\Gamma \backslash G} \langle Y_j \alpha_{i_1, \dots, i_r}, \rho(Y_j) \alpha_{i_1, \dots, i_r} \rangle_V d\mu_G, \quad j \in \{i_1, \dots, i_r\},$$

and on the other hand terms of the form

$$(-1)^{j+k} \int_{\Gamma \backslash G} \langle Y_{i_k} \alpha_{i_1, \dots, \hat{i}_j, \dots, i_k, \dots, i_r}, \rho(Y_{i_j}) \alpha_{i_1, \dots, i_j, \dots, \hat{i}_k, \dots, i_r} \rangle d\mu_G,$$

for $i_j \neq i_k$. By Lemma 2.11, this last term is the opposite of 7. Hence, it suffices to prove that for every $Y \in \mathfrak{m}$, and $f \in \mathcal{C}(\Gamma \setminus G; V)$, we have

$$\int_{\Gamma \setminus G} \langle Yf, \rho(Y)f \rangle_V d\mu_G = 0.$$

But it is also an immediate consequence of Lemma 2.11 and the symmetry of $\rho(Y)$. The lemma now follows from the fact that $(D^*\alpha, T^*\alpha) = (D(*\alpha), T(*\alpha))$. \square

Corollary 2.12 (Matsushima–Murakami formula). *Assume the inner product on V is symmetric respect to the action of \mathfrak{m} . Then*

$$\Delta_\rho = \Delta + H_\rho,$$

where $\Delta = DD^* + D^*D$, and $H_\rho = TT^* + T^*T$.

Proof. We have $\Delta_\rho = D_\rho D_\rho^* + D_\rho^* D_\rho = \Delta + H_\rho + S$, and Lemma 2.10. \square

Let's denote by \mathbf{T} , \mathbf{T}^* , \mathbf{H}_ρ the restriction to $V \otimes \wedge^p \mathfrak{m}^*$ of T , T^* and H_ρ respectively. Since T is an operator of degree zero, essentially all information of T , T^* and H_ρ is contained in \mathbf{T} , \mathbf{T}^* , \mathbf{H}_ρ . In particular, H_ρ is positive definite if and only \mathbf{H}_ρ is so.

Proposition 2.13. *Let $\alpha \in V \otimes \wedge^p \mathfrak{m}^*$. Then we have,*

$$(\mathbf{H}_\rho \alpha)_{i_1, \dots, i_r} = \sum_{j=1}^m \rho(Y_j)^2 \alpha_{i_1, \dots, i_r} + \sum_{k=1}^r \sum_{j=1}^m (-1)^{k+1} \rho([Y_{i_k}, Y_j]) \alpha_{j, i_1, \dots, \hat{i}_k, \dots, i_r}.$$

Proof. Put $\beta_{i_1, \dots, i_{r+1}} = (T\alpha)_{i_1, \dots, i_{r+1}}$ and $\gamma_{i_1, \dots, i_{r-1}} = (T^*\alpha)_{i_1, \dots, i_{r-1}}$. Then, on one hand we have

$$\begin{aligned} (TT^*\alpha)_{i_1, \dots, i_r} &= \sum_{k=1}^r (-1)^{k+1} \rho(Y_{i_k}) \gamma_{i_1, \dots, \hat{i}_k, \dots, i_r} \\ &= \sum_{k=1}^r (-1)^{k+1} \rho(Y_{i_k}) \sum_{j=1}^m \rho(Y_j) \alpha_{j, i_1, \dots, \hat{i}_k, \dots, i_r}, \end{aligned}$$

and on the other hand,

$$\begin{aligned} (T^*T\alpha)_{i_1, \dots, i_r} &= \sum_{j=1}^m \rho(Y_j) \beta_{j, i_1, \dots, i_r} \\ &= \sum_{j=1}^m \rho(Y_j) \left(\rho(Y_j) \alpha_{i_1, \dots, i_r} + \sum_{k=1}^r (-1)^k \rho(Y_{i_k}) \alpha_{j, i_1, \dots, \hat{i}_k, \dots, i_r} \right). \end{aligned}$$

And the proposition follows. □

2.2. Proof of Theorem 2.1. We want to apply the criterion of Andreotti–Vesentini of Theorem 2.3. For this purpose, we will use Matsushima–Murakami’s formula (Corollary 2.12) for the representation of $SL(2, \mathbb{C})$. Since for every compactly supported 1-form α

$$(\Delta(\alpha), \alpha) = (D(\alpha), D(\alpha)) + (D^*(\alpha), D^*(\alpha)) \geq 0,$$

using Corollary 2.12, the criterion of Theorem 2.3 reduces to show that $(H_\rho(\alpha), \alpha) \geq c(\alpha, \alpha)$ for some uniform $c > 0$ and every compactly supported 1-form α .

Notice that since the linear operator H_ρ on 1-forms is induced from a linear operator \mathbf{H}_ρ on $V \otimes \mathfrak{m}^*$, if \mathbf{H}_ρ is positive definite, then there is a positive constant c so that $(H_\rho(\alpha), \alpha) \geq c(\alpha, \alpha)$ holds for every compactly supported one form α . The proof will follow from Lemma 2.14.

In order to apply Matsushima–Murakami’s formula to the representations of $SL(2, \mathbb{C})$, first we need to choose an orthonormal basis for $\mathfrak{su}(2)$ respect to the Killing form (in fact, respect to a constant multiple of it). Let’s define

$$X_1 = \begin{pmatrix} \mathbf{i} & 0 \\ 0 & -\mathbf{i} \end{pmatrix}, \quad X_2 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad X_3 = \begin{pmatrix} 0 & \mathbf{i} \\ \mathbf{i} & 0 \end{pmatrix}.$$

Then (X_1, X_2, X_3) is an orthonormal basis for $\mathfrak{su}(2)$. The orthogonal complement to $\mathfrak{su}(2)$ with respect to the Killing form is given by $Y_k = \mathbf{i}X_k$, for $k = 1, 2, 3$. On the other hand, we have $[X_i, X_{i+1}] = 2X_{i+2}$, for $i = 1, 2, 3$, where the indices are taken modulo 3.

Lemma 2.14. *Let $\rho: \mathfrak{sl}(2, \mathbb{C}) \rightarrow \text{End}(V)$ a complex finite dimensional irreducible representation, $\dim(V) \geq 2$. Then the operator \mathbf{H}_ρ is positively defined on degree 1 and 2.*

Proof. Since $\mathbf{H}_\rho = \mathbf{T}_\rho \mathbf{T}_\rho^* + \mathbf{T}_\rho^* \mathbf{T}_\rho$, to show that \mathbf{H}_ρ is positive definite is equivalent to show that its kernel is trivial. Let $\alpha \in V \otimes \mathfrak{m}^*$. We have $\alpha = \sum_{i=1}^3 \alpha_i \otimes Y^i$, with $\alpha_i \in V$. Assume $\mathbf{H}_\rho \alpha = 0$. Then $\mathbf{T}_\rho \alpha = 0$ must vanish too, and from Proposition 2.6 (2) we obtain

$$(8) \quad 0 = (\mathbf{T}_\rho \alpha)(Y_i, Y_j) = \rho(Y_i)\alpha_j - \rho(Y_j)\alpha_i, \quad i, j = 1, 2, 3.$$

Proposition 2.13 yields

$$(\mathbf{H}_\rho \alpha)(Y_j) = \sum_{k=1}^3 (\rho(Y_k)^2 \alpha_j + \rho([Y_j, Y_k])\alpha_k).$$

Taking the indices modulo 3, and using the Lie algebra relations, we get

$$\begin{aligned} \sum_{k=1}^3 \rho([Y_j, Y_k])\alpha_k &= \rho([Y_j, Y_{j+1}])\alpha_{j+1} + \rho([Y_j, Y_{j+2}])\alpha_{j+2} \\ &= 2(\rho(-X_{j+2})\alpha_{j+1} + \rho(X_{j+1})\alpha_{j+2}) \\ &= 2i(\rho(Y_{j+2})\alpha_{j+1} - \rho(Y_{j+1})\alpha_{j+2}). \end{aligned}$$

Notice that in the last equality we have used the complex structure. Hence, using (8), we get $(\mathbf{H}_\rho\alpha)(Y_j) = \sum_{k=1}^3 \rho(Y_k)^2\alpha_j$, and then

$$\begin{aligned} 0 = \langle \mathbf{H}_\rho\alpha, \alpha \rangle &= \sum_{j=1}^3 \left\langle \sum_{k=1}^3 \rho(Y_k)^2\alpha_j, \alpha_j \right\rangle \\ &= \sum_{j,k=1}^3 \langle \rho(Y_k)\alpha_j, \rho(Y_k)\alpha_j \rangle, \end{aligned}$$

that implies $\rho(Y_j)\alpha_k = 0$ for $j, k = 1, 2, 3$. Hence, for a fixed k , we have $\rho(Z)\alpha_k = 0$ for every $Z \in \mathfrak{sl}(2, \mathbf{C})$. Since we are assuming that ρ is irreducible and nontrivial, we get $\alpha_k = 0$ for all k . It proves the lemma in degree 1. Since $\mathfrak{m}^* \cong \bigwedge^2 \mathfrak{m}^*$, the same proof holds true in degree 2. □

3. Cohomology of the ends and lifts of the holonomy

Assume that M is a noncompact, nonelementary, orientable hyperbolic manifold with finite topology, in particular it is the interior of a compact manifold with boundary $\partial\overline{M}$. The aim of this section is to analyse the cohomology groups of $H^*(\partial\overline{M}, E_{\rho_n})$. This will be done in Subsection 3.1. When the ends of the manifold are cusps, this cohomology happens to be related to the lift of the holonomy, that we study in Subsection 3.2. Finally, this is used to prove Theorem 0.2.

3.1. Cohomology of the ends.

DEFINITION. Let G be a group acting on a vector space V . The *subspace of invariants* of V , denoted by V^G , is the subspace consisting of elements of V that are fixed by G . That is,

$$V^G = \{v \in V \mid g \cdot v = v, \text{ for all } g \in G\}.$$

Lemma 3.1. *Let F be a connected component of $\partial\overline{M}$. For every $n > 1$ we have,*

$$\begin{aligned} \dim_{\mathbf{C}} H^0(F; E_{\rho_n}) &= \dim_{\mathbf{C}} V_n^{\pi_1(F)}, \\ \dim_{\mathbf{C}} H^1(F; E_{\rho_n}) &= 2 \dim_{\mathbf{C}} V_n^{\pi_1(F)} - n\chi(F), \\ \dim_{\mathbf{C}} H^2(F; E_{\rho_n}) &= \dim_{\mathbf{C}} V_n^{\pi_1(F)}. \end{aligned}$$

Proof. Since F is a $K(\pi_1(F), 1)$ space, $H^0(F; E_{\rho_n}) = H^0(\pi_1(F); E_{\rho_n})$, and this is identified with $V_n^{\pi_1(F)}$. It proves the first equality. The third one follows from Poincaré duality, and the second one from an Euler characteristic argument. \square

Therefore, all the cohomological information comes from the subspace of invariants $V_n^{\pi_1(F)}$. We distinguish two cases according to whether F has genus $g \geq 2$, or F is a torus. In order to analyse the case when F is a torus, we make the following definition. If we have a torus $T^2 \subset \partial\overline{M}$, then the holonomy maps $\pi_1(T^2)$ to a parabolic subgroup; hence, up to conjugation every element in $\pi_1(T^2)$ is mapped by a lift of the holonomy representation to

$$\pm \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix}.$$

DEFINITION. Let us fix a lift to $SL(2, \mathbf{C})$ of the holonomy representation. We say that this lift is *positive* on $\pi_1(T^2)$ if every element of $\pi_1(T^2)$ has trace $+2$.

Proposition 3.2. *Let F a connected component of $\partial\overline{M}$, and $n > 1$. If F has genus $g \geq 2$, then $V_n^{\pi_1(F_g)} = 0$.*

If F is a torus T^2 , then we have the following cases,

$$V_n^{\pi_1(T^2)} = \begin{cases} 0 & \text{for } n \text{ even and a nonpositive lift;} \\ \mathbf{C} & \text{for } n \text{ even and a positive lift;} \\ \mathbf{C} & \text{for } n \text{ odd.} \end{cases}$$

Before proving it, we need the following lemmas. The first one can be found in standard references about Kleinian groups (cf. [18]):

Lemma 3.3. *Let M be a hyperbolic three manifold. Then the following are equivalent:*

- M is elementary (its holonomy is reducible in $PSL(2, \mathbf{C})$).
- $\pi_1(M)$ is abelian.
- M is homeomorphic to either the product of the plane with a circle, $\mathbf{R}^2 \times S^1$, or to the product of a 2-torus with a line, $S^1 \times S^1 \times \mathbf{R}$.

Lemma 3.4. *Let F be a connected component of $\partial\overline{M}$. If F has genus $g \geq 2$, then $\widetilde{Hol}(\pi_1(F))$ is an irreducible subgroup of $SL(2, \mathbf{C})$.*

Proof. When F is π_1 -injective (i.e. when $\pi_1(F)$ injects into $\pi_1(M)$) then the holonomy restricts to a discrete and faithful representation of $\pi_1(F)$, and irreducibility follows because $\pi_1(F)$ is nonabelian. Otherwise, when F is not π_1 -injective, according to Bonahon [5] and McCullough–Miller [22] there are two possibilities: either M is a *handlebody* or F is a boundary component of a *characteristic compression body* $C \subseteq$

M . A handlebody is the result of attaching one handles to a 3-ball; in particular when M is a handlebody then $\pi_1(F)$ surjects onto $\pi_1(M)$, thus $\text{Hol}(\pi_1(F)) = \text{Hol}(\pi_1(M))$ and irreducibility comes from the hypothesis that M is nonelementary. Next, assume that F is the positive boundary of a characteristic compression body C , namely $C \subseteq M$ is a codimension 0 closed submanifold, whose boundary splits as a union $\partial C = \partial_- C \cup \partial_+ C$, so that $\partial_+ C = F$, the components of $\partial_- C$ are π_1 -injective in M , and C is the result of gluing 1-handles to $\partial_- C \times [0, 1]$ along $\partial_- C \times \{1\}$. In particular $\pi_1(F)$ surjects onto $\pi_1(C)$ and $\text{Hol}(\pi_1(F)) = \text{Hol}(\pi_1(C))$. Thus, if $F = \partial_+ C$ and one of the components of $\partial_- C$ has genus ≥ 2 , then we are done by the π_1 -injective case. Finally if $F = \partial_+ C$ and all components of $\partial_- C$ are tori, since incompressible tori in M are boundary parallel, then the inclusion $C \subseteq M$ is a homotopy equivalence. Thus $\pi_1(F)$ surjects onto $\pi_1(M)$ and irreducibility follows again because M is nonelementary. \square

Lemma 3.5. *Let M be a nonelementary, orientable and hyperbolic three manifold. Then, for $n \geq 2$ the subspace of invariants of V_n is trivial:*

$$V_n^{\pi_1(M)} = 0.$$

Proof. Let us fix a basis for V_n . Let $e_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $e_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$, so that $\{e_1, e_2\}$ is the standard basis for $V_2 = \mathbf{C}^2$. Thus

$$\{e_1^{n-1}, e_1^{n-2}e_2, \dots, e_2^{n-1}\}$$

is a basis for $V_n = \text{Sym}^{n-1}(V_2)$.

Since M is nonelementary, there exists at least one element $\gamma \in \pi_1(M)$ whose holonomy is nonparabolic (cf. [18, Corollary 3.25]). Up to conjugation, it is

$$\pm \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix},$$

for some $\lambda \in \mathbf{C}$, with $|\lambda| > 1$. This means that the vectors e_1 and e_2 of the standard basis for \mathbf{C}^2 are eigenvectors. Since V_n is the $(n - 1)$ -symmetric power of \mathbf{C}^2 , for n even the only element of V_n γ -invariant is zero. For n odd, the subspace of γ -invariants of V_n is the line generated by $e_1^{(n-1)/2}e_2^{(n-1)/2}$. Any other matrix of $\text{SL}(2, \mathbf{C})$ that fixes $e_1^{(n-1)/2}e_2^{(n-1)/2}$ is either diagonal or antidiagonal (zero entries in the diagonal). Antidiagonal matrices have trace zero, hence they have order four, so they cannot occur because the holonomy of M has no torsion elements. Also, any element $\gamma' \in \pi_1(M)$ that does not commute with γ has nondiagonal holonomy, thus 0 is the only element of V_n invariant by both γ and γ' . \square

Proof of Proposition 3.2. When F has genus $g \geq 2$, then by Lemma 3.4 $\text{Hol}(\pi_1(F)) \setminus \mathbf{H}^3$ is a nonelementary hyperbolic 3-manifold. We apply Lemma 3.5 to conclude that $V_n^{\pi_1(F)} = 0$.

Assume now that F is a torus T^2 . After conjugation, elements of $\pi_1(T^2)$ have holonomy

$$\pm \begin{pmatrix} 1 & \tau \\ 0 & 1 \end{pmatrix} \in \text{SL}(2, \mathbf{C}).$$

The previous matrix maps $e_1^{n-i-1}e_2^i$ to $(\pm 1)^{n-1}e_1^{n-i-1}(e_2 + \tau e_1)^i$, and it follows easily that there is no invariant subspace when n is even and the lift is nonpositive or it is generated by e_1^{n-1} otherwise. \square

Applying Lemma 3.1, Proposition 3.2, Theorem 0.1 and Lemma 3.5, we get the following corollaries.

Corollary 3.6. *Let M be a hyperbolic manifold with k cusps and l ends of infinite volume of genus g_1, \dots, g_l , and let $n \geq 2$. Then*

$$\begin{aligned} \dim_{\mathbf{C}} H^0(\partial\bar{M}; E_{\rho_n}) &= a, \\ \dim_{\mathbf{C}} H^1(\partial\bar{M}; E_{\rho_n}) &= \sum_{i=1}^l 2n(g_i - 1) + 2a, \\ \dim_{\mathbf{C}} H^2(\partial\bar{M}; E_{\rho_n}) &= a, \end{aligned}$$

where a is equal to k if n is odd, and equals to the number of cusps for which the lift of the holonomy is positive if n is even.

Corollary 3.7. *Let M be as in Corollary 3.6. Then $H^0(M; E_{\rho_n}) = 0$,*

$$\dim_{\mathbf{C}} H^1(M; E_{\rho_n}) = \sum_{i=1}^l n(g_i - 1) + a,$$

and $\dim_{\mathbf{C}} H^2(M; E_{\rho_n}) = a$.

3.2. Lifts of the holonomy representation.

Proposition 3.8 ([10]). *The holonomy representation of a hyperbolic 3-manifold M lifts to $\text{SL}(2, \mathbf{C})$. In addition, there is a natural bijection between the set of lifts and the set of spin structures.*

This is proved in Section 2 of [10]. Essentially the idea is that a spin structure on M has a section, because M is parallelizable, and this section lifts to an equivariant section of the spin bundle on the universal covering of M . Identifying the universal covering of M with \mathbf{H}^3 , the spin bundle corresponds to $\text{SL}(2, \mathbf{C})$, and equivariance of

the section gives the lifted representation of $\pi_1(M)$ in $\mathrm{SL}(2, \mathbf{C})$. Notice that on both sets, the set of spin structures and the set of lifts, there is a simply transitive action of $H^1(M; \mathbf{Z}/2\mathbf{Z})$. We view elements in $H^1(M; \mathbf{Z}/2\mathbf{Z})$ as homomorphisms $\pi_1(M) \rightarrow \mathbf{Z}/2\mathbf{Z}$ that describe the difference between signs of two different lifts.

Assume that M has k cusps, and choose $\gamma_1, \dots, \gamma_k \in \pi_1(M)$ k elements so that each γ_i is represented by a simple closed curve in one of the torus of the cusp, and different curves go to different cusps.

Lemma 3.9. *For any choice of curves as above, there exists a lift*

$$\rho: \pi_1(M) \rightarrow \mathrm{SL}(2, \mathbf{C})$$

of the holonomy representation such that $\mathrm{trace}(\rho(\gamma_i)) = -2$, for $i = 1, \dots, k$.

Proof. We denote the peripheral torus by T_1^2, \dots, T_k^2 . Let $\mu_i \in \pi_1(T_i^2)$ be represented by a simple closed curve intersecting γ_i in one point, so that γ_i and μ_i generate $\pi_1(T_i^2)$. We can replace γ_i by $\gamma_i \mu_i^{2n_i}$, for any integer n_i , as multiplying by an even power of μ_i does not change the sign of the trace. We chose the n_i sufficiently large so that Thurston's hyperbolic Dehn filling applies to these slopes. More precisely, we require that there is a continuous path of cone manifold structures with cone angle $\alpha \in [0, 2\pi]$, so that $\alpha = 0$ is the complete structure on M and $\alpha = 2\pi$ is the filled manifold (cf. [26, 17]). Now we chose the lift of the hyperbolic structure on the filled manifold, using Culler's theorem [10], and consider the induced lifts corresponding to changing continuously the cone angle. The map $X(M, \mathrm{SL}(2, \mathbf{C})) \rightarrow X(M, \mathrm{PSL}(2, \mathbf{C}))$ is a local homeomorphism except at characters of reducible representations or representations that preserve a (unoriented) geodesic of \mathbf{H}^3 [14]. Thus we get a continuous path of representations in $X(M, \mathrm{SL}(2, \mathbf{C}))$ parametrized by the cone angle $\alpha \in [0, 2\pi]$, cf. [10, Theorem 4.1].

The holonomy of γ_i is conjugate to

$$\pm \begin{pmatrix} \exp\left(\frac{i\alpha}{2}\right) & 0 \\ 0 & \exp\left(-\frac{i\alpha}{2}\right) \end{pmatrix}$$

and its trace is $\pm 2 \cos(\alpha/2)$. The sign \pm must be constant by continuity. This is clear when $\alpha \neq \pi$ because then the trace is nonzero. When $\alpha = \pi$, we use the local rigidity theorem of [16, 28], that says that this path is locally parametrized by α , and since the derivative of $\pm 2 \cos(\alpha/2)$ at $\alpha = \pi$ is $\pm \sin(\pi/2) = \pm 1$, the trace is monotonic on α when $\alpha = \pi$.

Finally, since we have chosen a lift that is trivial on γ when $\alpha = 2\pi$, the choice of sign is

$$-2 \cos \frac{\alpha}{2},$$

and when $\alpha = 0$ we get the result. □

We obtain the following well known result, proved by Calegari in [8], that applies for instance to the longitude of a knot.

Corollary 3.10. *Let γ be a simple closed curve in a torus of $\partial\overline{M}$ homotopically nontrivial. If γ is homologous to zero in $H_1(M; \mathbf{Z}/2\mathbf{Z})$, then, for every lift $\varphi: \pi_1(M) \rightarrow \text{SL}(2, \mathbf{C})$ of the holonomy representation,*

$$\text{trace}(\varphi(\gamma)) = -2.$$

Proof. The proof follows from the fact that the sign of $\varphi(\gamma)$ cannot be changed by taking different lifts, and by applying Lemma 3.9. □

Corollary 3.11. *Let M be a hyperbolic manifold with a single cusp. Then all lifts of the holonomy representation are nonpositive on $\pi_1(\partial M)$.*

Proof. Since the inclusion in homology

$$H_1(U; \mathbf{Z}/2\mathbf{Z}) \rightarrow H_1(M; \mathbf{Z}/2\mathbf{Z})$$

has rank one, there exists a simple closed curve representing a nontrivial element in $H_1(T^2; \mathbf{Z}/2\mathbf{Z}) \cong H_1(U; \mathbf{Z}/2\mathbf{Z})$ that is $\mathbf{Z}/2\mathbf{Z}$ -homologous to zero in M . Thus Corollary 3.10 applies here, and every lift of the holonomy restricted to the peripheral group is nonpositive. □

Proof of Theorem 0.2. Apply Corollaries 3.7 and 3.11. □

4. Infinitesimal rigidity

Here we prove Theorem 0.3, that we restate.

Theorem 4.1. *Let M be a complete hyperbolic 3-manifold that is topologically finite. If $\partial\overline{M}$ is the union of k tori and l surfaces of genus $g_1, \dots, g_l \geq 2$, and $n \geq 2$, then*

$$\dim_{\mathbf{C}} H^1(M; E_{\text{Ad}\circ\rho_n}) = k(n-1) + \sum (g_i - 1)(n^2 - 1).$$

In particular, if M is closed then $H^1(M; E_{\text{Ad}\circ\rho_n}) = 0$. In addition, all nontrivial elements in $H^1(M; E_{\text{Ad}\circ\rho_n})$ are nontrivial in $H^1(\partial\overline{M}, E_{\text{Ad}\circ\rho_n})$ and have no L^2 representative.

Proof. By Lemma 1.4 we have $\mathfrak{sl}(n, \mathbf{C}) \cong V_{2n-1} \oplus V_{2n-3} \cdots \oplus V_3$. Hence,

$$(9) \quad H^1(M; E_{\text{Ad} \circ \rho_n}) \cong H^1(M; E_{\rho_{2n-1}}) \oplus H^1(M; E_{\rho_{2n-3}}) \oplus \cdots \oplus H^1(M; E_{\rho_3}).$$

The theorem now follows from this isomorphism, Corollary 3.6 and Theorem 0.1. \square

Next we want to prove Theorem 0.4. See [20] for basic results about representation and character varieties. The variety of representations of $\pi_1(M)$ in $\text{SL}(n, \mathbf{C})$ is

$$R(M, \text{SL}(n, \mathbf{C})) = \text{hom}(\pi_1(M), \text{SL}(n, \mathbf{C})).$$

Since $\pi_1(M)$ is finitely generated, this is an algebraic affine set. The group $\text{SL}(n, \mathbf{C})$ acts by conjugation on $R(M, \text{SL}(n, \mathbf{C}))$ algebraically, and the quotient in the algebraic category is the variety of characters:

$$X(M, \text{SL}(n, \mathbf{C})) = R(M, \text{SL}(n, \mathbf{C})) // \text{SL}(n, \mathbf{C}).$$

For a representation $\rho \in R(M, \text{SL}(n, \mathbf{C}))$ its character is the map

$$\begin{aligned} \chi_\rho: \pi_1(M) &\rightarrow \mathbf{C} \\ \gamma &\mapsto \text{trace}(\rho(\gamma)). \end{aligned}$$

The projection $R(M, \text{SL}(n, \mathbf{C})) \rightarrow X(M, \text{SL}(n, \mathbf{C}))$ maps each representation ρ to its character χ_ρ .

Weil's construction gives a natural isomorphism between the Zariski tangent space to a representation $T_\rho^{\text{Zar}} R(M, \text{SL}(n, \mathbf{C}))$ and $Z^1(\pi_1(M), V_{\text{Ad} \circ \rho})$, the space of group cocycles valued in the lie algebra $\mathfrak{sl}(n, \mathbf{C})$, which as $\pi_1(M)$ -module is also written as $V_{\text{Ad} \circ \rho}$. Namely, $Z^1(\pi_1(M), V_{\text{Ad} \circ \rho})$ is the set of maps $d: \pi_1(M) \rightarrow V_{\text{Ad} \circ \rho}$ that satisfy the cocycle relation

$$d(\gamma_1 \gamma_2) = d(\gamma_1) + \text{Ad}_{\rho(\gamma_1)} d(\gamma_2), \quad \forall \gamma_1, \gamma_2 \in \pi_1(M).$$

Notice that $R(M, \text{SL}(n, \mathbf{C}))$ may be a non reduced algebraic set, so the Zariski tangent space may be larger than the Zariski tangent space of the underlying algebraic variety.

The space of coboundaries $B^1(\pi_1(M), V_{\text{Ad} \circ \rho})$ is the set of cocycles that satisfy $d(\gamma) = \text{Ad}_{\rho(\gamma)} m - m$ for all $\gamma \in \pi_1(M)$ and for some fixed $m \in V_{\text{Ad} \circ \rho}$. The space of coboundaries is the tangent space to the orbit by conjugation, so under some hypothesis the cohomology may be identified with the tangent space of the variety of characters (Proposition 4.2). Since M is aspherical, the group cohomology of $\pi_1(M)$

$$H^1(\pi_1(M); V_{\text{Ad} \circ \rho}) = Z^1(\pi_1(M), V_{\text{Ad} \circ \rho}) / B^1(\pi_1(M), V_{\text{Ad} \circ \rho})$$

is naturally isomorphic to $H^1(M; E_{\text{Ad} \circ \rho})$.

DEFINITION. A representation $\rho: \pi_1(M) \rightarrow \mathrm{SL}(n, \mathbf{C})$ is *semisimple* if every subspace of \mathbf{C}^n invariant by $\rho(\pi_1(M))$ has an invariant complement.

Thus a semisimple representation decomposes as direct sum of simple representations, where simple means without proper invariant subspaces.

The following summarizes the relation between tangent spaces and cohomology. See [20] for a proof.

Proposition 4.2. *Let $\rho \in R(M, \mathrm{SL}(n, \mathbf{C}))$.*

1. *There is a natural isomorphism*

$$Z^1(\pi_1(M), V_{\mathrm{Ad} \circ \rho}) \cong T_\rho^{\mathrm{Zar}} R(M, \mathrm{SL}(n, \mathbf{C})).$$

2. *If ρ is semisimple, then it induces an isomorphism*

$$H^1(\pi_1(M); V_{\mathrm{Ad} \circ \rho}) \cong T_\rho^{\mathrm{Zar}} X(M, \mathrm{SL}(n, \mathbf{C})).$$

3. *If ρ is semisimple and a smooth point of $R(M, \mathrm{SL}(n, \mathbf{C}))$, then its character χ_ρ is a smooth point of $X(M, \mathrm{SL}(n, \mathbf{C}))$.*

A point in an algebraic affine set is smooth iff it has the same dimension that its Zariski tangent space. So to prove smoothness we need to compute these dimensions.

Lemma 4.3. *Let ρ_n be as in Theorem 0.4, and T^2 a component of $\partial \overline{M}$ corresponding to a cusp. Then the restriction of ρ_n to $\pi_1(T^2)$ is a smooth point of $R(T^2, \mathrm{SL}(n, \mathbf{C}))$.*

Proof. Knowing that $\dim R(T^2, \mathrm{SL}(n, \mathbf{C})) \leq \dim Z^1(T^2, V_{\mathrm{Ad} \circ \rho_n})$, we want to show that equality of dimensions holds. Before the cocycle space, we first compute the dimension of the cohomology group. By Equation (9) in the proof of Theorem 4.1:

$$\dim H^1(T^2; E_{\mathrm{Ad} \circ \rho_n}) = \sum_{i=2}^n \dim H^1(T^2; E_{\rho_{2i-1}}).$$

Hence, by Corollary 3.6,

$$\dim H^1(T^2; E_{\mathrm{Ad} \circ \rho_n}) = 2(n - 1).$$

We apply the same splitting for computing the dimension of the coboundary space. It is the sum of terms $\dim B^1(T^2, E_{\rho_k})$, for k odd from 3 to $2n - 1$. Since we have an exact sequence

$$0 \rightarrow V_k^{\pi_1(T^2)} \rightarrow V_k \rightarrow B^1(T^2, E_{\rho_k}) \rightarrow 0,$$

$\dim B^1(T^2, E_{\rho_k}) = k - \dim V_k^{\pi_1(T^2)} = k - 1$, by Lemma 3.2. Thus

$$\dim B^1(T^2, E_{\text{Ad}\circ\rho_n}) = (2n - 2) + (2n - 4) + \dots + 2 = n^2 - n.$$

Hence as $H^1(T^2; E_{\text{Ad}\circ\rho_n}) = Z^1(T^2, E_{\text{Ad}\circ\rho_n})/B^1(T^2, E_{\text{Ad}\circ\rho_n})$, we have:

$$\begin{aligned} \dim Z^1(T^2, E_{\text{Ad}\circ\rho_n}) &= \dim H^1(T^2, E_{\text{Ad}\circ\rho_n}) + \dim B^1(T^2, E_{\text{Ad}\circ\rho_n}) \\ &= n^2 + n - 2. \end{aligned}$$

Now we look for a lower bound of $\dim R(T^2, \text{SL}(n, \mathbf{C}))$. Fix $\{\gamma_1, \gamma_2\}$ a generating set of $\pi_1(T^2)$. The representation ρ_n restricted to $\pi_1(T^2)$ has eigenvalues equal to ± 1 . By deforming the representation of $\pi_1(T^2)$ to $\text{SL}(2, \mathbf{C})$, and by composing it with the representation of $\text{SL}(2, \mathbf{C})$ to $\text{SL}(n, \mathbf{C})$, there exists a representation $\rho' \in R(T^2, \text{SL}(n, \mathbf{C}))$ arbitrarily close to ρ_n such that all eigenvalues of $\rho'(\gamma_1)$ are different, in particular $\rho'(\gamma_1)$ diagonalises. Now, to find deformations of ρ' , notice that $\rho'(\gamma_1)$ can be deformed with $n^2 - 1 = \dim(\text{SL}(n, \mathbf{C}))$ parameters, and having all eigenvalues different is an open condition. As $\rho'(\gamma_2)$ has to commute with $\rho'(\gamma_1)$, it has the same eigenspaces, but one can still chose $n - 1$ eigenvalues for $\rho'(\gamma_2)$. This proves that the dimension of some irreducible component of $R(T^2, \text{SL}(n, \mathbf{C}))$ that contains ρ_n is at least

$$n^2 - 1 + n - 1 = n^2 + n - 2.$$

As this is $\dim Z^1(T^2, E_{\text{Ad}\circ\rho_n})$, it is a smooth point. □

Proof of Theorem 0.4. Using Proposition 4.2, we just prove that ρ_n is a smooth point of the variety of representations.

Given a Zariski tangent vector $v \in Z^1(M, V_{\text{Ad}\circ\rho_n})$, we have to show that it is integrable, i.e. that here is a path in the variety of representations whose tangent vector is v . For this, we use the algebraic obstruction theory, see [13, 15]. There exist an infinite sequence of obstructions that are cohomology classes in $H^2(M, V_{\text{Ad}\circ\rho_n})$, each obstruction being defined only if the previous one vanishes. These are related to the analytic expansion in power series of a deformation of a representation, and to Kodaira's theory of infinitesimal deformations. Our aim is to show that this infinite sequence vanishes. This gives a formal power series, that does not need to converge, but this is sufficient for v to be a tangent vector by a theorem of Artin [3] (see [15] for details). We do not give the explicit construction of these obstructions, we just use that they are natural and that they live in the second cohomology group.

By Theorem 0.1 we have an isomorphism:

$$(10) \quad H^2(M; E_{\text{Ad}\circ\rho_n}) \cong H^2(\partial\overline{M}; E_{\text{Ad}\circ\rho_n}).$$

Now, $H^2(\partial\overline{M}; E_{\text{Ad}\circ\rho_n})$ decomposes as the sum of the connected components of $\partial\overline{M}$. If F_g has genus $g \geq 2$ then $H^2(F_g; E_{\text{Ad}\circ\rho_n}) = 0$. Thus, only the components of $\partial\overline{M}$

that are tori appear in $H^2(\partial\overline{M}; E_{\text{Ad}\circ\rho_n})$. By Lemma 4.3 and naturality, the obstructions vanish when restricted to $H^2(T^2; E_{\text{Ad}\circ\rho_n})$, hence they vanish in $H^2(M; E_{\text{Ad}\circ\rho_n})$ by the isomorphism (10). \square

A. Some results on principal bundles

Throughout this section P will denote a G -principal bundle over a manifold M .

REMARK. We will follow the convention that the action of G is on the right.

Assume we have a connection on P with connection form $\omega \in \Omega^1(P; \mathfrak{g})$. This connection defines a horizontal vector bundle H on P . The differential of the bundle projection $\pi_P: P \rightarrow M$ is an isomorphism when restricted to H . Hence, given $X_p \in TM$ and $u \in \pi_P^{-1}(p)$, there exists a unique $\tilde{X}_u \in H_u$ that is projected to X_p . The vector \tilde{X}_u is called the horizontal lift of X_p at u . A vector field on P is called horizontal if it is tangent to H .

All these definitions can be extended in a natural way to the cotangent bundle, exterior powers, tensor powers, etc. Therefore, it makes sense to talk about horizontal forms, horizontal tensors, etc.

Let's recall a common construction. Let F be a differentiable manifold on which G acts on the left. The associated bundle, denoted by $P \times_G F$, is the quotient of $P \times F$ by the diagonal right action of G (i.e. if $(u, x) \in P \times F$, then $(u, x) \cdot g = (ug, g^{-1}x)$). The space $P \times_G F$ has in a natural way a structure of fiber bundle over M with typical fiber F .

OBSERVATION. The definition of $P \times_G F$ allows us to interpret a point u in P as an isomorphism between F and the fiber of $P \times_G F$ at $\pi_P(u)$. Let's say, if π denotes the quotient map $P \times F \rightarrow P \times_G F$, then $\pi(u, \cdot)$ is an isomorphism. Note that $\pi(u \cdot g, x) = \pi(u, gx)$.

We can generalize the notion of associated bundle just "twisting F "; that is, we can take as a starting point an arbitrary bundle over P with typical fiber F , instead of just the product bundle $P \times F$. Let $\pi_Q: Q \rightarrow P$ be a bundle over P with typical fiber F . Assume that we have a fiber-preserving action (on the right) of G on Q that is compatible with the action on P (i. e. $\pi_Q(q \cdot g) = \pi_Q(q) \cdot g$). The quotient Q/G is in a natural way a fiber bundle over M with typical fiber F . In this case, a point $u \in P$ can be interpreted as an isomorphism between the fiber of Q at u , and the fiber of Q/G at $\pi(u)$.

Proposition A.1. *There is a canonical isomorphism between the space of G -equivariant sections of Q , and the space of sections of the associated bundle Q/G .*

Now we want to specialize all these things to the case $Q = \bigwedge^r H^* \otimes V$, where V is a fixed vector space. Let's fix a linear representation $\rho: G \rightarrow \text{Aut}(V)$, in such a way that V becomes a left G -module. We then let G act on Q on the right as follows: if $\alpha_p \otimes w_p$ belongs to Q_p , then $(\alpha_p \otimes w_p) \cdot g = R_{g^{-1}}^* \alpha_p \otimes \rho(g)^{-1} w_p \in Q_{pg}$. Using horizontal lifts we can identify Q/G with $\bigwedge^r T^*M \otimes E$. More precisely, let $p \in M$, $u \in \pi^{-1}(p)$, and $H_u: T_pM \rightarrow H_u$ the horizontal lift map. Then, if we interpret u as an isomorphism between V and E_p , we obtain the isomorphism $\varphi_u: H_u^* \otimes u: Q_u \rightarrow \bigwedge^r T_p^*M \otimes E_p$. Since horizontal lift and u commute with the action of G , we have $\varphi_u(v) = \varphi_{ug}(vg)$, for all $v \in Q$. Therefore, we get an isomorphism φ between Q/G and $\bigwedge^r T^*M \otimes E$.

We will denote by $\Omega_{\text{Hor}}^*(P; V)^G$ the space of horizontal V -valued differential forms over P that are G -equivariant, or, equivalently, the space of G -equivariant sections of the bundle $\bigwedge^r H^* \otimes V$.

OBSERVATION. A form α is horizontal if, and only if, it vanishes on vertical directions, that is, $i_X \alpha = 0$ for any vertical vector field X . Also, α is G -equivariant if, and only if, $R_g^* \alpha = \rho(g^{-1}) \alpha$ for all $g \in G$. Therefore, $\alpha \in \Omega^r(P; V)$ belongs to $\Omega_{\text{Hor}}^r(P; V)^G$ if, and only if,

$$(11) \quad R_g^* \alpha = \rho(g)^{-1} \alpha, \quad \text{for all } g \in G,$$

$$(12) \quad i_Y \alpha = 0, \quad \text{for all } Y \in \mathfrak{g}.$$

Note that we are identifying \mathfrak{g} with the space of G -invariant vertical vectors over a fixed fiber of P .

The connection on P defines an exterior covariant differential on G -equivariant horizontal forms. Namely,

$$D\alpha = (d\alpha) \circ \pi_h, \quad \text{for } \alpha \in \Omega_{\text{Hor}}^r(P; V)^G$$

where π_h is projection on the horizontal distribution defined by the connection. On the other hand, a connection on P induces a connection on the vector bundle $P \times_\rho V$, and hence an exterior covariant differential d_ρ on $\Omega^r(M; E)$. It is easily verified that the canonical isomorphism between the spaces $\Omega_{\text{Hor}}^*(P; V)^G$ and $\Omega^*(M; E)$, "commute" with exterior covariant differentiation (see [19, p.76]).

Proposition A.2. *Let $\omega \in \Omega^1(P; \mathfrak{g})$ be the connection form of the connection defined on P . Then the following formula holds*

$$D\alpha = d\alpha + \rho(\omega) \wedge \alpha.$$

REMARK. If V_1, \dots, V_{p+1} are vector fields on P , by definition,

$$(\rho \wedge \alpha)(V_1, \dots, V_{p+1}) = \sum_{i=1}^{p+1} (-1)^{i+1} \rho(Y_i)(\alpha(V_1, \dots, \hat{V}_i, \dots, V_{p+1})).$$

Taking a base of V , $\rho(\omega)$ is just a matrix of 1-forms, α a column vector of p -forms, and the product $\rho(\omega) \wedge \alpha$ is just the product of a matrix by a vector.

Proof of Proposition A.2. We must prove the form $d\alpha + \rho(\omega) \wedge \alpha$ is horizontal, and that on horizontal vectors coincides with $D\alpha$. The second fact is obvious from the definition of D and the fact that ω vanishes on horizontal vectors. Hence we only need to prove that $d\alpha + \rho(\omega) \wedge \alpha$ vanishes on vertical vectors. Let be X^* the fundamental vector field associated to $X \in \mathfrak{g}$, using Cartan’s identity ($L_{X^*} = di_{X^*} + i_{X^*}d$) we get $i_{X^*}(d\alpha + \rho(\omega) \wedge \alpha) = L_{X^*}\alpha - d(i_{X^*}\alpha) + \rho(X)\alpha$. The infinitesimal version of the G -equivariance of α states that $L_{X^*}\alpha = -\rho(X)\alpha$. Then we conclude that $d\alpha + \rho(\omega) \wedge \alpha$ is vertical. \square

Now assume that M is a Riemannian manifold, and that we have a metric on the vector bundle $E = P \times_G V$. These metrics induce an inner product on the space of E -valued forms over M .

$$(\alpha, \beta) = \int_M \langle \alpha(x), \beta(x) \rangle_x \omega_M.$$

On the other hand, the Riemannian metric on M defines a metric tensor on the horizontal bundle H , in such a way that horizontal lifts are isometries. Also, the metric defined on E defines a metric on the trivial vector bundle $P \times V$. A right invariant volume form ω_G on G defines a right invariant volume form along the fibers of P . Therefore, we can define an inner product on $\Omega_{\text{Hor}}^r(P; V)^G$ by

$$(\tilde{\alpha}, \tilde{\beta}) = \int_P \langle \tilde{\alpha}(u), \tilde{\beta}(u) \rangle_u \pi_P^*(\omega_M) \wedge \omega_G^*.$$

We want to study how the metrics defined on $\Omega^r(M; E)$ and $\Omega_{\text{Hor}}^r(P; V)^G$ are related by the canonical isomorphism. However, this comparison doesn’t make sense if G is not assumed to be compact (if $\alpha \in \Omega^r(M; E)$ has compact support, then the corresponding form $\tilde{\alpha}$ in $\Omega_{\text{Hor}}^r(P; V)^G$ has compact support if, and only if, G is compact). From now on we will assume that G is compact. In order to avoid confusions we will denote G by K in this case. In this case we can simplify things a little bit. First, take a K -invariant metric on V , and use it to define a “constant” metric on $P \times V$. Since this metric is K -invariant, we get a metric on the vector bundle E . Under these hypothesis, we get a nice relation between these two metrics. In order to get this relation, we need the following lemma.

Proposition A.3. *Let ω_M be a volume form on M , and ω_K a right invariant volume form on K . Denote by ω_K^* the right invariant volume form on the fibers of P defined by ω_K . If f is a function defined on P , then the function $\tilde{f}(u) = \int_K f(ug)\omega_K$ is invariant along the fibers, and hence can be seen as a function on M . With these hypothesis, we have*

$$\int_P f(u)\pi_P^*(\omega_M) \wedge \omega_K^* = \int_M \tilde{f}(x)\omega_M.$$

Proof. Take an open set $U \subset M$ that trivializes P , and a trivializing map $\psi: U \times K \rightarrow \pi_P^{-1}(U)$. Let's denote by π_U and π_K the projection of $U \times K$ on the first and on the second factor respectively. We have $\omega_K^* = (\psi^*)^{-1}(\pi_K^*(\omega_K))$. The change of variable formula gives

$$\int_{\pi_P^{-1}(U)} f(u)\pi_P^*(\omega_M) \wedge \omega_K^* = \int_{U \times K} f(\psi(x, g))\pi_U^*(\omega_M) \wedge \pi_K^*(\omega_K).$$

By Fubini's theorem, the last integral is,

$$\int_U \left(\int_K f(\psi(x, g))\omega_K \right) \omega_M = \int_U \tilde{f}(x)\omega_M.$$

The result follows by taking a partition of unity subordinated to a trivializing open cover. □

The function $\langle \tilde{\alpha}(u), \tilde{\beta}(u) \rangle_V$ is constant along the fibers, and equals to $\langle \alpha(x), \beta(x) \rangle_x$, where $x = \pi_P(u)$. The above lemma then implies the following proposition.

Proposition A.4. *With the above notation,*

$$\langle \tilde{\alpha}, \tilde{\beta} \rangle = \mu(K)\langle \alpha, \beta \rangle,$$

where μ denotes the measure defined by the volume form ω_K .

Consider the pairing

$$\begin{aligned} \Omega_{\text{Hor}}^r(P; V)^K \times \Omega_{\text{Hor}}^{m-r}(P; V)^K &\rightarrow \mathbf{R} \\ (\alpha, \beta) &\mapsto \int_P (\alpha \wedge \beta) \wedge \omega_K, \end{aligned}$$

where the wedge product of a V -valued is defined using the usual wedge product on scalar-valued forms, and the inner product on V . On the other hand, the metric on the horizontal bundle, and the orientation we have on it, allow us to define a Hodge star operator on the space of horizontal forms,

$$* : \Omega_{\text{Hor}}^r(P; V)^K \rightarrow \Omega_{\text{Hor}}^{m-r}(P; V)^K.$$

Note that we have $(\alpha, \beta) = \phi(\alpha, * \beta)$.

Proposition A.5. *Let $T: \Omega_{\text{Hor}}^r(P; V)^K \rightarrow \Omega_{\text{Hor}}^{r+k}(P; V)^K$ be a linear operator that decreases supports. Assume we have a linear operator*

$$S: \Omega_{\text{Hor}}^{m-(r+k)}(P; V)^K \rightarrow \Omega_{\text{Hor}}^{m-r}(P; V)^K$$

such that $\phi(T\alpha, \beta) = \phi(\alpha, S\beta)$. Then, the formal adjoint of T is

$$T^* = (-1)^{r(m-r)} * S*: \Omega_{\text{Hor}}^{r+k}(P; V)^K \rightarrow \Omega_{\text{Hor}}^r(P; V)^K.$$

Proof. Let's denote $\Omega_{\text{Hor}}^r(P; V)^K$ by M_r . We have the following commutative diagram,

$$\begin{array}{ccc} M_{r+k}^* & \xrightarrow{T'} & M_r^* \\ \uparrow & & \uparrow \\ M_{r+k} & \xrightarrow{T^*} & M_r \end{array}$$

where the vertical arrows are the isomorphisms given by the metrics, T' is the dual map of T , and T^* its adjoint. We can factor the metric isomorphism as $\phi(*, \cdot)$. We have the following commutative diagram

$$\begin{array}{ccc} M_{r+k}^* & \xrightarrow{T'} & M_r^* \\ \phi(\cdot, \cdot) \uparrow & & \uparrow \phi(\cdot, \cdot) \\ M_{m-(r+k)} & \xrightarrow{S} & M_{m-r} \\ * \uparrow & & \uparrow * \\ M_{r+k} & \xrightarrow{T^*} & M_r \end{array}$$

The proposition now follows from the fact that on degree r we have $*^{-1} = (-1)^{r(m-r)}*$. □

References

- [1] I. Agol: *Tameness of hyperbolic 3-manifolds*, preprint, arXiv:math/0405568.
- [2] A. Andreotti and E. Vesentini: *Carleman estimates for the Laplace–Beltrami equation on complex manifolds*, Inst. Hautes Études Sci. Publ. Math. **25** (1965), 81–130.
- [3] M. Artin: *On the solutions of analytic equations*, Invent. Math. **5** (1968), 277–291.

- [4] M. Boileau and J. Porti: Geometrization of 3-Orbifolds of Cyclic Type, *Astérisque* **272**, 2001.
- [5] F. Bonahon: *Cobordism of automorphisms of surfaces*, *Ann. Sci. École Norm. Sup. (4)* **16** (1983), 237–270.
- [6] A. Borel and N.R. Wallach: *Continuous Cohomology, Discrete Subgroups, and Representations of Reductive Groups*, *Annals of Mathematics Studies* **94**, Princeton Univ. Press, Princeton, NJ, 1980.
- [7] K. Bromberg: *Rigidity of geometrically finite hyperbolic cone-manifolds*, *Geom. Dedicata* **105** (2004), 143–170.
- [8] D. Calegari: *Real places and torus bundles*, *Geom. Dedicata* **118** (2006), 209–227.
- [9] D. Calegari and D. Gabai: *Shrinkwrapping and the taming of hyperbolic 3-manifolds*, *J. Amer. Math. Soc.* **19** (2006), 385–446.
- [10] M. Culler: *Lifting representations to covering groups*, *Adv. in Math.* **59** (1986), 64–70.
- [11] W. Fulton and J. Harris: *Representation Theory*, *Graduate Texts in Mathematics* **129**, Springer, New York, 1991.
- [12] H. Garland: *A rigidity theorem for discrete subgroups*, *Trans. Amer. Math. Soc.* **129** (1967), 1–25.
- [13] W.M. Goldman: *Invariant functions on Lie groups and Hamiltonian flows of surface group representations*, *Invent. Math.* **85** (1986), 263–302.
- [14] M. Heusener and J. Porti: *The variety of characters in $PSL_2(\mathbb{C})$* , *Bol. Soc. Mat. Mexicana (3)* **10** (2004), Special Issue, 221–237.
- [15] M. Heusener and J. Porti: *Deformations of reducible representations of 3-manifold groups into $PSL_2(\mathbb{C})$* , *Algebr. Geom. Topol.* **5** (2005), 965–997.
- [16] C.D. Hodgson and S.P. Kerckhoff: *Rigidity of hyperbolic cone-manifolds and hyperbolic Dehn surgery*, *J. Differential Geom.* **48** (1998), 1–59.
- [17] C.D. Hodgson and S.P. Kerckhoff: *Universal bounds for hyperbolic Dehn surgery*, *Ann. of Math. (2)* **162** (2005), 367–421.
- [18] M. Kapovich: *Hyperbolic Manifolds and Discrete Groups*, *Progress in Mathematics* **183**, Birkhäuser Boston, Boston, MA, 2001.
- [19] S. Kobayashi and K. Nomizu: *Foundations of Differential Geometry, I*, reprint of the 1963 original, *Wiley Classics Library*, Wiley, New York, 1996.
- [20] A. Lubotzky and A.R. Magid: *Varieties of Representations of Finitely Generated Groups*, *Mem. Amer. Math. Soc.* **58**, 1985.
- [21] Y. Matsushima and S. Murakami: *On vector bundle valued harmonic forms and automorphic forms on symmetric riemannian manifolds*, *Ann. of Math. (2)* **78** (1963), 365–416.
- [22] D. McCullough and A. Miller: *Homeomorphisms of 3-Manifolds with Compressible Boundary*, *Mem. Amer. Math. Soc.* **61**, 1986.
- [23] K. Nomizu: *Lie Groups and Differential Geometry*, *The Mathematical Society of Japan*, 1956.
- [24] M.S. Raghunathan: *On the first cohomology of discrete subgroups of semisimple Lie groups*, *Amer. J. Math.* **87** (1965), 103–139.
- [25] M.S. Raghunathan: *Discrete Subgroups of Lie Groups*, Springer, New York, 1972.
- [26] W.P. Thurston. *The Geometry and Topology of Three-Manifolds*, Princeton University, <http://www.msri.org/publications/books/gt3m/>.
- [27] A. Weil: *Remarks on the cohomology of groups*, *Ann. of Math. (2)* **80** (1964), 149–157.
- [28] H. Weiss: *Local rigidity of 3-dimensional cone-manifolds*, *J. Differential Geom.* **71** (2005), 437–506.

Pere Menal-Ferrer
Departament de Matemàtiques
Universitat Autònoma de Barcelona
08193 Cerdanyola del Vallès
Spain
e-mail: pmenal@mat.uab.cat

Joan Porti
Departament de Matemàtiques
Universitat Autònoma de Barcelona
08193 Cerdanyola del Vallès
Spain
e-mail: porti@mat.uab.cat