

THE ROTATION NUMBER OF PRIMITIVE VECTOR SEQUENCES

YUSUKE SUYAMA

(Received October 10, 2013, revised April 4, 2014)

Abstract

We give a formula on the rotation number of a sequence of primitive vectors, which is a generalization of the formula on the rotation number of a unimodular sequence in [2].

1. Introduction

Let $v_1, \dots, v_d \in \mathbb{Z}^2$ be a sequence of primitive vectors such that $\varepsilon_i = \det(v_i, v_{i+1}) \neq 0$ for all $i = 1, \dots, d$, and let $a_i = \varepsilon_{i-1}^{-1} \varepsilon_i^{-1} \det(v_{i+1}, v_{i-1})$, where $v_0 = v_d$ and $v_{d+1} = v_1$. The *rotation number* $\text{Rot}(v_1, \dots, v_d)$ of the sequence v_1, \dots, v_d around the origin is defined by

$$\frac{1}{2\pi} \sum_{i=1}^d \int_{L_i} \frac{-y dx + x dy}{x^2 + y^2},$$

where L_i is the line segment from v_i to v_{i+1} . The sequence is called *unimodular* if $|\varepsilon_i| = 1$ for all $i = 1, \dots, d$. Recently A. Higashitani and M. Masuda [2] proved the following:

Theorem A ([2]). *If v_1, \dots, v_d is a unimodular sequence, then*

$$\text{Rot}(v_1, \dots, v_d) = \frac{1}{12} \sum_{i=1}^d (3\varepsilon_i + a_i).$$

When $\varepsilon_i = 1$ for all i and the rotation number is one, Theorem A is well known and formulated as $3d + \sum_{i=1}^d a_i = 12$. It can be proved in an elementary way, but interestingly it can also be proved using toric geometry, to be more precise, by applying Nöther's formula to complete non-singular toric varieties of complex dimension two, see [1]. When $\varepsilon_i = 1$ for all i but the rotation number is not necessarily one, Theorem A was proved in [4] using toric topology. The proof is a generalization of the proof above using toric geometry. The original proof of Theorem A by Higashitani and

Masuda was a slight modification of the proof in [4] but then they found an elementary proof. Another elementary proof of Theorem A is given by R.T. Živaljević [3].

Theorem A does not hold when the unimodularity condition is dropped. In this paper, we give a formula on the rotation number of a (not necessarily unimodular) sequence of primitive vectors v_1, \dots, v_d with $\varepsilon_i \neq 0$ for all i , see Theorem 6. In this case, a_i 's are rational numbers but are not always integers. The proof is done by adding primitive vectors in an appropriate way to the given sequence so that the enlarged sequence is unimodular and then by applying Theorem A to the enlarged unimodular sequence. This combinatorial process, that is, making the given sequence to a unimodular sequence by adding primitive vectors corresponds to resolution of singularity by blow-up in geometry, see [1].

The structure of the paper is as follows: In Section 2, we discuss Hirzebruch–Jung continued fractions used in our proof of the main theorem. In Section 3, we state the main theorem and give an example. In Section 4, we give a proof of the main theorem.

2. Continued fractions

Let $m \geq 2$ and $x (< m)$ be a positive integer prime to m , and let

$$\frac{m}{x} = n_1 - \frac{1}{n_2 - \frac{1}{\ddots - \frac{1}{n_{L(m,x)}}}}, \quad n_j \geq 2$$

be the continued fraction expansion. This continued fraction expansion is unique, and is called a *Hirzebruch–Jung continued fraction*. We define $L(1, 0) = 0$.

Lemma 1. *Let $m \geq 2$ and $x (< m)$ be a positive integer prime to m , and let*

$$\frac{m}{x} = n_1 - \frac{1}{n_2 - \frac{1}{\ddots - \frac{1}{n_{L(m,x)}}}}, \quad n_j \geq 2$$

be the continued fraction expansion. Let $y (< m)$ be a unique positive integer such that $xy \equiv 1 \pmod{m}$. Then the following identity holds:

$$\begin{pmatrix} 0 & -1 \\ 1 & n_1 \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & n_2 \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{L(m,x)} \end{pmatrix} = \begin{pmatrix} \frac{1-xy}{m} & -x \\ y & m \end{pmatrix}.$$

Proof. We prove this by induction on $L(m, x)$.

If $L(m, x) = 1$, then we must have $x = 1$, $n_1 = m$ and $y = 1$. So the lemma holds when $L(m, x) = 1$.

Suppose that $L(m, x) \geq 2$ and the lemma holds for $L(m, x) - 1$. We have

$$\frac{x}{n_1x - m} = n_2 - \frac{1}{n_3 - \frac{1}{\dots - \frac{1}{n_{L(m,x)}}}}$$

Since $L(m, x) \geq 2$, we have $x \geq 2$. Since the right hand side in the identity above is greater than 1, we have $0 < n_1x - m < x$. Since m and x are relatively prime, x and $n_1x - m$ are relatively prime. Moreover $(xy - 1)/m$ is a positive integer less than x and $(n_1x - m)(xy - 1)/m \equiv 1 \pmod{x}$. Hence by the hypothesis of induction, we obtain

$$\begin{aligned} \begin{pmatrix} 0 & -1 \\ 1 & n_2 \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{L(m,x)} \end{pmatrix} &= \begin{pmatrix} \frac{1 - (n_1x - m)(xy - 1)/m}{x} & -(n_1x - m) \\ \frac{xy - 1}{m} & x \end{pmatrix} \\ &= \begin{pmatrix} y - n_1 \frac{xy - 1}{m} & -(n_1x - m) \\ \frac{xy - 1}{m} & x \end{pmatrix}. \end{aligned}$$

Therefore we have

$$\begin{aligned} \begin{pmatrix} 0 & -1 \\ 1 & n_1 \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{L(m,x)} \end{pmatrix} \\ = \begin{pmatrix} 0 & -1 \\ 1 & n_1 \end{pmatrix} \begin{pmatrix} y - n_1 \frac{xy - 1}{m} & -(n_1x - m) \\ \frac{xy - 1}{m} & x \end{pmatrix} &= \begin{pmatrix} \frac{1 - xy}{m} & -x \\ y & m \end{pmatrix}, \end{aligned}$$

proving the lemma for $L(m, x)$. □

Proposition 2. *The following identity holds:*

$$\frac{m}{y} = n_{L(m,x)} - \frac{1}{n_{L(m,x)-1} - \frac{1}{\dots - \frac{1}{n_1}}}$$

Proof. Let $f: M_2(\mathbb{Z}) \rightarrow M_2(\mathbb{Z})$ be the antihomomorphism defined by

$$f\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \begin{pmatrix} a & -c \\ -b & d \end{pmatrix}.$$

By Lemma 1, we have

$$\begin{aligned} \begin{pmatrix} 0 & -1 \\ 1 & n_{L(m,x)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_1 \end{pmatrix} &= f\left(\begin{pmatrix} 0 & -1 \\ 1 & n_1 \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{L(m,x)} \end{pmatrix}\right) \\ &= f\left(\begin{pmatrix} \frac{1-xy}{m} & -x \\ y & m \end{pmatrix}\right) = \begin{pmatrix} \frac{1-xy}{x} & -y \\ m & m \end{pmatrix}, \end{aligned}$$

proving the proposition. □

REMARK 3. A similar assertion holds for regular continued fractions. Let

$$\frac{m}{x} = n_1 + \frac{1}{n_2 + \frac{1}{\ddots + \frac{1}{n_l}}}, \quad n_j \geq 1$$

be a continued fraction expansion, and let $y (< m)$ be a unique positive integer such that $xy \equiv (-1)^{l+1} \pmod{m}$. Then the following identity holds:

$$\begin{pmatrix} 0 & 1 \\ 1 & n_1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & n_2 \end{pmatrix} \cdots \begin{pmatrix} 0 & 1 \\ 1 & n_l \end{pmatrix} = \begin{pmatrix} \frac{xy + (-1)^l}{m} & x \\ y & m \end{pmatrix}.$$

The proof is similar to Lemma 1. The following identity can be deduced by taking transpose at the identity above:

$$\frac{m}{y} = n_l + \frac{1}{n_{l-1} + \frac{1}{\ddots + \frac{1}{n_1}}}.$$

3. The main theorem

Let $v_1, \dots, v_d \in \mathbb{Z}^2$ be a sequence of vectors such that $\varepsilon_i = \det(v_i, v_{i+1}) \neq 0$ for all $i = 1, \dots, d$. We define $v_0 = v_d$ and $v_{d+1} = v_1$. We assume that each vector is *primitive*, i.e. its components are relatively prime.

Lemma 4. *For each $i = 1, \dots, d$, there exists a unique non-negative integer $x_i < |\varepsilon_i|$ such that x_i and $|\varepsilon_i|$ are relatively prime and*

$$P_i = (v_i, v_{i+1}) \begin{pmatrix} 1 & -x_i \\ 0 & |\varepsilon_i| \end{pmatrix}^{-1}$$

is a unimodular matrix.

Proof. Let $v_i = \begin{pmatrix} a \\ b \end{pmatrix}$ and $v_{i+1} = \begin{pmatrix} c \\ d \end{pmatrix}$. We assume that $\varepsilon_i > 0$. Since v_i is primitive, there exist $p, q \in \mathbb{Z}$ such that $ap + bq = 1$. Then we have

$$\begin{pmatrix} p & q \\ -b & a \end{pmatrix}(v_i, v_{i+1}) = \begin{pmatrix} 1 & cp + dq \\ 0 & |\varepsilon_i| \end{pmatrix}.$$

There exists a unique $n \in \mathbb{Z}$ satisfying $-|\varepsilon_i| < cp + dq + n|\varepsilon_i| \leq 0$. So we put $x_i = -(cp + dq + n|\varepsilon_i|)$. Then we have

$$\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \begin{pmatrix} p & q \\ -b & a \end{pmatrix} (v_i, v_{i+1}) = \begin{pmatrix} 1 & -x_i \\ 0 & |\varepsilon_i| \end{pmatrix}.$$

Hence

$$P_i = \begin{pmatrix} a & -q \\ b & p \end{pmatrix} \begin{pmatrix} 1 & -n \\ 0 & 1 \end{pmatrix}$$

is a unimodular matrix. When $\varepsilon_i < 0$, we can show that the assertion holds by a similar argument. Since $\begin{pmatrix} -x_i \\ |\varepsilon_i| \end{pmatrix} = P_i^{-1}v_{i+1}$ is primitive, x_i and $|\varepsilon_i|$ are relatively prime. \square

Note that $\det(P_i) = \varepsilon_i/|\varepsilon_i|$. Similarly, there exists a unique non-negative integer $y_i < |\varepsilon_i|$ such that

$$(3.1) \quad Q_i = (v_{i+1}, v_i) \begin{pmatrix} 1 & -y_i \\ 0 & |\varepsilon_i| \end{pmatrix}^{-1}$$

is a unimodular matrix.

Lemma 5. Let $a_i = \varepsilon_{i-1}^{-1} \varepsilon_i^{-1} \det(v_{i+1}, v_{i-1})$. Then a_i satisfies

$$(3.2) \quad \varepsilon_{i-1}^{-1} v_{i-1} + \varepsilon_i^{-1} v_{i+1} + a_i v_i = 0.$$

Proof. It is easy to check that

$$\det(v_i, v_{i+1})v_{i-1} + \det(v_{i-1}, v_i)v_{i+1} + \det(v_{i+1}, v_{i-1})v_i = 0.$$

Dividing both sides by $\varepsilon_{i-1}\varepsilon_i = \det(v_{i-1}, v_i) \det(v_i, v_{i+1})$, we obtain (3.2). \square

Let $m \geq 2$ and x ($< m$) be a positive integer prime to m , and let

$$\frac{m}{x} = n_1 - \frac{1}{\frac{1}{n_2 - \frac{1}{\ddots - \frac{1}{n_{L(m,x)}}}}}, \quad n_j \geq 2$$

be the Hirzebruch–Jung continued fraction expansion. We set $N(m, x) = n_1 + \dots + n_{L(m,x)}$ and $N(1, 0) = 0$. The following is our main theorem:

Theorem 6. *Let v_1, \dots, v_d be a sequence of primitive vectors and $\varepsilon_i = \det(v_i, v_{i+1})$, $a_i = \varepsilon_{i-1}^{-1} \varepsilon_i^{-1} \det(v_{i+1}, v_{i-1})$. Let x_i and y_i be the integers defined in Lemma 4 and (3.1). Then $\text{Rot}(v_1, \dots, v_d)$ is given by*

$$(3.3) \quad \frac{1}{12} \sum_{i=1}^d \left((3(L(|\varepsilon_i|, x_i) + 1) - N(|\varepsilon_i|, x_i)) \frac{\varepsilon_i}{|\varepsilon_i|} + a_i - \frac{x_i + y_i}{\varepsilon_i} \right).$$

In particular, the value of (3.3) is an integer.

For i such that $|\varepsilon_i| \geq 2$, let

$$\frac{|\varepsilon_i|}{x_i} = n_1^{(i)} - \frac{1}{n_2^{(i)} - \frac{1}{\dots - \frac{1}{n_{L(|\varepsilon_i|, x_i)}^{(i)}}}}, \quad n_j^{(i)} \geq 2,$$

be the Hirzebruch–Jung continued fraction expansion.

EXAMPLE 7. Let $d = 5$ and

$$v_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 1 \\ 3 \end{pmatrix}, \quad v_3 = \begin{pmatrix} -2 \\ -1 \end{pmatrix}, \quad v_4 = \begin{pmatrix} -2 \\ 1 \end{pmatrix}, \quad v_5 = \begin{pmatrix} 5 \\ -3 \end{pmatrix}.$$

Then we have the following:

i	ε_i	a_i	x_i	y_i	$L(\varepsilon_i , x_i)$	$n_1^{(i)}$	$n_2^{(i)}$	$N(\varepsilon_i , x_i)$
1	3	-2	2	2	2	2	2	4
2	5	1/15	2	3	2	2	3	5
3	-4	7/20	1	1	1	4		4
4	1	11/4	0	0	0			0
5	3	1/3	1	1	1	3		3




Fig. 1. A sequence of primitive vectors.

So we have

$$\sum_{i=1}^d (3(L(|\varepsilon_i|, x_i) + 1) - N(|\varepsilon_i|, x_i)) \frac{\varepsilon_i}{|\varepsilon_i|}$$

$$= (3(2 + 1) - 4) + (3(2 + 1) - 5) - (3(1 + 1) - 4) + 3 + (3(1 + 1) - 3) = 13,$$

$$\sum_{i=1}^d a_i = -2 + \frac{1}{15} + \frac{7}{20} + \frac{11}{4} + \frac{1}{3} = \frac{3}{2},$$

$$\sum_{i=1}^d \frac{x_i + y_i}{\varepsilon_i} = \frac{2 + 2}{3} + \frac{2 + 3}{5} + \frac{1 + 1}{-4} + \frac{0 + 0}{1} + \frac{1 + 1}{3} = \frac{5}{2}.$$

Therefore the value (3.3) is $(1/12)(13 + 3/2 - 5/2) = 1$, while the rotation number of the sequence v_1, \dots, v_5 in Fig. 1 is clearly one.

4. Proof of Theorem 6

In this section, we give a proof of Theorem 6. We will use the notation in Section 3 freely. We need the following lemma.

Lemma 8. For each $i = 1, \dots, d$, the following identity holds:

$$\begin{pmatrix} 0 & -1 \\ 1 & n_1^{(i)} \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & n_2^{(i)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{L(|\varepsilon_i|, x_i)}^{(i)} \end{pmatrix} = \begin{pmatrix} \frac{1 - x_i y_i}{|\varepsilon_i|} & -x_i \\ y_i & |\varepsilon_i| \end{pmatrix}.$$

Proof. If $|\varepsilon_i| = 1$, then $x_i = y_i = L(|\varepsilon_i|, x_i) = 0$ and the left hand side above is understood to be the identity matrix. Assume $|\varepsilon_i| \geq 2$. By Lemma 4, x_i and $|\varepsilon_i|$ are relatively prime. Since

$$\begin{aligned} Q_i^{-1} P_i &= \begin{pmatrix} 1 & -y_i \\ 0 & |\varepsilon_i| \end{pmatrix} (v_{i+1}, v_i)^{-1} (v_i, v_{i+1}) \begin{pmatrix} 1 & -x_i \\ 0 & |\varepsilon_i| \end{pmatrix}^{-1} \\ &= \frac{1}{|\varepsilon_i|} \begin{pmatrix} 1 & -y_i \\ 0 & |\varepsilon_i| \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} |\varepsilon_i| & x_i \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} -y_i & \frac{1 - x_i y_i}{|\varepsilon_i|} \\ |\varepsilon_i| & x_i \end{pmatrix} \end{aligned}$$

is a unimodular matrix, $x_i y_i$ is congruent to 1 modulo $|\varepsilon_i|$. Therefore the lemma follows from Lemma 1. □

Proof of Theorem 6. For $j = 0, \dots, L(|\varepsilon_i|, x_i) + 1$, we define

$$(4.1) \quad w_j^{(i)} = \begin{cases} P_i \begin{pmatrix} 0 & -1 \\ 1 & n_1^{(i)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_j^{(i)} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} & (0 \leq j \leq L(|\varepsilon_i|, x_i)), \\ P_i \begin{pmatrix} 0 & -1 \\ 1 & n_1^{(i)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{j-1}^{(i)} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} & (1 \leq j \leq L(|\varepsilon_i|, x_i) + 1). \end{cases}$$

Note that both expressions at the right hand side of (4.1) are equal if $1 \leq j \leq L(|\varepsilon_i|, x_i)$. By the definition of $w_j^{(i)}$, it follows that

$$(4.2) \quad \det(w_j^{(i)}, w_{j+1}^{(i)}) = \det(P_i) \det \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right) = \frac{\varepsilon_i}{|\varepsilon_i|} \in \{\pm 1\}$$

for any $j = 0, \dots, L(|\varepsilon_i|, x_i)$. So the sequence

$$(4.3) \quad \dots, v_i = w_0^{(i)}, w_1^{(i)}, \dots, w_{L(|\varepsilon_i|, x_i)+1}^{(i)} = v_{i+1}, \dots$$

is unimodular.




Fig. 2. Adding $w_j^{(i)}$ to the given vector sequence.

Hence by Theorem A, the rotation number of v_1, \dots, v_d is given by

$$\begin{aligned}
 (4.4) \quad & \frac{1}{4} \sum_{i=1}^d \sum_{j=0}^{L(|\varepsilon_i|, x_i)} \det(w_j^{(i)}, w_{j+1}^{(i)}) + \frac{1}{12} \sum_{i=1}^d \frac{\det(w_1^{(i)}, w_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)})}{\det(w_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)}, v_i) \det(v_i, w_1^{(i)})} \\
 & + \frac{1}{12} \sum_{i=1}^d \sum_{j=1}^{L(|\varepsilon_i|, x_i)} \frac{\det(w_{j+1}^{(i)}, w_{j-1}^{(i)})}{\det(w_{j-1}^{(i)}, w_j^{(i)}) \det(w_j^{(i)}, w_{j+1}^{(i)})}.
 \end{aligned}$$

As for the first summand in (4.4), it follows from (4.2) that

$$\sum_{j=0}^{L(|\varepsilon_i|, x_i)} \det(w_j^{(i)}, w_{j+1}^{(i)}) = (L(|\varepsilon_i|, x_i) + 1) \frac{\varepsilon_i}{|\varepsilon_i|}.$$

As for the second summand in (4.4), we first observe that it follows from Lemma 5 that

$$\begin{aligned}
 P_{i-1}^{-1} P_i &= \begin{pmatrix} 1 & -x_{i-1} \\ 0 & |\varepsilon_{i-1}| \end{pmatrix} (v_{i-1}, v_i)^{-1} (v_i, v_{i+1}) \begin{pmatrix} 1 & -x_i \\ 0 & |\varepsilon_i| \end{pmatrix}^{-1} \\
 &= \frac{1}{|\varepsilon_i|} \begin{pmatrix} 1 & -x_{i-1} \\ 0 & |\varepsilon_{i-1}| \end{pmatrix} (v_{i-1}, v_i)^{-1} (v_i, -\varepsilon_i (\varepsilon_{i-1}^{-1} v_{i-1} + a_i v_i)) \begin{pmatrix} |\varepsilon_i| & x_i \\ 0 & 1 \end{pmatrix} \\
 &= \frac{1}{|\varepsilon_i|} \begin{pmatrix} 1 & -x_{i-1} \\ 0 & |\varepsilon_{i-1}| \end{pmatrix} \begin{pmatrix} 0 & -\varepsilon_i \varepsilon_{i-1}^{-1} \\ 1 & -a_i \varepsilon_i \end{pmatrix} \begin{pmatrix} |\varepsilon_i| & x_i \\ 0 & 1 \end{pmatrix} \\
 &= \frac{1}{|\varepsilon_i|} \begin{pmatrix} -|\varepsilon_i| x_{i-1} & -\varepsilon_i \varepsilon_{i-1}^{-1} - x_{i-1} x_i + a_i \varepsilon_i x_{i-1} \\ |\varepsilon_{i-1}| |\varepsilon_i| & |\varepsilon_{i-1}| (x_i - a_i \varepsilon_i) \end{pmatrix}.
 \end{aligned}$$

So it follows from (4.3), (4.1), (4.2), and Lemma 8 that

$$\begin{aligned}
 & \frac{\det(w_1^{(i)}, w_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)})}{\det(w_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)}, v_i) \det(v_i, w_1^{(i)})} \\
 &= \frac{\det(w_1^{(i)}, w_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)})}{\det(w_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)}, w_{L(|\varepsilon_{i-1}|, x_{i-1})+1}^{(i-1)}) \det(w_0^{(i)}, w_1^{(i)})} \\
 &= \frac{|\varepsilon_{i-1}| |\varepsilon_i|}{\varepsilon_{i-1} \varepsilon_i} \det \left(P_i \begin{pmatrix} 0 \\ 1 \end{pmatrix}, P_{i-1} \begin{pmatrix} 0 & -1 \\ 1 & n_1^{(i-1)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{L(|\varepsilon_{i-1}|, x_{i-1})}^{(i-1)} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) \\
 &= \frac{|\varepsilon_{i-1}| |\varepsilon_i|}{\varepsilon_{i-1} \varepsilon_i} \det(P_{i-1}) \det \left(P_{i-1}^{-1} P_i \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 - x_{i-1} y_{i-1} \\ |\varepsilon_{i-1}| \\ y_{i-1} \end{pmatrix} \right) \\
 &= \frac{1}{\varepsilon_i} \det \begin{pmatrix} -\frac{\varepsilon_i}{\varepsilon_{i-1}} - x_{i-1} x_i + a_i \varepsilon_i x_{i-1} & 1 - x_{i-1} y_{i-1} \\ |\varepsilon_{i-1}| (x_i - a_i \varepsilon_i) & y_{i-1} \end{pmatrix} \\
 &= a_i - \frac{x_i}{\varepsilon_i} - \frac{y_{i-1}}{\varepsilon_{i-1}}.
 \end{aligned}$$

As for the last summand in (4.4), it follows from (4.1) and (4.2) that

$$\begin{aligned}
 & \frac{\det(w_{j+1}^{(i)}, w_{j-1}^{(i)})}{\det(w_{j-1}^{(i)}, w_j^{(i)}) \det(w_j^{(i)}, w_{j+1}^{(i)})} \\
 &= \det(P_i) \det \left(\begin{pmatrix} 0 & -1 \\ 1 & n_1^{(i)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_j^{(i)} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 & -1 \\ 1 & n_1^{(i)} \end{pmatrix} \cdots \begin{pmatrix} 0 & -1 \\ 1 & n_{j-1}^{(i)} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) \\
 &= \det(P_i) \det \left(\begin{pmatrix} 0 & -1 \\ 1 & n_j^{(i)} \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix} \right) \\
 &= \frac{\varepsilon_i}{|\varepsilon_i|} \det \begin{pmatrix} -1 & 1 \\ n_j^{(i)} & 0 \end{pmatrix} = -n_j^{(i)} \frac{\varepsilon_i}{|\varepsilon_i|}.
 \end{aligned}$$

Therefore (4.4) reduces to

$$\begin{aligned}
 & \frac{1}{4} \sum_{i=1}^d (L(|\varepsilon_i|, x_i) + 1) \frac{\varepsilon_i}{|\varepsilon_i|} + \frac{1}{12} \sum_{i=1}^d \left(a_i - \frac{x_i}{\varepsilon_i} - \frac{y_{i-1}}{\varepsilon_{i-1}} \right) + \frac{1}{12} \sum_{i=1}^d \sum_{j=1}^{L(|\varepsilon_i|, x_i)} \left(-n_j^{(i)} \frac{\varepsilon_i}{|\varepsilon_i|} \right) \\
 &= \frac{1}{12} \sum_{i=1}^d \left((3(L(|\varepsilon_i|, x_i) + 1) - N(|\varepsilon_i|, x_i)) \frac{\varepsilon_i}{|\varepsilon_i|} + a_i - \frac{x_i + y_i}{\varepsilon_i} \right),
 \end{aligned}$$

proving the theorem. □

REMARK 9. It sometimes happens that a sequence of primitive vectors v_1, \dots, v_d is not unimodular but is unimodular with respect to the sublattice of \mathbb{Z}^2 generated by vectors v_1, \dots, v_d . Such a sequence is called an l -reflexive loop and studied in [5]. Theorem A can be applied to an l -reflexive loop with respect to the sublattice generated by the vectors in the l -reflexive loop, but it is unclear whether the resulting formula can be obtained from Theorem 6.

ACKNOWLEDGEMENT. The author wishes to thank Professor Mikiya Masuda for his valuable advice and continuing support.

References

- [1] W. Fulton: Introduction to Toric Varieties, Annals of Mathematics Studies **131**, Princeton Univ. Press, Princeton, NJ, 1993.
- [2] A. Higashitani and M. Masuda: *Lattice multi-polygons*, arXiv:1204.0088v3.
- [3] R.T. Živaljević: *Rotation number of a unimodular cycle: an elementary approach*, arXiv:1209.4981v4, to appear in Discrete Math.
- [4] M. Masuda: *Unitary toric manifolds, multi-fans and equivariant index*, Tohoku Math. J. (2) **51** (1999), 237–265.
- [5] A.M. Kasprzyk and B. Nill: *Reflexive polytopes of higher index and the number 12*, Electron. J. Combin. **19** (2012).

Department of Mathematics
Graduate School of Science
Osaka City University
3-3-138 Sugimoto, Sumiyoshi-ku
Osaka 558-8585
Japan
e-mail: m13saU0r13@ex.media.osaka-cu.ac.jp