

# A LOGARITHMICALLY IMPROVED REGULARITY CRITERION OF SMOOTH SOLUTIONS FOR THE 3D BOUSSINESQ EQUATIONS

ZHUAN YE

(Received September 17, 2014, revised January 27, 2015)

## Abstract

In this note, we consider the three-dimensional (3D) incompressible Boussinesq equations. We obtain the logarithmically improved regularity criterion of smooth solutions in terms of the velocity field. This result improves some previous works.

## 1. Introduction

This note is devoted to the study of the regularity criterion of smooth solutions for the 3D Boussinesq equations

$$(1.1) \quad \begin{cases} \partial_t u + (u \cdot \nabla)u - \nu \Delta u + \nabla p = \theta e_3, & x \in \mathbb{R}^3, t > 0, \\ \partial_t \theta + (u \cdot \nabla)\theta - \kappa \Delta \theta = 0, & x \in \mathbb{R}^3, t > 0, \\ \nabla \cdot u = 0, & x \in \mathbb{R}^3, t > 0, \\ u(x, 0) = u_0(x), \quad \theta(x, 0) = \theta_0(x), & x \in \mathbb{R}^3, \end{cases}$$

where  $u = u(x, t) \in \mathbb{R}^3$  is the velocity,  $p = p(x, t) \in \mathbb{R}$  is the scalar pressure,  $\theta = \theta(x, t) \in \mathbb{R}$  is the temperature and  $e_3 = (0, 0, 1)^T$ .  $\nu \geq 0$  denotes the viscosity,  $\kappa \geq 0$  denotes the thermal diffusivity. The Boussinesq equations are of relevance to study a number of models coming from atmospheric or oceanographic turbulence where rotation and stratification play an important role (see e.g. [10, 12]).

Local existence and uniqueness theories of solutions to the Boussinesq equations have been studied by many mathematicians and physicists (see, e.g., [1, 2, 9]). But due to the presence of Navier–Stokes equations in the system (1.1) whether this unique local solution can exist globally is an outstanding challenge problem. For this reason, there have been a lot of literatures devoted to finding sufficient conditions to ensure the smoothness of the solutions; see [4, 3, 5, 6, 14, 15, 13, 16, 17, 18, 19] and so forth.

Motivated by the above cited works, our aim is to establish a logarithmically improved regularity criterion of smooth solutions in terms of the velocity field which significantly extends the result in [19]. For the sake of simplicity, we set  $\nu = \kappa = 1$ . More precisely, we will prove

**Theorem 1.1.** *Assume that  $(u_0, \theta_0) \in H^3(\mathbb{R}^3) \times H^3(\mathbb{R}^3)$ . Let  $(u, \theta)$  be a local smooth solution of the system (1.1). If the following condition holds*

$$(1.2) \quad \int_0^T \frac{\|\nabla u(t)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(t)\|_{\dot{B}_{\infty,\infty}^{-1}})} dt < \infty,$$

*then the solution pair  $(u, \theta)$  remains smooth on  $[0, T]$ .*

As a consequence of the fact  $\|\nabla u\|_{\dot{B}_{\infty,\infty}^{-1}} \approx \|u\|_{\dot{B}_{\infty,\infty}^0}$ , we have the following result.

**Corollary 1.2.** *Assume that  $(u_0, \theta_0) \in H^3(\mathbb{R}^3) \times H^3(\mathbb{R}^3)$ . Let  $(u, \theta)$  be a local smooth solution of the system (1.1). If the following condition holds*

$$(1.3) \quad \int_0^T \frac{\|u(t)\|_{\dot{B}_{\infty,\infty}^0}^2}{\ln(e + \|u(t)\|_{\dot{B}_{\infty,\infty}^0})} dt < \infty,$$

*then the solution pair  $(u, \theta)$  remains smooth on  $[0, T]$ .*

REMARK 1.3. As the case  $\theta = 0$ , the system (1.1) reduces to the classical Navier–Stokes equations. It is easy to see that the Corollary 1.2 is a refined improvement of that Theorem 1 in [8] due to the well-known embedding  $BMO \hookrightarrow \dot{B}_{\infty,\infty}^0$ .

## 2. The proof of the Theorem 1.1

This section is devoted to the proof of the Theorem 1.1. Throughout the paper,  $C$  stands for some real positive constants which may be different in each occurrence.

Proof of Theorem 1.1. If (1.2) holds, one can deduce that for any small  $\epsilon > 0$ , there exists  $T_0 = T_0(\epsilon) < T$  such that

$$(2.1) \quad \int_{T_0}^T \frac{\|\nabla u(t)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(t)\|_{\dot{B}_{\infty,\infty}^{-1}})} dt \leq \epsilon.$$

Consequently, the main goal of this section is to establish the following a priori estimate

$$\limsup_{t \rightarrow T^-} (\|\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3 \theta(t)\|_{L^2}^2) < \infty.$$

Thanks to the divergence-free condition  $\nabla \cdot u = 0$ , from the temperature  $\theta$  equation, we immediately have the global a priori bound for  $\theta$  in any Lebesgue space

$$(2.2) \quad \|\theta(t)\|_{L^p} \leq \|\theta_0\|_{L^p}, \quad \forall p \in [2, \infty],$$

for any  $t \in [0, T]$ .

We also have the following basic  $L^2$  energy estimate

$$(2.3) \quad \frac{1}{2} \frac{d}{dt} (\|u(t)\|_{L^2}^2 + \|\theta(t)\|_{L^2}^2) + \|\nabla u\|_{L^2}^2 + \|\nabla \theta\|_{L^2}^2 \leq \|u\|_{L^2} \|\theta\|_{L^2},$$

which together with (2.2) implies that for any  $t \in [0, T]$

$$(2.4) \quad \|u(t)\|_{L^2}^2 + \|\theta(t)\|_{L^2}^2 + \int_0^t (\|\nabla u\|_{L^2}^2 + \|\nabla \theta\|_{L^2}^2)(\tau) d\tau \leq C < \infty.$$

Multiplying the equation of (1.1)<sub>1</sub> and (1.1)<sub>2</sub> by  $\Delta u$  and  $\Delta \theta$ , respectively, integration by parts and taking the divergence free property into account, one concludes that

$$(2.5) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|\nabla u(t)\|_{L^2}^2 + \|\nabla \theta(t)\|_{L^2}^2) + \|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2 \\ &= - \int_{\mathbb{R}^3} \theta e_3 \cdot \Delta u \, dx + \int_{\mathbb{R}^3} (u \cdot \nabla u) \cdot \Delta u \, dx + \int_{\mathbb{R}^3} (u \cdot \nabla \theta) \cdot \Delta \theta \, dx \\ &:= N_1 + N_2 + N_3. \end{aligned}$$

Integrating by parts and Young inequality, it yields

$$(2.6) \quad N_1 \leq C \|\nabla u\|_{L^2} \|\nabla \theta\|_{L^2}.$$

In order to deal with the terms  $N_2$  and  $N_3$ , we need the following interpolation inequality due to Meyer–Gerard–Oru [11]

$$(2.7) \quad \|f\|_{L^4} \leq C \|\nabla f\|_{L^2}^{1/2} \|f\|_{\dot{B}_{\infty,\infty}^{-1}}^{1/2}, \quad \forall f \in \dot{H}^1(\mathbb{R}^3) \cap \dot{B}_{\infty,\infty}^{-1}(\mathbb{R}^3).$$

By the above interpolation inequality (2.7) and Young inequality, we can bound the terms  $N_2$  and  $N_3$  as

$$(2.8) \quad \begin{aligned} N_2 &= - \int_{\mathbb{R}^3} (\partial_k u \cdot \nabla u) \cdot \partial_k u \, dx \\ &\leq C \|\nabla u\|_{L^2} \|\nabla u\|_{L^4}^2 \leq C \|\nabla u\|_{L^2} \|\nabla^2 u\|_{L^2} \|\nabla u\|_{\dot{B}_{\infty,\infty}^{-1}} \\ &\leq \frac{1}{2} \|\nabla^2 u\|_{L^2}^2 + C \|\nabla u\|_{\dot{B}_{\infty,\infty}^{-1}}^2 \|\nabla u\|_{L^2}^2, \end{aligned}$$

and

$$(2.9) \quad \begin{aligned} N_3 &= - \int_{\mathbb{R}^3} (\partial_k u \cdot \nabla \theta) \cdot \partial_k \theta \, dx \\ &\leq C \|\nabla u\|_{L^2} \|\nabla \theta\|_{L^4}^2 \leq C \|\nabla u\|_{L^2} \|\nabla^2 \theta\|_{L^2} \|\nabla \theta\|_{\dot{B}_{\infty,\infty}^{-1}} \\ &\leq C \|\nabla u\|_{L^2} \|\nabla^2 \theta\|_{L^2} \|\theta\|_{\dot{B}_{\infty,\infty}^0} \leq C \|\nabla u\|_{L^2} \|\nabla^2 \theta\|_{L^2} \|\theta\|_{L^\infty} \\ &\leq \frac{1}{2} \|\nabla^2 \theta\|_{L^2}^2 + C \|\theta\|_{L^\infty}^2 \|\nabla \theta\|_{L^2}^2, \end{aligned}$$

where we have used the following fact in (2.9)

$$\|\nabla\theta\|_{\dot{B}_{\infty,\infty}^{-1}} \leq C\|\theta\|_{\dot{B}_{\infty,\infty}^0} \leq C\|\theta\|_{L^\infty}.$$

Substituting (2.6), (2.8), and (2.9) into (2.5), we arrive at

$$(2.10) \quad \begin{aligned} & \frac{d}{dt}(\|\nabla u(t)\|_{L^2}^2 + \|\nabla\theta(t)\|_{L^2}^2) + \|\nabla^2 u\|_{L^2}^2 + \|\nabla^2\theta\|_{L^2}^2 \\ & \leq C(1 + \|\theta\|_{L^\infty}^2 + \|\nabla u\|_{\dot{B}_{\infty,\infty}^{-1}}^2)(\|\nabla u\|_{L^2}^2 + \|\nabla\theta\|_{L^2}^2). \end{aligned}$$

For any  $t \in (T_0, T)$ , we denote

$$y(t) := \max_{\tau \in (T_0, t]} (\|u(\tau)\|_{H^3}^2 + \|\theta(\tau)\|_{H^3}^2).$$

It should be noted that  $y(t)$  is a nondecreasing function.

Thanks to the Gronwall inequality, it follows from (2.10) that for any  $T_0 \leq t < T$

$$(2.11) \quad \begin{aligned} & \|\nabla u(t)\|_{L^2}^2 + \|\nabla\theta(t)\|_{L^2}^2 + \int_{T_0}^t (\|\nabla^2 u(\tau)\|_{L^2}^2 + \|\nabla^2\theta(\tau)\|_{L^2}^2) d\tau \\ & \leq (\|\nabla u(T_0)\|_{L^2}^2 + \|\nabla\theta(T_0)\|_{L^2}^2) \exp\left[\tilde{C} \int_{T_0}^t (1 + \|\theta\|_{L^\infty} + \|\nabla u\|_{\dot{B}_{\infty,\infty}^{-1}}^2)(\tau) d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2 d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \frac{\|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}})} \ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}) d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \frac{\|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}})} \ln(e + \|u(\tau)\|_{\dot{B}_{\infty,\infty}^0}) d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \frac{\|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}})} \ln(e + \|u(\tau)\|_{L^\infty}) d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \frac{\|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}})} \ln(e + \|u(\tau)\|_{H^3}) d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \frac{\|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}})} \ln(e + y(\tau)) d\tau\right] \\ & \leq M \exp\left[\tilde{C} \int_{T_0}^t \frac{\|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}}^2}{\ln(e + \|\nabla u(\tau)\|_{\dot{B}_{\infty,\infty}^{-1}})} d\tau \ln(e + y(t))\right]. \end{aligned}$$

We want to state here that from the above observation  $\tilde{C}$  is an absolute constant and  $M$  depends on  $\|\nabla u(T_0)\|_{L^2}$ ,  $\|\nabla\theta(T_0)\|_{L^2}$ ,  $T_0$ ,  $T$  and  $\theta_0$ .

It follows from the condition (2.1) that

$$(2.12) \quad \|\nabla u(t)\|_{L^2}^2 + \|\nabla\theta(t)\|_{L^2}^2 + \int_{T_0}^t (\|\nabla^2 u(\tau)\|_{L^2}^2 + \|\nabla^2\theta(\tau)\|_{L^2}^2) d\tau \leq C(e + y(t))^{\tilde{C}\epsilon}.$$

Applying  $\nabla^3$  to the equation (1.1)<sub>1</sub> and (1.1)<sub>2</sub>, multiplying the resulting equations by  $\nabla^3 u$  and  $\nabla^3\theta$  respectively and adding them up, we have

$$(2.13) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} (\|\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3\theta(t)\|_{L^2}^2) + \|\nabla^4 u\|_{L^2}^2 + \|\nabla^4\theta\|_{L^2}^2 \\ &= \int_{\mathbb{R}^3} \nabla^3\theta e_3 \cdot \nabla^3 u \, dx - \int_{\mathbb{R}^3} \nabla^3(u \cdot \nabla u) \cdot \nabla^3 u \, dx - \int_{\mathbb{R}^3} \nabla^3(u \cdot \nabla\theta) \cdot \nabla^3\theta \, dx \\ &:= K_1 + K_2 + K_3. \end{aligned}$$

It follows from Young inequality that

$$(2.14) \quad \begin{aligned} K_1 &\leq C \|\nabla^3 u\|_{L^2} \|\nabla^3\theta\|_{L^2} \\ &\leq C \|\nabla^2 u\|_{L^2}^{1/2} \|\nabla^4 u\|_{L^2}^{1/2} \|\nabla^2\theta\|_{L^2}^{1/2} \|\nabla^4\theta\|_{L^2}^{1/2} \\ &\leq \frac{1}{4} \|\nabla^4 u\|_{L^2}^2 + \frac{1}{4} \|\nabla^4\theta\|_{L^2}^2 + C(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2\theta\|_{L^2}^2), \end{aligned}$$

where we have applied the following Gagliardo–Nirenberg inequality

$$\|\nabla^3 f\|_{L^2} \leq C \|\nabla^2 f\|_{L^2}^{1/2} \|\nabla^4 f\|_{L^2}^{1/2}.$$

Now we recall the following commutator estimate (see [7])

$$(2.15) \quad \|[\Lambda^s, f]g\|_{L^p} \leq C(\|\nabla f\|_{L^{p_1}} \|\Lambda^{s-1}g\|_{L^{p_2}} + \|\Lambda^s f\|_{L^{p_3}} \|g\|_{L^{p_4}}),$$

with  $s > 0$ ,  $p_2, p_3 \in (1, \infty)$  such that  $1/p = 1/p_1 + 1/p_2 = 1/p_3 + 1/p_4$ .

From the divergence-free condition and the commutator estimate (2.15), we obtain

$$\begin{aligned} K_2 &= - \int_{\mathbb{R}^3} [\nabla^3, u \cdot \nabla] u \cdot \nabla^3 u \, dx \\ &\leq C \|\nabla u\|_{L^2} \|\nabla^3 u\|_{L^4}^2 \\ &\leq C \|\nabla u\|_{L^2} \|\nabla^2 u\|_{L^2}^{1/4} \|\nabla^4 u\|_{L^2}^{7/4} \\ &\leq \frac{1}{8} \|\nabla^4 u\|_{L^2}^2 + C \|\nabla u\|_{L^2}^8 \|\nabla^2 u\|_{L^2}^2, \end{aligned}$$

where we have used the following Gagliardo–Nirenberg inequality

$$\|\nabla^3 u\|_{L^4} \leq C \|\nabla^2 u\|_{L^2}^{1/8} \|\nabla^4 u\|_{L^2}^{7/8}.$$

Similar to the estimate of  $K_2$ , the term  $K_3$  can be bounded as

$$\begin{aligned}
 (2.16) \quad K_3 &= - \int_{\mathbb{R}^3} [\nabla^3, u \cdot \nabla] \theta \cdot \nabla^3 \theta \, dx \\
 &\leq C \|\nabla u\|_{L^2} \|\nabla^3 \theta\|_{L^4}^2 + \|\nabla \theta\|_{L^2} \|\nabla^3 u\|_{L^4} \|\nabla^3 \theta\|_{L^4} \\
 &\leq C \|\nabla u\|_{L^2} \|\nabla^2 \theta\|_{L^2}^{1/4} \|\nabla^4 \theta\|_{L^2}^{7/4} + \|\nabla \theta\|_{L^2} \|\nabla^2 u\|_{L^2}^{1/8} \|\nabla^4 u\|_{L^2}^{7/8} \|\nabla^2 \theta\|_{L^2}^{1/8} \|\nabla^4 \theta\|_{L^2}^{7/8} \\
 &\leq \frac{1}{8} \|\nabla^4 u\|_{L^2}^2 + \frac{1}{4} \|\nabla^4 \theta\|_{L^2}^2 + C(\|\nabla u\|_{L^2}^8 + \|\nabla \theta\|_{L^2}^8)(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2).
 \end{aligned}$$

Combining the previous estimates, we get

$$\begin{aligned}
 (2.17) \quad \frac{d}{dt} (\|\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3 \theta(t)\|_{L^2}^2) \\
 \leq C(\|\nabla u\|_{L^2}^8 + \|\nabla \theta\|_{L^2}^8)(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2) + C(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2).
 \end{aligned}$$

Integrating the inequality (2.17) over  $(T_0, t)$ , we easily get

$$\begin{aligned}
 (2.18) \quad &\|\nabla^3 u(t)\|_{L^2}^2 + \|\nabla^3 \theta(t)\|_{L^2}^2 - (\|\nabla^3 u(T_0)\|_{L^2}^2 + \|\nabla^3 \theta(T_0)\|_{L^2}^2) \\
 &\leq C \int_{T_0}^t (\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2) \, d\tau + C \int_{T_0}^t (\|\nabla u\|_{L^2}^8 + \|\nabla \theta\|_{L^2}^8)(\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2) \, d\tau \\
 &\leq C(e + y(t))^{\tilde{C}\epsilon} + C \int_{T_0}^t (e + y(\tau))^{4\tilde{C}\epsilon} (\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2)(\tau) \, d\tau \\
 &\leq C(e + y(t))^{\tilde{C}\epsilon} + C(e + y(t))^{4\tilde{C}\epsilon} \int_{T_0}^t (\|\nabla^2 u\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2)(\tau) \, d\tau \\
 &\leq C(e + y(t))^{\tilde{C}\epsilon} + C(e + y(t))^{5\tilde{C}\epsilon} \\
 &\leq C(e + y(t))^{5\tilde{C}\epsilon},
 \end{aligned}$$

which immediately implies that

$$e + y(t) \leq C_{T_0} + C(e + y(t))^{5\tilde{C}\epsilon}, \quad C_{T_0} = \|\nabla^3 u(T_0)\|_{L^2}^2 + \|\nabla^3 \theta(T_0)\|_{L^2}^2.$$

By appropriately selecting  $\epsilon < 1/(5\tilde{C})$ , the above inequality allows us to show

$$y(t) \leq C(\|\nabla^3 u(T_0)\|_{L^2}, \|\nabla^3 \theta(T_0)\|_{L^2}, T_0, T) < \infty, \quad \forall t \in [T_0, T].$$

As a consequence, we get the boundness of  $H^3 \times H^3$ -norm of  $(u, \theta)$  for all  $t \in [0, T]$ . Consequently, The proof of Theorem 1.1 is completed.  $\square$

**ACKNOWLEDGEMENTS.** The author would like to thank the referees for their valuable comments and suggestions, which are very helpful for author to improve this paper.

## References

- [1] D. Chae and H.-S. Nam: *Local existence and blow-up criterion for the Boussinesq equations*, Proc. Roy. Soc. Edinburgh Sect. A **127** (1997), 935–946.
- [2] D. Chae, S.-K. Kim and H.-S. Nam: *Local existence and blow-up criterion of Hölder continuous solutions of the Boussinesq equations*, Nagoya Math. J. **155** (1999), 55–80.
- [3] J. Fan and T. Ozawa: *Regularity criteria for the 3D density-dependent Boussinesq equations*, Nonlinearity **22** (2009), 553–568.
- [4] J. Fan and Y. Zhou: *A note on regularity criterion for the 3D Boussinesq system with partial viscosity*, Appl. Math. Lett. **22** (2009), 802–805.
- [5] S. Gala, Z. Guo and M.A. Ragusa: *A remark on the regularity criterion of Boussinesq equations with zero heat conductivity*, Appl. Math. Lett. **27** (2014), 70–73.
- [6] Z. Guo and S. Gala: *Regularity criterion of the Newton–Boussinesq equations in  $\mathbb{R}^3$* , Commun. Pure Appl. Anal. **11** (2012), 443–451.
- [7] T. Kato and G. Ponce: *Commutator estimates and the Euler and Navier–Stokes equations*, Comm. Pure Appl. Math. **41** (1988), 891–907.
- [8] H. Kozono and Y. Taniuchi: *Bilinear estimates in BMO and the Navier–Stokes equations*, Math. Z. **235** (2000), 173–194.
- [9] A.J. Majda and A.L. Bertozzi: *Vorticity and Incompressible Flow*, Cambridge Univ. Press, Cambridge, 2002.
- [10] A. Majda: *Introduction to PDEs and Waves for the Atmosphere and Ocean*, Courant Lecture Notes in Mathematics **9**, New York Univ., Courant Inst. Math. Sci., New York, 2003.
- [11] P. Gerard, Y. Meyer and F. Oru: *Inégalités de Sobolev précisées*; in Séminaire sur les Équations aux Dérivées Partielles, 1996–1997, Exp. IV, 11 pp, École Polytech., Palaiseau.
- [12] J. Pedlosky: *Geophysical fluid dynamics*, New York, Springer-Verlag, 1987.
- [13] Y. Qin, X. Yang, Y. Wang, X. Liu: *Blow-up criteria of smooth solutions to the 3D Boussinesq equations*, Math. Methods Appl. Sci. **35** (2012), 278–285.
- [14] Y. Taniuchi: *A note on the blow-up criterion for the inviscid 2-D Boussinesq equations*; in The Navier–Stokes Equations: Theory and Numerical Methods (Varenna, 2000), Lecture Notes in Pure and Appl. Math. **223**, Dekker, New York, 131–140.
- [15] H. Qiu, Y. Du and Z. Yao: *A blow-up criterion for 3D Boussinesq equations in Besov spaces*, Nonlinear Anal. **73** (2010), 806–815.
- [16] Z. Xiang: *The regularity criterion of the weak solution to the 3D viscous Boussinesq equations in Besov spaces*, Math. Methods Appl. Sci. **34** (2011), 360–372.
- [17] F. Xu, Q. Zhang and X. Zheng: *Regularity criteria of the 3D Boussinesq equations in the Morrey–Campanato space*, Acta Appl. Math. **121** (2012), 231–240.
- [18] Z. Ye: *Blow-up criterion of smooth solutions for the Boussinesq equations*, Nonlinear Anal. **110** (2014), 97–103.
- [19] Z. Zhang: *Some regularity criteria for the 3D Boussinesq equations in the class  $L^2(0, T; \dot{B}_{\infty, \infty}^{-1})$* , ISRN Appl. Math. (2014), Art. ID 564758, 4 pp.

School of Mathematical Sciences  
 Beijing Normal University  
 Laboratory of Mathematics and Complex Systems, Ministry of Education  
 Beijing 100875  
 People's Republic of China  
 e-mail: yezhuan815@126.com