

# CURVES WITH MAXIMALLY COMPUTED CLIFFORD INDEX

TAKAO KATO and GERRIET MARTENS

(Received November 17, 2017)

## Abstract

We say that a curve  $X$  of genus  $g$  has maximally computed Clifford index if the Clifford index  $c$  of  $X$  is, for  $c > 2$ , computed by a linear series of the maximum possible degree  $d < g$ ; then  $d = 2c + 3$  resp.  $d = 2c + 4$  for odd resp. even  $c$ . For odd  $c$  such curves have been studied in [6]. In this paper we analyze if/how far analogous results hold for such curves of even Clifford index  $c$ .

## 1. Introduction

Let  $X$  denote a smooth irreducible projective curve defined over the complex numbers, and let  $g \geq 4$  resp.  $c \geq 0$  denote its genus resp. its Clifford index. We say that a (complete and base point free) linear series  $g_d^r$  on  $X$ , or a divisor in it, computes  $c$  if  $d < g$ ,  $r > 0$  and  $d - 2r = c$ . It is well known ([5, Thm. C]) that in this case we have  $d \leq 2c + 4$  if  $X$  is neither hyper- nor bi-elliptic (which certainly holds for  $c > 2$ ). For  $c > 2$  we say that the Clifford index  $c$  of  $X$  is *maximally computed* if  $X$  has a  $g_d^r$  computing  $c$  of the maximal possible degree, i.e.  $d = 2c + 3$  resp.  $d = 2c + 4$  if  $c$  is odd resp. even. Such curves exist for every  $c > 2$  ([5, 3.3]) and examples are constructed on K3 surfaces.

Let  $X$  be such a curve. Then we have  $g = d + 1$  ([5, 3.2.5]).

For odd  $c$  we also know:  $X$  has gonality  $c + 3$  and infinitely many pencils  $g_{c+3}^1$  ([5, 3.2.2 and 2.3]), and by [6], 3.6 and 3.7 the  $g_d^r$  is the only series on  $X$  computing  $c$  (in particular, it is half-canonical, i.e.  $|2g_d^r|$  is the canonical series of  $X$ , and very ample); moreover, the  $g_d^r$  is even normally generated.

For even  $c$  our knowledge on  $X$  is less complete ([5], [10]) mainly because a basic Diophantine equation ([6, sections 1 and 2]) valid for  $X$  in the case of odd  $c$  is not available if  $X$  has even Clifford index. One knows, for even  $c$ :

- $X$  has gonality  $c + 2$ ,
- for every pencil  $|D|$  of degree  $c + 2$  on  $X$  there is a pencil  $|D'|$  of degree  $c + 2$  on  $X$  such that  $g_d^r = |D + D'|$  ([5, 3.2.3 and 3.2.4]),
- $X$  has no base point free pencil of degree  $c + 3$  ([5, 3.2.1]),
- $X$  has no series computing  $c$  of degree  $e$  with  $3(c + 2)/2 < e < 2(c + 2) = d$  ([13, Cor. 1]); note that this implies that our  $g_d^r$  must be very ample.

In [5, 3.3.2] the following "recognition theorem" is proved: On any  $k$ -gonal curve ( $k \geq 3$ ) having only finitely many base point free pencils of degree  $k$  and  $k + 1$ , a linear series  $g_d^r$ ,

---

2010 Mathematics Subject Classification. Primary 14H45; Secondary 14H51.

The first named author expresses his thanks to Univ. Erlangen-Nürnberg for the hospitality during his stay.

$r \geq 2$ , computing its Clifford index  $c$  computes  $c$  maximally and is the only linear series computing  $c$  which is not a pencil. (Note that  $c$  is even, then, and the  $g_d^r$  is half-canonical.) Moreover, it follows that the  $g_d^r$  is even normally generated: Since there are (by assumption) only finitely many  $g_{c+2}^1$  the curve embedded into  $\mathbb{P}^r$  by the  $g_d^r$  lies on only finitely many quadrics of rank  $\leq 4$  which implies (cf. [2, III, ex. D-1 and V, ex. C-7]) that it is quadratically normal, and to see that it is  $n$ -normal for all other integers  $n \geq 1$  we can use Green's results on Koszul-cohomology, as is done in [6, proof of Theorem 3.6].

However, there are curves whose (even) Clifford index  $c$  is computed maximally which have infinitely many pencils of degree  $c + 2$ ; this will be shown in the next section where we discuss the case  $c = 4$  in greater detail. So the recognition theorem does not always answer the

QUESTION. Is, on our  $X$ , the  $g_d^r$  the only linear series computing  $c$  which is not a pencil?

In this paper we deal with this Question. For  $c \equiv 0 \pmod 4$  we prove in Section 3 that every effective divisor of  $X$  computing  $c$  is contained in a divisor of the  $g_d^r$ ; in particular, the  $g_d^r$  is then the only linear series on  $X$  computing  $c$  maximally. And for  $c = 4, c = 6$  and  $c = 8$  we answer our Question in the affirmative. Finally, for  $X$  lying, via the  $g_d^r$ , on a K3 surface of degree  $2r - 2$  we check if the divisor theory of the surface may be helpful to provide a negative answer.

NOTATION. The basic reference is [2]. For any curve  $X$ ,  $\text{Div}(X)$  denotes its group of divisors and the symbol  $\sim$  means the linear equivalence of divisors. For  $D, E \in \text{Div}(X)$  we write  $D \leq E$  (and say that  $D$  is contained in  $E$ ) if  $E - D$  is effective, i.e.  $E - D \geq 0$ , and for linear series  $g_d^r, g_e^s$  on  $X$  the notation  $g_d^r \subset g_e^s$  means that every divisor in  $g_d^r$  is contained in a divisor of  $g_e^s$  (equivalently,  $|g_e^s - g_d^r| \neq \emptyset$ ). We sometimes identify a complete  $g_d^r$  on  $X$  with the point in the variety  $W_d^r = W_d^r(X)$  corresponding to it via the Abel-Jacobi map. (Specifically, for a canonical divisor  $K_X$  of  $X$  the canonical series  $|K_X|$  likewise is the only point in  $W_{2g-2}^{g-1}$ , for  $g > 0$ .)

## 2. Clifford index $c = 4$

For  $c = 4$  we construct a curve whose Clifford index  $c$  is maximally computed and satisfies  $\dim(W_6^1) > 0$ .

EXAMPLE. Let  $E$  denote a smooth elliptic curve and  $S \rightarrow E$  be a ruled surface with invariant  $e \geq 0$ . Using the notations of [9, V, 2] we can find a smooth elliptic curve  $H$  in the numerical equivalence class of  $C_0 + e \cdot f$  ( $C_0^2 = -e, f$  a fibre); we have  $h^0(H) = e + 1$ , and  $-C_0 - H$  is a canonical divisor of  $S$  ([8, 3.3]). Observe that  $H^2 = e$  and  $C_0 \cdot H = 0$ . For  $e > 0$  we consider the divisor  $D := 3H$  of  $S$ ; then  $|D|$  is base point free and so a general member  $X$  in  $|D|$  is a smooth curve, by Bertini's theorem. Writing  $X = X_1 + X_2$  with effective divisors  $X_1, X_2$  of  $S$ , we thus must have  $X_1 \cdot X_2 = 0$ . If  $X_1 \equiv \alpha C_0 + \beta f$  (here  $\equiv$  denotes numerical equivalence) we have  $X_2 \equiv (3 - \alpha)C_0 + (3e - \beta)f$  with integers  $\alpha, \beta \geq 0, \alpha \leq 3, \beta \leq 3e$  ([8, 3.1]), and  $X_1 \cdot X_2 = 0$  implies the relation  $(2\beta - e\alpha)(2\alpha - 3) = 3e\alpha$  leading to  $X_1 \equiv 0$  for  $\alpha = 0$  resp.  $X_2 \equiv 0$  for  $\alpha = 3$  and  $\beta = -e < 0$  for  $\alpha = 1$  resp.  $\beta = 4e > 3e$  for  $\alpha = 2$ . Thus it

follows that  $X$  is irreducible, and its genus is, by adjunction,  $g = 3e + 1$ .

Now, let  $e = 4$  and view the base curve  $E$  as an elliptic normal curve in  $\mathbb{P}^{e-1} = \mathbb{P}^3$  (of degree  $e = 4$ ); let  $S_0$  denote the cone over  $E$  in  $\mathbb{P}^4$ . Blowing up the vertex of the elliptic cone  $S_0$  we obtain a ruled surface  $S \rightarrow E$  of invariant  $e = 4$  as above ([9, V, 2.11.4]), and the blow down  $S \rightarrow S_0 \subset \mathbb{P}^4$  is defined by  $|H|$ . Our curve  $X \subset S$  from above blows down to a curve  $X' \subset S_0$  of degree  $X \cdot H = 3H^2 = 3e = 12 = g - 1$  and  $X'$  is smooth since it misses the vertex of  $S_0$ . Since  $h^0(S, H - X) = h^0(S, -2H) = 0$  the linear series  $|H|$  of  $S$  cuts out on  $X$  a (maybe, incomplete) linear series of degree 12 and dimension  $h^0(H) - 1 = e = 4$ . Hence  $X$  has Clifford index  $c \leq 12 - 2 \cdot 4 = 4$ . To see that  $c = 4$  we recall that on a curve of genus 13 its Clifford index can be computed by pencils; so we have to show that the gonality  $k$  of  $X$  is 6. Since the natural map  $\pi : X \subset S \rightarrow E$  has degree  $X \cdot f = 3$  our curve  $X$  is a triple covering of an elliptic curve; in particular,  $X$  has infinitely many  $g_6^1$ . If  $k < 6$  we obtain, according to Castelnuovo's genus formula for curves with independent morphisms ([2, VIII, ex. C-1]), that  $g \leq (k - 1)(3 - 1) + 3g(E) = 2k + 1 \leq 11$ , a contradiction. So  $k = 6$ ,  $c = 4$ , and the series  $|H|_X$  is a complete (and very ample)  $g_{12}^4$  on  $X$  thus computing  $c = 4$  maximally. Since  $W_6^1(X)$  contains (at least) the one-dimensional irreducible component  $\pi^*W_2^1(E)$  we clearly have  $\dim(W_6^1(X)) > 0$ .

**Proposition 2.1.** *Let  $X$  be a curve whose Clifford index  $c = 4$  is computed maximally. Assume that  $\dim(W_6^1) > 0$ . Then  $X$  admits a triple covering  $\pi : X \rightarrow E$  over an elliptic curve  $E$ ,  $\pi^*(W_2^1(E))$  is the only infinite irreducible component of  $W_6^1$ , and this component is singular with finitely many singularities. Furthermore,  $X$  has only one series  $g_{12}^4$  (computing  $c$  maximally), and the variety  $W_{12}^4$  is not reduced.*

Proof. By de Franchis' theorem, on any  $k$ -gonal curve  $X$  with an infinite set  $S$  of  $g_k^1$  either infinitely many  $g_k^1$  in  $S$  are compounded of the same irrational involution or there are only finitely many compounded  $g_k^1$  in  $S$ . For  $k = 6$ , in the latter case such a curve is a smooth plane septic ( $g = 15$ ) or we have  $g \leq 11$  ([4]), and in the first case infinitely many  $g_6^1$  in  $S$  are induced by a covering  $\rho : X \rightarrow Y$  over a non-hyperelliptic curve  $Y$  of genus 3 or by a triple covering  $\pi : X \rightarrow E$  over an elliptic curve  $E$ . Now, let  $X$  be a curve whose Clifford index  $c = 4$  is computed maximally and admitting infinitely many  $g_6^1$ . Since  $g = 13$  we then are in the first case from above.

Assume that  $\rho : X \rightarrow Y$  is a double covering of  $X$  over a curve  $Y$  of genus 3. Then  $Y$  is a smooth plane quartic, and every  $g_6^1$  on  $X$  is induced by  $\rho$  since otherwise we would have  $g \leq (6 - 1)(2 - 1) + 2g(Y) = 11$ , by Castelnuovo's genus formula for curves with independent morphisms. Hence we have  $W_6^1(X) = \rho^*(W_3^1(Y)) = \rho^*K_Y - \rho^*(W_1(Y))$ . Since we know that there are pencils  $g_6^1, h_6^1$  on  $X$  such that  $g_{12}^4 = |g_6^1 + h_6^1|$  we thus have pencils  $L_1, L_2$  of degree 3 on  $Y$  such that  $g_{12}^4 = |\rho^*(L_1) + \rho^*(L_2)|$ . But (cf. [12, p. 1797])

$$h^0(X, \rho^*(L_1 + L_2)) = h^0(Y, L_1 + L_2) + h^0(Y, (L_1 + L_2) - D) = 4 + h^0(Y, L_1 + L_2 - D)$$

for a divisor  $D$  of  $Y$  such that  $2D$  is linearly equivalent to the branch divisor  $B$  of  $\rho$  (i.e.  $B$  is made up by the points of  $Y$  over which  $\rho$  ramifies). So  $2\deg(D) = \deg(B) = 2g - 2 - 2(2g(Y) - 2) = 16$ , i.e.  $\deg(D) = 8 > 6 = \deg(L_1 + L_2)$  which implies that  $h^0(Y, L_1 + L_2 - D) = 0$ . Thus we obtain  $h^0(X, \rho^*(L_1 + L_2)) = 4$  which contradicts  $|\rho^*(L_1 + L_2)| = g_{12}^4$ .

So  $X$  admits a triple covering  $\pi : X \rightarrow E$  over an elliptic curve  $E$ . Our very ample  $g_{12}^4$

embeds  $X$  as a curve of degree 12 in  $\mathbb{P}^4$ . Assume that there is another series on  $X$  computing  $c$  maximally, i.e. a  $h_{12}^4 \neq g_{12}^4$ . Then  $|h_{12}^4 - g_{12}^4| = \emptyset$ , and, according to a refinement of the base point free pencil trick ([2, III, ex. B-6]) we have:  $\dim(|h_{12}^4 + g_{12}^4|) \geq 2 \cdot 4 - \dim(|h_{12}^4 - g_{12}^4|) + 4 - 1 = 12 = g - 1$  whence  $h_{12}^4 = |K_X - g_{12}^4|$  and so  $|2g_{12}^4| \neq |K_X|$ . Thus it follows that  $\dim(|2g_{12}^4|) = g - 2 = 11 = 3 \cdot 4 - 1$ , and so a result of Castelnuovo ([2, p. 120]) implies that  $X$  lies on a non-degenerate surface  $S$  of minimal degree in  $\mathbb{P}^4$ , i.e. on a cubic rational normal scroll. But this is impossible: By Segre’s formula for curves on a rational normal scroll whose ruling consists of  $n$ -secant lines for the curve, we obtain  $13 = g = (n-1)(\deg(X)-1-(n/2)\deg(S)) = (n-1)(12-1-(n/2)\cdot 3)$  which cannot hold. Consequently, we see that a  $h_{12}^4 \neq g_{12}^4$  cannot exist on  $X$ , i.e.  $W_{12}^4$  is a point, and this point is not a smooth point of  $W_{12}^4$  since the tangent space to  $W_{12}^4$  at it has positive dimension ([2, IV, ex. A-2]; observe that the unique  $g_{12}^4$  on  $X$  is half-canonical).

$W_6^1(X)$  has the irreducible component  $\pi^*(W_2^1(E))$ . The argument in the beginning of this proof shows that a further infinite irreducible component of  $W_6^1(X)$  gives rise to a second triple covering  $\pi^* : X \rightarrow E'$  over an elliptic curve  $E'$ ; but applying Castelnuovo’s genus bound for curves admitting independent morphisms to the pair  $(\pi, \pi')$  of coverings we get the contradiction  $g \leq (3 - 1)(3 - 1) + 3g(E) + 3g(E') = 10$ .

For simplicity we identify our  $g_{12}^4$  on  $X$  with the point  $\ell$  of  $W_{12}^4(X)$  corresponding to it. Then the irreducible component  $\ell - \pi^*(W_2^1(E))$  of  $W_6^1(X)$  coincides with  $\pi^*(W_2^1(E))$ . Hence there are four points  $p_1, \dots, p_4 \in E$  such that  $\ell = |\pi^*(p_1 + \dots + p_4)|$ . Since, on  $E$ ,  $p_1 + \dots + p_4 \sim 2q_1 + 2q_2$  for two points  $q_1, q_2 \in E$  there exists a  $g_6^1 = |\pi^*(q_1 + q_2)|$  on  $X$  such that  $|2g_6^1| = \ell$ , and since  $X$  has only finitely many 2-torsion points  $X$  has only a finite number of such  $g_6^1$ . Recall that the embedding series  $\ell$  is the only  $g_{12}^4$  on  $X$ . Hence  $|2g_6^1| = \ell$  is equivalent with  $\dim|2g_6^1| \geq 4$ , and it follows ([2, IV, 4.2]) that the  $g_6^1$  in  $\pi^*(W_2^1(E))$  satisfying  $|2g_6^1| = \ell$  correspond to the singularities of the component  $\pi^*(W_2^1(E))$  of  $W_6^1(X)$ .  $\square$

Though  $\dim(W_6^1) > 0$  is possible, on every curve  $X$  whose Clifford index  $c = 4$  is computed maximally only the unique  $g_{12}^4$  and the pencils of degree 6 compute  $c$ . To see this, recall that  $X$  has no series computing  $c$  of degree  $d$  with  $3(c + 2)/2 < d < 2(c + 2)$ , i.e. no  $g_{10}^3$ . A  $g_8^2$  on  $X$  (computing  $c$ ) cannot be simple since we know that  $W_7^1 = W_6^1 + W_1$  which implies that  $|g_8^2 - P|$  has a base point, for every point  $P \in X$ . So a  $g_8^2$  on  $X$  is compounded thus inducing a double covering  $\rho : X \rightarrow Y$  over a smooth plane quartic, i.e. over a non-hyperelliptic curve of genus 3. But in the proof of the Proposition we observed already that this is impossible.

Finally, we just note that one can show that the curve  $X$  of Proposition 2.1 is as in the example. (In fact, viewing  $X$  as being embedded by the  $g_{12}^4$  it lies in the intersection of two irreducible quadrics in  $\mathbb{P}^4$ , i.e. on a surface of degree 4 which turns out to be an elliptic cone.)

### 3. The main result

The following general result is elementary but useful, for our purposes.

**Lemma 3.1.** *On any curve  $Y$  of genus  $g$  and Clifford index  $c$  let  $D, E$  be effective divisors computing  $c$ . Then the greatest common divisor  $(D, E)$  of  $D$  and  $E$  has Clifford index  $\text{cliff}((D, E)) \leq c$ , and if  $\dim |(D, E)| > 0$  then  $(D, E)$  and one of the divisors  $D + E - (D, E)$*

(the "least common multiple" of  $D$  and  $E$ ) resp. its dual  $K_Y - (D + E - (D, E))$  compute  $c$ .

Proof. Recall that, for a divisor  $\Delta$  of  $Y$ , we have  $\text{cliff}(\Delta) = \text{deg}(\Delta) - 2h^0(\Delta) + 2$ ,  $\text{cliff}(K_X - \Delta) = \text{cliff}(\Delta)$ , and that the Clifford index  $c$  of  $Y$  is the minimum of all  $\text{cliff}(\Delta)$  such that  $h^0(\Delta) \geq 2$  and  $h^1(\Delta) \geq 2$  holds.

It is easy to prove the inequality (cf. [14, 2.21])

$$\text{cliff}(D) + \text{cliff}(E) \geq \text{cliff}((D, E)) + \text{cliff}(D + E - (D, E)).$$

Since  $\text{cliff}(D) = c = \text{cliff}(E)$  the first claim of the Lemma follows from this inequality provided that  $\text{cliff}(D + E - (D, E)) \geq c$ . So assume that  $\text{cliff}(D + E - (D, E)) < c$ . Since  $h^0(D + E - (D, E)) \geq h^0(D) \geq 2$  we then must have  $h^1(D + E - (D, E)) \leq 1$ , and so we obtain  $c > \text{cliff}(D + E - (D, E)) = \text{cliff}(K_Y - (D + E - (D, E))) = 2g - 2 - (\text{deg}(D) + \text{deg}(E) - \text{deg}((D, E))) - 2h^1(D + E - (D, E)) + 2 \geq \text{deg}((D, E))$  (recall that  $\text{deg}(D) < g$  and  $\text{deg}(E) < g$ ). But  $\text{deg}((D, E)) < c$  implies that  $h^0((D, E)) = 1$  whence it follows that  $\text{cliff}((D, E)) = \text{deg}((D, E)) < c$ .

Assume that  $h^0((D, E)) \geq 2$ . We then have  $\text{cliff}((D, E)) \geq c$ , and by the (just proved) first claim of the Lemma we see that  $(D, E)$  computes  $c$ . Hence the inequality at the beginning of this proof shows that  $\text{cliff}(D + E - (D, E)) \leq c$ . Since  $h^0(D + E - (D, E)) \geq 2$  it follows that  $|D + E - (D, E)|$  or its dual series computes  $c$  (depending on which of these two series has degree  $< g$ ) provided that  $h^1(D + E - (D, E)) \geq 2$ , too. But for  $h^1(D + E - (D, E)) \leq 1$  we obtain  $c \geq \text{cliff}(K_Y - (D + E - (D, E))) \geq 2g - 2 - (\text{deg}(D) + \text{deg}(E) - \text{deg}((D, E))) \geq \text{deg}((D, E))$  whence  $h^0((D, E)) \leq 1$ , a contradiction.  $\square$

From now on we use the following notation:  $X$  always denotes a curve of genus  $g$  whose Clifford index  $c$  is even and computed maximally. We set  $d_0 := g - 1 = 2c + 4$ ,  $r_0 := (d_0 - c)/2 = (c + 4)/2$ , and  $g_{d_0}^{r_0}$  is an arbitrary but fixed series on  $X$  (computing  $c$  maximally). Finally,  $I$  denotes the set of effective divisors  $D$  of  $X$  computing  $c$  such that  $\text{deg}(D) > c + 2$ . (Clearly,  $I \neq \emptyset$  since it contains the  $g_{d_0}^{r_0}$ .)

**Theorem 3.2.** *Assume that there is a divisor  $D \in I$  which is not contained in a divisor of the  $g_{d_0}^{r_0}$ . Then  $c \equiv 2 \pmod{4}$ ,  $D$  computes  $c$  maximally and  $W_{d_0}^{r_0}$  is infinite.*

Proof. For a divisor  $D \in I$  let  $d := \text{deg}(D)$ , and  $r := \dim(|D|) = (d - c)/2 \geq 2$ . Using a notation of [5], for any integer  $e \geq r - 1$  the set

$$V_e^{r-2}(|D|) := \{E \in \text{Div}(X) : E \geq 0, \text{deg}(E) = e \text{ and } \dim|D - E| \geq 1\}$$

is the variety of  $e$ -secant  $(r - 2)$ -plane divisors of  $X$ ; if  $V_e^{r-2}(|D|) \neq \emptyset$  every irreducible component  $Z$  of it has dimension  $\dim(Z) \geq 2(r - 1) - e$ . By [5, 1.2] we know that  $V_{2r-3}^{r-2}(|D|) \neq \emptyset$ , and for  $E \in V_{2r-3}^{r-2}(|D|)$  we have  $|D - E| \in W_{c+3}^1 = W_{c+2}^1 + W_1$ . Hence for every  $E \in V_{2r-3}^{r-2}(|D|)$  there is exactly one point  $P_E \in X$  such that  $E + P_E \in V_{2r-2}^{r-2}(|D|)$ . So the assignment  $E \mapsto E + P_E$  defines a surjection  $V_{2r-3}^{r-2}(|D|) \rightarrow V_{2r-2}^{r-2}(|D|)$  with finite fibres whence  $\dim V_{2r-2}^{r-2}(|D|) = \dim V_{2r-3}^{r-2}(|D|) \geq 2(r - 1) - (2r - 3) = 1$ . Let  $i : V_{2r-2}^{r-2}(|D|) \rightarrow W_{c+2}^1$  be the natural map defined by  $F \mapsto |D - F|$  for  $F \in V_{2r-2}^{r-2}(|D|)$ .

For any pencil  $L$  in the image of  $i$  there is a divisor  $F \in V_{2r-2}^{r-2}(|D|)$  resp. a pencil  $L'$  of degree  $c + 2$  on  $X$  such that  $|D| = |L + F|$  resp.  $g_{d_0}^{r_0} = |L + L'|$ , and for any point  $P$  in the

support of  $F$  we can find a divisor  $E' \in L'$  containing  $P$ . Hence for any  $E \in L$  the greatest common divisor  $G := (E + F, E + E')$  of  $E + F \in |D|$  and  $E + E' \in g_{d_0}^{r_0}$  contains the divisor  $E + P$ . So  $\deg(G) > \deg(E) = c + 2$ , and by Lemma 3.1 we know that  $\text{cliff}(G) \leq c$ . Since  $\dim|G| \geq \dim|E| = 1$  we see that  $G$  computes  $c$ , i.e.  $G \in I$ .

Now assume that  $D$  is not contained in a divisor of the  $g_{d_0}^{r_0}$ . Then  $G$  is properly contained in  $E + F \in |D|$ , and so  $\deg(G) < d$ . Thus the divisor  $H := (E + E') + (E + F) - G$  has degree  $g - 1 + d - \deg(G) \geq g$ , and, again by Lemma 3.1,  $|K_X - H|$  is a linear series of degree at most  $g - 2 = 2c + 3$  computing  $c$  which implies that  $\deg(K_X - H) \leq 3(c + 2)/2$ , i.e. we have  $2(c + 2) - d + \deg(G) = \deg(K_X - H) \leq 3(c + 2)/2$ . Hence  $\deg(G) \leq d - (c + 2)/2$ , and since  $\deg(G) > c + 2$  we obtain  $d > 3(c + 2)/2$ . It follows that  $d = 2c + 4 = g - 1$ , i.e.  $|D|$  is a  $g_{2c+4}^{(c+4)/2}$  on  $X$  different from our chosen  $g_{d_0}^{r_0}$ .

**CLAIM.** Assume that  $X$  has a linear series computing  $c$  maximally which is different from our  $g_{d_0}^{r_0}$ . Then  $W_{d_0}^{r_0}$  is infinite, and  $X$  has linear series of degree  $3(c + 2)/2$  computing  $c$ .

To prove this claim let  $h_{d_0}^{r_0}$  be a  $g_{2c+4}^{(c+4)/2}$  on  $X$  different from our  $g_{d_0}^{r_0}$ . For any  $L \in W_{c+2}^1$  there is a unique pair  $(L', L'')$  of different pencils  $L', L''$  of degree  $c + 2$  on  $X$  such that  $g_{d_0}^{r_0} = |L + L'|$  and  $h_{d_0}^{r_0} = |L + L''|$ . Let  $L = |E|$ .

Assume that  $L'$  and  $L''$  are not compounded of the same involution. Then the General Position Theorem ([1, 4.1]) implies that there is a divisor  $E' \in L'$  having with every divisor  $E'' \in L''$  at most one point in common, and for every point  $P$  in the support of  $E'$  we can find a divisor  $E'' \in L''$  containing  $P$ . With this choice we see, by Lemma 3.1, that  $G := (E + E', E + E'') = E + (E', E'') = E + P$  is a divisor computing  $c$  which is impossible since  $\deg(G) = c + 3$ .

Hence the two pencils  $L' = |g_{d_0}^{r_0} - L|, L'' = |h_{d_0}^{r_0} - L|$  are compounded of the same (irrational) involution. Then there is a covering  $\pi : X \rightarrow Y$  of maximum possible degree  $n$  such that  $L', L''$  are induced from pencils of degree  $(c + 2)/n$  on the curve  $Y$  (in particular,  $n$  divides  $c + 2$ ). For this pair  $(L', L'')$  specified by  $L = |E|$  we can choose, for any point  $P \in X$ , unique divisors  $E'_P \in L', E''_P \in L''$  having the point  $P$  in common. Then the greatest common divisor  $(E'_P, E''_P)$  of  $E'_P$  and  $E''_P$  is the divisor  $\pi^*(\pi(P))$  of degree  $n$  of  $X$ . (Clearly,  $\dim|(E'_P, E''_P)| = 0$ . Choosing  $E'_Q \in L', E''_Q \in L''$  having another point  $Q \in X$  in common we either have  $(E'_Q, E''_Q) = (E'_P, E''_P)$  - which happens only in the case  $\pi(Q) = \pi(P)$  - or that  $(E'_Q, E''_Q)$  and  $(E'_P, E''_P)$  have no point in common.) The divisor  $G_P := (E + E'_P, E + E''_P) = E + (E'_P, E''_P)$  has degree  $\deg(G_P) = c + 2 + n = ((\lambda + 1)/\lambda)(c + 2)$  if  $2 \leq \lambda := (c + 2)/n$ , and according to Lemma 3.1 it computes  $c$ . We will show that  $\lambda = 2$ , i.e.  $\deg(G_P) = 3(c + 2)/2$ ; then  $Y$  is an elliptic curve.

For  $m \geq 2$  points  $P_1, \dots, P_m$  of  $X$  such that  $(E'_{P_i}, E''_{P_i})$  and  $(E'_{P_j}, E''_{P_j})$  have disjoint support for  $1 \leq i < j \leq m$  we set  $G_{P_1, \dots, P_m} := E + (E'_{P_1}, E''_{P_1}) + \dots + (E'_{P_m}, E''_{P_m})$ . Then  $(G_{P_1, \dots, P_{m-1}}, G_{P_m}) = E$  computes  $c$ , and we have  $G_{P_1, \dots, P_m} = G_{P_1, \dots, P_{m-1}} + G_{P_m} - E = G_{P_1, \dots, P_{m-1}} + G_{P_m} - (G_{P_1, \dots, P_{m-1}}, G_{P_m})$ . Inductively applying Lemma 3.1 we see that  $G_{P_1, \dots, P_m}$  computes  $c$  as long as  $\deg(G_{P_1, \dots, P_m}) = c + 2 + mn = c + 2 + m(c + 2)/\lambda = (1 + (m/\lambda))(c + 2)$  is strictly smaller than  $g$ , i.e. for  $m \leq \lambda$ . If  $\lambda \geq 3$  we choose  $m = \lambda - 1$  and obtain that  $G_{P_1, \dots, P_{\lambda-1}}$  is a divisor computing  $c$  of degree strictly between  $3(c + 2)/2$  and  $2(c + 2)$ ; this is not possible. Hence we have  $\lambda = 2$ . Then we choose  $m = \lambda$  whence  $\deg(G_{P_1, P_2}) = 2c + 4 = d_0$ . Since, for  $Q \in X$ , we have  $G_{P_1, P_2} \sim G_{P_1, Q}$  iff  $(E'_{P_2}, E''_{P_2}) = (E'_Q, E''_Q)$  (i.e.  $\pi(P_2) = \pi(Q)$ ) we see

that - fixing  $P_1$  but varying  $P_2$  - we obtain this way infinitely many linear series on  $X$  which compute  $c$  maximally. This proves the claim.

Finally, we observe that  $3(c + 2)/2 = \deg(G_P) \equiv c \equiv 0 \pmod 2$  implies that  $c \equiv 2 \pmod 4$ . □

**Corollary 3.3.** *In the case  $c \equiv 0 \pmod 4$  the  $g_{d_0}^{r_0}$  is the only linear series on  $X$  computing  $c$  maximally.*

REMARK. Let  $V_e^n(g_{d_0}^{r_0}) := \{E \in \text{Div}(X) : E \geq 0, \deg(E) = e \text{ and } \dim(|g_{d_0}^{r_0} - E|) \geq r_0 - 1 - n\}$ ; here  $n \in \mathbb{Z}$  with  $n \leq e - 1$  and  $n \leq r_0 - 1$ . Choose an integer  $r$  such that  $1 < r < r_0$  and set  $d = c + 2r$  (note that  $d_0 - d = 2(r_0 - r)$ ). The upshot of the Theorem, then, is that  $V_{2(r_0-r)}^{r_0-1-r}(g_{d_0}^{r_0}) \cong W_d^r$  (via  $E \mapsto |g_{d_0}^{r_0} - E|$ ). For  $r = 1$  (i.e.  $d = c + 2$ ) this bijection is wrong since  $V_{2r_0-2}^{r_0-2}(g_{d_0}^{r_0})$  is the set of all effective divisors of degree  $2r_0 - 2 = c + 2$  of  $X$  which move in a non-trivial linear series, i.e.  $V_{2r_0-2}^{r_0-2}(g_{d_0}^{r_0}) = \{0 \leq E \in \text{Div}(X) : |E| = g_{c+2}^1\}$ ; so  $V_{2r_0-2}^{r_0-2}(g_{d_0}^{r_0})$  is a  $\mathbb{P}^1$ -bundle over  $W_{c+2}^1$ .

The Theorem thus relates the question if  $W_d^r \neq \emptyset$  ( $1 < r < r_0$ ) to the existence of a  $2(r_0 - r)$ -secant  $(r_0 - 1 - r)$ -plane for the curve  $X$  viewed as imbedded into  $\mathbb{P}^{r_0}$  by the  $g_{d_0}^{r_0}$ . And for  $2(c + 2) > d > 3(c + 2)/2$  (i.e. for  $0 < r_0 - r < (c + 2)/4$ ) we know that there is no such plane.

**Corollary 3.4.** *Assume that there exists a divisor  $D \in I$  of degree  $d < g - 1$ . Then  $W_{c+2}^1$  contains a one-dimensional irreducible component  $W$  such that for every pencil  $L \in W$  we have  $\dim |D - L| = 0$ , and the unique divisor in  $|D - L|$  is contained in a divisor of the pencil  $|g_{d_0}^{r_0} - L|$  of degree  $c + 2$ .*

Proof. We use the notation from the proof of the Theorem. Let  $r := \dim(|D|)$  and  $i|_Z : Z \rightarrow W_{c+2}^1$  be the natural map from an irreducible component  $Z$  of  $V_{2r-2}^{r-2}(|D|)$  into  $W_{c+2}^1$ ; recall that  $\dim(Z) \geq 1$ . Since there is no pencil of degree  $2r - 2 = d - c - 2 < c + 2$  on  $X$  the map  $i$  is injective whence we have  $\dim(i(Z)) \geq 1$ . But since  $\dim(W_{c+2}^1) \leq 1$  ([2, VII, ex. C-2]) it follows that  $\dim(i(Z)) = 1 = \dim(Z)$ . (In particular,  $V_{2r-2}^{r-2}(|D|)$  is equi-dimensional of dimension 1.)

Let  $W := i(Z)$ . Then  $W$  is an infinite irreducible component of  $W_{c+2}^1$ , and for every  $L \in W$  there is a divisor  $F \in Z$  such that  $|D| = |L + F|$ . Since  $\deg(F) = 2r - 2 = d - (c + 2) < c + 2$  we have  $|D - L| = \{F\}$ , and, by the Theorem,  $F$  is contained in a divisor of the pencil  $|g_{d_0}^{r_0} - L|$ . □

Recall that  $D \in I$ ,  $\deg(D) < g - 1 = 2c + 4$  implies that  $\deg(D) \leq 3(c + 2)/2$ , and for  $c \equiv 0 \pmod 4$  we even have  $d < 3(c + 2)/2$  since  $d \equiv c \equiv 0 \pmod 2$ . We add the following observation.

**Corollary 3.5.** *In Corollary 3.4, if  $d < 3(c + 2)/2$  then  $W_{c+2}^1$  contains a one-dimensional irreducible component (namely  $g_{d_0}^{r_0} - W$ ) such that no two different pencils in it are compounded of the same involution.*

Proof. In Corollary 3.4 we have  $|g_{d_0}^{r_0} - D| \subset |g_{d_0}^{r_0} - L|$  for any  $L \in W$ . Setting  $d = \deg(D)$  we clearly have  $\deg(|g_{d_0}^{r_0} - D|) = d_0 - d$ , and we know that  $(c + 2)/2 = 2(c + 2) - 3(c + 2)/2 \leq d_0 - d \leq (2c + 4) - (c + 4) = c$ . In particular,  $|g_{d_0}^{r_0} - D|$  consists of a single divisor  $E \geq 0$ .

Assume that two pencils  $L' \neq L''$  in  $g_{d_0}^{r_0} - W$  are compounded of the same involution thus giving rise to a covering  $\pi : X \rightarrow Y$  of degree  $n \geq 2$  such that  $L', L''$  are induced from pencils of degree  $(c + 2)/n$  on the curve  $Y$ . We can choose divisors  $E' \in L', E'' \in L''$  whose greatest common divisor  $(E', E'')$  contains  $E$ . We may assume that  $n = \deg((E', E''))$ ; then  $n \geq \deg(E) \geq (c + 2)/2$ , and so we obtain  $n = (c + 2)/2 = \deg(E)$ . Thus  $d = 3(c + 2)/2$ ;  $Y$  is an elliptic curve, then, and  $g_{d_0}^{r_0} - W = \pi^*(W_2^1(Y))$ . However, for  $d < 3(c + 2)/2$  this does not occur. □

We see that the divisor  $D \in I$  in Corollary 3.5 endows  $X$  with a feature of its pencils of minimal degree which - observing that their Brill-Noether number is negative - is apparently only known to be shared by the smooth plane curves (of degree  $\geq 6$ ). Cf. Remark 3.8 in [6].

**Corollary 3.6.** *For integers  $d, r$  such that  $c + 2 \leq d \leq g - 1$  and  $d - 2r = c$  we have  $\dim(W_d^r) \leq 1$ .*

Proof. We have  $\dim(W_{c+2}^1) \leq 1$  ([2, VII, ex. C-2]), and since  $W_{d_0}^{r_0} \subset g_{c+2}^1 + W_{c+2}^1$  for a fixed pencil  $g_{c+2}^1$  on  $X$  it follows that  $\dim(W_{d_0}^{r_0}) \leq 1$ . So we assume that  $c + 2 < d < d_0 = g - 1$ . Let  $K$  be an irreducible component of maximal dimension of  $W_d^r$ . Then  $\bigcup_{g_d^r \in K} i(V_{2r-2}^{r-2}(g_d^r)) \subset W_{c+2}^1$  is a union of one-dimensional irreducible components  $W_1, \dots, W_n$  of  $W_{c+2}^1$ . If  $K_j := \{g_d^r \in K \mid i(V_{2r-2}^{r-2}(g_d^r)) \supset W_j\}$  ( $j = 1, \dots, n$ ) we thus have  $K = K_1 \cup \dots \cup K_n$ . Fixing  $L_j \in W_j$  we have, by Corollary 3.4, a map  $\gamma_j : K_j \rightarrow \mathbb{P}^1$  which assigns to  $g_d^r \in K_j$  that divisor of the pencil  $|g_{d_0}^{r_0} - L_j|$  which contains the (unique) divisor  $E = |g_{d_0}^{r_0} - g_d^r|$ . Since  $E$  specifies  $g_d^r$  (and since the divisor  $\gamma_j(g_d^r)$  of degree  $c + 2$  contains only a finite number of effective divisors of degree  $d_0 - d \leq c$ ) the fibres of  $\gamma_j$  are finite. Choosing  $j$  such that  $\dim(K_j) = \dim(K) = \dim(W_d^r)$  it follows that  $\dim(W_d^r) \leq \dim(\mathbb{P}^1) = 1$ . □

**Corollary 3.7.** *If the  $g_{d_0}^{r_0}$  on  $X$  is not unique then every pencil of degree  $c + 2$  on  $X$  is induced by a pencil of degree 2 on a smooth elliptic curve (which is covered by  $X$  with  $(c + 2)/2$  sheets), and  $I$  consists of divisors of degree  $3(c + 2)/2$  and  $2(c + 2) = d_0$ .*

Proof. Let  $L \in W_{c+2}^1$ . There are pencils  $L', L'' \in W_{c+2}^1$  with  $L'' \neq L$  such that  $\dim(|L' + L|) = r_0 = \dim(|L' + L''|)$ , and from the proof of the Claim in the proof of the Theorem we see that  $L$  and  $L''$  are compounded of the same elliptic involution of order  $(c + 2)/2$ . The remaining assertion follows from Corollary 3.5. □

**Lemma 3.8.**  *$X$  has no net computing  $c$  if  $c > 8$ .*

Proof. Assume that  $X$  has a net  $g_{c+4}^2$ . Then for every point  $P \in X$  the pencil  $g_{c+4}^2(-P)$  of degree  $c + 3$  has a base point since  $W_{c+3}^1 = W_{c+2}^1 + W_1$ . Hence the  $g_{c+4}^2$  is not simple. Then it induces a morphism  $X \rightarrow Y$  of degree  $m > 1$  upon an integral plane curve  $Y$  of degree  $(c + 4)/m$ . If  $m > 2$  or if  $Y$  has singularities the normalization of  $Y$  has a pencil of degree  $d < (c + 2)/m$  which induces a pencil of degree  $md < c + 2$  on  $X$  which cannot exist. Hence  $m = 2$  and  $Y$  is a smooth plane curve of degree  $(c + 4)/2$ . Then  $Y$  has genus  $g(Y) = (1/2)((c + 4)/2 - 1)((c + 4)/2 - 2) = c(c + 2)/8$ , and by the Riemann-Hurwitz genus formula for coverings we obtain  $2c + 5 = g \geq 2g(Y) - 1 = c(c + 2)/4 - 1$ , i.e.  $(c - 3)^2 \leq 33$  which implies  $c \leq 8$ . □



For  $c = 6$  and  $c = 8$  we don't know yet if  $X$  has no net computing  $c$ .

**4. Clifford index  $c = 6$  and  $c = 8$**

In this section we turn to the Question posed in the Introduction, for  $c = 6$  and  $c = 8$ . In these cases the series computing  $c$ , besides those computing  $c$  maximally, are at most pencils, nets and webs. First, we reduce to pencils and nets, by the

**Lemma 4.1.** *Let  $c = 6$  or  $c = 8$ . If  $X$  has a web computing  $c$  then it also has a net computing  $c$ .*

Proof. Assume that  $X$  has a  $g^3_{c+6}$ . Then this series is base point free and simple thus inducing a birational morphism onto an integral space curve  $X'$  of degree  $c + 6$ .

Let  $D \in g^3_{c+6}$ . The number  $\rho_2$  of conditions imposed on quadrics in  $\mathbb{P}^2$  by a general plane section of  $X'$  is at most  $h^0(2D) - h^0(D)$ , and from the proof of Corollary 1 in [13] we know that  $h^0(2D) \geq 4 \cdot 3 - 2 = 10$ , i.e.  $|2D| = g^r_{2c+12}$  with  $r \geq 9$ . If  $r \geq 10$  then  $X$  has a  $g^{10}_{24} = |K_X - g^2_8|$  for  $c = 6$  which is impossible resp.  $X$  has a  $g^{10}_{28} = |K_X - g^2_{12}|$  for  $c = 8$  in which case there is a net computing  $c = 8$  on  $X$ . So we may assume that  $r = 9$  whence  $\rho_2 \leq 10 - 4 = 6 = 2\dim(|D|)$ . By a lemma of Castelnuovo and Fano's extension of it ([3, 1.10 and 3.1]) this implies that  $X'$  lies on a surface  $S$  of degree at most 3 in  $\mathbb{P}^3$ . The proof of Corollary 1 in [13] shows that  $X' \subset \mathbb{P}^3$  cannot lie on a quadric; so  $S$  is a cubic surface.

The projection  $\pi : X' \rightarrow \mathbb{P}^2$  with center a smooth point of  $X'$  is birational onto its image  $Y$  since  $c + 5$  is a prime number for  $c = 6$  and  $c = 8$ . Hence  $Y$  is a plane curve of degree  $c + 5$  which cannot be smooth. Since  $X$  has no base point free  $g^1_{c+3}$  all singular points of  $Y$  are triple points (points of multiplicity 3). Thus the fibre of  $\pi$  at a singular point of  $Y$  consists of 3 points of  $X'$ . Consequently,  $X'$  has a quadriseccant line through every smooth point. Clearly, then, all these lines must lie on the cubic  $S$ ; since our  $g^3_{c+6}$  is complete this is only possible if  $S$  is an elliptic cone. The ruling of the cone makes  $X$  a 4-fold covering of an elliptic curve. In particular,  $X$  has infinitely many  $g^1_8$  which is impossible for  $c = 8$ . For  $c = 6$  we use Segre's formula for the arithmetic genus of a curve on an elliptic scroll whose ruling are  $n$ -secant lines for the curve,

$p_a(X') = (n-1)(\deg(X')-1-(1/2)n\deg(S))+n = 3(12-1-(1/2)\cdot 4\cdot 3)+4 = 19 > g = 17$ . So  $X'$  has at least one singular point; taking the projection  $X' \rightarrow \mathbb{P}^2$  with center this point we obtain a net of degree  $m \leq \deg(X') - 2 = c + 4 = 10$  on  $X$ . Since  $c = 6$  we must have  $m = 10$ , and so we are done. □

**Theorem 4.2.** *For  $c = 6$  and  $c = 8$  the  $g^r_{d_0}$  is the only non-pencil on  $X$  computing  $c$ .*

Proof. By Corollary 3.7, Lemma 4.1 for  $c = 6$  resp. Corollary 3.3 for  $c = 8$ , the  $g^r_{d_0}$  on  $X$  is unique (and so, in particular, half-canonical). By Lemma 4.1 it remains to show the non-existence of nets on  $X$  computing  $c$ . So assume there is a  $g^2_{c+4}$  on  $X$ . As in the proof of Lemma 3.8 we see that this net induces a double covering  $\pi : X \rightarrow Y$  over a smooth plane curve  $Y$  of degree  $(c + 4)/2$ . Let  $\sigma$  ( $\sigma^2 = id$ .) denote the unique automorphism of  $X/Y$ .

By Theorem 3.2 there is an effective divisor  $D_c$  of  $X$  of degree  $d_0 - (c + 4) = c$  such that  $g^2_{c+4} = |g^r_{d_0} - D_c|$ . Since the  $g^2_{c+4}$  is base point free the support of a general divisor  $D' \in g^2_{c+4}$  consists of pairwise different points (is "separable") and is disjoint to the support of  $D_c$ . Since all divisors in our  $g^2_{c+4}$  are of the form  $\pi^*(\delta)$  for a divisor  $\delta$  in the unique net  $g^2_{(c+4)/2}$

on  $Y$  the divisor  $D'$  (being separable) contains no ramification point of  $\pi$  and is  $\sigma$ -invariant (i.e.  $\sigma D' = D'$ ).

Let  $D_0 := D' + D_c$ . Then  $D_0 \in g_{d_0}^{r_0}$ . Since the  $g_{d_0}^{r_0}$  on  $X$  is unique we have  $\sigma(g_{d_0}^{r_0}) = g_{d_0}^{r_0}$ . In particular,  $D' + D_c = D_0 \sim \sigma D_0 = \sigma D' + \sigma D_c = D' + \sigma D_c$ , i.e.  $\sigma D_c \sim D_c$ . But  $\dim|D_c| = 0$ , and so it follows that  $\sigma D_c = D_c$  and, then,  $\sigma D_0 = D_0$ .

Let  $R_1, \dots, R_n$  be the ramification points of  $\pi$ ; then  $R := R_1 + \dots + R_n \in \text{Div}(X)$  is the ramification divisor of  $\pi$ , and we have  $n = 12$  for  $c = 6$ ,  $n = 4$  for  $c = 8$ . For a  $\sigma$ -invariant divisor  $D = \sum_{i=1}^n k_i R_i + \sum_j l_j (P_j + \sigma(P_j)) \in \text{Div}(X)$  with  $P_j \neq R_i$  for all  $i, j$  we define a divisor  $\pi_0 D$  of  $Y$  by  $\pi_0 D := \sum_{i=1}^n [k_i/2] \pi(R_i) + \sum_j l_j \pi(P_j) \in \text{Div}(Y)$ , and we let  $V_e(D) := \{f \in H^0(D) | f \circ \sigma = f\}$  resp.  $V_o(D) := \{f \in H^0(D) | f \circ \sigma = -f\}$  be the even resp. odd part of  $H^0(D)$ . Then  $\deg(\pi_0 D) \leq (1/2) \deg(D)$ , and we have equality here iff  $\pi^*(\pi_0 D) = D$ . Furthermore,  $V_e(D) \cong H^0(Y, \pi_0 D)$  (since  $f \in V_e(D)$  has a pole of even order at every ramification point  $R_i$  of  $\pi$ ), and  $H^0(D) = V_e(D) \oplus V_o(D)$ .

Let  $V_e := V_e(D_0)$ ,  $V_o := V_o(D_0)$ . Since  $H^0(Y, \pi_0 D') \cong V_e(D') \subset V_e$  we have  $\dim(V_e) \geq h^0(\pi_0 D') = 3$ . Furthermore,  $\dim(V_e) = h^0(\pi_0 D_0)$  with  $\deg(\pi_0 D_0) \leq d_0/2 = c+2 = 2\deg(Y) - 2$ . Since  $Y$  is a smooth plane curve it follows that  $h^0(\pi_0 D_0) \leq 4$ , and if  $h^0(\pi_0 D_0) = 4$  holds then  $\deg(\pi_0 D_0) = c+2$ . So we see that  $\dim(V_e) \leq 4$ , and if  $\dim(V_e) = 4$  then  $\pi^*(\pi_0 D_0) = D_0$ .

We first consider the case  $\dim(V_e) = 3$ , i.e.  $V_e = V_e(D')$ . Then  $\dim(V_o) = h^0(D_0) - 3 = (((c+4)/2) + 1) - 3 = c/2$ .

Let  $D_c \leq R$  (i.e.  $\pi_0 D_c = 0$ ); this is only possible for  $c = 6$ . By adjunction we have  $K_X \sim \pi^*(K_Y) + R \sim \pi^*(2\delta) + R \sim 2D' + R$  for a divisor  $\delta$  in the net  $g_{(c+4)/2}^2$  on  $Y$ , and since  $|D_0|$  is half-canonical we have  $K_X \sim 2D_0 = 2D' + 2D_c$ . Hence we have  $2D_c \sim R$ . For a suitable numbering of the ramification points  $R_1, \dots, R_{12}$  of  $\pi$  we thus have  $2(R_1 + \dots + R_6) \sim R_1 + \dots + R_6 + R_7 + \dots + R_{12}$ , i.e.  $R_1 + \dots + R_6 \sim R_7 + \dots + R_{12}$ . But  $X$  has no  $g_6^1$ ; hence it follows that  $R_1 + \dots + R_6 = R_7 + \dots + R_{12}$  which is not true.

So we have  $2R_i \leq D_c$  for some  $i$  or  $P + \sigma(P) \leq D_c$  for a non-ramification point  $P \in X$ . Let  $k_i \geq 2$  resp.  $l \geq 1$  be the multiplicity of  $R_i$  resp.  $P$  in  $D_c$ ; note that  $k_i$  is odd. Choose a basis  $f_1, \dots, f_{c/2}$  of  $V_o$  such that  $R_i$  resp.  $P$  is a pole of order  $k_i$  resp.  $l$  of these functions. Then there are  $a_1, \dots, a_{(c/2)-1} \in \mathbb{C}$  such that the functions  $g_j := f_{c/2} - a_j f_j \in V_o$  ( $j = 1, \dots, (c/2) - 1$ ) have a pole of order  $k_i - 2$  at  $R_i$  resp.  $l - 1$  at  $P$  (and  $\sigma(P)$ ). Then the vector space  $V_e \oplus \text{span}(g_1, \dots, g_{(c/2)-1})$  of dimension  $\dim(V_e) + ((c/2) - 1) = (c/2) + 2$  gives rise to a linear series on  $X$  of dimension  $(c/2) + 1$  and degree  $\deg(D') + 2((c/2) - 1) = 2c + 2$ . Since this series computes  $c$  we obtain a contradiction.

So we have  $\dim(V_e) = 4$ , i.e.  $h^0(\pi_0 D_0) = 4$ . Then  $\pi^*(\pi_0 D_0) = D_0$  whence ([12, p. 1797])

$$((c+4)/2) + 1 = h^0(X, D_0) = h^0(X, \pi^*(\pi_0 D_0)) = h^0(Y, \pi_0 D_0) + h^0(Y, \pi_0 D_0 - E)$$

for a divisor  $E$  of  $Y$  such that  $2E$  is linearly equivalent to the branch divisor  $\pi_*(R)$  of  $\pi$ .

Thus we obtain  $h^0(\pi_0 D_0 - E) = (c+6)/2 - 4 = (c-2)/2$ , i.e.  $h^0(\pi_0 D_0 - E) = 2$  for  $c = 6$  and  $h^0(\pi_0 D_0 - E) = 3$  for  $c = 8$ . But for  $c = 6$  we have  $\deg(E) = n/2 = 6$  and so  $\deg(\pi_0 D_0 - E) = (1/2) \deg(D_0) - \deg(E) = (c+2) - 6 = 2$ , i.e.  $|\pi_0 D_0 - E|$  is a  $g_2^1$  on  $Y$  which is impossible. Let  $c = 8$ . Then we have  $\deg(E) = n/2 = 2$  whence  $\deg(\pi_0 D_0 - E) = 8$ , i.e.  $|\pi_0 D_0 - E|$  is a  $g_8^2$  on  $Y$ . Let  $\delta$  be a divisor in the unique net  $g_8^2$  on  $Y$ . Then there are points  $p_1, p_2, q_1, q_2$  of  $Y$  such that  $\pi_0 D_0 \sim 2\delta - p_1 - p_2$  and  $\pi_0 D_0 - E \sim \delta + q_1 + q_2$ . (In

fact, it is well known that  $W_8^2(Y) = W_6^2(Y) + W_2(Y) = |\delta| + W_2(Y)$  for a smooth plane sextic  $Y$  whence  $W_{10}^3(Y) = |K_Y| - W_8^2(Y) = |3\delta| - (|\delta| + W_2(Y)) = |2\delta| - W_2(Y)$ . So we obtain  $\delta + q_1 + q_2 \sim \pi_0 D_0 - E \sim 2\delta - p_1 - p_2 - E$ , i.e.  $\delta - E \sim p_1 + p_2 + q_1 + q_2$  which implies that  $h^0(\delta - E) \geq 1$ . But we have  $3 = h^0(X, \pi^*(\delta)) = h^0(Y, \delta) + h^0(Y, \delta - E) = 3 + h^0(Y, \delta - E)$  which shows that  $h^0(Y, \delta - E) = 0$ , and this contradiction proves the Theorem.  $\square$

If a smooth curve in  $\mathbb{P}^5$  on a cone over a 4-gonal canonical curve of genus 5 is cut out there by a quadric hypersurface it has maximally computed Clifford index 6 and infinitely many  $g_8^1$ ; so Theorem 4.2 is, for  $c = 6$ , not merely a consequence of the recognition theorem stated in the Introduction.

**5.  $X$  on a K3 surface**

Viewing  $X$  as being embedded into  $\mathbb{P}^{r_0}$  by our  $g_{d_0}^{r_0}$  it possibly lies on a smooth projective K3 surface  $S$  of degree  $2r_0 - 2$  in  $\mathbb{P}^{r_0}$ . (In fact, the examples of curves with maximally computed Clifford index have been constructed in this way, cf. [5, 3.2.6, 3.2.7].) If so, observing that  $c < [(g - 1)/2] = c + 2$  there exists an effective divisor  $D$  of  $S$  such that its restriction  $D|_X$  to  $X$  computes  $c$  ([7]). Hence one may ask if it is possible to find an (unexpected)  $g_{c+2r}^r$  with  $1 < r < r_0$  on  $X \subset S$  with the aid of a suitable divisor of  $S$ . As a consequence of an interesting result of Knutsen for curves on a K3 surface ([11, 3.4]) we have the

**Theorem 5.1.** *Assume that  $X$  lies, as a curve of degree  $d_0$ , on a K3 surface  $S$  of degree  $2r_0 - 2$  in  $\mathbb{P}^{r_0}$ . Then for every complete linear series  $|D|$  of  $S$  without a base curve such that  $D|_X$  computes  $c$  we have  $\deg(D|_X) = 2c + 4$  or  $\deg(D|_X) = c + 2$ .*

Proof. Let  $H$  be a hyperplane section of  $S$ . We have  $H^2 = \deg(S) = 2r_0 - 2 = c + 2$ ,  $X^2 = 2g - 2 = 4c + 8$  and  $H \cdot X = d_0 = 2c + 4$ , i.e.  $(H \cdot X)^2 = 4(c + 2)^2 = H^2 X^2$  which implies, by the Hodge index theorem ([9, V, 1.9 and ex. 1.9]), that  $X \sim ((H \cdot X)/H^2)H = 2H$ . Since the canonical series of  $S$  is trivial we have  $h^0(H - X) = h^0(-H) = h^2(H) = 0$  and  $h^1(H - X) = h^1(X - H) = h^1(H) = 0$  ([15, 2.2]) whence by a standard exact sequence and by the Riemann-Roch theorem ([9, V, 1.6]) it follows that  $h^0(X, H|_X) = h^0(H) = 2 + (1/2)H^2 = r_0 + 1$ , i.e.  $|H|_X = g_{d_0}^{r_0}$ .

Let  $D$  be an effective divisor of  $S$  such that  $|D|$  has no base curve and  $D|_X$  computes  $c$ . Then  $D^2 \geq 0$ , and since  $\deg(D) = D \cdot H = (1/2)D \cdot X = (1/2) \deg(D|_X) < g - 1 = d_0 = \deg(X)$  we have  $h^0(D - X) = 0$ .

Assume that  $h^1(D) = 0$ . Then a standard exact sequence shows that  $h^0(X, D|_X) = h^0(D) + h^1(D - X)$ . Likewise, if  $X_0$  is an arbitrary smooth irreducible curve in  $|2H|$  we have  $h^0(X_0, D|_{X_0}) = h^0(D) + h^1(D - X_0)$ . Clearly,  $D - X_0 \sim D - X$  implies that  $h^1(D - X_0) = h^1(D - X)$  whence  $h^0(X_0, D|_{X_0}) = h^0(X, D|_X)$ . Since, by [7],  $X_0$  has the same Clifford index  $c$  as  $X$ , we see that  $D|_{X_0}$  computes the Clifford index of  $X_0$ .

Choose  $X_0$  general in  $|2H|$ . Then  $X_0$  has only finitely many pencils  $g_{c+2}^1$ , according to a theorem of Knutsen ([11, 3.4]), and since the Clifford index  $c$  of  $X_0$  is maximally computed (by  $H|_{X_0}$ ) there are no base point free  $g_{c+3}^1$  on  $X_0$ . Consequently, the recognition theorem (applied to  $X_0$ ) shows that  $D|_{X_0}$  computes  $c$  maximally or  $|D|_{X_0} = g_{c+2}^1$ . Hence, for  $X$ , we have  $h^0(X, D|_X) = r_0 + 1$  or (provided that  $D^2 = 0$ )  $h^0(X, D|_X) = 2$ .

Assume that  $h^1(D) \neq 0$ . Then  $D \sim kE_0$  for an irreducible curve  $E_0$  with  $E_0^2 = 0$  and some integer  $k \geq 2$  ([15, 2.6]). We have  $k \deg(E_0|_X) = \deg(D|_X) \leq g - 1 = 2c + 4$ , and since  $h^0(X, E_0|_X) \geq h^0(E_0) \geq 2 + (1/2)E_0^2 = 2$  we have  $\deg(E_0|_X) \geq c + 2$ . Thus we obtain  $k = 2$  and  $\deg(D|_X) = 2c + 4$ .  $\square$

---

### References

- [1] R.D.M. Accola: *On Castelnuovo's inequality for algebraic curves. I*, Trans. Amer. Math. Soc. **251** (1979), 355–373.
- [2] E. Arbarello, M. Cornalba, P. Griffiths and J. Harris: *Geometry of Algebraic Curves. I*, Grundlehren Math. Wiss., **267**, Springer, New York 1985.
- [3] C. Ciliberto: *Hilbert functions of finite sets of points and the genus of a curve in a projective space*; in *Lecture Notes in Mathematics* **1266**, Springer, New York, 1987, 24–73.
- [4] M. Coppens: *Smooth curves having infinitely many linear systems  $g_d^1$* , Bull. Math. Soc. Belg., Ser. B **40** (1988), 153–176.
- [5] M. Coppens and G. Martens: *Secant spaces and Clifford's theorem*, Compositio Math. **78** (1991), 193–212.
- [6] D. Eisenbud, H. Lange, G. Martens and F.-O. Schreyer: *The Clifford dimension of a projective curve*, Compositio Math. **72** (1989), 173–204.
- [7] M. Green and R. Lazarsfeld: *Special divisors on curves on a K3 surface*, Invent. Math. **89** (1987), 357–370.
- [8] F.J. Gallego and B.P. Purnaprajna: *Normal presentation on elliptic ruled surfaces*, J. Algebr **186** (1996), 597–625.
- [9] R. Hartshorne: *Algebraic Geometry*, Graduate Texts in Mathematics, **52**, Springer, New York 1977.
- [10] S. Kim: *On the algebraic curve of Clifford index  $c$* , Comm. Korean Math. Soc. **14** (1999), 693–698.
- [11] A.L. Knutsen: *On two conjectures for curves on K3 surfaces*, Int. J. Math. **20** (2009), 1547–1560.
- [12] C. Keem and G. Martens: *Curves without plane models of small degree*, Math. Nachr. **281** (2008), 1791–1798.
- [13] C. Keem, S. Kim and G. Martens: *On a result of Farkas*, J. Reine Angew. Math. **405** (1990), 112–116.
- [14] H.H. Martens: *Varieties of special divisors on a curve. II*, J. Reine Angew. Math. **233** (1968), 89–100.
- [15] B. Saint-Donat: *Projective models of K3 surfaces*, Amer. J. Math. **96** (1974), 602–639.

Takao Kato  
 13–1 Ogori-Ozaki-cho, Yamaguchi, 754–0006  
 Japan  
 e-mail: kato@yamaguchi-u.ac.jp

Gerriet Martens  
 Department Mathematik  
 Univ. Erlangen-Nürnberg  
 Cauerstr. 11, D–91058 Erlangen  
 Germany  
 e-mail: martens@mi.uni-erlangen.de