

## 2-STRATIFOLD SPINES OF CLOSED 3-MANIFOLDS

J.C. GÓMEZ-LARRAÑAGA, F. GONZÁLEZ-ACUÑA and WOLFGANG HEIL

(Received August 6, 2018, revised November 21, 2018)

### Abstract

2-stratifolds are a generalization of 2-manifolds in that there are disjoint simple closed branch curves. We obtain a list of all closed 3-manifolds that have a 2-stratifold as a spine.

### 1. Introduction

2-stratifolds form a special class of 2-dimensional stratified spaces. A (closed with empty 0-stratum) 2-stratifold is a compact connected 2-dimensional cell complex  $X$  that contains a 1-dimensional subcomplex  $X^{(1)}$ , consisting of branch curves, such that  $X - X^{(1)}$  is a (not necessarily connected) 2-manifold. The exact definition is given in section 2.  $X$  can be constructed from a disjoint union  $X^{(1)}$  of circles and compact 2-manifolds  $W^2$  by attaching each component of  $\partial W^2$  to  $X^{(1)}$  via a covering map  $\psi : \partial W^2 \rightarrow X^{(1)}$ , with  $\psi^{-1}(x) > 2$  for  $x \in X^{(1)}$ . A slightly more general class of 2-dimensional stratified spaces, called *multibranched surfaces* and which have been defined and studied in [13], is obtained by allowing boundary curves, i.e. considering a covering map  $\psi : \partial W' \rightarrow X^{(1)}$ , where  $\partial W'$  is a sub collection of the components of  $\partial W^2$ .

2-stratifolds arise as the nerve of certain decompositions of 3-manifolds into pieces where they determine whether the  $\mathcal{G}$ -category of the 3-manifold is 2 or 3 ([6]). They are related to *foams*, which include special spines of 3-dimensional manifolds and which have been studied by Khovanov [10] and Carter [4]. Simple 2-dimensional stratified spaces arise in Topological Data Analysis [2], [11].

Matsuzaki and Ozawa [13] show that 2-stratifolds can be embedded in  $\mathbb{R}^4$ . Furthermore they show that they can be embedded into some orientable closed 3-manifold if and only if their branch curves satisfy a certain regularity condition. However, the embeddings are not  $\pi_1$ -injective, i.e. the induced homomorphism of fundamental groups is not injective. In fact, there are many 2-stratifolds whose fundamental group is not isomorphic to a 3-manifold group; for example there are infinitely many 2-stratifolds with (Baumslag-Solitar) non-Hopfian fundamental groups. These can not be embedded as  $\pi_1$ -injective subcomplexes into 3-manifolds since 3-manifold groups are residually finite.

Further comparing properties of 2-stratifold groups with 3-manifold groups we note that a 2-stratifold group  $G$  is the fundamental group of a graph of groups where each edge group is cyclic and each vertex group is an  $F$ -group or a free product of cyclic groups. (This is described in detail in section 2). If  $G$  is torsion free then the vertex groups are surface groups. Since the latter (except for  $Z_2$ ) are left-orderable it follows from Corollary 3.6 of [5] that  $G$

is left-orderable. On the other hand, some torsion free 3-manifold groups are left orderable and some are not. For example it is shown in [3] that groups of compact  $P^2$ -irreducible 3-manifolds  $M$  with first Betti number  $> 0$  are left-orderable, but not all Haken manifolds have left orderable groups. Thus the following question arises:

QUESTION 1. Which 3-manifolds  $M$  have fundamental groups isomorphic to the fundamental group of a 2-stratifold?

The fundamental group of a closed 2-manifold  $S$  is isomorphic to the fundamental group of a closed 3-manifold  $M$  if and only if  $S$  is the 2-sphere or projective plane and  $M$  is  $S^3$  or  $P^3$ , respectively. Since  $S^2$  is not a spine of  $S^3$ , the only closed 3-manifold with a (closed) 2-manifold spine is  $P^3$ . This motivates the next question:

QUESTION 2. Which closed 3-manifolds  $M$  have spines that are 2-stratifolds?

The main results of this paper are Theorem 1 which answers question 1 for closed 3-manifolds and Theorem 2, which answers question 2 by showing that a closed 3-manifold  $M$  has a 2-stratifold spine if and only if  $M$  is a connected sum of lens spaces,  $S^2$ -bundles over  $S^1$ , and  $P^2 \times S^1$ 's.

## 2. 2-stratifolds and their graphs.

In this section we review the definitions of a 2-stratifold  $X$  and its associated graph  $G_X$  given in [7].

A (closed) 2-stratifold is a compact 2-dimensional cell complex  $X$  that contains a 1-dimensional subcomplex  $X^{(1)}$ , such that  $X - X^{(1)}$  is a 2-manifold ( $X^{(1)}$  and  $X - X^{(1)}$  need not be connected). A component  $C \approx S^1$  of  $X^1$  has a regular neighborhood  $N(C) = N_\pi(C)$  that is homeomorphic to  $(Y \times [0, 1]) / (y, 1) \sim (h(y), 0)$ , where  $Y$  is the closed cone on the discrete space  $\{1, 2, \dots, d\}$  (for  $d \geq 3$ ) and  $h : Y \rightarrow Y$  is a homeomorphism whose restriction to  $\{1, 2, \dots, d\}$  is the permutation  $\pi : \{1, 2, \dots, d\} \rightarrow \{1, 2, \dots, d\}$ . The space  $N_\pi(C)$  depends only on the conjugacy class of  $\pi \in S_d$  and therefore is determined by a partition of  $d$ . A component of  $\partial N_\pi(C)$  corresponds then to a summand of the partition determined by  $\pi$ . Here the neighborhoods  $N(C)$  are chosen sufficiently small so that for disjoint components  $C$  and  $C'$  of  $X_1$ ,  $N(C)$  is disjoint from  $N(C')$ .

Note that  $X$  may also be described as a quotient space  $W \cup_\psi X^{(1)}$ , where  $\psi : \partial W \rightarrow X^{(1)}$  is a covering map (and  $|\psi^{-1}(x)| > 2$  for every  $x \in X^{(1)}$ ).

We construct an associated bicolored graph  $G = G_X$  of  $X = X_G$  by letting the white vertices  $w$  of  $G_X$  be the components  $W$  of  $M := \overline{X - \cup_j N(C_j)}$  where  $C_j$  runs over the components of  $X^1$ ; the black vertices  $b_j$  are the  $C_j$ 's. An edge  $e$  is a component  $S$  of  $\partial M$ ; it joins a white vertex  $w$  corresponding to  $W$  with a black vertex  $b$  corresponding to  $C_j$  if  $S = W \cap N(C_j)$ . The number of boundary components of  $W$  is the number of adjacent edges of  $W$ .  $G_X$  embeds naturally as a retract into  $X_G$ .

We label the white vertices  $w$  with the genus  $g$  of  $W$ ; here we use Neumann's [16] convention of assigning negative genus  $g$  to nonorientable surfaces; for example the genus  $g$  of the projective plane or the Moebius band is  $-1$ , the genus of the Klein bottle is  $-2$ . We orient all components  $C_j$  and  $S$  of  $X^{(1)}$  and  $\partial W$ , resp., and assign a label  $m$  to an edge  $e$ , where  $|m|$

is the summand of the partition  $\pi$  corresponding to the component  $S \subset \partial N_\pi(C)$ ; the sign of  $m$  is determined by the orientation of  $C_j$  and  $S$ . In terms of attaching maps,  $m$  is the degree of the covering map  $\psi : S \rightarrow C_j$  for the corresponding components of  $\partial W$  and  $X^{(1)}$ .

(Note that the partition  $\pi$  of a black vertex is determined by the labels of its adjacent edges).

### 3. Structure of $\pi_1(X_G)$

In this section we obtain a natural presentation for the fundamental group of a 2-stratifold  $X_G$  with associated bicolored graph  $G = G_X$  and describe  $\pi_1(X_G)$  as the fundamental group of a graph of groups  $\mathcal{G}$  with the same underlying graph  $G$ .

For a given white vertex  $w$ , the compact 2-manifold  $W$  has conveniently oriented boundary curves  $s_1, \dots, s_p$  such that

$$(*) \quad \pi_1(W) = \langle s_1, \dots, s_p, y_1, \dots, y_n : s_1 \cdots s_p \cdot q = 1 \rangle$$

where  $q = [y_1, y_2] \dots [y_{2g-1}, y_{2g}]$ , if  $W$  is orientable of genus  $g$  and  $n = 2g$ ,  $q = y_1^2 \dots y_n^2$ , if  $W$  is non-orientable of genus  $-n$ .

Let  $\mathcal{B}$  be the set of black vertices,  $\mathcal{W}$  the set of white vertices and choose a fixed maximal tree  $T$  of  $G$ . Choose orientations of the black vertices and of all boundary components of  $M$  such that all labels of edges in  $T$  are positive.

Then  $\pi_1(X_G)$  has a natural presentation with *generators*:

$$\{b\}_{b \in \mathcal{B}}$$

$\{s_1, \dots, s_p, y_1, \dots, y_n\}$ , one set for each  $w \in \mathcal{W}$ , as in (\*)

$\{t_i\}$ , one  $t_i$  for each edge  $c_i \in G - T$  between  $w$  and  $b$

and *relations*:

$s_1 \cdots s_p \cdot q = 1$ , one for each  $w \in \mathcal{W}$ , as in (\*)

$b^m = s_i$ , for each edge  $s_i \in T$  between  $w$  and  $b$  with label  $m \geq 1$

$t_i^{-1} s_i t_i = b^{m_i}$ , for each edge  $s_i \in G - T$  between  $w$  and  $b$  with label  $m_i \in \mathbb{Z}$ .

As an example we show in Figure 1 (the graph of) a 2-stratifold  $X_G$  with  $\pi_1(X_G) = \mathcal{F}$ , an  $F$ -group as in Proposition (III)5.3 of [12], with presentation

$$(\mathcal{F}) \quad \mathcal{F} = \langle c_1, \dots, c_p, y_1, \dots, y_n : c_1^{m_1}, \dots, c_p^{m_p}, c_1 \cdots c_p \cdot q = 1 \rangle$$

where  $p, n \geq 0$ , all  $m_i > 1$  and  $q = [y_1, y_2] \dots [y_{2g-1}, y_{2g}]$  or  $q = y_1^2 \dots y_n^2$ .

Here we have denoted the generators corresponding to the black vertices by  $c_i$ , rather than  $b_i$ , to indicate that the finite order elements correspond to attaching disks along the boundary curves of  $W$ .

The fundamental group of  $X_G$  is best described as the fundamental group of a graph of




Fig. 1. F-group

groups [8].

If  $\pi_1(X_G)$  has no elements of finite order, then  $\pi_1(X_G)$  is the fundamental group of a graph of groups  $\mathcal{G}$ , with underlying graph  $G$ , the groups of white vertices are the fundamental groups of the  $W$ 's, the groups of the black vertices and edges are (infinite) cyclic.

Elements of finite order occur when a generator  $b$  of a black vertex has finite order  $o(b) \geq 1$ . In this case we attach 2-cells  $d_b$  and  $d_e$  to  $C_b$ , the circle corresponding to  $b$ , as follows:  $d_b$  is attached by a map of degree  $o(b)$ . If  $e$  is an edge joining  $b$  to  $w$  with label  $m$ , attach  $d_e$  with degree  $o(c) = o(b)/(o(b), m)$ . Letting  $\hat{X}_b = N(C_b) \cup d_b \cup (\cup d_e)$ , where  $e$  runs over the edges having  $b$  as an endpoint,  $\hat{X}_w = W \cup (\cup d_e)$ , where  $e$  runs over the edges incident to  $w$ , and  $\hat{X}_e = (\hat{X}_b \cap \hat{X}_w)$ , for an edge  $e$  joining  $b$  to  $w$ , we obtain a graph of CW-complexes that determines a graph of groups  $\mathcal{G}$  with the same underlying graph as  $G_X$ .

The vertex groups are  $G_b = \pi_1(\hat{X}_b)$  and  $G_w = \pi_1(\hat{X}_w)$ , the edge groups are  $G_e = \pi_1(\hat{X}_e)$ , the monomorphisms  $\delta : G_e \rightarrow G_b$  (resp.  $G_e \rightarrow G_w$ ) are induced by inclusion. Then (see for example [17],[18])  $\pi_1 \mathcal{G} \cong \pi_1(\hat{X})$ .

Note that the groups  $G_b$  of the black vertices and the groups  $G_e$  of the edges are cyclic. For a white vertex  $w$  with edges  $e_1, \dots, e_p$  labelled  $m_1, \dots, m_p$  with associated vertex space  $X_w = W \cup_{i=1}^r d_{e_i}$  we obtain

$$G_w = \langle c_1, \dots, c_p, y_1, \dots, y_n : c_1 \cdots c_p \cdot q = 1, c_1^{m_1} = \dots = c_r^{m_r} = 1 \rangle$$

where  $q$  is as in (F),  $1 \leq r \leq p$  and  $k_i \geq 1$ .

If all  $k_i \geq 2$  and  $r = p$  then  $G_w$  is an  $F$ -group ([12] p. 126-127). If  $r < p$  it is a free product of cyclic groups.

#### 4. Necessary Conditions

In this section we show that a 2-stratifold group that is a closed 3-manifold group is a free product of cyclic or  $\mathbb{Z} \times \mathbb{Z}_2$  groups.

First consider an  $F$ -group  $\mathcal{F}$  as in (F).

**Proposition 1** ([12] Proposition (III)7.4). *Let  $H$  be a subgroup of an  $F$ -group. If  $H$  has finite index then  $H$  is an  $F$ -group. If  $H$  has infinite index then  $H$  is a free product of cyclic groups.*

**Proposition 2** ([12] p.132). (a)  $\mathcal{F}$  is finite non-cyclic if and only if  $n = 0$ ,  $p = 3$  and  $(m_1, m_2, m_3) = (2, 2, m)$  ( $m \geq 2$ ) (the dihedral group of order  $2m$ ) or  $(m_1, m_2, m_3) = (2, 3, k)$

for  $k = 3, 4$  or  $5$  (the tetrahedral, octahedral, dodecahedral groups). In each case,  $c_1$  is a non-central element of order 2.

(b)  $\mathcal{F}$  is finite cyclic if and only if  $n = 0, p \leq 2$  (the 2-sphere orbifold with at most two cone points) or  $n = 1, p \leq 1$  (the projective plane orbifold with at most one cone point).

**Lemma 1.**  $\mathcal{F}$  is not a non-trivial free product.

Proof. If  $\mathcal{F} = A * B$  with  $A, B$  non-trivial, then  $A$  and  $B$  have infinite index and so, by Proposition 1,  $A, B$  and  $\mathcal{F}$  are free products of cyclic groups. However,  $\mathcal{F}$  is not such a group since it contains a subgroup isomorphic to the fundamental group of an orientable closed surface of genus  $\geq 1$  (see the remark after Proposition (III)7.12 in [12]).  $\square$

The following remark is easy to see.

REMARK 1. If  $\mathcal{F} \neq \mathbb{Z}_2$  then  $\mathcal{F}$  has no elements of finite order if and only if  $\mathcal{F}$  is a surface group.

**Lemma 2.** If  $M$  is an orientable (not necessarily closed or compact) 3-manifold with  $\pi_1(M) \cong \mathcal{F}$  then  $\pi_1(M)$  is cyclic or a surface group.

Proof. We may assume that  $\partial M$  contains no 2-spheres. By Scott's Core Theorem we may assume that  $M$  is compact and by Lemma 1 that  $M$  is irreducible.

If  $\pi_1(M)$  is infinite then  $M$  is aspherical (see e.g. [1]). It follows that  $\pi_1(M)$  is torsion-free and from Remark 1 that  $\pi_1(M)$  is a surface group.

If  $\pi_1(M)$  is finite then  $M$  is closed. If  $\pi_1(M)$  is also non-cyclic then by Proposition 2,  $\pi_1(M)$  contains a non-central element of order 2. This can not happen by Milnor [15].  $\square$

We now consider a 2-stratifold  $X_G$  with  $\pi_1(X_G) = \pi(\mathcal{G})$  as in section 3.

Up to conjugacy, the only elements of finite order of  $\pi_1(X_G)$  are contained in the vertex groups; they correspond to black vertices of finite order and elements of white vertices  $w$  whose corresponding group in  $\mathcal{G}$  is finite. The latter are described in Proposition 2. It is also shown in [12] (proof of Proposition (III)7.12) that in an infinite F-group the only elements of finite order are the obvious ones, namely conjugates of powers of  $c_1, \dots, c_p$ .

For a group  $H$ , denote by  $QH$  be the quotient group of  $H$  modulo the smallest subgroup of  $H$  containing all elements of finite order of  $H$ .

Let  $w$  be a white vertex in  $G_X$ . We say that  $w$  is a *white hole*, if  $w$  has label  $-1$ , all of its (black) neighbors have finite order and at most one of its neighbors has order  $> 1$ .

If  $G_X$  has more than one vertex, note that  $Q\pi_1(X_G)$  is obtained from  $\pi_1(X_G)$  by killing the open stars of all the black vertices representing elements of finite order  $\geq 1$  of  $\pi_1(X_G)$  and deleting the white holes. In the example of Figure 1, when genus  $g = -1$  (and so  $n = 1$ ),  $Q\pi_1(X_G) = \mathbb{Z}_2$ . (Note that the white vertex of genus  $-1$  is not a white hole if  $p \geq 2, m_i > 1$ ).

**Proposition 3.** If  $Q(\pi_1(X_G))$  has no elements of order 2, then  $H_3(Q\pi_1(X_G)) = 0$ .

Proof. Let  $G'$  be the labelled subgraph of  $G_X$  obtained by deleting the open stars of all black vertices representing elements of finite order of  $\pi_1(X_G)$  and all white holes. ( $\pi_1(X_0) = 1$  by definition). Let  $C$  be a component of  $G'$ . Then  $Q\pi(X_G) = L * (*_C(\pi(X_C)))$ , the free product of a free group  $L$  with the free product of the  $\pi(X_C)$  where  $C$  runs over the components of  $G'$ .

If  $C$  consists of only one (white) vertex, then  $X_C$  is a closed 2-manifold, different from  $P^2$ , since by assumption  $Q(\pi_1(X_G))$  has no elements of order 2. We may ignore the  $C$ 's consisting of spheres, since they do not contribute to  $Q\pi(X_G)$ . (A nonseparating 2-sphere only changes the rank of  $L$ ). In all other cases  $X_C$  is the total space of a bicolored graph of spaces with white vertex spaces 2-manifolds with boundary, edge spaces circles, and black vertex spaces homotopy equivalent to circles.

Thus every vertex and edge space of  $X_C$  is aspherical (with free fundamental group) of dimension  $\leq 2$ . By Proposition 3.6 (ii) of [17],  $X_C$  is aspherical. It follows that  $Q\pi_1(X_G)$  has (co)homological dimension  $\leq 2$  and so  $H_3(Q\pi_1(X_G)) = 0$ .  $\square$

The assumption that  $Q(\pi_1(X_G))$  has no elements of finite order is satisfied if  $\pi_1(X_G)$  is a 3-manifold group: We claim that  $Q\pi_1(M)$  is torsion free if  $M$  is a closed orientable 3-manifold.

For let  $M = M_1 \# \dots \# M_k$  be its prime decomposition. If  $M_i$  is irreducible with infinite fundamental group, then  $M_i$  is aspherical and so  $\pi_1(M_i)$  is torsion free; if  $M_i$  has finite fundamental group, then  $Q\pi_1(M_i) = 1$ . Now the claim follows since  $Q\pi_1(M) = Q\pi_1(M_1) * \dots * Q\pi_1(M_k)$ .

**Lemma 3.** *Let  $M$  be a closed orientable 3-manifold with prime decomposition  $M = M_1 \# \dots \# M_k$ . If  $\pi_1(M) \cong \pi_1(X_G)$ , then each  $\pi_1(M_i)$  is infinite cyclic or finite.*

Proof. If there is some  $M_i$  with  $\pi_1(M_i) \neq \mathbb{Z}$ , then  $M_i$  is irreducible. If  $\pi_1(M_i)$  is infinite then  $M_i$  is aspherical and hence  $H_3(Q\pi_1(M_i)) = H_3(\pi_1(M_i)) = H_3(M_i) \neq 0$ . Since  $Q\pi_1(M) = Q\pi_1(M_1) * \dots * Q\pi_1(M_k)$  it follows that  $H_3(Q\pi_1(M)) \neq 0$ , which contradicts Proposition 3.  $\square$

REMARK 2. One can give an alternate proof of Lemma 3 by using the fact that 3-manifold groups are virtually torsion free. Using this fact one can show (by looking at torsion-free subgroups instead of torsion-free quotients) that if  $G$  is both a 2-stratifold group and a 3-manifold group, then the virtual cohomological dimension of  $G$  is at most 2. By a slight modification of the above proof of Lemma 3 this implies that no connected sum summand of a closed orientable 3-manifold  $M$  is aspherical.

**Lemma 4.** *Let  $M$  be a closed 3-manifold and suppose  $\pi_1(M) = \pi_1(X_G)$ . Then any finite subgroup  $H$  of  $\pi_1(M)$  is cyclic.*

Proof.  $\pi_1(X_G) \cong \pi_1(\mathcal{G})$  where  $\mathcal{G}$  is a graph of groups in which the groups of black vertices are cyclic and the groups of white vertices are  $F$ -groups or free products of finitely many cyclic groups. The finite group  $H$  is non-splittable (i.e. not a non-trivial HHN extension or free product with amalgamation). By Corollary 3.8 and the Remark after Theorem 3.7 of [17],  $H$  is a cyclic group or isomorphic to a subgroup of an  $F$ -group. If  $H$  is not cyclic, then (since  $H$  is not a non-trivial free product of cyclic groups),  $H$  is itself an  $F$ -group by

Proposition 1. Since  $H$  is a 3-manifold group it follows from Lemma 2 that this case can not occur. □

**Corollary 1.** *Let  $M$  be a closed orientable 3-manifold. If  $\pi_1(M) \cong \pi_1(X_G)$ , then  $\pi_1(M)$  is a free product of cyclic groups.*

Proof. This follows from Lemmas 3 and 4. □

**Theorem 1.** *Let  $M$  be a closed 3-manifold. If  $\pi_1(M) \cong \pi_1(X_G)$ , then  $\pi_1(M)$  is a free product of groups, where each factor is cyclic or  $\mathbb{Z} \times \mathbb{Z}_2$ .*

Proof. If  $M$  is orientable this is Corollary 1 (with each factor cyclic). Thus assume  $M$  is non-orientable and let  $p : \tilde{M} \rightarrow M$  be the 2-fold orientable cover of  $M$ . Then  $\pi_1(\tilde{M}) = \pi_1(X_{\tilde{G}})$  for the 2-stratifold  $X_{\tilde{G}}$ , which is the 2-fold cover of  $X_G$  corresponding to the orientation subgroup of  $\pi_1(M)$ . Hence  $\pi_1(\tilde{M})$  is a free product of cyclic groups.

Let  $M = \hat{M}_1 \# \dots \# \hat{M}_k$  be a prime decomposition of  $M = M_1 \cup \dots \cup M_k$ . If  $M_i$  is orientable, then  $M_i$  lifts to two homeomorphic copies  $\tilde{M}_{i1}, \tilde{M}_{i2}$  of  $M_i$ , with each  $\hat{M}_{ij}$  a factor of the prime decomposition of  $\tilde{M}$  and it follows that  $\pi_1(M_i)$  is cyclic.

If  $\hat{M}_i$  is non-orientable and  $P^2$ -irreducible, then  $M_i$  lifts to  $\tilde{M}_i$ , where  $\hat{M}_i$  is irreducible. Then  $\pi_1(\hat{M}_i)$ , being a factor of the free product decomposition of  $\pi_1(\tilde{M})$ , is finite cyclic, which can not occur since  $\pi_1(M_i)$  is infinite.

If  $\hat{M}_i$  is non-orientable irreducible, contains  $P^2$ 's, but is not  $P^2 \times S^1$ , then by Proposition (2.2) of [19],  $M_i$  splits along two-sided  $P^2$ 's into 3-manifolds  $N_1, \dots, N_m$  such that the fundamental group of the lifts  $\tilde{N}_i$  is indecomposable, torsion free and not isomorphic to  $\mathbb{Z}$ . Since  $\pi_1(\tilde{N}_i)$  is a factor of the free product decomposition of  $\pi_1(\tilde{M})$ , this can not happen.

Therefore each non-orientable  $M_i$  is either the  $S^2$ -bundle over  $S^1$  or  $P^2 \times S^1$ , which proves the Theorem. □

### 5. Realizations of spines.

Recall that a subpolyhedron  $P$  of a 3-manifold  $M$  is a *spine* of  $M$ , if  $M - \text{Int}(B^3)$  collapses to  $P$ , where  $B^3$  is a 3-ball in  $M$ .

An equivalent definition is that  $M - P$  is homeomorphic to an open 3-ball (Theorem 1.1.7 of [14]).

We first construct 2-stratifold spines of lens spaces (different from  $S^3$ ), the non-orientable  $S^2$ -bundle over  $S^1$ , and  $P^2 \times S^1$ .

EXAMPLE 1. Lens space  $L(0, 1) = S^3$ .

$S^3$  does not have a 2-stratifold spine. Otherwise such a spine  $X$  would be a deformation retract of the 3-ball and therefore contractible. However there are no contractible 2-stratifolds [7].

EXAMPLE 2. Lens spaces  $L(p, q)$  with  $q \neq 0, 1$ .

Let  $r$  be the rotation of the disk  $D^2$  about its center  $c$  with angle  $2\pi p/q$ , let  $1 \in S^1 \subset D^2$  and let  $x_i = r^{i-1}(1)$ ,  $i = 1, \dots, q$ . Let  $Y \subset D^2$  be the cone of  $\{x_1, \dots, x_q\}$  with cone point  $c$ . Embed  $Y \times I / (x_i, 0) \sim (x_{i+1}, 1)$  into the solid torus  $V = D^2 \times I / (x, 0) \sim (r(x), 1)$ .

The punctured lens space  $L(p, q)$  is obtained from  $V$  by attaching a 2-handle  $D \times I$  with  $\partial D$  attached to the boundary curve of  $(Y \times I) / \sim$ . Then  $L(p, q)$  deformation retracts to  $(Y \times I) / \sim \cup D$ , which is the 2-stratifold with one white vertex of genus 0, one black vertex, and one edge with label  $q$ .

EXAMPLE 3. Lens space  $L(1, 0) = S^2 \times S^1$  and non-orientable  $S^2$ -bundle over  $S^1$ .

Consider  $S^2 \tilde{\times} S^1$ , the non-orientable  $S^2$ -bundle over  $S^1$ , as the quotient space  $q(S^2 \times I)$  under the quotient map  $q : S^2 \times I \rightarrow S^2 \tilde{\times} S^1$  that identifies  $(x, 0)$  with  $(\alpha(x), 1)$ ,  $x \in S^2$ , where  $\alpha$  is the antipodal map.

Let  $D_0 \subset S^2 \times \{0\}$  be a disk and  $B_1$  be the 3-ball  $D_0 \times I \subset S^2 \times I$ , let  $D_1$  be the disk  $B_1 \cap S^2 \times \{1\}$ , let  $A$  be the annulus  $\partial B_1 - (Int(D_0) \cup Int(D_1))$ , and let  $B_2$  be the ball  $S^2 \times I - Int(B_1)$ , see Figure 2. Then  $S^2 \times I - Int(B_2) = S^2 \times \{0\} \cup B_1 \cup S^2 \times \{1\}$  and  $S^2 \tilde{\times} S^1 - Int(B_2) = q(S^2 \times I - Int(B_2)) = S^2 \cup q(B_1)$ , where  $S^2 = q(S^2 \times \{0\}) = q(S^2 \times \{1\})$ . Collapsing the ball  $q(B_1)$  across the free face  $q(D_1)$  onto  $q(A) \cup q(D_0)$  we obtain a collapse of  $S^2 \tilde{\times} S^1 - Int(B_2)$  onto  $(S^2 - Int(q(D_1))) \cup q(A)$ , which is a torus with a disk attached. This is a 2-stratifold  $X_G$  with graph  $G_X$  in Figure 3(b). (The white vertices have genus 0).




Fig.2.  $S^2 \tilde{\times} S^1 - Int(B_2) \searrow S^1 \tilde{\times} S^1 \cup D^2$

A similar construction, considering  $S^2 \times S^1$  as the obvious quotient space of  $q : S^2 \times I \rightarrow S^2 \times S^1$  and first isotoping the ball  $B_1$  such that  $D_0 \cap D_1 = \emptyset$ , we obtain a collapse of  $S^2 \times S^1 - Int(B_2)$  onto a Kleinbottle with a disk attached. This is a 2-stratifold  $X_G$  with graph  $G_X$  in Figure 3(a).




Fig.3. 2-stratifold spines of punctured  $S^2 \tilde{\times} S^1$  and  $P^2 \times S^1$

We would like to thank Mario Eudave-Muñoz for pointing out that in this example the spine for the non-orientable (resp. orientable  $S^2$ -bundle over  $S^1$ ) is a torus (resp. Kleinbottle) with a disk attached, rather than a Kleinbottle (resp. torus) with a disk attached.



EXAMPLE 4.  $P^2 \times S^1$ .

For a one-sided simple closed curve  $c$  in  $P^2$  and a point  $t_0$  in  $S^1$  let  $X = P^2 \times \{t_0\} \cup c \times S^1 \subset P^2 \times S^1$ . Observe that the boundary of a regular neighborhood  $N$  of  $X$  in  $P^2 \times S^1$  is a 2-sphere. Since  $P^2 \times S^1$  is irreducible,  $\partial N$  bounds a 3-ball  $B^3$  and therefore  $P^2 \times S^1 - \text{Int}(B^3) = N$ , which collapses onto  $X = X_G$ , a 2-stratifold with graph in Figure 3(c).

**Proposition 4.** *If the closed 3-manifold  $M_i$  ( $i = 1, 2$ ) has a 2-stratifold spine and  $M$  is a connected sum of  $M_1$  and  $M_2$ , then  $M$  has a 2-stratifold spine.*

Proof. Let  $K_i$  be a 2-stratifold spine of  $M_i$ . Let  $K_1 \vee K_2$  be obtained by identifying, in the disjoint union of  $K_1$  and  $K_2$  a nonsingular point of  $K_1$  with a nonsingular point of  $K_2$ . By Lemma 1 of [9],  $K_1 \vee K_2$  is a spine of  $M$ . Though  $K_1 \vee K_2$  is not a 2-stratifold, by performing the operation explained below (replacing the wedge point by a disk) we will change  $K_1 \vee K_2$  to a 2-stratifold spine  $K_1 \Delta K_2$ .




Fig.4.  $K_1 \Delta K_2$

A 3-ball neighborhood  $B^3$  of the wedge point of  $K_1 \vee K_2$  intersects  $K_1 \vee K_2$  in the double cone shown in Fig.4. Replace, in  $K_1 \vee K_2$ ,  $K_1 \vee K_2 \cap B^3$  by  $A \cup D$ , as shown in Fig. 4, where  $A = S^1 \times [0, 1]$  is an horizontal cylinder,  $\partial A = (K_1 \vee K_2) \cap \partial B^3$ ,  $D$  is a vertical 2-disk with  $A \cap D = \partial D = S^1 \times (1/2)$ . The result is a 2-stratifold  $K_1 \Delta K_2$ . There is a homeomorphism from  $B^3 - (A \cup D)$  onto  $B^3 - K_1 \vee K_2$  which is the identity on the boundary (roughly collapse  $D$  to a point) and so  $M - K_1 \Delta K_2$  is homeomorphic to  $M - K_1 \vee K_2$  which is homeomorphic to  $R^3$ .

Therefore  $K_1 \Delta K_2$  is a 2-stratifold spine of  $M$ . □

Now Theorem 1 together with the examples and Proposition 4 yields our main Theorem. Here we do not consider  $S^3$  to be a lens space.

**Theorem 2.** *A closed 3-manifold  $M$  has a 2-stratifold as a spine if and only if  $M$  is a connected sum of lens spaces,  $S^2$ -bundles over  $S^1$ , and  $P^2 \times S^1$ 's.*

ACKNOWLEDGEMENTS. J. C. Gómez-Larrañaga would like to thank LAISLA and the TDA project from CIMAT for financial support and IST Austria for their hospitality.

---

### References

- [1] M. Aschenbrenner, S. Friedel and H. Wilson: *Decision Problems for 3-manifolds and their fundamental groups*; in Interactions between low-dimensional topology and mapping class groups, Geometry & Topology Monographs **19**, Geon. Topol. Publ., Coventry, 201–236, 2015.
- [2] P. Bendich, E. Gasparovicy, C.J. Traliez and J. Harer: *Scaffoldings and Spines: Organizing High-Dimensional Data Using Cover Trees, Local Principal Component Analysis, and Persistent Homology*, Research in computational topology, 93–114, Assoc. Women Math. Ser. **13**, Springer, Cham, 2018.
- [3] S. Boyer, D. Rolfsen and B. Wiest: *Orderable 3-manifold groups*, Ann. Inst. Fourier (Grenoble) **55** (2005), 243–288.
- [4] J.S. Carter: *Reidemeister/Roseman-type moves to embedded foams in 4-dimensional space*; in New Ideas in low Dimensional Topology, Series on Knots and Everything **56**, World Sci. Publ., Hackensack, NJ, 1–30, 2015.
- [5] I.M. Chiswell: *Right orderability and graphs of groups*, J. Group Theory **14** (2011), 589–601.
- [6] J.C. Gómez-Larrañaga, F. González-Acuña and Wolfgang Heil: *Categorical group invariants of 3-manifolds*, Manuscripta Math. **145** (2014), 433–448.
- [7] J.C. Gómez-Larrañaga, F. González-Acuña and Wolfgang Heil: *2-dimensional stratifolds*; in A Mathematical Tribute to José María Montesinos Amilibia, Universidad Complutense de Madrid, 395–405, 2016.
- [8] J.C. Gómez-Larrañaga, F. González-Acuña and Wolfgang Heil: *2-stratifold groups have solvable Word Problem*, Rev. R. Acad. Cienc. Exactas, Fís. Nat. Ser. A Mat. RACSAM **12** (2018), 803–810.
- [9] G.W. Knutson: *A characterization of closed 3-manifolds with spines containing no wild arcs*, Proc. Amer. Math. Soc. **21** (1969), 310–314.
- [10] M. Khovanov: *sl(3) link homology*, Algebr. Geom. Topol. **4** (2004), 1045–1081.
- [11] P. Lum, G. Singh, J. Carlsson, A. Lehman, T. Ishkhanov, M. Vejdemo-Johansson, M. Alagappan and G. Carlsson: *Extracting insights from the shape of complex data using topology*, Nature Scientific Reports **3** (2013), Article ID 12–36.
- [12] R.C. Lyndon and P.E. Schupp: *Combinatorial Group Theory*, Modern Surveys in Math. **89**, Springer Verlag, Berlin, 1977.
- [13] S. Matsuzaki and M. Ozawa: *Genera and minors of multibranching surfaces*, Topology Appl. **230** (2017), 621–638.
- [14] S. Matveev: *Algorithmic Topology and Classification of 3-Manifolds*, Algorithms and Computation in Mathematics **9**, Springer, Berlin, 2007.
- [15] J. Milnor: *Groups which act on  $S^n$  without fixed points*, Amer. J. Math. **79** (1957), 623–630.
- [16] W. Neumann: *A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves*, Trans. Amer. Math. Soc. **268** (1981), 299–344.
- [17] P. Scott and C.T.C. Wall: *Topological Methods in Group Theory*; in Homological Group Theory, London Math. Soc. Lecture Notes Ser. **36**, Cambridge Univ. Press, Cambridge-New York, 1979.
- [18] J.P. Serre: *Trees*, Springer-Verlag, Berlin-New York, 1980.
- [19] G.A. Swarup: *Projective planes in irreducible 3-manifolds*, Math. Z. **132** (1973), 305–317.

J.C. Gómez-Larrañaga  
Centro de Investigación en Matemáticas  
A.P. 402, Guanajuato 36000, Gto.  
México  
e-mail: jcarlos@cimat.mx

F. González-Acuña  
Instituto de Matemáticas  
UNAM, 62210 Cuernavaca, Morelos  
México and Centro de Investigación en Matemáticas  
A.P. 402, Guanajuato 36000, Gto.  
México  
e-mail: fico@math.unam.mx

Wolfgang Heil  
Department of Mathematics  
Florida State University  
Tallahassee, FL 32306  
USA  
e-mail: heil@math.fsu.edu