

## INVARIANT SUBSPACES FOR FINITE MAXIMAL SUBDIAGONAL ALGEBRAS

KICHI-SUKE SAITO

Let  $M$  be a von Neumann algebra with a faithful, normal, tracial state  $\tau$  and  $H^\infty$  a finite, maximal, subdiagonal algebra in  $M$ . If  $1 \leq p < s \leq \infty$ , then there is a one-to-one correspondence between left-(resp. right-) invariant subspaces of the noncommutative Lebesgue space  $L^p(M, \tau)$  and those of  $L^s(M, \tau)$ .

1. Introduction. Let  $M$  be a von Neumann algebra with a faithful, normal, tracial state  $\tau$  and let  $H^\infty$  be a finite, maximal, subdiagonal algebra in  $M$ . A number of authors have investigated the structure of the invariant subspaces for  $H^\infty$  acting on the noncommutative Lebesgue space  $L^p(M, \tau)$  (cf. [3], [4], [5] and [6]). In [6], we showed that, if  $\mathfrak{M}$  is a left-(resp. right-) invariant subspace of  $L^p(M, \tau)$ ,  $1 \leq p < \infty$ , then  $\mathfrak{M}$  is the closure of the space of bounded elements it contains.

In this paper, we shall show that, if  $1 \leq p < s \leq \infty$ , then there is a one-to-one correspondence between left- (resp. right-) invariant subspaces  $\mathfrak{M}_p$  of  $L^p(M, \tau)$  and left- (resp. right-) invariant subspaces  $\mathfrak{M}_s$  of  $L^s(M, \tau)$ , such that  $\mathfrak{M}_s = \mathfrak{M}_p \cap L^s(M, \tau)$  and  $\mathfrak{M}_p$  is the closure in  $L^p(M, \tau)$  of  $\mathfrak{M}_s$ . This is of course true in the weak\*-Dirichlet algebras setting (cf. [2, p. 131]) and this is attractive to study the structure of the invariant subspaces of  $L^p(M, \tau)$ .

2. Let  $M$  be a von Neumann algebra with a faithful, normal, tracial state  $\tau$ . We shall denote the noncommutative Lebesgue spaces associated with  $M$  and  $\tau$  by  $L^p(M, \tau)$ ,  $1 \leq p < \infty$  (cf. [7]). As is customary,  $M$  will be identified with  $L^\infty(M, \tau)$ . The closure of a subset  $S$  of  $L^p(M, \tau)$  in the  $L^p$ -norm will be denoted by  $[S]_p$ ;  $[S]_\infty$  will denote the closure of  $S$  in the  $\sigma$ -weak topology on  $L^\infty(M, \tau)$ .

DEFINITION 1. Let  $H^\infty$  be a  $\sigma$ -weakly closed subalgebra of  $M$  containing the identity operator 1 and let  $\Phi$  be a faithful, normal expectation from  $M$  onto  $D = H^\infty \cap H^{\infty*}$  ( $H^{\infty*} = \{x^*: x \in H^\infty\}$ ). Then  $H^\infty$  is called a finite, maximal, subdiagonal algebra in  $M$  with respect to  $\Phi$  and  $\tau$  in case the following conditions are satisfied: (1)  $H^\infty + H^{\infty*}$  is  $\sigma$ -weakly dense in  $M$ ; (2)  $\Phi(xy) = \Phi(x)\Phi(y)$ , for all  $x, y \in H^\infty$ ; (3)  $H^\infty$  is maximal among those subalgebras of  $M$  satisfying (1) and (2); and (4)  $\tau \circ \Phi = \tau$ .

For  $1 \leq p < \infty$ , the closure of  $H^\infty$  in  $L^p(M, \tau)$  is denoted by  $H^p$  and the closure of  $H_0^\infty = \{x \in H^\infty: \Phi(x) = 0\}$  is denoted by  $H_0^p$ .

**DEFINITION 2.** Let  $\mathfrak{M}$  be a closed (resp.  $\sigma$ -weakly closed) subspace of  $L^p(M, \tau)$  (resp.  $L^\infty(M, \tau)$ ). We shall say that  $\mathfrak{M}$  is left- (resp. right-) invariant if  $H^\infty \mathfrak{M} \subseteq \mathfrak{M}$  (resp.  $\mathfrak{M} H^\infty \subseteq \mathfrak{M}$ ).

The following theorem shows that, in considering left- (resp. right-) invariant subspaces, it suffices to consider left- (resp. right-) invariant subspaces of  $L^2(M, \tau)$ , or alternatively,  $\sigma$ -weakly closed left- (resp. right-) invariant subspaces of  $L^\infty(M, \tau)$ . The method in the proof is based on a factorization theorem, that is, if  $k$  is in  $M$  with inverse lying in  $L^2(M, \tau)$ , then there are unitary operators  $u_1, u_2$  in  $M$  and operators  $a_1, a_2$  in  $H^\infty$  with inverses lying in  $H^2$  such that  $k = u_1 a_1 = a_2 u_2$  ([6, Proposition 1]).

**THEOREM 1.** Suppose  $1 \leq p < s \leq \infty$ .

(1) If  $\mathfrak{M}$  is a left- (resp. right-) invariant subspace of  $L^p(N, \tau)$ , then  $\mathfrak{M} \cap L^s(M, \tau)$  is a left- (resp. right-) invariant subspace of  $L^s(M, \tau)$  and  $\mathfrak{M} = [\mathfrak{M} \cap L^s(M, \tau)]_p$ .

(2) If  $\mathfrak{M}$  is a left- (resp. right-) invariant subspace of  $L^s(M, \tau)$ , then  $[\mathfrak{M}]_p$  is a left- (resp. right-) invariant subspace of  $L^p(M, \tau)$  and  $\mathfrak{M} = [\mathfrak{M}]_p \cap L^s(M, \tau)$ .

*Proof.* It suffices to consider the assertion for left- invariant subspaces.

(1) Let  $\mathfrak{M}$  be a left-invariant subspace of  $L^p(M, \tau)$ . It is clear that  $\mathfrak{M} \cap L^s(M, \tau)$  is a left-invariant subspace of  $L^s(M, \tau)$ . By [6, Theorem], we have  $\mathfrak{M} = [\mathfrak{M} \cap L^\infty(M, \tau)]_p$  and so

$$\mathfrak{M} = [\mathfrak{M} \cap L^\infty(M, \tau)]_p \subseteq [\mathfrak{M} \cap L^s(M, \tau)]_p \subseteq \mathfrak{M}.$$

Therefore  $\mathfrak{M} = [\mathfrak{M} \cap L^s(M, \tau)]_p$ . This completes the proof of (1).

(2) Let  $\mathfrak{M}$  be a left-invariant subspace of  $L^s(M, \tau)$ . It is clear that  $[\mathfrak{M}]_p$  is a left-invariant subspace of  $L^p(M, \tau)$ . Now, if the assertion (2) in the case  $s = \infty$  is proved, then  $[\mathfrak{M} \cap L^\infty(M, \tau)]_p \cap L^\infty(M, \tau) = \mathfrak{M} \cap L^\infty(M, \tau)$ . By (1),

$$\begin{aligned} [\mathfrak{M}]_p \cap L^s(M, \tau) &= [[\mathfrak{M}]_p \cap L^\infty(M, \tau)]_s = [[\mathfrak{M} \cap L^\infty(M, \tau)]_p \cap L^\infty(M, \tau)]_s \\ &= [\mathfrak{M} \cap L^\infty(M, \tau)]_s = \mathfrak{M}. \end{aligned}$$

Therefore, suppose that  $s = \infty$ . Let  $\mathfrak{M}$  be a left-invariant subspace of  $L^\infty(M, \tau)$  and put  $\tilde{\mathfrak{M}} = [\mathfrak{M}]_p \cap L^\infty(M, \tau)$ . It is clear that  $\mathfrak{M} \subseteq \tilde{\mathfrak{M}}$ . If  $\mathfrak{M} \subsetneq \tilde{\mathfrak{M}}$ , then there exist  $x \in \tilde{\mathfrak{M}} \setminus \mathfrak{M}$  and  $a \in L^1(M, \tau)$  such that  $\tau(ax) = 1$  and  $\tau(ay) = 0$  for every  $y \in \mathfrak{M}$ .

(i) Case  $2 \leq p < \infty$ . Define the number  $q$  by the equation  $1/p + 1/q = 1$ . Let  $a = v|a|$  be the polar decomposition of  $a$ . Let  $f$  be the function on  $[0, \infty)$  defined by the formula  $f(t) = 1, 0 \leq t \leq 1, f(t) = t^{-1}, t > 1$ , and define  $k$  to be  $f(|a|^{1/p})$  through the functional calculus. Then note that  $k \in L^\infty(M, \tau)$  and  $k^{-1} \in L^p(M, \tau) \subset L^2(M, \tau)$ . By [6, Proposition 1], we may choose a unitary operator  $u$  in  $L^\infty(M, \tau)$  and an operator  $b \in H^\infty$  such that  $k = bu$  and  $b^{-1} \in H^2$ . Since  $k^{-1} \in L^p(M, \tau)$ , by [6, Proposition 2],  $b^{-1} \in L^p(M, \tau) \cap H^2 = H^p$  and note that  $ab = v|a|^{1/q}|a|^{1/p}ku^* \in L^q(M, \tau)$ , because  $|a|^{1/p}k \in L^\infty(M, \tau)$ . Since  $\mathfrak{M}$  is left-invariant,  $\tau(aby) = 0$  for every  $y \in \mathfrak{M}$  and so  $\tau(aby) = 0$  for every  $y \in [\mathfrak{M}]_p$ . On the other hand,  $b^{-1}x \in H^p \widetilde{\mathfrak{M}} \subset [\widetilde{\mathfrak{M}}]_p = [\mathfrak{M}]_p$  and so  $\tau(ax) = \tau(abb^{-1}x) = 0$ . This is a contradiction. Thus  $\mathfrak{M} = \widetilde{\mathfrak{M}}$ .

(ii) Case  $1 \leq p < 2$ . Define the numbers  $q$  and  $r$  by the equations  $1/p + 1/q = 1$  and  $1/r + 1/2 = 1/p$ . Put  $k = f(|a|^{1/2})$ , where  $f$  is the function in (i). By [6, Proposition 1], there are a unitary operator  $u$  in  $L^\infty(M, \tau)$  and an operator  $b \in H^\infty$  with inverse lying in  $H^2$  such that  $k = bu$  and note that  $ab$  is a nonzero element in  $L^2(M, \tau)$ . Also, let  $ab = v'|ab|$  be the polar decomposition of  $ab$ . Put  $k' = f(|ab|^{2/r})$ , where  $f$  is the function in (i). Since  $|ab|^{2/r} \in L^r(M, \tau) \subset L^2(M, \tau)$ , by [6, Proposition 1], there exists an operator  $c$  in  $H^\infty$  with inverse lying in  $H^r$  such that  $abc$  is a nonzero element in  $L^q(M, \tau)$ . Since  $\mathfrak{M}$  is left-invariant, we have  $\tau((abc)y) = \tau(a(bcy)) = 0$ , for every  $y \in \mathfrak{M}$ , and so  $\tau(abcy) = 0$  for every  $y \in [\mathfrak{M}]_2$ . On the other hand, since  $(bc)^{-1} = c^{-1}b^{-1} \in H^r H^2 \subset H^p$ ,  $(bc)^{-1}x \in H^p \mathfrak{M} \subset [\widetilde{\mathfrak{M}}]_p = [\mathfrak{M}]_p$  and so  $\tau(ax) = \tau(abc(bc)^{-1}x) = 0$ . This is a contradiction. Thus  $\mathfrak{M} = \widetilde{\mathfrak{M}}$ .

This completes the proof of (2).

Next we shall consider the structure of doubly invariant subspaces and simply invariant subspaces of  $L^p(M, \tau), 1 \leq p \leq \infty$ .

DEFINITION 3. Let  $\mathfrak{M}$  be a closed subspace of  $L^p(M, \tau), 1 \leq p \leq \infty$ .

(1)  $\mathfrak{M}$  is said to be left (resp. right) doubly invariant if  $(H^\infty + H^{\infty'})\mathfrak{M} \subseteq \mathfrak{M}$  (resp.  $\mathfrak{M}(H^\infty + H^{\infty'}) \subseteq \mathfrak{M}$ ).

(2)  $\mathfrak{M}$  is said to be left (resp. right) simply invariant if  $[H_0^\infty \mathfrak{M}]_p \subsetneq \mathfrak{M}$  (resp.  $[\mathfrak{M} H_0^\infty]_p \subsetneq \mathfrak{M}$ ).

By [5, Theorem 4.1] and Theorem 1, we have the following theorem.

THEOREM 2. Let  $\mathfrak{M}$  be a closed subspace of  $L^p(M, \tau), 1 \leq p \leq \infty$ . Then  $\mathfrak{M}$  is a left (resp. right) doubly invariant subspace of  $L^p(M, \tau)$  if and only if there exists a projection  $e$  in  $M$  such that  $\mathfrak{M} = L^p(M, \tau)e$  (resp.  $eL^p(M, \tau)$ ).

In [3], Kamei has shown the simply invariant subspace theorem for antisymmetric finite subdiagonal algebras in case  $p = 1, 2$ . Also, in [5], we characterized the simply invariant subspace for  $H^\infty$  in  $L^p(M, \tau)$ ,  $1 \leq p \leq \infty$ , when  $H^\infty$  is determined by a trace preserving ergodic flow. However, by Theorem 1 and [3], we have the following theorem.

**THEOREM 3.** *Let  $\mathfrak{M}$  be a closed subspace of  $L^p(M, \tau)$ ,  $1 \leq p \leq \infty$ . If  $H^\infty$  is antisymmetric, that is,  $D = C1$ , then  $\mathfrak{M}$  is a left (resp. right) simply invariant subspace of  $L^p(M, \tau)$  if and only if there is a unitary operator  $u$  in  $M$  such that  $\mathfrak{M} = H^p u$  (resp.  $uH^p$ ).*

#### REFERENCES

1. W. B. Arveson, *Analyticity in operator algebras*, Amer. J. Math., **89** (1967), 578-642.
2. T. W. Gamelin, *Uniform Algebras*, Prentice-Hall, Englewood Cliffs, N.J., 1969.
3. N. Kamei, *Simply invariant subspaces theorems for antisymmetric finite subdiagonal algebras*, Tôhoku Math. J., **21** (1969), 467-473.
4. M. McAsey, P. S. Muhly and K.-S. Saito, *Nonselfadjoint crossed products (Invariant subspaces and maximality)*, Trans. Amer. Math. Soc., **248** (1979), 381-409.
5. K.-S. Saito, *On non-commutative Hardy spaces associated with flows on finite von Neumann algebras*, Tôhoku Math. J., **29** (1977), 585-595.
6. ———, *A note on invariant subspaces for finite maximal subdiagonal algebras*, Proc. Amer. Math. Soc., **77** (1979), 348-352.
7. I. E. Segal, *A non-commutative extension of abstract integration*, Ann. of Math., **57** (1953), 401-457.

Received September 11, 1979.

NIGATA UNIVERSITY  
 IKARASHI 2-8050,  
 NIGATE 950-21, JAPAN