

CONSTRUCTIVE VERSIONS OF TARSKI'S FIXED POINT THEOREMS

PATRICK COUSOT AND RADHIA COUSOT

Let F be a monotone operator on the complete lattice L into itself. Tarski's lattice theoretical fixed point theorem states that the set of fixed points of F is a nonempty complete lattice for the ordering of L . We give a constructive proof of this theorem showing that the set of fixed points of F is the image of L by a lower and an upper preclosure operator. These preclosure operators are the composition of lower and upper closure operators which are defined by means of limits of stationary transfinite iteration sequences for F . In the same way we give a constructive characterization of the set of common fixed points of a family of commuting operators. Finally we examine some consequences of additional semi-continuity hypotheses.

1. Introduction. Let $L(\subseteq, \perp, \top, \cup, \cap)$ be a nonempty complete lattice with partial ordering \subseteq , least upper bound \cup , greatest lower bound \cap . The infimum \perp of L is $\cap L$, the supremum \top of L is $\cup L$. (Birkhoff's standard reference book [3] provides the necessary background material.) Set inclusion, union and intersection are respectively denoted by \subseteq , \cup and \cap .

Let F be a monotone operator on $L(\subseteq, \perp, \top, \cup, \cap)$ into itself (i.e., $\forall X, Y \in L, \{X \subseteq Y\} \Rightarrow \{F(X) \subseteq F(Y)\}$).

The fundamental theorem of Tarski [19] states that the set $fp(F)$ of fixed points of F (i.e., $fp(F) = \{X \in L: X = F(X)\}$) is a nonempty complete lattice with ordering \subseteq . The proof of this theorem is based on the definition of the least fixed point $lfp(F)$ of F by $lfp(F) = \cap \{X \in L: F(X) \subseteq X\}$. The least upper bound of $S \subseteq fp(F)$ in $fp(F)$ is the least fixed point of the restriction of F to the complete lattice $\{X \in L: (\cup S) \subseteq X\}$. An application of the duality principle completes the proof.

This definition is not constructive and many applications of Tarski's theorem (specially in computer science (Cousot [5]) and numerical analysis (Amann [2])) use the alternative characterization of $lfp(F)$ as $\cup \{F^i(\perp): i \in \mathbb{N}\}$. This iteration scheme which originates from Kleene [10]'s first recursion theorem and which was used by Tarski [19] for complete morphisms, has the drawback to require the additional assumption that F is semi-continuous ($F(\cup S) = \cup F(S)$) for every increasing nonempty chain S , see e.g., Kolodner [11]).

The purpose of this paper is to give a constructive proof of Tarski's theorem without using the continuity hypothesis. The set of fixed points of F is shown to be the image of L by preclosure operations defined by means of limits of stationary transfinite iteration sequences. Then the set of common fixed points of a family of commuting monotone operators on a complete lattice into itself is characterized in the same way. The advantage of characterizing fixed points by iterative schemes is that they lead to practical computation or approximation procedures. Also the definition of fixed points as limits of stationary iteration sequences allows the use of transfinite induction for proving properties of these fixed points.

Finally some consequences of the additional and less general continuity hypothesis are examined.

2. Definitions.

DEFINITION 2.1. (*Upper iteration sequence.*) Let $L(\subseteq, \perp, \top, \cup, \cap)$ be a complete lattice, μ the smallest ordinal such that the class $\{\delta: \delta \in \mu\}$ has a cardinality greater than the cardinality $\text{Card}(L)$ of L and F a monotone operator on L into itself. The μ -termed upper iteration sequence for F starting with $D \in L$ is the μ -termed sequence $\langle X^\delta, \delta \in \mu \rangle$ of elements of L defined by transfinite recursion in the following way:

$$(a) \quad X^0 = D$$

$$(b) \quad X^\delta = F(X^{\delta-1}) \text{ for every successor ordinal } \delta \in \mu$$

$$(c) \quad X^\delta = \bigcup_{\alpha < \delta} X^\alpha \text{ for every limit ordinal } \delta \in \mu$$

(the dual lower iteration sequence is defined by:

$$(c') \quad X^\delta = \bigcap_{\alpha < \delta} X^\alpha \text{ for every limit ordinal } \delta \in \mu).$$

DEFINITION 2.2. (*Limit of a stationary transfinite sequence.*) We say that the sequence $\langle X^\delta, \delta \in \mu \rangle$ is *stationary* if and only if $\{\exists \varepsilon \in \mu: \{\forall \beta \in \mu, \{\beta \geq \varepsilon\} \Rightarrow \{X^\varepsilon = X^\beta\}\}\}$ in which case the *limit* of the sequence is X^ε . We denote by $\text{lwis}(F)(D)$ the limit of a stationary upper iteration sequence for F starting with D (dually $\text{llis}(F)(D)$).

In the following the class of ordinals, the ordinal addition, the ordinal multiplication and the first infinite limit ordinal are respectively denoted by Ord , $+$, \cdot and ω (the definition of $+$ and \cdot shall be used in the form stated by Birkhoff [3]).

The set of *prefixed points* of F is $\text{pref}p(F) = \{X \in L: X \subseteq F(X)\}$. Dually $\text{post}fp(F) = \{X \in L: F(X) \subseteq X\}$. Therefore $\text{fp}(F) = \text{pref}p(F) \cap \text{post}fp(F)$.

We use Church [4]'s lambda notation (so that F is $\lambda X. F(X)$).

3. Behavior of an upper iteration sequence.

LEMMA 3.1. Let $\langle X^\delta, \delta \in \text{Ord} \rangle$ be the Ord-termed upper iteration sequence for the monotone operator F on the complete lattice $L(\subseteq, \perp, \top, \cup, \cap)$ into itself starting with $D \in L$,

- (1) $\forall P \in L, \{D \subseteq P\} \text{ and } \{F(P) \subseteq P\} \Rightarrow \{\forall \delta \in \text{Ord}, X^\delta \subseteq P\}$,
- (2) $\{D \in \text{postfp}(F)\} \Rightarrow \{\forall \delta \in \text{Ord}, X^\delta \subseteq D\}$.

Proof. Let $P \in L$ be such that $D \subseteq P$ and $F(P) \subseteq P$, then by Definition 2.1(a) $D = X^0 \subseteq P$. Assume that $\forall \alpha \in \text{Ord}, \{\alpha < \delta\} \Rightarrow \{X^\alpha \subseteq P\}$. If δ is a successor ordinal, then we have $X^{\delta-1} \subseteq P$ so that by monotony $F(X^{\delta-1}) \subseteq F(P) \subseteq P$ proving by Definition 2.1(b) that $X^\delta \subseteq P$. If δ is a limit ordinal then by induction hypothesis and definition of least upper bounds $\bigcup_{\alpha < \delta} X^\alpha \subseteq P$ proving by Definition 2.1(c) that $X^\delta \subseteq P$. By transfinite induction $\forall \delta \in \text{Ord}, X^\delta \subseteq P$. In particular when $D \in \text{postfp}(F)$ we have $D \subseteq D$ and $F(D) \subseteq D$ which imply $\forall \delta \in \text{Ord}, X^\delta \subseteq D$.

THEOREM 3.2. Let $\langle X^\delta, \delta \in \text{Ord} \rangle$ be the Ord-termed upper iteration sequence for the monotone operator F on the complete lattice $L(\subseteq, \perp, \top, \cup, \cap)$ into itself starting with $D \in L$,

(1) $\forall \delta \in \text{Ord}$, let $\beta \leq \delta$ and $n < \omega$ be respectively the quotient and remainder of the ordinal division of δ by ω (i.e., $\delta = \beta \cdot \omega + n$), $\forall \beta' : \beta' > \beta, \forall \gamma : \beta' \cdot \omega \leq \gamma \leq \beta' \cdot \omega + n, X^\beta \subseteq X^\gamma$.

(2) The subsequence $\langle X^{\alpha \cdot \omega}, \alpha \in \mu \rangle$ is a stationary increasing chain, its limit $X^{\gamma \cdot \omega}$ is the least postfixed point of F greater than or equal to D .

(3) There exists a smallest limit ordinal ξ such that $\xi \leq \eta \cdot \omega$ and $X^\xi \in \text{prefp}(F) \cup \text{postfp}(F)$.

(4) If $X^\xi \in \text{prefp}(F)$ then the subsequence $\langle X^\delta, \xi \leq \delta < \xi + \mu \rangle$ (as well as $\langle X^\delta, \xi \leq \delta \rangle$) is a stationary increasing chain of elements of $\text{prefp}(F)$, its limit $\text{luis}(F)(X^\xi)$ is equal to $X^{\gamma \cdot \omega}$ which is the least of the fixed points of F greater than or equal to D .

(5) If $X^\xi \in \text{postfp}(F)$ then $\langle X^{\xi+n}, n \in \omega \rangle$ is a decreasing chain of elements of $\text{postfp}(F)$ and $\forall \delta \in \text{Ord}, X^{\xi+\delta} = X^{\xi+m}$ where m is the remainder of the ordinal division of δ by ω .

Proof.

(1) $\forall \delta \in \text{Ord}$, there exist unique β and n such that $\delta = \beta \cdot \omega + n$ and $\beta \leq \delta, n < \omega$. If δ is a limit ordinal then $n = 0$ and $\forall \beta' > \beta, \beta' \cdot \omega > \beta \cdot \omega = \delta$ and $\beta' \cdot \omega$ is an infinite limit ordinal so that by Definition 2.1(c) $X^\delta \subseteq \bigcup_{\alpha < \beta' \cdot \omega} X^\alpha = X^{\beta' \cdot \omega}$. If $n \neq 0$ then δ is a successor ordinal and $(\delta - 1) = \beta \cdot \omega + (n - 1)$. Assume that $\forall \beta'$ such that $\beta' > \beta$ and $\forall \gamma$ such that $\beta' \cdot \omega \leq \gamma \leq \beta' \cdot \omega + (n - 1)$ we have

$X^{\delta-1} \subseteq X^\gamma$. According to Definition 2.1(b) and by monotony $X^\delta = F(X^{\delta-1}) \subseteq F(X^\gamma) = X^{\gamma+1}$. Also $X^\delta \subseteq X^{\beta' \cdot \omega}$ therefore with $\gamma' = \gamma + 1$ we get $\forall \beta': \beta' > \beta, \forall \gamma': \beta' \cdot \omega \leq \gamma' \leq \beta' \cdot \omega + n, X^\delta \subseteq X^{\gamma'}$. By transfinite induction on δ Theorem 3.2(1) is proved.

(2) By 3.2(1) the subsequence $\langle X^{\alpha \cdot \omega}, \alpha \in \mu \rangle$ is an increasing chain. Assume that $\{\forall \eta \in \text{Ord}, \{\eta \in \mu \text{ and } (\eta + 1) \in \mu\} \Rightarrow \{X^{\eta \cdot \omega} \neq X^{(\eta+1) \cdot \omega}\}\}$. This implies that $\langle X^{\alpha \cdot \omega}, \alpha \in \mu \rangle$ is a strictly increasing chain so that the class $\{X^{\alpha \cdot \omega}: \alpha \in \mu\}$ is equipotent with the class $\{\alpha \cdot \omega: \alpha \in \mu\}$. Since $\lambda \alpha. (\alpha \cdot \omega)$ is a one-one function mapping $\{\alpha: \alpha \in \mu\}$ onto $\{\alpha \cdot \omega: \alpha \in \mu\}$ the class $\{X^{\alpha \cdot \omega}: \alpha \in \mu\}$ is equipotent with the class $\{\alpha: \alpha \in \mu\}$. Therefore by definition of μ we have $\text{Card}(\{X^{\alpha \cdot \omega}: \alpha \in \mu\}) > \text{Card}(L)$ and also by $\{\forall \alpha \in \mu, X^{\alpha \cdot \omega} \in L\}$ we obtain the contradiction $\text{Card}(\{X^{\alpha \cdot \omega}: \alpha \in \mu\}) \leq \text{Card}(L)$. By reductio ad absurdum $\{\exists \eta: (\eta \in \mu) \text{ and } ((\eta + 1) \in \mu) \text{ and } X^{\eta \cdot \omega} = X^{(\eta+1) \cdot \omega}\}$.

Since $(\eta \cdot \omega) + 1 < (\eta + 1) \cdot \omega$ and $(\eta + 1) \cdot \omega$ is an infinite limit ordinal Definitions 2.1(b) and 2.1(c) imply that $F(X^{\eta \cdot \omega}) = X^{(\eta \cdot \omega) + 1} \subseteq \bigcup_{\alpha < (\eta+1) \cdot \omega} X^\alpha = X^{(\eta+1) \cdot \omega} = X^{\eta \cdot \omega}$. Also $D = X^0 \subseteq \bigcup_{\alpha < \eta \cdot \omega} X^\alpha = X^{\eta \cdot \omega}$ so that $X^{\eta \cdot \omega}$ is a postfixed point of F greater than or equal to D . Let $P \in L$ be such that $F(P) \subseteq P$ and $D \subseteq P$. Then Lemma 3.1(1) implies that $X^{\eta \cdot \omega} \subseteq P$ proving that $X^{\eta \cdot \omega}$ is the least postfixed point of F greater than or equal to D .

$\forall \alpha \in \text{Ord}, \alpha > \eta$ implies $\alpha \cdot \omega > \eta \cdot \omega$ and therefore by Definition 2.1(c) $X^{\alpha \cdot \omega} = \bigcup_{\beta < \alpha \cdot \omega} X^\beta = X^{\eta \cdot \omega} \cup (\bigcup_{\eta \cdot \omega \leq \beta < \alpha \cdot \omega} X^\beta)$. But $X^{\eta \cdot \omega} \in \text{postfp}(F)$ so that according to Lemma 3.1(2), $\forall \beta \geq \eta \cdot \omega$ we have $X^\beta \subseteq X^{\eta \cdot \omega}$ proving that $X^{\alpha \cdot \omega} = X^{\eta \cdot \omega}$ and that $\langle X^{\alpha \cdot \omega}, \alpha \in \mu \rangle$ and $\langle X^{\alpha \cdot \omega}, \alpha \in \text{Ord} \rangle$ are stationary.

(The following Theorem 4.1 will show that $X^{\eta \cdot \omega}$ can be constructed more directly as $\text{luis}(\lambda X. X \cup F(X))(D) = \text{luis}(\lambda X. D \cup F(X))(D)$.)

(3) Since $X^{\eta \cdot \omega} \in \text{postfp}(F)$ and Ord is well-ordered there exists a smallest limit ordinal $\xi \leq \eta \cdot \omega$ such that X^ξ and $F(X^\xi)$ are comparable.

(4) If $X^\xi \in \text{prefp}(F)$ then by monotony of F , Definition 2.1 and transfinite induction, it is easy to prove that $\{\forall \delta, \beta \in \text{Ord}, \{\xi \leq \delta \leq \beta\} \Rightarrow \{D \subseteq X^\delta \subseteq X^\beta \subseteq F(X^\beta)\}\}$. By definition of μ the increasing subchain $\langle X^\delta, \xi \leq \delta < \xi + \mu \rangle$ of elements of L cannot be strictly increasing so that $\{\exists \varepsilon \in \text{Ord}: (\xi \leq \varepsilon < \varepsilon + 1 < \xi + \mu) \text{ and } (X^\varepsilon = X^{\varepsilon+1})\}$. Then by transfinite induction using Definition 2.1 it is immediate that $\langle X^\delta, \xi \leq \delta < \xi + \mu \rangle$ and $\langle X^\delta, \xi \leq \delta \rangle$ are stationary of limit X^ξ . Since $D \subseteq X^\xi = X^{\xi+1} = F(X^\xi)$, X^ξ is a fixed (and postfixed) point of F greater than or equal to D . Let $P \in L$ be such that $D \subseteq P$ and $F(P) \subseteq P$. By Lemma 3.1(1) we have $X^\xi \subseteq P$ proving that X^ξ is the least fixed (and postfixed) point of F greater than or equal to D . Moreover $X^\xi = X^{\eta \cdot \omega}$ by 3.2(2).

(5) When $F(X^\xi) \subseteq X^\xi$ it is easy, using the monotony of F , to

prove by finite induction that the subsequence $\langle X^{\xi+n}, n \in \omega \rangle$ is a decreasing chain. If $\delta = 0$ then $\delta = 0 \cdot \omega + 0$ and obviously $X^{\xi+\delta} = X^\xi = X^{\xi+0}$. Assume that $\forall \alpha \in \text{Ord}, \{\alpha < \delta\} \Rightarrow \{X^{\xi+\alpha} = X^{\xi+m}\}$ where m is the remainder of the ordinal division of α by ω . If δ is a successor ordinal then $\exists \beta \in \text{Ord}, \exists n \in \omega$ such that $\delta = \beta \cdot \omega + n$ with $n \neq 0$. Hence $\delta - 1 = \beta \cdot \omega + (n - 1)$ so that by induction hypothesis $X^{(\xi+\delta)-1} = X^{\xi+(\delta-1)} = X^{\xi+(n-1)} = X^{(\xi+n)-1}$. By Definition 2.1(b), $X^{\xi+\delta} = F(X^{(\xi+\delta)-1}) = F(X^{(\xi+n)-1}) = X^{\xi+n}$. If δ is a limit ordinal then $\xi + \delta$ is a limit ordinal because ξ is a limit ordinal. Hence by Definition 2.1(c) $X^{\xi+\delta} = \bigcup_{r < \xi+\delta} X^r = (\bigcup_{r < \xi} X^r) \cup (\bigcup_{\xi \leq r < \xi+\delta} X^r) = X^\xi \cup (\bigcup_{r < \delta} X^{\xi+r}) = X^\xi$ since $X^\xi \in \text{postfp}(F)$ implies according to Lemma 3.1(2) that $\forall \gamma, X^{\xi+r} \subseteq X^\xi$. By transfinite induction, $\forall \delta \in \text{Ord}, X^{\xi+\delta} = X^{\xi+n}$ where n is the remainder of the ordinal division of δ by ω .

The following corollary is immediate from 3.2(4):

COROLLARY 3.3. (*Behavior of an upper iteration sequence starting from a prefixed point of F .*) A μ -termed upper iteration sequence $\langle X^\delta, \delta \in \mu \rangle$ for F starting with $D \in \text{prefp}(F)$ is a stationary increasing chain, its limit $\text{luis}(F)(D)$ is the least of the fixed points of F greater than or equal to D .

An upper closure operator $\bar{\rho}$ on L into L is monotone, extensive ($\forall X \in L, X \subseteq \bar{\rho}(X)$) and idempotent ($\forall X \in L, \bar{\rho}(\bar{\rho}(X)) = \bar{\rho}(X)$). Dually, a lower closure operator ρ on L into L is monotone, reductive ($\forall X \in L, \rho(X) \subseteq X$) and idempotent.

COROLLARY 3.4. *The restriction of $\text{luis}(F)$ to $\text{prefp}(F)$ is an upper closure operator.*

Proof. $\forall D \in \text{prefp}(F)$, we have $\text{luis}(F)(D) \in \text{fp}(F) \subseteq \text{prefp}(F)$. By 3.3, $D \subseteq \text{luis}(F)(D)$. By transfinite induction it is easy to show that the upper iteration sequence $\langle X^\delta, \delta \in \mu \rangle$ for F starting with a fixed point P of F is such that $\{\forall \delta \in \mu, P = X^\delta\}$ so that in particular for $P = \text{luis}(F)(D)$ we have $\text{luis}(F)(\text{luis}(F)(D)) = \text{luis}(F)(D)$. Finally by transfinite induction it is easy to show that the upper iteration sequences $\langle X^\delta, \delta \in \mu \rangle$ and $\langle Y^\delta, \delta \in \mu \rangle$ starting respectively by prefixed points D and E of L satisfying $D \subseteq E$ are such that $\{\forall \delta \in \mu, X^\delta \subseteq Y^\delta\}$. Therefore by Theorem 3.3, $\exists \varepsilon \in \mu, \exists \varepsilon' \in \mu$ such that $\text{luis}(F)(D) = X^\varepsilon = X^{\max(\varepsilon, \varepsilon')} \subseteq Y^{\max(\varepsilon, \varepsilon')} = Y^{\varepsilon'} = \text{luis}(F)(E)$.

Applying the duality principle, we get:

COROLLARY 3.5. *The restriction of $\text{llis}(F)$ to $\text{postfp}(F)$ is a*

lower closure operator.

4. Constructive characterization of the sets of pre- and post-fixed points of F .

THEOREM 4.1. *The μ -termed upper iteration sequences $\langle X^\delta, \delta \in \mu \rangle$ and $\langle Y^\delta, \delta \in \mu \rangle$ for $\lambda X \cdot X \cup F(X)$ and $\lambda X \cdot D \cup F(X)$ respectively, starting with an arbitrary element D of the complete lattice L are stationary increasing chains such that $\forall \delta \in \mu, X^\delta = Y^\delta$. Their limits $\text{luis}(\lambda X \cdot X \cup F(X))(D)$ and $\text{luis}(\lambda X \cdot D \cup F(X))(D)$ are equal to the least of the postfixed points of F greater than or equal to D .*

Proof. 4.1.1. $\forall D \in L$, D is a prefixed point of $\lambda X \cdot X \cup F(X)$ and $\lambda X \cdot D \cup F(X)$ which are monotone operators on the complete lattice L into itself. Hence Theorem 3.3 implies that $\langle X^\delta, \delta \in \mu \rangle$ and $\langle Y^\delta, \delta \in \mu \rangle$ are stationary increasing chains.

4.1.2. $\forall \delta \in \mu, X^\delta = Y^\delta$.

By Definitions 2.1(a) and 2.1(b) the lemma is true for $\delta = 0$ and $\delta = 1$. Assume it is true for every γ such that $2 \leq \gamma < \delta < \mu$. If δ is the successor of a successor ordinal then $X^\delta = X^{\delta-1} \cup F(X^{\delta-1}) = Y^{\delta-1} \cup F(Y^{\delta-1}) = D \cup F(Y^{\delta-2}) \cup F(Y^{\delta-1}) = D \cup F(Y^{\delta-1}) = Y^\delta$ by Definition 2.1(b), induction hypothesis, 4.1.1 and monotony of F . If δ is the successor of a limit ordinal then Definition 2.1(b), induction hypothesis, 4.1.1, Definition 2.1(c) and definition of least upper bounds imply $X^\delta = X^{\delta-1} \cup F(X^{\delta-1}) = Y^{\delta-1} \cup F(Y^{\delta-1}) = \bigcup_{\alpha < \delta-1} Y^\alpha \cup F(Y^{\delta-1}) = \bigcup_{\alpha < \delta-1} (Y^{\alpha+1} \cup F(Y^{\delta-1})) = \bigcup_{\alpha < \delta-1} (D \cup F(Y^\alpha) \cup F(Y^{\delta-1})) = D \cup F(Y^{\delta-1}) = Y^\delta$. If δ is a limit ordinal then Definition 2.1(c) and induction hypothesis imply $X^\delta = \bigcup_{\alpha < \delta} X^\alpha = \bigcup_{\alpha < \delta} Y^\alpha = Y^\delta$. By transfinite induction the lemma is true for every $\delta \in \mu$.

4.1.3. By 4.1.1 and 4.1.2 the limits $\text{luis}(\lambda X \cdot X \cup F(X))(D)$ and $\text{luis}(\lambda X \cdot D \cup F(X))(D)$ exist and are equal. By 3.3 $\text{luis}(\lambda X \cdot X \cup F(X))(D)$ is the least of the fixed points of $\lambda X \cdot X \cup F(X)$ greater than or equal to D so that $\{\forall P \in L, \{P = P \cup F(P)\} \Leftrightarrow \{F(P) \subseteq P\}\}$ implies that $\text{luis}(\lambda X \cdot X \cup F(X))(D)$ and $\text{luis}(\lambda X \cdot D \cup F(X))(D)$ are equal to the least of the postfixed points of F greater than or equal to D .

COROLLARY 4.2. *The set of postfixed points of F is a nonempty complete lattice:*

$$\text{postfp}(F)(\subseteq, \text{lfp}(F), \top, \lambda S \cdot \text{luis}(\lambda Z \cdot Z \cup F(Z))(US), \cap)$$

where the least fixed point of F is $\text{lfp}(F) = \text{luis}(F)(D) = \cap \{X \in L: F(X) \subseteq X\}$ for every $D \in L$ such that $D \subseteq \text{lfp}(F)$.

Proof. By 4.1 and 3.4 the image of the nonempty complete lattice L by the upper closure operator $\bar{\rho} = \text{luis}(\lambda Z \cdot Z \cup F(Z))$ is included in $\text{postfp}(F)$. Reciprocally, $\forall P \in \text{postfp}(F)$ we know that $P \in \text{fp}(\lambda Z \cdot Z \cup F(Z))$ so that the upper iteration sequence $\langle X^\delta, \delta \in \mu \rangle$ for $\lambda Z \cdot Z \cup F(Z)$ starting with P is such that $\{\forall \delta \in \mu, P = X^\delta\}$. Hence $\bar{\rho}(P) = P$ that is $\text{postfp}(F) \subseteq \bar{\rho}(L)$ and by antisymmetry we have $\text{postfp}(F) = \bar{\rho}(L)$.

By Ward [21]'s theorem $\bar{\rho}(L)$ is a nonempty complete lattice $(\subseteq, \bar{\rho}(\perp), \top, \lambda S \cdot \bar{\rho}(\cup S), \cap)$.

Also by 4.1 $\text{luis}(\lambda Z \cdot Z \cup F(Z))(\perp) = \text{luis}(\lambda Z \cdot \perp \cup F(Z))(\perp) = \text{luis}(F)(\perp) = \cap \text{postfp}(F)$ by definition of the infimum of a complete lattice. By 3.3 $\text{luis}(F)(\perp)$ is the least of the fixed points of F greater than or equal to \perp , therefore it is the least fixed point of F .

Finally let $D \in L$, be such that $D \subseteq \text{lf}p(F)$ and $\langle X^\delta, \delta \in \mu \rangle$, $\langle Y^\delta, \delta \in \mu \rangle$, $\langle Z^\delta, \delta \in \mu \rangle$ be the upper iteration sequences for F respectively starting with \perp , D , and $\text{lf}p(F)$. By transfinite induction it is immediate that $\{\forall \delta \in \mu, X^\delta \subseteq Y^\delta \subseteq Z^\delta = \text{lf}p(F)\}$. According to 3.3, $\langle X^\delta, \delta \in \mu \rangle$ is stationary and its limit $\text{luis}(F)(\perp)$ is $\text{lf}p(F)$. Therefore $\langle Y^\delta, \delta \in \mu \rangle$ is stationary of limit $\text{lf}p(F)$.

Applying the duality principle, we obtain:

COROLLARY 4.3. *The set of prefixed points of F is a nonempty complete lattice:*

$$\text{prefp}(F)(\subseteq, \perp, \text{gfp}(F), \cup, \lambda S \cdot \text{llis}(\lambda Z \cdot Z \cap F(Z))(\cap S))$$

where the greatest fixed point of F is $\text{gfp}(F) = \text{llis}(F)(D) = \cup \{X \in L : X \subseteq F(X)\}$ for every $D \in L$ such that $\text{gfp}(F) \subseteq D$.

Let $\{F_i : i \in I\}$ be a family of monotone maps from L into L . The *unary polynomials* of the algebra $\langle L; \cup, \cap, \{F_i : i \in I\} \rangle$ are mappings on L into L defined as follows:

- (i) The identity mapping $\lambda X \cdot X$ is a unary polynomial.
- (ii) For every $i \in I$, if P is an unary polynomial then so is $\lambda X \cdot F_i(P(X))$.
- (iii) If $\{P_\gamma : \gamma \in J\}$ is a family of unary polynomials then so are $\lambda X \cdot \bigcup_{\gamma \in J} P_\gamma(X)$ and $\lambda X \cdot \bigcap_{\gamma \in J} P_\gamma(X)$.
- (iv) Unary polynomials are those and only those which we get from (i), (ii), and (iii).

Since polynomials are functions of L into L they are ordered by the pointwise ordering $\{F \subseteq G\} \Leftrightarrow \{\forall X \in L, F(X) \subseteq G(X)\}$.

COROLLARY 4.4. *Every unary polynomial of $\langle L; \cup, \cap, \{F_i : i \in I\} \rangle$*

is less than or equal to $\lambda X \cdot \text{luis}(\lambda Z \cdot Z \cup (\bigcup_{i \in I} F_i(Z)))(X)$ and greater than or equal to $\lambda X \cdot \text{llis}(\lambda Z \cdot Z \cap (\bigcap_{i \in I} F_i(Z)))(X)$.

Proof. Let \bar{F} be $\lambda Z \cdot (Z \cup (\bigcup_{i \in I} F_i(Z)))$ and \underline{F} be $\lambda Z \cdot (Z \cap (\bigcap_{i \in I} F_i(Z)))$, \bar{F} and \underline{F} are monotone maps on L into L . The proof is by induction on the structure of unary polynomials:

(i) $\text{luis}(\bar{F})$ is extensive and $\text{llis}(\underline{F})$ is reductive so that for every X of L we have $\text{llis}(\underline{F})(X) \subseteq X \subseteq \text{luis}(\bar{F})(X)$.

(ii) Let P be a unary polynomial such that for every X of L we have $\text{llis}(\underline{F})(X) \subseteq P(X) \subseteq \text{luis}(\bar{F})(X)$. Then for every monotone F_i , we have $F_i(\text{llis}(\underline{F})(X)) \subseteq F_i(P(X)) \subseteq F_i(\text{luis}(\bar{F})(X))$. But $\text{llis}(\underline{F})(X) = \underline{F}(\text{llis}(\underline{F})(X)) \subseteq F_i(\text{llis}(\underline{F})(X))$ and dually $F_i(\text{luis}(\bar{F})(X)) \subseteq \bar{F}(\text{luis}(\bar{F})(X)) = \text{luis}(\bar{F})(X)$ so that by transitivity $\text{llis}(\underline{F})(X) \subseteq F_i(P(X)) \subseteq \text{luis}(\bar{F})(X)$.

(iii) Let $\{P_\gamma : \gamma \in J\}$ be a family of unary polynomials such that for every $X \in L$, $\text{llis}(\underline{F})(X) \subseteq P_\gamma(X) \subseteq \text{luis}(\bar{F})(X)$ then by definition of least upper bounds we have $\text{llis}(\underline{F})(X) \subseteq \bigcup_{\gamma \in J} P_\gamma(X) \subseteq \text{luis}(\bar{F})(X)$ and by definition of greatest lower bounds we have $\text{llis}(\underline{F})(X) \subseteq \bigcap_{\gamma \in J} P_\gamma(X) \subseteq \text{luis}(\bar{F})(X)$.

The generalization of 4.4 to n -ary polynomials is immediate.

5. Constructive characterization of the set of fixed points of F .

THEOREM 5.1. (*Constructive version of Tarski's lattice theoretical fixed point theorem.*) *The set of fixed points of F is a nonempty complete lattice with ordering \subseteq , infimum $\text{luis}(F)(\perp)$, supremum $\text{llis}(F)(\top)$, least upper bound $\lambda S \cdot \text{luis}(F)(\cup S)$ and greatest lower bound $\lambda S \cdot \text{llis}(F)(\cap S)$.*

Proof. By Theorems 3.3 and 3.4, $\text{fp}(F)$ is the image of $\text{prefp}(F)$ by the upper closure operator $\text{luis}(F)$ and by Theorem 4.3 $\text{prefp}(F)$ is a nonempty complete lattice so that by Ward [21]'s theorem $\text{fp}(F)$ is a nonempty complete lattice with ordering \subseteq , infimum $\text{luis}(F)(\perp)$ and least upper bound $\lambda S \cdot \text{luis}(F)(\cup S)$. By duality, $\text{fp}(F)$ is the image of the nonempty complete lattice $\text{postfp}(F)$ by the lower closure operator $\text{llis}(F)$ so that the supremum of F is $\text{llis}(F)(\top)$ and the greatest lower bound $\lambda S \cdot \text{llis}(F)(\cap S)$.

The construction of extremal fixed points of monotone operators as limits of stationary transfinite iteration sequences may be found in Devidé [7] (where $\text{lfp}(\lambda Z \cdot D \cup F(Z))$ is the limit of the sequence $X^0 = D$, $X^\delta = X^{\delta-1} \cup F(X^{\delta-1})$ for successor ordinals and $X^\delta = \bigcup_{\alpha < \delta} X^\alpha$

for limit ordinals) in Hitchcock and Park [8] (where $lfp(F)$ is the limit of $X^0 = \perp$, $X^\beta = \bigcup_{\alpha < \beta} F(X^\alpha)$ for every nonzero ordinal) and in Pasini [15] (where transfinite sequences are defined as in Definition 2.1).

COROLLARY 5.2. *Let D be an arbitrary element of L . $luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))(D)$ and $llis(F) \circ luis(\lambda Z \cdot Z \cup F(Z))(D)$ are fixed points of F greater than or equal to any fixed point of F less than or equal to D and less than or equal to any fixed point of F greater than or equal to D . Moreover $luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))(D) \subseteq llis(F) \circ luis(\lambda Z \cdot Z \cup F(Z))(D)$.*

Proof. Assume that A is a fixed point of F less than or equal to D and B a fixed point of F greater than or equal to D , that is $F(A) = A \subseteq D \subseteq B = F(B)$. Then by monotony (3.4, 3.5) and fixed point property $A = luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))(A) \subseteq luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))(D) \subseteq luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))(B) = B$. The same way, $A \subseteq llis(F) \circ luis(\lambda Z \cdot Z \cup F(Z))(D) \subseteq B$.

Let P be $llis(\lambda Z \cdot Z \cap F(Z))(D)$ and Q be $luis(\lambda Z \cdot Z \cup F(Z))(D)$. Let S be $\{X \in L : P \subseteq X \subseteq Q\}$. S is a complete sublattice of L with infimum P and supremum Q . By 4.1 and its dual $P \subseteq F(P)$ and $F(Q) \subseteq Q$ so that $F(S) \subseteq S$. Then by 5.1 the least fixed point of F restricted to S is $luis(F)(P)$ and the greatest fixed point of F restricted to S is $llis(F)(Q)$ proving that $luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))(D) \subseteq llis(F) \circ luis(\lambda Z \cdot Z \cup F(Z))(D)$.

A lower preclosure operator ρ on L is monotone, idempotent and satisfies the lower connectivity axiom $\{\forall X \in L, \rho(X \cap \rho(X)) = \rho(X)\}$. An upper preclosure operator $\bar{\rho}$ on L is monotone, idempotent and satisfies the upper connectivity axiom $\{\forall X \in L, \bar{\rho}(X \cup \bar{\rho}(X)) = \bar{\rho}(X)\}$.

COROLLARY 5.3. *The set $fp(F)$ of fixed points of F is the image of L by the lower preclosure operator $luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))$ and the image of L by the upper preclosure operator $llis(F) \circ luis(\lambda Z \cdot Z \cup F(Z))$.*

Proof. $luis(F) \circ llis(\lambda Z \cdot Z \cap F(Z))$ is a lower preclosure operator since it is the composition of the upper closure operator $luis(F)$ and the lower closure operator $llis(\lambda Z \cdot Z \cap F(Z))$ (3.4, 4.1 and 3.5, Ladegaillierie [12]). By duality $llis(F) \circ luis(\lambda Z \cdot Z \cup F(Z))$ is an upper preclosure operator.

Cousot and Cousot [5] already used the idea of constructing (or approximating) the fixed points of monotone operators by means of

an upper iteration sequence followed by a lower iteration sequence. This idea was also used by Manna and Shamir [13] and our results 3.3, 4.1, 5.2, and 5.3 improve their results obtained on the more restricted model of continuous functional equations on functions of flat lower semi-lattices.

6. Constructive characterization of the set of fixed points of a family of commuting operators.

LEMMA 6.1. *Let \underline{F} and \bar{F} be monotone operators on the non-empty complete lattice $L(\subseteq, \perp, \top, \cup, \cap)$ into itself such that $\bar{F} \circ \underline{F} \subseteq \underline{F} \circ \bar{F}$ and $\underline{F} \subseteq \bar{F}$ (i.e., $\forall X \in L, \bar{F}(\underline{F}(X)) \subseteq \underline{F}(\bar{F}(X))$ and $\underline{F}(X) \subseteq \bar{F}(X)$). The set of common fixed points of \bar{F} and \underline{F} is a nonempty complete lattice:*

$fp(\underline{F}, \bar{F})(\subseteq, lfp(\bar{F}), gfp(\underline{F}), \lambda S \cdot luis(\bar{F})(\cup S), \lambda S \cdot llis(\underline{F})(\cap S))$ which is the image of L by $luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cap \underline{F}(Z))$ and the image of L by $llis(\underline{F}) \circ luis(\lambda Z \cdot Z \cup \bar{F}(Z))$.

Proof.

6.1.1. $\forall D \in pref\,fp(\underline{F}), \underline{F}(luis(\bar{F})(D)) = luis(\bar{F})(D)$.

Since $D \in pref\,fp(\underline{F})$ and $\underline{F} \subseteq \bar{F}$ we have $D \subseteq \underline{F}(D) \subseteq \bar{F}(D)$ so that the upper iteration sequence $\langle X^\delta, \delta \in \mu \rangle$ for \bar{F} starting with D is stationary, its limit $luis(\bar{F})(D)$ is a fixed point of \bar{F} (3.3). Again since $\underline{F} \subseteq \bar{F}$ we have $\underline{F}(luis(\bar{F})(D)) \subseteq \bar{F}(luis(\bar{F})(D)) = luis(\bar{F})(D)$.

Let us show that $\{\forall \delta \in \mu, X^\delta \subseteq \underline{F}(X^\delta)\}$. For $\delta = 0$ we have $X^0 = D \subseteq \underline{F}(X^0)$ since $D \in pref\,fp(\underline{F})$. Assume that the lemma is true for all $\alpha < \delta < \mu$. If δ is a successor ordinal then in particular $X^{\delta-1} \subseteq \underline{F}(X^{\delta-1})$. Since \bar{F} is monotone and $\bar{F} \circ \underline{F} \subseteq \underline{F} \circ \bar{F}$ we have by Definition 2.1(b), $X^\delta = \bar{F}(X^{\delta-1}) \subseteq \bar{F}(\underline{F}(X^{\delta-1})) \subseteq \underline{F}(\bar{F}(X^{\delta-1})) = \underline{F}(X^\delta)$. If δ is a limit ordinal then $X^\alpha \subseteq \underline{F}(X^\alpha)$ for every $\alpha < \delta$. By 2.1(c) and monotony, $X^\delta = \bigcup_{\alpha < \delta} X^\alpha \subseteq \bigcup_{\alpha < \delta} \underline{F}(X^\alpha) \subseteq \underline{F}(\bigcup_{\alpha < \delta} X^\alpha) = \underline{F}(X^\delta)$. By transfinite induction the lemma is true for every $\delta \in \mu$.

By 3.3, $luis(\bar{F})(D)$ is the limit of $\langle X^\delta, \delta \in \mu \rangle$ so that $luis(\bar{F})(D) \subseteq \underline{F}(luis(\bar{F})(D))$. By antisymmetry we conclude that $luis(\bar{F})(D) = \underline{F}(luis(\bar{F})(D))$.

6.1.2. Let D be an arbitrary element of L , then by the dual of Theorem 4.1, $llis(\lambda Z \cdot Z \cap \underline{F}(Z))(D) \in pref\,fp(\underline{F}) \subseteq pref\,fp(\bar{F})$ so that Theorem 3.3 implies that $luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cup \bar{F}(Z))(D) \in fp(\bar{F})$. Also by 6.1.1 $luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cap \underline{F}(Z))(D) \in fp(\underline{F})$. Consequently $luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cap \underline{F}(Z))(L) \subseteq fp(\underline{F}) \cap fp(\bar{F}) = fp(\underline{F}, \bar{F})$ and $fp(\underline{F}, \bar{F})$ is not empty (take D equal to \perp).

Let $P \in fp(\underline{F}, \bar{F})$ then $P \in L$ and $luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cap \underline{F}(Z))(P)$ is equal to P since $\underline{F}(P) = P$ and $\bar{F}(P) = P$. Therefore $fp(\underline{F}, \bar{F}) \subseteq luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cap \underline{F}(Z))(L)$ so that by antisymmetry we conclude

$fp(\underline{F}, \bar{F}) = luis(\bar{F}) \circ llis(\lambda Z \cdot Z \cap \underline{F}(Z))(L)$.

6.1.3. By 4.3 $llis(\lambda Z \cdot Z \cap \underline{F}(Z))(L)$ is a nonempty complete lattice $prefp(\underline{F})(\subseteq, \perp, gfp(\underline{F}), \cup, \lambda S \cdot llis(\lambda Z \cdot Z \cap \underline{F}(Z))(\cap S))$. By 3.4 $luis(\bar{F})$ is an upper closure operator so that by 6.1.2 and Ward [21]'s theorem $fp(\underline{F}, \bar{F})$ is a nonempty complete lattice with ordering \subseteq , infimum $luis(\bar{F})(\perp) = lfp(\bar{F})$ and least upper bound operation $\lambda S \cdot luis(\bar{F})(\cup S)$.

The remaining parts of Lemma 6.1 are obtained by duality, $fp(\underline{F}, \bar{F})$ is the image of the nonempty complete lattice $postfp(\bar{F})(\subseteq, lfp(\bar{F}), \top, \lambda S \cdot luis(\lambda Z \cdot Z \cup \bar{F}(Z))(\cup S), \cap)$ by the lower closure operation $llis(\underline{F})$ so that the supremum of $fp(\underline{F}, \bar{F})$ is $llis(\underline{F})(\top) = gfp(\underline{F})$ and the greatest lower bound operation is $\lambda S \cdot llis(\underline{F})(\cap S)$.

THEOREM 6.2. (*Constructive version of Tarski's generalized lattice theoretical fixed point theorem.*) *Let $\{F_i: i \in I\}$ be a nonempty family of monotone commuting operators on the nonempty complete lattice $L(\subseteq, \perp, \top, \cup, \cap)$ into itself. The set of all common fixed points $fp(\{F_i: i \in I\})$ of all the operators $\{F_i: i \in I\}$ is a nonempty complete lattice with ordering \subseteq , infimum $lfp(\lambda Z \cdot \bigcup_{i \in I} F_i(Z))$, supremum $gfp(\lambda Z \cdot \bigcap_{i \in I} F_i(Z))$, least upper bound operation $\lambda S \cdot luis(\lambda Z \cdot \bigcup_{i \in I} F_i(Z))(\cup S)$ and greatest lower bound operation $\lambda S \cdot llis(\lambda Z \cdot \bigcap_{i \in I} F_i(Z))(\cap S)$.*

Proof.

6.2.1. Let \bar{F} be $\lambda Z \cdot \bigcup_{i \in I} F_i(Z)$ and \underline{F} be $\lambda Z \cdot \bigcap_{i \in I} F_i(Z)$. \bar{F} and \underline{F} are monotone operators on L into itself such that $\underline{F} \subseteq \bar{F}$. $\forall X \in L, \forall i \in I$, we have $\bar{F}(F_i(X)) = \bigcup_{j \in I} F_j(F_i(X)) = \bigcup_{j \in I} F_i(F_j(X)) \subseteq F_i(\bigcup_{j \in I} F_j(X)) = F_i(\bar{F}(X))$ by monotony and the commuting property. Therefore $\forall X \in L, \bar{F}(\underline{F}(X)) = \bar{F}(\bigcap_{i \in I} F_i(X)) \subseteq \bigcap_{i \in I} \bar{F}(F_i(X)) \subseteq \bigcap_{i \in I} F_i(\bar{F}(X)) = \underline{F}(\bar{F}(X))$.

6.2.2. Clearly $fp(\{F_i: i \in I\}) \subseteq fp(\bar{F}, \underline{F})$ since $\{\forall i \in I, F_i(X) = X\}$ implies $\bar{F}(X) = \bigcup_{i \in I} F_i(X) = \bigcup_{i \in I} X = X$ and dually $\underline{F}(X) = X$. Whenever $X \in fp(\bar{F}, \underline{F})$ we have $\forall i \in I, X = \underline{F}(X) = \bigcap_{j \in I} F_j(X) \subseteq F_i(X)$ and dually $F_i(X) \subseteq \bigcup_{j \in I} F_j(X) = \bar{F}(X) = X$ so that by antisymmetry $X = F_i(X)$ and $fp(\bar{F}, \underline{F}) \subseteq fp(\{F_i: i \in I\})$. By antisymmetry $fp(\bar{F}, \underline{F}) = fp(\{F_i: i \in I\})$ so that by Lemma 6.1, $fp(\{F_i: i \in I\})$ is a complete lattice $(\subseteq, lfp(\bar{F}), gfp(\underline{F}), \lambda S \cdot luis(\bar{F})(\cup S), \lambda S \cdot llis(\underline{F})(\cap S))$.

COROLLARY 6.3. *Let D be an arbitrary element of L , then $luis(\lambda Z \cdot \bigcup_{i \in I} F_i(Z)) \circ llis(\lambda Z \cdot Z \cap (\bigcap_{i \in I} F_i(Z)))(D)$ and $llis(\lambda Z \cdot \bigcap_{i \in I} F_i(Z)) \circ luis(\lambda Z \cdot Z \cup (\bigcup_{i \in I} F_i(Z)))(D)$ are common fixed points of the $F_i, i \in I$ which are greater than or equal to any common fixed point of the F_i less than or equal to D and which are less than or*

equal to any common fixed point of the F_i greater than or equal to D . Moreover

$$\begin{aligned} & \text{luis}(\lambda Z \cdot \bigcup_{i \in I} F_i(Z)) \circ \text{llis}(\lambda Z \cdot Z \cap (\bigcap_{i \in I} F_i(Z)))(D) \\ & \subseteq \text{llis}(\lambda Z \cdot \bigcap_{i \in I} F_i(Z)) \circ \text{luis}(\lambda Z \cdot Z \cup (\bigcup_{i \in I} F_i(Z)))(D). \end{aligned}$$

COROLLARY 6.4. *The set $\text{fp}(\{F_i: i \in I\})$ of common fixed points of the family $\{F_i: i \in I\}$ is the image of L by the lower preclosure operator $\text{luis}(\lambda Z \cdot \bigcup_{i \in I} F_i(Z)) \circ \text{llis}(\lambda Z \cdot Z \cap (\bigcap_{i \in I} F_i(Z)))$ and the image of L by the upper preclosure operator $\text{llis}(\lambda Z \cdot \bigcap_{i \in I} F_i(Z)) \circ \text{luis}(\lambda Z \cdot Z \cup (\bigcup_{i \in I} F_i(Z)))$.*

Let $\{F_i: i \in I\}$ be a finite family of monotone commuting operators on the complete lattice L into itself. If we assume that I is well-ordered (i.e., $I = \{i_\alpha: \alpha \leq \gamma\}$ where $\gamma \in \omega$) then we denote $\lambda Z \cdot F_{i_0}(F_{i_1}(\dots F_{i_\gamma}(Z) \dots))$ by $\bigcirc_{i \in I} F_i$.

Applying Theorem 5.1 to $\bigcirc_{i \in I} F_i$ and Theorem 6.2 to $\{F_i: i \in I\}$ a natural question is whether $\text{fp}(\bigcirc_{i \in I} F_i) = \text{fp}(\{F_i: i \in I\})$. The answer is affirmative thanks to the following:

THEOREM 6.5.

$$\begin{aligned} & \text{luis}(\lambda Z \cdot \bigcup_{i \in I} F_i(Z)) \circ \text{llis}(\lambda Z \cdot Z \cap (\bigcap_{i \in I} F_i(Z))) \\ & = \text{luis}(\bigcirc_{i \in I} F_i) \circ \text{llis}(\lambda Z \cdot Z \cap (\bigcirc_{i \in I} F_i)(Z)) \\ & \text{llis}(\lambda Z \cdot \bigcap_{i \in I} F_i(Z)) \circ \text{luis}(\lambda Z \cdot Z \cup (\bigcup_{i \in I} F_i(Z))) \\ & = \text{llis}(\bigcirc_{i \in I} F_i) \circ \text{luis}(\lambda Z \cdot Z \cup (\bigcirc_{i \in I} F_i)(Z)). \end{aligned}$$

Proof. It is sufficient to prove that if D is a prefixed point of each F_i such that $i \in I$ then $\text{luis}(\lambda Z \cdot \bigcup_{i \in I} F_i(Z))(D) = \text{luis}(\bigcirc_{i \in I} F_i)(D)$. Since $\{\forall i \in I, D \subseteq F_i(D)\}$ we have by monotony and the commuting property $D \subseteq (\bigcirc_{i \in I} F_i)(D)$ and Theorem 3.3 implies that $P = \text{luis}(\bigcirc_{i \in I} F_i)(D) = (\bigcirc_{i \in I} F_i)(P)$ and $D \subseteq P$. For every $j \in I$ we have $D \subseteq F_j(D) \subseteq F_j(P) = F_j((\bigcirc_{i \in I} F_i)(P)) = (\bigcirc_{i \in I} F_i)(F_j(P))$. Therefore $F_j(P)$ is a fixed point of $\bigcirc_{i \in I} F_i$ greater than or equal to D so that by Theorem 3.3 $\{\forall j \in I, P \subseteq F_j(P)\}$. Then by monotony and transitivity $P \subseteq F_{i_1}(P) \subseteq F_{i_1}(\dots F_{i_\gamma}(P) \dots) \subseteq (\bigcirc_{i \in I} F_i)(P) = P$ so that $\forall j \in I, P = F_j(P)$. P is a common fixed point of the family $\{F_i: i \in I\}$ greater than or equal to D . Let Q be another common fixed point of $\{F_i: i \in I\}$ greater than or equal to D . Then $(\bigcirc_{i \in I} F_i)(Q) = Q$ so that by Theorem 3.3 we have $P \subseteq Q$. Hence P is the least common fixed point of the family $\{F_i: i \in I\}$ greater than or equal to D .

By Corollary 6.3, $R = \text{luis}(\lambda Z \cdot \bigcup_{i \in I} F_i(Z))(D)$ is a common fixed point of $\{F_i; i \in I\}$ greater than or equal to D . Let Q be another common fixed point of $\{F_i; i \in I\}$ greater than or equal to D . Then $\bigcup_{i \in I} F_i(Q) = Q$ so that by Theorem 3.3 we have $R \subseteq Q$. Hence R is the least common fixed point of the family $\{F_i; i \in I\}$ greater than or equal to D .

By existence and unicity of the least common fixed point of the family $\{F_i; i \in I\}$ greater than or equal to D , we conclude $\text{luis}(\bigcirc_{i \in I} F_i)(D) = P = R = \text{luis}(\lambda Z \cdot \bigcup_{i \in I} F_i(Z))(D)$.

7. Fixed point theorems for continuous operators. An operator F on the complete lattice L into itself is *upper-semi-continuous* if and only if for every ordinal $\delta \leq \omega$ and every δ -termed increasing chain $\langle C^\alpha, \alpha \in \delta \rangle$ of elements of L we have $F(\bigcup_{\alpha \in \delta} C^\alpha) = \bigcup_{\alpha \in \delta} F(C^\alpha)$. The dual notion is the one of *lower-semi-continuous* operator. An operator is *continuous* when it is lower and upper-semi-continuous.

Since semi-continuity implies monotony the results of paragraphs 3, 4, and 5 can be applied to continuous operators. However the proofs are simplified since one can consider $(\omega + 1)$ -termed iteration sequences. For example, Theorem 3.3 can be reformulated as follows:

THEOREM 7.1. *Let F be an upper-semi-continuous operator on the complete lattice L into itself. An upper iteration sequence $\langle X^\delta, \delta \in \min(\mu, \omega + 1) \rangle$ for F starting with $D \in \text{prefp}(F)$ is a stationary increasing chain, its limit $\text{luis}(F)(D)$ is the least of the fixed points of F greater than or equal to D .*

Proof. When $\mu > \omega + 1$ Definition 2.1, Theorem 3.3 and upper-semi-continuity imply $X^{\omega+1} = F(X^\omega) = F(\bigcup_{\alpha < \omega} X^\alpha) = \bigcup_{\alpha < \omega} F(X^\alpha) = \bigcup_{\alpha < \omega} X^{\alpha+1} \subseteq \bigcup_{\alpha < \omega} X^\alpha = X^\omega$. Also by Theorem 3.3, $X^\omega \subseteq X^{\omega+1}$ so that by antisymmetry $X^\omega = X^{\omega+1}$. Then by transfinite induction it is easy to show that $\{\forall \beta: \omega \leq \beta < \mu, X^\omega = X^\beta\}$.

When considering a family of commuting monotone operators the results of paragraph 6 can be perfected as follows:

LEMMA 7.2. *Let \underline{F} and \bar{F} be upper-semi-continuous operators on the complete lattice L into itself such that $\bar{F} \circ \underline{F} \subseteq \underline{F} \circ \bar{F}$ and $\underline{F} \subseteq \bar{F}$. Then for every prefixed point D of F we have:*

$$\{\underline{F}(D) = \bar{F}(D)\} \implies \{\text{luis}(\underline{F})(D) = \text{luis}(\bar{F})(D)\}.$$

Proof. Let $\langle X^\delta, \delta \in \min(\omega + 1, \mu) \rangle$ and $\langle Y^\delta, \delta \in \min(\omega + 1, \mu) \rangle$ be respectively the upper iteration sequences for \underline{F} and \bar{F} starting with the prefixed point D of \underline{F} and \bar{F} .

For $\delta = 0$ we know by hypothesis and 2.1(a) that $D = X^0 = Y^0 \subseteq \bar{F}(X^0) = \underline{F}(Y^0)$.

Assume that $\delta \in \min(\omega + 1, \mu)$ is a successor ordinal such that $X^{\delta-1} = Y^{\delta-1}$ and $\bar{F}(X^{\delta-1}) = \underline{F}(Y^{\delta-1})$. Then by 2.1(b) $X^\delta = \underline{F}(X^{\delta-1}) = \underline{F}(Y^{\delta-1})$ and $\bar{F}(X^{\delta-1}) = \bar{F}(Y^{\delta-1}) = Y^\delta$ so that by induction hypothesis and transitivity $X^\delta = Y^\delta$. Also since $\underline{F} \subseteq \bar{F}$ and $X^\delta = Y^\delta$ we know that $\underline{F}(Y^\delta) \subseteq \bar{F}(X^\delta)$. Since $\bar{F} \circ \underline{F} \subseteq \underline{F} \circ \bar{F}$ and $X^{\delta-1} = Y^{\delta-1}$ we know that $\bar{F}(\underline{F}(X^{\delta-1})) \subseteq \underline{F}(\bar{F}(Y^{\delta-1}))$ so that by Definition 2.1(b) we get $\bar{F}(X^\delta) \subseteq \underline{F}(Y^\delta)$. By antisymmetry we conclude $\bar{F}(X^\delta) = \underline{F}(Y^\delta)$.

Assume that $\delta \in \min(\omega + 1, \mu)$ is a limit ordinal then $\delta = \omega$. If by induction hypothesis $\{\forall \beta < \omega, X^\beta = Y^\beta \text{ and } \bar{F}(X^\beta) = \underline{F}(Y^\beta)\}$ then by 2.1(c) and definition of least upper bounds we have $X^\omega = \bigcup_{\alpha < \omega} X^\alpha = \bigcup_{\alpha < \omega} Y^\alpha = Y^\omega$. The same way by upper-semi-continuity, $\bar{F}(X^\omega) = \bar{F}(\bigcup_{\alpha < \omega} X^\alpha) = \bigcup_{\alpha < \omega} \bar{F}(X^\alpha) = \bigcup_{\alpha < \omega} \underline{F}(Y^\alpha) = \underline{F}(\bigcup_{\alpha < \omega} Y^\alpha) = \underline{F}(Y^\omega)$.

By transfinite induction and Theorem 7.1 we conclude $lwis(\underline{F})(D) = lwis(\bar{F})(D)$.

As an application of Lemma 7.2 for $D = \perp$, we get:

THEOREM 7.3. *Let $\{F_i; i \in I\}$ be a family of commuting operators on the complete lattice L into itself. Then $\{\{\forall i \in I, F_i \text{ is upper-semi-continuous}\} \text{ and } \{\forall i, j \in I, F_i(\perp) = F_j(\perp)\}\} \Rightarrow \{\forall i, j \in I, lfp(F_i) = lfp(F_j)\}$.*

8. Remark. In our proofs it is the existence of lower or upper bounds of chains and not the existence of lower or upper bounds of arbitrary sets that is crucial. The same remark was made by numerous authors who generalized Tarski's fixed point theorem to weaken the completeness hypothesis (see among others Abian and Brown [1], Höft [9], Pasini [15], Pelczar [16], Markowsky [14], Ward [20], Wolk [22]). This was also the case for Tarski's fixed point theorem on commuting maps (see a.o., DeMarr [6], Markowsky [14], Pelczar [17], Smithson [18], Wong [23]). Along the same lines our results could be strengthened to be applicable to partially ordered sets which are not complete lattices.

ACKNOWLEDGMENT. The authors thank the referee in particular for strengthening the initial version of Theorem 4.1.

REFERENCES

1. S. Abian and A. B. Brown, *A theorem on partially ordered sets with applications to fixed point theorems*, *Canad. J. Math.*, **13** (1961), 78-82.
2. H. Amann, *Fixed point equations and non-linear eigenvalue problems in ordered Banach spaces*, *SIAM Review* 18, **4** (October 1976), 620-709.
3. G. Birkhoff, *Lattice theory*, Amer. Math. Soc. Colloquium Publications 25, 3rd ed., Providence, Rhode Island, (1976).
4. A. Church, *The calculi of lambda-conversion*, *Annals of Math. Studies*, **6** (1951), Princeton University Press, Princeton, N.J.
5. P. Cousot and R. Cousot, *Abstract interpretation: a unified lattice model for static analysis of programs by construction or approximation of fixpoints*, *Conf. Rec. of the Fourth ACM Symp. on Principles of Programming Languages*, Los Angeles, Calif. (Jan. 1977), 238-252.
6. R. DeMarr, *Common fixed points for isotone mappings*, *Colloquium Math.*, **13** (1964), 45-48.
7. V. Devidé, *On monotonous mappings of complete lattices*, *Fundamenta Mathematicae*, **LIII** (1964), 147-154.
8. P. Hitchcock and D. Park, *Induction rules and proofs of termination*, *Proc. of a Symp. on Automata, Languages and Programming*, North-Holland Pub. Co., Amsterdam, (1973), 225-251.
9. H. Höft and M. Höft, *Some fixed point theorems for partially ordered sets*, *Canad. J. Math.*, **5** (1976), 992-997.
10. S. C. Kleene, *Introduction to metamathematics*, North-Holland Pub. Co., Amsterdam, (1952).
11. I. I. Kolodner, *On completeness of partially ordered sets and fixpoints theorems for isotone mappings*, *Amer. Math. Monthly*, **75** (1968), 48-49.
12. Y. Ladegallerie, *Prefermeture sur un ensemble ordonné*, *RAIRO*, **1** (Avril 1973), 35-43.
13. Z. Manna and A. Shamir, *The convergence of functions to fixed points of recursive definitions*, *Theoretical Computer Science*, **6** (1978), 109-141.
14. G. Markowsky, *Chain-complete posets and directed sets with applications*, *Algebra Univ.*, **6** (1976), 53-58, Birkhäuser Verlag, Basel.
15. A. Pasini, *Some fixed point theorems of the mappings of partially ordered sets*, *Rend. Sem. Mat. Univ. Padova*, **51** (1974), 167-177.
16. A. Pelczar, *On the invariant points of a transformation*, *Ann. Polon. Math.*, **11** (1961), 199-202.
17. ———, *Remarks on commutings mappings in partially ordered spaces*, *Zeszyty Nauk. Univ. Jagiello.*, *Prace Mat. Zeszyt*, **15** (1971), 131-133.
18. R. E. Smithson, *Fixed points in partially ordered sets*, *Pacific J. Math.*, **1** (1973), 363-367.
19. A. Tarski, *A lattice theoretical fixpoint theorem and its applications*, *Pacific J. Math.*, **5** (1955), 285-310.
20. L. E. Ward Jr., *Completeness in semi-lattices*, *Canad. J. Math.*, **9** (1957), 578-582.
21. M. Ward, *The closure operators of a lattice*, *Annals Math.*, **43** (1942), 191-196.
22. E. S. Wolk, *Dedekind completeness and a fixed point theorem*, *Canad. J. Math.*, **9** (1957), 400-405.
23. J. S. W. Wong, *Common fixed points of commuting monotone mapping*, *Canad. J. Math.*, **19** (1967), 617-620.

Received December 5, 1977 and in revised form June 26, 1978. Work of the first author was supported in part by CNRS, Laboratoire associé n°7 and ATP-Informatique D3119 and the second author in part by IRIA-SESORI under grant 76160.

LABORATOIRE I.M.A.G.
U. S. M. G., BP. 53X
38041 GRENOBLE CEDEX, FRANCE

