

## REMARK ON A PAPER OF STUX CONCERNING SQUAREFREE NUMBERS IN NON-LINEAR SEQUENCES

G. J. RIEGER

**Stux studied squarefree numbers of the form  $[f(n)]$ ; his most interesting application is  $f(n)=n^c$  for real  $c$  with  $1 < c < 4/3$ . We would like to point out that a stronger result follows immediately from estimates of Deshouillers.**

Let  $1 < c < 2$ ,  $x \geq 1$ ; denote by  $N_c(x; k, l)$  the number of natural numbers  $n \leq x$  with  $[n^c] \equiv 1 \pmod k$ . According to [1], we have

$$(1) \quad N_c(x; k, l) = \frac{x}{k} + O_c((x^{1+c}k^{-1})^{1/3}) \quad \text{for } x^{c-5/4} \leq k < x^{c-1/2},$$

$$(2) \quad N_c(x; k, l) = \frac{x}{k} + O_c((x^{4+c}k^{-1})^{1/7}) \quad \text{for } k < x^{c-5/4}.$$

Denote by  $S_c(x)$  the number of squarefree numbers of the form  $[n^c]$  with natural  $n \leq x$ ; the inclusion-exclusion principle in the form  $|\mu(n)| = \sum_{d^2 | n, d > 0} \mu(d)$  gives

$$(3) \quad S_c(x) = \sum_{d^2 \leq x^c} \mu(d) N_c(x; d^2, 0) \quad (x \geq 1).$$

For  $d^2 \geq x^{c-1/2}$  we use the trivial estimate  $N_c(x; d^2, 0) = O(x^c d^{-2})$ ; using

$$(4) \quad \sum_{d > t} d^{-2} = O(t^{-1}) \quad (t \geq 1),$$

we obtain

$$(5) \quad S_c(x) = \sum_{d^2 < x^{c-1/2}} \mu(d) N_c(x; d^2, 0) + O(x^{(2c+1)/4}).$$

In case  $c \leq 5/4$ , we use (1) and

$$(6) \quad \sum_{0 < d \leq t} d^{-2/3} = O(t^{1/3}) \quad (t \geq 1)$$

in (5); this gives

$$(7) \quad S_c(x) = \sum_{d^2 < x^{c-1/2}} \mu(d) d^{-2} x + O_c(x^{(2c+1)/4}).$$

In case  $c > 5/4$ , we split the sum in (5) according to  $d^2 < \text{or } \geq x^{c-5/4}$  and apply (2) and (1); using  $\sum_{0 < d \leq t} d^{-2/7} = O(t^{5/7})$  ( $t \geq 1$ ) and (6), we obtain again (7). But (7),  $\sum_{d > 0} \mu(d) d^{-2} = 6\pi^{-2}$ , and (4) give immediately

THEOREM 1. For real  $c$  with  $1 < c < 3/2$ , we have

$$S_c(x) = 6\pi^{-2}x + O_c(x^{(2c+1)/4}) \quad (x \geq 1).$$

Looking at  $m - [n^c]$  instead of  $[n^c]$  we obtain similarly

THEOREM 2. For real  $c$  with  $1 < c < 3/2$ , the number of representations of the natural number  $m$  as  $m = q + [n^c]$  with squarefree  $q$  and natural  $n$  equals

$$6\pi^{-2}m^{1/c} + O_c(m^{(2c+1)/4c}).$$

This can easily be generalized to  $r$ -free instead of squarefree. It should not be difficult to extend the method of [1] to cover the function class studied in [2].

#### REFERENCES

1. Jean-Marc Deshouillers, *Sur la répartition des nombres  $[n^c]$  dans les progressions arithmétiques*, C.R. Acad. Sci. Paris, **277** (1973), Ser. A., 647-650.
2. Ivan F. Stux, *Distribution of squarefree integers in non-linear sequences*, Pacific J. Math., **59** (1975), 577-584.

Received June 26, 1977 and in revised form December 16, 1977.

TECHNISCHE UNIVERSITÄT  
D-3000 HANNOVER, GERMANY