

## SEQUENCES OF BOUNDED SUMMABILITY DOMAINS

R. M. DEVOS AND F. W. HARTMANN

**C. Goffman and G. N. Wollan conjectured that the bounded summability field of a regular matrix  $A$  is so thin that the union of countably many such sets is not dense in  $m$ . G. M. Petersen proved this conjecture. This result is strengthened by showing if  $A$  is a noncoercive matrix whose summability field contains all the finite sequences then its bounded summability field is so thin that the union of countably many such sets is not dense in  $m$ . An example is given to show that the condition of containing the finite sequences is necessary.**

**Preliminaries.** Let  $m$  and  $c$  be respectively the Banach spaces of bounded and convergent sequences,  $x = \{x_n\}$ , of complex numbers with norm  $\|x\|_\infty = \sup_n |x_n|$ ,  $B(x, r) = \{z \in m: \|x + z\|_\infty < r\}$ . Denote the  $n$ th section of  $x$  by  $P_n(x) = (x_1, \dots, x_n, 0, 0, \dots)$ . For each infinite matrix  $A$  the set of  $x$  transformed by  $A$  to convergent sequences is called the summability field of  $A$  and denoted by  $c_A$ . The set of bounded sequences in  $c_A$  is called the bounded summability field of  $A$  and is denoted by  $\mathcal{A}$ .  $A$  is called conservative if and only if  $c_A \supset c$ , regular if and only if  $A$  is conservative and limits are preserved, coercive if and only if  $c_A \supset m$ . If  $A = (a_{nk})$ , then the  $A$  transform of  $x$  is designated by  $Ax = \{(Ax)_n\} = \{\sum_k a_{nk}x_k\}$ .  $A$  is conservative if and only if  $\|A\|_\infty = \sup_n \sum_k |a_{nk}| < \infty$ ,  $a_k = \lim_n a_{nk}$  exists for each  $k$  and  $\lim_n \sum_k a_{nk}$  exists [5, p. 165].  $A$  is coercive if and only if  $\sum_k |a_{nk}|$  converges uniformly in  $n$  and  $a_k$  exists for each  $k$  [5, p. 169]. Define the essential norm of  $A$  by  $\|A\|_c = \limsup_n \sum_k |a_{nk} - a_k|$  whenever  $a_k$  exists for each  $k$ . (Note  $\|\cdot\|_c$  is not a true norm, since  $\|\cdot\|_c$  may be infinite.)

Let  $E^\infty$  be the set of all finite sequences and  $N_0$  the set of all sequences of 0's and 1's. Using binary expansions there is a natural injective mapping of  $(0, 1)$  onto all but a countable subset of  $N_0$ .

**MAIN RESULTS.** C. Goffman and G. N. Wollan conjectured [4] that the bounded summability field of regular  $A$  is so thin that the union of countably many such sets is not dense in  $m$ . G. M. Petersen proved this conjecture [6]. We strengthen that result and show that in a certain sense our result is best possible.

**THEOREM.** *Let  $\{A_i\}$  be a countable collection of noncoercive matrices with  $\mathcal{A}_i \supset E^\infty$ ,  $i = 1, 2, \dots$ , then  $\bigcup_{i=1}^\infty \mathcal{A}_i$  is not dense in  $m$ .*

We prove the theorem through a series of lemmas. Since we

want  $E^\infty \subset \mathcal{A}$ , we shall assume all  $A$  in the sequel have convergent columns.

LEMMA 1. *Let  $\|A\|_\infty < \infty$  then  $\|A\|_c = 0$  if and only if  $A$  is coercive.*

*Proof.* Suppose  $A$  is coercive. Let  $\varepsilon > 0$ . There exists  $k_0$

$$\sum_{k=k_0+1}^\infty |a_{nk}| < \varepsilon/3$$

for all  $n$ . Since  $\{a_k\} \in \mathcal{C}^1$ , there is a  $k_1$  such that  $k > k_1$  implies

$$\sum_{k=k_1+1}^\infty |a_k| < \varepsilon/3.$$

Let  $k_2 = \max(k_1, k_0)$ . There exists  $n_0 = n_c(k_2)$  such that  $n > n_0$  implies

$$\sum_{k=1}^{k_2} |a_{nk} - a_k| < \varepsilon/3.$$

Let  $n > n_0$  then

$$\begin{aligned} \sum_{k=1}^\infty |a_{nk} - a_k| &= \sum_{k=1}^{k_2} |a_{nk} - a_k| + \sum_{k=k_2+1}^\infty |a_{nk} - a_k| \\ &\leq \sum_{k=1}^{k_2} |a_{nk} - a_k| + \sum_{k=k_2+1}^\infty |a_{nk}| + \sum_{k=k_2+1}^\infty |a_k| \\ &< \varepsilon/3 + \varepsilon/3 + \varepsilon/3 = \varepsilon. \end{aligned}$$

Conversely assume  $A$  is noncoercive. There exists  $\varepsilon > 0$  and an increasing sequence of positive integers  $\{n(p)\}_{p=1}^\infty$  such that  $\sum_{k=p+1}^\infty |a_{n(p),k}| > \varepsilon$ . There exists  $k_0$  such that  $\sum_{k=k_0+1}^\infty |a_k| < \varepsilon/2$ . Pick  $p$  with  $p > k_0$  then

$$\begin{aligned} \sum_{k=1}^\infty |a_{n(p),k} - a_k| &\geq \sum_{k=k_0+1}^\infty |a_{n(p),k} - a_k| \\ &\geq \sum_{k=k_0+1}^\infty |a_{n(p),k}| - \sum_{k=k_0+1}^\infty |a_k| \\ &\geq \varepsilon - \varepsilon/2 = \varepsilon/2. \end{aligned}$$

Therefore  $\|A\|_c > 0$ .

Let  $\Gamma(c, c)$  be the Banach algebra of conservative matrices and  $\mathcal{K}$  be the ideal of compact operators. It is well known [8] that  $A \in \mathcal{K}$  if and only if  $A$  is coercive.  $\Gamma(c, c)/\mathcal{K}$  is a Banach algebra and is called a Calkin algebra [2]. It is easily seen that  $\|\cdot\|_c$  is the norm in the Calkin algebra.

LEMMA 2. *Let  $\|A\|_c < \infty$  and  $a$  and  $b$  be cluster points of  $Ax$ ,*

$x \in m$ , then  $|a - b| \leq 2\|A\|_c \|x\|_\infty$ .

*Proof.* Let  $a$  and  $b$  be cluster points of  $Ax$  and  $\varepsilon > 0$ . There exist increasing sequences of positive integers  $\{n(i)\}$ ,  $\{m(j)\}$  and  $N_0$  such that for  $n(i), m(j) > N_0$

$$\left| \sum_k a_{n(i),k} x_k - a \right| < \varepsilon$$

and

$$\left| \sum_k a_{m(j),k} x_k - b \right| < \varepsilon.$$

There exists  $N_1$  such that  $n > N_1$  implies

$$\sum_k |a_{nk} - a_k| < \|A\|_c + \varepsilon.$$

Let  $n(i), m(j) > \max(N_0, N_1)$  then

$$\begin{aligned} |a - b| &\leq \left| \sum_k a_{n(i),k} x_k - \sum_k a_{m(j),k} x_k \right| + 2\varepsilon \\ &\leq \sum_k |a_{n(i),k} - a_{m(j),k}| |x_k| + 2\varepsilon \\ &\leq \|x\|_\infty \sum_k |(a_{n(i),k} - a_k) - (a_{m(j),k} - a_k)| + 2\varepsilon \\ &\leq \|x\|_\infty (\|A\|_c + \varepsilon + \|A\|_c + \varepsilon) + 2\varepsilon. \end{aligned}$$

Since  $\varepsilon$  is arbitrary the conclusion follows.

The next lemma is due to Bennett and Kalton and appears as Lemma 7 of [1, p. 577].

**LEMMA 3.** (*Bennett and Kalton*). *If  $z_1, z_2, \dots, z_n$  is any finite collection of complex numbers then there exists a subset  $J(n)$  of  $\{1, \dots, n\}$  such that*

$$\left| \sum_{j \in J(n)} z_j \right| \geq \frac{1}{4} \sum_{i=1}^n |z_i|.$$

**LEMMA 4.** *If  $\|A\| = \infty$ , then there exists  $E(A)$  with  $E(A) \subset N_0$ ,  $N_0 \setminus E(A)$  of first category and if  $u \in E(A)$  then  $B(u, 1/32) \cap \mathcal{A} = \emptyset$ .*

*Proof.* **Case 1.** Assume all the rows of  $A$  are in  $\ell^1$ . Let  $\|A\| = \infty$ . Pick sequences  $n(k)$  and  $q(k)$  inductively such that  $n(1) = 1$  and

- (i)  $\sum_{i=q(k)+1}^\infty |a_{n(k),i}| < 2^{-k}$
- (ii)  $\sum_{i=q(k-1)+1}^{q(k)} |a_{n(k),i}| > (65/7) \sup_j \{ \sum_{i=1}^{q(k-1)} |a_{ji}| \}$ .

By Lemma 3 select  $J(k) \subset \{q(k-1) + 1, \dots, q(k)\}$  with

$$\left| \sum_{i \in J(k)} a_{n^{(k)}, i} \right| \geq \frac{1}{4} \sum_{i=q^{(k-1)+1}}^{q^{(k)}} |a_{n^{(k)}, i}|.$$

For each natural number  $k$  define the sequence  $u^k$  by  $u_i^k = 1$  if  $i \in J(k)$ ,  $u_i^k = 0$  if  $i \notin J(k)$ . Let

$$O_k = \{u \in N_0 : (P_{q^{(k)}} - P_{q^{(k-1)}})(u - u^k) = 0\}.$$

If  $E(A) = \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} O_k$ , then  $E(A)$  is of second category. [ $\bigcup_{k=n}^{\infty} O_k$  is open and dense, hence by the Baire theorem  $E(A)$  is of second category.] Let  $u \in E(A)$  and  $\|z\|_{\infty} < 1/32$ .  $u$  is in an infinite number of the  $O_k$ . Let  $u \in O_r$ . Then

$$\begin{aligned} |(A(u+z))_{n^{(r)}}| &\geq |(Au)_{n^{(r)}}| - |(Az)_{n^{(r)}}| \\ &\geq \left| \sum_{i=q^{(r-1)+1}}^{q^{(r)}} a_{n^{(r)}, i} u_i \right| - \left| \sum_{i=1}^{q^{(r-1)}} a_{n^{(r)}, i} u_i \right| - \left| \sum_{i=q^{(r)+1}}^{\infty} a_{n^{(r)}, i} u_i \right| \\ &\quad - \frac{1}{32} \sum_{i=1}^{\infty} |a_{n^{(r)}, i}| \\ &\geq \frac{1}{4} \sum_{i=q^{(r-1)+1}}^{q^{(r)}} |a_{n^{(r)}, i}| - \frac{33}{32} \sum_{i=1}^{q^{(r-1)}} |a_{n^{(r)}, i}| \\ &\quad - \frac{33}{32} \sum_{i=q^{(r)+1}}^{\infty} |a_{n^{(r)}, i}| - \frac{1}{32} \sum_{i=q^{(r-1)+1}}^{q^{(r)}} |a_{n^{(r)}, i}| \\ &\geq \frac{7}{32} \sum_{i=q^{(r-1)+1}}^{q^{(r)}} |a_{n^{(r)}, i}| - \frac{33}{32} \sum_{i=1}^{q^{(r-1)}} |a_{n^{(r)}, i}| - \frac{33}{32} 2^{-r} \\ &\geq \frac{7}{32} \frac{65}{7} \sup_j \left\{ \sum_{i=1}^{q^{(r-1)}} |a_{ji}| \right\} - \frac{33}{32} \sup_j \left\{ \sum_{i=1}^{q^{(r-1)}} |a_{ji}| \right\} - 2^{1-r} \\ &\geq \sup_j \left\{ \sum_{i=1}^{q^{(r-1)}} |a_{ji}| \right\} - 2^{1-r} \longrightarrow \infty \text{ as } r \longrightarrow \infty. \end{aligned}$$

Hence the  $A$  transform of  $u+z$  is unbounded.

*Case 2.* Let  $A$  have one row,  $x$ , not in  $\mathcal{A}^1$ . Let  $B = (b_{nk})$  where  $b_{nk} = P_n(x)$ ,  $n = 1, 2, \dots$ . Then  $\mathcal{A} \subset \mathcal{B}$  and  $B$  satisfies the hypothesis of Case 1. Let  $E(A) = E(B)$  then  $E(A) \cap \mathcal{A} = \emptyset$  and  $E(A)$  satisfies the other conditions of the lemma's conclusion.

**LEMMA 5.** *If  $\|A\| < \infty$ , and  $A$  is noncoercive then there is  $E(A)$  with  $E(A) \subseteq N_0$ ,  $N_0 \setminus E(A)$  is of first category and if  $u \in E(A)$ , then  $B(u, 1/32) \cap \mathcal{A} = \emptyset$ .*

*Proof. Case 1.* Assume  $a_k = 0$ ,  $k = 1, 2, \dots$ . Let  $\alpha^n$  be the  $n$ th row of  $A$ . Using an argument similar to that of Petersen and Baker [6] (see also the construction of Lemma 4) it can be shown that without loss of generality one may assume that the rows and columns of  $A$  are in  $E^{\infty}$  and moving to the right, (if  $P_j \alpha^n = 0$  then

$P_j \alpha^m = 0$  for  $m \geq n$ ). By Lemma 1  $\|A\|_c > 0$ . Hence there exists increasing sequences  $n(j)$  and  $r(j)$  of positive integers such that

- (i)  $\sum_{k=r(j-1)+1}^{r(j)} |a_{n(j),k}| > \|A\|_c/2$
- (ii)  $(P_{r(j)} - P_{r(j-1)})\alpha^{n(j)} = \alpha^{n(j)}$ .

Let  $J(2j)$  be a subset of  $r(2j - 1)$  to  $r(2j) - 1$  with

$$\left| \sum_{k \in J(2j)} a_{n(j),k} \right| \geq \frac{1}{4} \sum_{j=r(2j-1)+1}^{r(2j)} |a_{n(j),k}| \geq \frac{\|A\|_c}{8}$$

(see Lemma 3). Define  $O_j = \{u \in N_0: u_k = 1 \text{ if } k \in J(2j), u_k = 0 \text{ if } r(2j - 2) + 1 \leq k \leq r(2j), k \notin J(2j)\}$ . Since only a finite number of coordinates are specified for elements of  $O_j$ ,  $O_j$  is open. For each  $k$ ,  $\bigcup_{j=k}^\infty O_j$  is open and dense, hence by the Baire category theorem.  $\bigcap_{k=1}^\infty \bigcup_{j=k}^\infty O_j$  is of second category. Let  $E(A) = \{u \in N_0: Au \text{ has cluster points } \alpha, b, \text{ with } |\alpha - b| \geq \|A\|_c/8\}$ . By construction each element of  $\bigcap_{k=1}^\infty \bigcup_{j=k}^\infty O_j$  has 0 and  $\alpha$  ( $|\alpha| > \|A\|_c/8$ ) as cluster points thus  $E(A)$  is of second category. Let  $u \in E(A)$  and  $\|z\|_\infty < 1/32$  and consider  $A(u + z)$ .  $Au$  has two cluster points separated in distance by at least  $\|A\|_c/8$ , and  $A(z)$  has cluster points separated by at most  $2(1/32)\|A\|_c$  (Lemma 2). Therefore  $A(u + z)$  has at least two cluster points; hence  $u + z \notin \mathcal{A}$ .

*Case 2.* Let  $a_k \neq 0$  for some  $k$ . Define  $B = (b_{nk})$  where  $b_{nk} = a_k, n, k = 1, 2, \dots$ .  $B$  transforms every bounded sequence to a constant sequence, thus the cluster points of  $(A - B)u, u \in m$ , are a shift of those of  $Au$ , and  $A - B$  satisfies the hypothesis of Case 1. Thus the conclusion follows in a manner similar to Case 1.

*Proof of Theorem.* Let  $A_i$  be a countable collection of non-coercive matrices with  $\mathcal{A}_i \supset E^\infty, i = 1, 2, \dots$ . By Lemmas 4 and 5 for each  $i$  there exists  $E(A_i) \subseteq N_0, E(A_i)$  of second category, and if  $u \in E(A_i), B(u, 1/32) \cap \mathcal{A}_i = \emptyset$ . Thus  $\bigcap_{i=1}^\infty E(A_i) \neq \emptyset$  and if  $u \in \bigcap_{i=1}^\infty E(A_i)$ , then  $B(u, 1/32) \cap (\bigcup_{i=1}^\infty \mathcal{A}_i) = \emptyset$ . Hence  $\bigcup_{i=1}^\infty \mathcal{A}_i$  is not dense in  $m$ .

Goffman and Wollan in [4] gave an example of a countable family of  $FK$  spaces contained in  $m$  whose union is dense in  $m$ . They can be realized as summability domains in the following manner. Let  $\{r_i\}$  be a denumeration of the nonzero rationals. Define  $A_i = (a_{nk}^{(i)})$  by

- (i)  $a_{n1}^{(i)} = r_i, a_{n2}^{(i)} = -1, n = 1, 3, 5, \dots$
- (ii)  $a_{n1}^{(i)} = -1, a_{n2}^{(i)} = r_i^{-1}, n = 2, 4, 6, \dots$   
 $a_{nk} = 0, k \geq 3, n = 1, 2, 3, \dots$

Then  $\mathcal{A}_i = \{(x_n)_{n=1}^\infty: x_1 = x, x_2 = r_i x, x_k \text{ arbitrary for } k \geq 3 \text{ and } x \text{ complex}\} \cap m$ . Each  $\mathcal{A}_i$  is nowhere dense in  $m$ , but  $\bigcup_{i=1}^\infty \mathcal{A}_i$  is dense. Note, however, that  $\mathcal{A}_i \not\supset E^\infty$ . Hence the hypothesis that each

$\mathcal{A}_i \supseteq E^\infty$  cannot be removed and our result is in some sense best possible.

Although we have proved our result only for  $\mathcal{A}_i$ , we conjecture that the following more general result holds:

**Conjecture.** If  $\{F_i\}$  is a countable collection of *FK*-spaces each containing  $E^\infty$  but not  $m$ , then  $\bigcup_{i=1}^\infty F_i$  is not dense in  $m$ . (See [8] for definitions and basic results.)

#### REFERENCES

1. G. Bennett and N. J. Kalton, *FK-spaces containing  $c_0$* , Duke Math. J., **39** (1972), 561-582.
2. S. R. Caradus, W. E. Pfaffenberger and Bertram Yood, *Calkin Algebras and Algebras of Operators on Banach Spaces*, Marcel Dekker, New York, 1974.
3. R. DeVos, *Category of sequences of 0's and 1's in some FK spaces*, to appear in the Glasgow Math. J.
4. C. Goffman and G. N. Wollan, *Sequences of regular summability matrices*, Monats. für Math., **76** (1972), 118-120.
5. I. J. Maddox, *Elements of Functional Analysis*, Cambridge Univ. Press, 1970.
6. G. M. Petersen, *Summability fields which span the bounded sequences densely*, Bull. London Math. Soc., **5** (1973), 187-191.
7. ———, *Addendum: Summability fields which span the bounded sequences densely*, Bull. London Math. Soc., **7** (1975), 105.
8. A. Wilansky, *Functional Analysis*, Blaisdell, New York, 1964.

Received April 4, 1977 and in revised form August 8, 1977.

VILLANOVA UNIVERSITY  
VILLANOVA, PA 19085