

Values of Brownian intersection exponents, II: Plane exponents

by

GREGORY F. LAWLER

*Duke University
Durham, NC, U.S.A.*

ODED SCHRAMM

*Microsoft Research
Redmond, WA, U.S.A.*

and

WENDELIN WERNER

*Université Paris-Sud
Orsay, France*

and

*The Weizmann Institute of Science
Rehovot, Israel*

1. Introduction

This paper is the follow-up of the paper [27], in which we derived the exact value of intersection exponents between Brownian motions in a half-plane. In the present paper, we will derive the value of intersection exponents between planar Brownian motions (or simple random walks) in the whole plane.

This problem is very closely related to the more general question of the existence and value of critical exponents for a wide class of two-dimensional systems from statistical physics, including percolation, self-avoiding walks and other random processes. Theoretical physics predicts that these systems behave in a conformally invariant way in the scaling limit, and uses this fact to predict certain critical exponents associated to these systems. We refer to [27] for a more detailed account on this link and for more references on this subject.

Let us now briefly describe some of the results that we shall derive in the present paper. Suppose that B^1, \dots, B^n are $n \geq 2$ independent planar Brownian motions started from n different points in the plane. Then it is easy to see (using a subadditivity argument) that there exists a constant ζ_n such that

$$\mathbf{P}[\forall i \neq j \in \{1, \dots, n\}, B^i[0, t] \cap B^j[0, t] = \emptyset] = t^{-\zeta_n + o(1)} \quad (1.1)$$

when $t \rightarrow \infty$. We shall prove

THEOREM 1.1. *For all $n \geq 2$,*

$$\zeta_n = \frac{1}{24}(4n^2 - 1).$$

This result had been conjectured by Duplantier–Kwon [14] (see also, more recently, Duplantier [11]), using ideas from theoretical physics (conformal field theory, quantum gravity and analogies with some other models for which exponents had also been conjectured).

It was shown in [5] that the exponent ζ_2 equals the corresponding exponent for simple random walks (see also [10]). This result was sharpened in [21], [26], where estimates were derived up to multiplicative constants. It follows from these results and Theorem 1.1 that if S and S' denote two independent simple random walks started from neighboring vertices in \mathbf{Z}^2 , then for some constant $c > 0$,

$$c^{-1}k^{-5/8} \leq \mathbf{P}[S[0, k] \cap S'[0, k] = \emptyset] \leq ck^{-5/8}$$

for all $k \geq 1$. Similarly, [20] and Theorem 1.1 imply

$$c^{-1}t^{-5/8} \leq \mathbf{P}[B^1[0, t] \cap B^2[0, t] = \emptyset] \leq ct^{-5/8}$$

for all $t \geq 1$, assuming that the distance between $B^1(0)$ and $B^2(0)$ is 1, say.

One can define more general exponents, allowing intersection between some Brownian motions, but forbidding intersection between different packets of Brownian motions. For instance, there exists a constant $\zeta = \zeta(n, m)$ such that if B^1, \dots, B^n and B'^1, \dots, B'^m denote $n+m$ independent planar Brownian motions started from points such that $B^i(0) \neq B'^j(0)$ for all $i \leq n$ and $j \leq m$, then

$$\mathbf{P}[\forall i \leq n, \forall j \leq m, B^i[0, t] \cap B'^j[0, t] = \emptyset] = t^{-\zeta + o(1)} \quad (1.2)$$

when $t \rightarrow \infty$. Similarly, one can define exponents $\zeta(n_1, \dots, n_k)$ corresponding to non-intersection between k packets of Brownian motions.

It is easy to see (e.g. [23]) that there is a natural extension of the definition of $\zeta(n, m)$ to pairs (n, λ) , where n is a positive integer and λ is any positive real. In [31], it is shown that there is also a natural definition of $\zeta(\lambda_1, \dots, \lambda_k)$ where the λ_j are positive reals with $\lambda_1, \lambda_2 \geq 1$. In the present paper, we shall derive the value of the exponents for a certain class of k -tuples $(\lambda_1, \dots, \lambda_k)$ (see Theorem 5.3). In particular, we shall prove

THEOREM 1.2. *For all real $\lambda \geq 2$,*

$$\zeta(2, \lambda) = \frac{1}{96} \left((5 + \sqrt{24\lambda + 1})^2 - 4 \right). \quad (1.3)$$

It has been shown by Lawler [20], [22], [23], [25] that some of these critical exponents are closely related to the Hausdorff dimension of exceptional subsets of a planar Brownian path. Recall that a *cut point* of a connected set K is the set of points $x \in K$ such that $K \setminus \{x\}$ is disconnected. The Hausdorff dimension of the set of cut points of the Brownian path $B[0, 1]$ is $2 - 2\zeta_2$ almost surely [20]. Consequently, we get the following corollary from Theorem 1.1.

COROLLARY 1.3. *Let B be Brownian motion in the plane. Then the Hausdorff dimension of the set of cut points of $B[0, 1]$ is $\frac{3}{4}$ almost surely.*

Recall that the *frontier* of a bounded set $K \subset \mathbf{R}^2$ is its outer boundary, i.e., the boundary of the unbounded connected component of $\mathbf{R}^2 \setminus K$. The exponents $\zeta(2, \lambda)$ are closely related to the multifractal spectrum of the Brownian frontier [23]. In particular [22], the Hausdorff dimension d_F of the frontier of $B[0, 1]$ is almost surely $d_F = 2 - \eta_2$, where $\eta_2 := \lim_{\lambda \searrow 0} 2\zeta(2, \lambda)$ is called the *disconnection exponent* for two Brownian paths. (This is not the definition of the disconnection exponent used in [22]; however, the two definitions are equivalent; see [24], [30].) It had been conjectured by Mandelbrot [35] (by analogy with the conjectures for planar self-avoiding walks) that $d_F = \frac{4}{3}$. Upper and lower bounds for η_2 from [6], [40], combined with the fact that $d_F = 2 - \eta_2$ [22], showed that $1.015 < d_F < 1.5$. (See also [4] for another proof of $d_F > 1$.) In the present paper, we derive the values of $\zeta(2, \lambda)$ only for $\lambda \geq 2$, so that we cannot directly apply our results to show that $\eta_2 = \frac{2}{3}$. However, in the subsequent paper [29], we prove

THEOREM 1.4. *The function $\lambda \mapsto \zeta(2, \lambda)$ is real analytic on $(0, \infty)$.*

Combining this with Theorem 1.2 shows immediately that (1.3) holds for all $\lambda > 0$, and therefore $\eta_2 = \frac{2}{3}$. This completes the proof of Mandelbrot’s conjecture:

COROLLARY 1.5. *The Hausdorff dimension of the Brownian frontier is almost surely $\frac{4}{3}$.*

The formula for the multifractal spectrum of the Brownian frontier also follows. This formula has been conjectured in [31] as a consequence of the conjectures of Duplantier–Kwon [14] and of the functional relations between generalized Brownian exponents derived in [31]; see also recent physics work on this subject by Duplantier [11], [12].

The *pioneer points* of B are the image under B of the set of times t such that $B(t)$ is in the frontier of $B[0, t]$. It has been shown [25] that the dimension of the set of pioneer points of B is $2 - \eta_1$, where $\eta_1 := \lim_{\lambda \searrow 0} 2\zeta(1, \lambda)$. Below, we show that

$$\zeta(1, \lambda) = \frac{1}{96} \left((3 + \sqrt{24\lambda + 1})^2 - 4 \right)$$

for all sufficiently large λ (see (5.10)). In [29] it will be proved that $\zeta(1, \lambda)$ is analytic for $\lambda > 0$. Consequently, by analytic continuation of the above formula for $\zeta(1, \lambda)$, it follows that $\eta_1 = \frac{1}{4}$. Hence, using the above-mentioned result of [25], we obtain

COROLLARY 1.6. *The Hausdorff dimension of the set of pioneer points of Brownian motion is $\frac{7}{4}$ almost surely.*

Let us briefly mention that there is a (non-rigorous) link between our results and the conjectures concerning two-dimensional self-avoiding walks. For instance, there is a heuristic argument (see [32]) which uses the Brownian intersection exponents and explains why the number of self-avoiding walks of length N on a planar lattice increases asymptotically like $N^{11/32} \mu^N$, for some (lattice-dependent) constant $\mu > 1$, as conjectured by Nienhuis [36] (see also [34] for a mathematical account).

Just as in [27], a central role in the present paper will be played by SLE_6 , the stochastic Löwner evolution process with parameter 6, which is conjectured [39] to correspond to the scaling limit of two-dimensional critical percolation cluster boundaries. In [39] the processes SLE_κ were introduced, and it was shown that SLE_2 is the scaling limit of loop-erased random walk, assuming the conjecture that the latter has a conformally invariant scaling limit.

Actually, there are two versions of SLE_κ . In the first version, which we now call *radial* SLE_κ , one has a set K_t growing from a boundary point of the unit disk to the interior point 0, while in *chordal* SLE_κ , the set K_t grows from a point in \mathbf{R} to ∞ within the upper half-plane. (The precise definitions will be recalled in §2.2.) By applying conformal maps, these processes can be defined in any simply-connected proper subdomain of the plane.

After recalling the definition of SLE_κ , we study in §3 some of its properties. In particular, the SLE_κ -analogues of the exponents $\zeta(1, \lambda)$, $\lambda \geq 1$, are computed. From Cardy's formula for SLE_6 (that we proved in [27]) we then derive the asymptotic decay of the probability that SLE_6 crosses a long rectangle without touching the upper and lower boundaries of this rectangle, and show that chordal and radial SLE_6 are very closely related.

We then turn our attention to the Brownian intersection exponents. In §5, the definition and some properties of the exponents are recalled. In particular, it is explained how to formulate these exponents in terms of non-intersection between two-dimensional Brownian excursions. Then, these facts (properties of SLE_6 , exponents for SLE_6 , description of the Brownian exponents in terms of Brownian excursions, properties of these Brownian excursions) are combined to derive the value of the Brownian intersection exponents.

2. Preliminaries

2.1. Notation

Throughout this paper, \mathbf{U} will denote the unit disk in the complex plane \mathbf{C} . $C=C_1=\partial\mathbf{U}$ will denote the unit circle. For any $r>0$, $C_r=rC_1$ will denote the circle of radius r centered at 0. \mathbf{H} will denote the upper half-plane $\mathbf{H}=\{x+iy:y>0\}$. When $w\neq w'$ are two points on the unit circle C , then $a(w,w')$ will denote the counterclockwise arc of C from w to w' .

Just as in [31], [32], [27], it will be convenient to use π -extremal distances of quadrilaterals: this just means π times the usual extremal distance. (Extremal distance is also known as extremal length. See, e.g., [1] for the definition and basic properties of extremal length and conformal maps.) The π -extremal distance between sets A and B in a set O will be denoted by $\ell(A,B,O)$.

Let f and g be functions, and let $l\in\mathbf{R}$ or $l=\infty$. We say that $f(x)\sim g(x)$ when $x\rightarrow l$, if $f(x)/g(x)\rightarrow 1$. We write $f(x)\approx g(x)$, if $\log f(x)/\log g(x)\rightarrow 1$, and we write $f(x)\asymp g(x)$, if $f(x)/g(x)$ is bounded above and below by positive finite constants when x is sufficiently close to l .

2.2. Radial and chordal SLE

In [27], we studied chordal SLE_κ as a random increasing family $(K_t, t\geq 0)$ of bounded subsets of the upper half-plane \mathbf{H} (or, more generally, their image under a conformal map). As t increases, the set K_t grows, and $\bigcup_t K_t$ is unbounded. One can say that K_t is growing towards ∞ , which we think of as a boundary point of \mathbf{H} .

Similarly, when looking at the conformal image of $(K_t, t\geq 0)$ under the map $\varphi(z)=(z-i)/(z+i)$ that maps \mathbf{H} onto the unit disk and ∞ to 1, we get an increasing family of subsets of the unit disk that is growing towards 1.

In the present paper, we will mainly use a variant of this process, called *radial* SLE_κ , where $(K_t, t\geq 0)$ is an increasing family of subsets of the unit disk that grows towards 0. The main distinction is that 0 is an interior point of \mathbf{U} , instead of a boundary point.

Suppose that $(\delta_t, t\geq 0)$ is a continuous function taking values on the unit circle $C_1=\partial\mathbf{U}$. Consider for each $z\in\mathbf{U}$ the solution $g_t=g_t(z)$ of the ordinary differential equation

$$\partial_t g_t = g_t \frac{\delta_t + g_t}{\delta_t - g_t}, \quad t \geq 0, \tag{2.1}$$

with $g_0=z$. This equation (and the corresponding equation for g_t^{-1}) was first considered by Löwner (see [33], also [37]) and is called Löwner's differential equation. For each $z\in\mathbf{U}$, it is well defined up to the time τ_z where $\lim_{t\uparrow\tau_z} g_t = \delta_{\tau_z}$, if there is such a τ_z ,

and otherwise $\tau_z = \infty$. Let D_t be the set of $z \in \mathbf{U}$ such that $t < \tau_z$ (i.e., the set on which g_t is defined), and $K_t = \mathbf{U} \setminus D_t$. It is easy to check that g_t is the unique conformal map from D_t onto \mathbf{U} such that $g_t(0) = 0$ and $g_t'(0)$ is a positive real number. (The notation g' refers to differentiation with respect to z .) It is also easy to verify that $g_t'(0) = \exp(t)$ by differentiating both sides of (2.1) with respect to z at $z = 0$ and noting that $g_t(0) = 0$.

Let $(B_t, t \geq 0)$ be Brownian motion on the real line, starting at some point $B_0 \in \mathbf{R}$, and let $\kappa \geq 0$. Set $\delta_t := \exp(i\sqrt{\kappa} B_t)$ and consider the solution to Löwner's differential equation as defined above. (Note that δ_t is just Brownian motion on $\partial\mathbf{U}$ with time scaled by κ .) The resulting process was defined in [39] and called SLE_κ (this acronym stands for stochastic Löwner evolution with parameter κ). The set K_t is called the hull of SLE_κ , and $(\delta_t)_{t \geq 0}$ its driving process.

If $f: D \rightarrow \mathbf{U}$ is a conformal map, then radial SLE_κ in D starting from f can be defined as the composition $g_t \circ f$, where g_t is radial SLE_κ in \mathbf{U} . Its hull is $f^{-1}(K_t)$, where K_t is the hull of g_t . If ∂D is sufficiently tame, then f^{-1} extends continuously to $\partial\mathbf{U}$, and hence $f^{-1}(\delta_0)$ is well defined, where δ is the driving parameter of g_t . We may then refer to the resulting SLE_κ -process in D as SLE_κ from $f^{-1}(\delta_0)$ to $f^{-1}(0)$ in D .

In [39], another variant SLE was also defined (see also [27]), which we now call *chordal* SLE_κ . Let \mathbf{H} be the upper half-plane, let $\delta_t = \sqrt{\kappa} B_t$, and consider for each $z \in \mathbf{H}$ the solution $\tilde{g}_t = \tilde{g}_t(z)$ of the differential equation

$$\partial_t \tilde{g}_t = \frac{-2}{\delta_t - \tilde{g}_t}, \quad t \geq 0, \quad (2.2)$$

with $\tilde{g}_0(z) = z$. As before, let \tilde{D}_t be the set of points $z \in \mathbf{H}$ for which (2.2) has a solution in some interval $[0, t']$, $t' > t$, and let $\tilde{K}_t := \mathbf{H} \setminus \tilde{D}_t$. Then the process $(\tilde{g}_t)_{t \geq 0}$ is chordal SLE_κ in \mathbf{H} , and \tilde{K}_t is its hull. Recall that \tilde{K}_t is bounded for each t , but $\bigcup_{t \geq 0} \tilde{K}_t$ is unbounded. If $f: D \rightarrow \mathbf{U}$ is a conformal map, then $f \circ \tilde{g}_t$ is called chordal SLE_κ in D . Its hull is $f^{-1}(K_t)$, where K_t is the hull of the process \tilde{g}_t . If ∂D is sufficiently tame, $f^{-1}(\infty)$ and $f^{-1}(\delta_0)$ are well defined. In this case, we refer to the process $f \circ \tilde{g}_t$ as SLE_κ from $f^{-1}(\delta_0)$ to $f^{-1}(\infty)$ in D . This terminology is explained more fully in [27].

For the main results of this paper, only the case $\kappa = 6$ will be used. As we shall see in §4.1, radial SLE_6 is essentially the same process as chordal SLE_6 . For $\kappa \neq 6$, the radial and chordal SLE_κ -processes are closely related, but the equivalence is weaker.

3. Annulus-crossing exponents for SLE

3.1. Statement

Consider a radial SLE_κ -process (with $\kappa \geq 0$) in the unit disk \mathbf{U} , with driving element δ_t starting at $\delta_0 = 1$. As described before, let g_t be the conformal map $g_t: D_t \rightarrow \mathbf{U}$, with

$g_t(0)=0$ and $g'_t(0)=\exp(t)$. Let

$$A_t := \partial\mathbf{U} \setminus \bar{K}_t.$$

It is easy to see that A_t is either an arc on $\partial\mathbf{U}$, or $A_t=\emptyset$.

Let $b \geq 0$, and set

$$\nu = \nu(\kappa, b) := \frac{1}{16} (8b + \kappa - 4 + \sqrt{(\kappa - 4)^2 + 16b\kappa}). \tag{3.1}$$

THEOREM 3.1. *Let $\kappa > 0$ and $r \in (0, 1)$, and let $T(r)$ be the least $t > 0$ such that K_t intersects the circle C_r . Let $\mathfrak{L}(r)$ be the π -extremal distance from C_r to C_1 in $\mathbf{U} \setminus K_{T(r)}$ (we set $\mathfrak{L}(r) := \infty$ if $A_t = \emptyset$). Then for $b \geq 1$, as $r \rightarrow 0$,*

$$\mathbf{E}[\exp(-b\mathfrak{L}(r))] \asymp r^\nu.$$

This theorem gives an analogue of Theorem 3.1 in [27] for crossing an annulus. Its proof and usage will also be similar.

Remarks. The constants implicit in the \asymp -notation may depend on κ and b .

One can also show that the theorem holds when $b \in (0, 1)$, but we only need the case $b \geq 1$ (and $\kappa = 6$) in the present paper. Observe that the case $\kappa = 0$ is also correct, and easy to verify, for then $K_{T(r)} = [r, 1]$.

One should also note that the values of the exponents for $\kappa = 2$ fit with the conjecture that SLE_2 is the scaling limit of two-dimensional loop-erased random walks [39] and the exponents computed by Kenyon [16], [17], [18] for loop-erased walks. For instance, the definition of loop-erased random walks suggests that the number of vertices in the loop erasure of an N^2 -step walk is roughly $N^{2-\nu(2,1)} = N^{5/4}$ (loosely speaking, in order for one of the N^2 steps of the simple random walk to remain in the loop erasure, the future of the random walk beyond that step has to avoid the past loop-erased walk), in accordance with Kenyon’s results.

Similarly, combining this result (with $\kappa = 6$) with the non-intersection exponents between SLE_6 in a rectangle (see the discussion at the end of [27]) and the restriction property for SLE_6 (which was proved for chordal SLE_6 in [27] and will be proved for radial SLE_6 in §4) leads to the value of non-intersection exponents between independent SLE_6 in an annulus that agree with predictions for annulus crossings in a critical percolation cluster (see [15], [9], [2]).

The proof of Theorem 3.1 will consist of three steps. First, we will obtain an estimate on $\mathbf{E}[|g'_t(e^{ix})|^b]$ for large (deterministic) times. We deduce from it a result concerning the large-time behavior of the arc length of $g_t(A_t)$, and then show that this implies Theorem 3.1.

3.2. Derivative exponents

LEMMA 3.2. *Assume that $\kappa > 0$ and $b > 0$. Let $\mathcal{H}(x, t)$ denote the event $\{\exp(ix) \in A_t\}$, and set*

$$f(x, t) := \mathbf{E}[|g'_t(\exp(ix))|^b 1_{\mathcal{H}(x, t)}],$$

$$q = q(\kappa, b) := \frac{\kappa - 4 + \sqrt{(\kappa - 4)^2 + 16b\kappa}}{2\kappa},$$

$$h^*(x, t) := \exp(-t\nu) \left(\sin\left(\frac{1}{2}x\right)\right)^q,$$

where ν is as in (3.1). Then there is a constant $c > 0$ such that

$$\forall t \geq 1, \forall x \in (0, 2\pi), \quad h^*(x, t) \leq f(x, t) \leq ch^*(x, t).$$

Proof. Let $\delta_t = \exp(i\sqrt{\kappa} B_t)$ be the driving process of the SLE $_{\kappa}$, with $B_0 = 0$. For all $x \in (0, 2\pi)$, let Y_t^x be the continuous real-valued function of t which satisfies

$$g_t(e^{ix}) = \delta_t \exp(iY_t^x)$$

and $Y_0^x = x$. The function Y_t^x is defined on the set of pairs (x, t) such that $\mathcal{H}(x, t)$ holds. Since g_t satisfies Löwner’s differential equation

$$\partial_t g_t(z) = g_t(z) \frac{\delta_t + g_t(z)}{\delta_t - g_t(z)}, \tag{3.2}$$

we find that

$$dY_t^x = \sqrt{\kappa} dB_t + \cot\left(\frac{1}{2}Y_t^x\right) dt. \tag{3.3}$$

Let

$$\tau^x := \inf\{t \geq 0 : Y_t^x \in \{0, 2\pi\}\}$$

denote the time at which $\exp(ix)$ is absorbed by \bar{K}_t , and define for all $t < \tau^x$

$$\Phi_t^x := |g'_t(\exp(ix))|.$$

On $t \geq \tau^x$ set $\Phi_t^x := 0$. Note that on $t < \tau^x$,

$$\Phi_t^x = \partial_x Y_t^x.$$

By differentiating (3.2) with respect to z , we find that for $t < \tau^x$,

$$\partial_t \log \Phi_t^x = -\frac{1}{2 \sin^2\left(\frac{1}{2}Y_t^x\right)}, \tag{3.4}$$

and hence (since $\Phi_0^x=1$),

$$(\Phi_t^x)^b = \exp\left(-\frac{1}{2}b \int_0^t \frac{ds}{\sin^2(\frac{1}{2}Y_s^x)}\right) \tag{3.5}$$

for $t < \tau^x$.

We now show that the right-hand side of (3.5) is 0 when $t = \tau^x$. For all $x \in (0, 2\pi)$, choose $m \in \mathbf{N}$ such that $x \in (2^{-m}, 2\pi - 2^{-m})$, and define for all $n \geq m$ the stopping times

$$\begin{aligned} \varrho_n &:= \inf \{t > 0 : Y_t^x \notin (2^{-n}, 2\pi - 2^{-n})\}, \\ \varrho'_n &:= \inf \{t > \varrho_n : |Y_t^x - Y_{\varrho_n}^x| \geq 2^{-n-1}\}, \end{aligned}$$

and the event $\mathcal{R}_n := \{\varrho'_n - \varrho_n > 4^{-n}\}$. It is easy to see (for instance by comparing Y^x with two Bessel processes and using their scaling property, or alternatively, by comparing with Brownian motions with constant drifts; see, e.g., [38] for the definition of Bessel processes) that there is some constant $c' > 0$ such that for every $n \geq m$,

$$\mathbf{P}[\mathcal{R}_n] > c'.$$

The strong Markov property shows that the events $(\mathcal{R}_m, \mathcal{R}_{m+1}, \dots)$ are independent, so that almost surely there exist infinitely many values of $n \in \mathbf{N}$ such that \mathcal{R}_n holds. For these values of n ,

$$\int_{\varrho_n}^{\varrho_{n+1}} \frac{dt}{\sin^2(\frac{1}{2}Y_t^x)} \geq \frac{4^{-n}}{\sin^2(2^{-n})} \geq 1.$$

Hence, almost surely,

$$\int_0^{\tau^x} \frac{dt}{\sin^2(\frac{1}{2}Y_t^x)} = \infty. \tag{3.6}$$

A similar argument shows that

$$\lim_{x \searrow 0} f(x, t) = \lim_{x \nearrow 2\pi} f(x, t) = 0 \tag{3.7}$$

holds for all fixed $t > 0$. Suppose, for instance, that $x \leq 2^{-n_0} \min(\sqrt{t}, \frac{1}{2}\pi)$, define $\varrho''_0 = 0$, $x_0 = x$, and for all $n \geq 1$,

$$\varrho''_n = \inf \{s \geq \varrho''_{n-1} : s = \varrho''_{n-1} + (x_{n-1})^2 \text{ or } |Y_s^x - x_{n-1}| \geq \frac{1}{2}x_{n-1}\}$$

and $x_n = Y_{\varrho''_n}^x$. Clearly, for all $n \leq n_0$, $0 < x_n < \frac{1}{2}\pi$ and $\varrho''_n < t$. By comparing Y^x with Bessel processes or Brownian motions with constant drift, it is easy to check that for some (sufficiently small) $c > 0$ (independent of n and x), if we define the events $\mathcal{R}'_n :=$

$\{\varrho_n'' = \varrho_{n-1}'' + (x_{n-1})^2\}$, then for all $n \leq n_0$, $\mathbf{P}[\mathcal{R}'_n | \mathcal{F}_{n-1}] \geq c'$, where \mathcal{F}_{n-1} denotes the σ -field generated by the events $\mathcal{R}'_1, \dots, \mathcal{R}'_{n-1}$. It therefore easily follows that when $x \searrow 0$,

$$\int_0^{\min\{\tau^x, t\}} \frac{ds}{\sin^2(\frac{1}{2}Y_s^x)} \rightarrow \infty$$

in probability, and therefore also that $f(x, t) \rightarrow 0$.

Let $F: [0, 2\pi] \rightarrow \mathbf{R}$ be a continuous function with $F(0) = F(2\pi) = 0$, which is smooth in $(0, 2\pi)$, and set

$$h(x, t) = h_F(x, t) := \mathbf{E}[(\Phi_t^x)^b F(Y_t^x)].$$

By (3.5) and the general theory of diffusion Markov processes (see, e.g., [3]), we know that h is smooth in $(0, 2\pi) \times \mathbf{R}_+$. From the Markov property for Y_t^x and (3.5), it follows that $h(Y_t^x, t' - t)(\Phi_t^x)^b$ is a local martingale on $t < \min\{\tau^x, t'\}$. Consequently, the drift term of the stochastic differential $d(h(Y_t^x, t' - t)(\Phi_t^x)^b)$ is zero at $t = 0$. By Itô's formula, this means

$$\partial_t h = \Lambda h, \tag{3.8}$$

where

$$\Lambda h := \frac{1}{2}\kappa \partial_x^2 h + \cot(\frac{1}{2}x) \partial_x h - \frac{b}{2 \sin^2(\frac{1}{2}x)} h.$$

It can be verified directly that h^* from the statement of the lemma solves (3.8). We therefore choose

$$F(x) := (\sin(\frac{1}{2}x))^q,$$

and claim that $h^* = h_F$. Indeed, both satisfy (3.8) on $[0, 2\pi] \times [0, \infty)$, and $h^*(x, 0) = F(x) = h_F(x, 0)$ on $[0, 2\pi]$. Moreover, $F \leq 1$ implies that $h_F \leq f$ everywhere, so $h^* - h_F = 0$ on $\{0, 2\pi\} \times (0, \infty)$, by (3.7). It is also immediate to verify that $h_F(x, t) \rightarrow 0$ as $(x, t) \rightarrow (0, 0)$ or $(x, t) \rightarrow (2\pi, 0)$. Set $M = h_F - h^*$. Then M is smooth in $(0, 2\pi) \times (0, \infty)$, continuous on $[0, 2\pi] \times [0, \infty)$, and satisfies $\partial_t M = \Lambda M$.

The proof that $M = 0$ can be viewed as a straightforward application of the maximum principle. Fix some $\varepsilon > 0$, and suppose that $M \geq \varepsilon$ at some point $(x, t) \in [0, 2\pi] \times [0, \infty)$. Among such points, let (x_0, t_0) be a point with t_0 minimal. It is clear that there must be such a minimal point and that $x_0 \in (0, 2\pi)$, $t_0 > 0$. At (x_0, t_0) we must have $\partial_t M \geq 0$, by minimality of t_0 . Similarly, $\partial_x M(x_0, t_0) = 0$, $\partial_x^2 M(x_0, t_0) \leq 0$ and $M(x_0, t_0) = \varepsilon$. However, this gives $0 \leq \partial_t M(x_0, t_0) = (\Lambda M)(x_0, t_0) \leq -b\varepsilon / 2 \sin^2(\frac{1}{2}x_0)$, by the definition of the operator Λ , a contradiction. Since ε was arbitrary, this gives $M \leq 0$. The same argument applied to $-M$ shows that $M \geq 0$, which verifies (the subscript will henceforth be omitted from h_F)

$$h(x, t) = h^*(x, t) = \exp(-t\nu) (\sin(\frac{1}{2}x))^q. \tag{3.9}$$

As mentioned above, $F \leq 1$ implies $h(x, t) \leq f(x, t)$. Therefore, it remains to prove that for all $t \geq 1$ and $x \in (0, 2\pi)$, $f(x, t) \leq ch(x, t)$ for some fixed $c > 0$. By the Markov property at time $t-1$, we have for $t > 1$,

$$h(x, t) = \mathbf{E}[(\Phi_{t-1}^x)^b h(Y_{t-1}^x, 1)],$$

and similarly with f replacing h on both sides. Hence, it suffices to prove $ch(x, 1) \geq f(x, 1)$; i.e.,

$$c\mathbf{E}[(\Phi_1^x)^b F(Y_1^x)] \geq \mathbf{E}[(\Phi_1^x)^b].$$

Let $\sigma_y = \sigma_y^x$ be the first time $s \geq 0$ such that $Y_s^x = y$, and if no such s exists, set $\sigma_y = \infty$. By (3.9), $h(x, t)$ is a decreasing function of t . This and the strong Markov property give

$$\mathbf{E}[F(Y_1^x) 1_{\{\sigma_y < 1\}} (\Phi_1^x)^b] \geq h(y, 1) \mathbf{E}[1_{\{\sigma_y < 1\}} (\Phi_{\sigma_y}^x)^b]. \tag{3.10}$$

Let

$$a := \min\{y > 0 : y - \cot(\frac{1}{2}y) = -2\pi\},$$

and consider the event

$$\mathcal{A} := \bigcap_{s \in [0, 1]} \{Y_s^x \in (0, a)\}.$$

From (3.3) it then follows that on the event \mathcal{A} , $\sqrt{\kappa} B_1 \leq a - \cot(\frac{1}{2}a) = -2\pi$. Define the Brownian motion \tilde{B} on $[0, 1]$ by $\tilde{B}_s := B_s - 2sB_1$, and define \tilde{Y}_s^x by the equation (3.3), but with \tilde{B} replacing B . Note that on \mathcal{A} we have $B_1 < 0$, and hence

$$\forall s \in [0, 1], \quad \tilde{Y}_s^x \geq Y_s^x. \tag{3.11}$$

Moreover, given \mathcal{A} , there is a minimal $s_0 \in [0, 1]$ with $\tilde{Y}_{s_0}^x = \pi$. Since (3.11) holds on \mathcal{A} , it follows from (3.5) that

$$\tilde{\Phi}_{s_0}^x \geq \Phi_{s_0}^x \geq \Phi_1^x,$$

where $\tilde{\Phi}$ is the analogue of Φ for the process \tilde{Y}^x . Since \tilde{Y}^x has the same law as Y^x , with (3.10) this implies

$$h(x, 1) \geq h(\pi, 1) \mathbf{E}[1_{\{\sigma_\pi < 1\}} (\Phi_{\sigma_\pi}^x)^b] \geq h(\pi, 1) \mathbf{E}[1_{\mathcal{A}} (\Phi_1^x)^b].$$

The same proof gives this relation with \mathcal{A} replaced by the event

$$\mathcal{A}' := \bigcap_{s \in [0, 1]} \{Y_s^x \in (2\pi - a, 2\pi)\}.$$

However, for the event

$$\mathcal{A}'' := \{\exists s \in [0, 1] : Y_s^x \in [a, 2\pi - a]\},$$

we have

$$h(x, 1) \geq h(a, 1) \mathbf{E}[1_{\mathcal{A}''}(\Phi_1^x)^b].$$

Since $\mathcal{A} \cup \mathcal{A}' \cup \mathcal{A}'' \supset \mathcal{H}(x, 1)$ and $h(a, 1) \leq h(\pi, 1)$, we get

$$h(x, 1) \geq \frac{1}{3} h(a, 1) f(x, 1).$$

This completes the proof of the lemma. □

3.3. Harmonic measure exponents

By conformal invariance, the harmonic measure from 0 of A_t in $D_t = \mathbf{U} \setminus K_t$ is $L_t/2\pi$, where L_t is the length of the arc $g_t(A_t)$.

THEOREM 3.3. *Suppose that $\kappa > 0$ and $b \geq 1$. Then, when $t \rightarrow \infty$,*

$$\mathbf{E}[(L_t)^b] \asymp \exp(-\nu t).$$

Proof. We have to relate the behavior of $|g'_t(e^{ix})|^b$, which we have studied above, to the behavior of

$$(L_t)^b = \left(\int_0^{2\pi} |g'_t(e^{ix})| dx \right)^b = \left(\int_0^{2\pi} \Phi_t^x dx \right)^b,$$

where we set $\Phi_t^x = 0$ if $\tau^x \leq t$. By convexity of the function $a \mapsto a^b$, it is clear that

$$\begin{aligned} \frac{1}{(2\pi)^b} \mathbf{E}[(L_t)^b] &= \mathbf{E} \left[\left(\frac{1}{2\pi} \int_0^{2\pi} \Phi_t^x dx \right)^b \right] \leq \mathbf{E} \left[\frac{1}{2\pi} \int_0^{2\pi} (\Phi_t^x)^b dx \right] \\ &= \frac{1}{2\pi} \int_0^{2\pi} f(x, t) dx \leq c \exp(-\nu t), \end{aligned}$$

where we have used Lemma 3.2 for the last inequality. Consequently, we only need to prove the lower bound for $\mathbf{E}[(L_t)^b]$.

We will find constants $c_1, c_2 > 0$ and an event \mathcal{U}_t^* such that

$$\mathbf{E}[(\Phi_t^\pi)^b 1_{\mathcal{U}_t^*}] \geq c_1 e^{-\nu t},$$

and on the event \mathcal{U}_t^* ,

$$|\log \Phi_t^\pi - \log \Phi_t^y| \leq c_2, \quad \forall y \in [\pi, \pi + c_1].$$

Then

$$\mathbf{E}[(L_t)^b] \geq \mathbf{E} \left[1_{\mathcal{U}_t^*} \left(\int_\pi^{\pi+c_1} \Phi_t^x dx \right)^b \right] \geq (c_1 e^{-c_2})^b \mathbf{E}[1_{\mathcal{U}_t^*} (\Phi_t^\pi)^b] \geq c_3 e^{-\nu t},$$

which will prove the theorem.

Assume (with no loss of generality) that $t > 3$, and let t' be the integer in $(t-2, t-1]$. Define the event

$$\mathcal{V}_t = \left\{ \frac{1}{2}\pi \leq Y_s^x \leq \frac{3}{2}\pi, \forall s \in [t', t] \right\}.$$

It follows from Lemma 3.2 that there is some constant $c_4 > 0$ such that

$$\mathbf{E}[(\Phi_{t'-1}^x)^b 1_{\{Y_{t'-1}^x \in [c_4, 2\pi - c_4]\}}] \geq c_4 e^{-\nu t}.$$

This clearly implies

$$\mathbf{E}[(\Phi_t^\pi)^b 1_{\mathcal{V}_t}] \geq c_5 e^{-\nu t}$$

for some $c_5 > 0$.

Let $\mathcal{U}_t = \mathcal{U}_t(\alpha)$ be the event

$$\mathcal{U}_t := \{ \alpha e^{-s/8} \leq Y_s^\pi \leq 2\pi - \alpha e^{-s/8}, \forall s \in [0, t] \}.$$

We claim that for some $\alpha > 0$, and every $t > 3$,

$$\mathbf{E}[(\Phi_t^\pi)^b 1_{\mathcal{U}_t} 1_{\mathcal{V}_t}] \geq \frac{1}{2} c_5 e^{-\nu t}. \tag{3.12}$$

To prove this it suffices to show that for some $\alpha > 0$,

$$\mathbf{E}[(\Phi_t^x)^b 1_{\neg \mathcal{U}_t} 1_{\mathcal{V}_t}] \leq \frac{1}{2} c_5 e^{-\nu t}. \tag{3.13}$$

For $u = 0, \dots, t' - 1$, and $\alpha \in (0, \frac{1}{5})$, let $\mathcal{W}_u = \mathcal{W}_u(\alpha)$ be the event

$$\mathcal{W}_u := \{ \alpha e^{-u/8} \leq Y_s^\pi \leq 2\pi - \alpha e^{-u/8}, \forall s \in [u, u+1] \}.$$

Note that

$$\bigcup_{u=0}^{t'-1} \neg \mathcal{W}_u \supset \mathcal{V}_t \cap \neg \mathcal{U}_t.$$

Hence,

$$\mathbf{E}[(\Phi_t^x)^b 1_{\neg \mathcal{U}_t} 1_{\mathcal{V}_t}] \leq \sum_{u=0}^{t'-1} \mathbf{E}[(\Phi_t^x)^b 1_{\neg \mathcal{W}_u}]. \tag{3.14}$$

Note that for $u = 0, \dots, t' - 1$, the strong Markov property shows that

$$\mathbf{E}[(\Phi_t^x)^b 1_{\neg \mathcal{W}_u} | \mathcal{F}_u] \leq (\Phi_u^x)^b f(\alpha e^{-u/8}, t-u-1)$$

(here \mathcal{F}_u denotes the σ -field generated by $(Y_s, s \leq u)$). Hence, by Lemma 3.2,

$$\mathbf{E}[(\Phi_t^x)^b 1_{\neg \mathcal{W}_u}] \leq c_6 \alpha^q e^{-uq/8} e^{-\nu t}.$$

Now (3.14) gives

$$\mathbf{E}[(\Phi_t^x)^b 1_{\mathcal{U}_t} 1_{\mathcal{V}_t}] \leq c_7 \alpha^q e^{-\nu t},$$

and hence by choosing α sufficiently small, we get (3.13) and therefore (3.12). Fix such an $\alpha \in (0, \frac{1}{5})$, and let $\mathcal{U}_t^* := \mathcal{U}_t \cap \mathcal{V}_t$.

Observe that (3.3) implies that if $0 < x < y < 2\pi$,

$$\partial_t(Y_t^y - Y_t^x) = \cot(\frac{1}{2}Y_t^y) - \cot(\frac{1}{2}Y_t^x) \leq -\frac{1}{2}(Y_t^y - Y_t^x)$$

(since $\cot'(u) \leq -1$ in the range $u \in (0, \pi)$) as long as $t < \min\{\tau^x, \tau^y\}$, so that

$$|Y_t^y - Y_t^x| \leq |x - y| e^{-t/2}. \tag{3.15}$$

Let $y \in (\pi, \pi + \frac{1}{2}\alpha)$. Then

$$0 < Y_s^y - Y_s^\pi \leq e^{-s/2}(y - \pi) \leq \frac{1}{2}\alpha e^{-s/2}, \quad \forall s \leq \min\{\tau^\pi, \tau^y\}.$$

On the event \mathcal{U}_t^* , we must therefore have $t < \min\{\tau^x, \tau^y\}$ and

$$Y_s^\pi, Y_s^y \in [\frac{1}{2}\alpha e^{-s/8}, 2\pi - \frac{1}{2}\alpha e^{-s/8}]$$

for all $s \in [0, t]$. By (3.4), this shows that on the event \mathcal{U}_t^* , for all $s \leq t$,

$$\begin{aligned} |\partial_s(\log \Phi_s^y - \log \Phi_s^\pi)| &\leq |Y_s^y - Y_s^\pi| \max\{\frac{1}{2}|\partial_x(\sin^{-2}(\frac{1}{2}x))| : x \in [\frac{1}{2}\alpha e^{-s/8}, 2\pi - \frac{1}{2}\alpha e^{-s/8}]\} \\ &\leq c_7 |Y_s^y - Y_s^\pi| e^{3s/8}. \end{aligned}$$

Now (3.15) gives

$$|\partial_s(\log \Phi_s^y - \log \Phi_s^\pi)| \leq c_8 e^{-s/8}.$$

Therefore, on the event \mathcal{U}_t^* ,

$$|\log(\Phi_t^y / \Phi_t^\pi)| \leq 8c_8.$$

This completes the proof of the theorem. □

3.4. Extremal distance exponents

Proof of Theorem 3.1. Let $\varrho(t) := \inf\{|z| : z \in K_t\}$. Recall that

$$T(r) = \inf\{t : \varrho(t) = r\},$$

$A_t = \partial\mathbf{U} \setminus \overline{K_t}$ and that L_t is the length of the arc $g_t(A_t)$. Recall that the Schwarz lemma says that if $G: \mathbf{U} \rightarrow \mathbf{U}$ is analytic, then $|G'(0)| \leq 1$, and the Koebe $\frac{1}{4}$ -theorem says that

if $G: \mathbf{U} \rightarrow \mathbf{C}$ is conformal with $G(0)=0$, then $G(\mathbf{U}) \supset \frac{1}{4}|G'(0)|\mathbf{U}$. (See, e.g., [1].) Since $g'_t(0)=\exp(t)$, applying the former to $z \mapsto g_t(\varrho(t)z)$ and the latter to g_t^{-1} give

$$\frac{1}{4}e^{-t} \leq \varrho(t) \leq e^{-t}.$$

In particular, if we fix the radius $r < \frac{1}{8}$ and define the deterministic times

$$t' = t'(r) := \log \frac{1}{r}, \quad t = t(r) := \log \frac{1}{8r},$$

then

$$t < T(r) \leq t' \leq T(\frac{1}{4}r)$$

and

$$\varrho(t) \geq 2r \geq r \geq \varrho(t') \geq \frac{1}{4}r \geq \frac{1}{8}r.$$

In the following lines, $\ell(S, S'; U)$ will stand for the π -extremal distance between the sets S and S' in U . Recall that $\mathfrak{L}(r) = \ell(C_r, C_1; \mathbf{U} \setminus K_{T(r)})$ and define $l_r = \ell(C_r, C_1; \mathbf{U} \setminus K_{t(r)})$. It follows from the above that

$$l_r \leq \mathfrak{L}(r) \leq l_{r/8}. \tag{3.16}$$

Hence, it will be sufficient to study the asymptotic behavior of $\mathbf{E}[\exp(-bl_r)]$.

Note that $g_t: D_t \rightarrow \mathbf{U}$ is a conformal map defined on $D_t \supset 2r\mathbf{U}$, that $g_t(0)=0$ and that $g'_t(0)=1/8r$. Hence, it follows immediately from the Schwarz lemma and the Koebe $\frac{1}{4}$ -theorem that for any $r < \frac{1}{8}$,

$$2^{-5}\mathbf{U} \subset g_t(r\mathbf{U}) \subset \frac{1}{2}\mathbf{U}.$$

Hence,

$$\ell(C_{2^{-5}}, g_t(A_t); \mathbf{U}) \geq l_r \geq \ell(C_{1/2}, g_t(A_t); \mathbf{U}),$$

and this implies easily that

$$\exp(-l_r) \asymp L_{t(r)}.$$

Theorem 3.1 now follows from Theorem 3.3 and (3.16). □

4. Properties of SLE₆

We now turn our attention towards specific properties of SLE₆.

4.1. Equivalence of chordal and radial SLE₆

The following result shows that chordal SLE₆ and radial SLE₆ are nearly the same process. (When $\kappa \neq 6$, a weaker form of equivalence holds.) A consequence of the equivalence

of radial and chordal SLE_6 is that radial SLE_6 satisfies a restriction property, since in [27] a restriction property for chordal SLE_6 has been established. The significance of the restriction property to the Brownian intersection exponents has been evident since [32].

THEOREM 4.1. *Let $\theta \in \partial\mathbf{U} \setminus \{1\}$, and let K_t be the hull of a radial SLE_6 -process in the unit disk \mathbf{U} with driving process δ_t satisfying $\delta_0 = \theta$. Set*

$$T := \sup\{t \geq 0 : 1 \notin \bar{K}_t\}.$$

Let \tilde{K}_u be the hull of a chordal SLE_6 -process in \mathbf{U} starting also at θ and growing towards 1, and let

$$\tilde{T} := \sup\{u \geq 0 : 0 \notin \tilde{K}_u\}.$$

Then, up to a random time change, the process $t \mapsto K_t$ restricted to $[0, T)$ has the same law as the process $u \mapsto \tilde{K}_u$ restricted to $[0, \tilde{T})$.

Note that T (resp. \tilde{T}) is the first time where K_t (resp. \tilde{K}_u) disconnects 0 from 1.

Proof. In order to point out where the assumption $\kappa=6$ is important, we let $(K_t, t \geq 0)$ and $(\tilde{K}_u, u \geq 0)$ be SLE_κ -processes, without fixing the value of κ for the moment.

Let us first briefly recall (see, e.g., [27]) how \tilde{K}_u is defined. Let ψ be the Möbius transformation that satisfies $\psi(\mathbf{U}) = \mathbf{H}$, $\psi(1) = \infty$, $\psi(-1) = 0$ and $\psi(0) = i$; i.e.,

$$\psi(z) = i \frac{1+z}{1-z}.$$

Suppose that $u \mapsto \tilde{B}_u$ is a real-valued Brownian motion such that $\sqrt{\kappa} \tilde{B}_0 = \psi(e^{i\theta})$. For all $z \in \mathbf{U}$, define the function $\tilde{g}_u = \tilde{g}_u(z)$ such that $\tilde{g}_0(z) = \psi(z)$ and

$$\partial_u \tilde{g}_u = \frac{2}{\tilde{g}_u - \sqrt{\kappa} \tilde{B}_u}.$$

This function is defined up to the (possibly infinite) time σ_z where $\tilde{g}_u(z)$ hits $\sqrt{\kappa} \tilde{B}_u$. Then, \tilde{K}_u is defined by $\tilde{K}_u = \{z \in \mathbf{U} : \sigma_z \leq u\}$, so that \tilde{g}_u is a conformal map from $\mathbf{U} \setminus \tilde{K}_u$ onto the upper half-plane. This defines the process $(\tilde{K}_u, u \geq 0)$ (the scaling property of Brownian motion shows that the choice of the conformal map ψ only influences the law of $(\tilde{K}_u)_{u \geq 0}$ via a time change).

We are now going to study the radial SLE_κ . Let $g_t: \mathbf{U} \setminus K_t \rightarrow \mathbf{U}$ be the conformal map normalized by $g_t(0) = 0$ and $g'_t(0) > 0$. Recall that

$$\partial_t g = g \frac{\delta + g}{\delta - g}, \tag{4.1}$$

where $\delta_t = \exp(i\sqrt{\kappa} B_t)$, and B is Brownian motion on \mathbf{R} with $\exp(iB_0) = \theta$. Let ψ be the Möbius transformation as before, and define

$$\begin{aligned} e_t &:= g_t(1), \\ f_t(z) &:= \psi(g_t(z)/e_t), \\ \gamma_t &:= \psi(\delta_t/e_t). \end{aligned}$$

These are well defined, as long as $t < T$. Note that f_t is a conformal map from $\mathbf{U} \setminus K_t$ onto the upper half-plane, $f_t(1) = \infty$ and $\gamma_t \in \mathbf{R}$. From (4.1) it follows that

$$\partial_t f = -\frac{(1+\gamma^2)(1+f^2)}{2(\gamma-f)}.$$

Let

$$\phi_t(z) = a(t)z + b(t)$$

where

$$a(0) = 1, \quad \partial_t a = -\frac{1}{2}(1+\gamma^2)a$$

and

$$b(0) = 0, \quad \partial_t b = -\frac{1}{2}(1+\gamma^2)a\gamma.$$

Set

$$\begin{aligned} h_t &:= \phi_t \circ f_t, \\ \beta_t &:= \phi_t(\gamma(t)). \end{aligned}$$

Then (and this is the reason for the choice of the functions a and b)

$$\partial_t h = -\frac{1}{2}a \frac{(1+\gamma^2)^2}{\gamma-f} = -\frac{\frac{1}{2}(1+\gamma^2)^2 a^2}{\beta-h}.$$

h_t is also a conformal map from $\mathbf{U} \setminus K_t$ onto the upper half-plane with $h_t(1) = \infty$. Note also that $h_0(z) = \psi(z)$. We introduce a new time parameter $u = u(t)$ by setting

$$\partial_t u = \frac{1}{4}(1+\gamma^2)^2 a^2, \quad u(0) = 0.$$

Then

$$\frac{\partial h}{\partial u} = \frac{-2}{\beta-h}.$$

Since this is the equation defining the chordal SLE₆-process, it remains to show that $u \mapsto \beta_{t(u)}/\sqrt{\kappa}$ is Brownian motion (stopped at some random time). This is a direct but tedious application of Itô's formula:

$$d\gamma_t = \frac{1}{2}(1+\gamma^2)\sqrt{\kappa} dB_t + \frac{1}{2}\gamma(1+\gamma^2)(\frac{1}{2}\kappa - 1) dt$$

and

$$d\beta_t = \frac{1}{2}(1+\gamma^2)a(\sqrt{\kappa} dB_t + (-3 + \frac{1}{2}\kappa)\gamma dt).$$

This proves the claim, and establishes the theorem. \square

Note that when $\kappa \neq 6$, even though $u \rightarrow \beta$ is not a local martingale, its law is absolutely continuous with respect to that of $\sqrt{\kappa}$ times a Brownian motion, as long as γ and u remain bounded. More precisely:

PROPOSITION 4.2. *Let $(K_t, t \geq 0)$, $(\tilde{K}_u, u \geq 0)$, T and \tilde{T} be defined just as in Theorem 4.1, except that they are SLE_κ with general $\kappa > 0$. There exist two non-decreasing families of stopping times, $(T_n, n \geq 1)$ and $(\tilde{T}_n, n \geq 1)$, such that almost surely $T_n \rightarrow T$ and $\tilde{T}_n \rightarrow \tilde{T}$ when $n \rightarrow \infty$, and such that for each $n \geq 1$, the laws of $(K_t, t \in [0, T_n])$ and $(\tilde{K}_u, u \in [0, \tilde{T}_n])$ are equivalent (in the sense that they have a positive density with respect to each other) modulo increasing time change.*

Proof. It suffices to take

$$T_n = \min\{n, \inf\{t > 0 : |\delta_t - e_t| < 1/n\}\}.$$

Then, it is easy to see that before T_n , $|\gamma|$ remains bounded, a is bounded away from 0 (note also that $a \leq 1$ always), so that t/u is bounded and bounded away from 0. Hence, $u(T_n)$ is also bounded (since $T_n \leq n$).

It now follows directly from Girsanov's theorem (see, e.g., [38]) that the law of $(\beta(u)/\sqrt{\kappa})_{u \leq u(T_n)}$ is equivalent to that of Brownian motion up to some (bounded) stopping time, and Proposition 4.2 follows. \square

4.2. The crossing exponent for SLE_6

In this section, we are going to study the probability that chordal SLE_6 started at some point on the left-hand side of a rectangle crosses the rectangle from the left to the right without touching the upper and lower boundaries of the rectangle. As we shall see, the estimate obtained is a direct consequence of Cardy's formula for SLE_6 proved in [27].

The notation turns out to be simpler when considering crossings of a quadrilateral in the unit disk, which is equivalent to the rectangle case by conformal invariance. We now describe the setup more precisely. Recall that when $w, w' \in \partial\mathbf{U}$, the counterclockwise




Fig. 4.1. A successful crossing.

arc from w to w' is denoted $a(w, w')$. Let $\theta \in (0, \frac{1}{2}\pi)$, $\alpha \in (-1, 1)$, and define the points

$$\begin{aligned} w_1 &:= \exp(i(\pi - \theta)), \\ w'_1 &:= \exp(i(\pi + \alpha\theta)), \\ w_2 &:= \exp(i(\pi + \theta)), \\ w_3 &:= \exp(-i\theta), \\ w_4 &:= \exp(i\theta), \end{aligned}$$

and note that they appear in counterclockwise order on $\partial\mathbf{U}$. See Figure 4.1. Let w'_3 be any point in $a(w_3, w_4)$. Consider a chordal SLE_6 in \mathbf{U} started from w'_1 and growing towards w'_3 . Let K_t be the hull, and let T be the first time t such that \bar{K}_t intersects the arc $a(w_3, w_4)$. Set

$$\mathfrak{K} := \bigcup_{t < T} \bar{K}_t.$$

As shown in [27], the restriction property for SLE_6 shows that up to a monotone time change, the law of the process $(K_t)_{t < T}$ does not depend on the choice of w'_3 . Since we use the restriction property, the result derived in this subsection is specific to $\kappa=6$.

We are interested in the event

$$\mathcal{E} := \{\mathfrak{K} \cap (a(w_2, w_3) \cup a(w_4, w_1)) = \emptyset\}.$$

LEMMA 4.3. *Suppose that $\alpha_0 \in (0, 1)$ is fixed. When $\theta \searrow 0$,*

$$\mathbf{P}[\mathcal{E}] \asymp \theta^2,$$

and this estimate is uniform for $\alpha \in (-\alpha_0, \alpha_0)$. Moreover, when $\theta \searrow 0$,

$$\max_{\alpha \in (-1, 1)} \mathbf{P}[\mathcal{E}] \asymp \theta^2.$$

Recall also that $\theta^2 \sim \exp(-\ell)$ as $\theta \searrow 0$, where $\ell = \ell(a(w_1, w_2), a(w_3, w_4); \mathbf{U})$ denotes the π -extremal distance between $a(w_1, w_2)$ and $a(w_3, w_4)$ in \mathbf{U} .

The lemma is hardly surprising. Suppose for a moment that we take $w'_1 = -1$, and let T' be the first time such that \bar{K}_t intersects $a(-i, i)$. Let z be the point in $a(-i, i)$ which is on the boundary of $\bigcup_{t < T'} K_t$. It is then easy to believe that the probability density for the location of z should be bounded away from 0 and infinity in every compact subset of $a(-i, i)$. By conformal invariance, this implies the first part of the lemma for the case $\alpha = 0$. Although it should not be too hard to prove the lemma with some general arguments such as these, we find it easier to rely on the more refined results from [27].

Proof. Define the events

$$\begin{aligned} \mathcal{E}' &= \{ \bar{K}_t \text{ hits } a(w_2, w_4) \text{ before } a(w_4, w_1) \}, \\ \mathcal{E}'' &= \{ \bar{K}_t \text{ hits } a(w_2, w_3) \text{ before } a(w_3, w_1) \}. \end{aligned}$$

Note that $\mathcal{E}'' \subset \mathcal{E}'$, and that $\mathcal{E} = \mathcal{E}' \setminus \mathcal{E}''$. Cardy's formula for SLE₆ derived in [27] (Theorem 3.2 in the case where $b = 0$) gives the exact value of $\mathbf{P}(\mathcal{E}')$ and $\mathbf{P}(\mathcal{E}'')$, as follows. Define the cross ratios

$$c' := \frac{(w_1 - w'_1)(w_4 - w_2)}{(w_4 - w'_1)(w_1 - w_2)}, \quad c'' := \frac{(w_1 - w'_1)(w_3 - w_2)}{(w_3 - w'_1)(w_1 - w_2)},$$

and set

$$G(x) := {}_2F_1\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}; x\right) \frac{\sqrt{\pi}}{2^{1/3} \Gamma(\frac{1}{3}) \Gamma(\frac{7}{6})} x^{1/3}$$

(where ${}_2F_1$ denotes the hypergeometric function). Then

$$\mathbf{P}[\mathcal{E}'] = G(c'), \quad \mathbf{P}[\mathcal{E}''] = G(c''), \quad \mathbf{P}[\mathcal{E}] = G(c') - G(c'').$$

(To compute $\mathbf{P}[\mathcal{E}']$, we view K_t as an SLE₆ from w'_1 to w_4 , while to compute $\mathbf{P}[\mathcal{E}'']$, we view K_t as an SLE₆ from w'_1 to w_3 . As remarked above, the choice of $w'_3 \in a(w_3, w_4)$ does not matter.) Note that

$$c' = \cot \theta \tan\left(\frac{1}{2}(1 + \alpha)\theta\right), \quad c'' = \frac{\sin\left(\frac{1}{2}(1 + \alpha)\theta\right)}{\sin \theta \cos\left(\frac{1}{2}(1 - \alpha)\theta\right)}.$$

Both c' and c'' converge to $\frac{1}{2}(1 + \alpha)$ when $\theta \searrow 0$ and $c' - c'' \sim \frac{1}{4}(1 - \alpha^2)\theta^2$. Since $G'(x) = (\frac{1}{3}\sqrt{\pi}) 2^{-1/3} \Gamma(\frac{1}{3})^{-1} \Gamma(\frac{7}{6})^{-1} ((1-x)x)^{-2/3}$, it follows that

$$\mathbf{P}[\mathcal{E}] = G(c') - G(c'') \sim \left(\frac{1}{6}\sqrt{\pi}\right) \Gamma\left(\frac{1}{3}\right)^{-1} \Gamma\left(\frac{7}{6}\right)^{-1} (1 - \alpha^2)^{1/3} \theta^2,$$

as $\theta \searrow 0$, and the lemma follows. □

5. Brownian intersection exponents

5.1. Definitions and statement of results

This section begins with a review of the definitions and some general facts concerning intersection exponents between planar Brownian paths, and proceeds with a statement of some theorems. For more details concerning the background results on intersection exponents, as well as some references to the literature, see [31], [32].

Suppose that $n+m$ independent Brownian motions B^1, \dots, B^n and B'^1, \dots, B'^m are started from points $B^1(0)=\dots=B^n(0)=i$ and $B'^1(0)=\dots=B'^m(0)=1$ in the complex plane, and let $\mathfrak{B}_r^j, \mathfrak{B}_r'^k$ denote the traces (i.e., images) of the paths up to the first time they reach the circle C_r . Consider the probability $f_{n,m}(r)$ that the \mathfrak{B} -traces do not intersect the \mathfrak{B}' -traces, i.e.,

$$f_{n,m}(r) := \mathbf{P} \left[\left(\bigcup_{j=1}^n \mathfrak{B}_r^j \right) \cap \left(\bigcup_{l=1}^m \mathfrak{B}_r'^l \right) = \emptyset \right].$$

It is easy to see that as $r \rightarrow \infty$ this probability decays like a power law; the (n, m) -intersection exponent $\xi(n, m)$ is defined by

$$f_{n,m}(r) = r^{-\xi(n,m)+o(1)}, \quad r \rightarrow \infty.$$

We call $\xi(n, m)$ the intersection exponent between one packet of n Brownian motions and one packet of m Brownian motions. It is easy to see (e.g. [31]) that the exponent $\zeta(n, m)$ described in the introduction is equal to $\frac{1}{2}\xi(n, m)$.

By using the conformal map $z \mapsto 1/z$ and invariance of planar Brownian motion under conformal mapping, it is clear that

$$f_{n,m}(r) = f_{n,m}(1/r),$$

so that the exponents also measure the decay of $f_{n,m}$ when $r \searrow 0$.

Similarly, one can define corresponding probabilities for intersection exponents in a half-plane,

$$\tilde{f}_{n,m}(r) := \mathbf{P} \left[\left(\bigcup_{j=1}^n \mathfrak{B}_r^j \right) \cap \left(\bigcup_{l=1}^m \mathfrak{B}_r'^l \right) = \emptyset \text{ and } \left(\bigcup_{j=1}^n \mathfrak{B}_r^j \right) \cup \left(\bigcup_{l=1}^m \mathfrak{B}_r'^l \right) \subset H \right],$$

where H is some half-plane through the origin containing 1 and i . ($\tilde{f}_{n,m}(r)$ does depend on H .) In plain words, we are looking at the probability that all Brownian motions stay in the half-plane H and that all the \mathfrak{B} -traces avoid all the \mathfrak{B}' -traces. It is also easy to see that there exists a $\tilde{\xi}(n, m)$ (which does not depend on H) such that

$$\tilde{f}_{n,m}(r) = r^{-\tilde{\xi}(n,m)+o(1)}$$

when $r \rightarrow \infty$.

These intersection exponents can be generalized in a number of ways. For example, set

$$Z_r := \mathbf{P}\left[\mathfrak{B}_r^{1/1} \cap \bigcup_{j=1}^n \mathfrak{B}_r^j = \emptyset \mid \bigcup_{j=1}^n \mathfrak{B}_r^j\right].$$

Then $f_{n,m}(r) = \mathbf{E}[Z_r^m]$, and define $\xi(n, \lambda)$ for all $\lambda > 0$ by the relation

$$\mathbf{E}[Z_r^\lambda] \approx r^{-\xi(n,\lambda)}, \quad r \rightarrow \infty.$$

It has been proved by Lawler [24] that in fact

$$\mathbf{E}[Z_r^\lambda] \asymp r^{-\xi(2,\lambda)}.$$

The same proof, with minor notational modifications, applies to the other exponents as well. See also [30] for a new self-contained proof of this result.

One can also define the exponents $\xi(n_1, n_2, \dots, n_k)$ and $\tilde{\xi}(n_1, n_2, \dots, n_k)$ describing the probability of non-intersection of k packets of Brownian motions. In fact, [31] proves that there is a natural and rigorous way to generalize these definitions to the case where the numbers n_j are positive reals (i.e., not required to be integers). For the definition of $\xi(\lambda_1, \dots, \lambda_k)$ one only needs to assume that at least two of the numbers $\lambda_1, \dots, \lambda_k$ are at least 1, and for $\tilde{\xi}$ even this assumption is not necessary. What makes these extended definitions natural is that they are uniquely determined by certain identities and relations. For one, the functions ξ and $\tilde{\xi}$ are invariant under permutations of their arguments. Moreover, they satisfy the so-called cascade relations [31]: for any $1 \leq q \leq m-1$ and $(\lambda_1, \dots, \lambda_m)$ such that $\lambda_1 \geq 1$ and $\max\{\lambda_2, \dots, \lambda_m\} \geq 1$,

$$\xi(\lambda_1, \dots, \lambda_m) = \xi(\lambda_1, \dots, \lambda_q, \tilde{\xi}(\lambda_{q+1}, \dots, \lambda_m)). \tag{5.1}$$

In [31] it was also established that the cascade relations imply the existence of a continuous increasing function $\eta: [\tilde{\xi}(1, 1), \infty) \rightarrow [\xi(1, 1), \infty)$ such that for all $(\lambda_1, \dots, \lambda_m) \in \mathbf{R}_+^m$ such that at least two of the λ_j 's are at least 1,

$$\xi(\lambda_1, \dots, \lambda_m) = \eta(\tilde{\xi}(\lambda_1, \dots, \lambda_m)). \tag{5.2}$$

In [27], we have determined the exact value of exponents $\tilde{\xi}(\lambda_1, \dots, \lambda_m)$ for a certain class of numbers $(\lambda_1, \dots, \lambda_m)$; namely, for all $m \geq 2$, and all $(\lambda_1, \dots, \lambda_m) \in \{\frac{1}{6}l(l+1) : l \in \mathbf{N}\}^{m-1} \times \mathbf{R}_+$,

$$\tilde{\xi}(\lambda_1, \dots, \lambda_m) = \frac{1}{24} \left((\sqrt{24\lambda_1+1} + \dots + \sqrt{24\lambda_m+1} - (m-1))^2 - 1 \right). \tag{5.3}$$

In particular,

$$\tilde{\xi}(1, 1, \lambda) = \frac{1}{24} \left((8 + \sqrt{24\lambda+1})^2 - 1 \right), \tag{5.4}$$

$$\tilde{\xi}(1, 1, 1, \lambda) = \frac{1}{24} \left((12 + \sqrt{24\lambda+1})^2 - 1 \right). \tag{5.5}$$

In the next section, we will prove the following two results:

THEOREM 5.1.

$$\xi(1, 1) = \frac{5}{4}. \quad (5.6)$$

THEOREM 5.2. For every $\lambda \geq 0$,

$$\xi(1, 1, 1, \lambda) = \frac{1}{48} \left((11 + \sqrt{24\lambda + 1})^2 - 4 \right). \quad (5.7)$$

Let us now see what these two results directly imply:

THEOREM 5.3. For all $x \geq 7$,

$$\eta(x) = \frac{1}{48} \left((\sqrt{24x + 1} - 1)^2 - 4 \right). \quad (5.8)$$

Moreover, for all $m \geq 2$ and for all $(\lambda_1, \dots, \lambda_m) \in \{\frac{1}{6}l(l+1) : l \in \mathbf{N}\}^{m-1} \times \mathbf{R}_+$,

$$\xi(\lambda_1, \dots, \lambda_m) = \frac{1}{48} \left((\sqrt{24\lambda_1 + 1} + \dots + \sqrt{24\lambda_m + 1} - m)^2 - 4 \right), \quad (5.9)$$

provided that at least two of the numbers $\lambda_1, \dots, \lambda_m$ are at least 1, and the right-hand side of (5.9) is at least $\frac{35}{12}$.

In particular, for all $\lambda \geq \frac{10}{3}$ and $\lambda' \geq 2$,

$$\xi(1, \lambda) = \frac{1}{48} \left((3 + \sqrt{24\lambda + 1})^2 - 4 \right), \quad (5.10)$$

$$\xi(2, \lambda') = \frac{1}{48} \left((5 + \sqrt{24\lambda' + 1})^2 - 4 \right), \quad (5.11)$$

and for all integers $m \geq 2$ (using (5.6) for the case $m=2$),

$$\xi(1^{\otimes m}) = \frac{1}{12} (4m^2 - 1), \quad (5.12)$$

where $1^{\otimes m} := (1, \dots, 1) \in \mathbf{N}^m$.

Proof of Theorem 5.3 (assuming (5.6) and (5.7)). Combining (5.5), (5.7) and (5.2) gives (5.8). Hence, we get (5.9) from (5.3), (5.2) and (5.8). \square

Proof of Theorems 1.1 and 1.2 (assuming (5.6) and (5.7)). In view of the fact that the time exponents differ by a factor of 2 from the space exponents, the theorems follow from Theorem 5.3. \square

Remark. In [28], we will show that (5.3) holds for all $(\lambda_1, \dots, \lambda_m) \in \mathbf{R}_+^m$. The proofs in the present paper can then be very easily adapted to show that (5.9) holds for all $(\lambda_1, \dots, \lambda_m) \in \mathbf{R}_+^m$ such that at least two of the λ_j 's are at least 1.

5.2. Excursion measures

In [31], [32] a characterization of the intersection exponents was given in terms of excursions. The Brownian excursion measures are natural and interesting objects. The utility of the excursion measures in [31], [32] arises from the fact that in the context of excursion measures, one can study Brownian motions without specifying the starting point. This significantly simplifies some arguments and estimates.

Roughly speaking, Brownian excursions in a domain D are Brownian motions started on the boundary, conditioned to immediately enter D , and stopped upon leaving. Since Brownian excursions stay in the domain D , conformal transformations of D can be applied to the excursions.

Let us now describe these Brownian excursions more precisely. For any bounded simply-connected open domain D , there exists a Brownian excursion measure μ_D in D . This is an infinite measure on the set of paths $(B(t), t \leq \tau)$ in D such that $B(0, \tau) \subset D$ and $B(0), B(\tau) \in \partial D$ (these endpoints can be viewed as prime ends if necessary). $x_s := B(0)$ and $x_e := B(\tau)$ will denote the starting point and terminal point of the excursion. In this discussion, we will identify two paths (two excursions) when one is obtained by an increasing time change of the other.

One possible definition of μ_D is the following. Consider first $D = \mathbf{U}$, the unit disc. For every $s \in (0, 1)$ let P^s be the law of a Brownian motion started uniformly on the circle of radius s , and stopped when it exits \mathbf{U} (modulo continuous increasing time change). Since $z \mapsto \log |z|$ is harmonic, for any $r \in (0, s)$,

$$P^s[B \text{ hits } C_r] = \frac{\log(1/s)}{\log(1/r)}.$$

Set

$$\mu_{\mathbf{U}} := \lim_{s \nearrow 1} \frac{2\pi}{\log(1/s)} P^s,$$

as a weak limit. Note that the $\mu_{\mathbf{U}}$ -measure of the set of paths that hit the circle C_r is $2\pi/\log(1/r)$.

One can then check that for any Möbius transformation ϕ from \mathbf{U} onto \mathbf{U} , $\phi(\mu_{\mathbf{U}}) = \mu_{\mathbf{U}}$. This makes it possible to extend the definition of μ_D to any simply-connected domain D , by conformal invariance. These Brownian excursions also have a “restriction” property [32], which is a result of the fact that the Brownian paths only feel the boundary of D when they hit it (and then stop).

In [27], we made an extensive use of the Brownian excursion measure in rectangles $R_L = (0, L) \times (0, \pi)$. It is easy to see that the measure μ_{R_L} , restricted to those excursions with starting point on the left-hand side of the rectangle $[0, i\pi]$, is obtained as the limit when $s \rightarrow 0$ of $\pi s^{-1} P_L^s$, where P_L^s is the law of a Brownian motion with uniform starting

point on $[s, s+i\pi]$ which is stopped when it exits R_L . In particular, this leads to the following result:

LEMMA 5.4. *Let \mathcal{E}_L denote the event that the Brownian excursion B in R_L crosses the rectangle from the left to the right (i.e., $x_s \in (0, i\pi)$ and $x_e \in (L, L+i\pi)$). Then, when $L \rightarrow \infty$,*

$$\mu_{R_L}[\mathcal{E}_L] \asymp e^{-L}. \tag{5.13}$$

Proof. Let h_z denote harmonic measure from z on ∂R_L , where $z \in R_L$. Since $\text{Im}(\exp z)$ is a harmonic function, it easily follows that for all $L > 1$ and $z \in (1, 1+i\pi)$,

$$\sin\left(\frac{1}{4}\pi\right) h_z\left(\left[L + \frac{1}{4}i\pi, L + \frac{3}{4}i\pi\right]\right) \leq e^{-L} \leq \frac{h_z([L, L+i\pi])}{\text{Im}(\exp(z)) - 1}.$$

It is easy to verify (e.g., by conformal invariance or by a reflection argument) that there is a constant c such that $h_z([L, L+i\pi]) \leq ch_z([L + \frac{1}{4}i\pi, L + \frac{3}{4}i\pi])$ holds for all $L > 2$ and $z \in (1, 1+i\pi)$. Hence, for such L and z ,

$$c_1 h_z([L, L+i\pi]) \leq e^{-L} \leq c_2 h_z([L, L+i\pi]) 1_{\{\text{Im}(z) \in [\frac{1}{4}\pi, \frac{3}{4}\pi]\}},$$

with some constants $c_1, c_2 > 0$. Since the μ_{R_L} -measure of the excursions which reach the line $\{\text{Re}(z)=1\}$ does not depend on L , and from these a fixed proportion first hit $\{\text{Re}(z)=1\}$ in $[1 + \frac{1}{4}\pi, 1 + \frac{3}{4}\pi]$, (5.13) now follows from the Markov property, which is valid for the excursion measures. □

Similarly, one can define Brownian excursions in non-simply-connected domains. For instance, consider the annulus $A(r, 1)$ bounded between the circles C_r and C_1 (where $r \in (0, 1)$). Rather than defining the excursion in $A(r, 1)$ directly, we base the definition on the excursions in \mathbf{U} . If γ is a path, let $\Psi_r(\gamma)$ be the initial segment of γ , until the first hit of C_r , or all of γ , if γ does not hit C_r . Now set $\mu_1^r := \Psi_r(\mu_{\mathbf{U}})$. This will be called the Brownian excursion measure on $A(r, 1)$ for excursions started on C . It is clear that $\mu_{\mathbf{U}} = \lim_{r \searrow 0} \mu_1^r$.

The measures μ_D and μ_1^r are also related by restriction and conformal invariance. Suppose that O is a simply-connected subset of $A(r, 1)$ such that each of the sets $\bar{O} \cap C_r$ and $\bar{O} \cap C$ is an arc of positive length. Let L denote the π -extremal distance between these two arcs in O , and let ϕ denote the conformal map from O onto $R_L = (0, L) \times (0, \pi)$ such that $\bar{O} \cap C$ corresponds to $(0, i\pi)$ and $\bar{O} \cap C_r$ corresponds to $(L, L+i\pi)$ under ϕ .

Let $\hat{\mathcal{E}}_1$ be the set of paths starting in C that reach C_r without exiting \bar{O} . Consider the image under ϕ of the measure μ_1^r restricted to $\hat{\mathcal{E}}_1$. Then (up to time change) the

image measure is exactly μ_{R_L} restricted to the set of excursions that cross the rectangle from the left to the right. (5.13) therefore shows that when $L \rightarrow \infty$,

$$\mu_1^r[\hat{\mathcal{E}}_1] \asymp \exp(-L). \tag{5.14}$$

This may also be easily verified directly.

The mapping $z \mapsto r/z$ maps $A(r, 1)$ conformally onto itself. Let μ_r^1 be the image of μ_1^r under this map; this is a measure on Brownian excursions started on C_r and stopped upon leaving $A(r, 1)$. By symmetry, it follows that $\mu_r^1[\hat{\mathcal{E}}_1] = \mu_1^r[\hat{\mathcal{E}}_1] \asymp \exp(-L)$ when $L \rightarrow \infty$.

Although we will not use this here, it is worthwhile to note that the measures μ_r^1 and μ_1^r agree on the set of paths crossing the annulus, up to time reversal of the path.

5.3. Exponents and excursions

We now describe the intersection exponents in terms of excursions (referring to [31], [32] for the proofs). Let $(B(t), t \leq \tau)$ be an excursion in R_L , and, as above, let

$$\mathcal{E}_L := \{\text{Re}(x_s) = 0 \text{ and } \text{Re}(x_e) = L\}$$

be the event that B crosses R_L from left to right. Let \mathfrak{B} denote the image of B . When \mathcal{E}_L holds, let O_B^+ be the component of $R_L \setminus \mathfrak{B}$ above \mathfrak{B} , and let O_B^- be the component of $R_L \setminus \mathfrak{B}$ below \mathfrak{B} . Let \mathfrak{L}_B^- (resp. \mathfrak{L}_B^+) denote the π -extremal distance between $[0, x_s]$ and $[L, x_e]$ in O_B^- (resp. $[x_s, i\pi]$ and $[x_e, L+i\pi]$ in O_B^+). (We use script fonts for these \mathfrak{L} to indicate that they are random variables.) Then, for any $\alpha \geq 0$ and $\alpha' \geq 0$, the exponent $\tilde{\xi}(\alpha, 1, \alpha') = \tilde{\xi}(1, \tilde{\xi}(\alpha, \alpha'))$ is characterized by

$$\int_{\mathcal{E}_L} \exp(-\alpha \mathfrak{L}_B^+ - \alpha' \mathfrak{L}_B^-) d\mu_{R_L}(B) = \exp(-\tilde{\xi}(\alpha', 1, \alpha)L + o(L)), \tag{5.15}$$

as $L \rightarrow \infty$.

Let \mathcal{E}_L^2 be the set of pairs of paths $(B, B') \in \mathcal{E}_L \times \mathcal{E}_L$ such that the trace \mathfrak{B}' of B' is contained in O_B^- . It follows from (5.15), Lemma 5.4 and conformal invariance of the excursion measures that

$$\int_{\mathcal{E}_L^2} \exp(-\alpha \mathfrak{L}_B^+) d\mu_{R_L}(B) d\mu_{R_L}(B') = \exp(-\tilde{\xi}(1, 1, \alpha)L + o(L)), \tag{5.16}$$

as $L \rightarrow \infty$. On \mathcal{E}_L^2 , let $\mathfrak{L}_{B'}^B$ be the π -extremal distance from $[0, i\pi]$ to $[L, L+i\pi]$ in the domain between \mathfrak{B} and \mathfrak{B}' . Given B' , it is clear by conformal invariance and the restriction property of the excursion measure that $1_{\mathcal{E}_L^2} \mathfrak{L}_{B'}^B$ has the same law as $1_{\mathcal{E}_L^2} \mathfrak{L}_B^+$. Consequently, (5.16) gives

$$\int_{\mathcal{E}_L^2} \exp(-\alpha \mathfrak{L}_{B'}^B) d\mu_{R_L}(B) d\mu_{R_L}(B') = \exp(-\tilde{\xi}(1, 1, \alpha)L + o(L)). \tag{5.17}$$

Similarly, one can characterize the exponents ξ in terms of the excursion measure μ_r^1 . For any $r < 1$, consider two independent excursions B and B' of the annulus $A(r, 1)$. Define the events

$$\begin{aligned} \mathcal{E} = \mathcal{E}(r) &:= \{\mathfrak{B} \text{ crosses the annulus without separating } C_r \text{ from } C\}, \\ \tilde{\mathcal{E}} = \tilde{\mathcal{E}}(r) &:= \{\mathfrak{B} \text{ and } \mathfrak{B}' \text{ are disjoint and both cross the annulus}\}. \end{aligned}$$

When \mathcal{E} holds, let \mathfrak{L}_B be the π -extremal distance between C_r and C in $A(r, 1) \setminus \mathfrak{B}$. Similarly, when $\tilde{\mathcal{E}}$ is satisfied, let O_1 and O_2 be the two components of $A(r, 1) \setminus (\mathfrak{B} \cup \mathfrak{B}')$ which have arcs of C on their boundaries, in such a way that the sequence $\mathfrak{B}, O_1, \mathfrak{B}', O_2$ is in counterclockwise order around $A(r, 1)$. Let \mathfrak{L}_1 be the π -extremal distance from C_r to C in O_1 , and let \mathfrak{L}_2 be the corresponding quantity for O_2 . Then for $\lambda, \lambda_1, \lambda_2 \in \mathbf{R}_+$, the exponents $\xi(1, \lambda)$ and $\xi(1, \lambda_1, 1, \lambda_2)$ can be described [32] as

$$\int_{\mathcal{E}} \exp(-\lambda \mathfrak{L}) d\mu_r^1(B) \approx r^{\xi(1, \lambda)} \tag{5.18}$$

and

$$\int_{\tilde{\mathcal{E}}} \exp(-\lambda_1 \mathfrak{L}_1 - \lambda_2 \mathfrak{L}_2) d\mu_r^1(B) d\mu_r^1(B') \approx r^{\xi(1, \lambda_1, 1, \lambda_2)}, \tag{5.19}$$

as $r \searrow 0$.

5.4. A useful technical lemma

We now derive a technical refinement of (5.18) and (5.19) (in the case $\lambda_1 = 1$) that will be useful to identify the Brownian intersection exponents with those computed for SLE_6 .

Keep the same notation as above, and on \mathcal{E} , let ϕ denote the conformal map that maps O onto $R_{\mathcal{E}}$ in such a way that the images of $C_1 \cap \bar{O}$ and $C_r \cap \bar{O}$ are mapped onto the left and right edges of the rectangle, respectively. Similarly, on $\tilde{\mathcal{E}}$, let ϕ_1 be the corresponding conformal map from O_1 onto $R_{\mathcal{E}_1}$. For all $\alpha > 0$ set

$$\begin{aligned} \mathcal{H}_\alpha &:= \mathcal{E} \cap \{i \in \bar{O} \text{ and } \phi(i) \in [i\alpha, i(\pi - \alpha)]\}, \\ \tilde{\mathcal{H}}_\alpha &:= \tilde{\mathcal{E}} \cap \{i \in \bar{O}_1 \text{ and } \phi_1(i) \in [i\alpha, i(\pi - \alpha)]\}. \end{aligned}$$

LEMMA 5.5. *Let $\lambda > 0$. Then there are sequences $x_n \searrow 0$ and $y_n \searrow 0$, and an $\alpha > 0$, such that*

$$\int_{\mathcal{H}_\alpha} \exp(-\lambda \mathfrak{L}_B) d\mu_{x_n}^1(B) \approx (x_n)^{\xi(1, \lambda)}, \quad n \rightarrow \infty, \tag{5.20}$$

$$\int_{\tilde{\mathcal{H}}_\alpha} \exp(-\mathfrak{L}_1 - \lambda \mathfrak{L}_2) d\mu_{y_n}^1(B) d\mu_{y_n}^1(B') \approx (y_n)^{\xi(1, 1, 1, \lambda)}, \quad n \rightarrow \infty. \tag{5.21}$$

Actually (see, e.g., [30]), much stronger statements hold: In the above, \approx may be replaced by \asymp , and these statements hold for every sequence tending to ∞ . But the present statement will be sufficient to determine the values of the Brownian intersection exponents, and it can be easily proved as follows.

Proof. We will only give the detailed proof of (5.21). The proof of (5.20) is easier, follows exactly the same lines, and is safely left to the reader.

Because of (5.19), it suffices to find the lower bound for the left-hand side of (5.21). Let us first introduce some notation. Let $r \in (0, \frac{1}{4})$, and consider the measure $\mu_r^1 \times \mu_r^1$ on the space of pairs (B, B') . Let

$$f(r) := \int_{\tilde{\mathcal{E}}(r)} \exp(-\mathcal{L}_1 - \lambda \mathcal{L}_2) d\mu_r^1(B) d\mu_r^1(B').$$

Let B^* be the path B stopped when it hits $C_{1/4}$, if it does, and $B^* = B$, if it does not hit $C_{1/4}$. Similarly, define B'^* from B' . Let \mathfrak{B} and \mathfrak{B}' be the traces of B and B' , respectively. Let $\tilde{\mathcal{E}}^*$ be the event that the traces of B^* and B'^* are disjoint. On $\tilde{\mathcal{E}}^*$, let O_1^* and O_2^* be the domains defined for B^* and B'^* , as O_1 and O_2 were defined for B and B' . Then $O_j^* \subset O_j$ on $\tilde{\mathcal{E}}$, $j=1, 2$. For $j=1, 2$, on $\tilde{\mathcal{E}}^*$, let \mathcal{L}_j^* be the π -extremal distance between C_r and $C_{1/4}$ in O_j^* . Otherwise, set $\mathcal{L}_j^* = \infty$.

For $a > 0$, let \mathcal{D}_a be the event that the distance between $\mathfrak{B} \cap A(\frac{1}{2}, 1)$ and $\mathfrak{B}' \cap A(\frac{1}{2}, 1)$ is at least a . Suppose that $a \in (0, \frac{1}{5})$. Observe that for $(B, B') \notin \mathcal{D}_a$, there is for $j=1$ or $j=2$ a path of length at most a in $O_j \cap A(\frac{1}{2}, 1)$ which separates C_r from C_1 in O_j . It then follows that $\mathcal{L}_j \geq \mathcal{L}_j^* + c_1 \log(1/a)$, for some constant $c_1 > 0$. Consequently,

$$\begin{aligned} & \int_{\tilde{\mathcal{E}}(r) \setminus \mathcal{D}_a} \exp(-\mathcal{L}_1 - \lambda \mathcal{L}_2) d\mu_r^1(B) d\mu_r^1(B') \\ & \leq a^{c_1 \min\{\lambda, 1\}} \int_{\tilde{\mathcal{E}}^*} \exp(-\mathcal{L}_1^* - \lambda \mathcal{L}_2^*) d\mu_r^1(B) d\mu_r^1(B') \quad (5.22) \\ & = a^{c_1 \min\{\lambda, 1\}} f(4r), \end{aligned}$$

since the image of μ_r^1 under the map $B \mapsto 4B^*$ is μ_{4r}^1 .

By (5.19), we have $f(r) \approx r^{\xi(1,1,\lambda,1)}$ (and f is non-decreasing). Consequently, if a is chosen sufficiently small, there is a sequence $y_n \searrow 0$ (for instance a subsequence of 4^{-n}) such that $f(y_n) \geq 2a^{c_1 \min\{\lambda, 1\}} f(4y_n)$. For these y_n , (5.22) gives

$$\int_{\tilde{\mathcal{E}}(y_n) \cap \mathcal{D}_a} \exp(-\mathcal{L}_1 - \lambda \mathcal{L}_2) d\mu_{y_n}^1(B) d\mu_{y_n}^1(B') \geq a^{c_1 \min\{\lambda, 1\}} f(4y_n). \quad (5.23)$$

Fix such an a and such a y_n . Let \mathcal{I}_a be the event that $i \in \bar{O}_1$ and the distance from i to $\mathfrak{B} \cup \mathfrak{B}'$ is at least $\frac{1}{10}a$. Observe that if we apply an independent random uniform rotation

about 0 to a pair $(B, B') \in \mathcal{D}_a \cap \tilde{\mathcal{E}}$, then with probability at least $a/10\pi$ the rotated pair is in \mathcal{I}_a . Since the integrand in (5.23) is invariant under rotations, (5.23) and (5.19) give

$$\int_{\tilde{\mathcal{E}}(y_n) \cap \mathcal{I}_a} \exp(-\mathcal{L}_1 - \lambda \mathcal{L}_2) d\mu_{y_n}^1(B) d\mu_{y_n}^1(B') \approx (y_n)^{\xi(1,1,\lambda,1)}.$$

It therefore suffices to show that when $\alpha > 0$ is small, we have $\mathcal{I}_a \cap \tilde{\mathcal{E}} \subset \tilde{\mathcal{H}}_\alpha$. To prove this, consider a pair $(B, B') \in \mathcal{I}_a$. Let A be the subarc of $\bar{O}_1 \cap C_1$ that has i as one endpoint and the other endpoint is in \mathfrak{B} , and let A' be the subarc of $\bar{O}_1 \cap C_1$ that has i as one endpoint and the other endpoint is in \mathfrak{B}' . Then the extremal distance from A to \mathfrak{B}' in O_1 is bounded from below by a positive constant depending only on a , as is the extremal distance from A' to \mathfrak{B} in O_1 . Conformal invariance of extremal distance therefore shows that the distance from $\phi(i)$ to $\{0, i\pi\}$ is bounded from below by a constant depending only on a , which proves that $(B, B') \in \tilde{\mathcal{H}}_\alpha$, where $\alpha > 0$ depends only on a . □

6. The universality argument

We are now ready to combine the results derived so far to prove our main theorems. As in [27], we follow the universality ideas presented in [32]. First, Theorem 5.6 ($\xi(1, 1) = \frac{5}{4}$) will be proved, followed by Theorem 5.7 (giving $\xi(1, 1, 1, \lambda)$). As we have seen, Theorems 1.1 and 1.2 are immediate consequences.

6.1. Proof of $\xi(1, 1) = \frac{5}{4}$

Let $r > 0$, let K_t be a radial SLE₆-process in \mathbf{U} starting at i , and let $T = T(r)$ be the least t such that $\bar{K}_t \cap C_r \neq \emptyset$. Set $\mathfrak{K} := \bar{K}_T$, let \mathbf{P}_r denote the law of \mathfrak{K} , and let \mathbf{E}_r denote expectation with respect to this measure. As before, we let μ_r^1 denote the Brownian excursion measure in $A(r, 1)$ started from C_r , and let \mathfrak{B} denote the trace of the excursion B . We are interested in the event \mathcal{E}^* in which $\mathfrak{K} \cap \mathfrak{B} = \emptyset$ and \mathfrak{B} crosses $A(r, 1)$ (i.e., $\mathfrak{B} \cap C_1 \neq \emptyset$).

The proof will proceed by computing $(\mathbf{P}_r \times \mu_r^1)[\mathcal{E}^*]$ in two different ways. In the first computation, we begin by conditioning on \mathfrak{K} and then taking the expectation, while the second computation begins by conditioning on B .

When \mathfrak{B} does not separate C_r from C in \mathbf{U} , let O_B denote the connected component of $A(r, 1) \setminus \mathfrak{B}$ that touches both circles C_r and C , and \mathcal{L}_B the corresponding π -extremal distance.

When \mathfrak{K} does not disconnect C_r from C in \mathbf{U} , let $O_{\mathfrak{K}}$ denote the connected component of $A(r, 1) \setminus \mathfrak{K}$ that touches both circles, and $\mathcal{L}_{\mathfrak{K}}$ the corresponding π -extremal distance.

Suppose first that \mathfrak{R} is given and that $\mathfrak{L}_{\mathfrak{R}} < \infty$. Note that $\mathfrak{L}_{\mathfrak{R}} \rightarrow \infty$ as $r \searrow 0$. By the restriction property and conformal invariance of the Brownian excursion measure (see §5.2) and Lemma 5.4,

$$\mu_r^1[B \text{ crosses the annulus in } O_{\mathfrak{R}}] = \mu_{R_{\mathfrak{L}_{\mathfrak{R}}}}[\mathcal{E}_{\mathfrak{L}_{\mathfrak{R}}}] \asymp \exp(-\mathfrak{L}_{\mathfrak{R}}), \quad r \searrow 0,$$

where \mathcal{E}_L is as in Lemma 5.4. On the other hand, we know from Theorem 3.1, with $b=1$, $\kappa=6$, that

$$\mathbf{E}_r[\exp(-\mathfrak{L}_{\mathfrak{R}})] \approx r^{5/4}, \quad r \searrow 0.$$

Combining these two facts implies that

$$(\mathbf{P}_r \times \mu_r^1)[\mathcal{E}^*] \approx r^{5/4}, \quad r \searrow 0. \quad (6.1)$$

Suppose now that B is given and that it does cross the annulus without separating the disk C_r from C . The second part of Lemma 4.3 shows that there exists a constant $c > 0$ such that

$$\mathbf{P}_r[\mathcal{E}^*] \leq c \exp(-\mathfrak{L}_B).$$

Hence, combining this with (5.18) shows that

$$(\mathbf{P}_r \times \mu_r^1)[\mathcal{E}^*] \leq r^{\xi(1,1)+o(1)}. \quad (6.2)$$

On the other hand, suppose now that B is given and that $B \in \mathcal{H}_\alpha$, where \mathcal{H}_α is as in Lemma 5.5. Then by the first part of Lemma 4.3, there exists a constant c' such that

$$\mathbf{P}_r[\mathcal{E}^*] \geq c' \exp(-\mathfrak{L}_B).$$

Combining this with Lemma 5.5, we find that

$$(\mathbf{P}_{x_n} \times \mu_{x_n}^1)[\mathcal{E}^*] \geq (\mathbf{P}_{x_n} \times \mu_{x_n}^1)[\mathcal{H}_\alpha \cap \mathcal{E}^*] \geq (x_n)^{\xi(1,1)+o(1)}, \quad n \rightarrow \infty.$$

Comparing with (6.2) gives

$$(\mathbf{P}_{x_n} \times \mu_{x_n}^1)[\mathcal{E}^*] \approx (x_n)^{\xi(1,1)}, \quad n \rightarrow \infty.$$

Now, by (6.1),

$$(x_n)^{5/4} \approx (\mathbf{P}_{x_n} \times \mu_{x_n}^1)[\mathcal{E}^*] \approx (x_n)^{\xi(1,1)}$$

when $n \rightarrow \infty$, which proves $\xi(1,1) = \frac{5}{4}$.

6.2. The determination of $\xi(1, 1, 1, \lambda)$

The goal now is to prove (5.7). As $\lambda \mapsto \xi(1, 1, 1, \lambda) = \xi(1, 1, \tilde{\xi}(1, \lambda))$ is continuous on $[0, \infty)$ (see [31]), we can restrict ourselves to the case where $\lambda > 0$. The proof goes along similar lines as the proof of $\xi(1, 1) = \frac{5}{4}$. This time, we will consider two independent Brownian excursions B and B' in the annulus $A(r, 1)$, and one SLE_6 \mathfrak{K} as before. We are interested in the event that B, B' and \mathfrak{K} all cross the annulus, that they remain disjoint, and that B, \mathfrak{K} and B' are in clockwise order. Let us call this event l . Note that in this case, \mathfrak{K} crosses the annulus in O_1 , where we use the notations of §5.3.

We shall compute in two different ways the quantity

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mu_r^1(B) d\mu_r^1(B') d\mathbf{P}_r(\mathfrak{K}).$$

On the one hand, we know from conformal invariance and the restriction property of Brownian excursions and from (5.17) that when \mathfrak{K} crosses the annulus without separating C_r from C ,

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mu_r^1(B) d\mu_r^1(B') \approx \exp(-\mathcal{L}_{\mathfrak{K}} \tilde{\xi}(1, 1, \lambda)), \quad r \searrow 0. \tag{6.3}$$

But we know from Theorem 3.1 that

$$\int \exp(-b \mathcal{L}_{\mathfrak{K}}) d\mathbf{P}_r \approx r^\nu, \tag{6.4}$$

where

$$\nu = \nu(b) = \frac{1}{8} (4b + 1 + \sqrt{1 + 24b}).$$

Note that ν is a continuous increasing function of b on $(0, \infty)$. Consequently, (6.3) and (6.4) give

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mu_r^1(B) d\mu_r^1(B') d\mathbf{P}_r(\mathfrak{K}) \approx r^{\nu(\tilde{\xi}(1, 1, \lambda))}.$$

Combining this with (5.4) shows that

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mu_r^1(B) d\mu_r^1(B') d\mathbf{P}_r(\mathfrak{K}) \approx r^a, \tag{6.5}$$

where

$$a = a(\lambda) := \nu(\tilde{\xi}(1, 1, \lambda)) = \frac{1}{48} ((11 + \sqrt{24\lambda + 1})^2 - 4).$$

Suppose now that B and B' are given and that $\tilde{\mathcal{E}}$ is satisfied; i.e., that B and B' cross the annulus without intersecting each other. Then Lemma 4.3 shows that

$$\mathbf{P}_r[\mathfrak{K} \subset O_1] \leq c \exp(-\mathcal{L}_1),$$

for some constant c . Hence, combining this with (5.19) shows that

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mathbf{P}_r(\mathfrak{K}) d\mu_r^1(B) d\mu_r^1(B') \leq r^{\xi(1,1,1,\lambda)+o(1)}. \quad (6.6)$$

Suppose now that B and B' are given and that $(B, B') \in \tilde{\mathcal{H}}_\alpha$. Then Lemma 4.3 shows that there exists a constant $c' > 0$ such that

$$\mathbf{P}_r[\mathfrak{K} \subset O_1] \geq c' \exp(-\mathcal{L}_1).$$

Combining this with Lemma 5.5 gives

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mathbf{P}_{y_n}(\mathfrak{K}) d\mu_{y_n}^1(B) d\mu_{y_n}^1(B') \geq (y_n)^{\xi(1,1,1,\lambda)+o(1)}.$$

Comparing with (6.6) implies that

$$\int_l \exp(-\lambda \mathcal{L}_2) d\mathbf{P}_{y_n}(\mathfrak{K}) d\mu_{y_n}^1(B) d\mu_{y_n}^1(B') \approx (y_n)^{\xi(1,1,1,\lambda)},$$

when $n \rightarrow \infty$. Consequently, by (6.5), $\xi(1, 1, 1, \lambda) = a(\lambda)$. □

References

- [1] AHLFORS, L. V., *Conformal Invariants: Topics in Geometric Function Theory*. McGraw-Hill, New York, 1973.
- [2] AIZENMAN, M., DUPLANTIER, B. & AHARONY, A., Path crossing exponents and the external perimeter in 2D percolation. *Phys. Rev. Lett.*, 83 (1999), 1359–1362.
- [3] AZENCOTT, R., Behaviour of diffusion semi-groups at infinity. *Bull. Soc. Math. France*, 102 (1974), 193–240.
- [4] BISHOP, C. J., JONES, P. W., PEMANTLE, R. & PERES, Y., The dimension of the Brownian frontier is greater than 1. *J. Funct. Anal.*, 143 (1997), 309–336.
- [5] BURDZY, K. & LAWLER, G. F., Non-intersection exponents for Brownian paths. Part I: Existence and an invariance principle. *Probab. Theory Related Fields*, 84 (1990), 393–410.
- [6] — Non-intersection exponents for Brownian paths. Part II: Estimates and applications to a random fractal. *Ann. Probab.*, 18 (1990), 981–1009.
- [7] CARDY, J. L., Conformal invariance and surface critical behavior. *Nuclear Phys. B*, 240 (1984), 514–532.
- [8] — Critical percolation in finite geometries. *J. Phys. A*, 25 (1992), L201–L206.
- [9] — The number of incipient spanning clusters in two-dimensional percolation. *J. Phys. A*, 31 (1998), L105.
- [10] CRANSTON, M. & MOUNTFORD, T., An extension of a result of Burdzy and Lawler. *Probab. Theory Related Fields*, 89 (1991), 487–502.
- [11] DUPLANTIER, B., Random walks and quantum gravity in two dimensions. *Phys. Rev. Lett.*, 81 (1998), 5489–5492.
- [12] — Two-dimensional copolymers and exact conformal multifractality. *Phys. Rev. Lett.*, 82 (1999), 880–883.
- [13] — Harmonic measure exponents for two-dimensional percolation. *Phys. Rev. Lett.*, 82 (1999), 3940–3943.
- [14] DUPLANTIER, B. & KWON, K.-H., Conformal invariance and intersection of random walks. *Phys. Rev. Lett.*, 61 (1988), 2514–2517.
- [15] DUPLANTIER, B. & SALEUR, H., Exact determination of the percolation hull exponent in two dimensions. *Phys. Rev. Lett.*, 58 (1987), 2325–2328.
- [16] KENYON, R., Conformal invariance of domino tiling. *Ann. Probab.*, 28 (2000), 759–795.
- [17] — Long-range properties of spanning trees. Probabilistic techniques in equilibrium and nonequilibrium statistical physics. *J. Math. Phys.*, 41 (2000), 1338–1363.
- [18] — The asymptotic determinant of the discrete Laplacian. *Acta Math.*, 185 (2000), 239–286.
- [19] LAWLER, G. F., *Intersections of Random Walks*. Birkhäuser Boston, Boston, MA, 1991.
- [20] — Hausdorff dimension of cut points for Brownian motion. *Electron. J. Probab.*, 1:2 (1996), 1–20 (electronic).
- [21] — Cut times for simple random walk. *Electron. J. Probab.*, 1:13 (1996), 1–24 (electronic).
- [22] — The dimension of the frontier of planar Brownian motion. *Electron. Comm. Probab.*, 1:5 (1996), 29–47 (electronic).
- [23] — The frontier of a Brownian path is multifractal. Preprint, 1997.
- [24] — Strict concavity of the intersection exponent for Brownian motion in two and three dimensions. *Math. Phys. Electron. J.*, 4:5 (1998), 1–67 (electronic).
- [25] — Geometric and fractal properties of Brownian motion and random walk paths in two and three dimensions, in *Random Walks* (Budapest, 1998), pp. 219–258. Bolyai Soc. Math. Stud., 9. János Bolyai Math. Soc., Budapest, 1999.
- [26] LAWLER, G. F. & PUCKETTE, E. E., The intersection exponent for simple random walk. *Combin. Probab. Comput.*, 9 (2000), 441–464.

- [27] LAWLER, G. F., SCHRAMM, O. & WERNER, W., Values of Brownian intersection exponents, I: Half-plane exponents. *Acta Math.*, 187 (2001), 237–273.
- [28] — Values of Brownian intersection exponents, III: Two-sided exponents. To appear in *Ann. Inst. H. Poincaré Probab. Statist.* <http://arxiv.org/abs/math.PR/0005294>.
- [29] — Analyticity of intersection exponents for planar Brownian motion. To appear in *Acta Math.*, 188 (2002). <http://arxiv.org/abs/math.PR/0005295>.
- [30] — Sharp estimates for Brownian non-intersection probabilities. To appear in *In and Out of Equilibrium. Probability with a Physics Flavor*. Progr. Probab. Birkhäuser Boston, Boston, MA.
- [31] LAWLER, G. F. & WERNER, W., Intersection exponents for planar Brownian motion. *Ann. Probab.*, 27 (1999), 1601–1642.
- [32] — Universality for conformally invariant intersection exponents. *J. Eur. Math. Soc. (JEMS)*, 2 (2000), 291–328.
- [33] LÖWNER, K., Untersuchungen über schlichte konforme Abbildungen des Einheitskreises, I. *Math. Ann.*, 89 (1923), 103–121.
- [34] MADRAS, N. & SLADE, G., *The Self-Avoiding Walk*. Birkhäuser Boston, Boston, MA, 1993.
- [35] MANDELBROT, B. B., *The Fractal Geometry of Nature*. Freeman, San Francisco, CA, 1982.
- [36] NIENHUIS, B., Critical behavior of two-dimensional spin models and charge asymmetry in the Coulomb gas. *J. Statist. Phys.*, 34 (1984), 731–761.
- [37] POMMERENKE, CH., On the Loewner differential equation. *Michigan Math. J.*, 13 (1966), 435–443.
- [38] REVUZ, D. & YOR, M., *Continuous Martingales and Brownian Motion*. Grundlehren Math. Wiss., 293. Springer-Verlag, Berlin, 1991.
- [39] SCHRAMM, O., Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118 (2000), 221–288.
- [40] WERNER, W., Bounds for disconnection exponents. *Electron. Comm. Probab.*, 1:4 (1996), 19–28 (electronic).

GREGORY F. LAWLER
 Department of Mathematics
 Duke University
 Box 90320
 Durham, NC 27708-0320
 U.S.A.
 jose@math.duke.edu

ODED SCHRAMM
 Microsoft Research
 1, Microsoft Way
 Redmond, WA 98052
 U.S.A.
 schramm@microsoft.com

WENDELIN WERNER
 Département de Mathématiques
 Université Paris-Sud
 Bât. 425
 FR-91405 Orsay Cedex
 France
 wendelin.werner@math.u-psud.fr

Received May 18, 2000