

# Contracting automorphisms and $L^p$ -cohomology in degree one

Yves Cornulier and Romain Tessera

**Abstract.** We characterize those Lie groups, and algebraic groups over a local field of characteristic zero, whose first reduced  $L^p$ -cohomology is zero for all  $p > 1$ , extending a result of Pansu. As an application, we obtain a description of Gromov-hyperbolic groups among those groups. In particular we prove that any non-elementary Gromov-hyperbolic algebraic group over a non-Archimedean local field of zero characteristic is quasi-isometric to a 3-regular tree. We also extend the study to general semidirect products of a locally compact group by a cyclic group acting by contracting automorphisms.

## 1. Introduction

Let  $G$  be a locally compact group, endowed with a *left* Haar measure. Let  $\rho = \rho_G$  denote the *right* regular representation of  $G$  on the space  $\mathbf{R}^G$  of all real-valued functions on  $G$ , defined by  $(\rho(g)f)(h) = f(hg)$  for  $g, h \in G$  and  $f \in \mathbf{R}^G$ . Given  $p \in [1, \infty[$ , let  $D^p(G)$  denote the set of  $p$ -Dirichlet functions on  $G$ , namely measurable functions  $f$  on  $G$  such that  $b(g) = f - \rho(g)f$  belongs to  $L^p(G)$  for all  $G$ , and such that  $g \mapsto b(g)$  is continuous from  $G$  to  $L^p(G)$ . This space contains constant functions and  $L^p$  functions (indeed, the continuity of  $g \mapsto b(g)$  is clear for continuous compactly supported functions, which form a dense subspace of  $L^p(G)$ ). For every compact subset  $Q \subset G$ , we define a seminorm on  $D^p(G)$  by

$$\|f\|_{D^p, Q} = \sup_{g \in Q} \|f - \rho(g)f\|_p.$$

We equip  $D^p(G)$  with the topology induced from the seminorms  $\|\cdot\|_{D^p, Q}$  for all  $Q$ . Then it follows from [T2, Lemma 5.2] that  $D^p(G)/\mathbf{R}$  is isomorphic, as a topological vector space with a  $G$ -action, to the space of 1-cocycles  $Z^1(G, \rho_G^p)$ , where  $\rho_G^p$  denotes the right regular representation on  $L^p(G)$ .

---

The first author was supported by ANR quantiT JCO8\_318197.

Define the topological quotient vector space

$$H_p^1(G) = D^p(G)/(L^p(G) + \mathbf{R}) \simeq H^1(G, \rho_G^p)$$

and the Banach space

$$\overline{H_p^1}(G) = D^p(G)/\overline{(L^p(G) + \mathbf{R})} \simeq \overline{H^1}(G, \rho_G^p).$$

The first  $L^p$ -cohomology and the reduced one coincide if and only if the norms  $\|f\|_p + \|f\|_{D^p, Q}$  and  $\|f\|_{D^p, Q}$  are equivalent on  $L^p(G)$  for some compact subset  $Q$ . If  $G$  is non-compact, this happens if and only if  $G$  satisfies the Sobolev inequality

$$\|f\|_p \leq C\|f\|_{D^p, Q},$$

i.e. if and only if [T1, Proposition 11.9]  $G$  is either non-unimodular or non-amenable. This can be reformulated as

- if  $G$  is non-compact, amenable and unimodular, then the topological vector space  $H_p^1(G)$  is non-Hausdorff (and in particular is non-zero);
- otherwise,  $H_p^1(G) = \overline{H_p^1}(G)$ .

A definition of the first  $L^p$ -cohomology of a locally compact group in the context of metric measured spaces is given in [Pa2] (see also [T2, Section 3] and Appendix B); the equivalence between the two definitions is obtained in [T2, Section 5].

*Definition 1.1.* Throughout the paper, *Lie groups* refer to real Lie groups.

- The unit component of a locally compact group  $G$  is denoted by  $G_0$ .
- Let  $\mathbf{Z}$  act by automorphisms on a locally compact group  $H$ . A *vacuum subset* is a subset  $U \subset H$  such that for every compact subset  $M \subset H$ , we have  $kM \subset U$  for  $k$  large enough. We say that  $\mathbf{Z}$  *contracts*  $H$  if there exists a compact vacuum subset  $U$ . The action is *strictly contracting* if any neighborhood of 1 is a vacuum subset (in some papers this is simply referred to as *contracting*).
- We say that a Lie group  $G$  with finitely many connected components is of *Heintze type* if it is isomorphic to a semidirect product  $S \rtimes N$ , where  $N$  is a non-compact simply connected nilpotent Lie group, and  $S$  contains  $\mathbf{Z}$  as a cocompact subgroup which contracts  $N$ .
- We say that a locally compact group  $G$  is of *non-Archimedean Heintze type* if it is isomorphic to a semidirect product  $S \rtimes N$ , where  $N$  is a non-compact, totally disconnected, locally compact nilpotent group, and  $S$  contains  $\mathbf{Z}$  as a cocompact subgroup which contracts  $N$ .
- We say that a locally compact group  $G$  is of *rank-one type* (resp. *non-Archimedean rank-one type*) if for some (unique) compact normal subgroup  $W$  in  $G$ , some finite index subgroup of  $G/W$  is isomorphic to the quotient of a simple group of rank one over the reals (resp. over some non-Archimedean local field) by its center.

*Example 1.2.* Let  $G=(K \times \mathbf{R}) \ltimes N$  be a Lie group with finitely many connected components, with  $K$  compact, and assume that for every positive  $t \in \mathbf{R} \subset G$ , every eigenvalue  $\lambda$  of  $\text{Ad}(t)$  acting on the Lie algebra  $\mathfrak{n}$  of  $N$  satisfies  $|\lambda| < 1$ . Then  $N$  is nilpotent and simply connected,  $G$  is of Heintze type, and conversely every Lie group with finitely many connected components of Heintze type is of this form. Besides, Heintze’s main result in [He] is that a connected Lie group of dimension  $\geq 2$  has a left-invariant Riemannian metric of negative curvature if and only if it is a simply connected solvable Heintze Lie group.

Similarly, if  $G=(K \times \mathbf{K}^*) \ltimes N$  is an algebraic group over a non-Archimedean local field with  $K$  compact, and for every  $t \in \mathbf{K}^*$  with  $|t| > 1$ , all eigenvalues  $\lambda$  of  $\text{Ad}(t)$  acting on  $\mathfrak{n}$  satisfy  $|\lambda| < 1$ . Then  $G$  is of non-Archimedean Heintze type. When  $\mathbf{K}$  has characteristic zero, every connected linear algebraic  $\mathbf{K}$ -group of non-Archimedean Heintze type, is of this form.

Our main result is the following theorem. We include the statement (1) in order to give a complete picture, but only prove the other ones.

**Theorem 1.** *Consider  $p \in [1, \infty)$ . Let  $G$  be a connected Lie group, or a linear algebraic group over a non-Archimedean local field of characteristic zero.*

- (1) ([T2] and [T3]) *If  $G$  is amenable and unimodular, then  $\overline{H}_p^1(G) = 0$  for all  $p > 1$ .*
- (2) *Suppose that  $G$  is Heintze or rank-one type (Lie or non-Archimedean). Then  $\overline{H}_p^1(G) \neq 0$  for  $p$  large enough.*
- (3) *Otherwise,  $\overline{H}_p^1(G) = 0$  for all  $p \geq 1$ .*

Pansu [Pa3] obtains (2) and (3) when  $G$  is a connected solvable Lie group. His approach is based on a definition of  $L^p$ -cohomology for Riemannian manifolds, involving differentiation. This gives rise to some technical issues, related to the fact that the gradient of an  $L^p$ -function need not be  $L^p$ . The equivalence between the two definitions of  $L^p$ -cohomology is given in [T2, Chapters 4 and 5]. The discrete version, given here, allows a unified proof of (2) and (3) in the Lie and non-Archimedean case. Theorem 7 below provides a more general and more precise statement, which contains (2) as a particular case. We first focus on some corollaries on Theorem 1.

**Corollary 2.** *The following are equivalent:*

- (2.1) *for some  $p > 1$ , we have  $\overline{H}_p^1(G) \neq 0$ ;*
- (2.2)  *$G$  is Heintze or rank-one type (Lie or non-Archimedean).*

Lie groups admitting a left-invariant Riemannian metric with negative sectional curvature have been described algebraically in [He]. An application of our main

result is to provide a characterization of Gromov-hyperbolicity for Lie groups and algebraic groups over a local field of zero characteristic. This can be understood as a “large-scale” version of Heintze’s theorem [He]. Note that one advantage of focusing on the large-scale geometry of a Lie group is that the statements are independent of a choice of Riemannian metric and also make sense when we consider the word length with respect to a compact generating set. In general, a locally compact group is said to be *Gromov-hyperbolic* if it is compactly generated and, viewed as a metric space with the word metric to some compact generating set, is a Gromov-hyperbolic metric space.

By [CT, Theorem 1.2], a connected Lie group (or an algebraic group over a local field with characteristic zero) with exponential growth has a bi-Lipschitz embedded 3-regular tree. It then follows from [T2, Theorem 6] that if such a group is Gromov-hyperbolic, it has non-trivial first reduced  $L^p$ -cohomology for large enough  $p$ . Accordingly, we get the following corollary.

**Corollary 3.** *The following are equivalent:*

- (3.1) *the group  $G$  is non-elementary Gromov-hyperbolic;*
- (3.2) *the group  $G$  is of Heintze or rank-one type (Lie or non-Archimedean).*

If we distinguish the Lie and the non-Archimedean cases, we get the two following additional corollaries.

**Corollary 4.** *If  $G$  is a Lie group with finitely many connected components, we have the following equivalences:*

- (4.1)  *$G$  is [non-elementary] Gromov-hyperbolic;*
- (4.2)  *$G$  is quasi-isometric to a simply connected homogeneous manifold of negative curvature [of dimension at least two];*
- (4.3)  *$G$  acts properly transitively by isometries on a simply connected homogeneous manifold of negative curvature [of dimension at least two].*

The construction of a left-invariant metric on a simply connected Heintze Lie group is one of the principal results in [He]. We need slightly more to obtain the third statement of Corollary 4 for an arbitrary Heintze Lie group, but it turns out that what we need follows from Heintze’s construction, namely [He, Theorem 2]. This amounts to proving that if  $G$  is a simply connected Heintze group and  $K$  is a compact group of automorphisms of  $G$ , then  $G$  possesses a left-invariant Riemannian metric which has negative curvature and is  $K$ -invariant.

**Corollary 5.** *If  $G$  is an algebraic group over a non-Archimedean local field of characteristic 0, we have the following equivalences:*

- (5.1)  $G$  is [non-elementary] Gromov-hyperbolic;
- (5.2)  $G$  is quasi-isometric to a regular tree of finite degree [of degree at least three];
- (5.3)  $G$  acts properly, cocompactly (i.e. with finitely many orbits) by isometries on some regular tree of finite degree [of degree at least three].

Note that all regular trees of degree  $3 \leq d < \infty$  are quasi-isometric to each other. In order to get the third condition, we make use of the following general proposition.

**Proposition 6.** *Consider a contracting action  $\sigma$  of  $\mathbf{Z}$  on some non-compact, totally disconnected, locally compact group  $H$ . There exists a proper length function  $\ell$  on the semidirect product  $G = \mathbf{Z} \ltimes H$  such that the pseudo-metric space  $(G, d)$ , where  $d(g, g') = \ell(g^{-1}g')$ , is isometric to the vertex set of an  $r$ -regular tree for some  $r \geq 3$ .*

Turning back to Theorem 1(2), we have the following more general result.

**Theorem 7.** *Let  $H$  be a locally compact group whose unit component  $H_0$  is non-compact, with an action of a locally compact group  $S$ , which is contracting in restriction to some cocompact subgroup  $\mathbf{Z} = \langle \xi \rangle$  of  $S$ , and set  $G = S \ltimes H$ . Let  $\xi^{-1}$  multiply the Haar measure of  $N$  by  $\delta > 1$ , and let  $\lambda > 1$  be the smallest modulus of eigenvalues greater than 1 of  $\xi^{-1}$  on  $H_0$ . Set*

$$p_0 = p_0(G) = \frac{\log \delta}{\log \lambda} \geq 1.$$

Then for all  $p \geq 1$ ,

$$\overline{H}_p^1(G) \neq 0 \quad \text{if and only if} \quad p > p_0.$$

Theorem 7 is proved by Pansu ([Pa1] and [Pa3]) when  $H$  is a simply connected solvable Lie group. Theorem 7 applies to “mixed” groups such as  $\mathbf{Z} \ltimes_{(t^{-1}, \ell)} (\mathbf{R} \times \mathbf{Q}_\ell)$ , where  $t > 1$  and  $\ell$  prime, for which  $p_0 = \log(t\ell)/\log t$ . Unlike the case of Lie groups, this provides examples where  $p_0$  is (any number) in  $]1, 2[$ . Besides, the assumption in Theorem 7 of  $H_0$  being non-compact is no restriction, since otherwise when  $H_0$  is compact (but not  $H$ ), Proposition 6 applies and  $\overline{H}_p^1(G) \neq 0$  for all  $p \geq 1$  (see Proposition 3.4 for a direct proof).

In Section 2, we collect some results relating some properties of  $L^p$ -cocycles to the modular function of the group. These observations, crucial in the non-unimodular case, are largely adapted from [Pa3]. At the additional cost of some structural results on Heintze Lie groups, we then obtain Theorem 1(3).

In Section 3, we prove Theorem 7 and in particular deduce (2) of Theorem 1.

In Section 4, we complete the proof of the corollaries of Theorem 1. In particular, we prove the existence of a  $G$ -invariant metric of negative curvature on  $G/K$  for general Heintze Lie groups, and prove Proposition 6.

Finally, in Appendix A, we give a general structure theorem for locally compact groups with a contraction, and in Appendix B, we prove the quasi-isometric invariance of  $L^p$ -cohomology in degree one, a particular case of an unpublished result of Pansu [Pa2].

## 2. Vanishing of the first $L^p$ -cohomology and the modular function

In all this section,  $p \in [1, \infty)$ . The aim of this section is to prove Theorem 1(3).

### 2.1. Generalities

Recall that  $\rho$  denotes the *right* regular representation of  $G$  on  $L^p(G)$ . Let  $\Delta$  be the modular function on  $G$ . For every measurable function  $f$  on  $G$  and  $g \in G$ , we have  $\|\rho(g)f\| = \Delta(g)^{-1/p} \|f\|$ , where  $\|\cdot\|$  is always assumed to denote the  $L^p$ -norm. For every  $\xi \in G$ , define

$$W_\xi = \{h \in G \mid \{\xi^{-n}h\xi^n\}_{n \geq 0} \text{ is bounded}\}.$$

This is a subgroup of  $G$ .

**Lemma 2.1.** *Fix  $\xi \in G$  such that  $\Delta(\xi) > 1$  (such  $\xi$  exists if and only if  $G$  is non-unimodular). Suppose that  $u \in D^p(G)$ . Then there exists  $u_\infty \in D^p(G)$  such that  $u - u_\infty \in L^p(G)$  and  $\rho(\xi)u_\infty = u_\infty$ .*

*Proof.* We have

$$\|\rho(\xi^{n+1})u - \rho(\xi^n)u\| = \Delta(\xi)^{-n/p} \|\rho(\xi)u - u\|.$$

Therefore the sequence  $\{\rho(\xi^n)u - u\}_{n \geq 0}$  converges in  $L^p(G)$  to some function  $v$ ; we set  $u_\infty = v + u$ . In particular, the sequence  $\{\rho(\xi^n)u\}_{n \geq 0}$  converges almost surely to  $u_\infty$ . In particular,  $\rho(\xi)u_\infty = u_\infty$  almost everywhere. Moreover,  $u - u_\infty \in L^p(G)$ , so that  $u_\infty \in D^p(G)$  and defines the same class as  $u$  in  $H^1_p(G)$ .  $\square$

**Lemma 2.2.** *Let  $G$  be any locally compact group, and suppose that  $u \in D^p(G)$  satisfies  $\rho(H)u = u$  for some non-compact closed subgroup  $H$  of  $G$ . Let  $V$  be the centralizer of  $H$  in  $G$ . Then  $\rho(V)u = u$ .*

*Proof.* For any measurable function  $f$  on  $G$  and any measurable subset  $X$  of  $G$ , denote by  $\|f\|_X$  the  $L^p$ -norm of  $f\mathbf{1}_X$ .

Fix any compact subset  $X$  of  $G$ . As  $H$  is non-compact, its subset  $H_+ = H \cap \{g \mid \Delta(g) \leq 1\}$  is non-compact. As the right action of  $G$  on itself is proper, there exists a sequence  $\{h_i\}_{i \geq 0}$  in  $H_+$  such that the subsets  $Bh_i$  are pairwise disjoint. Fix  $g \in V$ . We have

$$\begin{aligned} \|u - \rho(g)u\|_X &= \|\rho(h_i)u - \rho(g)\rho(h_i)u\|_X && \text{(by the } H\text{-invariance of } u) \\ &= \|\rho(h_i)u - \rho(h_i)\rho(g)u\|_X && \text{(as } u \text{ and } h_i \text{ commute)} \\ &= \Delta(h_i)^{-1/p} \|u - \rho(g)u\|_{Xh_i}, \end{aligned}$$

which tends to 0, as  $i \rightarrow \infty$ , since  $u - \rho(g)u$  is  $L^p$ . Hence,  $\|u - \rho(g)u\|_X = 0$  for every compact subset  $X \subset G$ , so that  $\|u - \rho(g)u\| = 0$ , i.e.  $u = \rho(g)u$  almost everywhere.  $\square$

Fix  $\xi \in G$  and  $u \in D^p(G)$  satisfying  $\rho(\xi)u = u$ . For  $g \in G$  write  $b(g) = u - \rho(g)u$ .

**Lemma 2.3.** *If  $g \in G$  then*

$$\Delta(\xi)^{1/p} \|b(g)\| = \|b(\xi^{-1}g\xi)\|.$$

*Proof.* This follows from the formula

$$u - \rho(g)u = \rho(\xi)(u - \rho(\xi^{-1}g\xi)u). \quad \square$$

**Lemma 2.4.** *Suppose that  $\Delta(\xi) \geq 1$ . If  $u \in D^p(G)$  and  $\rho(\xi)u = u$ , then  $u$  is invariant by  $\rho(W_\xi)$ .*

*Proof.* Using that  $b$  is bounded on bounded subsets of  $G$ , this follows from Lemma 2.3.  $\square$

**Lemma 2.5.** *Let  $G$  be a locally compact group.*

(1) *If  $f \in L^p(G)$ ,  $p < \infty$ , then  $f$  cannot be left- or right-invariant under a non-compact closed subgroup  $H$  unless  $f = 0$ .*

(2) *If  $f \in D^p(G)$  is invariant under a non-compact closed normal subgroup  $N$ , then  $f$  is constant.*

*Proof.* (1) Otherwise, there exists  $\varepsilon > 0$  such that  $W = \{f \mid |f| > \varepsilon\}$  has non-zero finite measure  $m$ . Take a compact subset  $K$  such that the measure of  $K \cap W$  has measure  $> m/2$ . There exists  $g$  such that  $K$  and  $gK$  (resp.  $Kg$ ) are disjoint; we can suppose that  $\Delta(g) \geq 1$ . Then as  $gK$  (resp.  $Kg$ ) are contained in  $W$ , we get a contradiction.

(2) Then  $b(g)=f-\rho(g)f$  is (left)-invariant by  $N$ , and so is zero, i.e.  $f$  is constant.  $\square$

**2.2. A criterion for vanishing of  $L^p$ -cohomology**

**Proposition 2.6.** *Suppose that  $G$  contains*

- (i) *an element  $\xi$  satisfying  $\Delta(\xi)>1$ ;*
- (ii) *two non-compact closed subgroups  $Z$  and  $Y$ , with  $Z$  normal in  $G$ ;*

*and assume that  $Y \subset W_\xi$  and  $Y$  centralizes  $Z$ . Then  $H_p^1(G)=0$  for all  $p \geq 1$ .*

*Proof.* Take  $u \in D^p(G)$  and let us show that it is in the cohomology class of 0. By Lemma 2.1 we can suppose that  $\rho(\xi)u=u$ . By Lemma 2.4,  $u$  is then invariant by  $\rho(Y)$ . By Lemma 2.2,  $u$  is invariant by  $Z$ , and therefore by Lemma 2.5,  $u$  is constant.  $\square$

*Remark 2.7.* In general, it is not true that if  $H$  is a closed non-compact subgroup contained in  $\text{Ker}\Delta$  and if  $u \in D^p(G)$  is  $\rho(H)$ -invariant, then  $u$  is necessarily zero in  $H_p^1(G)=0$ . It is shown in [CTV, Proposition 4.3] that if  $H$  is any infinite discrete group and  $K$  is any non-trivial discrete group, then the free product  $G=H * K$  is a counterexample.

**2.3. Application to non-unimodular amenable Lie or  $p$ -adic groups**

We say that a connected Lie group, resp. connected algebraic group over a local field, is *triangulable* if it embeds as a closed subgroup of upper triangular real matrices (resp. upper triangular matrices over the ground field).

**Proposition 2.8.** *Let  $G$  be*

- (i) *either a non-unimodular triangulable Lie group; or*
- (ii) *a non-unimodular amenable connected linear algebraic group over a non-Archimedean local field of characteristic 0.*

*Then  $G$  satisfies the hypotheses of Proposition 2.6 if (and only if)  $G$  is not of Heintze type (Lie or non-Archimedean).*

*Proof.* The “only if” part is not the point of this section; it follows from Propositions 2.6 and 3.4. Let us focus on the “if” part, assuming that  $G$  is not of Heintze type.

If  $G$  is non-Archimedean, removing a maximal anisotropic torus in a Levi decomposition, we see that it has a cocompact Zariski-closed normal subgroup which



is triangulable (and necessarily non-Heintze as well), so we can also suppose in that case that  $G$  is triangulable (the group  $Z$  will be chosen characteristic so will remain normal in the whole group).

Let  $N$  be the nilpotent radical and  $Z$  be its center; the assumptions imply that  $Z$  is not compact. Write a Cartan decomposition  $G=AN$ ; this means that  $A$  is nilpotent and  $G=AN$  (in the non-Archimedean case we can have moreover  $G=A \times N$  and  $A$  abelian).

Consider the adjoint action of  $A$  on  $\mathfrak{g}$ . It defines a homomorphism  $A \rightarrow \text{GL}(\mathfrak{g})$ . Its Zariski closure is connected nilpotent, so decomposes as a direct product  $DU$  with  $D$  diagonalizable (over  $\mathbf{K}$ ) and  $U$  unipotent. If  $a \in A$ , we can thus decompose the corresponding automorphism of  $\mathfrak{g}$  as  $du$  and write (after choice of a suitable basis)  $d = \text{diag}(\lambda_1(a), \dots, \lambda_m(a))$ . Define the *weights* as the homomorphisms  $\omega_i: A \rightarrow \mathbf{R}$  defined by  $\omega_i(a) = \log |\lambda_i(a)| \in \mathbf{R}$ . Set  $B = \bigcap_{i=1}^m \text{Ker}(\omega_i)$ , so  $A/B$  is isomorphic to  $\mathbf{R}^k$  (Lie case) or  $\mathbf{Z}^k$  (non-Archimedean case) for some  $k$ . For every  $a \in A$ , we have  $\Delta(a) = \exp(\sum_{i=1}^m \omega_i(a))$ . Set

$$E_+ = \{a \in A \mid \Delta(a) > 1\} = \left\{ a \in A \mid \sum_{i=1}^m \omega_i(a) \geq 0 \right\},$$

which can be viewed as a “half-space” in  $A/B$ . Set

$$E_{\text{dil}} = \{a \in A \mid \omega_i(a) \geq 0 \text{ for all } i\}.$$

Clearly,  $E_{\text{dil}} \subset E_+$ , but in  $A/B$ ,  $E_{\text{dil}}/B$  identifies with  $\mathbf{R}_+^k$  or  $\mathbf{N}^k$ .

Therefore, if  $k \geq 2$ , then there exists an element  $\xi$  in  $E_+$  which is not in  $E_{\text{dil}}$ . So  $\omega_i(\xi) < 0$  for some  $i$ . This means that  $W_\xi$  is not compact and the hypotheses are fulfilled.

Let us now suppose that  $k=1$  ( $k=0$  is ruled out as it would force  $G$  to be unimodular (and even nilpotent)). Notice that this forces  $\mathfrak{n}$  to have codimension one. Pick  $\xi$  with  $\Delta(\xi) > 1$ . Since the action of  $\xi$  on  $\mathfrak{n}$  has at least one eigenvalue of modulus  $\leq 1$  (because  $G$  is not Heintze), we have that  $W_\xi$  non-compact.  $\square$

*Remark 2.9.* Let  $\mathbf{R}^2$  act on  $\mathbf{C}^2$  by  $(r, \theta) \cdot (z_1, z_2) = r(e^{i\theta} z_1, e^{\pi i \theta} z_2)$ . Then the semidirect product  $\mathbf{R}^2 \rtimes \mathbf{C}^2$  does not satisfy the hypotheses of Proposition 2.6, yet is non-Heintze. Note that this group is not triangulable.

**Lemma 2.10.** *Let  $G$  be a Lie group with  $\pi_0(G)$  finite, and assume that  $G$  is of Heintze type with  $G=S \times N$  as in Definition 1.1. Set  $K = \text{Ker}(\Delta_G) \cap S$ . Then  $K$  is a compact normal subgroup of  $S$  and has a direct factor in  $S$ , isomorphic to  $\mathbf{R}$ . Moreover, any element  $g$  in  $S \setminus K$  generates a cocompact subgroup contracting  $N$ .*

*Proof.* As  $S_0$  contains a cocompact cyclic subgroup, we have  $K \cap S_0$  compact, and hence  $K$  is compact. Since both  $S$  and  $N$  are unimodular,  $\Delta_G$  is non-trivial in restriction to  $S$ , and hence to  $S_0$ . Note that  $K$  is a maximal compact subgroup in  $S$  and in particular  $K \cap S_0 = K_0$ . As  $K_0$  is a connected compact Lie group, its inner automorphism group is open. In particular, the action of  $S_0$  by conjugation on  $K_0$  is by inner automorphisms and therefore its kernel  $S_1$  satisfies  $S_0 = K_0 S_1$ , in particular  $S_1$  is cocompact in  $S$ . Let  $L$  denote a one-parameter subgroup of  $S_1$ , not contained in  $\text{Ker}(\Delta_G)$ . In particular,  $L$  is closed and non-compact; by construction it centralizes  $K$ . As  $\Delta_G$  is surjective in restriction to  $L$ , we have  $S = KL$  and as  $[K, L] = 1$ ,  $K$  is compact and  $K \cap L = 1$ , this is a (topological) direct product.

Let  $\xi$  be an element of  $S$  contracting  $N$ . For some suitable norm, the adjoint action of  $K$  on  $\mathfrak{n}$  is isometric, and the action of the element  $\xi$  is strictly contracting. For some  $\gamma \in L$ , we have  $\gamma k = \xi$  for some  $k \in K$ . Hence  $\gamma$  contracts  $\mathfrak{n}$ . So it has all its eigenvalues on  $\mathfrak{n}$  of modulus  $< 1$ . Therefore all  $\{\gamma^t\}_{t > 0}$  is contracting, and therefore any element of the form  $\gamma^t k$  with  $t > 0$  and  $k \in K$  is contracting; each cyclic subgroup of  $S$  not contained in  $K$  contains such an element.  $\square$

**Lemma 2.11.** *Let  $G_1$  be a Lie group of Heintze type and  $G_2$  be a connected, cocompact normal subgroup. Then  $G_2$  is of Heintze type.*

*Proof.* Write  $G_1 = S \times N$  as in the definition of Heintze type and let  $\xi$  be a contracting element. Then both  $\mathfrak{s}$  and  $\mathfrak{n}$  are sums of characteristic subspaces for the adjoint action of  $\xi$ . Therefore,  $\mathfrak{g}_2 = \mathfrak{s}_2 \times \mathfrak{n}_2$  for some subspaces (necessarily ideals)  $\mathfrak{s}_2$  and  $\mathfrak{n}_2$  of  $\mathfrak{s}$  and  $\mathfrak{n}$ . Hence  $G_2 = S_2 \times N_2$  with  $S_2$  and  $N_2$  being normal subgroups of  $S$  and  $N$ , respectively. As  $G_2$  is cocompact and connected, necessarily  $N_2$  is cocompact and connected, and hence  $N_2 = N$ . In view of Lemma 2.10,  $S_2$  contains a cocompact cyclic subgroup contracting  $N$ , so  $G_2$  is of Heintze type.  $\square$

**Lemma 2.12.** *Let  $G$  be a Lie group with  $\pi_0(G)$  finite. Let  $G_1$  be a cocompact, normal, contractible subgroup of  $G$ . Then  $G = K \times G_1$  for some compact subgroup (actually, any maximal compact subgroup).*

*Proof.* Let  $K$  be a maximal compact subgroup of  $G$ . From the exact sequence associated with a fibration, we see that the natural map  $G \rightarrow G/G_1$  is a homotopy equivalence. Also, the inclusion  $K \subset G$  is a homotopy equivalence. So  $K \rightarrow G/G_1$  is a homomorphism between compact Lie groups which is a homotopy equivalence. Again using the long exact sequence associated with a fibration, we obtain that both the kernel and the cokernel of  $K \rightarrow G/G_1$  are contractible. As these are compact manifolds, they are necessarily points, that is, the map  $K \rightarrow G/G_1$  is an isomorphism. So  $G = K \times G_1$ .  $\square$

**Lemma 2.13.** *Let  $G$  be a Lie group with  $\pi_0(G)$  finite. Assume that  $G$  is of Heintze type with  $G=S \times N$  as in Definition 1.1. Then  $G/N$  is the largest quotient of  $G$  with polynomial growth (also known as the “exponential radical of  $G$ ”). In particular,  $N$  is a characteristic subgroup of  $G$ .*

*Proof.* Obviously  $G/N$  has polynomial growth. Conversely, let  $M$  be a normal, closed subgroup of  $G$  such that  $G/M$  has polynomial growth and let us show that  $N \subset M$ . By Guivarc’h [Gu], this means that in the adjoint representation of  $G$  on  $\mathfrak{g}/\mathfrak{m}$ , all eigenvalues have modulus one. This forces  $\mathfrak{n} \subset \mathfrak{m}$ , so  $N \subset M$ .  $\square$

**Lemma 2.14.** *Let  $G$  be a Lie group with  $\pi_0(G)$  finite. Assume that  $G$  has a normal, cocompact subgroup  $G_1$  which is of Heintze type. Then  $G$  is of Heintze type.*

*Proof.* Write  $G_1=S \times N$  as in Definition 1.1. By Lemma 2.13,  $N$  is characteristic in  $G_1$  and therefore normal in  $G$ .

By Lemma 2.10,  $S=K \times T$  with  $T \simeq \mathbf{R}$  and  $K$  compact. Let us prove that  $T \times N$  is normal in  $G$ . The subgroup  $T \times N$  is contained in the radical  $R_1$  of  $G_1$ . Now  $R_1$  is characteristic in  $G_1$ , so is normal in  $G$ , and thus is contained in the radical  $R$  of  $G$ . Hence  $T \times N \subset R$ . Therefore the image of  $T$  in  $G/N$ , which we still denote by  $T$ , is contained in the radical  $R/N$  of  $G/N$ . Since  $G_1$  is normal and cocompact, the restriction of  $\Delta_G$  to  $G_1$  is  $\Delta_{G_1}$ . So the map  $\Delta_G$  can be viewed as a homomorphism on  $G/N$ , which is non-trivial on  $T$ , and hence on  $R$ . Set  $M=R \cap \text{Ker}(\Delta)$ , so  $R/M \simeq \mathbf{R}$  and  $M/N$  is compact. Since  $R/N$  is connected and  $M/N$  is its maximal compact subgroup,  $M/N$  is connected. As a solvable, connected compact normal subgroup of  $R/N$ ,  $M/N$  is a central torus. Since it has codimension one in  $R/N$ , there is a one-dimensional factor, and hence a direct factor of  $M/N$  in  $R/N$ . Thus  $R/N$  is abelian. In particular the action of  $G$  on  $R/N$  by conjugation factors through the compact group  $G/R$ . This group preserves a direct product decomposition  $R/N=M/N \oplus V$ ; since  $G/R$  is compact, its action on the compact torus  $M/N$  is trivial. Moreover, its action leaves invariant the function  $\Delta_G$  and therefore acts trivially on  $V$ . Thus  $R/N$  is central in  $G/N$ , and hence  $T$  is normal in  $G/N$ . Therefore  $T \times N$  is normal in  $G$ .  $\square$

**Lemma 2.15.** *Let  $G$  be a connected amenable Lie group and  $W$  be a compact normal subgroup. If  $G/W$  is of Heintze type, then so is  $G$ .*

*Proof.* First observe that the result is clear when  $W$  is locally a direct factor. In general, we can suppose that  $W$  is either semisimple, finite, or a circle. When  $W$  is semisimple, as its outer automorphism group is discrete and  $W$  has finite center, its centralizer is locally a direct factor.

Suppose that  $W$  is finite. In particular,  $W$  centralizes  $G_0$ . Using Lemma 2.10, write  $G/W = ((T/W) \times (K/W)) \times (N/W)$ . As  $T/W$  and  $N/W$  are both simply connected, we have  $T = T_0 \times W$  and  $N = N_0 \times W$ . So  $G = TK \times N$  is of Heintze type.

Finally, suppose that  $W$  is a circle and that  $G/W$  is of Heintze type. By Lemmas 2.11 and 2.14, we can suppose that  $G/W = (S/W) \times (N/W)$  with  $S/W \simeq \mathbf{R}$ . As  $S/W$  is one-dimensional, we can lift it to a one-parameter subgroup  $T$  of  $G$ , so  $G = T \times N$  and  $N \supset W$ .

Let  $\xi \in T$  be an element contracting  $\mathfrak{n}$ . Consider the adjoint action of  $\xi$  on  $\mathfrak{n}$ , and denote by  $\mathfrak{c}$  be the sum of the characteristic subspaces for eigenvalues of modulus  $< 1$ . Since  $\mathfrak{c} = \{x \in \mathfrak{n} \mid \lim_{n \rightarrow \infty} \text{Ad}(\xi^n)x \rightarrow 0\}$ , we see that  $\mathfrak{c}$  is a Lie subalgebra. Note that since  $G/W$  is Heintze,  $\mathfrak{c}$  projects onto  $\mathfrak{n}/\mathfrak{w}$ , that is,  $\mathfrak{c} + \mathfrak{w} = \mathfrak{n}$ . Clearly the intersection is trivial (since  $\mathfrak{w}$  is central), so  $\mathfrak{c} \oplus \mathfrak{w} = \mathfrak{n}$ . Again using that  $\mathfrak{w}$  is central, we obtain that this is a decomposition as a direct product of Lie subalgebras. Let  $C$  be the Lie subgroup corresponding to  $\mathfrak{c}$ . The projection  $C \rightarrow N/W$  inducing the isomorphism  $\mathfrak{c} \rightarrow \mathfrak{n}/\mathfrak{w}$  is a covering; as  $G/W$  is simply connected, it is an isomorphism. In particular,  $C$  is closed, nilpotent and simply connected. This implies in particular that  $C \cap W = 1$ . So  $N$  is the topological direct product of  $C$  and  $W$ . As  $T$  normalizes both  $C$  and  $W$ , we obtain that  $W$  is a direct factor in  $G$ , hence clearly  $G$  is of Heintze type.  $\square$

**Theorem 2.16.** *Let  $G$  be a non-unimodular amenable Lie group with finitely many components, or a non-unimodular amenable linear algebraic group over a local field of characteristic zero. Assume that  $G$  is not of Heintze type (Lie or non-Archimedean). Then  $H_p^1(G) = 0$  for all  $p \geq 1$ .*

*Proof.* If  $G$  is triangulable or non-Archimedean, this is Proposition 2.8. The rest of the proof concerns connected Lie groups, and consists in reducing to the triangulable case, using the fact that the  $L^p$ -cohomology is a quasi-isometry invariant (see Appendix B).

Assume that  $G$  is an amenable Lie group with  $\pi_0(G)$  finite and  $G$  not of Heintze type. Then the radical  $G_1$  of  $G$  is cocompact and by Lemma 2.14 is not of Heintze type. Let  $W$  be the maximal normal compact subgroup in the connected solvable Lie group  $G_1$ . By Lemma 2.15,  $G_2 = G_1/W$  is not of Heintze type. Note that  $[G_2, G_2]$  is simply connected. By the *trigshadow* construction [C, Lemma 2.4], there exist connected Lie groups with normal cocompact inclusions  $G_2 \subset G_3 \supset G_4$  with  $G_4$  triangulable. By Lemmas 2.11 and 2.14,  $G_4$  is not of Heintze type and we can conclude by Proposition 2.8, using that cocompact inclusions are quasi-isometries.  $\square$

## 2.4. The case of non-amenable Lie or $p$ -adic groups

**Lemma 2.17.** *Let  $G$  be a connected non-amenable Lie group which is not of rank-one type. Let  $W$  be the maximal compact normal subgroup in  $G$ . Then  $G$  has a cocompact subgroup  $L$ , containing  $W$ , such that  $L/W$  is a simply connected solvable Lie group which is not of Heintze type.*

*Proof.* Let  $M$  be the connected amenable radical of  $G$ . Assume that  $M$  is not compact and  $G/M$  has finite center. Let  $R$  be the radical of  $G$  ( $R$  is cocompact in  $M$ ). Modding out if necessary, we can suppose that  $W=1$ . So the derived subgroup  $[R, R]$  of  $R$  is simply connected. Consider the action by conjugation of  $G/R$  on  $R/[R, R]$ . As  $G/R$  is semisimple, using the action on the universal covering, we obtain that  $R/[R, R]$  decomposes, under the action of  $G$ , as the direct sum of its maximal compact subgroup and some vector space  $V_0$ . Let  $V$  be the inverse image of  $V_0$  in  $R$ . So  $V$  is normal in  $G$ , simply connected, and cocompact in  $M$ . The group  $G/V$  is the direct product, up to some normal finite subgroup, of a compact group and a semisimple group with finite center. Therefore it has a simply connected solvable cocompact subgroup; let  $P$  be its inverse image in  $G$ , which is a simply connected solvable cocompact subgroup of  $G$ . Assume by contradiction that  $P$  is Heintze. We know that the derived subgroup of  $P/V$  has codimension at least one in  $P/V$ . As the Heintze assumption implies that  $[P, P]$  has codimension one, this forces  $[P, P]$  to contain  $V$ . As  $P$  is Heintze, it contains an element  $\xi$  contracting  $[P, P]$ . In particular,  $\xi$  contracts  $V$ . If  $g \in G$ , let the action of  $g$  by conjugation on  $V$  multiply the Haar measure by  $q(g)$ . So, using that  $V$  is non-compact,  $q(\xi) \neq 1$ . Because of the existence of this contraction,  $V \subset [G, G]$  and in particular  $V \subset \text{Ker}(q)$ , i.e.  $q$  factors through  $G/V$ . Now  $G/V$  has compact abelianization because it is compact-by-semisimple. So  $q$  is trivial, a contradiction.

Assume that either  $M$  is compact or  $G/M$  has infinite center. If  $M$  is compact, then by assumption  $G/M$  has rank at least two, so  $G/M$  either has infinite center or has rank at least two. Then by Lemmas 2.4 and 6.7 in [C], there exist cocompact inclusions

$$G/W \supset G_1 \subset G_2$$

with  $G_2$  being a solvable and simply connected Lie group. The assumptions imply that  $G/M$  contains  $\mathbf{Z}^2$  as a quasi-isometrically embedded subgroup, this copy can be lifted to a Levi factor of  $M$  in  $G/W$ , so  $G/W$  also contains a quasi-isometrically embedded subgroup isomorphic to  $\mathbf{Z}^2$ . Therefore  $G_2$  contains a quasi-isometrically embedded copy of  $\mathbf{Z}^2$  as well (actually, it follows from the construction in [C] that it can be realized as a subgroup, even if we can bypass it), so is not of Heintze type.  $\square$

If  $G$  is a linear algebraic group over a local field  $\mathbf{K}$ , the *rank* (or  $\mathbf{K}$ -rank) of  $G$  is the least  $k$  such that  $G$  contains a  $\mathbf{K}$ -split torus of rank  $k$ .

**Lemma 2.18.** *Let  $G$  be a connected linear algebraic group over a local field  $\mathbf{K}$  of characteristic zero. Suppose that  $G$  is not amenable, and is not reductive of rank one. Then  $G$  contains a cocompact subgroup of the form  $DU$  ( $D$  split torus and  $U$  unipotent) which is not of non-Archimedean Heintze type.*

*Proof.* This is a simplified analog of the proof of Lemma 2.17, so we only sketch it. The hypotheses mean that either  $G$  has rank at least two, or is not reductive.

Let  $N$  be the unipotent radical of  $G$ . The reductive group  $G/N$  has a cocompact subgroup of the form  $DU'$  with  $D$  being a split torus and  $U'$  unipotent. Let  $P=DU$  be the inverse image of this subgroup in  $G$ . If  $G$  has rank at least two, so does  $P$ , so that  $P$  is not of non-Archimedean Heintze type. Otherwise, since we assume that  $G$  is not amenable,  $G/N$  is non-abelian reductive of rank one. In particular,  $G/N$  has no non-trivial homomorphism to  $\mathbf{R}$ , as well as  $N$ . Therefore the action of  $G$  on  $N$  by conjugation preserves the Haar measure. As in the proof of Lemma 2.17, and as  $N$  is non-trivial, this prevents  $P$  from being a non-Archimedean Heintze group.  $\square$

**Proposition 2.19.** *Let  $G$  be either a connected Lie group, or a linear algebraic group over a local field of characteristic zero. Suppose that  $G$  is not amenable, and not of rank-one type. Then  $H_p^1(G)=0$  for all  $p \geq 1$ .*

*Proof.* The hypotheses exactly mean that Lemmas 2.17 and 2.18 do apply. So  $G$  is quasi-isometric to a non-Heintze simply connected solvable Lie group, or a non-Heintze triangulable group over a local field of characteristic zero. Again using the quasi-isometry invariance of the  $L^p$ -cohomology (Appendix B), the result then follows from Theorem 2.16.  $\square$

### 3. The $L^p$ -cohomology of Heintze groups

In this section, we prove (2) and (2') of Theorem 1.

We consider semidirect products  $D \ltimes N$  with the following convention:  $D$  acts on  $N$  on the right, and the group law is given by

$$(d_1, n_1)(d_2, n_2) = (d_1 d_2, (n_1 d_2) n_2).$$

Suppose that  $D$  is discrete. Then it is checked at once that if  $\lambda_0$  is a left Haar measure on  $N$ , and  $\delta$  is the counting measure on  $D$ , then  $\lambda = \delta \otimes \lambda_0$  is a left Haar measure on  $G = D \ltimes N$ .

Suppose now that  $u$  is a continuous function on  $G$  that is invariant for the right-regular action of  $D$  (i.e.  $u(gd)=u(g)$  for all  $g \in G$  and  $d \in D$ ). Then, if we set, for  $g \in N$ ,  $v(g)=u(1, g)$ , we have  $u(d, n)=v(nd^{-1})$  for all  $(d, n) \in D \times N$ .

**Lemma 3.1.** *If  $D$  is infinite and  $v$  is not constant, then  $u$  is not in  $\mathbf{R}+L^p(G)$ .*

*Proof.* For some  $\varepsilon > 0$ , there exist two disjoint measurable subsets of positive measure  $A_1$  and  $A_2$  of  $N$  such that for every  $(a_1, a_2) \in A_1 \times A_2$ ,  $v(a_1) \leq v(a_2) - \varepsilon$ . As  $D$  is infinite, the subset  $D_+$  of  $D$  consisting of elements such that the automorphism  $g \mapsto gd$  of  $N$  dilates the measure, is infinite. If  $d \in D_+$  and  $a \in A_i$ , then  $u(d, a_i d) = v(a_i)$ . Therefore, for every  $(b_1, b_2)$  with

$$b_i \in B_i = \bigcup_{d \in D_+} \{d\} \times (A_i d),$$

we have  $u(b_1) \leq u(b_2) - \varepsilon$ . As  $B_1$  and  $B_2$  both have infinite measure, this implies that  $u \notin \mathbf{R}+L^p(G)$ .  $\square$

**Lemma 3.2.** *Set, for  $g \in N$ ,  $b(g) = v - \rho(g)v$ . Then  $u \in D^p(G)$  if and only if  $v \in D^p(N)$  and*

$$(1) \quad \sum_{d \in D} \Delta(d) \|b(gd^{-1})\|_p^p$$

*is finite for all  $g \in N$  and tends to 0 when  $g \rightarrow 1$ .*

*Proof.* For  $g \in G$ , set  $B(g) = u - \rho(g)u$ . To check that  $B(g) \in L^p(G)$  for all  $g \in G$ , it is enough to check it for  $g$  ranging over a generating subset of  $G$ . As  $B(g) = 0$  for  $g \in D$ , it is enough to check it for  $g \in N$ .

For fixed  $d \in D$ , we have  $v_d(n) := u(d, n) = v(nd^{-1})$ , and the condition is that

$$\sum_{d \in D} \|v_d - \rho(g)v_d\|_p^p < \infty$$

for all  $g \in N$ . Now  $(v_d - \rho(g)v_d)(n) = b(gd^{-1})(nd^{-1})$ , so that (reminder:  $\int f(gh) dg = \Delta(h)^{-1} f(g) dg$ )

$$\|v_d - \rho(g)v_d\|_p^p = \Delta(d) \|b(gd^{-1})\|_p^p.$$

Thus, for  $g \in N$  we have

$$\|B(g)\|_p^p = \sum_{d \in D} \Delta(d) \|b(gd^{-1})\|_p^p. \quad \square$$

Let us now specialize to the case when  $D$  is cyclic and generated by an element  $\xi$  satisfying  $\delta = \Delta(\xi) > 1$ .

Let  $W$  be the largest compact normal subgroup in the unit component  $H_0$  of  $H$ , look at the eigenvalues of  $\xi$  on the Lie algebra on the non-trivial simply connected Lie group  $N_0/W$ , and define  $\lambda$  as the minimal modulus of its (complex) eigenvalues.

**Theorem 3.3.** *Consider a locally compact group  $G = S \rtimes H$ . Assume that some cyclic cocompact subgroup of  $S$  contracts  $H$ , and that  $H_0$  is non-compact. Define  $\delta$  and  $\lambda$  as above and  $G = \mathbf{Z} \rtimes H$ . Then  $H_p^1(G) = 0$  if and only if*

$$p \leq p_0(G) := \frac{\log \delta}{\log \lambda}.$$

*Proof.* Using the quasi-isometry invariance of  $L^p$ -cohomology (Appendix B), we can suppose that  $S = \mathbf{Z}$  contracts  $H$ .

By Corollary A.6,  $H/W(H_0)$  has a characteristic open subgroup  $L$  decomposing canonically as a direct product of a connected Lie group and a totally disconnected characteristic subgroup. Therefore the theorem is a combination of the following two propositions.  $\square$

**Proposition 3.4.** *Consider contracting actions  $\sigma$  and  $\sigma'$  of  $\mathbf{Z}$  on a connected Lie group  $N$  and on a totally disconnected locally compact group  $R$ , and  $H = N \times R$  with the diagonal (contracting) action of  $\mathbf{Z}$ . Define  $G = \mathbf{Z} \rtimes H$ . Pick  $p \geq 1$ . If  $N$  is non-compact, assume in addition that  $p > p_0(G)$ . Then  $H_p^1(G) \neq 0$ .*

*Proof.* On  $H = N \times R$ , we define  $v(n, r) = w(n)\beta(r)$ , where  $w$  is non-zero Lipschitz and compactly supported on  $N$  (Lipschitz referring to the intrinsic Riemannian distance on  $N$ ), and  $\beta$  being the indicator function of some clopen neighborhood of 1.

Since  $b$  is bounded, as  $d \rightarrow -\infty$ , the sum (1) converges in  $L^p$ -norm, uniformly on  $g$ .

Then for fixed  $g = (h, s) \in N \times R$ ,

$$\begin{aligned} b(g\xi^{-d})(n, r) &= v(n, r) - v((n, r)(g^{-1}\xi^{-d})) \\ &= w(n)\beta(r) - w((n(h^{-1}\xi^{-d}))\beta(r(s^{-1}\xi^{-d}))). \end{aligned}$$

When  $d \rightarrow \infty$ , we have that  $s^{-1}\xi^{-d} \rightarrow 1$ , so eventually, say for  $d \geq d_0 = d_0(g)$ , we obtain that  $\beta(r(g^{-1}\xi^{-d})) = \beta(r)$ . Thus for  $d \geq d_0$ ,

$$b(g\xi^{-d})(n, r) = \beta(r)[w(n) - w(n(h^{-1}\xi^{-d}))].$$



Pick  $\lambda' < \lambda$  such that  $\delta < (\lambda')^p$ . Then as  $\lambda' < \lambda$ , the Riemannian length  $h^{-1}\xi^{-d}$  is bounded above by  $(\lambda')^{-d}$  for  $d \gg 0$ . As  $w$  is Lipschitz, this implies that for some constant  $C$ ,

$$b(g\xi^{-d})(n, r) \leq C(\lambda')^{-d}.$$

As  $b(g')$  has support of bounded measure (independently of  $g'$ ), this implies that  $\|b(g\xi^{-d})\| \leq C'(\lambda')^{-d}$  for  $d \geq 0$ , for some suitable constant  $C'$ . So

$$\Delta(\xi)^d \|b(g\xi^{-d})\|^p \leq C'(\delta/(\lambda')^{-p})^d$$

and (1) holds.

We still need the continuity at 1. First note that  $d_0 = d_0(g)$ , as defined above, can be chosen bounded when  $g$  is bounded. We pick  $d_0$  working for some neighborhood  $V$  of 1. From the Lipschitz condition, we actually have

$$|b(g\xi^{-d})(n, r)| \leq C|h^{-1}\xi^{-d}|,$$

where  $|\cdot|$  denotes the Riemannian length in  $N_0$ , and  $|h^{-1}\xi^{-d}| \leq C'|h|(\lambda')^{-d}$  for all  $d \geq d_1$ ,  $d_1$  being independent of  $h \in V$ , which lies in the given neighborhood of 1. This implies that the sum

$$\sum_{d \geq d_1} \Delta(\xi)^d \|b(g\xi^{-d})\|^p$$

is continuous at 1, and therefore so is the entire sum, indexed by  $\mathbf{Z}$ .  $\square$

**Proposition 3.5.** *Consider contracting actions  $\sigma$  and  $\sigma'$  of  $\mathbf{Z}$  on a non-compact connected Lie group  $N$  and on a totally disconnected group  $R$ , and  $H = N \times R$  with the diagonal (contracting) action of  $\mathbf{Z}$ . Define  $G = \mathbf{Z} \ltimes H$  and  $p_0(G)$  as above. Then  $H_p^1(G) = 0$  for all  $p \leq p_0(G)$ .*

*Proof.* The first step is to modify the action of  $\mathbf{Z}$  on  $N$  so as to have a triangulable action with positive real eigenvalues.

Consider the tangent action of  $\sigma(\mathbf{Z})$  on  $\mathfrak{n}$ . Then we have  $\mathfrak{n} = \mathfrak{k} \oplus \mathfrak{m}$ , where  $\mathfrak{m}$ , resp.  $\mathfrak{k}$  is the sum of characteristic subspaces for eigenvalues of  $\sigma(\mathbf{Z})$  of modulus less than 1, resp. equal to 1. Moreover these are Lie subalgebras, and  $[\mathfrak{k}, \mathfrak{m}] \subset \mathfrak{m}$ , so  $\mathfrak{m}$  is an ideal. Let  $M$  and  $K$  be the corresponding Lie subgroups of  $G$ . Then  $M$  is strictly contracted by  $\sigma(\mathbf{Z})$ , and so is nilpotent and simply connected. Since  $M$  is contained in the nilpotent radical of  $N$ , which is simply connected,  $M$  is closed. Moreover  $\sigma$  contracts the quotient  $N/M$ , so that the tangent action has all its eigenvalues of modulus one. This implies (for instance) that  $\sigma(\mathbf{Z})$  preserves the measure on  $N/M$ , and therefore  $N/M$  is compact. So  $\mathbf{Z} \ltimes M \times R$  is cocompact in  $G$ , and hence quasi-isometric to it. In view of the quasi-isometry invariance of

the  $L^p$ -cohomology (see Appendix B), we can replace  $N$  by  $M$  if necessary, and we can suppose that  $N$  is nilpotent and simply connected.

Then the group  $\text{Aut}(N)=\text{Aut}(\mathfrak{n})$  is a linear algebraic group; and thus we can write  $\sigma(1)=g_+k=kg_+$  with  $k$  elliptic and  $g_+$  having all eigenvalues real and positive. If we define a new action  $\sigma_1$  of  $\mathbf{Z}$  on  $N$  by replacing  $\sigma(n)$  by  $g_+^n$ , define  $G_1=\mathbf{Z}\rtimes_{\sigma_1,\sigma'}(N\times R)$ . Let  $K$  denote the closure of the subgroup  $\langle k \rangle$  of  $\text{Aut}(N)$ . We can make  $\mathbf{Z}\times K$  act on  $N\times R$ , the action on  $N$  being the original action, the action of  $\mathbf{Z}$  on  $R$  being the original action, and the action of  $K$  on  $R$  being trivial. Then both  $G$  and  $G_1$  embed into  $(\mathbf{Z}\times K)\rtimes(N\times R)$  as cocompact subgroups. Therefore  $G_1$  is quasi-isometric to  $G$ . Again using the quasi-isometry invariance of  $L^p$ -cohomology, we can henceforth assume that  $N$  is simply connected and nilpotent, and  $\mathbf{Z}$  acts on it with all eigenvalues real and positive.

Assume that  $H_p^1(G)\neq 0$ . By Lemma 2.1, there exist  $u\in D^p(G)$  which is  $\rho(\xi)$ -invariant, and therefore  $u$  can be written as above. As the eigenvalues of  $\xi$  on  $N$  are real, there exists a one-parameter subgroup  $\{\gamma(t)\}_{t\in\mathbf{R}}$  of  $N$  on which  $\xi$  acts by multiplication by  $\lambda$ . Then (1) reads as

$$(2) \quad \sum_{n\in\mathbf{Z}} \Delta(\xi)^n \|b(\gamma(\lambda^{-n}))\|_p^p < \infty.$$

Set  $g_0=\gamma(1)$ . For  $n\geq 0$ , write  $\lambda^n=m_n+\varepsilon_n$ , where  $m_n=\lfloor \lambda^n \rfloor$ . Then

$$\|b(g_0)\| = \|b(\gamma(1))\| = \|b(\gamma((m_n+\varepsilon_n)\lambda^{-n}))\| \leq m_n \|b(u(\lambda^{-n}))\| + \|b(\varepsilon_n\lambda^{-n})\|.$$

In the last inequality, we use that  $\rho|_N$  is an isometric action. Set

$$e_n = \sup\{\|b(\gamma(t))\| \mid t \in [0, \lambda^{-n}]\}.$$

Then

$$\|b(g_0)\| \leq \lambda^n \|b(\gamma(\lambda^{-n}))\| + e_n.$$

We have  $e_n\rightarrow 0$ , as  $n\rightarrow\infty$ , since  $\lambda>1$  and the cocycle  $b$  is continuous. Assume that  $\|b(g_0)\|\neq 0$ . Then for  $n\gg 0$  we have

$$0 \leq \|b(g_0)\| - e_n \leq \lambda^n \|b(\gamma(\lambda^{-n}))\|,$$

so (2) implies that

$$\sum_{n\in\mathbf{N}} \left(\frac{\Delta(\xi)}{\lambda^p}\right)^n (\|b(g_0)\| - e_n)^p < \infty.$$

As  $\|b(g_0)\| - e_n \rightarrow \|b(g_0)\| > 0$ , this implies that  $|\Delta(\xi)/\lambda^p| < 1$ , that is,

$$p > \frac{\log \Delta(\xi)}{\log \lambda}.$$

Now assume by contradiction that  $b(g_0)=0$ , that is,  $\rho(g_0)u=u$ . Let  $Z$  be the center of  $N_0$ , which is non-compact. By Lemma 2.2,  $\rho(Z)u=u$ . Thus, by Lemma 2.5,  $u$  is constant on  $G$ , a contradiction.  $\square$

#### 4. On Gromov-hyperbolic groups

In the section, we prove the corollaries of Theorem 1. First, Corollary 2 is immediate from the theorem.

The following proposition is essentially proved, in the even more general context of metric groups by Gromov.

**Proposition 4.1.** *Let  $G$  be a non-elementary Gromov-hyperbolic locally compact group. Then  $G$  contains a quasi-isometrically embedded free subsemigroup on two generators.*

*Proof.* Gromov provides [Gr1, Section 8.2.D] two hyperbolic elements  $\gamma_1$  and  $\gamma_2$  with origins  $o_1$  and  $o_2$ , targets  $t_1$  and  $t_2$ , and  $\#\{o_i, t_1, t_2\}=3$  for  $i=1, 2$ . It then follows from the quasi-isometric ping-pong lemma [CT, Lemma 2.1] that suitable powers of  $\gamma_1$  and  $\gamma_2$  generate a free subsemigroup.  $\square$

Such a quasi-isometrically embedded subsemigroup provides a quasi-isometrically embedded tree in  $G$ , and by [T2, Theorem 6], this implies that the  $L^p$ -cohomology is non-zero for large  $p$ .

**Corollary 4.2.** *If  $G$  is a non-elementary Gromov-hyperbolic locally compact group, then  $H_p^1(G) \neq 0$  for  $p$  large enough.*

Note that for the groups considered in this paper, the results of [CT] are enough to provide a quasi-isometrically embedded subsemigroup without using Proposition 4.1, but we found it natural and useful to mention it. Anyway, (3.1) $\Rightarrow$ (2.1) and thus (3.1) $\Rightarrow$ (3.2) is proved. The converse (3.2) $\Rightarrow$ (3.1) is a particular case of Corollaries 4 and 5 (although the reader can prove it directly in a more straightforward way).

In Corollary 4, the implications (4.3) $\Rightarrow$ (4.2) $\Rightarrow$ (4.1) are straightforward. If (4.1) holds, by the already proved implication of Corollary 3, the unit component  $G_0$ , and therefore  $G$ , is a Lie group of Heintze type or rank-one type. So we have to prove the following proposition.

**Proposition 4.3.** *Let  $G$  be a Lie group with finitely many components, of Heintze type or rank-one type, and  $K$  be a maximal compact subgroup. Then the*

*connected manifold  $G/K$  has a  $G$ -invariant Riemannian metric of negative curvature.*

*Proof.* If  $G$  is of rank-one type,  $K$  acts irreducibly on the tangent space of the base-point of  $G/K$ , so the left-invariant Riemannian metric is unique up to scalar multiplication, and thus we get one of the simply connected irreducible symmetric spaces of rank one, which are negatively curved.

So assume that  $G$  is of Heintze type. Let  $H$  be a cocompact normal subgroup which is simply connected. Let  $K$  be a maximal compact subgroup of  $G$ . The adjoint action of  $K$  on  $\mathfrak{h}$  preserves the hyperplane  $[\mathfrak{h}, \mathfrak{h}]$ , and so preserves a complement line  $\mathfrak{a}$ , corresponding to some one-parameter  $A$  normalized by  $K$ . As the modular function  $\Delta$  is non-trivial on  $A$ , an element of  $A$  cannot be conjugate to its inverse, and therefore  $K$  centralizes  $A$ .

Let  $b$  be a  $K$ -invariant scalar product on  $\mathfrak{h}$ . Consider, for  $\lambda > 0$ , the linear automorphism of  $\mathfrak{h}$  mapping  $a+v$ ,  $a \in \mathfrak{a}$  and  $v \in [\mathfrak{h}, \mathfrak{h}]$ , to  $\lambda^{-1}a+v$ . Note that it commutes with the action of  $K$ , so  $(u_\lambda)_*b$  is also  $K$ -invariant. Then [He, Theorem 2] states that if  $\lambda$  is large enough, then the left-invariant metric on  $H$  obtained by translating  $(u_\lambda)_*b$  from the identity, is negatively curved. Moreover, this metric is  $K$ -invariant, and the group of isometries generated by left translations of  $H$  and conjugation by elements of  $K$  is naturally identified with  $G=K \ltimes H$ .  $\square$

Finally similarly, in Corollary 5, the implications (5.3) $\Rightarrow$ (5.2) $\Rightarrow$ (5.1) are straightforward. If (5.1) holds, by the already proved implication of Corollary 3,  $G$  is of non-Archimedean Heintze or rank-one type. In the latter case,  $G$  acts on the corresponding Bruhat–Tits tree, giving (5.3). Precisely, the action by conjugation of  $G$  on  $G_0$  provides a proper map  $G \rightarrow \text{Aut}(G_0)$ , and  $\text{Aut}(G_0)$  has a natural action on the Bruhat–Tits tree. For groups of non-Archimedean type, we have Proposition 4.6 below. We first need the following lemma.

**Lemma 4.4.** *Let  $H$  be a non-compact locally compact group with  $H_0$  compact, endowed with a contracting action of  $\mathbf{Z}$ . Then there exists a vacuum subset which is a compact open subgroup.*

*Proof.* Let  $U$  be a compact vacuum subset and set  $L = \bigcap_{k \geq 0} kU$ . Then it is easy to check that  $L$  is a compact subgroup. So, as  $H_0$  is compact,  $L$  is contained in a compact open subgroup  $V$  (by an easy argument; see if necessary [Pe, Theorem 2]), which is necessarily a vacuum subset.  $\square$

**Lemma 4.5.** *Let  $H$  be a non-compact locally compact group with  $H_0$  compact. Let  $S$  be a locally compact group with an action  $\pi$  on  $H$  by group automorphisms, and suppose that  $S$  possesses a cocompact copy of  $\mathbf{Z}$ , which contracts  $H$ . Then*

- (i)  *$S$  has a unique homomorphism  $p$  onto  $\mathbf{Z}$ , which is positive on the given copy of  $\mathbf{Z}$ , and  $W = \text{Ker}(p)$  is compact;*
- (ii) *there exists a compact open subgroup  $\Omega$  of  $H$  which is  $\pi(W)$ -invariant, and stable under  $\pi(p^{-1}(\mathbf{N}))$ , and for every compact subset  $K$  of  $H$  there exists  $k$  such that  $\pi(p^{-1}(\mathbf{N}_{\geq k}))K \subset \Omega$ .*

*Proof.* Let us first check that  $p$  is unique. Let  $p$  be a surjective homomorphism  $S \rightarrow \mathbf{Z}$ . As  $S$  contains a cocompact copy of  $\mathbf{Z}$ ,  $W = \text{Ker}(p)$  has to be compact and is thus determined as the unique maximal normal compact subgroup of  $S$ . This gives only two possibilities for  $p$ , and uniqueness follows from the positivity assumption.

If  $g \in S$ , let the automorphism  $\pi(s)$  of  $H$  multiply the Haar measure of  $H$  by  $q(s) \in \mathbf{R}_+^*$ . As  $H$  is non-compact, this is a non-trivial homomorphism  $S \rightarrow \mathbf{R}_+^*$ . By Lemma 4.4, we can choose a vacuum subset  $L$  which is a compact open subgroup of  $H$ . Then  $W_+ = \{g \in S \mid \pi(g)L \subset L\}$  contains 1 in its interior. Indeed, otherwise there exists a net  $g_i \rightarrow 1$  in  $S$  such that  $g_i \notin W_+$ . So there exists  $h_i \in L$  with  $\pi(g_i)h_i \notin L$ . As  $L$  is compact, we can suppose that  $\{h_i\}_{i \geq 0}$  has a limit  $h \in L$ . Since the action of  $G$  on  $H$  is continuous,  $\pi(g_i)h_i \notin L$  converges to  $h \in L$ , a contradiction. Similarly  $W_- = \{g \in S \mid \pi(g^{-1})L \subset L\}$  contains 1 in its interior, and therefore  $W_0 = W_+ \cap W_- = \{g \in S \mid \pi(g)L = L\}$  is an open subgroup of  $S$ . Clearly,  $W_0 \subset W = \text{Ker}(q)$ . Therefore  $W$  is open in  $S$ . The group  $S/W$  is discrete, is embedded into  $\mathbf{R}$  (by  $\log \circ q$ ), and contains a cocompact copy of  $\mathbf{Z}$ , so it is infinite cyclic as well. This yields the desired homomorphism  $p$ .

Let us now turn to the second assertion. It follows from the algebraic contractibility assumption that every compact subset of  $H$  generates a relatively compact subgroup of  $G$ . Therefore the subgroup  $L_1$ , defined as the closed subgroup generated by the compact subset  $\{\pi(g)h \mid g \in W \text{ and } h \in L\}$  of  $H$ , is compact. Now  $L_1$  is  $\pi(W)$ -invariant.

Fix  $t \in p^{-1}(\{1\})$ . Let  $t'$  be the positive generator of the given copy of  $\mathbf{Z}$ . As  $\pi(t)L_1$  is compact, there exists  $k \geq 0$  such that  $\pi((t')^k)\pi(t)L_1 \subset L_1$ . We can write  $(t')^k t = t^\ell w$  with  $\ell \geq 1$  and  $w \in W$ . So  $\pi(t^\ell)L_1 \subset L_1$ . Define  $\Omega = \bigcap_{i=0}^{\ell-1} \pi(t^i)L_1$ . Then  $\Omega$  is compact and open, and  $\pi(t)\Omega \subset \Omega$ . Moreover, as  $W$  is normalized by  $S$ ,  $\pi(t^i)L_1$  is  $\pi(W)$ -invariant for all  $i$ , and so  $\Omega$  is  $\pi(W)$ -invariant as well. Now  $t$  and  $W$  generate the semigroup  $p^{-1}(\mathbf{N})$ , and thus  $\pi(p^{-1}(\mathbf{N}))\Omega \subset \Omega$ .

As  $\bigcup_{i=0}^{\ell-1} \pi(t^i)L_1$  is a compact subset of  $H$ , there exists  $\varkappa$  such that

$$\pi(t^\varkappa) \left( \bigcup_{i=1}^{\ell} \pi(t^i)L_1 \right) \subset L \subset L_1.$$

So

$$\pi(t^{\varkappa+i})L_1 \subset L_1 \quad \text{for all } i = 1, \dots, \ell,$$

that is,

$$\pi(t^\varkappa)L_1 \subset \pi(t^{-i})L_1 \quad \text{for all } i = 1, \dots, \ell,$$

and thus

$$\pi(t^{\varkappa+\ell})L_1 \subset \bigcap_{j=0}^{\ell-1} \pi(t^j)L_1 = \Omega.$$

Now if  $K$  is a compact subset of  $H$ , there exists  $k \geq 0$  such that  $\pi(t^k)K \subset L \subset L_1$ . Therefore  $\pi(t^{k+\varkappa+\ell})K \subset \Omega$ , and hence  $\pi(g)K \subset \Omega$  whenever  $p(g) \geq k + \varkappa + \ell$ .  $\square$

**Proposition 4.6.** *Keep the assumptions as in Lemma 4.5. Then there exists a proper length function  $\ell$  on the semidirect product  $G = S \rtimes H$  such that the pseudo-metric space  $(G, d)$ , where  $d(g, g') = \ell(g^{-1}g')$ , is isometric to the vertex set of an  $r$ -regular tree for some  $r \geq 3$ .*

*Proof.* Let  $W$ ,  $p$  and  $H_0$  be as given by Lemma 4.5, and fix  $t \in p^{-1}(\{1\})$ . Replacing if necessary  $S$  by  $\mathbf{Z}$  (but we still write it multiplicatively) and  $H$  by  $W \rtimes H$  (and  $H_0$  by  $W \rtimes H_0$ ), we can suppose that  $G = \mathbf{Z} \rtimes H$ , so now  $p$  is just the identity.

Set  $M = H_0\{t, t^{-1}\}H_0$ . Consider the Cayley graph of  $G$  with respect to  $M$ . As  $M$  is invariant under conjugation by  $H_0$ , the right action of  $H_0$  on  $G$  preserves this Cayley graph structure. So we get a graph structure on the quotient  $G/H_0$ ; moreover as the original right action of  $H_0$  commutes with the left action of  $G$ , we get a left action of  $G$  on the graph  $G/H_0$ . As  $M$  generates  $G$ , this graph is connected. To check that it is a tree, it is enough to check that  $-p$  behaves like a Busemann function, i.e. for any vertex  $v$  with  $p(v) = n$  there is only one vertex  $v'$  adjacent to  $v$  with  $p(v') = n + 1$ . (Indeed, if we have an injective loop, it contains a vertex  $v$  with  $p(v)$  minimal, and the two adjacent vertices in the loop have  $p(v') = p(v) + 1$ .) By homogeneousness, it is enough to check this when  $v = (1, 1)$  is the identity. Let  $(t, u_1)$  and  $(t, u_2)$  be two neighbors of  $v$ , viewed in  $G$ . This means that  $(t, u_i)$  belong to  $H_0 t H_0$  for  $i = 1, 2$ . So there exist  $v_1, v'_1 \in H_0$  such that

$$(t, u_1) = (1, v_1)(t, 1)(1, v_2) = (1, v_1)(t, v_2) = (t, \pi(t)v_1 \cdot v_2).$$

As  $\pi(t)v_1 \in H_0$ , we obtain that  $u_1$  belongs to  $H_0$ . Similarly  $u_2 \in H_0$ . Now

$$(t, u_1)^{-1}(t, u_2) = (1, u_1^{-1}u_2) \in H_0,$$

so  $(t, u_1)$  and  $(t, u_2)$  are identified in  $G/H_0$ .  $\square$

### Appendix A. Direct decomposition for a contraction

This appendix is needed for the proof of Theorem 3.3 (and thus Theorem 7). It can also be of independent interest for the general study of contractions, as it generalizes [S, Proposition 4.2], which applies to strict contractions.

We refer to Definition 1.1 for the definition of a contraction and a vacuum subset. If a locally compact group  $G$  has a maximal normal compact subgroup, such a subgroup is unique and denoted by  $W(G)$ . Notably, such a subgroup exists when  $\pi_0(G) = G/G_0$  is compact [MZ, p. 175] and in particular when  $G$  is connected. We say that a locally compact group  $G$  is *elliptic* if every compact subset is contained in a compact subgroup. We have the following easy lemma.

**Lemma A.1.** *If a locally compact group  $G$  has a contraction, then  $G/G_0$  is elliptic.*

*Proof.* As  $\alpha$  induces a contraction of  $G/G_0$ , we can suppose that  $G$  is totally disconnected and we have to prove that  $G$  is elliptic. Let  $U$  be a compact vacuum subset. Define the limit set  $L$  as  $\bigcap_{n \geq 0} \alpha^n(U)$ . This is a compact subgroup. Note that  $\alpha$  (strictly) contracts  $G$  modulo  $L$ , in the sense of [HaS]. Since  $G$  is totally disconnected, there exists a compact open subgroup  $V$  containing  $L$ . If  $K$  is any compact subset of  $G$ , then  $\alpha^n(K)$  is contained in  $V$  for some  $n$ , and therefore  $K$  is contained in the compact subgroup  $\alpha^{-n}(V)$ .  $\square$

**Lemma A.2.** *Let  $G$  be a locally compact group with  $\pi_0(G)$  compact and  $W(G_0) = 1$ . Then the subgroup generated by  $G_0$  and  $W(G)$  is naturally isomorphic to the direct product  $G_0 \times W(G)$  and is open of finite index.*

*Proof.* As  $W(G_0) = 1$ , we have  $W(G) \cap G_0 = \{1\}$ . Hence  $W(G)$  and  $G_0$  centralize each other. As  $W(G)$  is compact, this is enough to ensure that the natural homomorphism  $G_0 \times W(G) \rightarrow G_0 W(G)$  is a topological isomorphism onto a closed subgroup.

Since  $\pi_0(G)$  is compact,  $G/W(G)$  is a Lie group with finitely many components. Now  $G/G_0 W(G)$  is both a quotient of  $G/W(G)$  and of  $G/G_0$ , which is totally discontinuous, and therefore  $G/G_0 W(G)$  is finite.  $\square$

Under the same assumption, we need to characterize  $W(G)$  without referring to any normality assumption (since in the sequel  $G$  will vary among open subgroups of a given group).

**Lemma A.3.** *Let  $G$  be a Lie group with  $\pi_0(G)$  finite and  $W(G)=1$ . Then  $G$  has no non-trivial compact subgroup centralizing  $G_0$ .*

*Proof.* Let  $K$  be a compact subgroup centralizing  $G_0$ ; let us assume that  $K$  is maximal for this property. Then  $K \cap G_0$  is compact and central in  $G_0$  and thus trivial. In particular,  $K$  is finite. Let  $S$  be a maximal compact subgroup of  $G$  containing  $K$ , and let  $N$  be the centralizer of  $G_0$  in  $G$ . Since  $G=SG_0$  [Mo, Theorem 3.1], any conjugate of  $K$  is contained in  $S$ . Moreover since  $K$  is contained in  $N$  which is normal, any conjugate of  $K$  is contained in  $S \cap N$ . Now  $S \cap N$  is a compact subgroup centralizing  $G_0$ , so by maximality of  $K$ , we have  $K=S \cap N$  and therefore  $K$  contains all its conjugates, and thus normal in  $G$ . As  $W(G)=1$ , this implies  $K=1$ .  $\square$

**Lemma A.4.** *Let  $G$  be a locally compact group with  $\pi_0(G)$  compact and  $W(G_0)=1$ . Then any compact subgroup of  $G$  centralizing  $G_0$  is contained in  $W(G)$ .*

*Proof.* Let  $K$  be a compact subgroup with  $[K, G_0]=1$ . By Lemma A.3, the image of  $K$  in  $G/W(G)$  is trivial.  $\square$

Therefore, when  $\pi_0(G)$  is compact and  $W(G_0)=1$  holds,  $W(G)$  appears as the maximal compact subgroup centralizing  $G_0$ .

**Theorem A.5.** *Let  $G$  be a locally compact group with  $\pi_0(G)$  elliptic and  $W(G_0)=1$ . Then  $G$  has a unique maximal subgroup  $W'(G)$  among those closed elliptic subgroups centralizing  $G_0$ . Moreover, the natural map  $G_0 \times W'(G) \rightarrow G_0 W'(G)$  is a topological isomorphism onto an open subgroup of  $G$ .*

*Proof.* Write  $G$  as the union of a net of open subgroups  $G_i$  with  $\pi_0(G_i)$  compact. If  $G_i \subset G_j$ , then

$$(A.1) \quad W(G_j) \cap G_i = W(G_i).$$

Indeed,  $W(G_j) \cap G_i \subset W(G_i)$  by definition of  $W$ , and the reverse inclusion follows from Lemma A.4 applied inside  $G_j$ . Set  $W'(G) = \bigcup_i W(G_i)$ . From (A.1) it follows that  $W'(G) \cap G_i = W(G_i)$  for all  $i$ , and in particular  $W'(G)$  is closed. Clearly it is elliptic and centralizes  $G_0$ . Conversely if  $H$  is closed, elliptic and centralizes  $G_0$ , then  $H \cap G_i$  is compact for all  $i$  and therefore by Lemma A.4 is contained in  $W(G_i)$ , and hence in  $W'(G)$ , so that  $H$  is contained in  $W'(G)$ .



To check that the natural map  $G_0 \times W'(G) \rightarrow G$  is a topological isomorphism onto its image, it is enough to check that it is a proper injective homomorphism. It is injective since  $G_0 \cap W'(G) = 1$ . To check properness, let  $\{(g_n, w_n)\}_{n \geq 0}$  be a sequence with  $g_n \in G_0$ ,  $w_n \in W'(G)$  and  $g_n w_n \rightarrow 1$ . Then since  $G_i$  is open, eventually  $g_n w_n \in G_i$ , so  $w_n \in W(G_i)$ , and therefore properness is reduced to properness in restriction to  $G_0 \times W(G_i)$ , which was established in Lemma A.2.

Finally  $G_0 W'(G)$  contains the open subgroup  $G_0 W(G_i)$  of the open subgroup  $G_i$ , so is open itself in  $G$ .  $\square$

**Corollary A.6.** *Let  $G$  be a locally compact group with  $W(G_0) = 1$ , and assume that  $G$  has a contraction. Then  $G$  has a characteristic subgroup  $W'(G)$  satisfying the properties stated in Theorem A.5, and  $G_0 W'(G)$  has finite index.*

*Proof.* By Lemma A.1,  $G/G_0$  is elliptic and Theorem A.5 applies. It is obvious from its statement that the subgroup  $W'(G)$  is characteristic, and in particular the open subgroup  $G_0 W'(G)$  is characteristic as well, and thus is stable under the given contraction. Therefore the discrete group  $G/G_0 W'(G)$  also has a contraction, and hence must be finite.  $\square$

This contains [S, Proposition 4.2] as a particular case.

**Corollary A.7.** *Let  $G$  be a locally compact group and assume that  $G$  has a strict contraction. Then  $G$  decomposes (canonically) as a product of characteristic subgroups  $G = G_0 \times W'(G)$ .*

*Proof.* The strict contraction  $\alpha$  strictly contracts the compact normal subgroup  $W(G_0)$  and therefore  $W(G_0) = 1$ , so Corollary A.6 applies. Moreover  $\alpha$  induces a strict contraction of the finite group  $G/G_0 W'(G)$ , so this is the trivial group, that is,  $G = G_0 W'(G)$ .  $\square$

## Appendix B. Quasi-isometry invariance

The invariance under quasi-isometries of the  $L^p$ -cohomology was obtained in all degrees in [Pa2]. As that preprint from 1995 is not yet published, and the case of degree one, used in the proof of the vanishing results of the  $L^p$ -cohomology in Theorem 1, is considerably simpler than in higher degree, we include a full proof.

The following coarse notion of (first)  $L^p$ -cohomology in degree one is essentially due to [Pa2] (see also the chapter about  $L^p$ -cohomology in [Gr2, Chapter 8], and [T2] for the case of degree 1). Here, we use the notation of [T2], but we extend

the definition to spaces which are not necessarily 1-geodesic. This will allow us to apply our result to non-compactly generated locally compact groups. The fact that this definition is equivalent to the one we gave in the introduction follows from the proof of [T2, Theorem 5.1].

Let  $X=(X, d, \mu)$  be a metric measure space, and let  $p \geq 1$ . For every  $s > 0$ , we write  $\Delta_s = \{(x, y) \in X^2 \mid d(x, y) \leq s\}$ .

First, let us introduce the  $p$ -Dirichlet space  $D_p(X)$ .

– The space  $D_p(X)$  is the set of measurable functions  $f$  on  $X$  such that

$$\int_{\Delta_s} |f(x) - f(y)|^p d\mu(x) d\mu(y) < \infty$$

for every  $s > 0$ .

– Let us equip  $D_p(X)$  with the topology induced by the following seminorms (for all  $s > 0$ ):

$$\|f\|_{D_{p,s}} = \left( \int_{\Delta_s} |f(x) - f(y)|^p d\mu(x) d\mu(y) \right)^{1/p}.$$

*Definition B.1.* The first  $L^p$ -cohomology of  $X$  is the space

$$H_p^1(X) = D_p(X) / (L^p(X) + \mathbf{R}),$$

and the first reduced  $L^p$ -cohomology of  $X$  is the space

$$\overline{H}_p^1(X) = D_p(X) / \overline{(L^p(X) + \mathbf{R})}^{D_p(X)}.$$

Let  $X=(X, d, \mu)$  be a metric measure space satisfying the following “bounded geometry” condition: for all  $x \in X$  and  $r > 0$ ,

$$v(r) \leq \mu(B_X(x, r)) \leq V(r),$$

where  $v$  and  $V$  are increasing positive functions on  $(0, \infty)$  and  $B_X(x, r) = \{y \in X \mid d(x, y) < r\}$ . It is not difficult to check that this condition is automatically satisfied if the group of measure-preserving isometries of  $X$  acts cocompactly. In particular, a locally compact group equipped with a left-invariant, proper metric has bounded geometry (properness is assumed here only for the balls to have finite volume).

We will use the notation  $f \preceq_T g$  when  $f \leq Cg$  for some constant  $C$  depending “only” on  $T$ .

**Theorem B.2.** *Suppose that two metric measure spaces  $X$  and  $Y$  with bounded geometry are coarse equivalent. Then there exists a topological isomorphism between their first  $L^p$ -cohomology.*

*Proof.* First, note that if the spaces are discrete (equipped with the counting measure) and if  $\varphi$  is a *bijective* coarse equivalence from  $X$  to  $Y$ , then  $\varphi$  induces a topological isomorphism on the first  $L^p$ -cohomology. Therefore, owing to the following lemma, we can suppose that  $Y$  is a *discretization* of  $X$ , i.e. a sub-metric space of  $X$  such that  $d(y, y') \geq 1$  for all  $y, y' \in Y, y \neq y'$ , and such that the inclusion  $Y \subset X$  is a coarse equivalence. This is equivalent to the existence of some  $R > 0$  such that  $X \subset \bigcup_{y \in Y} B_X(y, R)$ . Moreover, since  $X$  has bounded geometry, at most  $N = N(R)$  of these balls intersect.

**Lemma B.3.** *Let  $\varphi: X \rightarrow Y$  be a coarse equivalence. There exists a discretization  $X_d$  of  $X$ , such that the restriction of  $\varphi$  to  $X_d$  is one-to-one, and such that  $\varphi(X_d)$  is a discretization of  $Y$ .*

*Proof.* By definition of a coarse equivalence, there exists  $R > 0$  such that points at distance at least  $R$  in  $X$  are mapped to points at distance at least 1 in  $Y$ . Hence the lemma follows by letting  $X_d$  be a maximal  $R$ -separated net in  $X$ .  $\square$

The two main ingredients of the proof, that we will use thoroughly without mentioning them, are Hölder’s inequality and the fact that  $X$  and  $Y$  have bounded geometry.

For every  $T > 0$ , let us define an operator  $P_T$  on  $L^p(X)$  as follows:

$$P_T f(x_0) = \mathbf{E}_{B_X(x_0, T)} f,$$

where  $\mathbf{E}_A f := (1/\mu(A)) \int_A f(x) d\mu(x)$ . Also, we set  $\nabla f(x, x') := f(x) - f(x')$ .

**Lemma B.4.** *For every  $T \geq 0$ , the linear map  $\psi_T(f)(y) = P_T f(y)$ , from  $D_p(X)$  to  $D_p(Y)$ , induces a continuous map on the  $L^p$ -cohomology.*

*Proof.* Note that  $\psi_T$  obviously preserves constant functions and it is easy to see that it is bounded on  $L^p$ . Let  $s \geq T$ . We will sketch the proof that  $\psi_T$  is continuous from  $D_p(X)$  to  $D_p(Y)$  (the details are straightforward and therefore left to the reader),

$$\begin{aligned} \sum_{d(z, z') \leq s} |P_T f(z) - P_T f(z')|^p &\leq_s \sum_{d(z, z') \leq s} (\mathbf{E}_{B_X(z, 2s) \times B_X(z, 2s)} |\nabla f|)^p \\ &\leq \sum_{d(z, z') \leq s} \mathbf{E}_{B_X(z, 2s) \times B_X(z, 2s)} (|\nabla f|^p) \\ &\leq_s \int_{d(x, x') \leq 5s} |f(x) - f(x')|^p d\mu(x) d\mu(x'). \quad \square \end{aligned}$$

Now, let us choose  $T \geq R$ , so that  $X \subset \bigcup_{y \in Y} B_X(y, T)$ . Let

$$S_T(x) = \sum_{z \in Y} \mathbf{1}_{B_X(z, T)} \quad \text{for all } x \in X,$$

which satisfies  $1 \leq S_T \leq N(T) < \infty$ , and for all  $z \in Y$ ,

$$S_T^z = \mathbf{1}_{B_X(z, T)} / S_T.$$

Note that  $\{S_T^z\}_{z \in Y}$  forms a partition of unity for  $X$ .

**Lemma B.5.** *For every  $T \geq R$ , the linear map  $\phi_T(f) = \sum_{z \in Y} f(z) S_T^z$ , from  $D_p(Y)$  to  $D_p(X)$ , induces a continuous map on the  $L^p$ -cohomology.*

*Proof.* It is clear that  $\phi_T$  preserves constant functions, and the proof that it is bounded on  $L^p$  is easy and left to the reader. Let us sketch the proof that  $\phi_T$  is continuous on  $D_p$ . Let  $(x, x') \in \Delta_s(X)$  for some  $s \geq T$ . Then

$$\begin{aligned} |\phi_T(f)(x) - \phi_T(f)(x')|^p &= \left| \sum_{z, z' \in Y} (f(z) - f(z')) S_T^z(x) S_T^{z'}(x') \right|^p \\ &= \left| \sum_{z, z' \in B_X(x, 3s)} (f(z) - f(z')) S_T^z(x) S_T^{z'}(x') \right|^p \\ &\preceq_s \sum_{z, z' \in B_X(x, 3s)} |f(z) - f(z')|^p. \end{aligned}$$

Accordingly

$$\int_{d(x, x') \leq s} |\phi_T(f)(x) - \phi_T(f)(x')|^p \preceq_s \sum_{d(z, z') \leq 6s} |f(z) - f(z')|^p. \quad \square$$

**Lemma B.6.** *For every  $T \geq R$ , the maps  $\phi_T \circ \psi_T$  and  $\psi_T \circ \phi_T$  induce the identity on the  $L^p$ -cohomology of  $X$  and  $Y$ , respectively.*

*Proof.* For every  $x_0 \in X$ , we have

$$\begin{aligned} |f(x_0) - \phi_T \circ \psi_T(f)(x_0)|^p &= \left| \sum_{z \in Y} [f(x_0) - P_T f(z)] S_T^z(x_0) \right|^p \\ &= \left| \sum_{z \in Y} \mathbf{E}_{B_X(z, T)} [f(x_0) - f] S_T^z(x_0) \right|^p \end{aligned}$$

$$\begin{aligned}
 &= \left| \sum_{d(z,x_0) \leq T} \mathbf{E}_{B_X(z,T)} [f(x_0) - f] S_T^z(x_0) \right|^p \\
 &\preceq_T \sum_{d(z,x_0) \leq T} |\mathbf{E}_{B_X(z,T)} [f(x_0) - f]|^p \\
 &\leq \sum_{d(z,x_0) \leq T} \mathbf{E}_{B_X(z,T)} |f(x_0) - f|^p \\
 &\preceq_T \mathbf{E}_{B_X(x_0,2T)} |f(x_0) - f|^p.
 \end{aligned}$$

It follows that

$$\|f - \phi_T \circ \psi_T(f)\|_p \preceq_T \|f\|_{D_{p,2T}},$$

and therefore  $\phi_T \circ \psi_T$  induces the identity on  $H_p^1(X)$ .

For every  $z_0 \in Y$ ,

$$\begin{aligned}
 |f(z_0) - \psi_T \circ \phi_T(f)(z_0)|^p &= |P_T[f(z_0) - \phi_T(f)](z_0)|^p \\
 &= \left| P_T \left[ \sum_{z \in Y} (f(z_0) - f(z)) S_T^z \right] (z_0) \right|^p \\
 &= \left| \sum_{z \in Y} (f(z_0) - f(z)) \mathbf{E}_{B_X(z_0,T)} S_T^z \right|^p \\
 &\preceq_T \sum_{d(z,z_0) \leq 2T} |f(z_0) - f(z)|^p.
 \end{aligned}$$

We used the fact that  $\mathbf{E}_{B_X(z_0,T)} S_T^z \preceq_T \mathbf{1}_{B(z_0,2R)}(z)$ . We deduce from the above that

$$\|f - \psi_T \circ \phi_T(f)\|_p \preceq_T \|f\|_{D_{p,2T}},$$

which implies that  $\psi_T \circ \phi_T$  induces the identity on  $H_p^1(Y)$ .  $\square$

This also concludes the proof of Theorem B.2.  $\square$

### References

[C] CORNULIER, Y., Dimension of asymptotic cones of Lie groups, *Topology* **1** (2008), 343–361.  
 [CT] CORNULIER, Y. and TESSERA, R., Quasi-isometrically embedded free sub-semigroups, *Geom. Topol.* **12** (2008), 461–473.  
 [CTV] CORNULIER, Y., TESSERA, R. and VALETTE, A., Isometric group actions on Banach spaces and representations vanishing at infinity, *Transform. Groups* **13** (2008), 125–147.

- [Gr1] GROMOV, M., Hyperbolic groups, in *Essays in Group Theory*, Math. Sci. Res. Inst. Publ. **8**, pp. 75–263, Springer, New York, 1987.
- [Gr2] GROMOV, M., Asymptotic invariants of infinite groups, in *Geometric Group Theory*, London Math. Soc. Lecture Note Ser. **182**, pp. 11–12, Cambridge University Press, London, 1993.
- [Gu] GUIVARC'H, Y., Croissance polynomiale et périodes des fonctions harmoniques, *Bull. Soc. Math. France* **101** (1973), 333–379.
- [HaS] HAZOD, W. and SIEBERT, E., Continuous automorphism groups on a locally compact group contracting modulo a compact subgroup and applications to stable convolution semigroups, *Semigroup Forum* **33** (1986), 111–143.
- [He] HEINTZE, E., On homogeneous manifolds of negative curvature, *Math. Ann.* **211** (1974), 23–34.
- [MZ] MONTGOMERY, D. and ZIPPIN, L., *Topological Transformation Groups*, Interscience, New York, 1955.
- [Mo] MOSTOW, G., Self-adjoint groups, *Ann. of Math.* **62** (1955), 44–55.
- [Pa1] PANSU, P., Cohomologie  $L^p$  des variétés à courbure négative, cas du degré 1, in *Conference on Partial Differential Equations and Geometry (Torino, 1988)*, Rend. Sem. Mat. Univ. Politec. **1989**, Special Issue, pp. 95–120, Libreria Editrice Universitaria Levrotto & Bella, Turin, 1990.
- [Pa2] PANSU, P., Cohomologie  $L^p$  : invariance sous quasiisométrie, *Preprint*, 1995.
- [Pa3] PANSU, P., Cohomologie  $L^p$  en degré 1 des espaces homogènes, *Potential Anal.* **27** (2007), 151–165.
- [Pe] PEYROVIAN, M., Maximal compact normal subgroups, *Proc. Amer. Math. Soc.* **99** (1987), 389–394.
- [S] SIEBERT, E., Contractive automorphisms on locally compact groups, *Math. Z.* **191** (1986), 73–90.
- [T1] TESSERA, R., Large scale Sobolev inequalities on metric measure spaces and applications, *Rev. Mat. Iberoamericana* **24** (2008), 825–864.
- [T2] TESSERA, R., Vanishing of the first reduced cohomology with values in an  $L^p$ -representation, *Ann. Inst. Fourier (Grenoble)* **59** (2009), 851–876.
- [T3] TESSERA, R., Isoperimetric profile of subgroups and probability of return of random walks on elementary solvable groups, *Preprint*, 2009. [arXiv:0706.4257v3](https://arxiv.org/abs/0706.4257v3)

Yves Cornulier  
 Institut de recherche mathématique  
 de Rennes  
 Université de Rennes 1  
 263 avenue du Général Leclerc  
 FR-35042 Rennes Cedex  
 France  
[yves.decornulier@univ-rennes1.fr](mailto:yves.decornulier@univ-rennes1.fr)

Romain Tessera  
 Département de mathématiques (UMPA)  
 École normale supérieure de Lyon  
 46 allée d'Italie  
 FR-69364 Lyon Cedex 07  
 France  
[romain.tessera@umpa.ens-lyon.fr](mailto:romain.tessera@umpa.ens-lyon.fr)

*Received September 1, 2009*  
*published online September 22, 2010*