

## Research Article

# A Note on Some Properties of the Weighted $q$ -Genocchi Numbers and Polynomials

**L. C. Jang**

*Department of Mathematics and Computer Science, Konkuk University,  
Chungju 280-701, Republic of Korea*

Correspondence should be addressed to L. C. Jang, leechae.jang@kku.ac.kr

Received 30 July 2011; Revised 23 September 2011; Accepted 23 September 2011

Academic Editor: Mark A. Petersen

Copyright © 2011 L. C. Jang. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We consider the weighted  $q$ -Genocchi numbers and polynomials. From the construction of the weighted  $q$ -Genocchi numbers and polynomials, we investigate many interesting identities and relations satisfied by these new numbers and polynomials.

## 1. Introduction

Let  $p$  be a fixed odd prime number. Throughout this paper,  $\mathbb{Z}_p$ ,  $\mathbb{Q}_p$ ,  $\mathbb{C}$ , and  $\mathbb{C}_p$ , will, respectively, denote the ring of  $p$ -adic integers, the field, of  $p$ -adic rational numbers, the complex number field and the completion of algebraic closure of  $\mathbb{Q}_p$ . Let  $v_p$  be the normalized exponential valuation of  $\mathbb{C}_p$  such that  $|p|_p = p^{-v_p(p)} = 1/p$  (see [1–16]).

As well-known definition, the Euler numbers and Genocchi numbers are defined by

$$\frac{2}{e^t + 1} = e^{Et} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}, \quad (1.1)$$

with the usual convention of replacing  $E^n$  by  $E_n$  and

$$\frac{2t}{e^t + 1} = e^{Gt} = \sum_{n=0}^{\infty} G_n \frac{t^n}{n!}, \quad (1.2)$$

with the usual convention of replacing  $G^n$  by  $G_n$ . We assume that  $q \in \mathbb{C}_p$  with  $|1 - q|_p < 1$  and that the  $q$ -number of  $x$  is defined by

$$[x]_q = \frac{1 - q^x}{1 - q}, \quad [x]_{-q} = \frac{1 - (-q)^x}{1 + q} \quad (1.3)$$

(see [1–19]).

In [9], Kim introduced ordinary fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ , and he studied some interesting relations and identities related to  $q$ -extension of Euler numbers and polynomials. In [8], he also introduced the  $q$ -extension of the ordinary fermionic  $p$ -adic integral on  $\mathbb{Z}_p$  and he investigated many physical properties related to  $q$ -Euler numbers and polynomials. Recently, Kim firstly introduced the meaning of the weighted  $q$ -Euler numbers and polynomials associated with the weighted  $q$ -Bernstein polynomials by using the fermionic invariant  $p$ -adic integral on  $\mathbb{Z}_p$  (see [14, 15]). In [16], Ryoo tried to study the weighted  $q$ -Euler number and polynomials by the same method of Kim et al. in [14] and the  $q$ -extension of the fermionic  $p$ -adic invariant integrals on  $\mathbb{Z}_p$ . As well-known properties, the Genocchi numbers are integers. The first few Genocchi numbers for  $n = 2, 4, \dots$  are  $-1, 1, -3, 17, -155, 2073, \dots$ . The first few prime Genocchi numbers are  $-3$  and  $17$ , which occur for  $n = 6$  and  $8$ . There are no others with  $n < 10^5$ . These properties are very important to study in the area of fermionic distribution and  $p$ -adic numbers theory. By this reason, many mathematicians and physicians have studied Genocchi and Euler numbers which are in the different areas. By the same motivation, we consider weighted  $q$ -Genocchi polynomials and numbers by using the fermionic  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$  which are constructed by Kim and Ryoo (cf. [8, 16]).

In this paper, we consider the  $q$ -Genocchi numbers and polynomials with weighted  $\alpha$  ( $\alpha \in \mathbb{Q}$ ). From the construction of the weighted  $q$ -Genocchi numbers and polynomials, we investigate many interesting identities and relations satisfied by these new numbers and polynomials.

## 2. The Weighted $q$ -Genocchi Numbers and Polynomials

Let  $\text{UD}(\mathbb{Z}_p)$  be the space of uniformly differentiable functions and, for  $f \in \text{UD}(\mathbb{Z}_p)$ , the fermionic  $p$ -adic invariant integral of  $f$  on  $\mathbb{Z}_p$  is defined by Kim as follows:

$$I_{-1}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-1}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-1}} \sum_{x=0}^{p^N-1} f(x) (-1)^x \quad (2.1)$$

(see [1–16]). If we take  $f(x) = te^{xt}$ , then we get

$$t \int_{\mathbb{Z}_p} e^{xt} d\mu_{-1}(x) = \frac{2t}{e^t + 1}. \quad (2.2)$$

By (1.2) and (2.2), we get

$$\begin{aligned} \sum_{n=0}^{\infty} G_n \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} x^n d\mu_{-1}(x) \frac{t^{n+1}}{n!} \\ &= \sum_{n=0}^{\infty} (n+1) \int_{\mathbb{Z}_p} x^n d\mu_{-1}(x) \frac{t^{n+1}}{(n+1)!} \\ &= \sum_{n=1}^{\infty} n \int_{\mathbb{Z}_p} x^{n-1} d\mu_{-1}(x) \frac{t^n}{n!}. \end{aligned} \quad (2.3)$$

From (2.3),

$$G_0 = 0, \quad \frac{G_n}{n} = \int_{\mathbb{Z}_p} x^{n-1} d\mu_{-1}(x), \quad n \in \mathbb{N}. \quad (2.4)$$

For  $f \in \text{UD}(\mathbb{Z}_p)$ , the fermionic  $p$ -adic  $q$ -integral of  $f$  on  $\mathbb{Z}_p$  is defined by Kim as follows:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(x) d\mu_{-q}(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_{-q}} \sum_{x=0}^{p^N-1} f(x) (-q)^x \quad (2.5)$$

(see [1–16]). From (2.5), we note that

$$q^n I_{-q}(f_n) = (-1)^n I_{-q}(f) + [2]_q \sum_{l=0}^{n-1} (-1)^{n-1-l} q^l f(l), \quad (2.6)$$

where  $n \in \mathbb{N}$  and  $f_n(x) = f(x+n)$ .

For  $\alpha \in \mathbb{Q}$ , we consider the following fermionic  $p$ -adic  $q$ -integral on  $\mathbb{Z}_p$ :

$$t \int_{\mathbb{Z}_p} e^{[x]_q \alpha t} d\mu_{-q}(x) = \sum_{n=0}^{\infty} \tilde{G}_{n,q}^{(\alpha)} \frac{t^n}{n!}, \quad (2.7)$$

where  $\tilde{G}_{n,q}^{(\alpha)}$  are called the  $n$ th  $q$ -Genocchi numbers with weight  $\alpha$ . From (2.7), we get

$$\begin{aligned} t \int_{\mathbb{Z}_p} e^{[x]_q \alpha t} d\mu_{-q}(x) &= t \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} (n+1) \int_{\mathbb{Z}_p} [x]_q^n d\mu_{-q}(x) \frac{t^{n+1}}{(n+1)!} \\ &= \sum_{n=1}^{\infty} n \int_{\mathbb{Z}_p} [x]_q^{n-1} d\mu_{-q}(x) \frac{t^n}{n!}. \end{aligned} \quad (2.8)$$

By comparing the coefficients on the both sides of (2.7) and (2.8), we get

$$n \int_{\mathbb{Z}_p} [x]_{q^\alpha}^{n-1} d\mu_{-q}(x) = \tilde{G}_{n,q}^{(\alpha)}, \quad n \in \mathbb{N}, \quad \tilde{G}_{0,q}^{(\alpha)} = 0. \quad (2.9)$$

From (2.9), we obtain the following theorem.

**Theorem 2.1.** For  $n \in \mathbb{N}$  and  $\alpha \in \mathbb{Q}$ , one has

$$\int_{\mathbb{Z}_p} [x]_{q^\alpha}^{n-1} d\mu_{-q}(x) = \frac{\tilde{G}_{n,q}^{(\alpha)}}{n}, \quad \tilde{G}_{0,q}^{(\alpha)} = 0. \quad (2.10)$$

By the definition of fermionic  $p$ -adic  $q$ -integrals, we get

$$\begin{aligned} t \int_{\mathbb{Z}_p} [x]_{q^\alpha}^{n-1} d\mu_{-q}(x) &= \frac{1}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \int_{\mathbb{Z}_p} q^{\alpha l x} d\mu_{-q}(x) \\ &= \frac{[2]_q}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \frac{1}{1+q^{\alpha l+1}}. \end{aligned} \quad (2.11)$$

Therefore, we obtain the following theorem.

**Theorem 2.2.** For  $n \in \mathbb{N}$  and  $\alpha \in \mathbb{Q}$ , we have

$$\frac{\tilde{G}_{n,q}^{(\alpha)}}{n} = \frac{[2]_q}{(1-q^\alpha)^{n-1} [\alpha]_q^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \frac{1}{1+q^{\alpha l+1}}. \quad (2.12)$$

By Theorem 2.2, we have the generating function of  $\tilde{G}_{n,q}^{(\alpha)}$  as follows:

$$\begin{aligned} \sum_{n=0}^{\infty} \tilde{G}_{n,q}^{(\alpha)} \frac{t^n}{n!} &= [2]_q \sum_{n=0}^{\infty} \frac{n}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l \sum_{m=0}^{\infty} (-1)^m q^{\alpha l m + m} \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} \frac{n}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l q^{\alpha l m} \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} \frac{n}{(1-q^\alpha)^{n-1}} (1-q^{\alpha m})^{n-1} \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} n [m]_{q^\alpha}^{n-1} \frac{t^n}{n!} \end{aligned}$$

$$\begin{aligned}
&= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=1}^{\infty} [m]_{q^\alpha}^{n-1} \frac{t^n}{(n-1)!} \\
&= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} [m]_{q^\alpha}^n \frac{t^{n+1}}{n!} \\
&= [2]_q t \sum_{m=0}^{\infty} (-1)^m q^m e^{[m]_{q^\alpha} t}.
\end{aligned} \tag{2.13}$$

Let  $\tilde{F}_q^{(\alpha)}(t)$  be the generating function of  $\tilde{G}_{n,q}^{(\alpha)}$ . Then, by (2.9) and (2.13), we get

$$\begin{aligned}
\tilde{F}_q^{(\alpha)}(t) &= [2]_q t \sum_{m=0}^{\infty} (-1)^m q^m e^{[m]_{q^\alpha} t} \\
&= \sum_{n=0}^{\infty} \tilde{G}_{n,q}^{(\alpha)} \frac{t^n}{n!}.
\end{aligned} \tag{2.14}$$

The  $q$ -Genocchi polynomials with weight  $\alpha$  are defined by

$$\begin{aligned}
\tilde{F}_q^{(\alpha)}(t, x) &= t \int_{\mathbb{Z}_p} e^{[x+y]_{q^\alpha} t} d\mu_{-q}(y) \\
&= \sum_{n=0}^{\infty} \tilde{G}_{n,q}^{(\alpha)}(x) \frac{t^n}{n!}.
\end{aligned} \tag{2.15}$$

From (2.15), we get

$$\begin{aligned}
t \int_{\mathbb{Z}_p} e^{[x+y]_{q^\alpha} t} d\mu_{-q}(y) &= \sum_{n=0}^{\infty} \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^n d\mu_{-q}(y) \frac{t^{n+1}}{n!} \\
&= \sum_{n=0}^{\infty} (n+1) \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^n d\mu_{-q}(y) \frac{t^{n+1}}{(n+1)!} \\
&= \sum_{n=1}^{\infty} n \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^{n-1} d\mu_{-q}(y) \frac{t^n}{n!}.
\end{aligned} \tag{2.16}$$

By (2.15) and (2.16), we obtain the following theorem.

**Theorem 2.3.** For  $n \in \mathbb{N}$  and  $\alpha \in \mathbb{Q}$ , one has

$$n \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^{n-1} d\mu_{-q}(y) = \tilde{G}_{n,q}^{(\alpha)}(x), \quad \tilde{G}_{0,q}^{(\alpha)}(x) = 0. \tag{2.17}$$

We note that

$$\begin{aligned} \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^{n-1} d\mu_{-q}(y) &= \sum_{l=0}^{n-1} \binom{n-1}{l} [x]_{q^\alpha}^{n-1-l} q^{\alpha l x} \int_{\mathbb{Z}_p} [y]_{q^\alpha}^l d\mu_{-q}(y) \\ &= \sum_{l=0}^{n-1} \binom{n-1}{l} [x]_{q^\alpha}^{n-1-l} q^{\alpha l x} \frac{\tilde{G}_{l+1,q}^{(\alpha)}}{l+1}. \end{aligned} \quad (2.18)$$

From (2.17) and (2.18), we obtain the following theorem.

**Theorem 2.4.** For  $n \in \mathbb{N}$  and  $\alpha \in \mathbb{Q}$ , one has

$$\frac{\tilde{G}_{n,q}^{(\alpha)}(x)}{n} = \sum_{l=0}^{n-1} \binom{n-1}{l} [x]_{q^\alpha}^{n-1-l} q^{\alpha l x} \frac{\tilde{G}_{l+1,q}^{(\alpha)}}{l+1}. \quad (2.19)$$

From (2.15), we note that

$$\begin{aligned} \tilde{F}_q^{(\alpha)}(t, x) &= \sum_{n=0}^{\infty} \tilde{G}_{n,q}^{(\alpha)}(x) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} n \int_{\mathbb{Z}_p} [x+y]_{q^\alpha}^{n-1} d\mu_{-q}(y) \frac{t^n}{n!} \\ &= \sum_{n=0}^{\infty} n \left( \frac{1}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} q^{\alpha l x} (-1)^l \int_{\mathbb{Z}_p} q^{\alpha l y} d\mu_{-q}(y) \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{n=0}^{\infty} n \left( \frac{1}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} q^{\alpha l x} \frac{(-1)^l}{1+q^{\alpha(l+1)}} \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{n=0}^{\infty} n \left( \frac{1}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} q^{\alpha l x} (-1)^l \sum_{m=0}^{\infty} (-1)^m q^{\alpha(lm+m)} \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} n \left( \frac{1}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l q^{\alpha(x+m)l} \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} n \left( \frac{1}{(1-q^\alpha)^{n-1}} (1-q^{\alpha(x+m)})^{n-1} \right) \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} n [x+m]_{q^\alpha}^{n-1} \frac{t^n}{n!} \\ &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=1}^{\infty} [x+m]_{q^\alpha}^{n-1} \frac{t^n}{(n-1)!} \end{aligned}$$

$$\begin{aligned}
 &= [2]_q \sum_{m=0}^{\infty} (-1)^m q^m \sum_{n=0}^{\infty} [x+m]_{q^\alpha}^n \frac{t^{n+1}}{n!} \\
 &= [2]_q t \sum_{m=0}^{\infty} (-1)^m q^m e^{[x+m]_{q^\alpha} t}.
 \end{aligned}
 \tag{2.20}$$

Therefore, we obtain the following theorem.

**Theorem 2.5.** For  $\alpha \in \mathbb{Q}$ , one has

$$\tilde{F}_q^{(\alpha)}(t, x) = [2]_q t \sum_{m=0}^{\infty} (-1)^m q^m e^{[x+m]_{q^\alpha} t}.
 \tag{2.21}$$

From (2.15) and (2.21), we obtain that

$$\begin{aligned}
 \tilde{G}_{n,q}^{(\alpha)}(x) &= \left. \frac{d^n}{dt^n} \tilde{F}_q^{(\alpha)}(t, x) \right|_{t=0} \\
 &= n [2]_q \sum_{m=0}^{\infty} (-1)^m q^m [x+m]_{q^\alpha}^{m-1} \\
 &= n [2]_q \frac{1}{(1-q^\alpha)^{n-1}} \sum_{l=0}^{n-1} \frac{\binom{n-1}{l} q^{\alpha l x} (-1)^l}{1+q^{\alpha l+1}} \\
 &= \frac{n [2]_q}{(1-q)^{n-1} [\alpha]_q^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} (-1)^l q^{\alpha l x} \frac{1}{1+q^{\alpha l+1}}.
 \end{aligned}
 \tag{2.22}$$

Therefore, we obtain the following theorem.

**Theorem 2.6.** For  $n \in \mathbb{N}$  and  $\alpha \in \mathbb{Q}$ , one has

$$\tilde{G}_{n,q}^{(\alpha)}(x) = \frac{n [2]_q}{(1-q)^{n-1} [\alpha]_q^{n-1}} \sum_{l=0}^{n-1} \binom{n-1}{l} \frac{(-1)^l q^{\alpha l x}}{1+q^{\alpha l+1}}.
 \tag{2.23}$$

From (2.6), if we take  $f(x) = [x]_{q^\alpha}^m = ((1 - q^{\alpha x}) / (1 - q^\alpha))^m$ , then we get

$$q^n \int_{\mathbb{Z}_p} [x+n]_{q^\alpha}^m d\mu_{-q}(x) = (-1)^n \int_{\mathbb{Z}_p} [x]_{q^\alpha}^m d\mu_{-q}(x) + [2]_q \sum_{l=0}^{n-1} (-1)^l q^l [l]_{q^\alpha}^m.
 \tag{2.24}$$

By (2.17) and (2.24), we obtain the following theorem.

**Theorem 2.7.** For  $n \in \mathbb{N}$ ,  $m \in \mathbb{Z}_+ = \mathbb{N} \cup \{0\}$ , and  $\alpha \in \mathbb{Q}$ , one has

$$q^n \frac{\tilde{G}_{m+1,q}^{(\alpha)}(n)}{m+1} = (-1)^n \frac{\tilde{G}_{m+1,q}^{(\alpha)}}{m+1} + [2]_q \sum_{l=0}^{n-1} (-1)^l q^l [l]_{q^\alpha}^m. \quad (2.25)$$

We remark that if we take  $n = 2s$  ( $s \in \mathbb{Z}_+$ ) in Theorem 2.7, then we have

$$q^{2s} \frac{\tilde{G}_{m+1,q}^{(\alpha)}(2s)}{m+1} = \frac{\tilde{G}_{m+1,q}^{(\alpha)}}{m+1} + [2]_q \sum_{l=0}^{2s-1} (-1)^l q^l [l]_{q^\alpha}^m \quad (2.26)$$

and if we take  $n = 2s + 1$  ( $s \in \mathbb{Z}_+$ ) in Theorem 2.7, then we have

$$q^{2s+1} \frac{\tilde{G}_{m+1,q}^{(\alpha)}(2s+1)}{m+1} + \frac{\tilde{G}_{m+1,q}^{(\alpha)}}{m+1} = [2]_q \sum_{l=0}^{2s} (-1)^l q^l [l]_{q^\alpha}^m. \quad (2.27)$$

From (2.27) with  $s = 0$ , we obtain the following corollary.

**Corollary 2.8.** For  $\alpha \in \mathbb{Q}$  and  $m \in \mathbb{Z}_+$ , one has

$$q \frac{\tilde{G}_{m+1,q}^{(\alpha)}(1)}{m+1} + \frac{\tilde{G}_{m+1,q}^{(\alpha)}}{m+1} = \begin{cases} [2]_q & \text{if } m = 0, \\ 0 & \text{if } m > 0. \end{cases} \quad (2.28)$$

From (2.19), we note that

$$\begin{aligned} \frac{\tilde{G}_{m+1,q}^{(\alpha)}(n)}{m+1} &= \sum_{l=0}^m \binom{m}{l} [x]_{q^\alpha}^{m-l} \frac{\tilde{G}_{l+1,q}^{(\alpha)}}{l+1} q^{\alpha l x} \\ &= \frac{1}{m+1} \sum_{l=0}^m \binom{m+1}{l+1} [x]_{q^\alpha}^{m-l} \tilde{G}_{l+1,q}^{(\alpha)} q^{\alpha l x} \\ &= \frac{1}{m+1} \sum_{l=1}^m \binom{m+1}{l} [x]_{q^\alpha}^{m+1-l} \tilde{G}_{l,q}^{(\alpha)} q^{\alpha(l-1)x} \\ &= \frac{1}{q^\alpha} \frac{1}{m+1} \sum_{l=0}^m \binom{m+1}{l} [x]_{q^\alpha}^{m+1-l} \tilde{G}_{l,q}^{(\alpha)} q^{\alpha l x}. \end{aligned} \quad (2.29)$$

From (2.29), we get

$$\begin{aligned} q^\alpha \tilde{G}_{m+1,q}^{(\alpha)}(x) &= \sum_{l=0}^{m+1} \binom{m+1}{l} [x]_{q^\alpha}^{m+1-l} \tilde{G}_{l,q}^{(\alpha)} q^{\alpha l x} \\ &= \left( [x]_{q^\alpha} + q^{\alpha x} \tilde{G}_q^{(\alpha)} \right)^{m+1}, \end{aligned} \quad (2.30)$$

with the usual convention about replacing  $(\tilde{G}_q^{(\alpha)})^n$  by  $\tilde{G}_{n,q}^{(\alpha)}$ . By (2.28) and (2.30), we get

$$\frac{q^{1-\alpha} q^\alpha \tilde{G}_{m+1,q}^{(\alpha)}(1)}{m} + \frac{\tilde{G}_{m+1,q}^{(\alpha)}}{m+1} = \frac{q^{1-\alpha} (1 + q^\alpha \tilde{G}_q^{(\alpha)})^{m+1}}{m+1} + \frac{\tilde{G}_{m+1,q}^{(\alpha)}}{m+1}. \quad (2.31)$$

From (2.28) and (2.31), we obtain the following theorem.

**Theorem 2.9.** For  $\alpha \in \mathbb{Q}$  and  $m \in \mathbb{Z}_+$ , one has

$$q^{1-\alpha} (1 + q^\alpha \tilde{G}_q^{(\alpha)})^{m+1} + \tilde{G}_{m+1,q}^{(\alpha)} = \begin{cases} [2]_q & \text{if } m = 0, \\ 0 & \text{if } m > 0. \end{cases} \quad (2.32)$$

## Acknowledgment

This paper was supported by the Konkuk University in 2011.

## References

- [1] I. N. Cangul, V. Kurt, H. Ozden, and Y. Simsek, "On the higher-order  $w$ - $q$ -Genocchi numbers," *Advanced Studies in Contemporary Mathematics*, vol. 19, no. 1, pp. 39–57, 2009.
- [2] L.-C. Jang, "On multiple generalized  $w$ -Genocchi polynomials and their applications," *Mathematical Problems in Engineering*, vol. 2010, Article ID 316870, 8 pages, 2010.
- [3] L.-C. Jang, "A new  $q$ -analogue of Bernoulli polynomials associated with  $p$ -adic  $q$ -integrals," *Abstract and Applied Analysis*, vol. 2008, Article ID 295307, 6 pages, 2008.
- [4] L. Jang and T. Kim, " $q$ -Genocchi numbers and polynomials associated with fermionic  $p$ -adic invariant integrals on  $\mathbb{Z}_p$ ," *Abstract and Applied Analysis*, vol. 2008, Article ID 232187, 8 pages, 2008.
- [5] T. Kim, " $q$ -Volkenborn integration," *Russian Journal of Mathematical Physics*, vol. 9, no. 3, pp. 288–299, 2002.
- [6] T. Kim, "New approach to  $q$ -Euler polynomials of higher order," *Russian Journal of Mathematical Physics*, vol. 17, no. 2, pp. 218–225, 2010.
- [7] T. Kim, "New approach to  $q$ -Euler, Genocchi numbers and their interpolation functions," *Advanced Studies in Contemporary Mathematics*, vol. 18, no. 2, pp. 105–112, 2009.
- [8] T. Kim, "Barnes-type multiple  $q$ -zeta functions and  $q$ -Euler polynomials," *Journal of Physics A*, vol. 43, no. 25, Article ID 255201, 11 pages, 2010.
- [9] T. Kim, "Some identities on the  $q$ -Euler polynomials of higher order and  $q$ -Stirling numbers by the fermionic  $p$ -adic integral on  $\mathbb{Z}_p$ ," *Russian Journal of Mathematical Physics*, vol. 16, no. 4, pp. 484–491, 2009.
- [10] T. Kim, "On the  $q$ -extension of Euler and Genocchi numbers," *Journal of Mathematical Analysis and Applications*, vol. 326, no. 2, pp. 1458–1465, 2007.
- [11] T. Kim and B. Lee, "Some identities of the Frobenius-Euler polynomials," *Abstract and Applied Analysis*, vol. 2009, Article ID 639439, 7 pages, 2009.
- [12] T. Kim, L.-C. Jang, and H. Yi, "A note on the modified  $q$ -Bernstein polynomials," *Discrete Dynamics in Nature and Society*, vol. 2010, Article ID 706483, 12 pages, 2010.
- [13] T. Kim, J. Choi, and Y. H. Kim, " $q$ -Bernstein polynomials associated with  $q$ -Stirling numbers and Carlitz's  $q$ -Bernoulli numbers," *Abstract and Applied Analysis*, vol. 2010, Article ID 150975, 11 pages, 2010.
- [14] T. Kim, B. Lee, J. Choi, Y. H. Kim, and S. H. Rim, "On the  $q$ -Euler numbers and weighted  $q$ -bernstein polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 1, pp. 13–18, 2011.
- [15] T. Kim, "On the weighted  $q$ -Bernoulli numbers and polynomials," *Advanced Studies in Contemporary Mathematics (Kyungshang)*, vol. 21, no. 2, pp. 207–215, 2011.

- [16] C. S. Ryoo, "A note on the weighted  $q$ -Euler numbers and polynomials," *Advanced Studies in Contemporary Mathematics*, vol. 21, no. 1, pp. 47–54, 2011.
- [17] L. Carlitz, " $q$ -Bernoulli numbers and polynomials," *Duke Mathematical Journal*, vol. 15, pp. 987–1000, 1948.
- [18] B. A. Kupershmidt, "Reflection symmetries of  $q$ -Bernoulli polynomials," *Journal of Nonlinear Mathematical Physics*, vol. 12, pp. 412–422, 2005.
- [19] V. Kurt, "A further symmetric relation on the analogue of the Apostol-Bernoulli and the analogue of the Apostol-Genocchi polynomials," *Applied Mathematical Sciences*, vol. 3, no. 53–56, pp. 2757–2764, 2009.