

Research Article

Common Fixed Point Theorems of Altman Integral Type Mappings in G -Metric Spaces

Feng Gu and Hongqing Ye

Institute of Applied Mathematics and Department of Mathematics, Hangzhou Normal University, Zhejiang, Hangzhou 310036, China

Correspondence should be addressed to Feng Gu, gufeng99@sohu.com

Received 5 August 2012; Accepted 18 October 2012

Academic Editor: Yongfu Su

Copyright © 2012 F. Gu and H. Ye. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce the concept of φ -weakly commuting self-mapping pairs in G -metric space. Using this concept, we establish a new common fixed point theorem of Altman integral type for six self-mappings in the framework of complete G -metric space. An example is provided to support our result. The results obtained in this paper differ from the recent relative results in the literature.

1. Introduction and Preliminaries

Metric fixed point theory is an important mathematical discipline because of its applications in areas as variational and linear inequalities, optimization theory. Many results have been obtained by many authors considering different contractive conditions for self-mappings in metric space. In 1975, Altman [1] proved a fixed point theorem for a mapping which satisfies the condition $d(fx, fy) \leq Q(d(x, y))$, where $Q : [0, +\infty) \rightarrow [0, +\infty)$ is an increasing function satisfying the following conditions:

- (i) $0 < Q(t) < t, t \in (0, \infty)$;
- (ii) $p(t) = t/(t - Q(t))$ is a decreasing function;
- (iii) for some positive number t_1 , there holds $\int_0^{t_1} p(t)dt < +\infty$.

Remark 1.1. By condition (i) and that Q is increasing, we know that $Q(0) = 0$ and $Q(t) = t \Leftrightarrow t = 0$.

Gu and Deng [2], Liu [3], Zhang [4], and Li and Gu [5] discussed common fixed point theorems for Altman type mappings in metric space. In 2006, a new structure of generalized

metric space was introduced by Mustafa and Sims [6] as an appropriate notion of generalized metric space called G -metric space. Abbas and Rhoades [7] initiated the study of common fixed point in generalized metric space. Recently, many fixed point and common fixed point theorems for certain contractive conditions have been established in G -metric spaces, and for more details, one can refer to [8–42]. Coupled fixed point problems have also been considered in partially ordered G -metric spaces (see [43–56]). However, no one has discussed the common fixed point theorems for two or three pairs combining Altman type mappings recently.

Inspired by that, the purpose of this paper is to study common fixed point problem of Altman integral type for six self-mappings in G -metric space. We introduce a new concept of φ -weakly commuting self-mapping pairs in G -metric space, and a new common fixed point theorem for six self-mappings has been established through this concept. The results obtained in this paper differ from the recent relative results in the literature.

Throughout the paper, we mean by $\mathbb{N}$ the set of all natural numbers.

Definition 1.2 (see [6]). Let X be a nonempty set, and let $G : X \times X \times X \rightarrow R^+$ be a function satisfying the following axioms:

- (G1) $G(x, y, z) = 0$ if $x = y = z$,
- (G2) $0 < G(x, x, y)$, for all $x, y \in X$ with $x \neq y$,
- (G3) $G(x, x, y) \leq G(x, y, z)$, for all $x, y, z \in X$ with $z \neq y$,
- (G4) $G(x, y, z) = G(x, z, y) = G(y, z, x) = \dots$ (symmetry in all three variables),
- (G5) $G(x, y, z) \leq G(x, a, a) + G(a, y, z)$ for all $x, y, z, a \in X$ (rectangle inequality),

then the function G is called a generalized metric, or, more specifically a G -metric on X and the pair (X, G) is called a G -metric space.

Definition 1.3 (see [6]). Let (X, G) be a G -metric space and let $\{x_n\}$ be a sequence of points in X , a point x in X is said to be the limit of the sequence $\{x_n\}$ if $\lim_{m,n \rightarrow \infty} G(x, x_n, x_m) = 0$, and one says that sequence $\{x_n\}$ is G -convergent to x .

Thus, if $x_n \rightarrow x$ in a G -metric space (X, G) , then for any $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that $G(x, x_n, x_m) < \varepsilon$, for all $n, m \geq N$.

Proposition 1.4 (see [6]). Let (X, G) be a G -metric space, then the followings are equivalent:

- (1) $\{x_n\}$ is G -convergent to x ,
- (2) $G(x_n, x_n, x) \rightarrow 0$ as $n \rightarrow \infty$,
- (3) $G(x_n, x, x) \rightarrow 0$ as $n \rightarrow \infty$,
- (4) $G(x_n, x_m, x) \rightarrow 0$ as $n, m \rightarrow \infty$.

Definition 1.5 (see [6]). Let (X, G) be a G -metric space. A sequence $\{x_n\}$ is called G -Cauchy sequence if, for each $\varepsilon > 0$ there exists a positive integer $N \in \mathbb{N}$ such that $G(x_n, x_m, x_l) < \varepsilon$ for all $n, m, l \geq N$; that is, $G(x_n, x_m, x_l) \rightarrow 0$ as $n, m, l \rightarrow \infty$.

Definition 1.6 (see [6]). A G -metric space (X, G) is said to be G -complete, if every G -Cauchy sequence in (X, G) is G -convergent in X .

Proposition 1.7 (see [6]). Let (X, G) be a G -metric space. Then the followings are equivalent.

- (1) The sequence $\{x_n\}$ is G -Cauchy;
- (2) For every $\epsilon > 0$, there exists $k \in \mathbb{N}$ such that $G(x_n, x_m, x_m) < \epsilon$, for all $n, m \geq k$.

Proposition 1.8 (see [6]). Let (X, G) be a G -metric space. Then the function $G(x, y, z)$ is jointly continuous in all three of its variables.

Definition 1.9 (see [6]). Let (X, G) and (X', G') be G -metric space, and $f : (X, G) \rightarrow (X', G')$ be a function. Then f is said to be G -continuous at a point $a \in X$ if and only if for every $\epsilon > 0$, there is $\delta > 0$ such that $x, y \in X$, and $G(a, x, y) < \delta$ imply $G'(f(a), f(x), f(y)) < \epsilon$. A function f is G -continuous at X if and only if it is G -continuous at all $a \in X$.

Proposition 1.10 (see [6]). Let (X, G) and (X', G') be G -metric space. Then $f : X \rightarrow X'$ is G -continuous at $x \in X$ if and only if it is G -sequentially continuous at x , that is, whenever $\{x_n\}$ is G -convergent to x , $\{f(x_n)\}$ is G -convergent to $f(x)$.

Proposition 1.11 (see [6]). Let (X, G) be a G -metric space. Then, for any x, y, z, a in X it follows that:

- (i) if $G(x, y, z) = 0$, then $x = y = z$,
- (ii) $G(x, y, z) \leq G(x, x, y) + G(x, x, z)$,
- (iii) $G(x, y, y) \leq 2G(y, x, x)$,
- (iv) $G(x, y, z) \leq G(x, a, z) + G(a, y, z)$,
- (v) $G(x, y, z) \leq (2/3)(G(x, y, a) + G(x, a, z) + G(a, y, z))$,
- (vi) $G(x, y, z) \leq G(x, a, a) + G(y, a, a) + G(z, a, a)$.

Definition 1.12 (see [8]). Self-mappings f and g of a G -metric space (X, G) are said to be compatible if $\lim_{n \rightarrow \infty} G(fgx_n, gfx_n, gfx_n) = 0$ and $\lim_{n \rightarrow \infty} G(gfx_n, fgx_n, fgx_n) = 0$, whenever $\{x_n\}$ is a sequence in X such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = t$, for some $t \in X$.

In 2010, Manro et al. [9] introduced the concept of weakly commuting mappings, R -weakly commuting mappings into G -metric space as follows.

Definition 1.13 (see [9]). A pair of self-mappings (f, g) of a G -metric space is said to be weakly commuting if

$$G(fgx, gfx, gfx) \leq G(fx, gx, gx), \quad \forall x \in X. \quad (1.1)$$

Definition 1.14 (see [9]). A pair of self-mappings (f, g) of a G -metric space is said to be R -weakly commuting, if there exists some positive real number R such that

$$G(fgx, gfx, gfx) \leq RG(fx, gx, gx), \quad \forall x \in X. \quad (1.2)$$

Remark 1.15. If $R \leq 1$, then R -weakly commuting mappings are weakly commuting.

Now we introduce the new concept of φ -weakly commuting mappings as follow.

Definition 1.16. A pair of self-mappings (f, g) of a G -metric space is said to be φ -weakly commuting, if there exists a continuous function $\varphi : [0, \infty) \rightarrow [0, \infty)$, $\varphi(0) = 0$, such that

$$G(fgx, gfx, gfx) \leq \varphi(G(fx, gx, gx)), \quad \forall x \in X. \quad (1.3)$$

Remark 1.17. Commuting mappings are weakly commuting mappings, but the reverse is not true. For example: let $X = [0, 1/2]$, $G(x, y, z) = |x - y| + |y - z| + |z - x|$, for all $x, y, z \in X$, define $f(x) = x/2$, $g(x) = x^2/2$, through a straightforward calculation, we have: $fgx = x^2/4$, $gfx = x^2/8$, $G(fgx, gfx, gfx) = G(x^2/4, x^2/8, x^2/8) = x^2/4$, but $G(fx, gx, gx) = |x - x^2| = x - x^2$, hence, $G(fgx, gfx, gfx) \leq G(fx, gx, gx)$, but $fgx \neq gfx$.

Remark 1.18. Weakly commuting mappings are R -weakly commuting mappings, but the reverse is not true. For example: let $X = [-1, 1]$, define $G(x, y, z) = |x - y| + |y - z| + |z - x|$, for all $x, y, z \in X$, $f(x) = |x|$, $g(x) = |x| - 1$, then $gfx = |x| - 1$, $fgx = 1 - |x|$, $|fx - gx| = 1$, $|fgx - gfx| = 2(1 - |x|)$, $G(fgx, gfx, gfx) = 2|fgx - gfx| = 4(1 - |x|) \leq 4 = 4|fx - gx| = G(fx, gx, gx)$, when $R = 2$, we get that f and g are R -weakly commuting mappings, but not weakly commuting mappings.

Remark 1.19. R -weakly commuting mappings are φ -weakly commuting mappings but the reverse is not true. For example: let $X = [0, +\infty)$, $G(x, y, z) = |x - y| + |y - z| + |z - x|$, for all $x, y, z \in X$, $f(x) = x^2/4$, $g(x) = x^2$, thus, we have $fgx = x^4/4$, $gfx = x^4/16$, $G(fgx, gfx, gfx) = (3/8)x^4$, $G(fx, gx, gx) = (3/2)x^2$. Let $\varphi(x) = (1/2)x^2$, then

$$G(fgx, gfx, gfx) = \frac{3}{8}x^4 \leq \frac{9}{8}x^4 = \frac{1}{2} \left(\frac{3}{2}x^2 \right)^2 = \varphi \left(\frac{3}{2}x^2 \right) = \varphi(G(fx, gx, gx)). \quad (1.4)$$

For any given $R > 0$, since $\lim_{x \rightarrow +\infty} (1/4)x^2 = +\infty$, there exists $x \in X$ such that $(1/4)x^2 > R$, so we get $G(fgx, gfx, gfx) = (1/4)x^2 G(fx, gx, gx) > RG(fx, gx, gx)$. Therefore, f and g are φ -weakly commuting mappings, but not R -weakly commuting mappings.

Lemma 1.20. Let $\delta(t)$ be Lebesgue integrable, and $\delta(t) > 0$, for all $t > 0$, let $F(x) = \int_0^x \delta(t)dt$, then $F(x)$ is an increasing function in $[0, +\infty)$.

Definition 1.21. Let f and g be self-mappings of a set X . If $w = fx = gx$ for some x in X , then x is called a coincidence point of f and g , and w is called s point of coincidence of f and g .

2. Main Results

In this paper, we denote $\phi : [0, +\infty) \rightarrow [0, +\infty)$ the function satisfying $0 < \phi(t) < t$, for all $t > 0$.

Theorem 2.1. Let (X, G) be a complete G -metric space and let S, T, R, f, g , and h be six mappings of X into itself. If there exists an increasing function $Q : [0, +\infty) \rightarrow [0, +\infty)$ satisfying the conditions (i)~(iii) and the following conditions:

- (iv) $S(X) \subseteq g(X), T(X) \subseteq h(X), R(X) \subseteq f(X)$,
- (v) $\int_0^{G(Sx, Ty, Rz)} \delta(t)dt \leq \phi \left(\int_0^{Q(G(fx, gy, hz))} \delta(t)dt \right)$, for all $x, y, z \in X$,

where $\delta(t)$ is a Lebesgue integrable function which is summable nonnegative such that

$$\int_0^\epsilon \delta(t) dt > 0, \quad \forall \epsilon > 0. \quad (2.1)$$

Then,

- (a) one of the pairs (S, f) , (T, g) , and (R, h) has a coincidence point in X ,
- (b) if (S, f) , (T, g) , and (R, h) are three pairs of continuous φ -weakly commuting mappings, then the mappings S, T, R, f, g , and h have a unique common fixed point in X .

Proof. Let x_0 be an arbitrary point in X , from the condition (iv), there exist $x_1, x_2, x_3 \in X$ such that

$$y_1 = Sx_0 = gx_1, \quad y_2 = Tx_1 = hx_2, \quad y_3 = Rx_2 = fx_3. \quad (2.2)$$

By induction, there exist two sequences $\{x_n\}$, $\{y_n\}$ in X , such that

$$y_{3n+1} = Sx_{3n} = gx_{3n+1}, \quad y_{3n+2} = Tx_{3n+1} = hx_{3n+2}, \quad y_{3n+3} = Rx_{3n+2} = fx_{3n+3}, \quad n \in \mathbb{N}. \quad (2.3)$$

If $y_n = y_{n+1}$ for some n , with $n = 3m$, then $p = x_{3m+1}$ is a coincidence point of the pair (S, f) ; if $y_{n+1} = y_{n+2}$ for some n , with $n = 3m$, then $p = x_{3m+2}$ is a coincidence point of the pair (T, g) ; if $y_{n+2} = y_{n+3}$ for some n , with $n = 3m$, then $p = x_{3m+3}$ is a coincidence point of the pair (R, h) .

On the other hand, if there exists $n_0 \in \mathbb{N}$ such that $y_{n_0} = y_{n_0+1} = y_{n_0+2}$, then $y_n = y_{n_0}$ for any $n \geq n_0$. This implies that $\{y_n\}$ is a G-Cauchy sequence.

In fact, if there exists $p \in \mathbb{N}$ such that $y_{3p} = y_{3p+1} = y_{3p+2}$, then applying the contractive condition (v) with $x = y_{3p}$, $y = y_{3p+1}$, and $z = y_{3p+2}$, and the property of ϕ , we get

$$\begin{aligned} \int_0^{G(y_{3p+1}, y_{3p+2}, y_{3p+3})} \delta(t) dt &= \int_0^{G(Sx_{3p}, Tx_{3p+1}, Rx_{3p+2})} \delta(t) dt \\ &\leq \phi \left(\int_0^{Q(G(fx_{3p}, gx_{3p+1}, hx_{3p+2}))} \delta(t) dt \right) \\ &= \phi \left(\int_0^{Q(G(y_{3p}, y_{3p+1}, y_{3p+2}))} \delta(t) dt \right) \\ &\leq \int_0^{Q(G(y_{3p}, y_{3p+1}, y_{3p+2}))} \delta(t) dt. \end{aligned} \quad (2.4)$$

From Lemma 1.20 and the property of Q , we have

$$G(y_{3p+1}, y_{3p+2}, y_{3p+3}) \leq Q(G(y_{3p}, y_{3p+1}, y_{3p+2})) = Q(0) = 0. \quad (2.5)$$

Which implies that $y_{3p+3} = y_{3p+1} = y_{3p}$. So we find $y_n = y_{3p}$ for any $n \geq 3p$. This implies that $\{y_n\}$ is a G-Cauchy sequence. The same conclusion holds if $y_{3p+1} = y_{3p+2} = y_{3p+3}$, or $y_{3p+2} = y_{3p+3} = y_{3p+4}$ for some $p \in \mathbb{N}$.

Without loss of generality, we can assume that $y_n \neq y_m$ for all $n, m \in \mathbb{N}$ and $n \neq m$.

Now we prove that $\{y_n\}$ is a G-Cauchy sequence in X .

Let $t_n = G(y_n, y_{n+1}, y_{n+2})$, then we have

$$t_{n+1} \leq Q(t_n) < t_n \quad (2.6)$$

for all $n \in \mathbb{N}$. Actually, from the condition (v), (2.3) and the property of ϕ , we have

$$\begin{aligned} \int_0^{t_{3n}} \delta(t) dt &= \int_0^{G(y_{3n}, y_{3n+1}, y_{3n+2})} \delta(t) dt = \int_0^{G(Rx_{3n-1}, Sx_{3n}, Tx_{3n+1})} \delta(t) dt \\ &= \int_0^{G(Sx_{3n}, Tx_{3n+1}, Rx_{3n-1})} \delta(t) dt \leq \phi \left(\int_0^{Q(G(fx_{3n}, gx_{3n+1}, hx_{3n-1}))} \delta(t) dt \right) \\ &= \phi \left(\int_0^{Q(G(y_{3n}, y_{3n+1}, y_{3n-1}))} \delta(t) dt \right) = \phi \left(\int_0^{Q(t_{3n-1})} \delta(t) dt \right) \\ &\leq \int_0^{Q(t_{3n-1})} \delta(t) dt. \end{aligned} \quad (2.7)$$

By Lemma 1.20 and the property of Q , we have

$$t_{3n} \leq Q(t_{3n-1}) < t_{3n-1}. \quad (2.8)$$

Again, using condition (v), (2.3) and the property of ϕ , we get

$$\begin{aligned} \int_0^{t_{3n+1}} \delta(t) dt &= \int_0^{G(y_{3n+1}, y_{3n+2}, y_{3n+3})} \delta(t) dt = \int_0^{G(Sx_{3n}, Tx_{3n+1}, Rx_{3n+2})} \delta(t) dt \\ &\leq \phi \left(\int_0^{Q(G(fx_{3n}, gx_{3n+1}, hx_{3n+2}))} \delta(t) dt \right) = \phi \left(\int_0^{Q(G(y_{3n}, y_{3n+1}, y_{3n+2}))} \delta(t) dt \right) \\ &= \phi \left(\int_0^{Q(t_{3n})} \delta(t) dt \right) \leq \int_0^{Q(t_{3n})} \delta(t) dt. \end{aligned} \quad (2.9)$$

From Lemma 1.20 and the property of Q we have

$$t_{3n+1} \leq Q(t_{3n}) < t_{3n}. \quad (2.10)$$

Similarly, we can get

$$\begin{aligned}
 \int_0^{t_{3n+2}} \delta(t) dt &= \int_0^{G(y_{3n+2}, y_{3n+3}, y_{3n+4})} \delta(t) dt = \int_0^{G(Sx_{3n+3}, Tx_{3n+1}, Rx_{3n+2})} \delta(t) dt \\
 &\leq \phi \left(\int_0^{Q(fx_{3n+3}, gx_{3n+1}, hx_{3n+2}))} \delta(t) dt \right) = \phi \left(\int_0^{Q(y_{3n+3}, y_{3n+1}, y_{3n+2}))} \delta(t) dt \right) \quad (2.11) \\
 &= \phi \left(\int_0^{Q(t_{3n+1})} \delta(t) dt \right) \leq \int_0^{Q(t_{3n+1})} \delta(t) dt.
 \end{aligned}$$

From Lemma 1.20 and property of Q , we have

$$t_{3n+2} \leq Q(t_{3n+1}) < t_{3n+1}. \quad (2.12)$$

Combining (2.8), (2.10), and (2.12), we know that the (2.6) holds. This implies that $\{t_n\}$ is a nonnegative sequence which is strictly decreasing, hence, $\{t_n\}$ is convergent and $t_{n+1} \leq Q(t_n) < t_n$, for all $n \in \mathbb{N}$.

For any $n, m \in \mathbb{N}$, $m > n$, by combining (G5), (G3), and (2.6), we have

$$\begin{aligned}
 G(y_n, y_m, y_m) &\leq \sum_{i=n}^{m-1} G(y_i, y_{i+1}, y_{i+1}) \leq \sum_{i=n}^{m-1} G(y_i, y_{i+1}, y_{i+2}) = \sum_{i=n}^{m-1} t_i \\
 &= \sum_{i=n}^{m-1} \frac{t_i(t_i - t_{i+1})}{t_i - t_{i+1}} \leq \sum_{i=n}^{m-1} \frac{t_i(t_i - t_{i+1})}{t_i - Q(t_i)} \leq \sum_{i=n}^{m-1} \int_{t_{i+1}}^{t_i} \frac{t}{t - Q(t)} dt = \int_{t_m}^{t_n} p(t) dt. \quad (2.13)
 \end{aligned}$$

From the convergence of the sequence $\{t_n\}$ and the condition (iii) we assure that

$$\lim_{n, m \rightarrow \infty} \int_{t_m}^{t_n} p(t) dt = 0. \quad (2.14)$$

Thus, $\{y_n\}$ is a G -Cauchy sequence in X , since (X, G) is a complete G -metric space, there exists $u \in X$ such that $\lim_{n \rightarrow \infty} y_n = u$, hence

$$\begin{aligned}
 \lim_{n \rightarrow \infty} y_{3n+1} &= \lim_{n \rightarrow \infty} Sx_{3n} = \lim_{n \rightarrow \infty} gx_{3n+1} = u, \\
 \lim_{n \rightarrow \infty} y_{3n+2} &= \lim_{n \rightarrow \infty} Tx_{3n+1} = \lim_{n \rightarrow \infty} hx_{3n+2} = u, \\
 \lim_{n \rightarrow \infty} y_{3n+3} &= \lim_{n \rightarrow \infty} Rx_{3n+2} = \lim_{n \rightarrow \infty} fx_{3n+3} = u.
 \end{aligned} \quad (2.15)$$

Since (S, f) are φ -weakly commuting mappings, thus we have

$$G(Sfx_{3n}, fSx_{3n}, fSx_{3n}) \leq \varphi(G(Sx_{3n}, fx_{3n}, fx_{3n})). \quad (2.16)$$

On taking $n \rightarrow \infty$ at both sides, noting that S and f are continuous mappings, we have

$$G(Su, fu, fu) \leq \varphi(G(u, u, u)) = \varphi(0) = 0. \quad (2.17)$$

Which gives that $Su = fu$. Similarly, we can get $Tu = gu, Ru = hu$.

By using condition (v) and the property of ϕ , we get

$$\int_0^{G(Su, Tu, Ru)} \delta(t) dt \leq \phi \left(\int_0^{Q(G(fu, gu, hu))} \delta(t) dt \right) \leq \int_0^{Q(G(fu, gu, hu))} \delta(t) dt. \quad (2.18)$$

Thus, by Lemma 1.20, (G5) and the property of Q , noting that $Su = fu, Tu = gu, Ru = hu$, we have

$$\begin{aligned} G(Su, Tu, Ru) &\leq Q(G(fu, gu, hu)) \leq G(fu, gu, hu) \\ &\leq G(fu, Su, Su) + G(Su, gu, hu) \\ &\leq G(fu, Su, Su) + G(hu, Ru, Ru) + G(Ru, gu, Su) \\ &\leq G(fu, Su, Su) + G(hu, Ru, Ru) + G(Ru, Tu, Su) + G(Tu, Tu, gu) \\ &= G(Su, Tu, Ru). \end{aligned} \quad (2.19)$$

Which implies that

$$Q(G(fu, gu, hu)) = G(Su, Tu, Ru) = G(fu, gu, hu). \quad (2.20)$$

By Remark 1.1, we have $G(fu, gu, hu) = 0$, therefore, $fu = gu = hu$. So, immediately, we can have $Su = Tu = Ru = fu = gu = hu$. Setting

$$z = Su = Tu = Ru = fu = gu = hu. \quad (2.21)$$

Since (S, f) are φ -weakly commuting mappings, we have

$$G(Sz, fz, fz) = G(Sfu, fSu, fSu) \leq \varphi(G(Su, fu, fu)) = \varphi(0) = 0. \quad (2.22)$$

Which gives that $Sz = fz$. By the same argument, we can get $Tz = gz, Rz = hz$, So we have $Sfu = fSu, Tgu = gTu, Rhu = hRu$. Again, by condition (v), we have

$$\int_0^{G(Sz, z, z)} \delta(t) dt = \int_0^{G(S^2u, Tu, Ru)} \delta(t) dt \leq \phi \left(\int_0^{Q(G(fSu, gu, hu))} \delta(t) dt \right) \leq \int_0^{Q(G(fSu, gu, hu))} \delta(t) dt. \quad (2.23)$$

By the Lemma 1.20 and the property of Q , we have

$$G(Sz, z, z) \leq Q(G(fSu, gu, hu)) \leq G(fSu, gu, hu) = G(Sfu, gu, hu) = G(Sz, z, z). \quad (2.24)$$

Which implies that $Q(G(Sfu, gu, hu)) = G(Sfu, gu, hu)$. Thus, by the property of Q , we have $Sfu = gu = hu$, hence, $z = Sz = fz$. Similarly, we can prove that $z = Tz = gz$, $z = Rz = hz$, so we get $z = Sz = Tz = Rz = fz = gz = hz$, which means that z is a common fixed point of S, T, R, f, g , and h .

Now, we will show the common fixed point of S, T, R, f, g , and h is unique. Actually, assume $w \neq z$ is another common fixed point of S, T, R, f, g , and h , then by condition (v), we have

$$\int_0^{G(z,w,w)} \delta(t)dt = \int_0^{G(Sz,Tw,Rw)} \delta(t)dt \leq \phi \left(\int_0^{Q(G(fz,gw,hw))} \delta(t)dt \right) \leq \int_0^{Q(G(fz,gw,hw))} \delta(t)dt. \quad (2.25)$$

By Lemma 1.20 and the property of Q , we have

$$G(z,w,w) \leq Q(G(fz,gw,hw)) = Q(G(z,w,w)) < G(z,w,w). \quad (2.26)$$

It is a contradiction, unless $z = w$, that is, S, T, R, f, g , and h have a unique common fixed point in X . This completes the proof of Theorem 2.1. $\square$

Remark 2.2. If we take: (1) $S = T = R$; (2) $f = g = h$; (3) $f = g = h = I$ (I is identity mapping); (4) $T = R$ and $g = h$; (5) $T = R, g = h = I$; (6) $y = z$ in Theorem 2.1, then several new results can be obtained.

Corollary 2.3. Let (X, G) be a complete G -metric space and let S, T, R, f, g , and h be six mappings of X into itself. If there exists an increasing function $Q : [0, +\infty) \rightarrow [0, +\infty)$ satisfying the conditions (i)~(iii) and the following conditions:

$$(iv) \ S(X) \subseteq g(X), T(X) \subseteq h(X), R(X) \subseteq f(X),$$

$$(v) \ G(Sx, Ty, Rz) \leq \phi(Q(G(fx, gy, hz))), \forall x, y, z \in X.$$

Then,

(a) one of the pairs (S, f) , (T, g) , and (R, h) has a coincidence point in X .

(b) if (S, f) , (T, g) , and (R, h) are three pairs of continuous φ -weakly commuting mappings, then the mappings S, T, R, f, g , and h have a unique common fixed point in X .

Proof. Taking $\delta(t) = 1$ in Theorem 2.1, the conclusion of Corollary 2.3 can be obtained from Theorem 2.1 immediately. This completes the proof of Corollary 2.3. $\square$

Now we give an example to support Corollary 2.3.

Example 2.4. Let $X = [0, \infty)$, $G(x, y, z) = |x - y| + |y - z| + |z - x|$, for all $x, y, z \in X$. Let $S, T, R, f, g, h : X \rightarrow X$ be defined by $Sx = x/8$, $Tx = x/16$, $Rx = x/32$, $fx = x$, $gx = x/2$,

$hx = x/4$. Clearly, we can get $S(X) \subseteq g(X)$, $T(X) \subseteq h(X)$, $R(X) \subseteq f(X)$. Through calculation, we have

$$\begin{aligned}
 G(Sx, Ty, Rz) &= G\left(\frac{x}{8}, \frac{y}{16}, \frac{z}{32}\right) \\
 &= \left|\frac{x}{8} - \frac{y}{16}\right| + \left|\frac{y}{16} - \frac{z}{32}\right| + \left|\frac{x}{8} - \frac{z}{32}\right| \\
 &= \frac{1}{8} \left(\left|x - \frac{y}{2}\right| + \left|\frac{y}{2} - \frac{z}{4}\right| + \left|\frac{z}{4} - x\right|\right), \\
 G(fx, gy, hz) &= G\left(x, \frac{y}{2}, \frac{z}{4}\right) = \left|x - \frac{y}{2}\right| + \left|\frac{y}{2} - \frac{z}{4}\right| + \left|\frac{z}{4} - x\right|.
 \end{aligned} \tag{2.27}$$

Thus, we have

$$G(Sx, Ty, Rz) = \frac{1}{8}G(fx, gy, hz). \tag{2.28}$$

Now we choose $\phi(t) = 3t/4$, and $Q(t) = t/2$, then we have $\phi(t) < t$ and $Q(t)$ satisfies (i)~(iii). Thus, we have

$$\begin{aligned}
 G(Sx, Ty, Rz) &= \frac{1}{8}G(fx, gy, hz) \leq \frac{3}{4} \cdot \frac{1}{2}G(fx, gy, hz) \\
 &= \frac{3}{4}Q(G(fx, gy, hz)) = \phi(Q(G(fx, gy, hz))).
 \end{aligned} \tag{2.29}$$

On the other hand, let $\varphi(u) = u/2$ for all $u \in [0, \infty)$, we have

$$\begin{aligned}
 G(Sfx, fSx, fSx) &= G\left(\frac{x}{8}, \frac{x}{8}, \frac{x}{8}\right) \leq \frac{1}{2} \cdot \frac{7x}{4} = \frac{1}{2}G(Sx, fx, fx) = \varphi(G(Sx, fx, fx)), \\
 G(Tgx, gTx, gTx) &= G\left(\frac{x}{32}, \frac{x}{32}, \frac{x}{32}\right) \leq \frac{1}{2} \cdot \frac{7x}{8} = \frac{1}{2}G(Tx, gx, gx) = \varphi(G(Tx, gx, gx)), \\
 G(Rhx, hRx, hRx) &= G\left(\frac{x}{128}, \frac{x}{128}, \frac{x}{128}\right) \leq \frac{1}{2} \cdot \frac{7x}{16} = \frac{1}{2}G(Rx, hx, hx) = \varphi(G(Rx, hx, hx)),
 \end{aligned} \tag{2.30}$$

for all $x \in X$. Which means that (S, f) , (T, g) , and (R, h) are three pairs of continuous φ -weakly commuting mappings in X . So that all the conditions of Corollary 2.3 are satisfied. Moreover, 0 is the unique common fixed point for all of the mappings S, T, R, f, g , and h .

Corollary 2.5. *Let (X, G) be a complete G -metric space, $S, T, R : X \rightarrow X$ are three self-mappings in X , and function $Q : [0, +\infty) \rightarrow [0, +\infty)$ satisfies conditions (i)~(iii) and the following condition:*

$$G(Sx, Ty, Rz) \leq \phi(Q(G(x, y, z))), \quad \forall x, y, z \in X. \tag{2.31}$$

Then S, T , and R have a unique common fixed point in X .

Proof. Taking $h = g = f = I$ in Corollary 2.3, where I is an identity mapping. Then the conclusion of Corollary 2.5 can be obtained from Corollary 2.3 immediately. This completes the proof of Corollary 2.5. $\square$

Corollary 2.6. *Let (X, G) be a complete G-metric space, $S : X \rightarrow X$ is a self-mapping in X , and function $Q : [0, +\infty) \rightarrow [0, +\infty)$ satisfies conditions (i)~(iii) and the following condition:*

$$G(Sx, Sy, Sz) \leq \phi Q(G(x, y, z)), \quad \forall x, y, z \in X. \quad (2.32)$$

Then S has a unique fixed point in X .

Proof. Taking $S = T = R$ in Corollary 2.5, the conclusion of Corollary 2.6 can be obtained from Corollary 2.5 immediately. This completes the proof of Corollary 2.6. $\square$

Acknowledgments

This work was supported by the National Natural Science Foundation of China (11071169, 11271105), the Natural Science Foundation of Zhejiang Province (Y6110287, LY12A01030), and the Physical Experiment Center in Hangzhou Normal University.

References

- [1] M. Altman, "A fixed point theorem in compact metric spaces," *The American Mathematical Monthly*, vol. 82, no. 2, pp. 827–829, 1975.
- [2] F. Gu and B. Deng, "Common fixed point for Altman type mappings," *Journal of Harbin Normal University*, vol. 17, no. 5, pp. 44–46, 2001.
- [3] Z. Liu, "On common fixed points of Altman type mappings," *Journal of Liaoning Normal University (Natural Science Edition)*, vol. 16, no. 1, pp. 1–4, 1993.
- [4] S. Zhang, "Common fixed point theorem of Altman type mappings," *Journal of Yantai Normal University*, vol. 16, no. 2, pp. 95–97, 2000.
- [5] Y. Li and F. Gu, "Common fixed point theorem of Altman integral type mappings," *Journal of Nonlinear Science and Applications*, vol. 2, no. 4, pp. 214–218, 2009.
- [6] Z. Mustafa and B. Sims, "A new approach to generalized metric spaces," *Journal of Nonlinear and Convex Analysis*, vol. 7, no. 2, pp. 289–297, 2006.
- [7] M. Abbas and B. E. Rhoades, "Common fixed point results for noncommuting mappings without continuity in generalized metric spaces," *Applied Mathematics and Computation*, vol. 215, no. 1, pp. 262–269, 2009.
- [8] R. K. Vats, S. Kumar, and V. Sihag, "Some common fixed point theorems for compatible mappings of type(A) in complete G-metric space," *Advances in Fuzzy Mathematics*, vol. 6, no. 1, pp. 27–38, 2011.
- [9] S. Manro, S. S. Bhatia, and S. Kumar, "Expansion mappings theorems in G-metric spaces," *International Journal of Contemporary Mathematical Sciences*, vol. 5, no. 51, pp. 2529–2535, 2010.
- [10] Z. Mustafa, H. Obiedat, and F. Awawdeh, "Some fixed point theorem for mapping on complete G-metric spaces," *Fixed Point Theory and Applications*, vol. 2008, Article ID 189870, 12 pages, 2008.
- [11] Z. Mustafa, W. Shatanawi, and M. Bataineh, "Existence of fixed point results in G-metric spaces," *International Journal of Mathematics and Mathematical Sciences*, vol. 2009, Article ID 283028, 10 pages, 2009.
- [12] Z. Mustafa and B. Sims, "Fixed point theorems for contractive mappings in complete G-metric spaces," *Fixed Point Theory and Applications*, vol. 2009, Article ID 917175, 10 pages, 2009.
- [13] M. Abbas, T. Nazir, and R. Saadati, "Common fixed point results for three maps in generalized metric space," *Advances in Difference Equations*, vol. 49, no. 1, pp. 1–20, 2011.
- [14] W. Shatanawi, "Fixed point theory for contractive mappings satisfying Φ -maps in G-metric spaces," *Fixed Point Theory and Applications*, vol. 2010, Article ID 181650, 9 pages, 2010.

- [15] R. Chugh, T. Kadian, A. Rani, and B. E. Rhoades, "Property P in G -metric spaces," *Fixed Point Theory and Applications*, vol. 2010, Article ID 401684, 12 pages, 2010.
- [16] M. Abbas, T. Nazir, and S. Radenović, "Some periodic point results in generalized metric spaces," *Applied Mathematics and Computation*, vol. 217, no. 8, pp. 4094–4099, 2010.
- [17] Z. Mustafa, F. Awawdeh, and W. Shatanawi, "Fixed point theorem for expansive mappings in G -metric spaces," *International Journal of Contemporary Mathematical Sciences*, vol. 5, no. 50, pp. 2463–2472, 2010.
- [18] M. Abbas, S. H. Khan, and T. Nazir, "Common fixed points of R -weakly commuting maps in generalized metric spaces," *Fixed Point Theory and Applications*, vol. 2011, Article ID 784595, 11 pages, 2011.
- [19] H. Aydi, W. Shatanawi, and C. Vetro, "On generalized weak G -contraction mapping in G -metric spaces," *Computers & Mathematics with Applications*, vol. 62, no. 11, pp. 4223–4229, 2011.
- [20] M. Abbas, A. R. Khan, and T. Nazir, "Coupled common fixed point results in two generalized metric spaces," *Applied Mathematics and Computation*, vol. 217, no. 13, pp. 6328–6336, 2011.
- [21] W. Shatanawi, M. Abbas, and T. Nazir, "Common coupled coincidence and coupled fixed point results in two generalized metric spaces," *Fixed Point Theory and Applications*, vol. 2011, article 80, 2011.
- [22] H. Aydi, "A fixed point result involving a generalized weakly contractive condition in G -metric spaces," *Bulletin Mathematical Analysis and Applications*, vol. 3, no. 4, pp. 180–188, 2011.
- [23] Z. Mustafa, M. Khandagji, and W. Shatanawi, "Fixed point results on complete G -metric spaces," *Studia Scientiarum Mathematicarum Hungarica*, vol. 48, no. 3, pp. 304–319, 2011.
- [24] W. Shatanawi, "Coupled fixed point theorems in generalized metric spaces," *Haceteppe Journal of Mathematics and Statistics*, vol. 40, no. 3, pp. 441–447, 2011.
- [25] N. Hussain, M. H. Shah, and S. Radenović, "Fixed points of weakly contractions through occasionally weak compatibility," *Journal of Computational Analysis and Applications*, vol. 13, no. 3, pp. 532–543, 2011.
- [26] M. Abbas, T. Nazir, and D. Dorić, "Common fixed point of mappings satisfying (E.A) property in generalized metric spaces," *Applied Mathematics and Computation*, vol. 218, no. 14, pp. 7665–7670, 2012.
- [27] Z. Mustafa, H. Aydi, and E. Karapinar, "On common fixed points in G -metric spaces using (E.A) property," *Computers & Mathematics with Applications*, vol. 64, no. 6, pp. 1944–1956, 2012.
- [28] Lj. Gajić and M. Stojaković, "On Ćirić generalization of mappings with a contractive iterate at a point in G -metric spaces," *Applied Mathematics and Computation*, vol. 219, no. 1, pp. 435–441, 2012.
- [29] H. Ye and F. Gu, "Common fixed point theorems for a class of twice power type contraction maps in G -metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 736214, 19 pages, 2012.
- [30] F. Gu, "Common fixed point theorems for six mappings in generalized metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 379212, 2012.
- [31] W. Long, M. Abbas, T. Nazir, and S. Radenović, "Common fixed point for two pairs of mappings satisfying (E.A) property in generalized metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 394830, 15 pages, 2012.
- [32] N. Tahat, H. Aydi, E. Karapinar, and W. Shatanawi, "Common fixed points for single-valued and multi-valued maps satisfying a generalized contraction in G -metric spaces," *Fixed Point Theory and Applications*, vol. 2012, article 48, 2012.
- [33] Z. Mustafa, "Some new common fixed point theorems under strict contractive conditions in G -metric spaces," *Journal of Applied Mathematics*, vol. 2012, Article ID 248937, 21 pages, 2012.
- [34] V. Popa and A. M. Patriciu, "A general fixed point theorem for pairs of weakly compatible mappings in G -metric spaces," *Journal of Nonlinear Science and its Applications*, vol. 5, pp. 151–160, 2012.
- [35] M. Khandagji, S. Al-Sharif, and M. Al-Khaleel, "Property P and some fixed point results on (ψ, φ) -weakly contractive G -metric spaces," *International Journal of Mathematics and Mathematical Sciences*, vol. 2012, Article ID 675094, 11 pages, 2012.
- [36] Z. Mustafa, "Common fixed points of weakly compatible mappings in G -metric spaces," *Applied Mathematical Sciences*, vol. 6, no. 92, pp. 4589–4600, 2012.
- [37] M. Aggarwal, R. Chugh, and R. Kamal, "Suzuki-type fixed point results in G -metric spaces and applications," *International Journal of Computer Applications*, vol. 47, no. 12, pp. 14–17, 2012.
- [38] M. Gujani, M. Aggarwal, and R. Chugh, "Common fixed point results in G -metric spaces and applications," *International Journal of Computer Applications*, vol. 43, no. 11, pp. 38–42, 2012.
- [39] E. Karapinar, A. Yildiz-Ulus, and İ. M. Erhan, "Cyclic contractions on G -metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 182947, 15 pages, 2012.
- [40] H. Aydi, E. Karapinar, and W. Shatanawi, "Tripled fixed point results in generalized metric spaces," *Journal of Applied Mathematics*, vol. 2012, Article ID 314279, 10 pages, 2012.
- [41] H. Aydi, "A common fixed point of integral type contraction in generalized metric spaces," *Journal of Advanced Mathematical Studies*, vol. 5, no. 1, pp. 111–117, 2012.

- [42] K. P. R. Rao, K. B. Lakshmi, Z. Mustafa, and V. C. C. Raju, "Fixed and related fixed point theorems for three maps in G -metric spaces," *Journal of Advanced Studies in Topology*, vol. 3, no. 4, pp. 12–19, 2012.
- [43] R. Saadati, S. M. Vaezpour, P. Vetro, and B. E. Rhoades, "Fixed point theorems in generalized partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 52, no. 5-6, pp. 797–801, 2010.
- [44] B. S. Choudhury and P. Maity, "Coupled fixed point results in generalized G -metric spaces," *Mathematical and Computer Modelling*, vol. 54, no. 1-2, pp. 73–79, 2011.
- [45] H. Aydi, B. Damjanović, B. Samet, and W. Shatanawi, "Coupled fixed point theorems for nonlinear contractions in partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 54, no. 9-10, pp. 2443–2450, 2011.
- [46] W. Shatanawi, "Some fixed point theorems in ordered G -metric spaces and applications," *Abstract and Applied Analysis*, vol. 2011, Article ID 126205, 11 pages, 2011.
- [47] N. V. Luong and N. X. Thuan, "Coupled fixed point theorems in partially ordered G -metric spaces," *Mathematical and Computer Modelling*, vol. 55, no. 3-4, pp. 1601–1609, 2012.
- [48] H. Aydi, M. Postolache, and W. Shatanawi, "Coupled fixed point results for (ψ, φ) -weakly contractive mappings in ordered G -metric spaces," *Computers & Mathematics with Applications*, vol. 63, no. 1, pp. 298–309, 2012.
- [49] M. Abbas, T. Nazir, and S. Radenović, "Common fixed point of generalized weakly contractive maps in partially ordered G -metric spaces," *Applied Mathematics and Computation*, vol. 218, no. 18, pp. 9383–9395, 2012.
- [50] Y. J. Cho, B. E. Rhoades, R. Saadati, B. Samet, and W. Shatanawi, "Nonlinear coupled fixed point theorems in ordered generalized metric spaces with integral type," *Fixed Point Theory and Applications*, vol. 2012, article 8, 2012.
- [51] H. Aydi, E. Karapinar, and W. Shatanawi, "Tripled common fixed point results for generalized contractions in ordered generalized metric spaces," *Fixed Point Theory and Applications*, vol. 2012, article 101, 2012.
- [52] M. Abbas, W. Sintunavarat, and P. Kumam, "Coupled fixed point of generalized contractive mappings on partially ordered G -metric spaces," *Fixed Point Theory and Applications*, vol. 2012, article 31, 2012.
- [53] H. K. Nashine, "Coupled common fixed point results in ordered G -metric spaces," *Journal of Nonlinear Science and its Applications*, vol. 5, no. 1, pp. 1–13, 2012.
- [54] Z. Mustafa, H. Aydi, and E. Karapinar, "Mixed g -monotone property and quadruple fixed point theorems in partially ordered metric space," *Fixed Point Theory and Applications*, vol. 2012, article 71, 2012.
- [55] W. Shatanawi, M. Abbas, H. Aydi, and N. Tahat, "Common coupled coincidence and coupled fixed points in G -metric spaces," *Nonlinear Analysis and Application*, vol. 2012, Article ID jnaa-00162, 16 pages, 2012.
- [56] A. Razani and V. Parvaneh, "On generalized weakly G -contractive mappings in partially ordered G -metric spaces," *Abstract and Applied Analysis*, vol. 2012, Article ID 701910, 18 pages, 2012.