

Research Article

Iterative Algorithms for General Multivalued Variational Inequalities

Yonghong Yao,¹ Muhammad Aslam Noor,^{2,3} Yeong-Cheng Liou,⁴ and Shin Min Kang⁵

¹ Department of Mathematics, Tianjin Polytechnic University, Tianjin 300387, China

² Mathematics Department, College of Science, King Saud University, Riyadh 1145, Saudi Arabia

³ Mathematics Department, COMSATS Institute of Information Technology, Islamabad 44000, Pakistan

⁴ Department of Information Management, Cheng Shiu University, Kaohsiung 833, Taiwan

⁵ Department of Mathematics and RINS, Gyeongsang National University, Jinju 660-701, Republic of Korea

Correspondence should be addressed to Shin Min Kang, smkang@gnu.ac.kr

Received 21 October 2011; Accepted 1 November 2011

Academic Editor: Khalida Inayat Noor

Copyright © 2012 Yonghong Yao et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We introduce and study some new classes of variational inequalities and the Wiener-Hopf equations. Using essentially the projection technique, we establish the equivalence between these problems. This equivalence is used to suggest and analyze some iterative methods for solving the general multivalued variational inequalities in conjunction with nonexpansive mappings. We prove a strong convergence result for finding the common element of the set of fixed points of a nonexpansive mapping and the set of solutions of the general multivalued variational inequalities under some mild conditions. Several special cases are also discussed.

1. Introduction

Variational inequality problems were initially studied by Stampacchia in 1964. Variational inequalities have applications in diverse disciplines such as partial differential equations, optimal control, optimization, mathematical programming, mechanics, and finance, see [1–33] and the references therein. Variational inequalities have been extended and generalized in several directions using novel and innovative techniques. It is a common practice to study these variational inequalities in the setting of convexity. It has been observed that the optimality conditions of the differentiable convex functions can be characterized by the variational inequalities. In recent years, it has been shown that the minimum of the differentiable nonconvex functions can also be characterized by the variational inequalities. Motivated and inspired by these developments, Noor [19] has introduced a new type of variational inequality involving two nonlinear operators, which is called the general

variational inequality. It is worth mentioning that this general variational inequality is remarkable different from the so-called general variational inequality which was introduced by Noor [16] in 1988. Noor [19] proved that the general variational inequalities are equivalent to nonlinear projection equations and the Wiener-Hopf equations by using the projection technique. Using this equivalent formulation, Noor [19] suggested and analyzed some iterative algorithms for solving the special general variational inequalities and further proved these algorithms have strong convergence. In this paper, we introduce and consider a new class of variational inequalities, which is called the general multivalued variational inequality. Using essentially the projection technique, we establish the equivalence between the multivalued variational inequalities and the multivalued Wiener-Hopf equations.

Related to the variational inequalities, we have the problem of finding the fixed points of the nonexpansive mappings, which is the subject of current interest in functional analysis. It is natural to consider a unified approach to these two different problems. Noor and Huang [21] considered the problem of finding the common element of the set of the solutions of variational inequalities and the set of the fixed points of the nonexpansive mappings. We use the Wiener-Hopf technique to suggest and analyze some iterative methods for finding the common element the common element of the set of fixed points of a nonexpansive mapping and the set of solutions of the special general variational inequalities. We also consider the convergence criteria of the proposed algorithms under suitable conditions. Several special cases are also discussed.

2. Preliminaries

Let C be a nonempty closed convex subset of a real Hilbert space H . Let $A : H \rightarrow 2^H$ be a multivalued mapping. Let $F, g : H \rightarrow H$ be two nonlinear operators. We consider the problem of finding $u \in C$ and $w \in A(u)$ such that

$$\langle F(u) + w, g(v) - u \rangle \geq 0, \quad \forall v \in H, g(v) \in C. \quad (2.1)$$

Inequality of type (2.1) is called the general multivalued variational inequality. We will denote the set of solutions of the special general variational inequality (2.1) by $SGVI(F, A, g)$. The general multivalued variational inequality (2.1) can be written in the following equivalent form, that is, find $u \in C$, $w \in A(u)$ and $g(u) \in C$ such that

$$\langle \rho(F(u) + w) + u - g(u), g(v) - u \rangle \geq 0, \quad \forall v \in H, g(v) \in C. \quad (2.2)$$

This equivalent formulation is very important and plays a crucial role in the development of the iterative methods for solving the general multivalued variational inequalities.

We now discuss several special cases.

Special Cases

(A) If $w = 0$, then (2.1) reduces to: find $u \in C$ such that

$$\langle F(u), g(v) - u \rangle \geq 0, \quad \forall v \in H, g(v) \in C, \quad (2.3)$$

which is called the general variational inequality, introduced and studied by Noor [19]. It has been shown that the minimum of a class of differentiable functions can be characterized by the general variational inequality of type (2.3).

(B) If $g \equiv I$, the identity operator, then (2.1) reduces to find $u \in C$ and $w \in A(u)$ such that

$$\langle F(u) + w, v - u \rangle \geq 0, \quad \forall v \in C, \quad (2.4)$$

which is known as the mildly nonlinear multivalued variational inequality and has been studied extensively.

If F and A are single-valued nonlinear operators, then problem (2.1) is equivalent to finding $u \in C$ such that

$$\langle F(u) + A(u), v - u \rangle \geq 0, \quad \forall v \in C, \quad (2.5)$$

which is known as the mildly nonlinear variational inequality, the origin of which can be traced back to Noor [15].

(C) If $w = 0$ and $g \equiv I$, then (2.1) reduces to: find $u \in C$ such that

$$\langle F(u), v - u \rangle \geq 0, \quad \forall v \in C, \quad (2.6)$$

which is wellknown as the variational inequality, originally introduced and studied by Stampacchia [24] in 1964. It is clear from the above discussion that general multivalued variational inequality is quite general one. It has been shown that a wide class of problems arising in various discipline of mathematical and engineering sciences can be studied via the general multivalued variational inequalities (2.1) and its special cases.

In the sequel, we need the following well-known lemma.

Lemma 2.1. For a given $z \in H$, $u \in C$ satisfies the inequality

$$\langle u - z, v - u \rangle, \quad \forall v \in C, \quad (2.7)$$

if and only if

$$u = P_C z, \quad (2.8)$$

where P_C is the projection of H into the closed convex set C .

By using Lemma 2.1, one can prove that the general multivalued variational inequality (2.1) is equivalent to the following fixed point problem.

Lemma 2.2. $u \in C$ is a solution of the special general variational inequality (2.1) if and only if $u \in C$ satisfies the relation

$$u = P_C [g(u) - \rho(F(u) + w)], \quad (2.9)$$

where $\rho > 0$ is a constant.

Related to the general multivalued variational inequality (2.1), we consider the problem of solving the Wiener-Hopf equations. Let $F, g : H \rightarrow H$ be two nonlinear operators and $A : H \rightarrow 2^H$ be a multi-valued relaxed monotone operator. Let $Q_C = I - gP_C$, where I is the identity operator. We consider the problem of finding $y \in H : w \in A(y)$ such that

$$FP_C y + w + \rho^{-1}Q_C y = 0, \quad (2.10)$$

which is called the special general multivalued Wiener-Hopf equations. We use $SGWH(F, A, g)$ to denote the set of solutions of the special general multivalued Wiener-Hopf equations. For different and suitable choice of the operators F, A , we can obtain various forms of the Wiener-Hopf equations, which have been studied by Noor [17], Shi [22], and others.

Using essentially the technique of Noor [17, 18] and applying Lemma 2.2, one can establish the equivalence between the Wiener-Hopf equations and the general multivalued variational inequalities (2.1). To convey an idea of the technique and for the sake of completeness, we include its proof.

Lemma 2.3. *If $u \in SGVI(F, A, g)$, then $y \in H$ and $w \in A(y)$ satisfy the general Wiener-Hopf equations (2.10), where*

$$\begin{aligned} y &= g(u) - \rho(F(u) + w), \\ u &= P_C y, \end{aligned} \quad (2.11)$$

where $\rho > 0$ is a constant.

Proof. Let $u \in SGVI(F, A, g)$. Then, from Lemma 2.2, we have

$$u = P_C [g(u) - \rho(F(u) + w)]. \quad (2.12)$$

Let

$$y = g(u) - \rho(F(u) + w), \quad \forall w \in A(u). \quad (2.13)$$

Then, we have

$$u = P_C y. \quad (2.14)$$

Therefore, from (2.13), we obtain

$$y = gP_C y - \rho(FP_C y + w). \quad (2.15)$$

It follows that

$$FP_C y + w + \rho^{-1}Q_C y = 0, \quad \forall w \in AP_C y, \quad (2.16)$$

where $Q_C = I - gP_C$, which is exactly the general Wiener-Hopf equations (2.10). This completes the proof. \square

Remark 2.4. Let $S : C \rightarrow C$ be a nonexpansive mapping. If $u \in F(S) \cap \text{SGVI}(F, A, g)$, then one can easily see

$$u = Su = SP_C[g(u) - \rho(F(u) + w)], \quad (2.17)$$

which implies that

$$u = (1 - \alpha_n)u + \alpha_n SP_C[g(u) - \rho(F(u) + w)], \quad (2.18)$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$.

Using Remark 2.4 and Lemma 2.3, we can suggest the following algorithm for finding the common element of the solutions set of the variational inequalities and the set of fixed points of a nonexpansive mapping.

Algorithm 2.5. For a given $x_0 \in H$ arbitrarily, let the sequence $\{x_n\}$ be generated by

$$\begin{aligned} y_{n+1} &= (1 - \alpha_n)x_n + \alpha_n[g(x_n) - \rho(F(x_n) + w_n)], \\ x_n &= SP_C y_n, \end{aligned} \quad (2.19)$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $\rho > 0$ is some constant.

Note that, if $S \equiv I$, then Algorithm 2.5 reduces to the following iterative method for solving the general variational inequalities.

Algorithm 2.6. For a given $x_0 \in H$ arbitrarily, let the sequence $\{x_n\}$ be generated by

$$\begin{aligned} y_{n+1} &= (1 - \alpha_n)x_n + \alpha_n[g(x_n) - \rho(F(x_n) + w_n)], \\ x_n &= P_C y_n, \end{aligned} \quad (2.20)$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $\rho > 0$ is some constant.

If $\{w_n\} = 0$, then Algorithm 2.5 reduces to the following iterative method for solving the general variational inequalities (2.3), which was considered by Noor [19].

Algorithm 2.7. For a given $x_0 \in H$ arbitrarily, let the sequence $\{x_n\}$ be generated by

$$\begin{aligned} y_{n+1} &= (1 - \alpha_n)x_n + \alpha_n(g(x_n) - \rho F(x_n)), \\ x_n &= SP_C y_n, \end{aligned} \quad (2.21)$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $\rho > 0$ is some constant.

If $g = S \equiv I$ and $\{w_n\} = 0$, then Algorithm 2.5 reduces to the following iterative method for solving the variational inequalities (2.6).

Algorithm 2.8. For a given $x_0 \in H$ arbitrarily, let the sequence $\{x_n\}$ be generated by

$$\begin{aligned} y_{n+1} &= (1 - \alpha_n)x_n + \alpha_n(x_n - \rho F(x_n)), \\ x_n &= P_C y_n, \end{aligned} \quad (2.22)$$

where $\{\alpha_n\}$ is a sequence in $(0, 1)$ and $\rho > 0$ is some constant.

We recall the well-known concepts. The multivalued mapping A is said to be γ -Lipschitzian if there exists a constant $\gamma > 0$ such that

$$\|w_1 - w_2\| \leq \gamma \|u - v\|, \quad \forall w_1 \in A(u), w_2 \in A(v). \quad (2.23)$$

Recall that a mapping $S : C \rightarrow C$ is called *nonexpansive* if

$$\|Sx - Sy\| \leq \|x - y\|, \quad \forall x, y \in H. \quad (2.24)$$

We will use $F(S)$ to denote the set of fixed points of S .

A mapping $F : C \rightarrow H$ is called α -strongly monotone if there exists a constant $\alpha > 0$ such that

$$\langle F(x) - F(y), x - y \rangle \geq \alpha \|x - y\|^2, \quad \forall x, y \in H, \quad (2.25)$$

and β -Lipschitz continuous if there exists a constant $\beta > 0$ such that

$$\|F(x) - F(y)\| \leq \beta \|x - y\|, \quad \forall x, y \in H. \quad (2.26)$$

3. Main Results

Now we state and prove our main result.

Theorem 3.1. *Let C be a nonempty closed convex subset of a real Hilbert space H . Let $F : C \rightarrow H$ be an α -strongly monotone and β -Lipschitz continuous mapping, $g : C \rightarrow H$ an σ -strongly monotone and δ -Lipschitz continuous mapping and $A : H \rightarrow 2^H$ be a γ -Lipschitz continuous mapping. Let $S : C \rightarrow C$ be a nonexpansive mapping such that $F(S) \cap \text{SGVI}(F, A, g) \neq \emptyset$. Assume that*

$$\begin{aligned} \left| \rho - \frac{\alpha - \gamma(1 - k)}{\beta^2 - \gamma^2} \right| &< \frac{\sqrt{(\alpha - \gamma(1 - k))^2 - (\beta^2 - \gamma^2)k(2 - k)}}{\beta^2 - \gamma^2}, \\ \alpha &> \gamma(1 - k) + \sqrt{(\beta^2 - \gamma^2)k(2 - k)}, \quad \gamma\rho < 1 - k, \quad k < 1, \end{aligned} \quad (3.1)$$

where

$$k = \sqrt{1 - 2\sigma + \delta^2},$$

$$\sum_{n=0}^{\infty} \alpha_n = \infty, \tag{3.2}$$

then the approximate solution $\{y_{n+1}\}$ obtained from Algorithm 2.5 converges strongly to $y \in \text{SGWH}(F, A, g)$.

Proof. Let $x^* \in F(S) \cap \text{SGVI}(F, A, g)$. Then, from Remark 2.4, we have

$$x^* = SP_C y,$$

$$y = (1 - \alpha_n)x^* + \alpha_n [g(x^*) - \rho(F(x^*) + w)], \tag{3.3}$$

where $y \in H$ and $w \in A(y)$ satisfy the general Wiener-Hopf equations (2.10).

From (2.19) and (3.1), we have

$$\|x_{n+1} - x^*\| = \|SP_C y_n - SP_C y\| \leq \|y_n - y\|. \tag{3.4}$$

From (2.19), we have

$$\begin{aligned} \|y_{n+1} - y\| &\leq (1 - \alpha_n)\|x_n - x^*\| \\ &\quad + \alpha_n \|g(x_n) - \rho(F(x_n) + w_n) - [g(x^*) - \rho(F(x^*) + w)]\| \\ &\leq (1 - \alpha_n)\|x_n - x^*\| + \alpha_n \|x_n - x^* - (g(x_n) - g(x^*))\| \\ &\quad + \alpha_n \|x_n - x^* - \rho(F(x_n) - F(x^*))\| + \rho\alpha_n \|w_n - w\|. \end{aligned} \tag{3.5}$$

Since g is an σ -strongly monotone and δ -Lipschitz continuous mapping, we have

$$\begin{aligned} &\|x_n - x^* - (g(x_n) - g(x^*))\|^2 \\ &= \|x_n - x^*\|^2 - 2\langle g(x_n) - g(x^*), x_n - x^* \rangle + \|g(x_n) - g(x^*)\|^2 \\ &\leq \|x_n - x^*\|^2 - 2\sigma\|x_n - x^*\|^2 + \delta^2\|x_n - x^*\|^2 \\ &= (1 - 2\sigma + \delta^2)\|x_n - x^*\|^2 = k^2\|x_n - x^*\|^2, \end{aligned} \tag{3.6}$$

where $k = \sqrt{1 - 2\sigma + \delta^2}$.

At the same time, we note that F is an α -strongly monotone and β -Lipschitz continuous mapping, so we have

$$\begin{aligned} & \|x_n - x^* - \rho(F(x_n) - F(x^*))\|^2 \\ &= \|x_n - x^*\|^2 - 2\rho\langle F(x_n) - F(x^*), x_n - x^* \rangle + \rho^2\|F(x_n) - F(x^*)\|^2 \\ &= (1 - 2\rho\alpha + \rho^2\beta^2)\|x_n - x^*\|^2. \end{aligned} \quad (3.7)$$

From (3.5)–(3.7), we have

$$\begin{aligned} \|y_{n+1} - y\| &\leq (1 - \alpha_n)\|x_n - x^*\| + \alpha_n\left(k + \rho\gamma + \sqrt{1 - 2\rho\alpha + \rho^2\beta^2}\right)\|x_n - x^*\| \\ &= (1 - \alpha_n)\|x_n - x^*\| + \alpha_n\theta\|x_n - x^*\|, \end{aligned} \quad (3.8)$$

where

$$\theta = k + \rho\gamma + \sqrt{1 - 2\rho\alpha + \rho^2\beta^2}. \quad (3.9)$$

Using (3.1), we see that $\theta < 1$. Substituting (3.4) into (3.8), we have

$$\begin{aligned} \|y_{n+1} - y\| &\leq \left[(1 - \alpha_n) + \left(k + \rho\gamma + \sqrt{1 - 2\rho\alpha + \rho^2\beta^2} \right) \alpha_n \right] \|y_n - y\| \\ &= [1 - (1 - \theta)\alpha_n]\|x_n - x^*\| \\ &\leq \prod_{i=0}^n [1 - (1 - \theta)\alpha_i] \|y_0 - y\|. \end{aligned} \quad (3.10)$$

Since $\sum_{n=0}^{\infty} \alpha_n$ diverges and $1 - \theta > 0$, we have $\prod_{i=0}^n [1 - (1 - \theta)\alpha_i] = 0$. Consequently, the sequence $\{y_n\}$ converges strongly to y in H , the required result. \square

4. Conclusion

One of the most difficult and important problems in variational inequalities is the development of an efficient numerical methods. One of the technique is called the projection method and its variant forms. Projection method represent an important tool for finding the approximate solution of various types of variational inequalities. The projection type methods were developed in 1970s. The main idea in this techniques is to establish the equivalence between the variational inequalities and the fixed point problem using the concept of projection. These methods have been extended and modified in various ways. Shi [22] considered the problem of solving a system of nonlinear projections, which are called the Wiener-Hopf equations. It has been shown by Shi [22] that the Wiener-Hopf equations are equivalent to the variational inequalities. It turns out that this alternative formulation is more general and flexible. It has been shown that the Wiener-Hopf equations provide

us a simple, natural, elegant, and convenient device to develop some efficient numerical methods for solving variational and complementarity problems. In this paper, we introduce and study some new classes of variational inequalities and Wiener-Hopf equations. Using essentially the projection technique, we establish the equivalence between these problems. This equivalence is used to suggest and analyze some iterative methods for solving the general multivalued variational inequalities in conjunction with nonexpansive mappings. We prove a strong convergence result for finding the common element of the set of fixed points of a nonexpansive mapping and the set of solutions of the general multivalued variational inequalities under some mild conditions. Several special cases are also discussed. The ideas and techniques of this paper may be a starting point for a wide range of novel and innovative applications in various fields.

Acknowledgments

The authors thank the referees for useful comments and suggestions. The research of Professor M. Aslam Noor is supported by the Visiting Professor Program of King Saud University, Riyadh, Saudi Arabia, and Research Grant: KSU.VPP.108.

References

- [1] L. C. Ceng, Q. H. Ansari, and J. C. Yao, "Relaxed extragradient iterative methods for variational inequalities," *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 1112–1123, 2011.
- [2] L. C. Ceng, M. Teboulle, and J. C. Yao, "Weak convergence of an iterative method for pseudomonotone variational inequalities and fixed-point problems," *Journal of Optimization Theory and Applications*, vol. 146, no. 1, pp. 19–31, 2010.
- [3] S. S. Chang, H. W. J. Lee, C. K. Chan, and J. A. Liu, "A new method for solving a system of generalized nonlinear variational inequalities in Banach spaces," *Applied Mathematics and Computation*, vol. 217, no. 16, pp. 6830–6837, 2011.
- [4] F. Cianciaruso, G. Marino, L. Muglia, and Y. Yao, "On a two-step algorithm for hierarchical fixed point problems and variational inequalities," *Journal of Inequalities and Applications*, vol. 2009, Article ID 208692, 13 pages, 2009.
- [5] F. Facchinei and J. S. Pang, *Finite-Dimensional Variational Inequalities and Complementarity Problems*, vol. 1-2 of *Springer Series in Operations Research*, Springer, New York, NY, USA, 2003.
- [6] R. Glowinski, *Numerical Methods for Nonlinear Variational Problems*, Springer, New York, 1984.
- [7] B. S. He, "A new method for a class of linear variational inequalities," *Mathematical Programming*, vol. 66, no. 2, pp. 137–144, 1994.
- [8] A. N. Iusem and B. F. Svaiter, "A variant of Korpelevich's method for variational inequalities with a new search strategy," *Optimization*, vol. 42, no. 4, pp. 309–321, 1997.
- [9] P. Jaillet, D. Lambertson, and B. Lapeyre, "Variational inequalities and the pricing of American options," *Acta Applicandae Mathematicae*, vol. 21, no. 3, pp. 263–289, 1990.
- [10] E. N. Khobotov, "Modification of the extra-gradient method for solving variational inequalities and certain optimization problems," *USSR Computational Mathematics and Mathematical Physics*, vol. 27, no. 5, pp. 120–127, 1987.
- [11] G. M. Korpelevich, "An extragradient method for finding saddle points and for other problems," *Ekonomika i Matematicheskie Metody*, vol. 12, no. 4, pp. 747–756, 1976.
- [12] P. Kumam, N. Petrot, and R. Wangkeeree, "Existence and iterative approximation of solutions of generalized mixed quasi-variational-like inequality problem in Banach spaces," *Applied Mathematics and Computation*, vol. 217, no. 18, pp. 7496–7503, 2011.
- [13] X. Lu, H. K. Xu, and X. Yin, "Hybrid methods for a class of monotone variational inequalities," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 71, no. 3-4, pp. 1032–1041, 2009.
- [14] G. Marino, L. Muglia, and Y. Yao, "Viscosity methods for common solutions of equilibrium and variational inequality problems via multi-step iterative algorithms and common fixed points," *Nonlinear Analysis, Theory, Methods and Applications*. In press.

- [15] M. A. Noor, *On Variational Inequalities*, Brunel University, London, UK, 1975.
- [16] M. A. Noor, "General variational inequalities," *Applied Mathematics Letters*, vol. 1, no. 2, pp. 119–122, 1988.
- [17] M. A. Noor, "Wiener-Hopf equations and variational inequalities," *Journal of Optimization Theory and Applications*, vol. 79, no. 1, pp. 197–206, 1993.
- [18] M. A. Noor, "Some developments in general variational inequalities," *Applied Mathematics and Computation*, vol. 152, no. 1, pp. 199–277, 2004.
- [19] M. A. Noor, "Differentiable non-convex functions and general variational inequalities," *Applied Mathematics and Computation*, vol. 199, no. 2, pp. 623–630, 2008.
- [20] M. A. Noor and E. A. Al-Said, "Wiener-Hopf equations technique for quasimonotone variational inequalities," *Journal of Optimization Theory and Applications*, vol. 103, no. 3, pp. 705–714, 1999.
- [21] M. A. Noor and Z. Huang, "Wiener-Hopf equation technique for variational inequalities and nonexpansive mappings," *Applied Mathematics and Computation*, vol. 191, no. 2, pp. 504–510, 2007.
- [22] P. Shi, "Equivalence of variational inequalities with Wiener-Hopf equations," *Proceedings of the American Mathematical Society*, vol. 111, no. 2, pp. 339–346, 1991.
- [23] M. V. Solodov and B. F. Svaiter, "A new projection method for variational inequality problems," *SIAM Journal on Control and Optimization*, vol. 37, no. 3, pp. 765–776, 1999.
- [24] G. Stampacchia, "Formes bilineaires coercitives sur les ensembles convexes," *Comptes Rendus de l'Academie des Sciences*, vol. 258, pp. 4413–4416, 1964.
- [25] R. U. Verma, "Projection methods, algorithms, and a new system of nonlinear variational inequalities," *Computers & Mathematics with Applications*, vol. 41, no. 7-8, pp. 1025–1031, 2001.
- [26] H. K. Xu and T. H. Kim, "Convergence of hybrid steepest-descent methods for variational inequalities," *Journal of Optimization Theory and Applications*, vol. 119, no. 1, pp. 185–201, 2003.
- [27] J. C. Yao, "Variational inequalities with generalized monotone operators," *Mathematics of Operations Research*, vol. 19, no. 3, pp. 691–705, 1994.
- [28] Y. Yao, R. Chen, and H. K. Xu, "Schemes for finding minimum-norm solutions of variational inequalities," *Nonlinear Analysis. Theory, Methods & Applications*, vol. 72, no. 7-8, pp. 3447–3456, 2010.
- [29] Y. Yao, Y. C. Liou, and S. M. Kang, "Two-step projection methods for a system of variational inequality problems in Banach spaces," *Journal of Global Optimization*. In press.
- [30] Y. Yao, M. A. Noor, and Y. C. Liou, "Strong convergence of a modified extra-gradient method to the 4 minimum-norm solution of variational inequalities," *Abstract and Applied Analysis*, vol. 2012, Article ID 817436, 9 pages, 2012.
- [31] Y. Yao, M. A. Noor, K. I. Noor, Y. C. Liou, and H. Yaqoob, "Modified extragradient methods for a system of variational inequalities in Banach spaces," *Acta Applicandae Mathematicae*, vol. 110, no. 3, pp. 1211–1224, 2010.
- [32] Y. Yao and N. Shahzad, "New methods with perturbations for non-expansive mappings in Hilbert 5 spaces," *Fixed Point Theory and Applications*, vol. 2011, Article ID 79, 2011.
- [33] Y. Yao and N. Shahzad, "Strong convergence of a proximal point algorithm with general errors," *Optimization Letters*. In press.