

Research Article

On Convergence Results for Lipschitz Pseudocontractive Mappings

Shin Min Kang¹ and Arif Rafiq²

¹ Department of Mathematics and Research Institute of Natural Science, Gyeongsang National University, Jinju 660-701, Republic of Korea

² Hajvery University, 43-52 Industrial Area, Gulberg-III, Lahore, Pakistan

Correspondence should be addressed to Shin Min Kang, smkang@gnu.ac.kr

Received 4 June 2012; Revised 3 September 2012; Accepted 3 September 2012

Academic Editor: Alicia Cordero

Copyright © 2012 S. M. Kang and A. Rafiq. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We establish the strong convergence for the Ishikawa iteration scheme associated with Lipschitz pseudocontractive mappings in real Banach spaces. Moreover, our technique of proofs is of independent interest.

1. Introduction and Preliminaries

Let E be a real Banach space and K be a nonempty convex subset of E . Let J denote the normalized duality mapping from E to 2^{E^*} defined by

$$J(x) = \left\{ f^* \in E^* : \langle x, f^* \rangle = \|x\|^2, \|f^*\| = \|x\| \right\}, \quad \forall x \in E, \quad (1.1)$$

where E^* denotes the dual space of E and $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing. We will denote the single-valued duality mapping by j .

Let $T : D(T) \subset E \rightarrow E$ be a mapping with domain $D(T)$ in E .

Definition 1.1. T is said to be *Lipschitz* if there exists a constant $L > 1$ such that

$$\|Tx - Ty\| \leq L\|x - y\|, \quad \forall x, y \in D(T). \quad (1.2)$$

Definition 1.2. T is said to be *nonexpansive* if

$$\|Tx - Ty\| \leq \|x - y\|, \quad \forall x, y \in D(T). \quad (1.3)$$

Definition 1.3. T is said to be *pseudocontractive* if for all $x, y \in D(T)$, there exists $j(x - y) \in J(x - y)$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq \|x - y\|^2. \quad (1.4)$$

Remark 1.4. It is well known that every nonexpansive mapping is pseudocontractive. Indeed, if T is nonexpansive mapping, then for all $x, y \in D(T)$, there exists $j(x - y) \in J(x - y)$ such that

$$\langle Tx - Ty, j(x - y) \rangle \leq \|Tx - Ty\| \|x - y\| \leq \|x - y\|^2. \quad (1.5)$$

Rhoades [1] showed that the class of pseudocontractive mappings properly contains the class of nonexpansive mappings.

The class of pseudocontractions is, perhaps, the most important generalization of the class of nonexpansive mappings because of its strong relationship with the class of accretive mappings. A mapping $A : E \rightarrow E$ is accretive if and only if $I - A$ is pseudocontractive.

For a nonempty convex subset K of a normed space E and a mapping $T : K \rightarrow K$.

The Mann iteration scheme [2]: the sequence $\{x_n\}$ is defined by

$$\begin{aligned} x_1 &\in K, \\ x_{n+1} &= (1 - a_n)x_n + a_nTx_n, \quad n \geq 1, \end{aligned} \quad (1.6)$$

where $\{a_n\}$ is a sequence in $[0, 1]$.

The Ishikawa iteration scheme [3]: the sequence $\{x_n\}$ is defined by

$$\begin{aligned} x_1 &\in K, \\ x_{n+1} &= (1 - a_n)x_n + a_nTy_n, \\ y_n &= (1 - b_n)x_n + b_nTx_n, \quad n \geq 1, \end{aligned} \quad (1.7)$$

where $\{a_n\}$ and $\{b_n\}$ are sequences in $[0, 1]$.

In the last few years or so, numerous papers have been published on the iterative approximation of fixed points of Lipschitz strongly pseudocontractive mappings using the Ishikawa iteration scheme (e.g., [3]). Results which had been known only in Hilbert spaces and only for Lipschitz mappings have been extended to more general Banach spaces (e.g., [4-6] and the references cited therein).

In 1974, Ishikawa [3] introduced an iteration scheme which, in some sense, is more general than that of Mann and which converges, under this setting, to a fixed point of T . He proved the following result.

Theorem 1.5. *Let K be a compact convex subset of a Hilbert space H and let $T : K \rightarrow K$ be a Lipschitz pseudocontractive mapping. For arbitrary $x_1 \in K$, let $\{x_n\}_{n=1}^{\infty}$ be a sequence defined iteratively by*

$$\begin{aligned}x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n, \\y_n &= (1 - \beta_n)x_n + \beta_n T x_n, \quad n \geq 1,\end{aligned}\tag{1.8}$$

where $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\beta_n\}_{n=1}^{\infty}$ are sequences satisfying conditions

- (i) $0 \leq \alpha_n \leq \beta_n < 1$;
- (ii) $\lim_{n \rightarrow \infty} \beta_n = 0$;
- (iii) $\sum_{n=1}^{\infty} \alpha_n \beta_n = \infty$.

Then $\{x_n\}_{n=1}^{\infty}$ converges strongly to a fixed point of T .

In [4], Chidume extended the results of Schu [7] from Hilbert spaces to the much more general class of real Banach spaces and approximated the fixed points of strongly pseudocontractive mappings.

In this paper, we establish the strong convergence for the Ishikawa iteration scheme associated with Lipschitz pseudocontractive mappings in real Banach spaces. Moreover, our technique of proofs is of independent interest.

2. Main Results

We will need the following results.

Lemma 2.1 (see [8]). *Let $J : E \rightarrow 2^E$ be the normalized duality mapping. Then, for any $x, y \in E$, one has*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle, \quad \forall j(x + y) \in J(x + y).\tag{2.1}$$

Lemma 2.2 (see [9]). *If there exists a positive integer N such that for all $n \geq N$, $n \in \mathbb{N}$ (the set of all positive integers)*

$$\rho_{n+1} \leq (1 - \delta_n^2)\rho_n + b_n,\tag{2.2}$$

where $\delta_n \in [0, 1)$, $\sum_{n=1}^{\infty} \delta_n^2 = \infty$ and $b_n = o(\delta_n)$, then

$$\lim_{n \rightarrow \infty} \rho_n = 0.\tag{2.3}$$

We now prove our main results.

Theorem 2.3. Let K be a nonempty closed convex subset of a real Banach space E and $T : K \rightarrow K$ be a Lipschitz pseudocontractive mapping such that $p \in F(T) := \{x \in K : Tx = x\}$. Let $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\beta_n\}_{n=1}^{\infty}$ be sequences in $[0, 1]$ satisfying the conditions:

- (iv) $\sum_{n=1}^{\infty} \alpha_n^2 = \infty$;
- (v) $\lim_{n \rightarrow \infty} \alpha_n = 0$;
- (vi) $\lim_{n \rightarrow \infty} \beta_n = 0$.

For arbitrary $x_1 \in K$, let $\{x_n\}_{n=1}^{\infty}$ be defined iteratively by

$$\begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n, \\ y_n &= (1 - \beta_n)x_n + \beta_n T x_n, \quad n \geq 1. \end{aligned} \tag{2.4}$$

Then the following conditions are equivalent:

- (a) $\{x_n\}_{n=1}^{\infty}$ converges strongly to the fixed point p of T .
- (b) $\{T x_n\}_{n=1}^{\infty}$ and $\{T y_n\}_{n=1}^{\infty}$ are bounded.

Proof. Because p is a fixed point of T , then the set $F(T)$ of fixed points of T is nonempty. Suppose that $\lim_{n \rightarrow \infty} x_n = p$, then since T is Lipschitz, so

$$\begin{aligned} \lim_{n \rightarrow \infty} T x_n &= p, \\ \lim_{n \rightarrow \infty} y_n &= \lim_{n \rightarrow \infty} [(1 - \beta_n)x_n + \beta_n T x_n] = p, \end{aligned} \tag{2.5}$$

which implies that $\lim_{n \rightarrow \infty} T y_n = p$. Therefore $\{T x_n\}_{n=1}^{\infty}$ and $\{T y_n\}_{n=1}^{\infty}$ are bounded. Set

$$M_1 = \|x_0 - p\| + \sup_{n \geq 1} \|T x_n - p\| + \sup_{n \geq 1} \|T y_n - p\|. \tag{2.6}$$

Obviously $M_1 < \infty$.

It is clear that $\|x_0 - p\| \leq M_1$. Let $\|x_n - p\| \leq M_1$. Next we will prove that $\|x_{n+1} - p\| \leq M_1$. Consider

$$\begin{aligned} \|x_{n+1} - p\| &= \|(1 - \alpha_n)x_n + \alpha_n T y_n - p\| \\ &= \|(1 - \alpha_n)(x_n - p) + \alpha_n(T y_n - p)\| \\ &\leq (1 - \alpha_n) \|x_n - p\| + \alpha_n \|T y_n - p\| \\ &\leq M_1. \end{aligned} \tag{2.7}$$

So, from the above discussion, we can conclude that the sequence $\{x_n - p\}_{n=1}^{\infty}$ is bounded. Let $M_2 = \sup_{n \geq 1} \|x_n - p\|$.

Denote $M = M_1 + M_2$. Obviously $M < \infty$.

Now from Lemma 2.1 we obtain for all $n \geq 1$

$$\begin{aligned}
\|x_{n+1} - p\|^2 &= \|(1 - \alpha_n)x_n + \alpha_n T y_n - p\|^2 \\
&= \|(1 - \alpha_n)(x_n - p) + \alpha_n(T y_n - p)\|^2 \\
&\leq (1 - \alpha_n)^2 \|x_n - p\|^2 + 2\alpha_n \langle T y_n - p, j(x_{n+1} - p) \rangle \\
&= (1 - \alpha_n)^2 \|x_n - p\|^2 + 2\alpha_n \langle T x_{n+1} - p, j(x_{n+1} - p) \rangle \\
&\quad + 2\alpha_n \langle T y_n - T x_{n+1}, j(x_{n+1} - p) \rangle \\
&\leq (1 - \alpha_n)^2 \|x_n - p\|^2 + 2\alpha_n \|x_{n+1} - p\|^2 \\
&\quad + 2\alpha_n \|T y_n - T x_{n+1}\| \|x_{n+1} - p\| \\
&\leq (1 - \alpha_n)^2 \|x_n - p\|^2 + 2\alpha_n \|x_{n+1} - p\|^2 + 2\alpha_n \lambda_n,
\end{aligned} \tag{2.8}$$

where

$$\lambda_n = M \|T y_n - T x_{n+1}\|. \tag{2.9}$$

Using (2.4) we have

$$\begin{aligned}
\|y_n - x_{n+1}\| &\leq \|y_n - x_n\| + \|x_n - x_{n+1}\| \\
&= \beta_n \|x_n - T x_n\| + \alpha_n \|x_n - T y_n\| \\
&\leq 2M(\alpha_n + \beta_n).
\end{aligned} \tag{2.10}$$

From the conditions $\lim_{n \rightarrow \infty} \alpha_n = 0 = \lim_{n \rightarrow \infty} \beta_n$ and (2.10), we obtain

$$\lim_{n \rightarrow \infty} \|y_n - x_{n+1}\| = 0, \tag{2.11}$$

and since T is Lipschitz,

$$\lim_{n \rightarrow \infty} \|T y_n - T x_{n+1}\| = 0, \tag{2.12}$$

thus, we have

$$\lim_{n \rightarrow \infty} \lambda_n = 0. \tag{2.13}$$

The real function $f : [0, \infty) \rightarrow [0, \infty)$ defined by $f(t) = t^2$ is increasing and convex. For all $\lambda \in [0, 1]$ and $t_1, t_2 > 0$ we have

$$((1 - \lambda)t_1 + \lambda t_2)^2 \leq (1 - \lambda)t_1^2 + \lambda t_2^2. \tag{2.14}$$

Consider

$$\begin{aligned}
\|x_{n+1} - p\|^2 &= \|(1 - \alpha_n)x_n + \alpha_n T y_n - p\|^2 \\
&= \|(1 - \alpha_n)(x_n - p) + \alpha_n(T y_n - p)\|^2 \\
&\leq [(1 - \alpha_n)\|x_n - p\| + \alpha_n\|T y_n - p\|]^2 \\
&\leq (1 - \alpha_n)\|x_n - p\|^2 + \alpha_n\|T y_n - p\|^2 \\
&\leq (1 - \alpha_n)\|x_n - p\|^2 + M^2\alpha_n.
\end{aligned} \tag{2.15}$$

Substituting (2.15) in (2.8), we get

$$\begin{aligned}
\|x_{n+1} - p\|^2 &\leq [(1 - \alpha_n)^2 + 2\alpha_n(1 - \alpha_n)]\|x_n - p\|^2 \\
&\quad + 2\alpha_n(M^2\alpha_n + \lambda_n) \\
&= (1 - \alpha_n^2)\|x_n - p\|^2 + \varepsilon_n\alpha_n,
\end{aligned} \tag{2.16}$$

where $\varepsilon_n = 2(M^2\alpha_n + \lambda_n)$. Now, with the help of $\sum_{n=1}^{\infty} \alpha_n^2 = \infty$, $\lim_{n \rightarrow \infty} \alpha_n = 0$, (2.13), and Lemma 2.2, we obtain from (2.16) that

$$\lim_{n \rightarrow \infty} \|x_n - p\| = 0. \tag{2.17}$$

This completes the proof. □

Remark 2.4. Our technique of proofs is of independent interest.

Corollary 2.5. *Let K be a nonempty closed convex subset of a real Hilbert space E and let $T : K \rightarrow K$ be a Lipschitz pseudocontractive mapping such that $p \in F(T)$. Let $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\beta_n\}_{n=1}^{\infty}$ be sequences in $[0, 1]$ satisfying the conditions (iv), (v), and (vi).*

For arbitrary $x_1 \in K$, let $\{x_n\}_{n=1}^{\infty}$ be the sequence defined iteratively by (2.4). Then the following conditions are equivalent:

- (a) $\{x_n\}_{n=1}^{\infty}$ converges strongly to the fixed point p of T .
- (b) $\{Tx_n\}_{n=1}^{\infty}$ and $\{Ty_n\}_{n=1}^{\infty}$ are bounded.

The proof of the following result runs on the lines of proof of the Theorem 2.3, so is omitted.

Theorem 2.6. *Let K be a nonempty closed convex subset of a real Banach space E and let $T, S : K \rightarrow K$ be two Lipschitz pseudocontractive mappings such that $p \in F(T) \cap F(S) := \{x \in K : Tx = x =$*

Sx . Let $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\beta_n\}_{n=1}^{\infty}$ be sequences in $[0, 1]$ satisfying the conditions (iv), (v), and (vi). For arbitrary $x_1 \in K$, let $\{x_n\}_{n=1}^{\infty}$ be a sequence defined iteratively by

$$\begin{aligned}x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n, \\y_n &= (1 - \beta_n)x_n + \beta_n S x_n, \quad n \geq 1.\end{aligned}\tag{2.18}$$

Then the following conditions are equivalent:

- (a) $\{x_n\}_{n=0}^{\infty}$ converges strongly to the common fixed point p of T and S .
- (b) $\{T x_n\}_{n=0}^{\infty}$ and $\{S y_n\}_{n=0}^{\infty}$ are bounded.

Corollary 2.7. Let K be a nonempty closed convex subset of a real Hilbert space E and let $T, S : K \rightarrow K$ be two Lipschitz pseudocontractive mappings such that $p \in F(T) \cap F(S)$. Let $\{\alpha_n\}_{n=1}^{\infty}$ and $\{\beta_n\}_{n=1}^{\infty}$ be sequences in $[0, 1]$ satisfying conditions (iv), (v), and (vi).

For arbitrary $x_1 \in K$, let $\{x_n\}_{n=1}^{\infty}$ be the sequence defined iteratively by (2.18). Then the following conditions are equivalent:

- (a) $\{x_n\}_{n=0}^{\infty}$ converges strongly to the common fixed point p of T and S .
- (b) $\{T x_n\}_{n=0}^{\infty}$ and $\{S y_n\}_{n=0}^{\infty}$ are bounded.

Remark 2.8. It is worth to mentioning that we have the following.

- (1) The results of Chidume [4] and Zhou and Jia [10] depend on the geometry of the Banach space, whereas in our case we do not need such geometry.
- (2) We remove the boundedness assumption on K introduced in [4, 10].

Acknowledgments

The authors are grateful to the referees for their valuable comments and suggestions.

References

- [1] B. E. Rhoades, "A comparison of various definitions of contractive mappings," *Transactions of the American Mathematical Society*, vol. 226, pp. 257–290, 1977.
- [2] W. R. Mann, "Mean value methods in iteration," *Proceedings of the American Mathematical Society*, vol. 4, pp. 506–510, 1953.
- [3] S. Ishikawa, "Fixed points by a new iteration method," *Proceedings of the American Mathematical Society*, vol. 44, pp. 147–150, 1974.
- [4] C. E. Chidume, "Approximation of fixed points of strongly pseudocontractive mappings," *Proceedings of the American Mathematical Society*, vol. 120, no. 2, pp. 545–551, 1994.
- [5] C. E. Chidume and A. Udomene, "Strong convergence theorems for uniformly continuous pseudocontractive maps," *Journal of Mathematical Analysis and Applications*, vol. 323, no. 1, pp. 88–99, 2006.
- [6] X. Weng, "Fixed point iteration for local strictly pseudo-contractive mapping," *Proceedings of the American Mathematical Society*, vol. 113, no. 3, pp. 727–731, 1991.
- [7] J. Schu, "Approximating fixed points of Lipschitzian pseudocontractive mappings," *Houston Journal of Mathematics*, vol. 19, no. 1, pp. 107–115, 1993.
- [8] H. K. Xu, "Inequalities in Banach spaces with applications," *Nonlinear Analysis A*, vol. 16, no. 12, pp. 1127–1138, 1991.

- [9] A. Rafiq, "Some results on asymptotically pseudocontractive mappings," *Fixed Point Theory*, vol. 11, no. 2, pp. 355–360, 2010.
- [10] H. Zhou and Y. Jia, "Approximation of fixed points of strongly pseudocontractive maps without Lipschitz assumption," *Proceedings of the American Mathematical Society*, vol. 125, no. 6, pp. 1705–1709, 1997.