

Research Article

Tractable Approximation to Robust Nonlinear Production Frontier Problem

Lei Wang,¹ Xing Wang,² and Nan-jing Huang²

¹ Department of Economic Mathematics, South Western University of Finance and Economics, Sichuan, Chengdu 610074, China

² Department of Mathematics, Sichuan University, Chengdu 610064, China

Correspondence should be addressed to Nan-jing Huang, nanjinghuang@hotmail.com

Received 4 September 2012; Accepted 24 September 2012

Academic Editor: Jen-Chih Yao

Copyright © 2012 Lei Wang et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Robust optimization is a rapidly developing methodology for handling optimization problems affected by the uncertain-but-bounded data perturbations. In this paper, we consider the nonlinear production frontier problem where the traditional expected linear cost minimization objective is replaced by one that explicitly addresses cost variability. We propose a robust counterpart for the nonlinear production frontier problem that preserves the computational tractability of the nominal problem. We also provide a guarantee on the probability that the robust solution is feasible when the uncertain coefficients obey independent and identically distributed normal distributions.

1. Introduction

Optimization is a leading methodology in engineering design and control. Recently the *robust optimization* methodology has been introduced and studied [1–10] in order to deal with *uncertain* optimization problems: those for which part or all of their parameters are uncertain or inexact, or those for which the computed optimal solution cannot be physically implemented exactly. For such problems, a solution based on nominal values of the parameters may deviate severely from feasibility or optimality.

In robust optimization one is looking for a solution which satisfies the actual constraints for all possible realizations of the data within a fixed uncertainty set. This new problem is called the *robust counterpart* of the original problem. The robust counterpart is usually a semi-infinite optimization problem and in many cases can not be converted to explicitly convex optimization programs, solvable by high performance optimization techniques. For these cases the concept of an *approximate robust counterpart* was proposed by Ben-Tal and Nemirovski [1]. A solution of the approximate robust counterpart is always a robust solution of the original uncertain problem, and as shown in [1, 4], it can often be computed efficiently.

An increasingly active and challenging research area in growth and productivity studies is the nonsmooth frontier problems under a certain production technology as characterized with a production function, $y = f(x)$, $x \in X$, where $y \in \mathbb{R}$ is a scalar output and $X \subset \mathbb{R}_+^n$ is a given set of input possibilities. The non-smooth frontier problems are triggered by facts, as first noted by Solow [11], the production growth in many industries and countries follows a lagging and non-smooth (e.g., stepwise) trajectory as a function of production inputs, such as investments in IT capital and human capital. The interests in the study of nonlinear and non-smooth frontiers were further intensified by the reports at an FOMC meeting in late 1996 that the negative trends in measured productivity observed in many services industries seemed inconsistent with the fact that they ranked among the top computer-using industries (Carrado and Slifman [12]). Similarly, the issue of the nonsmoothness has been encountered in logistics and supply-chain systems. For example, differing container/trailer standards and heterogeneous cargo-handling characteristics of ports have been identified as a major issue of efficiency in ocean transport and short sea shipping as well (Perakis and Denisis [13]). In a recent econometric study on port logistics by Yan et al. [14], it is empirically identified that some heterogeneous and time-variant production factors, such as geographic location and business structure, affect significantly the efficiency of global container ports. With the devoted efforts over the past two decades, productivity researchers have collectively identified two key characteristics of sources of non-smooth growth, namely, degenerative input and nonlinear production cost. A degenerative input is defined as the input that generates stepwise output, and nonlinear production cost is expressed in the form of $\omega^T x + \varphi(x, D)$, where the nonlinear term $\varphi(x, D)$ represents intangible cost, in addition to the standard linear cost $\omega^T x$ and D is the vector of random coefficient. It shall be noted that production frontier problems studied so far assume a linear cost structure, that is, $\varphi(x, D) \equiv 0$. Under the aforementioned nonlinear cost, a nonlinear production frontier problem is constructed herein in the form of a nonlinear-cost minimization problem:

$$\begin{aligned} \min_x \quad & \omega^T x + \varphi(x, D) \\ \text{s.t.} \quad & x \in L(y) = \{x : f(x) \geq y\}, \quad \forall y \geq 0. \end{aligned} \tag{1.1}$$

We define the nominal problem to be problem (1.1) when the random coefficient D takes value equal to its expected value D_0 . In order to protect the solution against infeasibility of problem (1.1), we may formulate the problem using chance constraint as follows:

$$\begin{aligned} \min_{x, \tau} \quad & \tau \\ \text{s.t.} \quad & P(\omega^T x + \varphi(x, D) \leq \tau) \geq 1 - \epsilon, \\ & x \in L(y) = \{x : f(x) \geq y\}, \quad \forall y \geq 0. \end{aligned} \tag{1.2}$$

It is well known that such chance constraint is nonconvex and generally intractable. However, we will solve a tractable problem and obtain a *robust solution* that is feasible to the chance constraint problem (1.2) when ϵ is very small and without having to increase the objective function excessively. In order to address problem (1.1), Ben-Tal and Nemirovski [1, 3] and

independently by El-Ghaoui et al. [8, 9] propose to solve the following *robust optimization problem*

$$\begin{aligned} \min_{x, \tau} \quad & \tau \\ \text{s.t.} \quad & \omega^T x + \max_{D \in \mathcal{U}} \varphi(x, D) \leq \tau, \\ & x \in L(y) = \{x : f(x) \geq y\}, \quad \forall y \geq 0, \end{aligned} \quad (1.3)$$

where \mathcal{U} is a given uncertainty set. The motivation for solving problem (1.3) is to find a solution $x^* \in X$ that *immunizes* problem (1.1) against parameter uncertainty. That is, by selecting appropriate set \mathcal{U} , we can find a solution x^* to problem (1.3) that gives guarantee ϵ in problem (1.2). However, this is done at the expense of increasing the achievable objective. It is important to note that we describe uncertainty in problem (1.3) (using the set \mathcal{U}) in a deterministic manner. In selecting uncertainty set \mathcal{U} , we feel that two criteria are important.

- (a) Preserve the computational tractability both theoretically and most importantly practically of the nominal problem. From a theoretical perspective it is desirable that if the nominal problem is solvable in polynomial time, then the robust problem is also polynomially solvable.
- (b) Being able to find a guarantee on the probability that the robust solution is feasible, when the uncertain coefficients obey some natural probability distributions.

Our goal in this paper is to address (a) and (b) above for robust nonlinear production frontier problem. Specially, we propose a new robust counterpart of problem (1.1) that has the following properties. (a) It inherits the character of the nominal problem; (b) under reasonable probabilistic assumptions on data variation we establish probabilistic guarantee for feasibility that lead to explicit ways for selecting parameters that control the robustness. The structure of this paper is as follows. In Section 2, we describe the proposed robust model and in Section 3, we show that the robust nonlinear production frontier problem can be presented in a tractable manner. In Section 4, we prove probabilistic guarantee for feasibility for this problem.

2. Preliminaries

In this section, we outline the ingredients of the proposed framework for robust nonlinear production frontier problem.

2.1. Model for Parameter Uncertainty

The model of data uncertainty we consider is

$$D = D_0 + \sum_{j \in N} \Delta D_j u_j, \quad (2.1)$$

where D_0 is the nominal value of the data, $\Delta D_j, j \in N$ is a direction of data perturbation, and $u_j, j \in N$ are independent and identically distributed random variables with mean equal to

zero, so that $E[D] = D_0$. The cardinality of N may be small, modeling situations involving a small collection of primitive independent uncertainties, or large, potentially as large as the number of entries in the data. In the former case, the elements of D are strongly dependent, while in the latter case the elements of D are weakly dependent or even independent (when $|N|$ is equal to the number of entries in the data). The support of u_j , $j \in N$ can be unbounded or bounded. Ben-Tal and Nemirovski [3] and Bertsimas and Sim [6] have considered the case that $|N|$ is equal to the number of entries in the data.

2.2. Uncertainty Set and Related Norm

In the robust optimization framework (1.3), we consider the uncertainty set \mathcal{U} as follows:

$$\mathcal{U} = \left\{ D \mid \exists u \in \mathbb{R}^{|N|}: D = D_0 + \sum_{j \in N} \Delta D_j u_j, \|u\| \leq \Omega \right\}, \quad (2.2)$$

where Ω is a parameter, which Bertsimas and Sim [7] have showed, related to the probabilistic guarantee against infeasibility. Bertsimas and Sim [7] also restricted the vector norm $\|\cdot\|$ by imposing the condition:

$$\|u\| = \||u|\|, \quad (2.3)$$

where $|u| = (|u_1|, \dots, |u_{|N|}|)^T$ if $u = (u_1, \dots, u_{|N|})^T$ and they called such norm the *absolute norm*. Given a norm $\|\cdot\|$, we consider the dual norm $\|\cdot\|^*$ defined as

$$\|s\|^* = \max_{\|x\| \leq 1} s^T x. \quad (2.4)$$

We next show some basic properties of norms satisfying (2.3), which we will subsequently use in our development.

Proposition 2.1 (see [7]). *The absolute norm satisfies the following.*

- (1) $\|\omega\|^* = \||\omega|\|^*$;
- (2) for all v, ω such that $|v| \leq |\omega|$, $\|v\|^* \leq \|\omega\|^*$;
- (3) for all v, ω such that $|v| \leq |\omega|$, $\|v\| \leq \|\omega\|$.

2.3. The Class of Functions $\varphi(x, D)$

We impose the following restrictions on the function $\varphi(x, D)$ in problem (1.1).

Assumption 2.2. The function $\varphi(x, D)$ satisfies

- (a) the function $\varphi(x, D)$ is convex in D for all $x \in \mathbb{R}^n$;
- (b) $\varphi(x, kD) = k\varphi(x, D)$, for all $k \geq 0$, $x \in \mathbb{R}^n$.

Let the functions satisfy Assumption 2.2. Then it is easy to see that

$$\varphi(x, A + B) \leq \frac{1}{2}\varphi(x, 2A) + \frac{1}{2}\varphi(x, 2B) = \varphi(x, A) + \varphi(x, B). \quad (2.5)$$

3. The Proposed Robust Framework and Its Tractability

Specifically, under the model of data uncertainty in (2.1), we propose the constraint for controlling the feasibility of stochastic data uncertainty in constraint (1.3) as follows:

$$\omega^T x + \max_{(v,w) \in \mathcal{U}} \{ \varphi(x, D_0) + \sum_{j \in N} (\varphi(x, \Delta D_j) v_j + \varphi(x, -\Delta D_j) w_j) \} \leq \tau, \quad (3.1)$$

where

$$\mathcal{U} = \left\{ (v, w) \in \mathbb{R}_+^{|N| \times |N|} \mid \|v + w\| \leq \Omega \right\}, \quad (3.2)$$

and the norm $\|\cdot\|$ satisfies (2.3). We next show that under Assumption 2.2, (3.1) implies the classical definition of robustness:

$$\omega^T x + \varphi(x, D) \leq \tau, \quad \forall D \in \mathcal{U}, \quad (3.3)$$

where \mathcal{U} is defined in (2.2). Moreover, if the function $\varphi(x, D)$ is linear in D , then (3.1) is equivalent to (3.3).

Proposition 3.1. *Suppose the given norm $\|\cdot\|$ satisfies (2.3).*

- (a) *If $\varphi(x, A + B) = \varphi(x, A) + \varphi(x, B)$, then x satisfies (3.1) if and only if x satisfies (3.3).*
- (b) *Under Assumption 2.2, if x is feasible in problem (3.1), then x is feasible in problem (3.3).*

Proof. (a) Under the linearity assumption, (3.1) is equivalent to

$$\omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j (v_j - w_j)) \leq \tau, \quad \forall \|v + w\| \leq \Omega, \quad v, w \geq 0, \quad (3.4)$$

while (3.3) can be written as

$$\omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j u_j) \leq \tau, \quad \forall \|u\| \leq \Omega. \quad (3.5)$$

Suppose x is infeasible in (3.5). Then there exists u , $\|u\| \leq \Omega$ such that

$$\omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j u_j) > \tau. \quad (3.6)$$

For all $j \in N$, let $v_j = \max\{u_j, 0\}$ and $w_j = -\min\{u_j, 0\}$. Clearly, $r = v - w$ and since $v_j + w_j = |u_j|$, we have from (2.3) that

$$\|v + w\| = \|u\| \leq \Omega. \quad (3.7)$$

Hence, x is infeasible in (3.4) as well.

Conversely, suppose x is infeasible in (3.4), then there exists $v, w \geq 0$ and $\|v + w\| \leq \Omega$ such that

$$\omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j (v_j - w_j)) > \tau. \quad (3.8)$$

For all $j \in N$, we let $u_j = v_j - w_j$ and we observe that $|u_j| \leq v_j + w_j$. Therefore, for norms satisfying (2.3), we have

$$\|u\| = \||u\|| \leq \|v + w\| \leq \Omega, \quad (3.9)$$

and hence, x is infeasible in (3.5).

(b) Suppose x is feasible in problem (3.1). Then

$$\omega^T x + \varphi(x, D_0) + \sum_{j \in N} \{\varphi(x, \Delta D_j) v_j + \varphi(x, -\Delta D_j) w_j\} \leq \tau, \quad \forall \|v + w\| \leq \Omega, \quad v, w \geq 0. \quad (3.10)$$

From (2.5) and Assumption 2.2

$$\begin{aligned} \tau &\geq \omega^T x + \varphi(x, D_0) + \sum_{j \in N} \{\varphi(x, \Delta D_j) v_j + \varphi(x, -\Delta D_j) w_j\} \\ &\geq \omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j (v_j - w_j)) \end{aligned} \quad (3.11)$$

for all $\|v + w\| \leq \Omega$ and $v, w \geq 0$. In the proof of part (a), we showed that

$$\omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j u_j) \leq \tau, \quad \forall \|u\| \leq \Omega \quad (3.12)$$

is equivalent to

$$\omega^T x + \varphi(x, D_0 + \sum_{j \in N} \Delta D_j (v_j - w_j)) \leq \tau, \quad \forall \|v + w\| \leq \Omega, \quad v, w \geq 0. \quad (3.13)$$

Thus x satisfies (3.3). This completes the proof. \square

3.1. Tractability of the Proposed Framework

Unlike the classical definition of robustness (3.3), which cannot be represented in a tractable manner, we next show that (3.1) can be represented in a tractable manner.

Theorem 3.2. *For a norm satisfying (2.3) and a function $\varphi(x, D)$ satisfying Assumption 2.2, the following statements hold.*

(a) *Constraint (3.1) is equivalent to*

$$\omega^T x + \varphi(x, D_0) + \Omega \|s\|^* \leq \tau, \quad (3.14)$$

where

$$s_j = \max\{\varphi(x, \Delta D_j), \varphi(x, -\Delta D_j)\}, \quad \forall j \in N. \quad (3.15)$$

(b) Inequality (3.14) can be written as

$$\begin{aligned} \omega^T x + \varphi(x, D_0) + \Omega m &\leq \tau, \\ \varphi(x, \Delta D_j) - t_j &\leq 0, \quad \forall j \in N, \\ \varphi(x, -\Delta D_j) - t_j &\leq 0, \quad \forall j \in N, \\ \|t\|^* &\leq m, \\ m \in \mathbb{R}, \quad t &\in \mathbb{R}^{|N|}. \end{aligned} \quad (3.16)$$

Proof. (a) First, we introduce the following problems:

$$\begin{aligned} z_1 &= \max \quad a^T v + b^T w \\ \text{s.t.} \quad &\|v + w\| \leq \Omega, \\ &v, w \geq 0, \\ z_2 &= \max \quad \sum_{j \in N} \max\{a_j, b_j, 0\} u_j, \\ \text{s.t.} \quad &\|u\| \leq \Omega. \end{aligned} \quad (3.17)$$

In [7], Bertsimas and Sim showed that $z_1 = z_2$. Therefore, we observe that

$$\begin{aligned} &\max_{(v,w) \in \mathcal{U}} \sum_{j \in N} \{\varphi(x, \Delta D_j) v_j + \varphi(x, -\Delta D_j) w_j\} \\ &= \max_{\|u\| \leq \Omega} \sum_{j \in N} \{\max\{\varphi(x, \Delta D_j), \varphi(x, -\Delta D_j), 0\} u_j\} \end{aligned} \quad (3.18)$$

and using the definition of dual norm, $\|s\|^* = \max_{\|x\| \leq 1} s^T x$, we obtain that

$$\Omega \|s\|^* = \max_{\|x\| \leq \Omega} s^T x \quad (3.19)$$

and so (3.14) follows. Note that

$$s_j = \max\{\varphi(x, \Delta D_j), \varphi(x, -\Delta D_j)\} \geq 0, \quad (3.20)$$

since otherwise there exists an x such that $s_j < 0$, that is,

$$\varphi(x, \Delta D_j) < 0, \quad \varphi(x, -\Delta D_j) < 0. \quad (3.21)$$

From Assumption 2.2(b) $\varphi(x, 0) = 0$, contradicting the convexity of $\varphi(x, D)$ (Assumption 2.2(a)).

Suppose that x is feasible in Problem (3.14). Defining $t = s$ and $m = \|s\|^*$, we can easily check that (x, t, m) are feasible in problem (3.16). Conversely, suppose that x is infeasible in (3.14), that is,

$$\omega^T x + \varphi(x, D_0) + \Omega \|s\|^* > \tau. \quad (3.22)$$

Since

$$t_j \geq s_j = \max\{\varphi(x, \Delta D_j), \varphi(x, -\Delta D_j)\} \geq 0, \quad (3.23)$$

we apply Proposition 2.1(b) to obtain $\|t\|^* \geq \|s\|^*$ and so

$$\tau < \omega^T x + \varphi(x, D_0) + \Omega \|s\|^* \leq \omega^T x + \varphi(x, D_0) + \Omega \|t\|^* \leq \omega^T x + \varphi(x, D_0) + \Omega m. \quad (3.24)$$

Thus, x is infeasible in (3.16).

(b) It is immediate that (3.14) can be written in the form of (3.16).

This completes the proof. \square

Remark 3.3. In (1.1), let $\varphi(x, D) = 0$ and $L(y_i) = \{x \mid A_i x + y_i \in K_i\}$, $i = 1, \dots, m$. Then the optimization problem (1.1) will be in the conic form:

$$\begin{aligned} \min_x \quad & \omega^T x \\ \text{s.t.} \quad & A_i x + y_i \in K_i, \quad i = 1, \dots, m, \end{aligned} \quad (3.25)$$

where, for every i , K_i is

- (i) either a nonnegative orthant $\mathbb{R}_+^{m_i}$ (linear constraints),
- (ii) or the Lorentz cone $L^{m_i} = \{z \in \mathbb{R}^{m_i} \mid z_{m_i} \geq \sqrt{\sum_{j=1}^{m_i-1} z_j^2}\}$ (conic quadratic constraints),
- (iii) or a semidefinite cone $S_+^{m_i}$ —the cone of positive semidefinite matrices in the space S^{m_i} of $m_i \times m_i$ symmetric matrices (Linear Matrix Inequality constraints).

The class of problems which can be modeled in the form of (3.16) is extremely wide (see, e.g., [1–5]). It is also clear what is the structure and what are the data in (3.25)—the former is the design dimension n , the number of conic constraints m , and the list of the cones K_1, \dots, K_m ,

while the latter is the collection of matrices and vectors $\omega, \{A_i, y_i\}_{i=1}^m$ of appropriate sizes. Thus, an uncertain problem of (3.16) is a collection,

$$\begin{aligned} \min_{x, \tau} \quad & \tau \\ \text{s.t.} \quad & \tau - \omega^T x \in K_0 \equiv \mathbb{R}_+, \\ & A_i x + y_i \in K_i, \\ & (\omega, \{A_i, y_i\}_{i=1}^m) \in \mathcal{U}^*, \\ & i = 1, \dots, m, \end{aligned} \tag{3.26}$$

of instances (3.16) of a common structure $\{n, m, K_1, \dots, K_m\}$ and data $(\omega, \{A_i, y_i\}_{i=1}^m)$ varying in a given set \mathcal{U}^* . By specifying reasonable uncertainty sets \mathcal{U}^* in specific applications of K_i , Ben-Tal and Nemirovski [1–5] reformulate (3.26) as a computational tractable optimization problem, or at least approximate (3.26) by a tractable problem. Therefore, Theorem 3.2 generalizes model (3.16) from optimization problem with linear object function to a more widely nonlinear object function optimization problem.

3.2. Representation of the Function $\max\{\varphi(x, \Delta D_j), \varphi(x, -\Delta D_j)\}$

The function

$$g(x, \Delta D_j) = \max\{\varphi(x, \Delta D_j), \varphi(x, -\Delta D_j)\} \tag{3.27}$$

naturally arises in Theorem 3.2. Recall that a norm satisfies $\|A\| \geq 0$, $\|kA\| = |k| \cdot \|A\|$, $\|A+B\| \leq \|A\| + \|B\|$, and $\|A\| = 0$ implies that $A = 0$. We show next that the function $g(x, A)$ satisfies all these properties except the last one, that is, it behaves almost like a norm.

Proposition 3.4. *Under Assumption 2.2, the function*

$$g(x, A) = \max\{\varphi(x, A), \varphi(x, -A)\} \tag{3.28}$$

has the following properties:

- (a) $g(x, A) \geq 0$;
- (b) $g(x, kA) = |k|g(x, A)$;
- (c) $g(x, A + B) \leq g(x, A) + g(x, B)$.

Proof. (a) Suppose there exists x such that $g(x, A) < 0$, that is, $\varphi(x, A) < 0$ and $\varphi(x, -A) < 0$. From Assumption 2.2(b) $\varphi(x, 0) = 0$, contradicting the convexity of $\varphi(x, A)$ (Assumption 2.2(a)).

(b) For $k \geq 0$, we apply Assumption 2.2(b) and obtain

$$\begin{aligned} g(x, kA) &= \max\{\varphi(x, kA), \varphi(x, -kA)\} \\ &= k \max\{\varphi(x, A), \varphi(x, -A)\} \\ &= kg(x, A). \end{aligned} \quad (3.29)$$

Similarly, if $k < 0$ we have

$$\begin{aligned} g(x, kA) &= \max\{\varphi(x, -k(-A)), \varphi(x, -kA)\} \\ &= -kg(x, A). \end{aligned} \quad (3.30)$$

(c) Using (2.5) we obtain

$$\begin{aligned} g(x, A + B) &= g\left(x, \frac{1}{2}(2A + 2B)\right) \\ &\leq \frac{1}{2}g(x, 2A) + \frac{1}{2}g(x, 2B) \\ &= g(x, A) + g(x, B). \end{aligned} \quad (3.31)$$

This completes the proof. \square

Note that the function $g(x, A)$ does not necessarily define a norm for A , since $g(x, A) = 0$ does not necessarily imply $A = 0$. However, for specific direction of data perturbation, ΔD_j , we can map $g(x, \Delta D_j)$ to a function of norm such that

$$g(x, \Delta D_j) = \|\mathcal{L}(x, \Delta D_j)\|_{g'} \quad (3.32)$$

where $\mathcal{L}(x, \Delta D_j)$ is linear in ΔD_j .

3.3. Example

We consider the robust quadratically frontier problem, that is, $\varphi(x, D) = x^T Ax$. Then it is easy to see that $D = A$, $D_0 = A_0$, $\Delta D_j = \Delta A_j$, and the uncertainty set

$$\mathcal{U} = \left\{ A \mid \exists u \in \mathbb{R}^{|N|} \mid A = A_0 + \sum_{j \in N} \Delta A_j u_j, \|u\| \leq \Omega \right\}. \quad (3.33)$$

It follows that

$$g(x, \Delta D_j) = \max\{x^T \Delta A_j x, -x^T \Delta A_j x\} = |x^T \Delta A_j x| \quad (3.34)$$

and so

$$\|\mathcal{L}(x, \Delta D_j)\|_g = |x^T \Delta A_j x|. \quad (3.35)$$

By Theorem 3.2, we know that the robust quadratic frontier problem

$$\begin{aligned} \min_{x, \tau} \quad & \tau \\ \text{s.t.} \quad & \omega^T x + \max_{A \in \mathcal{U}} x^T A x \leq \tau, \\ & x \in L(y) = \{x : f(x) \geq y\}, \quad \forall y \geq 0, \end{aligned} \quad (3.36)$$

is equivalent to the following tractable optimization problem

$$\begin{aligned} \min_{x, \tau} \quad & \tau \\ \text{s.t.} \quad & \omega^T x + x^T A_0 x + \Omega \|s\|^* \leq \tau, \\ & x \in L(y) = \{x : f(x) \geq y\}, \quad \forall y \geq 0, \end{aligned} \quad (3.37)$$

where $s_j = |x^T \Delta A_j x|$.

4. Probabilistic Guarantee

In this section, we derive a guarantee on the probability that the robust solution is feasible, when the uncertain coefficients obey some natural probability distributions. An important component of our analysis is the relation among different norms. We denote by $\langle \cdot, \cdot \rangle$ the inner product on a vector space, R^M , or the space of m by m symmetric matrices, $S^{m \times m}$. The inner product induces a norm $\sqrt{\langle x, x \rangle}$. For a vector space, the natural inner product is the Euclidian inner product, $\langle x, y \rangle = x^T y$, and the induced norm is the Euclidian norm $\|x\|_2$.

We analyze the relation of the inner product norm $\sqrt{\langle x, x \rangle}$ with the norm $\|x\|_g$ defined in (3.32). Since $\|x\|_g$ and $\sqrt{\langle x, x \rangle}$ are valid norms in a finite dimensional space, there exist finite $\alpha_1, \alpha_2 > 0$ such that

$$\frac{1}{\alpha_1} \|r\|_g \leq \sqrt{\langle r, r \rangle} \leq \alpha_2 \|r\|_g \quad (4.1)$$

for all r in the relevant space.

Theorem 4.1. *Under the model of uncertainty in (2.1) and given a feasible solution x in (3.1), then*

$$P\left(\omega^T x + \varphi(x, D) \leq \tau\right) \leq P\left(\left\| \sum_{j \in N} r_j u_j \right\|_g > \Omega \|s\|^*\right), \quad (4.2)$$

where

$$r_j = \mathcal{L}(x, \Delta D_j), \quad s_j = \|r_j\|_g, \quad j \in N. \quad (4.3)$$

Proof. From the assumptions, it is easy to see that

$$\begin{aligned} & P(\omega^T x + \varphi(x, D) > \tau) \\ & \leq P(\omega^T x + \varphi(x, D_0) + \varphi(x, \sum_{j \in N} \Delta D_j u_j) > \tau) \quad (\text{from (2.5)}) \\ & \leq P(\varphi(x, \sum_{j \in N} \Delta D_j u_j) > \Omega \|s\|^*) \quad (\text{from (3.14), } s_j = \|\mathcal{L}(x, \Delta D_j)\|_g) \\ & \leq P(\max\{\varphi(x, \sum_{j \in N} \Delta D_j u_j), \varphi(x, -\sum_{j \in N} \Delta D_j u_j)\} > \Omega \|s\|^*) \\ & = P(g(x, \sum_{j \in N} \Delta D_j u_j) > \Omega \|s\|^*) \quad (4.4) \\ & = P(\|\mathcal{L}(x, \sum_{j \in N} \Delta D_j u_j)\|_g > \Omega \|s\|^*) \\ & = P(\|\sum_{j \in N} \mathcal{L}(x, \Delta D_j) u_j\|_g > \Omega \|s\|^*) \quad (\mathcal{L}(x, D) \text{ is linear in } D) \\ & = P\left(\left\|\sum_{j \in N} r_j u_j\right\|_g > \Omega \|s\|^*\right). \end{aligned}$$

This completes the proof. \square

We are naturally led to bound the probability $P(\|\sum_{j \in N} r_j u_j\|_g > \Omega \|s\|^*)$. When we use l_2 -norm in (3.2), that is, $\|s\|^* = \|s\|_2$, we have

$$P\left(\left\|\sum_{j \in N} r_j u_j\right\|_g > \Omega \|s\|^*\right) = P\left(\left\|\sum_{j \in N} r_j u_j\right\|_g > \Omega \sqrt{\sum_{j \in N} \|r_j\|_g^2}\right). \quad (4.5)$$

By the similar method used in Bertsimas and Sim [7], we can get the following result which provides a bound that is independent of the solution x .

Theorem 4.2. *Using the l_2 -norm in (3.2) and under the assumption that u_j are normally and independently distributed with mean zero and variance one, that is, $u \sim \mathcal{N}(0, 1)$, then*

$$P\left(\left\|\sum_{j \in N} r_j u_j\right\|_g > \Omega \sqrt{\sum_{j \in N} \|r_j\|_g^2}\right) \leq \frac{\sqrt{e} \Omega}{\alpha} \exp\left(-\frac{\Omega^2}{2\alpha^2}\right), \quad (4.6)$$

where $\alpha = \alpha_1 \alpha_2$ and α_1, α_2 derived in (4.1) with $\Omega > \alpha$.

Acknowledgments

The authors are grateful to the editor and the referees for their valuable comments and suggestions. This work was supported by the Key Program of NSFC (Grant no. 70831005) and the National Natural Science Foundation of China (11171237, 11126346).

References

- [1] A. Ben-Tal and A. Nemirovski, "Robust convex optimization," *Mathematics of Operations Research*, vol. 23, no. 4, pp. 769–805, 1998.
- [2] A. Ben-Tal and A. Nemirovski, "Robust solutions of uncertain linear programs," *Operations Research Letters*, vol. 25, no. 1, pp. 1–13, 1999.
- [3] A. Ben-Tal and A. Nemirovski, "Robust solutions of linear programming problems contaminated with uncertain data," *Mathematical Programming*, vol. 88, no. 3, pp. 411–424, 2000.
- [4] A. Ben-Tal and A. Nemirovski, "Robust optimization—methodology and applications," *Mathematical Programming*, vol. 92, no. 3, pp. 453–480, 2002.
- [5] A. Ben-Tal, A. Nemirovski, and C. Roos, "Robust solutions of uncertain quadratic and conic quadratic problems," *SIAM Journal on Optimization*, vol. 13, no. 2, pp. 535–560, 2002.
- [6] D. Bertsimas and M. Sim, "The price of robustness," *Operations Research*, vol. 52, no. 1, pp. 35–53, 2004.
- [7] D. Bertsimas and M. Sim, "Tractable approximations to robust conic optimization problems," *Mathematical Programming*, vol. 107, no. 1-2, pp. 5–36, 2006.
- [8] L. El Ghaoui and H. Lebret, "Robust solutions to least-squares problems with uncertain data," *SIAM Journal on Matrix Analysis and Applications*, vol. 18, no. 4, pp. 1035–1064, 1997.
- [9] L. El Ghaoui, F. Oustry, and H. Lebret, "Robust solutions to uncertain semidefinite programs," *SIAM Journal on Optimization*, vol. 9, no. 1, pp. 33–52, 1999.
- [10] L. Wang and N. J. Huang, "Robust solutions to uncertain weighted leastsquares problems," *Mathematical Communications*. In press.
- [11] R. M. Solow, *Book Review*, vol. 36, New York Times, 1987.
- [12] C. Carrado and L. Slifman, "A decomposition of productivity and costs," *American Economic Review*, vol. 89, pp. 328–332, 1999.
- [13] A. N. Perakis and A. Denisis, "A survey of short sea shipping and its prospects in the USA," *Maritime Policy and Management*, vol. 35, no. 6, pp. 591–614, 2008.
- [14] J. Yan, X. Sun, and J. Liu, "Assessing container operator efficiency with heterogeneous and time-varying production frontiers," *Transportation Research B*, vol. 43, no. 1, pp. 172–185, 2009.