

Research Article

On Subscularity of Some 2×2 M -Hyponormal Operator Matrices

Fei Zuo¹ and Junli Shen²

¹ College of Mathematics and Information Science, Henan Normal University, Xinxiang 453007, Henan, China

² School of Mathematical Science, Inner Mongolia University, Hohhot 010021, Inner Mongolia, China

Correspondence should be addressed to Fei Zuo; zuofei2008@sina.com

Received 25 December 2013; Accepted 16 January 2014; Published 25 February 2014

Academic Editor: Yisheng Song

Copyright © 2014 F. Zuo and J. Shen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We provide some conditions for 2×2 operator matrices whose diagonal entries are M -hyponormal operators to be subscalar. As a consequence, we obtain that Weyl type theorem holds for such operator matrices.

1. Introduction and Preliminaries

Let H be a complex separable Hilbert space and let $B(H)$ denote the algebra of all bounded linear operators on H . If $T \in B(H)$, we write $N(T)$, $R(T)$, $\sigma(T)$, and $\sigma_a(T)$ for the null space, the range space, the spectrum, and the approximate point spectrum of T , respectively. An operator T is called Fredholm if $R(T)$ is closed, $\alpha(T) := \dim N(T) < \infty$, and $\beta(T) := \dim N(T^*) < \infty$. The index of a Fredholm operator T is given by $i(T) = \alpha(T) - \beta(T)$. An operator T is called Weyl if it is Fredholm of index zero. The Weyl spectrum of T [1] is defined by $\omega(T) := \{\lambda \in \mathbb{C} : T - \lambda \text{ is not Weyl}\}$.

We consider the sets

$$\begin{aligned} \Phi_+(H) &:= \{T \in B(H) : R(T) \text{ is closed, } \alpha(T) < \infty\}; \quad (1) \\ \Phi_+^-(H) &:= \{T \in B(H) : T \in \Phi_+(H), i(T) \leq 0\} \end{aligned}$$

and define

$$\begin{aligned} \sigma_{ea}(T) &:= \{\lambda \in \mathbb{C} : T - \lambda \notin \Phi_+^-(H)\}; \\ \pi_{00}(T) &:= \{\lambda \in \text{iso } \sigma(T) : 0 < \alpha(T - \lambda) < \infty\}; \quad (2) \end{aligned}$$

$$\pi_{00}^a(T) := \{\lambda \in \text{iso } \sigma_a(T) : 0 < \alpha(T - \lambda) < \infty\},$$

where $\text{iso } \sigma(T)$ denotes the isolated points of $\sigma(T)$.

Following [2], we say that Weyl's theorem holds for T if $\sigma(T) \setminus \omega(T) = \pi_{00}(T)$ and that a -Weyl's theorem holds for T if $\sigma_a(T) \setminus \sigma_{ea}(T) = \pi_{00}^a(T)$.

Let $T \in B(H)$. As an easy extension of normal operators, hyponormal operators have been studied by many mathematicians. Though there are many unsolved interesting problems for hyponormal operators (e.g., the invariant subspace problem), one of recent trends in operator theory is studying natural extensions of hyponormal operators. So we introduce some of these nonhyponormal operators. An operator T is said to be M -hyponormal if there exists a real positive number M such that

$$M^2(T - \lambda)^* (T - \lambda) \geq (T - \lambda) (T - \lambda)^* \quad \forall \lambda \in \mathbb{C}. \quad (3)$$

Evidently,

$$T \text{ is hyponormal} \implies T \text{ is } M\text{-hyponormal}. \quad (4)$$

There is a vast literature concerning M -hyponormal operators (see [3–5], etc.). We also note that an operator T need not be hyponormal even though T and T^* are both M -hyponormal. To see this, consider the operator

$$T = \begin{pmatrix} U & K \\ 0 & U^* \end{pmatrix} : l_2 \oplus l_2 \longrightarrow l_2 \oplus l_2, \quad (5)$$

where U is the unilateral shift on l_2 and $K : l_2 \rightarrow l_2$ is given by $K(x_1, x_2, x_3, \dots) = (2x_1, 0, 0, \dots)$. Then a direct calculation shows that

$$\frac{1}{2} \|(T - z)x\| \leq \|(T - z)^*x\| \leq 2 \|(T - z)x\| \quad (6)$$

for all $z \in \mathbb{C}$ and for all $x \in l_2 \oplus l_2$, which says that T and T^* are both M -hyponormal. But since

$$T^*T = \begin{pmatrix} I & 0 \\ 0 & I + \frac{3}{2}K \end{pmatrix}, \tag{7}$$

while

$$TT^* = \begin{pmatrix} I + \frac{3}{2}K & 0 \\ 0 & I \end{pmatrix}, \tag{8}$$

T is not hyponormal.

Let z be the coordinate in the complex plane \mathbb{C} and let $d\mu(z)$ denote the planar Lebesgue measure. Fix a complex (separable) Hilbert space H and a bounded (connected) open subset U of \mathbb{C} . We will denote by $L^2(U, H)$ the Hilbert space of measurable functions $f : U \rightarrow H$, such that

$$\|f\|_{2,U} = \left(\int_U \|f(z)\|^2 d\mu(z) \right)^{1/2} < \infty. \tag{9}$$

The Bergman space for U is defined by $A^2(U, H) = L^2(U, H) \cap O(U, H)$, where $O(U, H)$ denotes the Fréchet space of H -valued analytic functions on U with respect to uniform topology. Note that $A^2(U, H)$ is a Hilbert space. Let us define now a special Sobolev type space. Let U be again a bounded open subset of \mathbb{C} and let m be a fixed nonnegative integer. The vector valued Sobolev space $W^m(U, H)$ with respect to $\bar{\partial}$ and of order m will be the space of those functions $f \in L^2(U, H)$ whose derivatives $\bar{\partial}f, \dots, \bar{\partial}^m f$ in the sense of distributions still belong to $L^2(U, H)$. Endowed with the norm

$$\|f\|_{W^m}^2 = \sum_{i=0}^m \|\bar{\partial}^i f\|_{2,U}^2 \tag{10}$$

$W^m(U, H)$ becomes a Hilbert space contained continuously in $L^2(U, H)$. A bounded linear operator S on H is called scalar of order m if it possesses a spectral distribution of order m , that is, if there is a continuous unital morphism of topological algebras

$$\Phi : C_0^m(\mathbb{C}) \longrightarrow B(H) \tag{11}$$

such that $\Phi(z) = S$, where z stands for the identity function on \mathbb{C} , and $C_0^m(\mathbb{C})$ stands for the space of compactly supported functions on \mathbb{C} , continuously differentiable of order m , $0 \leq m \leq \infty$. An operator is subscalar if it is similar to the restriction of a scalar operator to an invariant subspace. Let U be a (connected) bounded open subset of \mathbb{C} and let m be a nonnegative integer. The linear operator M_f of multiplication by f on $W^m(U, H)$ is continuous and it has a spectral distribution of order m , defined by the functional calculus

$$\Phi_M : C_0^m(\mathbb{C}) \longrightarrow B(W^m(U, H)), \quad \Phi_M(f) = M_f. \tag{12}$$

Therefore, M_z is a scalar operator of order m .

An operator $T \in B(H)$ is said to have the single-valued extension property (or SVEP) if for every open subset G of \mathbb{C}

and any analytic function $f : G \rightarrow H$ such that $(T-z)f(z) \equiv 0$ on G , we have $f(z) \equiv 0$ on G .

An operator $T \in B(H)$ is said to have Bishop's property (β) if for every open subset G of \mathbb{C} and every sequence $f_n : G \rightarrow H$ of H -valued analytic functions such that $(T-z)f_n(z)$ converges uniformly to 0 in norm on compact subsets of G , $f_n(z)$ converges uniformly to 0 in norm on compact subsets of G . It is well known that

$$\text{Bishop's property } (\beta) \implies \text{SVEP}. \tag{13}$$

In 1984, Putinar showed in [6] that every hyponormal operator is subscalar, and then in 1987, Brown used this result to prove that a hyponormal operator with rich spectrum has a nontrivial invariant subspace (see [7]). There have been a lot of generalizations of such beautiful consequences (see [8–11]). In this paper, we provide some conditions for 2×2 operator matrices whose diagonal entries are M -hyponormal operators to be subscalar. As a consequence, we obtain that Weyl type theorem holds for such operator matrices.

2. Subscalarity

Lemma 1 (see [6, Proposition 2.1]). *For a bounded open disk D in the complex plane \mathbb{C} , there is a constant C_D such that for an arbitrary operator $T \in B(H)$ and $f \in W^2(D, H)$ we have*

$$\|(I - P)f\|_{2,D} \leq C_D \left(\|(T - z)^* \bar{\partial}f\|_{2,D} + \|(T - z)^* \bar{\partial}^2 f\|_{2,D} \right), \tag{14}$$

where P denotes the orthogonal projection of $L^2(D, H)$ onto the Bergman space $A^2(D, H)$.

Corollary 2. *Let D be as in Lemma 1. If $T \in B(H)$ is an M -hyponormal operator, then there exists a constant C_D such that for all $z \in \mathbb{C}$ and $f \in W^2(D, H)$*

$$\begin{aligned} \|(I - P)f\|_{2,D} &\leq MC_D \\ &\times \left(\|(T - z) \bar{\partial}f\|_{2,D} + \|(T - z) \bar{\partial}^2 f\|_{2,D} \right), \end{aligned} \tag{15}$$

where P denotes the orthogonal projection of $L^2(D, H)$ onto the Bergman space $A^2(D, H)$.

Proof. This follows from Lemma 1 and the definition of M -hyponormal operator. \square

Lemma 3. *Let $T \in B(H)$ be an M -hyponormal operator and let D be a bounded disk in \mathbb{C} . If $\{f_n\}$ is a sequence in $W^m(D, H)$ ($m > 2$) such that*

$$\lim_{n \rightarrow \infty} \|(z - T) \bar{\partial}^i f_n\|_{2,D} = 0 \tag{16}$$

for $i = 1, 2, \dots, m$, then $\lim_{n \rightarrow \infty} \|\bar{\partial}^i f_n\|_{2,D_0} = 0$ for $i = 1, 2, \dots, m - 2$, where D_0 is a disk strictly contained in D .

Proof. Since T is an M -hyponormal operator, it follows from Corollary 2 that there exists a constant C_D such that

$$\begin{aligned} \|(I - P)\bar{\partial}^i f_n\|_{2,D} &\leq MC_D \left(\|(T - z)\bar{\partial}^{i+1} f_n\|_{2,D} \right. \\ &\quad \left. + \|(T - z)\bar{\partial}^{i+2} f_n\|_{2,D} \right) \end{aligned} \quad (17)$$

for $i = 0, 1, 2, \dots, m - 2$. From (17), we have

$$\lim_{n \rightarrow \infty} \|(I - P)\bar{\partial}^i f_n\|_{2,D} = 0 \quad (18)$$

for $i = 0, 1, 2, \dots, m - 2$. Hence,

$$\lim_{n \rightarrow \infty} \|(T - z)P\bar{\partial}^i f_n\|_{2,D} = 0 \quad (19)$$

for $i = 1, 2, \dots, m - 2$. Since T has Bishop's property (β) [12], we have

$$\lim_{n \rightarrow \infty} \|P\bar{\partial}^i f_n\|_{2,D_0} = 0 \quad (20)$$

for $i = 1, 2, \dots, m - 2$, where D_0 denotes a disk with $\sigma(T) \not\subseteq \overline{D_0} \not\subseteq D$. From (18) and (20), we get that

$$\lim_{n \rightarrow \infty} \|\bar{\partial}^i f_n\|_{2,D_0} = 0 \quad (21)$$

for $i = 1, 2, \dots, m - 2$. \square

Lemma 4. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, and both T_1 and T_4 are M -hyponormal operators. For any positive integer m and any bounded disk D in \mathbb{C} containing $\sigma(T)$, define the map $V_m : H \oplus H \rightarrow H(D)$ by

$$\begin{aligned} V_m h &= 1 \otimes h \\ &+ \overline{(T - z)W^m(D, H) \oplus W^m(D, H)} (= \widehat{1 \otimes h}), \end{aligned} \quad (22)$$

where

$$H(D) := \frac{W^m(D, H) \oplus W^m(D, H)}{(T - z)W^m(D, H) \oplus W^m(D, H)} \quad (23)$$

and $1 \otimes h$ denotes the constant function sending any $z \in D$ to $h \in H \oplus H$. Then the following statements hold.

- (i) If $T_2^r T_3^s = 0$ for some nonnegative integer r and s , where $T_2^0 = T_3^0 = I$, then V_{4N+2} is one-to-one and has closed range, where $N := \max\{r, s\}$.
- (ii) If $T_1 = T_4$, $T_2 = T_3$, and T_2 is algebraic of order k , then V_{4k+2} is one-to-one and has closed range.
- (iii) If $T_1 + T_4$ is an M -hyponormal operator and $T_1 T_4 = T_2 T_3$, then V_6 is one-to-one and has closed range.

Proof. Let $h_n = h_n^1 \oplus h_n^2 \in H \oplus H$ and

$$f_n = f_n^1 \oplus f_n^2 \in W^m(D, H) \oplus W^m(D, H) \quad (24)$$

be sequences such that

$$\lim_{n \rightarrow \infty} \|(z - T)f_n + 1 \otimes h_n\|_{W^m \oplus W^m} = 0. \quad (25)$$

Then (25) implies that

$$\begin{aligned} \lim_{n \rightarrow \infty} \|(z - T_1)f_n^1 + T_2 f_n^2 + 1 \otimes h_n^1\|_{W^m} &= 0, \\ \lim_{n \rightarrow \infty} \|T_3 f_n^1 + (z - T_4)f_n^2 + 1 \otimes h_n^2\|_{W^m} &= 0. \end{aligned} \quad (26)$$

By the definition of the norm of Sobolev space and (26) we get

$$\begin{aligned} \lim_{n \rightarrow \infty} \|(z - T_1)\bar{\partial}^i f_n^1 + T_2 \bar{\partial}^i f_n^2\|_{2,D} &= 0, \\ \lim_{n \rightarrow \infty} \|T_3 \bar{\partial}^i f_n^1 + (z - T_4)\bar{\partial}^i f_n^2\|_{2,D} &= 0 \end{aligned} \quad (27)$$

for $i = 1, 2, \dots, m$.

- (i) Set $m = 4N + 2$, where $N := \max\{r, s\}$. We may assume that $s \leq r$. Then $m = 4r + 2$.

We prove that for every $j = 0, 1, 2, \dots, s$, the following equations hold

$$\begin{aligned} \lim_{n \rightarrow \infty} \|T_2^{r-j} T_3^{s-j+1} \bar{\partial}^i f_n^1\|_{2,D_j} &= 0, \\ \lim_{n \rightarrow \infty} \|T_2^{r-j} T_3^{s-j} \bar{\partial}^i f_n^2\|_{2,D_j} &= 0 \end{aligned} \quad (28)$$

for $i = 1, 2, \dots, 4(r - j) + 2$, where $\sigma(T) \not\subseteq D_s \not\subseteq \dots \not\subseteq D_2 \not\subseteq D_1 \not\subseteq D$.

To prove (28), we will use the induction on j . Since $T_2^r T_3^s = 0$, then (28) holds when $j = 0$. Suppose that (28) is true for some $j < s$. From (27) and the inductions hypothesis, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \|(T_1 - z)T_2^{r-j-1} T_3^{s-j} \bar{\partial}^i f_n^1\|_{2,D_j} &= 0, \\ \lim_{n \rightarrow \infty} \|T_2^{r-j-1} T_3^{s-j} \bar{\partial}^i f_n^1 + (T_4 - z)T_2^{r-j-1} T_3^{s-j-1} \bar{\partial}^i f_n^2\|_{2,D_j} &= 0 \end{aligned} \quad (29)$$

for $i = 1, 2, \dots, 4(r - j) + 2$. Since T_1 is an M -hyponormal operator, by Lemma 3 we have

$$\lim_{n \rightarrow \infty} \|T_2^{r-j-1} T_3^{s-j} \bar{\partial}^i f_n^1\|_{2,D'_j} = 0 \quad (30)$$

for $i = 1, 2, \dots, 4(r - j)$, where $\sigma(T) \not\subseteq D'_j \not\subseteq D_j$. From (30) and the second equation of (27),

$$\lim_{n \rightarrow \infty} \|(T_4 - z)T_2^{r-j-1} T_3^{s-j-1} \bar{\partial}^i f_n^2\|_{2,D'_j} = 0 \quad (31)$$

for $i = 1, 2, \dots, 4(r - j)$. Since T_4 is an M -hyponormal operator, by Lemma 3 we derive

$$\lim_{n \rightarrow \infty} \|T_2^{r-j-1} T_3^{s-j-1} \bar{\partial}^i f_n^2\|_{2,D_{j+1}} = 0 \quad (32)$$

for $i = 1, 2, \dots, 4(r - j - 1) + 2$, where $\sigma(T) \not\subseteq D_{j+1} \not\subseteq D'_j$. Therefore, the proof of (28) is completed. Let $j = s$ in (28); we have

$$\lim_{n \rightarrow \infty} \|T_2^{r-s} T_3 \bar{\partial}^i f_n^1\|_{2, D_s} = 0, \quad (33)$$

$$\lim_{n \rightarrow \infty} \|T_2^{r-s} \bar{\partial}^i f_n^2\|_{2, D_s} = 0$$

for $i = 1, 2, \dots, 4(r - s) + 2$. From (27) and (33), it follows that

$$\lim_{n \rightarrow \infty} \|(T_1 - z) T_2^{r-s-1} \bar{\partial}^i f_n^1\|_{2, D_s} = 0 \quad (34)$$

for $i = 1, 2, \dots, 4(r - s) + 2$. Since T_1 is an M -hyponormal operator, from Lemma 3 we obtain that

$$\lim_{n \rightarrow \infty} \|T_2^{r-s-1} \bar{\partial}^i f_n^1\|_{2, D_s} = 0 \quad (35)$$

for $i = 1, 2, \dots, 4(r - s)$, where $\sigma(T) \not\subseteq D'_s \not\subseteq D_s$, and hence

$$\lim_{n \rightarrow \infty} \|(T_4 - z) T_2^{r-s-1} \bar{\partial}^i f_n^2\|_{2, D_s} = 0 \quad (36)$$

for $i = 1, 2, \dots, 4(r - s)$. Since T_4 is an M -hyponormal operator, Lemma 3 implies that

$$\lim_{n \rightarrow \infty} \|T_2^{r-s-1} \bar{\partial}^i f_n^2\|_{2, D_{s+1}} = 0 \quad (37)$$

for $i = 1, 2, \dots, 4(r - s - 1) + 2$, where $\sigma(T) \not\subseteq D_{s+1} \not\subseteq D'_s$. By repeating the process from (33) to (37), it holds for all $j = 0, 1, 2, \dots, r - s$ that

$$\lim_{n \rightarrow \infty} \|T_2^{r-s-j} \bar{\partial}^i f_n^2\|_{2, D_{s+j}} = 0 \quad (38)$$

for $i = 1, 2, \dots, 4(r - s - j) + 2$, where $\sigma(T) \not\subseteq D_{r+1} \not\subseteq D_r \not\subseteq D_{r-1} \not\subseteq \dots \not\subseteq D_{s+1} \not\subseteq D_s$. In particular, let $r = s + j$,

$$\lim_{n \rightarrow \infty} \|\bar{\partial}^i f_n^2\|_{2, D_r} = 0 \quad (39)$$

for $i = 1, 2$. Hence, from the first equation of (27), we have

$$\lim_{n \rightarrow \infty} \|(T_1 - z) \bar{\partial}^i f_n^1\|_{2, D_r} = 0 \quad (40)$$

for $i = 1, 2$. Applying Corollary 2, we have

$$\lim_{n \rightarrow \infty} \|(I - P) f_n^1\|_{2, D'_r} = 0, \quad (41)$$

where P denotes the orthogonal projection of $L^2(D'_r, H)$ onto $A^2(D'_r, H)$ and $\sigma(T) \not\subseteq D'_r \not\subseteq D_r$. By combining (26) with (39) and (41), we obtain that

$$\lim_{n \rightarrow \infty} \|(z - T) P f_n + 1 \otimes h_n\|_{2, D'_r} = 0, \quad (42)$$

where $P f_n := \begin{pmatrix} P f_n^1 \\ P f_n^2 \end{pmatrix}$.

Let Γ be a curve in D'_r surrounding $\sigma(T)$. Then for $z \in \Gamma$

$$\lim_{n \rightarrow \infty} \|P f_n(z) + (z - T)^{-1} (1 \otimes h_n)(z)\| = 0 \quad (43)$$

uniformly. Hence, by the Riesz functional calculus,

$$\lim_{n \rightarrow \infty} \left\| \frac{1}{2\pi i} \int_{\Gamma} P f_n(z) dz + h_n \right\| = 0. \quad (44)$$

But $(1/2\pi i) \int_{\Gamma} P f_n(z) dz = 0$ by Cauchy's theorem. Hence, $\lim_{n \rightarrow \infty} h_n = 0$, and so V_{4r+2} is one-to-one and has closed range.

(ii) Set $m = 4k + 2$. By the hypothesis and (27), we have

$$\lim_{n \rightarrow \infty} \|(T_1 - z) \bar{\partial}^i f_n^1 + T_2 \bar{\partial}^i f_n^2\|_{2, D} = 0, \quad (45)$$

$$\lim_{n \rightarrow \infty} \|T_2 \bar{\partial}^i f_n^1 + (T_1 - z) \bar{\partial}^i f_n^2\|_{2, D} = 0$$

for $i = 1, 2, \dots, 4k + 2$. Since T_2 is algebraic with order k , there exists a nonconstant polynomial $p(z) = (z - z_1)(z - z_2) \cdots (z - z_k)$ such that $p(T_2) = 0$. Set $q_j(z) = (z - z_{j+1}) \cdots (z - z_k)$ for $j = 0, 1, 2, \dots, k - 1$ and $q_k(z) = 1$.

Claim. For every $j = 0, 1, 2, \dots, k$, the following equations hold:

$$\lim_{n \rightarrow \infty} \|q_j(T_2) \bar{\partial}^i f_n^1\|_{2, D_j} = 0, \quad (46)$$

$$\lim_{n \rightarrow \infty} \|q_j(T_2) \bar{\partial}^i f_n^2\|_{2, D_j} = 0$$

for $i = 1, 2, \dots, 4(k - j) + 2$, where $\sigma(T) \not\subseteq D_k \not\subseteq \dots \not\subseteq D_2 \not\subseteq D_1 \not\subseteq D$.

To prove the claim, we use the induction on j . Since $q_0(T_2) = p(T_2) = 0$, then when $j = 0$ the claim holds. Suppose that the claim is true for some $j = r$, where $0 \leq r < k$. Multiplying (45) by $q_{r+1}(T_2)$, we obtain

$$\lim_{n \rightarrow \infty} \|(T_1 - z) q_{r+1}(T_2) \bar{\partial}^i f_n^1 + z_{r+1} q_{r+1}(T_2) \bar{\partial}^i f_n^2\|_{2, D_r} = 0,$$

$$\lim_{n \rightarrow \infty} \|z_{r+1} q_{r+1}(T_2) \bar{\partial}^i f_n^1 + (T_1 - z) q_{r+1}(T_2) \bar{\partial}^i f_n^2\|_{2, D_r} = 0 \quad (47)$$

for $i = 1, 2, \dots, 4(k - r) + 2$. From (47), we derive

$$\lim_{n \rightarrow \infty} \|(T_1 - (z - z_{r+1})) q_{r+1}(T_2) (\bar{\partial}^i f_n^1 + \bar{\partial}^i f_n^2)\|_{2, D_r} = 0 \quad (48)$$

for $i = 1, 2, \dots, 4(k - r) + 2$. Since T_1 is an M -hyponormal operator, from (48) and Lemma 3 we obtain

$$\lim_{n \rightarrow \infty} \|q_{r+1}(T_2) (\bar{\partial}^i f_n^1 + \bar{\partial}^i f_n^2)\|_{2, D'_r} = 0 \quad (49)$$

for $i = 1, 2, \dots, 4(k - r)$, where $\sigma(T) \not\subseteq D'_r \not\subseteq D_r$. Combining (49) with the first equation of (47), we have

$$\lim_{n \rightarrow \infty} \|(T_1 - (z + z_{r+1})) q_{r+1}(T_2) \bar{\partial}^i f_n^1\|_{2, D'_r} = 0 \quad (50)$$

for $i = 1, 2, \dots, 4(k - r)$. Since T_1 is an M -hyponormal operator, we obtain from Lemma 3 and (50) that

$$\lim_{n \rightarrow \infty} \|q_{r+1}(T_2) \bar{\partial}^i f_n^1\|_{2, D_{r+1}} = 0 \quad (51)$$

for $i = 1, 2, \dots, 4(k - r - 1) + 2$, where $\sigma(T) \not\subseteq D_{r+1} \not\subseteq D'_r$. From (49) and (51), we obtain

$$\lim_{n \rightarrow \infty} \left\| q_{r+1}(T_2) \bar{\partial}^i f_n^2 \right\|_{2, D_{r+1}} = 0 \quad (52)$$

for $i = 1, 2, \dots, 4(k - r - 1) + 2$. Therefore, the proof of the claim is completed.

From the claim with $j = k$, we have

$$\lim_{n \rightarrow \infty} \left\| \bar{\partial}^i f_n^1 \right\|_{2, D_k} = \lim_{n \rightarrow \infty} \left\| \bar{\partial}^i f_n^2 \right\|_{2, D_k} = 0 \quad (53)$$

for $i = 1, 2$. From Lemma 1 we derive that

$$\lim_{n \rightarrow \infty} \left\| (I - P) f_n^1 \right\|_{2, D_k} = \lim_{n \rightarrow \infty} \left\| (I - P) f_n^2 \right\|_{2, D_k} = 0, \quad (54)$$

where P denotes the orthogonal projection of $L^2(D_k, H)$ onto $A^2(D_k, H)$. By applying the proof of (i), we obtain that V_{4k+2} is one-to-one and has closed range.

(iii) Set $m = 6$. By (27), we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \left\| (T_1 T_3 - z T_3) \bar{\partial}^i f_n^1 + T_2 T_3 \bar{\partial}^i f_n^2 \right\|_{2, D} &= 0, \\ \lim_{n \rightarrow \infty} \left\| T_1 T_3 \bar{\partial}^i f_n^1 + (T_1 T_4 - z T_1) \bar{\partial}^i f_n^2 \right\|_{2, D} &= 0 \end{aligned} \quad (55)$$

for $i = 1, 2, \dots, 6$. Since $T_1 T_4 = T_2 T_3$, (55) implies that

$$\lim_{n \rightarrow \infty} \left\| z \left(T_1 \bar{\partial}^i f_n^2 - T_3 \bar{\partial}^i f_n^1 \right) \right\|_{2, D} = 0 \quad (56)$$

for $i = 1, 2, \dots, 6$. Since the zero operator is hyponormal operator, it follows from Lemma 3 and (56) that

$$\lim_{n \rightarrow \infty} \left\| T_1 \bar{\partial}^i f_n^2 - T_3 \bar{\partial}^i f_n^1 \right\|_{2, D_1} = 0 \quad (57)$$

for $i = 1, 2, \dots, 4$, where $\sigma(T) \not\subseteq D_1 \not\subseteq D$. Using (57) and the second equation of (27), we obtain

$$\lim_{n \rightarrow \infty} \left\| (T_1 + T_4 - z) \bar{\partial}^i f_n^2 \right\|_{2, D_1} = 0 \quad (58)$$

for $i = 1, 2, 3, 4$. Since $T_1 + T_4$ is an M -hyponormal operator, from Lemma 3 we have

$$\lim_{n \rightarrow \infty} \left\| \bar{\partial}^i f_n^2 \right\|_{2, D_2} = 0 \quad (59)$$

for $i = 1, 2$, where $\sigma(T) \not\subseteq D_2 \not\subseteq D_1$. From (59) and the first equation of (27) we obtain that

$$\lim_{n \rightarrow \infty} \left\| (T_1 - z) \bar{\partial}^i f_n^1 \right\|_{2, D_2} = 0 \quad (60)$$

for $i = 1, 2$. Since T_1 is an M -hyponormal operator, from Corollary 2 we get that

$$\lim_{n \rightarrow \infty} \left\| (I - P) f_n^1 \right\|_{2, D_3} = 0, \quad (61)$$

where P denotes the orthogonal projection of $L^2(D_3, H)$ onto $A^2(D_3, H)$ and $\sigma(T) \not\subseteq D_3 \not\subseteq D_2$. By applying the similar way of (i), we obtain that V_6 is one-to-one and has closed range. \square

Theorem 5. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, and both T_1 and T_4 are M -hyponormal operators. If $\{T_i\}_{i=1}^4$ satisfy one of the conditions in Lemma 4, then T is a subscalar operator of order m , where m is the appropriately chosen integer as in Lemma 4.

Proof. Let D be a bounded disk in \mathbb{C} containing $\sigma(T)$ and consider the quotient space

$$H(D) := \frac{W^m(D, H) \oplus \overline{W^m(D, H)}}{(T - z)W^m(D, H) \oplus W^m(D, H)} \quad (62)$$

endowed with the Hilbert space norm, where $m = 4N + 2$, $N := \max\{r, s\}$ for (i), $m = 4k + 2$ for (ii), and $m = 6$ for (iii) in Lemma 4. The class of a vector f or an operator S on $H(D)$ will be denoted by \tilde{f}, \tilde{S} , respectively. Let M be the operator of multiplication by z on $W^m(D, H) \oplus W^m(D, H)$. Then M is a scalar operator of order m and has a spectral distribution Φ . Since $R(T - z)$ is invariant under M , \tilde{M} can be well defined. Moreover, consider the spectral distribution $\Phi : C_0^m(\mathbb{C}) \rightarrow B(W^m(D, H) \oplus W^m(D, H))$ defined by the following relation: for $\varphi \in C_0^m(\mathbb{C})$ and $f \in W^m(D, H) \oplus W^m(D, H)$, $\Phi(\varphi)f = \varphi f$. Then the spectral distribution Φ of M commutes with $T - z$, and so \tilde{M} is still a scalar operator of order m with $\tilde{\Phi}$ as a spectral distribution. As in Lemma 4, if we define the map $V_m : H_1 \oplus H_2 \rightarrow H(D)$ by

$$\begin{aligned} V_m h &= 1 \otimes h \\ &+ \overline{(T - z)W^m(D, H) \oplus W^m(D, H)} (1 \otimes h), \end{aligned} \quad (63)$$

then V_m is one-to-one and has closed range. Since

$$V_m T h = \overline{1 \otimes T h} = \overline{z \otimes h} = \tilde{M} (1 \otimes h) = \tilde{M} V_m h \quad (64)$$

for all $h \in H \oplus H$, $V_m T = \tilde{M} V_m$. In particular, $R(V_m)$ is invariant under \tilde{M} and $R(V_m)$ is closed; it is a closed invariant subspace of the scalar operator \tilde{M} . Since T is similar to the restriction $\tilde{M}|_{R(V_m)}$ and \tilde{M} is scalar of order m , T is a subscalar operator of order m . \square

Corollary 6. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, both T_1 and T_4 are M -hyponormal operators, and $\{T_i\}_{i=1}^4$ satisfy one of the conditions in Lemma 4. Then T has property (β) and the single-valued extension property.

Proof. From section one, we need only to prove that T has property (β) . Since property (β) is transmitted from an operator to its restrictions to closed invariant subspaces, we are reduced by Theorem 5 to the case of a scalar operator. Since every scalar operator has property (β) (see [6]), T has property (β) . \square

Define the quasi-nilpotent part of $\lambda I - T$

$$H_0(\lambda I - T) := \left\{ x \in H : \lim_{n \rightarrow \infty} \|(\lambda I - T)^n x\|^{1/n} = 0 \right\}. \quad (65)$$

Definition 7. An operator $T \in B(H)$ is said to belong to the class $H(p)$ if there exists a natural number $p := p(\lambda)$ such that

$$H_0(\lambda I - T) = N(\lambda I - T)^p \quad \forall \lambda \in \mathbb{C}. \quad (66)$$

Theorem 8 (see [13]). Every subscalar operator $T \in B(H)$ is $H(p)$.

Definition 9. An operator $T \in B(H)$ is said to be polaroid if every $\lambda \in \text{iso } \sigma(T)$ is a pole of the resolvent of T .

Note that

$$T \text{ is polaroid} \iff T^* \text{ is polaroid}. \quad (67)$$

The condition of being polaroid may be characterized by means of the quasi-nilpotent part.

Theorem 10 (see [14]). An operator $T \in B(H)$ is polaroid if and only if there exists a natural number $p := p(\lambda)$ such that

$$H_0(\lambda I - T) = N(\lambda I - T)^p \quad \forall \lambda \in \text{iso } \sigma(T). \quad (68)$$

Corollary 11. Every $H(p)$ operator is polaroid.

Since a subscalar operator is $H(p)$, we have the following.

Corollary 12. Every subscalar operator is polaroid.

Corollary 13. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, both T_1 and T_4 are M -hyponormal operators, and $\{T_i\}_{i=1}^4$ satisfy one of the conditions in Lemma 4. Then T is polaroid.

If $T \in B(H)$ has SVEP, then T and T^* satisfy Browder's theorem. A sufficient condition for an operator T satisfying Browder's theorem to satisfy Weyl's theorem is that T is polaroid. Then we have the following result.

Corollary 14. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, both T_1 and T_4 are M -hyponormal operators, and $\{T_i\}_{i=1}^4$ satisfy one of the conditions in Lemma 4. Then Weyl's theorem holds for T and T^* .

Observe that if $T \in B(H)$ has SVEP, then $\sigma(T) = \overline{\sigma_a(T^*)}$. Hence, if T has SVEP and is polaroid, then T^* satisfies a -Weyl's theorem [15, Theorem 3.10].

Corollary 15. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, both T_1 and T_4 are M -hyponormal operators, and $\{T_i\}_{i=1}^4$ satisfy one of the conditions in Lemma 4. Then a -Weyl's theorem holds for T^* .

Proof. Since T is polaroid and has SVEP, then a -Weyl's theorem holds for T^* . \square

In the following, f is an analytic function on $\sigma(T)$ and f is not constant on each connected component of the open set U containing $\sigma(T)$.

Corollary 16. Let $T = \begin{pmatrix} T_1 & T_2 \\ T_3 & T_4 \end{pmatrix} \in B(H \oplus H)$, where T_i are mutually commuting, both T_1 and T_4 are M -hyponormal operators, and $\{T_i\}_{i=1}^4$ satisfy one of the conditions in Lemma 4. Then the following assertions hold:

- (i) Weyl's theorem holds for $f(T)$;
- (ii) a -Weyl's theorem holds for $f(T^*)$.

Proof. (i) Since T is polaroid and has SVEP, then $f(T)$ satisfies Weyl's theorem by [15, Theorem 3.14].

(ii) Since T is polaroid and has SVEP, then $f(T^*)$ satisfies a -Weyl's theorem by [15, Theorem 3.12]. \square

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

The authors would like to express their cordial gratitude to the referee for his valuable advice and suggestion. This work was partially supported by the National Natural Science Foundation of China (11201126), the Natural Science Foundation of the Department of Education, Henan Province (no. 14B110008), and the Youth Science Foundation of Henan Normal University (no. 2013QK01).

References

- [1] R. Harte, *Invertibility and Singularity for Bounded Linear Operators*, Marcel Dekker, New York, NY, USA, 1988.
- [2] R. Harte and W. Y. Lee, "Another note on Weyl's theorem," *Transactions of the American Mathematical Society*, vol. 349, no. 5, pp. 2115–2124, 1997.
- [3] X. Cao, M. Guo, and B. Meng, "Weyl type theorems for p -hyponormal and M -hyponormal operators," *Studia Mathematica*, vol. 163, no. 2, pp. 177–188, 2004.
- [4] R. L. Moore, D. D. Rogers, and T. T. Trent, "A note on intertwining M -hyponormal operators," *Proceedings of the American Mathematical Society*, vol. 83, no. 3, pp. 514–516, 1981.
- [5] A. Uchiyama and T. Yoshino, "Weyl's theorem for p -hyponormal or M -hyponormal operators," *Glasgow Mathematical Journal*, vol. 43, no. 3, pp. 375–381, 2001.
- [6] M. Putinar, "Hyponormal operators are subscalar," *Journal of Operator Theory*, vol. 12, no. 2, pp. 385–395, 1984.
- [7] S. W. Brown, "Hyponormal operators with thick spectra have invariant subspaces," *Annals of Mathematics*, vol. 125, no. 1, pp. 93–103, 1987.
- [8] S. Jung, Y. Kim, and E. Ko, "On subscalarity of some 2×2 class A operator matrices," *Linear Algebra and Its Applications*, vol. 438, no. 3, pp. 1322–1338, 2013.
- [9] S. Jung, E. Ko, and M.-J. Lee, "On class A operators," *Studia Mathematica*, vol. 198, no. 3, pp. 249–260, 2010.
- [10] S. Jung, E. Ko, and M.-J. Lee, "Subscalarity of (p, k) -quasi-hyponormal operators," *Journal of Mathematical Analysis and Applications*, vol. 380, no. 1, pp. 76–86, 2011.
- [11] E. Ko, " k th roots of p -hyponormal operators are subscalar operators of order $4k$," *Integral Equations and Operator Theory*, vol. 59, no. 2, pp. 173–187, 2007.

- [12] K. B. Laursen and M. M. Neumann, *An Introduction to Local Spectral Theory*, Clarendon Press, Oxford, UK, 2000.
- [13] M. Oudghiri, “Weyl’s and Browder’s theorems for operators satisfying the SVEP,” *Studia Mathematica*, vol. 163, no. 1, pp. 85–101, 2004.
- [14] P. Aiena, M. Chō, and M. González, “Polaroid type operators under quasi-affinities,” *Journal of Mathematical Analysis and Applications*, vol. 371, no. 2, pp. 485–495, 2010.
- [15] P. Aiena, E. Aponte, and E. Balzan, “Weyl type theorems for left and right polaroid operators,” *Integral Equations and Operator Theory*, vol. 66, no. 1, pp. 1–20, 2010.