

Solutions of Perturbed Fractional Schrödinger–Poisson System with Critical Nonlinearity in \mathbb{R}^3

Lvzhou Zheng

Abstract. In this paper, we study the existence of semiclassical solutions of the fractional Schrödinger–Poisson system with critical exponent in \mathbb{R}^3 . Under suitable assumptions, we show that the problem has at least one nontrivial solution provided that $\epsilon \leq \epsilon_\delta$ where ϵ_δ is sufficiently small positive number.

1. Introduction and main result

Let us consider the fractional Schrödinger–Poisson system of the form

$$(1.1) \quad \begin{cases} \epsilon^{2s}(-\Delta)^s u + V(x)u + \Phi(x)u = Q(x)|u|^{2^*(s)-2}u + f(x, u), & x \in \mathbb{R}^3, \\ (-\Delta)^t \Phi = u^2, & x \in \mathbb{R}^3, \end{cases}$$

where $\epsilon > 0$, $2^*(s) = 6/(3 - 2s)$, $0 < t < 1$, $3/4 < s < 1$. $V(x)$, $Q(x)$ and $f(x, u)$ satisfy the following assumptions:

(V₁) $V \in C(\mathbb{R}^3, \mathbb{R})$ and there is $b > 0$ such that the set $V^b := \{x \in \mathbb{R}^3 : V(x) < b\}$ has finite Lebesgue measure.

(V₂) $0 = V(0) = \min_{x \in \mathbb{R}^3} V(x) \leq V(x) < M$.

(Q) $Q \in C(\mathbb{R}^3, \mathbb{R})$, $0 < Q_1 := \inf_{x \in \mathbb{R}^3} Q(x) \leq Q_2 := \sup_{x \in \mathbb{R}^3} Q(x) < \infty$.

(f₁) $f \in C(\mathbb{R}^3 \times \mathbb{R}, \mathbb{R})$, $f(x, u) = o(|u|)$ uniformly in x as $u \rightarrow 0$.

(f₂) There are $c_0 > 0$ and $q \in (2, 2^*(s))$ such that

$$|f(x, u)| \leq c_0(1 + |u|^{q-1}) \quad \text{for all } (x, u).$$

(f₃) There are $a_0 > 0$ and $4 < \mu, l < 2^*(s)$ such that $F(x, u) \geq a_0|u|^l$ and $\mu F(x, u) \leq f(x, u)u$ for all (x, u) , where $F(x, u) = \int_0^u f(x, s) ds$.

Received August 20, 2020; Accepted December 7, 2020.

Communicated by François Hamel.

2020 *Mathematics Subject Classification.* 35J10, 35B99, 35J60.

Key words and phrases. fractional Schrödinger–Poisson system, critical nonlinearity, variational methods.

The fractional Schrödinger–Poisson system appears in an interesting physical context. In (1.1), the first equation is a nonlinear fractional Schrödinger equation in which the potential Φ satisfies a nonlinear fractional Poisson equation. For this reason, system (1.1) is called the fractional Schrödinger–Poisson system. Recently, a great attention has been focused on the study of problems involving the fractional Laplacian, since this operator arises in several areas such as probability, physics and finance. Also, the fractional Laplacian can be understood as the infinitesimal generator of a stable Lévy process.

When $s = t = 1$, the system reduces to the classical Schrödinger–Poisson system. The Schrödinger–Poisson system has been investigated in the literature, see [1, 4, 6, 10, 13–16, 32] and the references therein for existence results. In recent years, there have been a lot of works dealing with the fractional Schrödinger–Poisson equations. Zhang, Do Ó and Squassina in [34] obtained the existence of radial ground state solution to the following fractional Schrödinger–Poisson system with a general subcritical or critical nonlinearity

$$\begin{cases} (-\Delta)^s u + \lambda \Phi(y)u = g(u), & y \in \mathbb{R}^3, \\ (-\Delta)^t \Phi = \lambda u^2, & y \in \mathbb{R}^3. \end{cases}$$

Without assuming the Ambrosetti–Rabinowitz condition by using the fountain theorem, Zhang in [35] considered the existence of infinitely many large energy solutions to the following system

$$(1.2) \quad \begin{cases} (-\Delta)^s u + V(y)u + \Phi(y)u = f(y, u), & y \in \mathbb{R}^3, \\ (-\Delta)^t \Phi = \lambda u^2, & y \in \mathbb{R}^3. \end{cases}$$

Using the Lyapunov–Schmidt reduction method, Liu [17] obtained the existence of multi-bump solutions for the following fractional Schrödinger–Poisson system

$$\begin{cases} (-\Delta)^s u + u + \epsilon V(y)\Phi(y)u = |u|^{p-1}u, & y \in \mathbb{R}^3, \\ (-\Delta)^t \Phi = V(y)u^2, & y \in \mathbb{R}^3. \end{cases}$$

In [25], Shen and Yao obtained the existence of positive ground state solutions and concentration results for (1.2) via some new analytical skills and Nehari–Pohožaev identity. In [24], Shen concerned with the nontrivial solutions for fractional Schrödinger–Poisson system with the Bessel operator. Under certain assumptions on the nonlinearity, a nontrivial nonnegative solution is obtained by perturbation method. In [8], using the Non-Nehari manifold approach, the authors established the existence of the Nehari-type ground state solutions for fractional Schrödinger–Poisson system (1.2). By introducing some new tricks, in [9], Chen and Tang proved that the single problem admits a ground state solution of Pohožaev type and a least energy solution. For more related results, one can refer to [3, 5, 7, 18–22, 27, 28, 31, 33] and the references therein.

The purpose of this paper is to study the problem (1.1) and obtain the existence and multiplicity of semiclassical solutions. To the best of our knowledge, we cannot find any result in the literature that can be directly applied to obtain the existence and multiplicity of solutions to problem (1.1). In this work, we employ the methods of [2, 12, 23, 29, 30] to obtain our main result. Especially, in [12], Ding and Lin considered the single Schrödinger equation. Here, although the idea was used in [12], the adaptation to the procedure to our problem is not trivial at all, since the appearance of non-local term, we must consider our problem for suitable space and so we need more delicate estimates.

The main result of this paper is stated below.

Theorem 1.1. *Let (V_1) , (V_2) , (Q) , (f_1) , (f_2) and (f_3) hold, $0 < t < 1$, $3/4 < s < 1$. Then for any $\delta > 0$, there is $\varepsilon_\delta > 0$ such that if $\epsilon \leq \varepsilon_\delta$, problem (1.1) has at least one nontrivial solution u_ϵ satisfying*

- (i) $\left(\frac{1}{2} - \frac{1}{\mu}\right) \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_\epsilon|^2 + V(x)|u_\epsilon|^2) \leq \delta \lambda^{1-3/(2s)}$ and
- (ii) $\left(\frac{1}{4} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} Q(x)|u_\epsilon|^{2^*(s)} + \frac{\mu-4}{4} \int_{\mathbb{R}^3} F(x, u_\epsilon) \leq \delta \lambda^{-3/(2s)}.$

The paper is organized as below. In Section 2, we present some preliminary results. Section 3 is devoted to the behavior of Palais–Smale sequences. In Sections 4 and 5, we accomplish the proof of the main result.

2. Preliminaries

In this section, we recall some preliminary results which will be useful along the paper. First, we will give some useful facts of the fractional order Sobolev spaces. Then, we recall some useful facts of the homogeneous Sobolev space and some properties of the Poisson potential Φ .

For any $s \in (0, 1)$, the fractional Sobolev space $H^s(\mathbb{R}^3)$ is defined by

$$\begin{aligned} H^s(\mathbb{R}^3) &= \left\{ u \in L^2(\mathbb{R}^3) : \frac{|u(x) - u(y)|}{|x - y|^{3/2+s}} \in L^2(\mathbb{R}^3 \times \mathbb{R}^3) \right\} \\ &= \left\{ u \in L^2(\mathbb{R}^3) : \int_{\mathbb{R}^3} (1 + |\xi|^{2s}) |\widehat{u}(\xi)|^2 d\xi < \infty \right\} \end{aligned}$$

endowed with the natural norm

$$\|u\|_{H^s(\mathbb{R}^3)} = \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|u(x) - u(y)|^2}{|x - y|^{3+2s}} dx dy + \int_{\mathbb{R}^3} |u|^2 dx \right)^{1/2},$$

which is induced by the inner product

$$\begin{aligned} \langle u, v \rangle_{H^s(\mathbb{R}^3)} &= \langle u, v \rangle_s + \langle u, v \rangle_{L^2(\mathbb{R}^3)} \\ &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{3+2s}} dx dy + \int_{\mathbb{R}^3} u(x)v(x) dx. \end{aligned}$$

Here the term

$$[u]_{H^s(\mathbb{R}^3)} := \left(\int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|u(x) - u(y)|^2}{|x - y|^{3+2s}} dx dy \right)^{1/2}$$

is the so-called Gagliardo (semi-)norm of u and \widehat{u} means the Fourier transform of u . The following identity yields the relation between the fractional operator $(-\Delta)^s$ and the fractional Sobolev space $H^s(\mathbb{R}^3)$,

$$[u]_{H^s(\mathbb{R}^3)} = C \left(\int_{\mathbb{R}^3} |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi \right)^{1/2} = C \|(-\Delta)^{s/2} u\|_{L^2(\mathbb{R}^3)}$$

for a suitable positive constant C depending only on s .

For the reader's convenience, we review the main embedding result for this class of fractional Sobolev spaces.

Lemma 2.1. [11] *$H^s(\mathbb{R}^3)$ is continuously embedded into $L^q(\mathbb{R}^3)$ for $q \in [2, 6/(3 - 2s)]$, and the embedding is compact whenever $q \in [2, 6/(3 - 2s))$ on bounded domains.*

We introduce the space

$$E = \left\{ u \in H^s(\mathbb{R}^3) : \int_{\mathbb{R}^3} V(x)u^2 < \infty \right\},$$

which is a Hilbert space under the scalar product

$$\langle v_1, v_2 \rangle_E = \int_{\mathbb{R}^3} (-\Delta)^{s/2} v_1 (-\Delta)^{s/2} v_2 + \int_{\mathbb{R}^3} V(x)v_1 v_2.$$

The norm induced by the product $\langle \cdot, \cdot \rangle_E$ is

$$\|u\|_E = \sqrt{\langle u, u \rangle_E}, \quad u \in E.$$

The homogeneous Sobolev space $D^{t,2}(\mathbb{R}^3)$ is defined by

$$D^{t,2}(\mathbb{R}^3) = \left\{ u \in L^{2^*(t)}(\mathbb{R}^3) : \int_{\mathbb{R}^3} |\xi|^{2s} |\widehat{u}(\xi)|^2 d\xi < \infty \right\},$$

which is the completion of $C_0^\infty(\mathbb{R}^3)$ under the norm

$$\|u\|_{D^{t,2}} = \left(\int_{\mathbb{R}^3} |\xi|^{2t} |\widehat{u}(\xi)|^2 d\xi \right)^{1/2} = \|(-\Delta)^{t/2} u\|_{L^2(\mathbb{R}^3)}$$

and the inner product

$$(u, v)_{D^{t,2}} = \int_{\mathbb{R}^3} (-\Delta)^{t/2} u (-\Delta)^{t/2} v \, dx, \quad u, v \in D^{t,2}(\mathbb{R}^3).$$

By the assumption (V₁), we know that the embedding $E \hookrightarrow H^s(\mathbb{R}^3)$ is continuous. Note that the norm $\|\cdot\|_E$ is equivalent to the one $\|\cdot\|_\lambda$ defined by

$$\|u\|_\lambda = \left(\int_{\mathbb{R}^3} |(-\Delta)^{s/2} u|^2 + \lambda \int_{\mathbb{R}^3} V(x) u^2 \right)^{1/2}$$

for each $\lambda > 0$. It is obvious that for each $p \in [2, 2^*(s)]$, there is $c_p > 0$ such that if $\lambda \geq 1$,

$$(2.1) \quad |u|_p \leq c_p \|u\|_E \leq c_p \|u\|_\lambda.$$

To prove our results, the following Sobolev embedding result is necessary.

Lemma 2.2. [11] *For any $t \in (0, 1)$, $D^{t,2}(\mathbb{R}^3)$ is continuously embedded into $L^{2^*(t)}(\mathbb{R}^3)$, i.e., there exists $S_t > 0$ such that*

$$\left(\int_{\mathbb{R}^3} |u|^{2^*(t)} \, dx \right)^{2/2^*(t)} \leq S_t \int_{\mathbb{R}^3} |(-\Delta)^{t/2} u|^2 \, dx, \quad u \in D^{t,2}(\mathbb{R}^3).$$

By Lemma 2.1, if $0 < t < 1$, $3/4 < s < 1$, $H^s(\mathbb{R}^3) \hookrightarrow L^{12/(3+2t)}(\mathbb{R}^3)$. Then, for $u \in H^s(\mathbb{R}^3)$,

$$\int_{\mathbb{R}^3} u^2 v \, dx \leq \|u\|_{12/(3+2t)}^2 \|v\|_{2^*(t)} \leq C \|u\|_{H^s(\mathbb{R}^3)}^2 \|v\|_{D^{t,2}}.$$

Hence there exists a unique Φ_u^t such that $(-\Delta)^t \Phi_u^t = u^2$ and the t -Riesz potential

$$\Phi_u^t(x) = \frac{\Gamma(3/2 - 2t)}{\pi^{3/2} 2^{2t} \Gamma(t)} \int_{\mathbb{R}^3} \frac{u^2(y)}{|x - y|^{3-2t}} \, dy.$$

Furthermore, $\Phi_u^t \geq 0$ and we have

$$\|\Phi_u^t\|_{D^{t,2}} \leq C \|u\|_{H^s(\mathbb{R}^3)}^2 \quad \text{if } 0 < t < 1, \frac{3}{4} < s < 1.$$

Substituting Φ_u^t into (1.1), we are led to the equation

$$(2.2) \quad \epsilon^{2s} (-\Delta)^s u + V(x)u + \Phi_u^t(x)u = Q(x)|u|^{2^*(s)-2}u + f(x, u).$$

Let $\lambda = \epsilon^{-2s}$, then (2.2) becomes

$$(2.3) \quad (-\Delta)^s u + \lambda V(x)u + \lambda \Phi_u^t u = \lambda Q(x)|u|^{2^*(s)-2}u + \lambda f(x, u).$$

Let

$$g(x, u) = Q(x)|u|^{2^*(s)-2}u + f(x, u)$$

and

$$G(x, u) = \int_0^u g(x, s) ds = \frac{1}{2^*(s)} Q(x) |u|^{2^*(s)} + F(x, u).$$

Then, after the change of variables, we obtain the following functional

$$I_\lambda(u) = \frac{1}{2} \int_{\mathbb{R}^3} |(-\Delta)^{s/2} u|^2 + \frac{\lambda}{2} \int_{\mathbb{R}^3} V(x) |u|^2 + \frac{\lambda}{4} \int_{\mathbb{R}^3} \Phi_u^t |u|^2 - \lambda \int_{\mathbb{R}^3} G(x, u).$$

Now we can restate Theorem 1.1 as follows:

Theorem 2.3. *Let (V_1) , (V_2) , (Q) , (f_1) , (f_2) and (f_3) hold, $0 < t < 1$, $3/4 < s < 1$. Then for any $\delta > 0$, there is $\Lambda_\delta > 0$ such that if $\lambda \geq \Lambda_\delta$, problem (2.3) has at least one nontrivial solution v_λ satisfying*

- (i) $\left(\frac{1}{2} - \frac{1}{\mu}\right) \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_\epsilon|^2 + V(x) |u_\epsilon|^2) \leq \delta \lambda^{1-3/(2s)}$ and
- (ii) $\left(\frac{1}{4} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} Q(x) |u_\epsilon|^{2^*(s)} + \frac{\mu-4}{4} \int_{\mathbb{R}^3} F(x, u_\epsilon) \leq \delta \lambda^{-3/(2s)}.$

3. Behaviors of $(PS)_c$ sequences

Let E be a real Banach space and $I_\lambda: E \rightarrow \mathbb{R}$ be a function of class C^1 . We say that a sequence $\{u_n\} \subset E$ is a $(PS)_c$ sequence at level c if $I_\lambda(u_n) \rightarrow c$ and $I'_\lambda(u_n) \rightarrow 0$. I_λ is said to satisfy the $(PS)_c$ condition if any $(PS)_c$ sequence contains a convergent subsequence. The main result of this section is the following compactness result.

Lemma 3.1. *Let (V_1) , (V_2) , (Q) , (f_1) , (f_2) and (f_3) hold. If $\{u_n\}$ is a $(PS)_c$ sequence of I_λ , then $\{u_n\}$ is bounded in E and $c \geq 0$.*

Proof. If $\{u_n\} \subset E$ is a $(PS)_c$ sequence of I_λ , one has

$$(3.1) \quad I_\lambda(u_n) - \frac{1}{\mu} I'_\lambda(u_n) u_n = c + o(1) \|u_n\|_\lambda$$

as $n \rightarrow \infty$.

Gathering (Q) and (f_3) , we obtain

$$(3.2) \quad \begin{aligned} & I_\lambda(u_n) - \frac{1}{\mu} I'_\lambda(u_n) u_n \\ &= \left(\frac{1}{2} - \frac{1}{\mu}\right) \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_n|^2 + \lambda V(x) |u_n|^2) + \left(\frac{1}{\mu} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} \lambda Q(x) |u_n|^{2^*(s)} \\ & \quad + \frac{\lambda}{\mu} \int_{\mathbb{R}^3} f(x, u_n) u_n - \lambda \int_{\mathbb{R}^3} F(x, u_n) + \lambda \left(\frac{1}{4} - \frac{1}{\mu}\right) \int_{\mathbb{R}^3} \Phi_{u_n}^t |u_n|^2 \\ & \geq \left(\frac{1}{2} - \frac{1}{\mu}\right) \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_n|^2 + \lambda V(x) |u_n|^2). \end{aligned}$$

It follows from (3.1) and (3.2), we have

$$\left(\frac{1}{2} - \frac{1}{\mu}\right) \|u_n\|_\lambda^2 \leq c + o(1)\|u_n\|_\lambda.$$

This estimate implies that $\|u_n\|_\lambda$ is bounded as $n \rightarrow \infty$. Taking the limit in (3.2) shows that $c \geq 0$ and the lemma is proved. \square

Note that u_n is a $(PS)_c$ sequence. By Lemma 3.1, we may assume, without loss of generality, that $u_n \rightharpoonup u$ in E , $u_n \rightarrow u$ in $L_{\text{loc}}^t(\mathbb{R}^3)$ for $2 \leq t < 2^*(s)$ and $u_n \rightarrow u$ a.e. for $x \in \mathbb{R}^3$. Obviously, u is a critical point of I_λ .

Lemma 3.2. *Let $t \in [2, 2^*(s))$ and $\{u_n\}$ be a bounded $(PS)_c$ sequence. Then there is a subsequence $\{u_{n_j}\}$ such that, for each $\epsilon > 0$, there exists $r_\epsilon > 0$ such that*

$$\limsup_{j \rightarrow \infty} \int_{B_j \setminus B_r} |u_{n_j}|^t \leq \epsilon$$

for all $r \geq r_\epsilon$, where $B_k = \{x \in \mathbb{R}^N : |x| \leq k\}$.

Proof. Noting that $u_n \rightarrow u$ in $L_{\text{loc}}^t(\mathbb{R}^N)$ for $2 \leq t < 2^*(s)$ as $n \rightarrow \infty$, we have, for each $j \in \mathbb{N}$,

$$\int_{B_j} |u_n|^t \rightarrow \int_{B_j} |u|^t \quad \text{as } n \rightarrow \infty,$$

and there exists $\bar{n}_j \in \mathbb{N}$ such that

$$\int_{B_j} (|u_n|^t - |u|^t) < \frac{1}{j} \quad \text{for all } n = \bar{n}_j + j, j = 1, 2, \dots$$

Without loss of generality, we can assume $\bar{n}_{j+1} \geq \bar{n}_j$. For $n_j = \bar{n}_j + j$, we deduce

$$\int_{B_j} (|u_{n_j}|^t - |u|^t) < \frac{1}{j}.$$

Notice that there exists an r_ϵ such that $r \geq r_\epsilon$, and the following relation is satisfied:

$$\int_{B_r^c} |u|^t < \frac{1}{3}\epsilon.$$

We have

$$\begin{aligned} \int_{B_j \setminus B_r} |u_{n_j}|^t &= \int_{B_j} (|u_{n_j}|^t - |u|^t) + \int_{B_j \setminus B_r} |u|^t + \int_{B_r} (|u|^t - |u_{n_j}|^t) \\ &\leq \frac{1}{j} + \int_{B_r^c} |u|^t + \int_{B_r} (|u|^t - |u_{n_j}|^t) \\ &\leq \epsilon \quad \text{as } j \rightarrow \infty. \end{aligned}$$

\square

Remark 3.3. From the proof of Lemma 3.2, we can find the same subsequence $\{u_{n_j}\}$ such that the result of Lemma 3.2 holds for both $s = 2$ and $s = q$.

Let $\eta: [0, \infty) \rightarrow [0, 1]$ be a smooth function satisfying $\eta(t) = 1$ if $t \leq 1$, $\eta(t) = 0$ if $t \geq 2$. Define

$$\tilde{u}_j(x) = \eta\left(\frac{2|x|}{j}\right) u(x).$$

Using the same argument in [26], we obtain

$$(3.3) \quad \|u - \tilde{u}_j\|_\lambda \rightarrow 0 \quad \text{as } j \rightarrow \infty.$$

Then we have the following lemma which was proved in [29].

Lemma 3.4. [12, Lemma 3.4] *Let $\{u_{n_j}\}$ be defined in Lemma 3.2. Then we have*

$$\lim_{j \rightarrow \infty} \left| \int_{\mathbb{R}^3} (f(x, u_{n_j}) - f(x, u_{n_j} - \tilde{u}_j) - f(x, \tilde{u}_j)) \phi \right| = 0$$

uniformly in $\phi \in E$ with $\|\phi\|_\lambda \leq 1$.

Lemma 3.5. *Let $\{u_n\}$ be defined in Lemma 3.2. Then*

$$(i) \quad I_\lambda(u_n - \tilde{u}_n) \rightarrow c - I_\lambda(u);$$

$$(ii) \quad I'_\lambda(u_n - \tilde{u}_n) \rightarrow 0.$$

Proof. Since $V(x)$ and $Q(x)$ are bounded, by (3.3) and along the lines in the Brezis–Lieb lemma, we have

$$(3.4) \quad \begin{aligned} \int_{\mathbb{R}^3} V(x) (|u_n|^2 - |u_n - \tilde{u}_n|^2 - |\tilde{u}_n|^2) &\rightarrow 0, \\ \int_{\mathbb{R}^3} Q(x) (|u_n|^{2^*(s)} - |u_n - \tilde{u}_n|^{2^*(s)} - |\tilde{u}_n|^{2^*(s)}) &\rightarrow 0, \\ \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|u_n(x) - u_n(y)|^2}{|x - y|^{3+2s}} dx dy &= \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|u_n(x) - \tilde{u}_n(x) - u_n(y) + \tilde{u}_n(y)|^2}{|x - y|^{3+2s}} dx dy \\ &\quad + \int_{\mathbb{R}^3} \int_{\mathbb{R}^3} \frac{|\tilde{u}_n(x) - \tilde{u}_n(y)|^2}{|x - y|^{3+2s}} dx dy + o(1) \end{aligned}$$

and

$$\int_{\mathbb{R}^3} \Phi_{u_n - \tilde{u}_n} |u_n - \tilde{u}_n|^2 + \int_{\mathbb{R}^3} \Phi_{\tilde{u}_n} |\tilde{u}_n|^2 - \int_{\mathbb{R}^3} \Phi_{u_n} |u_n|^2 \rightarrow 0.$$

Similar to the proof of Lemma 3.4, it is not difficult to check that

$$(3.5) \quad \int_{\mathbb{R}^3} (F(x, u_n) - F(x, u_n - \tilde{u}_n) - F(x, \tilde{u}_n)) \rightarrow 0.$$

Gathering (3.4) to (3.5) and the facts that $I_\lambda(u_n) \rightarrow c$ and $I_\lambda(\tilde{u}_n) \rightarrow I_\lambda(u)$ as $n \rightarrow \infty$, we get Lemma 3.5(i).

To prove (ii), note that, for any $\phi \in E$,

$$\begin{aligned}
& I'_\lambda(u_n - \tilde{u}_n)\phi \\
&= I'_\lambda(u_n)\phi - I'_\lambda(\tilde{u}_n)\phi \\
&+ \lambda \int_{\mathbb{R}^3} Q(x)(|u_n|^{2^*(s)-2}u_n - |u_n - \tilde{u}_n|^{2^*(s)-2}(u_n - \tilde{u}_n) + |\tilde{u}_n|^{2^*(s)-2}\tilde{u}_n)\phi \\
&+ \lambda \int_{\mathbb{R}^3} (f(x, u_n) - f(x, u_n - \tilde{u}_n) - f(x, \tilde{u}_n))\phi \\
&+ \lambda \int_{\mathbb{R}^3} (\Phi_{u_n - \tilde{u}_n}(u_n - \tilde{u}_n) + \Phi_{\tilde{u}_n}\tilde{u}_n - \Phi_{u_n}u_n)\phi.
\end{aligned}$$

By standard argument, we can check that

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^3} Q(x)(|u_n|^{2^*(s)-2}u_n - |u_n - \tilde{u}_n|^{2^*(s)-2}(u_n - \tilde{u}_n) + |\tilde{u}_n|^{2^*(s)-2}\tilde{u}_n)\phi = 0$$

uniformly in $\|\phi\|_\lambda \leq 1$. By Lemma 3.4, we have

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}^3} (f(x, u_n) - f(x, u_n - \tilde{u}_n) - f(x, \tilde{u}_n))\phi = 0$$

uniformly in $\|\phi\|_\lambda \leq 1$. Hence, we have

$$I'_\lambda(u_n - \tilde{u}_n) \rightarrow 0. \quad \square$$

Proposition 3.6. *Assume that (V₁), (V₂), (Q), (f₁), (f₂) and (f₃) hold, $0 < t < 1$, $3/4 < s < 1$. If $\{u_n\}$ is a (PS)_c sequence of I_λ , then there exists a constant $\alpha_0 > 0$ independent of λ , either $u_n \rightarrow u$ in E , or*

$$c - I_\lambda(u) \geq \alpha_0 \lambda^{1-3/(2s)}.$$

Proof. Taking $v_n := u_n - \tilde{u}_n$. We have $u_n - u = v_n + (\tilde{u}_n - u)$, and by (3.3), $u_n \rightarrow u$ if and only if $v_n \rightarrow 0$. By Lemma 3.5, one has $I_\lambda(v_n) \rightarrow c - I_\lambda(u)$ and $I'_\lambda(v_n) \rightarrow 0$. Using (f₃), we get

$$\begin{aligned}
& I_\lambda(v_n) - \frac{1}{4}I'_\lambda(v_n)v_n \\
&= \frac{1}{4} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}v_n|^2 + \lambda V(x)|v_n|^2) \\
&+ \left(\frac{1}{4} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} \lambda Q(x)|v_n|^{2^*(s)} + \frac{\lambda}{4} \int_{\mathbb{R}^3} f(x, v_n)v_n - \lambda \int_{\mathbb{R}^3} F(x, v_n) \\
&\geq \left(\frac{1}{4} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} \lambda Q_1|v_n|^{2^*(s)}.
\end{aligned}$$

Therefore,

$$(3.6) \quad |v_n|_{2^*(s)}^{2^*(s)} \leq \frac{12(c - I_\lambda(u))}{(4s - 3)Q_1\lambda} + o(1).$$

Set

$$V_b := \max\{V(x), b\},$$

where the positive constant b is mentioned in (V_1) . Since the V^b has a finite measure and $v_n \rightarrow 0$ in $L^2_{\text{loc}}(\mathbb{R}^3)$, we see that

$$(3.7) \quad \int_{\mathbb{R}^3} V(x)|v_n|^2 = \int_{\mathbb{R}^3} V_b|v_n|^2 + o(1).$$

From (f_1) to (f_2) , we deduce for any fixed $\epsilon > 0$ that there exists C_ϵ such that

$$f(x, u)u \leq \epsilon|u|^2 + C_\epsilon|u|^{2^*(s)},$$

thus by (Q), we can find a constant C_b such that

$$(3.8) \quad Q(x)|u|^{2^*(s)} + f(x, u)u \leq b|u|^2 + C_b|u|^{2^*(s)}.$$

Let S be the best fractional Sobolev constant

$$S|u|_{2^*(s)}^2 \leq \int_{\mathbb{R}^3} |(-\Delta)^{s/2}u|^2 \quad \text{for all } u \in H^s(\mathbb{R}^3).$$

Assume that $\{v_n\}$ has no convergent subsequence. Then $\liminf_{n \rightarrow \infty} \|v_n\|_\lambda > 0$ and $c - I_\lambda(u) > 0$. From (3.7) and (3.8), we obtain

$$(3.9) \quad \begin{aligned} & S|v_n|_{2^*(s)}^2 \\ & \leq \int_{\mathbb{R}^3} |(-\Delta)^{s/2}v_n|^2 + \lambda \int_{\mathbb{R}^3} V(x)|v_n|^2 - \lambda \int_{\mathbb{R}^3} V(x)|v_n|^2 \\ & = \lambda \int_{\mathbb{R}^3} Q(x)|v_n|^{2^*(s)} + \lambda \int_{\mathbb{R}^3} f(x, v_n)v_n - \lambda \int_{\mathbb{R}^3} V_b|v_n|^2 + o(1) - \lambda \int_{\mathbb{R}^3} \Phi_{v_n}^t |v_n|^2 \\ & \leq \lambda \int_{\mathbb{R}^3} Q(x)|v_n|^{2^*(s)} + \lambda \int_{\mathbb{R}^3} f(x, v_n)v_n - \lambda \int_{\mathbb{R}^3} b|v_n|^2 + o(1) \\ & \leq \lambda C_b |v_n|_{2^*(s)}^{2^*(s)} + o(1). \end{aligned}$$

From (3.6) and (3.9), we have

$$S \leq \lambda C_b |v_n|_{2^*(s)}^{2^*(s)-2} + o(1) \leq \lambda C_b \left(\frac{12(c - I_\lambda(u))}{(4s - 3)Q_1 \lambda} \right)^{2s/3} + o(1).$$

Therefore, we get

$$\alpha_0 \lambda^{1-3/(2s)} \leq c - I_\lambda(u) + o(1),$$

where $\alpha_0 > 0$ is a constant. □

From above lemmas, we have the following conclusions.

Corollary 3.7. *Assume that (V_1) , (V_2) , (Q), (f_1) , (f_2) and (f_3) hold. Then I_λ satisfies the $(PS)_c$ condition for all $c < \alpha_0 \lambda^{1-3/(2s)}$.*

4. The mountain-pass structure

Lemma 4.1. *Let (V_1) , (V_2) , (Q) , (f_1) , (f_2) and (f_3) hold. There exist $\alpha_\lambda, \rho_\lambda > 0$ such that $I_\lambda(u) > 0$ if $u \in B_{\rho_\lambda} \setminus \{0\}$ and $I_\lambda(u) \geq \alpha_\lambda$ if $u \in \partial B_{\rho_\lambda}$, where $B_{\rho_\lambda} = \{u \in E : \|u\|_\lambda \leq \rho_\lambda\}$.*

Proof. By (f_1) , (f_2) and (f_3) , there is $C_\epsilon > 0$ such that

$$F(x, u) \leq \epsilon|u|^2 + C_\epsilon|u|^{2^*(s)}.$$

Thus, for $\epsilon \leq (4\lambda c_2^2)^{-1}$,

$$\begin{aligned} I_\lambda(u) &= \frac{1}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}u|^2 + \lambda V(x)|u|^2) + \frac{\lambda}{4} \int_{\mathbb{R}^3} \Phi_u^t |u|^2 - \lambda \int_{\mathbb{R}^3} F(x, u) \\ &\geq \frac{1}{2} \|u\|_\lambda^2 - \lambda \epsilon |u|_2^2 - \lambda C_\epsilon |u|_{2^*(s)}^{2^*(s)} \\ &\geq \left(\frac{1}{2} - \lambda \epsilon c_2^2 \right) \|u\|_\lambda^2 - \lambda C_\epsilon |u|_{2^*(s)}^{2^*(s)} \\ &\geq \frac{1}{4} \|u\|_\lambda^2 - \lambda C_\epsilon |u|_{2^*(s)}^{2^*(s)}, \end{aligned}$$

where c_2 is the embedding constant of (2.1). Since $2 < 2^*(s)$, we know that the conclusion of Lemma 4.1 holds. \square

Lemma 4.2. *Under the assumptions of Lemma 4.1, for any finite-dimensional subspace $E_1 \subset E$,*

$$I_\lambda(u) \rightarrow -\infty \quad \text{as } u \in E_1, \|u\|_\lambda \rightarrow \infty.$$

Proof. By (f_3) , we have

$$\begin{aligned} I_\lambda(u) &= \frac{1}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}u|^2 + \lambda V(x)|u|^2) + \frac{\lambda}{4} \int_{\mathbb{R}^3} \Phi_u^t |u|^2 - \lambda \int_{\mathbb{R}^3} F(x, u) \\ &\leq \frac{1}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}u|^2 + \lambda V(x)|u|^2) + C \frac{\lambda}{4} \|u\|_\lambda^4 - \lambda \int_{\mathbb{R}^3} a_0 |u|^l \end{aligned}$$

for all $u \in E_1$. Since all norms in a finite-dimensional space are equivalent and $l > 4$, one easily obtains the desired conclusion. \square

However, in general we do not know if I_λ satisfies $(PS)_c$ condition. Then, we will find a special finite-dimensional subspaces by which we construct sufficiently small minimax levels.

Note that

$$\inf \left\{ \int_{\mathbb{R}^3} |(-\Delta)^{s/2}\phi|^2 : \phi \in C_0^\infty(\mathbb{R}^3), |\phi|_r = 1 \right\} = 0.$$

For any $\sigma > 0$, we can choose $\phi_\sigma \in C_0^\infty(\mathbb{R}^3)$ with $|\phi_\sigma|_r = 1$ and $\text{supp } \phi_\sigma \subset B_{r_\sigma}(0)$ so that $|(-\Delta)^{s/2}\phi_\sigma|_2^2 < \sigma$. Denote

$$(4.1) \quad e_\lambda(x) = \phi_\sigma(\lambda^{1/(2s)}x).$$

Then $\text{supp } e_\lambda \subset B_{\lambda^{-1/(2s)}r_\sigma}(0)$.

Remark that for $k \geq 0$,

$$\begin{aligned}
I_\lambda(ke_\lambda) &= \frac{1}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}(ke_\lambda)|^2 + \lambda V(x)|ke_\lambda|^2) + \frac{\lambda}{4} \int_{\mathbb{R}^3} \Phi_{ke_\lambda}^t |ke_\lambda|^2 - \lambda \int_{\mathbb{R}^3} G(x, ke_\lambda) \\
&\leq \frac{k^2}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}e_\lambda|^2 + \lambda V(x)|e_\lambda|^2) + \frac{\lambda}{4} k^4 \int_{\mathbb{R}^3} \Phi_{e_\lambda}^t |e_\lambda|^2 - \lambda \int_{\mathbb{R}^3} F(x, ke_\lambda) \\
&\leq \frac{k^2}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}e_\lambda|^2 + \lambda V(x)|e_\lambda|^2) + \frac{\lambda}{4} k^4 \int_{\mathbb{R}^3} \Phi_{e_\lambda}^t |e_\lambda|^2 - \lambda a_0 k^l \int_{\mathbb{R}^3} |e_\lambda|^l \\
&= \lambda^{1-3/(2s)} \left(\frac{k^2}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}\phi_\sigma|^2 + V(\lambda^{-1/(2s)}x)|\phi_\sigma|^2) \right. \\
&\quad \left. + \frac{k^4}{4} \lambda^{-t/s} \int_{\mathbb{R}^3} \Phi_{\phi_\sigma}^t |\phi_\sigma|^2 - a_0 k^l \int_{\mathbb{R}^3} |\phi_\sigma|^l \right) \\
&:= \lambda^{1-3/(2s)} \Psi_\lambda(k\phi_\sigma),
\end{aligned}$$

where $\Psi_\lambda \in C^1(E, \mathbb{R})$ is defined by

$$\Psi_\lambda(v) = \frac{1}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}v|^2 + V(\lambda^{-1/(2s)}x)|v|^2) + \frac{1}{4} \lambda^{-t/s} \int_{\mathbb{R}^3} \Phi_v |v|^2 - a_0 \int_{\mathbb{R}^3} |v|^l.$$

Since $l > 4$, thus there exists finite number $k_0 \in [0, +\infty)$ such that

$$\begin{aligned}
&\max_{k \geq 0} \Psi_\lambda(k\phi_\sigma) \\
&= \frac{k_0^2}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}\phi_\sigma|^2 + V(\lambda^{-1/(2s)}x)|\phi_\sigma|^2) + \frac{k_0^4}{4} \lambda^{-t/s} \int_{\mathbb{R}^3} \Phi_{\phi_\sigma} |\phi_\sigma|^2 - a_0 k_0^l \int_{\mathbb{R}^3} |\phi_\sigma|^l \\
&\leq \frac{k_0^2}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2}\phi_\sigma|^2 + V(\lambda^{-1/(2s)}x)|\phi_\sigma|^2) + \frac{k_0^4}{4} C \lambda^{-t/s} \left(\int_{\mathbb{R}^3} |\phi_\sigma|^{12/5} \right)^{5/3}.
\end{aligned}$$

On the one hand, since $V(0) = 0$ and $\text{supp } \phi_\sigma \subset B_{r_\sigma}(0)$, there is Λ_σ such that

$$V(\lambda^{-1/(2s)}x) \leq \frac{\sigma}{|\phi_\sigma|_2^2} \quad \text{for all } |x| \leq r_\sigma \text{ and } \lambda \geq \Lambda_\sigma,$$

and $\lambda^{-t/s} < \sigma$. Then

$$\max_{k \geq 0} \Psi_\lambda(k\phi_\sigma) \leq \tilde{C}\sigma.$$

Hence, for any $\lambda \geq \Lambda_\sigma$,

$$(4.2) \quad \max_{k \geq 0} I_\lambda(ke_\lambda) \leq \tilde{C}\sigma \lambda^{1-3/(2s)}.$$

Now, we show the following lemma.

Lemma 4.3. *Under the assumptions of Lemma 4.1, for any $\delta > 0$ there exists $\bar{\Lambda}_\delta > 0$ such that, for each $\lambda \geq \bar{\Lambda}_\delta$, there is \bar{e}_λ with $\|\bar{e}_\lambda\| > \rho_\lambda$, $I_\lambda(\bar{e}_\lambda) \leq 0$ and*

$$\max_{t \geq 0} I_\lambda(t\bar{e}_\lambda) \leq \delta \lambda^{1-3/(2s)},$$

where ρ_λ is given by Lemma 4.1.

Proof. Choose $\sigma > 0$ so small that $\tilde{C}\sigma \leq \delta$, and the function $e_\lambda \in E$ is defined in (4.1). Take $\bar{\Lambda}_\delta = \Lambda_\sigma$. Let $\bar{t}_\lambda > 0$ be such that $\bar{t}_\lambda \|e_\lambda\|_\lambda > \rho_\lambda$ and $I_\lambda(te_\lambda) \leq 0$ for all $t \geq \bar{t}_\lambda$. Let $\bar{e}_\lambda := \bar{t}_\lambda e_\lambda$. As a consequence of (4.2), we know the conclusion of Lemma 4.3 holds. \square

For any $n^* \in \mathbb{N}$, we can choose n^* functions $\phi_\sigma^j \in C_0^\infty(\mathbb{R}^3)$ with $|\phi_\sigma^j|_r = 1$ and $\text{supp } \phi_\sigma^j \cap \text{supp } \phi_\sigma^k = \emptyset$, $j \neq k$ so that $|(-\Delta)^{s/2} \phi_\sigma^j|_2^2 < \sigma$. Let $r_\sigma^{n^*} > 0$ be such that $\text{supp } \phi_\sigma^j \subset B_{r_\sigma^{n^*}}(0)$ for $j = 1, 2, \dots, n^*$. Let

$$e_\lambda^j(x) = \phi_\sigma^j(\lambda^{1/(2s)}x) \text{ for } j = 1, 2, \dots, n^* \quad \text{and} \quad H_{\lambda\sigma}^{n^*} = \text{span}\{e_\lambda^1, \dots, e_\lambda^{n^*}\}.$$

Observe that for each

$$v = \sum_{j=1}^{n^*} c_j e_\lambda^j \in H_{\lambda\sigma}^{n^*},$$

we have

$$\int_{\mathbb{R}^N} |(-\Delta)^{s/2} v|^2 \leq C \sum_{j=1}^{n^*} |c_j|^2 \int_{\mathbb{R}^N} |(-\Delta)^{s/2} e_\lambda^j|^2$$

for some constant $C > 0$. Hence, we have

$$I_\lambda(v) \leq C \sum_{j=1}^{n^*} I_\lambda(c_j e_\lambda^j)$$

for some constant $C > 0$. It is the similar way to show that

$$I_\lambda(c_j e_\lambda^j) \leq \lambda^{1-3/(2s)} \Psi_\lambda(|c_j| e_\lambda^j).$$

Set

$$\beta_\sigma := \max\{|\phi_\sigma^j|_2^2, j = 1, 2, \dots, n^*\}$$

and choose $\Lambda_{n^*\sigma}$ such that

$$V(\lambda^{-1/(2s)}x) \leq \frac{\sigma}{\beta_\sigma} \quad \text{for all } |x| \leq r_\sigma^{n^*} \text{ and } \lambda \geq \Lambda_{n^*\sigma}.$$

As before, we can obtain the following

$$(4.3) \quad \max_{u \in H_{\lambda\sigma}^{n^*}} I_\lambda(v) \leq \tilde{C}\sigma \lambda^{1-3/(2s)}$$

for all $\lambda \geq \Lambda_{n^*\sigma}$.

As a consequence of this estimate, we obtain the following conclusion.

Lemma 4.4. *Assume that (V₁), (V₂), (Q), (f₁), (f₂) and (f₃) are satisfied. For any $n^* \in \mathbb{N}$ and $\delta > 0$ there exists $\bar{\Lambda}_{n^*\delta} > 0$ such that, for each $\lambda \geq \bar{\Lambda}_{n^*\delta}$, there exists an n^* -dimensional subspace $F_{\lambda n^*}$ satisfying*

$$\max_{u \in F_{\lambda n^*}} I_\lambda(v) \leq \delta \lambda^{1-3/(2s)}.$$

5. Proof of the main result

In section, we prove the main result.

Proof of Theorem 2.3. For any $0 < \delta < \alpha_0$, by Lemma 4.3 we choose Λ_δ and define for $\lambda \geq \Lambda_\delta$ the minimax value

$$c_\lambda := \inf_{\gamma \in \Gamma_\lambda} \max_{t \in [0,1]} I_\lambda(t\bar{e}_\lambda),$$

where

$$\Gamma_\lambda := \{\gamma \in C([0,1], E) : \gamma(0) = 0 \text{ and } \gamma(1) = \bar{e}_\lambda\}.$$

It follows from Lemma 4.1 and (4.3) that

$$\alpha_\lambda \leq c_\lambda \leq \delta \lambda^{1-3/(2s)}.$$

From Corollary 3.7, we know that I_λ satisfies the $(PS)_c$ condition. Using the Mountain-pass theorem, we get that there is $u_\lambda \in E$ such that $I'_\lambda(u_\lambda) = 0$ and $I_\lambda(u_\lambda) = c_\lambda$.

Because u_λ is a critical point of I_λ , for $\nu \in [4, 2^*(s)]$,

$$\begin{aligned} \delta \lambda^{1-3/(2s)} &\geq I_\lambda(u_\lambda) = I_\lambda(u_\lambda) - \frac{1}{\nu} I'_\lambda(u_\lambda) u_\lambda \\ &= \frac{1}{2} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_\lambda|^2 + \lambda V(x) |u_\lambda|^2) + \frac{\lambda}{4} \int_{\mathbb{R}^3} \Phi_{u_\lambda}^t |u_\lambda|^2 - \lambda \int_{\mathbb{R}^3} G(x, u_\lambda) \\ &\quad - \frac{1}{\nu} \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_\lambda|^2 + \lambda V(x) |u_\lambda|^2) - \frac{\lambda}{\nu} \int_{\mathbb{R}^3} \Phi_{u_\lambda}^t |u_\lambda|^2 + \frac{\lambda}{\nu} \int_{\mathbb{R}^3} g(x, u_\lambda) u_\lambda \\ &\geq \left(\frac{1}{2} - \frac{1}{\nu}\right) \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_\lambda|^2 + \lambda V(x) |u_\lambda|^2) + \lambda \left(\frac{1}{4} - \frac{1}{\nu}\right) \int_{\mathbb{R}^3} \Phi_{u_\lambda}^t |u_\lambda|^2 \\ &\quad + \lambda \left(\frac{1}{\nu} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} Q(x) |u_\lambda|^{2^*(s)} + \lambda \left(\frac{\mu}{\nu} - 1\right) \int_{\mathbb{R}^3} F(x, u_\lambda), \end{aligned}$$

where the constant μ is defined in (f₃). Choosing $\nu = \mu$, we have

$$\left(\frac{1}{2} - \frac{1}{\mu}\right) \int_{\mathbb{R}^3} (|(-\Delta)^{s/2} u_\lambda|^2 + V(x) |u_\lambda|^2) \leq \delta \lambda^{1-3/(2s)}$$

and taking $\nu = 4$ we obtain

$$\left(\frac{1}{4} - \frac{1}{2^*(s)}\right) \int_{\mathbb{R}^3} Q(x) |u_\lambda|^{2^*(s)} + \frac{\mu-4}{4} \int_{\mathbb{R}^3} F(x, u_\lambda) \leq \delta \lambda^{-3/(2s)}. \quad \square$$

Acknowledgments

The work was supported by the National Natural Science Foundation of China, No. 11601139, Hubei Provincial Natural Science Foundation of China, No. 2019CFB522 and Scientific Research and Innovation Team of Hubei Normal University, No. 2019CZ010.

References

- [1] C. O. Alves and M. A. S. Souto, *Existence of least energy nodal solution for a Schrödinger–Poisson system in bounded domains*, Z. Angew. Math. Phys. **65** (2014), no. 6, 1153–1166.
- [2] A. Ambrosetti and P. H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Functional Analysis **14** (1973), 349–381.
- [3] V. Ambrosio, *An existence result for a fractional Kirchhoff–Schrödinger–Poisson system*, Z. Angew. Math. Phys. **69** (2018), no. 2, Paper No. 30, 13 pp.
- [4] A. Azzollini, *Concentration and compactness in nonlinear Schrödinger–Poisson system with a general nonlinearity*, J. Differential Equations **249** (2010), no. 7, 1746–1763.
- [5] T. Bartsch and Z. Q. Wang, *Existence and multiplicity results for some superlinear elliptic problems on \mathbb{R}^N* , Comm. Partial Differential Equations **20** (1995), no. 9-10, 1725–1741.
- [6] G. Cerami and G. Vaira, *Positive solutions for some non-autonomous Schrödinger–Poisson systems*, J. Differential Equations **248** (2010), no. 3, 521–543.
- [7] X. Chang, *Ground state solutions of asymptotically linear fractional Schrödinger equations*, J. Math. Phys. **54** (2013), no. 6, 061504, 10 pp.
- [8] S. Chen, J. Peng and X. Tang, *Ground state solutions for asymptotically periodic fractional Schrödinger–Poisson problems with asymptotically cubic or super-cubic nonlinearities*, Math. Method Appl. Sci. **40** (2017), no. 13, 4948–4961.
- [9] S. Chen and X. Tang, *Berestycki–Lions conditions on ground state solutions for a nonlinear Schrödinger equation with variable potentials*, Adv. Nonlinear Anal. **9** (2020), no. 1, 496–515.
- [10] T. D’Aprile and J. Wei, *On bound states concentrating on spheres for the Maxwell–Schrödinger equation*, SIAM J. Math. Anal. **37** (2005), no. 1, 321–342.
- [11] E. Di Nezza, G. Palatucci and E. Valdinoci, *Hitchhiker’s guide to the fractional Sobolev spaces*, Bull. Sci. Math. **136** (2012), no. 5, 521–573.
- [12] Y. Ding and F. Lin, *Solutions of perturbed Schrödinger equations with critical nonlinearity*, Calc. Var. Partial Differential Equations **30** (2007), no. 2, 231–249.

- [13] X. He and W. Zou, *Existence and concentration of ground states for Schrödinger–Poisson equations with critical growth*, J. Math. Phys. **53** (2012), no. 2, 023702, 19 pp.
- [14] Y. Jiang and H.-S. Zhou, *Schrödinger–Poisson system with steep potential well*, J. Differential Equations **251** (2011), no. 3, 582–608.
- [15] G. Li, S. Peng and C. Wang, *Multi-bump solutions for the nonlinear Schrödinger–Poisson system*, J. Math. Phys. **52** (2011), no. 5, 053505, 19 pp.
- [16] G. Li, S. Peng and S. Yan, *Infinitely many positive solutions for the nonlinear Schrödinger–Poisson system*, Commun. Contemp. Math. **12** (2010), no. 6, 1069–1092.
- [17] W. Liu, *Existence of multi-bump solutions for the fractional Schrödinger–Poisson system*, J. Math. Phys. **57** (2016), no. 9, 091502, 17 pp.
- [18] ———, *Infinitely many positive solutions for the fractional Schrödinger–Poisson system*, Pacific J. Math. **287** (2017), no. 2, 439–464.
- [19] X. Mingqi, V. D. Rădulescu and B. Zhang, *Nonlocal Kirchhoff diffusion problems: Local existence and blow-up of solutions*, Nonlinearity **31** (2018), no. 7, 3228–3250.
- [20] ———, *A critical fractional Choquard–Kirchhoff problem with magnetic field*, Commun. Contemp. Math. **21** (2019), no. 4, 1850004, 36 pp.
- [21] ———, *Fractional Kirchhoff problems with critical Trudinger–Moser nonlinearity*, Calc. Var. Partial Differential Equations **58** (2019), no. 2, Paper No. 57, 27 pp.
- [22] E. G. Murcia and G. Siciliano, *Positive semiclassical states for a fractional Schrödinger–Poisson system*, Differential Integral Equations **30** (2017), no. 3-4, 231–258.
- [23] N. S. Papageorgiou, V. D. Rădulescu and D. D. Repovš, *Nonlinear Analysis—Theory and methods*, Springer Monographs in Mathematics, Springer, Cham, 2019.
- [24] L. Shen, *Existence result for fractional Schrödinger–Poisson systems involving a Bessel operator without Ambrosetti–Rabinowitz condition*, Comput. Math. Appl. **75** (2018), no. 1, 296–306.
- [25] L. Shen and X. Yao, *The existence and concentration of positive ground state solutions for a class of fractional Schrödinger–Poisson systems with steep potential wells*, arXiv:1703.07537.

- [26] Z. Shen and F. Gao, *On the existence of solutions for the critical fractional Laplacian equation in \mathbb{R}^N* , Abstr. Appl. Anal. **2014**, Art. ID 143741, 10 pp.
- [27] K. Teng, *Existence of ground state solutions for the nonlinear fractional Schrödinger–Poisson system with critical Sobolev exponent*, J. Differential Equations **261** (2016), no. 6, 3061–3106.
- [28] J. Vétois and S. Wang, *Infinitely many solutions for cubic nonlinear Schrödinger equations in dimension four*, Adv. Nonlinear Anal. **8** (2019), no. 1, 715–724.
- [29] C. Wang, *Solutions for perturbed biharmonic equations with critical nonlinearity*, Math. Methods Appl. Sci. **37** (2014), no. 6, 882–893.
- [30] C. Wang and J. Wang, *Solutions of perturbed p -Laplacian equations with critical nonlinearity*, J. Math. Phys. **54** (2013), no. 1, 013702, 16 pp.
- [31] L. Wang, V. D. Rădulescu and B. Zhang, *Infinitely many solutions for fractional Kirchhoff–Schrödinger–Poisson systems*, J. Math. Phys. **60** (2019), no. 1, 011506, 18 pp.
- [32] Z. Wang and H.-S. Zhou, *Positive solution for a nonlinear stationary Schrödinger–Poisson system in \mathbb{R}^3* , Discrete Contin. Dyn. Syst. **18** (2007), no. 4, 809–816.
- [33] Y.-F. Xue and C.-L. Tang, *Existence of a bound state solution for quasilinear Schrödinger equations*, Adv. Nonlinear Anal. **8** (2019), no. 1, 323–338.
- [34] J. Zhang, J. M. Do Ó and M. Squassina, *Fractional Schrödinger–Poisson systems with a general subcritical or critical nonlinearity*, Adv. Nonlinear Stud. **16** (2016), no. 1, 15–30.
- [35] J. Zhang, *Existence and multiplicity results for the fractional Schrödinger–Poisson systems*, arXiv: 1507.01205.

Lvzhou Zheng

School of Mathematics and Statistics, Hubei Normal University, Huangshi, 435002,
China

E-mail address: mathat66@163.com