

**STRONG CONVERGENCE THEOREMS FOR STRICTLY
PSEUDOCONTRACTIVE MAPPINGS OF
BROWDER-PETRYSHYN TYPE***

Lu-Chuan Zeng¹, Ngai-Ching Wong² and Jen-Chih Yao³

Abstract. Let $q > 1$ and E be a real q -uniformly smooth Banach space, K be a nonempty closed convex subset of E and $T : K \rightarrow K$ be a strictly pseudocontractive mapping in the sense of F. E. Browder and W.V. Pstryshyn [1]. Let $\{u_n\}$ be a bounded sequence in K and $\{\alpha_n\}$ and $\{\beta_n\}$ be real sequences in $[0, 1]$ satisfying some restrictions. Let $\{x_n\}$ be the sequence generated from an arbitrary $x_1 \in K$ by the Ishikawa iteration process with errors: $y_n = (1 - \beta_n)x_n + \beta_n T x_n$, $x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n + u_n$, $n \geq 1$. Sufficient and necessary conditions for the strong convergence $\{x_n\}$ to a fixed point of T is established.

1. INTRODUCTION AND PRELIMINARIES

Let E be an arbitrary real Banach space and let $J_q (q > 1)$ denote the generalized duality mapping from E into 2^{E^*} given by

$$J_q(x) = \{f \in E^* : \langle x, f \rangle = \|x\|^q = \|x\| \|f\|\},$$

where E^* denote the dual space of E and $\langle \cdot, \cdot \rangle$ denotes the generalized duality pairing between E and E^* . In particular, J_2 is called the normalized duality mapping and

Received February 12, 2004, revised November 12, 2004.

Communicated by Mau-Hsiang Shih.

2000 *Mathematics Subject Classification*: 47H09, 47H10, 47H17.

Key words and phrases: Fixed point, Strictly pseudocontractive mapping, Ishikawa iteration process with errors, q -Uniformly smooth Banach space.

*The authors thank the referee for his (her) useful and helpful comments and suggestions.

¹This research was partially supported by the Teaching and Research Award Fund for Outstanding Young Teachers in Higher Education Institutions of MOE, China and the Dawn Program Foundation in Shanghai.

²This research was partially supported by grant from the National Science Council of Taiwan.

³This research was partially supported by grant from the National Science Council of Taiwan.

it is usually denoted by J . It is well known (see [11]) that $J_q(x) = \|x\|^{q-2}J(x)$ if $x \neq 0$ and that if E^* is strictly convex, then J_q is single-valued.

Definition 1.1. A mapping T with domain $D(T)$ and range $R(T)$ in E is called strictly pseudocontractive [1] if there exists $\lambda > 0$ such that for all $x, y \in D(T)$ there exists $j(x - y) \in J(x - y)$ satisfying

$$(1.1) \quad \langle (I - T)x - (I - T)y, j(x - y) \rangle \geq \lambda \|(I - T)x - (I - T)y\|^2$$

where I denotes the identity operator.

Remark 1.1. Without loss of generality we may assume $\lambda \in (0, 1)$. In Hilbert spaces, (1.1) is equivalent to the following inequality

$$(1.2) \quad \|Tx - Ty\|^2 \leq \|x - y\|^2 + k\|(I - T)x - (I - T)y\|^2, \quad k = 1 - 2\lambda < 1.$$

The Mann iterative process (with errors) and the Ishikawa iterative process (with errors) have been extensively applied to approximate the solutions of nonlinear operator equations or fixed points of nonlinear mappings in Hilbert spaces or Banach spaces in the literature. See, e.g., [3-10]. In 1974, Rhoades [9] proved the following convergence theorem by using the Mann iterative process.

Theorem 1.1. *Let H be a real Hilbert space and K a nonempty compact convex subset of H . Let $T : K \rightarrow K$ be a strictly pseudocontractive mapping and let $\{\alpha_n\}$ be a real sequence satisfying the conditions: (i) $\alpha_0 = 1$; (ii) $0 < \alpha_n < 1, n \geq 1$; (iii) $\sum_{n=1}^{\infty} \alpha_n = \infty$; (iv) $\lim_{n \rightarrow \infty} \alpha_n = \alpha < 1$. Then the sequence $\{x_n\}$ generated from an arbitrary $x_0 \in K$ by the Mann iterative process*

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTx_n, \quad n \geq 0$$

converges strongly to a fixed point of T .

Let E be a real Banach space. The modulus of smoothness of E is defined as the function $\rho_E : [0, \infty) \rightarrow [0, \infty)$:

$$\rho_E(\tau) = \sup \left\{ \frac{1}{2}(\|x + y\| + \|x - y\|) - 1 : \|x\| \leq 1, \|y\| \leq \tau \right\}.$$

E is said to be uniformly smooth if and only if $\lim_{\tau \rightarrow 0^+} (\rho_E(\tau)/\tau) = 0$. Let $q > 1$. The space E is said to be q -uniformly smooth (or to have a modulus of smoothness of power type $q > 1$), if there exists a constant $c_q > 0$ such that $\rho_E(\tau) \leq c_q\tau^q$. It is well known that Hilbert spaces, L_p and l_p spaces, $1 < p < \infty$, as well as the Sobolev spaces, $W_m^p, 1 < p < \infty$, are q -uniformly smooth. Hilbert spaces are

2-uniformly smooth while if $1 < p \leq 2$, L_p , l_p and W_m^p are p -uniformly smooth. If $p \geq 2$, L_p , l_p and W_m^p are 2-uniformly smooth.

Theorem 1.2. [11] *Let $q > 1$ and E be a real smooth Banach space. Then the following are equivalent:*

- (1) E is q -uniformly smooth.
- (2) There exists a constant $c_q > 0$ such that for all $x, y \in E$

$$(1.3) \quad \|x + y\|^q \leq \|x\|^q + q\langle y, J_q(x) \rangle + c_q\|y\|^q.$$

- (3) There exists a constant d_q such that for all $x, y \in E$ and $t \in [0, 1]$

$$(1.4) \quad \|(1-t)x + ty\|^q \geq (1-t)\|x\|^q + t\|y\|^q - \omega_q(t)d_q\|x - y\|^q,$$

where $\omega_q(t) = t^q(1-t) + t(1-t)^q$.

Furthermore, it was shown in [12, Remark 5] that if E is q -uniformly smooth ($q > 1$), then for all $x, y \in E$, there exists a constant $L_* > 0$ such that

$$(1.5) \quad \|J_q(x) - J_q(y)\| \leq L_*\|x - y\|^{q-1}.$$

Recently, Osilike and Udomene [13] improved, unified and developed the above Theorem 1.1 and Browder and Petryshyn's corresponding result [1] in two aspects: (i) Hilbert spaces are extended to the setting of q -uniformly smooth Banach spaces ($q > 1$); (ii) Mann iterative process is extended to the case of Ishikawa iterative process.

Theorem 1.3. [13, Theorem 2]. *Let E be a real q -uniformly smooth Banach space which is also uniformly convex. Let K be a nonempty closed convex subset of E and $T : K \rightarrow K$ be a strictly pseudocontractive mapping with a nonempty fixed-point set $F(T)$. Let $\{\alpha_n\}$ and $\{\beta_n\}$ be real sequences in $[0, 1]$ satisfying the conditions:*

- (i) $0 < a \leq \alpha_n^{q-1} \leq b < (q\lambda^{q-1}/c_q)(1 - \beta_n), \forall n \geq 1$ and for some constants $a, b \in (0, 1)$;
- (ii) $\sum_{n=1}^{\infty} \beta_n^\tau < \infty$, where $\tau = \min\{1, (q-1)\}$.

If $\{x_n\}$ is the sequence generated from an arbitrary $x_1 \in K$ by the Ishikawa iterative process

$$\begin{cases} y_n = (1 - \beta_n)x_n + \beta_nTx_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTy_n, n \geq 1. \end{cases}$$

then $\{x_n\}$ converges weakly to a fixed point of T .

Definition 1.2. A mapping T with domain $D(T)$ and range $R(T)$ in E is called demiclosed at a point p if whenever $\{x_n\}$ is a sequence in $D(T)$ such that $\{x_n\}$ converges weakly to $x \in D(T)$ and $\{Tx_n\}$ converges strongly to p , then $Tx = p$. Furthermore, T is called demicompact if whenever $\{x_n\}$ is a bounded sequence in $D(T)$ such that $\{x_n - Tx_n\}$ converges strongly, then $\{x_n\}$ has a subsequence which converges strongly.

Theorem 1.4. [13, Corollary 2]. Let E be a real q -uniformly smooth Banach space, K be a nonempty closed convex subset of E , $T : K \rightarrow K$ be a demicompact strictly pseudocontractive mapping with a nonempty fixed-point set $F(T)$. Let $\{\alpha_n\}, \{\beta_n\}$ and $\{x_n\}$ be as in Theorem 1.3. Then $\{x_n\}$ converges strongly to a fixed point of T .

Let E be a real q -uniformly smooth Banach space, K be a nonempty closed convex (not necessarily bounded) subset of E with $K + K \subseteq K$, and $T : K \rightarrow K$ be a strictly pseudocontractive mapping with $F(T) \neq \emptyset$. Let $\{u_n\}$ be a bounded sequence in K and $\{\alpha_n\}, \{\beta_n\}$ be real sequences in $[0, 1]$ satisfying certain restrictions. Let $\{x_n\}$ be the sequence generated from $x_1 \in K$ by the Ishikawa iterative process with errors:

$$(1.6) \quad \begin{cases} y_n = (1 - \beta_n)x_n + \beta_nTx_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTy_n + u_n, n \geq 1. \end{cases}$$

In this paper, we establish the sufficient and necessary conditions for the strong convergence of $\{x_n\}$ to a fixed point of T . It is worth noting that comparing with [13, Theorem 2 and Corollary 2] our results have the following features: (i) The uniform convexity assumption on E is removed. (ii) The Ishikawa iterative process is replaced by the Ishikawa iterative process with errors. (iii) Our restrictions imposed on $\{\alpha_n\}$ are much weaker than those in [13, Theorem 2 and Corollary 2]. (iv) We establish the sufficient and necessary conditions on the strong convergence of the Ishikawa iterative process with errors. Furthermore, our results also improve and extend the corresponding results in [1, 9].

Now, we give some preliminaries which will be used in the sequel.

From (1.2) we have

$$\|x - y\| \geq \lambda\|x - y - (Tx - Ty)\| \geq \lambda\|Tx - Ty\| - \lambda\|x - y\|.$$

so that $\|Tx - Ty\| \leq L\|x - y\|, \forall x, y \in K$, where $L = (1 + \lambda)/\lambda$. Since

$$\|x - y\| \geq \lambda\|x - y - (Tx - Ty)\|,$$

we have

$$\begin{aligned} \langle x - Tx - (y - Ty), j_q(x - y) \rangle &= \|x - y\|^{q-2} \langle x - Tx - (y - Ty), j(x - y) \rangle \\ &\geq \lambda \|x - y\|^{q-2} \|x - Tx - (y - Ty)\|^2 \\ &\geq \lambda^{q-1} \|x - Tx - (y - Ty)\|^q. \end{aligned} \tag{1.7}$$

Lemma 1.1. [10]. *Let $\{a_n\}_{n=1}^\infty$ and $\{b_n\}_{n=1}^\infty$ be sequences of nonnegative real numbers such that $\sum_{n=1}^\infty b_n < \infty$ and $a_{n+1} \leq a_n + b_n, \forall n \geq 1$. Then $\lim_{n \rightarrow \infty} a_n$ exists.*

2. MAIN RESULTS

Throughout this section, λ denotes the constant appearing in (1.1). L stands for the Lipschitz constant of T , and $c_q, d_q, \omega_q(t)$, and L_* are the constants appearing in inequalities (1.3)-(1.5), respectively.

Lemma 2.1. *Let $q > 1$ and E be a real q -uniformly smooth Banach space and K be a nonempty convex subset of E with $K + K \subseteq K$, and $T : K \rightarrow K$ be a strictly pseudocontractive mapping with $F(T) \neq \emptyset$. Let $\{u_n\}_{n=1}^\infty$ be a bounded sequence in K , and $\{\alpha_n\}_{n=1}^\infty, \{\beta_n\}_{n=1}^\infty$ be real sequences in $[0, 1]$ satisfying the following conditions: (i) $\sum_{n=1}^\infty \|u_n\| < \infty$, (ii) $\alpha_n \leq \lambda(q/c_q)^{1/(q-1)}$, and $\sum_{n=1}^\infty \beta_n^\tau < \infty$ where $\tau = \min\{1, (q - 1)\}$. Let $\{x_n\}$ be the sequence generated from an arbitrary $x_1 \in K$ by the Ishikawa iterative process (1.6) with errors. Then*

$$(i) \|x_{n+1} - x^*\|^q \leq (1 + \delta_n) \|x_n - x^*\|^q + \theta_n, \forall n \geq 1, \forall x^* \in F(T),$$

where

$$\delta_n = 2\alpha_n\beta_n\lambda^{q-1}qd_q(1+L)^q + \alpha_n\beta_n^{q-1}qL_*(1+L)^{q+1} + \alpha_n\beta_nq\lambda^{q-1}(1+L^2)^q$$

and

$$\theta_n = q\|u_n\|\|x_{n+1} - u_n - x^*\|^{q-1} + c_q\|u_n\|^q.$$

(ii) *There exists a constant $M > 0$ (e.g., $M = e^{\sum_{n=1}^\infty \delta_n}$) such that*

$$\|x_{n+m} - x^*\|^q \leq M\|x_n - x^*\|^q + M \sum_{k=n}^{n+m-1} \theta_k, \forall n, m \geq 1, \forall x^* \in F(T).$$

Proof. (i) For each $n \geq 1$, from (1.6) we obtain

$$(2.1) \quad x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T y_n + u_n.$$

Let x^* be an arbitrary element in $F(T)$. Then it follows from (1.3) that

$$(2.2) \quad \begin{aligned} \|x_{n+1} - x^*\|^q &= \|(1 - \alpha_n)x_n + \alpha_n T y_n + u_n - x^*\|^q \\ &\leq \|(1 - \alpha_n)x_n + \alpha_n T y_n - x^*\|^q \\ &\quad + q \langle u_n, j_q(x_{n+1} - u_n - x^*) \rangle + c_q \|u_n\|^q \\ &\leq \|(1 - \alpha_n)x_n + \alpha_n T y_n - x^*\|^q \\ &\quad + q \|u_n\| \|x_{n+1} - u_n - x^*\|^{q-1} + c_q \|u_n\|^q. \end{aligned}$$

Observe that

$$(2.3) \quad \begin{aligned} \|(1 - \alpha_n)x_n + \alpha_n T y_n - x^*\|^q &= \|x_n - x^* - \alpha_n(x_n - T y_n)\|^q \\ &\leq \|x_n - x^*\|^q - q \alpha_n \langle x_n - T y_n, j_q(x_n - x^*) \rangle \\ &\quad + \alpha_n^q c_q \|x_n - T y_n\|^q, \end{aligned}$$

$$\begin{aligned} \langle x_n - T y_n, j_q(x_n - x^*) \rangle &= \langle x_n - y_n, j_q(x_n - x^*) \rangle + \langle y_n - T y_n, j_q(x_n - x^*) \rangle \\ &= \beta_n \langle x_n - T x_n - (x^* - T x^*), j_q(x_n - x^*) \rangle \\ &\quad + \langle y_n - T y_n, j_q(x_n - x^*) \rangle \\ &\geq \beta_n \lambda^{q-1} \|x_n - T x_n - (x^* - T x^*)\|^q \\ &\quad + \langle y_n - T y_n, j_q(x_n - x^*) \rangle \\ &= \beta_n \lambda^{q-1} \|x_n - T x_n\|^q + \langle y_n - T y_n, j_q(x_n - x^*) \rangle, \end{aligned}$$

and by (1.7)

$$(2.4) \quad \begin{aligned} \langle y_n - T y_n, j_q(x_n - x^*) \rangle &= \langle y_n - T y_n - (x^* - T x^*), j_q(x_n - x^*) - j_q(y_n - x^*) \rangle \\ &\quad + \langle y_n - T y_n - (x^* - T x^*), j_q(y_n - x^*) \rangle \\ &\geq \lambda^{q-1} \|y_n - T y_n - (x^* - T x^*)\|^q \\ &\quad + \langle y_n - T y_n - (x^* - T x^*), j_q(x_n - x^*) - j_q(y_n - x^*) \rangle \\ &= \lambda^{q-1} \|y_n - T y_n\|^q \\ &\quad + \langle y_n - T y_n - (x^* - T x^*), j_q(x_n - x^*) - j_q(y_n - x^*) \rangle. \end{aligned}$$

Moreover, by using (1.4), we have

$$(2.5) \quad \begin{aligned} \|y_n - Ty_n\|^q &= \|(1 - \beta_n)(x_n - Ty_n) + \beta_n(Tx_n - Ty_n)\|^q \\ &\geq (1 - \beta_n)\|x_n - Ty_n\|^q + \beta_n\|Tx_n - Ty_n\|^q \\ &\quad - \omega_q(\beta_n)d_q\|x_n - Tx_n\|^q. \end{aligned}$$

Then it follows from (2.2)-(2.5) that

$$\begin{aligned} \|x_{n+1} - x^*\|^q &\leq \|x_n - x^*\|^q - q\alpha_n\{\beta_n\lambda^{q-1}\|x_n - Tx_n\|^q \\ &\quad + \lambda^{q-1}(1 - \beta_n)\|x_n - Ty_n\|^q \\ &\quad + \lambda^{q-1}\beta_n\|Tx_n - Ty_n\|^q - \lambda^{q-1}\omega_q(\beta_n)d_q\|x_n - Tx_n\|^q \\ &\quad + \langle y_n - Ty_n, j_q(x_n - x^*) - j_q(y_n - x^*) \rangle\} \\ &\quad + \alpha_n^q c_q \|x_n - Ty_n\|^q + q\|u_n\|\|x_{n+1} - u_n - x^*\|^{q-1} + c_q\|u_n\|^q \\ &\leq \|x_n - x^*\|^q - \alpha_n(q\lambda^{q-1}(1 - \beta_n) - \alpha_n^{q-1}c_q)\|x_n - Ty_n\|^q \\ &\quad + qd_q\lambda^{q-1}\alpha_n\omega_q(\beta_n)\|x_n - Tx_n\|^q \\ &\quad + q\alpha_n\|y_n - Ty_n\|\|j_q(x_n - x^*) - j_q(y_n - x^*)\| \\ &\quad + q\|u_n\|\|x_{n+1} - u_n - x^*\|^{q-1} + c_q\|u_n\|^q. \end{aligned}$$

Also, observe that

$$\begin{aligned} \omega_q(\beta_n) &= \beta_n(1 - \beta_n)^q + \beta_n^q(1 - \beta_n) \leq 2\beta_n, \\ \|x_n - Tx_n\| &\leq (1 + L)\|x_n - x^*\|, \\ \|j_q(x_n - x^*) - j_q(y_n - x^*)\| &\leq L_*\beta_n^{q-1}\|x_n - Tx_n\|^{q-1} \quad (\text{using (1.5)}) \\ &\leq L_*(1 + L)^{q-1}\beta_n^{q-1}\|x_n - x^*\|^{q-1}, \end{aligned}$$

and

$$\begin{aligned} \|y_n - Ty_n\| &\leq (1 + L)\|y_n - x^*\| \\ &\leq (1 + L)[(1 - \beta_n)\|x_n - x^*\| + \beta_n L\|x_n - x^*\|] \\ &\leq (1 + L)^2\|x_n - x^*\|. \end{aligned}$$

Consequently, we have

$$\begin{aligned} \|x_{n+1} - x^*\|^q &\leq [1 + 2\alpha_n\beta_n\lambda^{q-1}qd_q(1 + L)^q + \alpha_n\beta_n^{q-1}qL_*(1 + L)^{q+1}]\|x_n - x^*\|^2 \\ &\quad - \alpha_n[q\lambda^{q-1}(1 - \beta_n) - \alpha_n^{q-1}c_q]\|x_n - Ty_n\|^q \\ &\quad + q\|u_n\|\|x_{n+1} - u_n - x^*\|^{q-1} + c_q\|u_n\|^q. \end{aligned}$$

Since $\alpha_n \leq \lambda(q/c_q)^{1/(q-1)}$, and

$$\begin{aligned} \|x_n - Ty_n\| &\leq \|x_n - x^*\| + L\|x^* - y_n\| \\ &\leq \|x_n - x^*\| + L[(1 - \beta_n)\|x_n - x^*\| + \beta_n L\|x_n - x^*\|] \\ &\leq (1 + L^2)\|x_n - x^*\|, \end{aligned}$$

we conclude that (i) is valid.

(ii) It follows from conclusion (i) that for all $n, m \geq 1$ and $x^* \in F(T)$

$$\begin{aligned} \|x_{n+m} - x^*\|^q &\leq (1 + \delta_{n+m-1})\|x_{n+m-1} - x^*\|^q + \theta_{n+m-1} \\ &\leq (1 + \delta_{n+m-1})(1 + \delta_{n+m-2})\|x_{n+m-2} - x^*\|^q \\ &\quad + (1 + \delta_{n+m-1})\theta_{n+m-2} + \theta_{n+m-1} \\ &\leq (1 + \delta_{n+m-1})(1 + \delta_{n+m-2})(1 + \delta_{n+m-3})\|x_{n+m-3} - x^*\|^q \\ &\quad + (1 + \delta_{n+m-1})(1 + \delta_{n+m-2})\theta_{n+m-3} + (1 + \delta_{n+m-1})\theta_{n+m-2} \\ &\quad + \theta_{n+m-1} \\ &\leq \dots \\ &\leq e^{\sum_{k=n}^{n+m-1} \delta_k} \|x_n - x^*\|^q + e^{\sum_{k=n}^{n+m-1} \delta_k} \sum_{k=n}^{n+m-1} \theta_k \\ &\leq M\|x_n - x^*\|^q + M \sum_{k=n}^{n+m-1} \theta_k, \end{aligned}$$

where $M = e^{\sum_{k=1}^{\infty} \delta_k}$. This shows that conclusion (ii) is also valid.

Theorem 2.1. *Let $q > 1$ and E be a real q -uniformly smooth Banach space, K be a nonempty closed convex subset of E with $K + K \subseteq K$, and $T : K \rightarrow K$ be a strictly pseudocontractive mapping with $F(T) \neq \emptyset$. Let $\{u_n\}$ be a bounded sequence in K . Let $\{\alpha_n\}$ and $\{\beta_n\}$ be real sequences in $[0, 1]$ satisfying the following conditions:*

- (i) $\sum_{n=1}^{\infty} \|u_n\| < \infty$;
- (ii) $\alpha_n \leq \lambda(q/c_q)^{1/(q-1)}$, and $\sum_{n=1}^{\infty} \beta_n^\tau < \infty$, where $\tau = \min\{1, (q-1)\}$.

Let $\{x_n\}$ be the sequence generated from an arbitrary $x_1 \in K$ by the Ishikawa iterative process (1.6) with errors. Then $\{x_n\}$ converges strongly to a fixed point of T if and only if $\{x_n\}$ is bounded and

$$\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$$

where $d(x_n, F(T))$ is the distance of x_n to set $F(T)$, i.e., $d(x_n, F(T)) = \inf_{u^* \in F(T)} d$

(x_n, u^*) .

Proof. “Necessity”. Suppose that $\{x_n\}$ converges strongly to a fixed point of T , say, $y^* \in F(T)$. Then it is clear that $\{x_n\}$ is bounded. Note that

$$d(x_n, F(T)) = \inf_{u^* \in F(T)} d(x_n, u^*) \leq d(x_n, y^*) \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Therefore,

$$\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$$

“Sufficiency”. Suppose that $\{x_n\}$ is bounded and that $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$. First, from Lemma 2.1(i), we obtain

$$\|x_{n+1} - x^*\|^q \leq (1 + \delta_n)\|x_n - x^*\|^q + \theta_n, \quad n \geq 1, x^* \in F(T),$$

where

$$\delta_n = 2\alpha_n\beta_n\lambda^{q-1}qd_q(1 + L)^q + \alpha_n\beta_n^{q-1}qL_*(1 + L)^{q+1} + \alpha_n\beta_nq\lambda^{q-1}(1 + L)^{2q},$$

and

$$\theta_n = q\|u_n\|\|x_{n+1} - u_n - x^*\|^{q-1} + c_q\|u_n\|^q, \quad \forall n \geq 1.$$

Since $\sum_{n=1}^{\infty} \|u_n\| < \infty$, $\sum_{n=1}^{\infty} \|u_n\|^q < \infty$. Note that $\{x_n\}$ and $\{u_n\}$ are both bounded. Thus, there is a number $\tilde{M} > 0$ such that $\|x_{n+1} - u_n - x^*\| \leq \tilde{M}$, and $\|x_n - x^*\| \leq \tilde{M}, \forall n \geq 1$. Hence,

$$\begin{aligned} \sum_{n=1}^{\infty} \theta_n &= \sum_{n=1}^{\infty} (q\|u_n\|\|x_{n+1} - u_n - x^*\|^{q-1} + c_q\|u_n\|^q) \\ &\leq q\tilde{M}^{q-1} \sum_{n=1}^{\infty} \|u_n\| + c_q \sum_{n=1}^{\infty} \|u_n\|^q < \infty. \end{aligned}$$

On the other hand, it follows from condition (ii) that $\sum_{n=1}^{\infty} \delta_n\tilde{M}^q < \infty$. Also, observe that

$$(2.6) \quad \|x_{n+1} - x^*\|^q \leq (1 + \delta_n)\|x_n - x^*\|^q + \theta_n \leq \|x_n - x^*\|^q + \delta_n\tilde{M}^q + \theta_n.$$

This implies that

$$[d(x_{n+1}, F(T))]^q \leq [d(x_n, F(T))]^q + \delta_n\tilde{M}^q + \theta_n.$$

By Lemma 1.1, we infer that $\lim_{n \rightarrow \infty} [d(x_n, F(T))]^q$ exists, that is, $\lim_{n \rightarrow \infty} d(x_n, F(T))$ exists. Since $\lim_{n \rightarrow \infty} \inf d(x_n, F(T)) = 0$, we have $\liminf_{n \rightarrow \infty} d(x_n, F(T)) = 0$.

Now we claim that $\{x_n\}$ is a Cauchy sequence. Indeed, according to Lemma 2.1(ii), we deduce that there exists a constant $M > 0$ such that

$$\|x_{n+m} - x^*\| \leq M\|x_n - x^*\|^q + M \sum_{k=n}^{n+m+1} \theta_k, \forall n, m \geq 1, x^* \in F(T).$$

Since $\lim_{n \rightarrow \infty} d(x_n, F(T)) = 0$ and $\sum_{n=1}^{\infty} \theta_n < \infty$, for an arbitrary $\varepsilon > 0$, there exists an integer $N_1 \geq 1$ such that for all $n \geq N_1$

$$d(x_n, F(T)) < \left(\frac{\varepsilon}{3M}\right)^{1/q} \cdot \frac{1}{2^{(q-1)/q}}, \text{ and } \sum_{k=n}^{\infty} \theta_k < \frac{\varepsilon}{6M} \cdot \frac{1}{2^{q-1}}.$$

Hence, $d(x_{N_1}, F(T)) < \left(\frac{\varepsilon}{3M}\right)^{1/q} \cdot \frac{1}{2^{(q-1)/q}}$. This implies that there exists an $x_1^* \in F(T)$ such that

$$d(x_{N_1}, x_1^*) < \left(\frac{\varepsilon}{3M}\right)^{1/q} \cdot \frac{1}{2^{(q-1)/q}}.$$

In view of Jensen's Inequality [2, p.183], we conclude that

$$(2.7) \quad \|x_{n+m} - x_n\|^q \leq 2^{q-1} (\|x_n - x_1^*\|^q + \|x_{n+m} - x_1^*\|^q).$$

Since for all $n \geq N_1$, we have

$$\begin{aligned} \|x_n - x_1^*\|^q &\leq M\|x_{N_1} - x_1^*\|^q + M \sum_{k=N_1}^n \theta_k \\ &\leq M\|x_{N_1} - x_1^*\|^q + M \sum_{k=N_1}^{\infty} \theta_k \\ &\leq M \frac{\varepsilon}{3M} \cdot \frac{1}{2^{(q-1)}} + M \frac{\varepsilon}{6M} \cdot \frac{1}{2^{q-1}} \\ &= \frac{\varepsilon}{2} \cdot \frac{1}{2^{q-1}}, \end{aligned}$$

and

$$\begin{aligned} \|x_{n+m} - x_1^*\|^q &\leq M\|x_{N_1} - x_1^*\|^q + M \sum_{k=N_1}^{n+m-1} \theta_k \\ &\leq M\|x_{N_1} - x_1^*\|^q + M \sum_{k=N_1}^{\infty} \theta_k \\ &\leq M \frac{\varepsilon}{3M} \cdot \frac{1}{2^{q-1}} + M \frac{\varepsilon}{6M} \cdot \frac{1}{2^{q-1}} \\ &= \frac{\varepsilon}{2} \cdot \frac{1}{2^{q-1}}, \end{aligned}$$

so, from (2.7), we get

$$\|x_{n+m} - x_n\|^q \leq 2^{q-1} \left(\frac{\varepsilon}{2} \cdot \frac{1}{2^{q-1}} + \frac{\varepsilon}{2} \cdot \frac{1}{2^{q-1}} \right) = \varepsilon, \quad \forall n \geq N_1, m \geq 1.$$

This shows that $\{x_n\}$ is Cauchy sequence. Since the space E is complete, $\lim_{n \rightarrow \infty} x_n$ exists. Thus, we may assume that $\lim_{n \rightarrow \infty} x_n = u^*$.

Next, we claim that u^* is a fixed point of T , i.e., $u^* \in F(T)$. Indeed, since $d(u^*, F(T)) = 0$, for any $\varepsilon > 0$, there is $z^* \in F(T)$ such that $\|u^* - z^*\| < \varepsilon$. Then we have

$$\begin{aligned} \|Tu^* - u^*\| &\leq \|Tu^* - Tz^*\| + \|u^* - z^*\| \\ &\leq (1 + L)\varepsilon. \end{aligned}$$

As ε is arbitrary, we conclude that $Tu^* = u^*$ and hence u^* is a fixed point of T . This completes the proof of Theorem 2.1.

Theorem 2.2. *Let $q > 1$ and E be a real q -uniformly smooth Banach space, K be a nonempty closed convex subset of E with $K + K \subseteq K$, and $T : K \rightarrow K$ be a strictly pseudocontractive mapping with $F(T) \neq \emptyset$. Let $\{u_n\}$ be a bounded sequence in K , and $\{\alpha_n\}, \{\beta_n\}$ be real sequences in $[0, 1]$ satisfying the following conditions:*

- (i) $\sum_{n=1}^{\infty} \|u_n\| < \infty$;
- (ii) $\alpha_n \leq \lambda(q/c_q)^{1/(q-1)}$, and $\sum_{n=1}^{\infty} \beta_n^\tau < \infty$, where $\tau = \min\{1, (q-1)\}$.

Let $\{x_n\}$ be the sequence generated from arbitrary $x_1 \in K$ by the Ishikawa iterative process (1.6) with errors. Then $\{x_n\}$ converges strongly to a fixed point u^* of T if and only if $\{x_n\}$ is bounded and $\{x_n\}$ has a subsequence which is strongly convergent to a fixed point u^* of T .

Proof. The conclusion of Theorem 2.2 follows immediately from Theorem 2.1.

Remark 2.1. If we take $\beta_n = 0, \forall n \geq 1$ in Theorems 2.1 and 2.2, then we can obtain the corresponding results on the strong convergence of the Mann iterative process with errors

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n T x_n + u_n, \quad \forall n \geq 1.$$

In addition, if we take $u_n = 0, \forall n \geq 1$ in (1.6), then Theorems 2.1 and 2.2 are still valid under the lack of the assumption that $\{x_n\}$ is bounded. Indeed, when $u_n = 0, \forall n \geq 1$, it follows from Lemma 2.1(i) that

$$\|x_{n+1} - x^*\|^q \leq (1 + \delta_n) \|x_n - x^*\|^q \leq e^{\sum_{j=1}^n \delta_j} \|x_1 - x^*\|^q \leq e^{\sum_{j=1}^{\infty} \delta_j} \|x_1 - x^*\|^q < \infty.$$

This shows that $\{x_n\}$ is bounded.

Remark 2.2. Recall that Ishikawa iterative process with errors introduced by Liu [3] is stated as follows: Let K be a nonempty convex subset of E with

$K + K \subseteq K$. For any given $x_1 \in K$, the sequence $\{x_n\}$ is defined by the iterative scheme

$$\begin{cases} y_n = (1 - \beta_n)x_n + \beta_nTx_n + v_n, \\ x_{n+1} = (1 - \alpha_n)x_n + \alpha_nTy_n + u_n, n \geq 1, \end{cases}$$

where $\{u_n\}$ and $\{v_n\}$ are bounded sequences in K , and $\{\alpha_n\}$ as well as $\{\beta_n\}$ are real sequences in $[0, 1]$. Naturally, one may ask the following open question.

Open Question: Are Theorems 2.1 and 2.2 extendable to the case of the Ishikawa iterative process with errors in the sense of Liu [3]?

REFERENCES

1. F. E. Browder and W. V. Petryshyn, Construction of fixed point of nonlinear mappings in Hilbert space, *J. Math. Anal. Appl.*, **20** (1967), 197-228.
2. M. Kuczma, *An Introduction to the Theory of Functional Equations and Inequalities: Cauchy's Equation and Jensen's Inequality*, PWN and US: Warszawa-Krakow-Katowice, 1985.
3. L. S. Liu, Ishikawa and Mann iterative process with errors for nonlinear strongly accretive mappings in Banach spaces, *J. Math. Anal. Appl.*, **194** (1995), 114-125.
4. L. Deng and X. P. Ding, Iterative approximation of Lipschitz strictly pseudocontractive mappings in uniformly smooth Banach spaces, *Nonlinear Anal. TMA*, **24** (1995), 981-987.
5. S. S. Chang, Y. J. Cho, B. S. Lee and S. M. Kang, Iterative approximation of fixed points and solutions for strongly accretive and strongly pseudocontractive mapping in Banach spaces, *J. Math. Anal. Appl.*, **224** (1998), 149-165.
6. L. C. Zeng, Iterative approximation of solutions to nonlinear equations of strongly accretive operators in Banach spaces, *Nonlinear Anal. TMA*, **31** (1998), 589-598.
7. L. C. Zeng, Iterative approximation of solutions to nonlinear equations involving m -accretive operators in Banach spaces, *J. Math. Anal. Appl.*, **270** (2002), 319-331.
8. Q. H. Liu, Iterative sequences for asymptotically quasi-nonexpansive mapping with error member, *J. Math. Anal. Appl.*, **259** (2001), 18-24.
9. B. E. Rhoades, Fixed point iterations using infinite matrices, *Trans. Amer. Math. Soc.*, **196** (1974), 161-176.
10. K. K. Tan and H. K. Xu, Approximating fixed points of nonexpansive mappings by the Ishikawa iteration process, *J. Math. Anal. Appl.*, **178** (1993), 301-308.
11. H. K. Xu, Inequalities in Banach spaces with applications, *Nonlinear Anal. TMA*, **116** (1991), 1127-1138.
12. Z. B. Xu and G. F. Roach, Characteristic inequalities of uniformly convex and uniformly smooth Banach spaces, *J. Math. Anal. Appl.*, **157** (1991), 198-210.

13. M. O. Osilike and A. Udomene, Demiclosedness principle and convergence theorems for strictly pseudocontractive mappings of Browder-Petryshyn type, *J. Math. Anal. Appl.*, **256** (2001), 431-445.

Lu-Chuan Zeng
Department of Mathematics,
Shanghai Normal University,
Shanghai 200234, China.

Ngai-Ching Wong
Department of Applied Mathematics,
National Sun Yat-sen University,
Kaohsiung 804, Taiwan.

Jen-Chih Yao
Department of Applied Mathematics,
National Sun Yat-sen University,
Kaohsiung 804, Taiwan.