

On parameter estimation for critical affine processes

Mátyás Barczy^{*,†}

*Faculty of Informatics, University of Debrecen
Pf. 12, H-4010 Debrecen, Hungary
e-mail: barczy.matyas@inf.unideb.hu*

Leif Döring[‡]

*Laboratoire de Probabilités et Modèles Aléatoires
Université Paris 6, 4 place Jussieu, 75252 Paris, France
e-mail: leif.doering@googlemail.com*

Zenghu Li^{*,§}

*School of Mathematical Sciences, Beijing Normal University
Beijing 100875, People's Republic of China
e-mail: lizh@bnu.edu.cn*

and

Gyula Pap^{*,†}

*Bolyai Institute, University of Szeged
Aradi vértanúk tere 1, H-6720 Szeged, Hungary
e-mail: papgy@math.u-szeged.hu*

Abstract: First we provide a simple set of sufficient conditions for the weak convergence of scaled affine processes with state space $\mathbb{R}_+ \times \mathbb{R}^d$. We specialize our result to one-dimensional continuous state branching processes with immigration. As an application, we study the asymptotic behavior of least squares estimators of some parameters of a two-dimensional critical affine diffusion process.

AMS 2000 subject classifications: Primary 60F05, 60J80; secondary 62F12, 91G70.

Keywords and phrases: Affine process, scaling theorem, least squares estimator.

Received October 2012.

*Supported by the Hungarian Chinese Intergovernmental S & T Cooperation Programme for 2011-2013 under Grant No. 10-1-2011-0079.

[†]Partially supported by the Hungarian Scientific Research Fund under Grant No. OTKA T-079128.

[‡]Supported by the Foundation Science *Matématiques de Paris*.

[§]Supported by NSFC under Grant No. 11131003 and 973 Program under Grant No. 2011CB808001.

Contents

| | | |
|---|---|-----|
| 1 | Introduction | 648 |
| 2 | A scaling theorem for affine processes | 649 |
| 3 | Least squares estimator for a critical two-dimensional affine diffusion process | 662 |
| | 3.1 Preparations and (sub)(super)criticality | 662 |
| | 3.2 Least squares estimator of θ when m is known | 667 |
| | 3.3 Least squares estimator of (θ, m) | 676 |
| | 3.4 Conditional least squares estimator of (θ, m) | 679 |
| A | The integrals in (2.2) | 684 |
| B | On consistency properties of the LSE and CLSE of (θ, m) | 687 |
| | References | 694 |

1. Introduction

In recent years quickly growing interest in pricing of credit-risky securities (e.g., defaultable bonds) has been seen in the mathematical finance literature. One of the basic models (for applications see for instance Chen and Joslin [8]) is the following two-dimensional affine diffusion process:

$$\begin{cases} dY_t = (a - bY_t) dt + \sqrt{Y_t} dW_t, \\ dX_t = (m - \theta X_t) dt + \sqrt{Y_t} dB_t, \end{cases} \quad t \geq 0, \quad (1.1)$$

where a, b, θ and m are real parameters such that $a > 0$ and B and W are independent standard Wiener processes. Note that Y is a Cox-Ingersoll-Ross (CIR) process. For practical use, it is important to estimate the appearing parameters from some discretely observed real data set. In the case of the one-dimensional CIR process, the parameter estimation of a and b goes back to Overbeck and Rydén [30], Overbeck [31], and see also the very recent papers of Ben Alaya and Kebaier [5, 6]. For asymptotic results on discrete time critical branching processes with immigration, one may refer to Wei and Winnicki [34] and [35].

The process (Y, X) given by (1.1) is a very special affine process. The set of affine processes contains a large class of important Markov processes such as continuous state branching processes and Orstein-Uhlenbeck processes. Further, a lot of models in financial mathematics are also special affine processes such as the Heston model [17], the model due to Barndorff-Nielsen and Shephard [4] or the model due to Carr and Wu [7]. A precise mathematical formulation and complete characterization of regular affine processes are due to Duffie et al. [11]. Later several authors have contributed to the study of properties of general affine processes: to name a few, Andersen and Piterbarg [1] (moment explosions in stochastic volatility models), Dawson and Li [10] (jump-type SDE representation for two-dimensional affine processes), Filipović and Mayerhofer [13] (applications to the pricing of bond and stock options), Glasserman and Kim

[15] (the range of finite exponential moments and the convergence to stationarity in affine diffusion models), Jena et al. [23] (long-term and blow-up behaviors of exponential moments in multi-dimensional affine diffusions), Keller-Ressel et al. [25, 26] (stochastically continuous, time-homogeneous affine processes with state space $\mathbb{R}_+^n \times \mathbb{R}^d$ or more general ones are regular). We also refer to the overview articles Cuchiero et al. [9] and Friz and Keller-Ressel [14].

To the best knowledge of the authors the parameter estimation problem for multi-dimensional affine processes has not been tackled so far. Since affine processes are being used in financial mathematics very frequently, the question of parameter estimation for them is of high importance. Our aim is to start the discussion with a simple non-trivial example: the two-dimensional affine diffusion process given by (1.1).

The article is divided into two parts and there are two appendices. In Section 2 we recall some notations, the definition of affine processes and some of their basic properties, and then a simple set of sufficient conditions for the weak convergence of scaled affine processes is presented. Roughly speaking, given a family of affine processes $(Y^{(\theta)}(t), X^{(\theta)}(t))_{t \geq 0}$, $\theta > 0$, such that the corresponding admissible parameters converge in an appropriate way (see Theorem 2.2), the scaled process $(\theta^{-1}Y^{(\theta)}(\theta t), \theta^{-1}X^{(\theta)}(\theta t))_{t \geq 0}$ converge weakly towards an affine diffusion process as $\theta \rightarrow \infty$. We specialize our result for one-dimensional continuous state branching processes with immigration which generalizes Theorem 2.3 in Huang et al. [20]. The scaling Theorem 2.2 is proved for quite general affine processes since it might have applications elsewhere later on. In Section 3 the scaling Theorem 2.2 is applied to study the asymptotic behavior of least squares and conditional least squares estimators of some parameters of a critical two-dimensional affine diffusion process given by (1.1), see Theorems 3.1, 3.2 and 3.3. In Appendix A we check that some integrals in the form of the infinitesimal generator of an affine process that we use are well-defined. Appendix B is devoted to show that the least squares estimator of m cannot be asymptotically weakly consistent.

2. A scaling theorem for affine processes

Let \mathbb{N} , \mathbb{Z}_+ , \mathbb{R} , \mathbb{R}_+ , \mathbb{R}_- , \mathbb{R}_{++} , and \mathbb{C} denote the sets of positive integers, non-negative integers, real numbers, non-negative real numbers, non-positive real numbers, positive real numbers and complex numbers, respectively. For $x, y \in \mathbb{R}$, we will use the notations $x \wedge y := \min(x, y)$ and $x \vee y := \max(x, y)$. For $x, y \in \mathbb{C}^k$, $k \in \mathbb{N}$, we write $\langle x, y \rangle := \sum_{i=1}^k x_i y_i$ (notice that this is not the scalar product on \mathbb{C}^k , however for $x \in \mathbb{C}^k$ and $y \in \mathbb{R}^k$, $\langle x, y \rangle$ coincides with the usual scalar product of x and y). By $\|x\|$ and $\|A\|$ we denote the Euclidean norm of a vector $x \in \mathbb{R}^p$ and the induced matrix norm of a matrix $A \in \mathbb{R}^{p \times p}$, respectively. Further, let $U := \{z_1 + iz_2 : z_1 \in \mathbb{R}_-, z_2 \in \mathbb{R}\} \times (i\mathbb{R}^d)$. By $C_c^2(\mathbb{R}_+ \times \mathbb{R}^d)$ ($C_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$) we denote the set of twice (infinitely) continuously differentiable complex-valued functions on $\mathbb{R}_+ \times \mathbb{R}^d$ with compact support, where $d \in \mathbb{N}$. The set of càdlàg functions from \mathbb{R}_+ to $\mathbb{R}_+ \times \mathbb{R}^d$ will

be denoted by $D(\mathbb{R}_+, \mathbb{R}_+ \times \mathbb{R}^d)$. For a bounded function $g : \mathbb{R}_+ \times \mathbb{R}^d \rightarrow \mathbb{R}^p$, let $\|g\|_\infty := \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} \|g(x)\|$. Convergence in distribution, in probability and almost sure convergence will be denoted by $\xrightarrow{\mathcal{L}}$, $\xrightarrow{\mathbb{P}}$ and $\xrightarrow{\text{a.s.}}$, respectively.

Next we briefly recall the definition of affine processes with state space $\mathbb{R}_+ \times \mathbb{R}^d$ based on Duffie et al. [11].

Definition 2.1. A transition semigroup $(P_t)_{t \in \mathbb{R}_+}$ with state space $\mathbb{R}_+ \times \mathbb{R}^d$ is called a (general) affine semigroup if its characteristic function has the representation

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} e^{\langle u, \xi \rangle} P_t(x, d\xi) = e^{\langle x, \psi(t, u) \rangle + \phi(t, u)} \tag{2.1}$$

for $x \in \mathbb{R}_+ \times \mathbb{R}^d$, $u \in U$ and $t \in \mathbb{R}_+$, where $\psi(t, \cdot) = (\psi_1(t, \cdot), \psi_2(t, \cdot)) \in \mathbb{C} \times \mathbb{C}^d$ is a continuous \mathbb{C}^{1+d} -valued function on U and $\phi(t, \cdot)$ is a continuous \mathbb{C} -valued function on U satisfying $\phi(t, 0) = 0$. The affine semigroup $(P_t)_{t \in \mathbb{R}_+}$ defined by (2.1) is called regular if it is stochastically continuous (equivalently, for all $u \in U$, the functions $\mathbb{R}_+ \ni t \mapsto \Psi(t, u)$ and $\mathbb{R}_+ \ni t \mapsto \phi(t, u)$ are continuous) and $\partial_1 \psi(0, u)$ and $\partial_1 \phi(0, u)$ exist for all $u \in U$ and are continuous at $u = 0$ (where $\partial_1 \psi$ and $\partial_1 \phi$ denote the partial derivatives of ψ and ϕ , respectively, with respect to the first variable).

Remark 2.1. We call the attention that Duffie et al. [11] in their Definition 2.1 assume only that Equation (2.1) hold for $x \in \mathbb{R}_+ \times \mathbb{R}^d$, $u \in \partial U = i\mathbb{R}^{1+d}$, $t \in \mathbb{R}_+$, i.e., instead of $u \in U$ they only require that u should be an element of the boundary ∂U of U . However, by Proposition 6.4 in Duffie et al. [11], one can formulate the definition of a regular affine process as we did. Note also that this kind of definition was already given by Dawson and Li [10, Definitions 2.1 and 3.3]. Finally, we remark that every stochastically continuous affine semigroup is regular due to Keller-Ressel et al. [25, Theorem 5.1]. \square

Definition 2.2. A set of parameters $(a, \alpha, b, \beta, m, \mu)$ is called admissible if

- (i) $a = (a_{i,j})_{i,j=1}^{1+d} \in \mathbb{R}^{(1+d) \times (1+d)}$ is a symmetric positive semidefinite matrix with $a_{1,1} = 0$ (hence $a_{1,k} = a_{k,1} = 0$ for all $k \in \{2, \dots, 1+d\}$),
- (ii) $\alpha = (\alpha_{i,j})_{i,j=1}^{1+d} \in \mathbb{R}^{(1+d) \times (1+d)}$ is a symmetric positive semidefinite matrix,
- (iii) $b = (b_i)_{i=1}^{1+d} \in \mathbb{R}_+ \times \mathbb{R}^d$,
- (iv) $\beta = (\beta_{i,j})_{i,j=1}^{1+d} \in \mathbb{R}^{(1+d) \times (1+d)}$ with $\beta_{1,j} = 0$ for all $j \in \{2, \dots, 1+d\}$,
- (v) $m(d\xi) = m(d\xi_1, d\xi_2)$ is a σ -finite measure on $\mathbb{R}_+ \times \mathbb{R}^d$ supported by $(\mathbb{R}_+ \times \mathbb{R}^d) \setminus \{(0, 0)\}$ such that

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} [\xi_1 + (\|\xi_2\| \wedge \|\xi_2\|^2)] m(d\xi) < \infty,$$

- (vi) $\mu(d\xi) = \mu(d\xi_1, d\xi_2)$ is a σ -finite measure on $\mathbb{R}_+ \times \mathbb{R}^d$ supported by $(\mathbb{R}_+ \times \mathbb{R}^d) \setminus \{(0, 0)\}$ such that

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\| \wedge \|\xi\|^2 \mu(d\xi) < \infty.$$

Remark 2.2. Note that our Definition 2.2 of the set of admissible parameters is not so general as Definition 2.6 in Duffie et al. [11]. Firstly, the set of admissible parameters is defined only for affine process with state space $\mathbb{R}_+ \times \mathbb{R}^d$, while Duffie et al. [11] consider affine processes with state space $\mathbb{R}_+^n \times \mathbb{R}^d$. We restrict ourselves to this special case, since our scaling Theorem 2.2 is valid only in this case. Secondly, our conditions (v) and (vi) of Definition 2.2 are stronger than that of (2.10) and (2.11) of Definition 2.6 in Duffie et al. [11]. Thirdly, according to our definition, a set of admissible parameters does not contain parameters corresponding to killing, while in Definition 2.6 in Duffie et al. [11] such parameters are included. Our definition of admissible parameters can be considered as a $(1 + d)$ -dimensional version of Definition 6.1 in Dawson and Li [10]. The reason for this definition is to have a more pleasant form of the infinitesimal generator of an affine process compared to that of Duffie et al. [11, formula (2.12)]. For more details, see Remark 2.3. \square

Theorem 2.1 (Duffie et al. [11, Theorem 2.7]). *Let $(a, \alpha, b, \beta, m, \mu)$ be a set of admissible parameters. Then there exists a unique regular affine semigroup $(P_t)_{t \in \mathbb{R}_+}$ with infinitesimal generator*

$$\begin{aligned}
 (\mathcal{A}f)(x) &= \sum_{i,j=1}^{1+d} (a_{i,j} + \alpha_{i,j}x_1) f''_{i,j}(x) + \langle f'(x), b + \beta x \rangle \\
 &+ \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'_{(2)}(x), \xi_2 \rangle) m(d\xi) \\
 &+ \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'(x), \xi \rangle) x_1 \mu(d\xi)
 \end{aligned} \tag{2.2}$$

for $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $f \in C_c^2(\mathbb{R}_+ \times \mathbb{R}^d)$, where f'_i , $i \in \{1, \dots, 1 + d\}$, and $f''_{i,j}$, $i, j \in \{1, \dots, 1 + d\}$, denote the first and second order partial derivatives of f with respect to its i -th and i -th and j -th variables, and $f'(x) := (f'_1(x), \dots, f'_{1+d}(x))^\top$, $f'_{(2)}(x) := (f'_2(x), \dots, f'_{1+d}(x))^\top$. Further, $C_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ is a core of \mathcal{A} .

Remark 2.3. Note that the form of the infinitesimal generator \mathcal{A} in Theorem 2.1 is slightly different from the one given in (2.12) in Duffie et al. [11]. Our formula (2.2) is in the spirit of Dawson and Li [10, formula (6.5)]. On the one hand, the point is that under the conditions (v) and (vi) of Definition 2.2, one can rewrite (2.12) in Duffie et al. [11] into the form (2.2), by changing the 2-nd, ..., $(1 + d)$ -th coordinates of $b \in \mathbb{R}_+ \times \mathbb{R}^d$ and the first column of $\beta \in \mathbb{R}^{(1+d) \times (1+d)}$, respectively, in appropriate ways (see Appendix A). To see this, it is enough to check that the integrals in (2.2) are well-defined (i.e., elements of \mathbb{C}) under the conditions (v) and (vi) of Definition 2.2. For further details, see also Appendix A. On the other hand, the killing rate (see page 995 in Duffie et al. [11]) of the affine semigroup $(P_t)_{t \in \mathbb{R}_+}$ in Theorem 2.1 is identically zero. This also implies that the affine processes that we will consider later on will have lifetime infinity. \square

Remark 2.4. In dimension 2 (i.e., if $d = 1$), by Theorem 6.2 in Dawson and Li [10] and Theorem 2.7 in Duffie et al. [11] (see also Theorem 2.1), for an

infinitesimal generator \mathcal{A} given by (2.2) with $d = 1$ one can construct a two-dimensional system of jump type SDEs of which there exists a pathwise unique strong solution $(Y(t), X(t))_{t \in \mathbb{R}_+}$ which is a regular affine Markov process with the given infinitesimal generator \mathcal{A} . \square

The next lemma is simple but very useful.

Lemma 2.1. *Let $(Z(t))_{t \in \mathbb{R}_+}$ be a time-homogeneous Markov process with state space $\mathbb{R}_+ \times \mathbb{R}^d$ and let us denote its infinitesimal generator by \mathcal{A}_Z . Suppose that $C_c^2(\mathbb{R}_+ \times \mathbb{R}^d)$ is a subset of the domain of \mathcal{A}_Z . Then for all $\theta \in \mathbb{R}_{++}$, the time-homogeneous Markov process $(Z_\theta(t))_{t \in \mathbb{R}_+} := (\theta^{-1}Z(\theta t))_{t \in \mathbb{R}_+}$ has infinitesimal generator*

$$(\mathcal{A}_{Z_\theta} f)(x) = \theta(\mathcal{A}_Z f_\theta)(\theta x), \quad x \in \mathbb{R}_+ \times \mathbb{R}^d, \quad f \in C_c^2(\mathbb{R}_+ \times \mathbb{R}^d),$$

where $f_\theta(x) := f(\theta^{-1}x)$, $x \in \mathbb{R}_+ \times \mathbb{R}^d$.

Proof. By definition, the infinitesimal generator of $(Z_\theta(t))_{t \in \mathbb{R}_+}$ takes the form

$$\begin{aligned} (\mathcal{A}_{Z_\theta} f)(x) &= \lim_{t \downarrow 0} \frac{\mathbb{E}(f(\theta^{-1}Z(\theta t)) \mid \theta^{-1}Z(0) = x) - f(x)}{t} \\ &= \lim_{t \downarrow 0} \frac{\theta \mathbb{E}(f_\theta(Z(\theta t)) \mid Z(0) = \theta x) - \theta f_\theta(\theta x)}{\theta t} \\ &= \theta \lim_{t' \downarrow 0} \frac{\mathbb{E}(f_\theta(Z(t')) \mid Z(0) = \theta x) - f_\theta(\theta x)}{t'} = \theta(\mathcal{A}_Z f_\theta)(\theta x) \end{aligned}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $f \in C_c^2(\mathbb{R}_+ \times \mathbb{R}^d)$. \square

Theorem 2.2. *For all $\theta \in \mathbb{R}_{++}$, let $(Y^{(\theta)}(t), X^{(\theta)}(t))_{t \in \mathbb{R}_+}$ be a $(1 + d)$ -dimensional affine process with state space $\mathbb{R}_+ \times \mathbb{R}^d$ and with admissible parameters $(a^{(\theta)}, \alpha^{(\theta)}, b^{(\theta)}, \beta^{(\theta)}, m, \mu)$ such that additionally*

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\| m(d\xi) < \infty \quad \text{and} \quad \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\|^2 \mu(d\xi) < \infty. \quad (2.3)$$

Let $a, \alpha, \beta \in \mathbb{R}^{(1+d) \times (1+d)}$, $b \in \mathbb{R}_+ \times \mathbb{R}^d$, and let $(Y(t), X(t))_{t \in \mathbb{R}_+}$ be a $(1 + d)$ -dimensional affine process with state space $\mathbb{R}_+ \times \mathbb{R}^d$ and with the set of admissible parameters $(a, \tilde{\alpha}, \tilde{b}, \beta, 0, 0)$, where

$$\tilde{\alpha} := \alpha + \frac{1}{2} \int_{\mathbb{R}_+ \times \mathbb{R}^d} \xi \xi^\top \mu(d\xi),$$

and $\tilde{b} = (\tilde{b}_i)_{i=1}^{1+d}$ with $\tilde{b}_i := b_i$ for $i \in \{2, \dots, 1 + d\}$ and

$$\tilde{b}_1 := b_1 + \int_{\mathbb{R}_+ \times \mathbb{R}^d} \xi_1 m(d\xi).$$

If

$$\begin{aligned} \theta^{-1} a^{(\theta)} &\rightarrow a, & \alpha^{(\theta)} &\rightarrow \alpha, & b^{(\theta)} &\rightarrow b, & \theta \beta^{(\theta)} &\rightarrow \beta, \\ \theta^{-1} (Y^{(\theta)}(0), X^{(\theta)}(0)) &\xrightarrow{\mathcal{L}} (Y(0), X(0)) \end{aligned}$$

as $\theta \rightarrow \infty$, then

$$\left(Y_\theta^{(\theta)}(t), X_\theta^{(\theta)}(t) \right)_{t \in \mathbb{R}_+} = \left(\theta^{-1} Y^{(\theta)}(\theta t), \theta^{-1} X^{(\theta)}(\theta t) \right)_{t \in \mathbb{R}_+} \xrightarrow{\mathcal{L}} (Y(t), X(t))_{t \in \mathbb{R}_+}$$

in $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+ \times \mathbb{R}^d)$ as $\theta \rightarrow \infty$.

Remark 2.5. (i) Note that the limit process $(Y(t), X(t))_{t \in \mathbb{R}_+}$ in Theorem 2.2 has continuous sample paths almost surely. However, this is not a big surprise, since in condition (2.3) of Theorem 2.2 we require finite second moment for the measure μ .

(ii) Note also that the matrix $\tilde{\alpha} \in \mathbb{R}^{(1+d) \times (1+d)}$ given in Theorem 2.2 is symmetric and positive semidefinite, since α is symmetric and positive semidefinite, and for all $z \in \mathbb{R}^{1+d}$,

$$\left\langle \int_{\mathbb{R}_+ \times \mathbb{R}^d} \xi \xi^\top \mu(d\xi) z, z \right\rangle = \int_{\mathbb{R}_+ \times \mathbb{R}^d} (z^\top \xi)^2 \mu(d\xi) \geq 0.$$

□

Proof of Theorem 2.2. By Duffie et al. [11, Theorem 2.7], $\mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ is a core of the infinitesimal generator $\mathcal{A}_{(Y,X)}$ of the process $(Y(t), X(t))_{t \in \mathbb{R}_+}$, and hence $\{(f, \mathcal{A}_{(Y,X)}f) : f \in \mathcal{D}(\mathcal{A}_{(Y,X)})\}$ coincides with the closure of $\{(f, \mathcal{A}_{(Y,X)}f) : f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)\}$, where $\mathcal{D}(\mathcal{A}_{(Y,X)})$ denotes the domain of $\mathcal{A}_{(Y,X)}$, see, e.g., Ethier and Kurtz [12, page 17]. In other words, the closure of $\{(f, \mathcal{A}_{(Y,X)}f) : f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)\}$ generates the affine semigroup corresponding to $\mathcal{A}_{(Y,X)}$.

Next we show that for all $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$, we have

$$\lim_{\theta \rightarrow \infty} \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} \left| (\mathcal{A}_{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})} f)(x) - (\mathcal{A}_{(Y,X)} f)(x) \right| = 0. \tag{2.4}$$

First note that it is enough to prove (2.4) for real-valued functions $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$, since if (2.4) holds for real-valued functions $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$, then, by decomposing f into real and imaginary parts, the linearity of the infinitesimal generators in question and triangular inequality yield (2.4) for complex-valued functions $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$. Hence in what follows without loss of generality we can assume that $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ is real-valued.

For all $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$, $\theta \in \mathbb{R}_{++}$, and $x \in \mathbb{R}_+ \times \mathbb{R}^d$, we have

$$\begin{aligned} f_\theta(x) &= f(\theta^{-1}x), \\ (f_\theta)'_i(x) &= \theta^{-1} f'_i(\theta^{-1}x), \quad i \in \{1, \dots, 1+d\}, \\ (f_\theta)''_{i,j}(x) &= \theta^{-2} f''_{i,j}(\theta^{-1}x), \quad i, j \in \{1, \dots, 1+d\}. \end{aligned} \tag{2.5}$$

Then, by Lemma 2.1, (2.2) and (2.5),

$$\begin{aligned} (\mathcal{A}_{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})} f)(x) &= \theta (\mathcal{A}_{(Y^{(\theta)}, X^{(\theta)})} f_\theta)(\theta x) \\ &= \theta \left[\sum_{i,j=1}^{1+d} (a_{i,j}^{(\theta)} + \alpha_{i,j}^{(\theta)} \theta x_1) \theta^{-2} f''_{i,j}(\theta^{-1}\theta x) + \langle \theta^{-1} f'(\theta^{-1}\theta x), b^{(\theta)} + \beta^{(\theta)} \theta x \rangle \right] \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(\theta^{-1}(\theta x + \xi)) - f(\theta^{-1}\theta x) - \langle \theta^{-1}f'_{(2)}(\theta^{-1}\theta x), \xi_2 \rangle) m(d\xi) \\
& + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(\theta^{-1}(\theta x + \xi)) - f(\theta^{-1}\theta x) - \langle \theta^{-1}f'(\theta^{-1}\theta x), \xi \rangle) \theta x_1 \mu(d\xi) \Big] \\
& = \sum_{i,j=1}^{1+d} (\theta^{-1}a_{i,j}^{(\theta)} + \alpha_{i,j}^{(\theta)} x_1) f''_{i,j}(x) + \langle f'(x), b^{(\theta)} + \theta\beta^{(\theta)}x \rangle \\
& + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \theta^{-1}\xi) - f(x) - \langle f'_{(2)}(x), \theta^{-1}\xi_2 \rangle) \theta m(d\xi) \\
& + x_1 \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \theta^{-1}\xi) - f(x) - \langle f'(x), \theta^{-1}\xi \rangle) \theta^2 \mu(d\xi)
\end{aligned}$$

for $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ and $x \in \mathbb{R}_+ \times \mathbb{R}^d$. Hence, for all $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$, using the triangular inequality and that $|\langle u, v \rangle| \leq \|u\| \|v\|$, $u, v \in \mathbb{R}^p$, we have

$$\begin{aligned}
& \left| (\mathcal{A}_{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})} f)(x) - (\mathcal{A}_{(Y, X)} f)(x) \right| \\
& \leq \sum_{i,j=1}^{1+d} (|\theta^{-1}a_{i,j}^{(\theta)} - a_{i,j}| + |\alpha_{i,j}^{(\theta)} - \alpha_{i,j}| x_1) |f''_{i,j}(x)| \\
& \quad + (\|b^{(\theta)} - b\| + \|\theta\beta^{(\theta)} - \beta\| \|x\|) \|f'(x)\| \\
& \quad + \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \theta^{-1}\xi) - f(x) - \theta^{-1}\langle f'(x), \xi \rangle) \theta m(d\xi) \right| \\
& \quad + x_1 \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \theta^{-1}\xi) - f(x) - \theta^{-1}\langle f'(x), \xi \rangle \right. \\
& \quad \quad \left. - \frac{1}{2}\theta^{-2}\langle f''(x)\xi, \xi \rangle) \theta^2 \mu(d\xi) \right|,
\end{aligned}$$

where

$$f''(x) := \begin{bmatrix} f''_{1,1}(x) & \cdots & f''_{1,1+d}(x) \\ \vdots & \ddots & \vdots \\ f''_{1+d,1}(x) & \cdots & f''_{1+d,1+d}(x) \end{bmatrix}.$$

Since $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$, we have

$$\begin{aligned}
\sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 |f''_{i,j}(x)| < \infty, & \quad \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} |f''_{i,j}(x)| < \infty, & \quad \forall i, j \in \{1, \dots, 1+d\}, \\
\sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} \|x\| \|f'(x)\| < \infty, & \quad \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} \|f'(x)\| < \infty,
\end{aligned}$$

and hence, by our assumptions, in order to prove (2.4) it is enough to check that

$$\lim_{\theta \rightarrow \infty} \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left(f(x + \theta^{-1}\xi) - f(x) - \theta^{-1} \langle f'(x), \xi \rangle \right) \theta m(d\xi) \right| = 0, \tag{2.6}$$

and

$$\lim_{\theta \rightarrow \infty} \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left(f(x + \theta^{-1}\xi) - f(x) - \theta^{-1} \langle f'(x), \xi \rangle - \frac{1}{2} \theta^{-2} \langle f''(x)\xi, \xi \rangle \right) \theta^2 \mu(d\xi) \right| = 0. \tag{2.7}$$

First we consider (2.6). Let $\varepsilon \in \mathbb{R}_{++}$ be fixed. Let us choose an $M \in \mathbb{R}_{++}$ such that

$$2\|f'\|_\infty \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus ([0, M] \times [-M, M]^d)} \|\xi\| m(d\xi) < \frac{\varepsilon}{2}. \tag{2.8}$$

In what follows, for abbreviation, $[0, M] \times [-M, M]^d$ will be denoted by D_M . Such an M can be chosen, since $f \in \mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ yields $\|f'\|_\infty < \infty$ and, by assumption (2.3), $\int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\| m(d\xi) < \infty$. By Taylor's theorem, for all $\theta \in \mathbb{R}_{++}$, $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in D_M$ there exists some $\tau = \tau(\theta, x, \xi) \in [0, 1]$ such that

$$f(x + \theta^{-1}\xi) - f(x) = \langle f'(x + \theta^{-1}\tau\xi), \theta^{-1}\xi \rangle. \tag{2.9}$$

Then

$$\begin{aligned} & \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left(f(x + \theta^{-1}\xi) - f(x) - \theta^{-1} \langle f'(x), \xi \rangle \right) \theta m(d\xi) \right| \\ & \leq \int_{\mathbb{R}_+ \times \mathbb{R}^d} |\langle f'(x + \theta^{-1}\tau\xi), \xi \rangle - \langle f'(x), \xi \rangle| m(d\xi) \leq A_{\theta, M}^{(1)}(x) + A_{\theta, M}^{(2)}(x) \end{aligned}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$, where

$$\begin{aligned} A_{\theta, M}^{(1)}(x) & := \int_{D_M} |\langle f'(x + \theta^{-1}\tau\xi) - f'(x), \xi \rangle| m(d\xi), \\ A_{\theta, M}^{(2)}(x) & := \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} (|\langle f'(x + \theta^{-1}\tau\xi), \xi \rangle| + |\langle f'(x), \xi \rangle|) m(d\xi). \end{aligned}$$

Here

$$\begin{aligned} A_{\theta, M}^{(1)}(x) & \leq \int_{D_M} \|f'(x + \theta^{-1}\tau\xi) - f'(x)\| \|\xi\| m(d\xi) \\ & \leq \sup_{\xi \in D_M} \|f'(x + \theta^{-1}\tau\xi) - f'(x)\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\| m(d\xi). \end{aligned}$$

The convexity of D_M implies $\tau\xi = \tau(\theta, x, \xi)\xi \in D_M$ for all $\theta \in \mathbb{R}_{++}$, $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in D_M$, and, hence,

$$\sup_{\xi \in D_M} \|f'(x + \theta^{-1}\tau\xi) - f'(x)\| \leq \sup_{\tilde{\xi} \in D_M} \|f'(x + \theta^{-1}\tilde{\xi}) - f'(x)\|.$$

Since f' is uniformly continuous on $\mathbb{R}_+ \times \mathbb{R}^d$ (which follows by mean value theorem using also that $\|f''\|_\infty < \infty$), there exists a $\theta_0 \in \mathbb{R}_{++}$ (depending on ε and M) such that

$$\sup_{\tilde{\xi} \in D_M} \|f'(x + \theta^{-1}\tilde{\xi}) - f'(x)\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\| m(d\xi) < \frac{\varepsilon}{2}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\theta \in [\theta_0, \infty)$. Further, by (2.8), we have

$$A_{\theta, M}^{(2)}(x) \leq 2\|f'\|_\infty \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} \|\xi\| m(d\xi) < \frac{\varepsilon}{2}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\theta \in \mathbb{R}_{++}$. Putting the pieces together we have (2.6).

Now we turn to prove (2.7) in a similar way. Let $\varepsilon \in \mathbb{R}_{++}$ be fixed again. Let us now choose an $M \in \mathbb{R}_{++}$ such that

$$2 \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 \|f''(x)\| \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} \|\xi\|^2 \mu(d\xi) < \frac{\varepsilon}{2}. \tag{2.10}$$

Such an M can be chosen, since $\sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 \|f''(x)\| < \infty$ for all $f \in C_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ and, by assumption (2.3), $\int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\|^2 \mu(d\xi) < \infty$. By Taylor's theorem, for all $\theta \in \mathbb{R}_{++}$, $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in D_M$ there exists some $\tau = \tau(\theta, x, \xi) \in [0, 1]$ such that

$$f(x + \theta^{-1}\xi) - f(x) - \langle f'(x), \theta^{-1}\xi \rangle = \frac{1}{2} \langle f''(x + \theta^{-1}\tau\xi) \theta^{-1}\xi, \theta^{-1}\xi \rangle. \tag{2.11}$$

Then

$$\begin{aligned} & x_1 \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left(f(x + \theta^{-1}\xi) - f(x) - \theta^{-1} \langle f'(x), \xi \rangle - \frac{1}{2} \theta^{-2} \langle f''(x) \xi, \xi \rangle \right) \theta^2 \mu(d\xi) \right| \\ & \leq \frac{1}{2} x_1 \int_{\mathbb{R}_+ \times \mathbb{R}^d} \left| \langle f''(x + \theta^{-1}\tau\xi) \xi, \xi \rangle - \langle f''(x) \xi, \xi \rangle \right| \mu(d\xi) \\ & \leq \frac{1}{2} (B_{\theta, M}^{(1)}(x) + B_{\theta, M}^{(2)}(x)) \end{aligned}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$, where

$$\begin{aligned} B_{\theta, M}^{(1)}(x) & := x_1 \int_{D_M} |\langle (f''(x + \theta^{-1}\tau\xi) - f''(x)) \xi, \xi \rangle| \mu(d\xi), \\ B_{\theta, M}^{(2)}(x) & := x_1 \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} (|\langle f''(x + \theta^{-1}\tau\xi) \xi, \xi \rangle| + |\langle f''(x) \xi, \xi \rangle|) \mu(d\xi). \end{aligned}$$

Here

$$\begin{aligned} B_{\theta, M}^{(1)}(x) &\leq x_1 \int_{D_M} \|f''(x + \theta^{-1}\tau\xi) - f''(x)\| \|\xi\|^2 \mu(d\xi) \\ &\leq \sup_{\xi \in D_M} x_1 \|f''(x + \theta^{-1}\tau\xi) - f''(x)\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\|^2 \mu(d\xi), \end{aligned}$$

and note that $\|A - B\| \geq \|A\| - \|B\|$, $A, B \in \mathbb{R}^{p \times p}$, yields that

$$\begin{aligned} x_1 \|f''(x + \theta^{-1}\tau\xi) - f''(x)\| &\leq \|(x_1 + \theta^{-1}\tau\xi_1)f''(x + \theta^{-1}\tau\xi) - x_1 f''(x)\| \\ &\quad + \theta^{-1}\tau\xi_1 \|f''(x + \theta^{-1}\tau\xi)\|. \end{aligned}$$

Further, we have again

$$\{\tau\xi = \tau(\theta, x, \xi)\xi : \theta \in \mathbb{R}_{++}, x \in \mathbb{R}_+ \times \mathbb{R}^d \text{ and } \xi \in D_M\} \subset D_M,$$

and hence

$$\begin{aligned} &\sup_{\xi \in D_M} \|(x_1 + \theta^{-1}\tau\xi_1)f''(x + \theta^{-1}\tau\xi) - x_1 f''(x)\| \\ &\leq \sup_{\tilde{\xi} \in D_M} \|(x_1 + \theta^{-1}\tilde{\xi}_1)f''(x + \theta^{-1}\tilde{\xi}) - x_1 f''(x)\|. \end{aligned}$$

Since the function $x \mapsto x_1 f''(x)$ is uniformly continuous on $\mathbb{R}_+ \times \mathbb{R}^d$, there exists a $\theta_1 \in \mathbb{R}_{++}$ (depending on ε and M) such that

$$\sup_{\tilde{\xi} \in D_M} \|(x_1 + \theta^{-1}\tilde{\xi}_1)f''(x + \theta^{-1}\tilde{\xi}) - x_1 f''(x)\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\|^2 \mu(d\xi) < \frac{\varepsilon}{4}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\theta \in [\theta_1, \infty)$. Moreover, there exists a $\theta_2 \in \mathbb{R}_{++}$ (depending on ε and M) such that

$$\begin{aligned} \theta^{-1} \sup_{\xi \in D_M} \tau\xi_1 \|f''(x + \theta^{-1}\tau\xi)\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\|^2 \mu(d\xi) \\ \leq \theta^{-1} \|f''\|_\infty \sup_{\xi \in D_M} \xi_1 \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\|^2 \mu(d\xi) < \frac{\varepsilon}{4} \end{aligned}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\theta \in [\theta_2, \infty)$. Consequently, $B_{\theta, M}^{(1)}(x) < \frac{\varepsilon}{2}$ for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\theta \in [\theta_1 + \theta_2, \infty)$. Further,

$$\begin{aligned} B_{\theta, M}^{(2)}(x) &\leq x_1 \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} (\|f''(x + \theta^{-1}\tau\xi)\| + \|f''(x)\|) \|\xi\|^2 \mu(d\xi) \\ &\leq \sup_{\xi \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 (\|f''(x + \theta^{-1}\tau\xi)\| + \|f''(x)\|) \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} \|\xi\|^2 \mu(d\xi). \end{aligned}$$

Here

$$x_1 (\|f''(x + \theta^{-1}\tau\xi)\| + \|f''(x)\|) \leq (x_1 + \theta^{-1}\tau\xi_1) \|f''(x + \theta^{-1}\tau\xi)\| + x_1 \|f''(x)\|,$$

hence

$$\sup_{\xi \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 (\|f''(x + \theta^{-1}\tau\xi)\| + \|f''(x)\|) \leq 2 \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 \|f''(x)\|.$$

Consequently, by (2.10), we have

$$B_{\theta, M}^{(2)}(x) \leq 2 \sup_{x \in \mathbb{R}_+ \times \mathbb{R}^d} x_1 \|f''(x)\| \int_{(\mathbb{R}_+ \times \mathbb{R}^d) \setminus D_M} \|\xi\|^2 \mu(d\xi) < \frac{\varepsilon}{2}$$

for all $x \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\theta \in \mathbb{R}_{++}$. Putting the pieces together we have (2.7).

Finally, Ethier and Kurtz [12, Corollary 8.7 on page 232] yields our assertion. Namely, with the notations of part (f) of this corollary (but replacing n by θ), let

- $G_\theta := E_\theta := E := \mathbb{R}_+ \times \mathbb{R}^d$ for all $\theta \in \mathbb{R}_{++}$,
- $C_a := C_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ which strongly separates points in $\mathbb{R}_+ \times \mathbb{R}^d$ (indeed, for every $(x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\delta \in \mathbb{R}_{++}$, the bump function defined by $h_1(u_1, u_2) := \exp\{-\frac{1}{1-(u_1-x_1)^2} - \frac{1}{1-\|u_2-x_2\|^2}\}$ if $|u_1 - x_1| < 1$ and $\|u_2 - x_2\| < 1$ with $(u_1, u_2) \in \mathbb{R}_+ \times \mathbb{R}^d$, and $h_1(u_1, u_2) := 0$ otherwise, satisfies (4.7) on page 113 in Ethier and Kurtz [12]),
- $\eta_\theta : E_\theta \rightarrow E$ with $\eta_\theta(x_1, x_2) := (x_1, x_2)$, $(x_1, x_2) \in E_\theta$ for all $\theta \in \mathbb{R}_{++}$,
- $\pi_\theta : E \rightarrow E_\theta$ with $\pi_\theta(x_1, x_2) := (x_1, x_2)$, $(x_1, x_2) \in E$ for all $\theta \in \mathbb{R}_{++}$,
- for each $f \in C_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$ one can choose $f_\theta := f$ and $g_\theta := \mathcal{A}_{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})} f$ for all $\theta \in \mathbb{R}_{++}$ (and hence $(f_\theta, g_\theta) \in \widehat{\mathcal{A}}_{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})}$ defined on page 24 in Ethier and Kurtz [12] by part (c) of Proposition 1.5 on page 9 in Ethier and Kurtz [12]),
- $(\mathcal{G}_t^\theta)_{t \in \mathbb{R}_+} := (\mathcal{F}_t^{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})})_{t \in \mathbb{R}_+}$, where $\mathcal{F}_t^{(Y_\theta^{(\theta)}, X_\theta^{(\theta)})}$ denotes the σ -algebra generated by $\{(Y_\theta^{(\theta)}(s), X_\theta^{(\theta)}(s)), s \in [0, t]\}$.

Then, by our assumptions, convergence of the initial distributions holds, condition (8.35) on page 232 in Ethier and Kurtz [12] is automatically satisfied, and (2.4) shows the validity of condition (8.36) on page 232 in Ethier and Kurtz [12]. □

Remark 2.6. By giving an example, we shed some light on why we consider only $(1+d)$ -dimensional affine processes with state space $\mathbb{R}_+ \times \mathbb{R}^d$ in Theorem 2.2 instead of $(n+d)$ -dimensional ones with state space $\mathbb{R}_+^n \times \mathbb{R}^d$, $n \in \mathbb{N}$. Let $(Y_t)_{t \in \mathbb{R}_+}$ be a two-dimensional continuous state branching process with infinitesimal generator

$$(\mathcal{A}_Y f)(y) = \sum_{i=1}^2 y_i \int_{\mathbb{R}_+^2 \setminus \{0\}} (f(y+u) - f(y) - f'_i(y)u_i) p_i(du),$$

for $f \in C_c^2(\mathbb{R}_+^2)$ and $y = (y_1, y_2) \in \mathbb{R}_+^2$, where $p_i, i = 1, 2$, are σ -finite

measures on $\mathbb{R}_+^2 \setminus \{0\}$ such that

$$\int_{\mathbb{R}_+^2 \setminus \{0\}} (u_1 + \|u\|^2) p_2(du) < \infty \quad \text{and} \quad \int_{\mathbb{R}_+^2 \setminus \{0\}} (u_2 + \|u\|^2) p_1(du) < \infty, \tag{2.12}$$

see, e.g., Duffie et al. [11, Theorem 2.7]. Note that Y can be considered as a two-dimensional affine process with state space \mathbb{R}_+^2 (formally with $d = 0$). Then, by a simple modification of Lemma 2.1, for all $\theta > 0$, $f \in \mathcal{C}_c^2(\mathbb{R}_+^2)$ and $y = (y_1, y_2) \in \mathbb{R}_+^2$,

$$\begin{aligned} (\mathcal{A}_{Y_\theta} f)(y) &= \theta (\mathcal{A}_Y f_\theta)(\theta y) \\ &= \theta \sum_{i=1}^2 \theta y_i \int_{\mathbb{R}_+^2 \setminus \{0\}} \left(f(\theta^{-1}(\theta y + u)) - f(\theta^{-1}\theta y) - \theta^{-1} f'_i(\theta^{-1}\theta y) u_i \right) p_i(du) \\ &= \theta^2 \sum_{i=1}^2 y_i \int_{\mathbb{R}_+^2 \setminus \{0\}} \left(f(y + \theta^{-1}u) - f(y) - \langle f'(y), \theta^{-1}u \rangle \right) p_i(du) \\ &\quad + \theta \sum_{i=1}^2 y_i f'_{3-i}(y) \int_{\mathbb{R}_+^2 \setminus \{0\}} u_{3-i} p_i(du), \end{aligned}$$

where the last equality follows by (2.12). Supposing that f is real-valued, by Taylor's theorem,

$$\begin{aligned} f(y + \theta^{-1}u) - f(y) - \langle f'(y), \theta^{-1}u \rangle &= \frac{1}{2} \langle f''(y + \tau\theta^{-1}u) \theta^{-1}u, \theta^{-1}u \rangle \\ &= \frac{\theta^{-2}}{2} \langle f''(y + \tau\theta^{-1}u) u, u \rangle \end{aligned}$$

with some $\tau = \tau(u, y) \in [0, 1]$. Hence, similarly to the proof of (2.7), we get

$$\begin{aligned} \lim_{\theta \rightarrow \infty} \theta^2 \sum_{i=1}^2 y_i \int_{\mathbb{R}_+^2 \setminus \{0\}} \left(f(y + \theta^{-1}u) - f(y) - \langle f'(y), \theta^{-1}u \rangle \right) p_i(du) \\ = \frac{1}{2} \sum_{i=1}^2 y_i \int_{\mathbb{R}_+^2 \setminus \{0\}} \langle f''(y) u, u \rangle p_i(du) \end{aligned}$$

for real-valued $f \in \mathcal{C}_c^2(\mathbb{R}_+^2)$ and $y = (y_1, y_2) \in \mathbb{R}_+^2$. However, $(\mathcal{A}_{Y_\theta} f)(y)$ does not converge as $\theta \rightarrow \infty$ provided that

$$\sum_{i=1}^2 y_i f'_{3-i}(y) \int_{\mathbb{R}_+^2 \setminus \{0\}} u_{3-i} p_i(du) \neq 0.$$

We also note that this phenomena is somewhat similar to that of Remark 2.1 in Ma [28]. □

In the next remark we formulate some special cases of Theorem 2.2.

Remark 2.7. (i) If $(Y(t), X(t))_{t \in \mathbb{R}_+}$ is a $(1 + d)$ -dimensional affine process on $\mathbb{R}_+ \times \mathbb{R}^d$ with admissible parameters $(a, \alpha, b, 0, m, \mu)$ such that condition (2.3) is satisfied, then the conditions of Theorem 2.2 are satisfied for $(Y^{(\theta)}(t), X^{(\theta)}(t))_{t \in \mathbb{R}_+} := (Y(t), X(t))_{t \in \mathbb{R}_+}$, $\theta \in \mathbb{R}_{++}$, and hence

$$(\theta^{-1}Y(\theta t), \theta^{-1}X(\theta t))_{t \in \mathbb{R}_+} \xrightarrow{\mathcal{L}} (\mathcal{Y}(t), \mathcal{X}(t))_{t \in \mathbb{R}_+} \quad \text{as } \theta \rightarrow \infty$$

in $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+ \times \mathbb{R}^d)$, where $(\mathcal{Y}(t), \mathcal{X}(t))_{t \in \mathbb{R}_+}$ is a $(1 + d)$ -dimensional affine process on $\mathbb{R}_+ \times \mathbb{R}^d$ with admissible parameters $(0, \tilde{\alpha}, \tilde{b}, 0, 0, 0)$, where $\tilde{\alpha}$ and \tilde{b} are given in Theorem 2.2.

(ii) If $(Y(t), X(t))_{t \in \mathbb{R}_+}$ is a $(1 + d)$ -dimensional affine process on $\mathbb{R}_+ \times \mathbb{R}^d$ with $(Y(0), X(0)) = (0, 0)$ and with admissible parameters $(0, \alpha, b, 0, 0, 0)$, then

$$(\theta^{-1}Y(\theta t), \theta^{-1}X(\theta t))_{t \in \mathbb{R}_+} \stackrel{\mathcal{L}}{=} (Y(t), X(t))_{t \in \mathbb{R}_+} \quad \text{for all } \theta \in \mathbb{R}_{++},$$

where $\stackrel{\mathcal{L}}{=}$ denotes equality in distribution. Indeed, by Proposition 1.6 on page 161 in Ethier and Kurtz [12], it is enough to check that the semigroups (on the Banach space of bounded Borel measurable functions on $\mathbb{R}_+ \times \mathbb{R}^d$) corresponding to the processes in question coincide. By the definition of a core, this follows from the equality of the infinitesimal generators of the processes in question on the core $\mathcal{C}_c^\infty(\mathbb{R}_+ \times \mathbb{R}^d)$, which has been shown in the proof of Theorem 2.2. \square

Next we present a corollary of Theorem 2.2 which states weak convergence of appropriately normalized one-dimensional continuous state branching processes with immigration. Our corollary generalizes Theorem 2.3 in Huang et al. [20] in the sense that we do not have to suppose that $\int_1^\infty \xi^2 m(d\xi) < \infty$, only that $\int_1^\infty \xi m(d\xi) < \infty$ (with the notations of Huang et al. [20]), and our proof defers from that of Huang et al. [20].

Corollary 2.1. *For all $\theta \in \mathbb{R}_{++}$, let $(Y^{(\theta)}(t))_{t \in \mathbb{R}_+}$ be a one-dimensional continuous state branching process with immigration on \mathbb{R}_+ with branching mechanism*

$$R^{(\theta)}(z) := \beta^{(\theta)}z + \alpha^{(\theta)}z^2 + \int_{\mathbb{R}_+} (e^{-zu} - 1 + zu) p(du), \quad z \in \mathbb{R}_+,$$

and with immigration mechanism

$$F^{(\theta)}(z) := b^{(\theta)}z + \int_{\mathbb{R}_+} (1 - e^{-zu}) n(du), \quad z \in \mathbb{R}_+,$$

where $\alpha^{(\theta)} \geq 0$, $b^{(\theta)} \geq 0$, $\beta^{(\theta)} \in \mathbb{R}$ and n and p are measures on $(0, \infty)$ such that

$$\int_{\mathbb{R}_+} u n(du) < \infty \quad \text{and} \quad \int_{\mathbb{R}_+} u^2 p(du) < \infty.$$

Let $\alpha, b, \beta \in \mathbb{R}$, and let $(Y(t))_{t \in \mathbb{R}_+}$ be a one-dimensional continuous state branching process with immigration on \mathbb{R}_+ with branching mechanism

$$R(z) := -\beta z + \left(\alpha + \frac{1}{2} \int_{\mathbb{R}_+} u^2 p(du) \right) z^2, \quad z \in \mathbb{R}_+,$$

and with immigration mechanism

$$F(z) := \left(b + \int_{\mathbb{R}_+} u n(du) \right) z, \quad z \in \mathbb{R}_+.$$

If

$$\lim_{\theta \rightarrow \infty} \alpha^{(\theta)} = \alpha, \quad \lim_{\theta \rightarrow \infty} b^{(\theta)} = b, \quad \lim_{\theta \rightarrow \infty} \theta \beta^{(\theta)} = \beta, \quad Y^{(\theta)}(0) \xrightarrow{\mathcal{L}} Y(0)$$

as $\theta \rightarrow \infty$, then

$$\left(\theta^{-1} Y^{(\theta)}(\theta t) \right)_{t \in \mathbb{R}_+} \xrightarrow{\mathcal{L}} (Y(t))_{t \in \mathbb{R}_+} \quad \text{as } \theta \rightarrow \infty$$

in $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+)$.

Proof. For each $\theta \in \mathbb{R}_{++}$ and $t \in \mathbb{R}_+$, let $X^{(\theta)}(t) := 0$. Then for each $\theta \in \mathbb{R}_{++}$, the process $(Y^{(\theta)}(t), X^{(\theta)}(t))_{t \in \mathbb{R}_+}$ is a two-dimensional affine process with admissible parameters $(0, \bar{\alpha}^{(\theta)}, \bar{b}^{(\theta)}, \bar{\beta}^{(\theta)}, m, \mu)$, where

$$\begin{aligned} \bar{\alpha}^{(\theta)} &:= \begin{bmatrix} \alpha^{(\theta)} & 0 \\ 0 & 0 \end{bmatrix}, & \bar{b}^{(\theta)} &:= \begin{bmatrix} b^{(\theta)} \\ 0 \end{bmatrix}, & \bar{\beta}^{(\theta)} &:= \begin{bmatrix} \beta^{(\theta)} & 0 \\ 0 & 0 \end{bmatrix}, \\ \mu(d\xi) &= \mu(d\xi_1, d\xi_2) := p(d\xi_1) \times \delta_0(d\xi_2), \\ m(d\xi) &= m(d\xi_1, d\xi_2) := n(d\xi_1) \times \delta_0(d\xi_2), \end{aligned}$$

where δ_0 denotes the Dirac measure concentrated on $0 \in \mathbb{R}$. Then, by Theorem 2.2, for the two-dimensional affine processes $(Y^{(\theta)}(t), X^{(\theta)}(t))_{t \in \mathbb{R}_+}$, $\theta \in \mathbb{R}_{++}$, we have

$$\left(\theta^{-1} Y^{(\theta)}(\theta t), \theta^{-1} X^{(\theta)}(\theta t) \right)_{t \in \mathbb{R}_+} \xrightarrow{\mathcal{L}} (Y(t), X(t))_{t \in \mathbb{R}_+} \quad \text{as } \theta \rightarrow \infty$$

in $\mathcal{D}(\mathbb{R}_+, \mathbb{R}_+ \times \mathbb{R})$, where $(Y(t), X(t))_{t \in \mathbb{R}_+}$ is a two-dimensional affine process on $\mathbb{R}_+ \times \mathbb{R}$ with infinitesimal generator

$$\begin{aligned} (\mathcal{A}_{(Y,X)} f)(x) &= \left(\alpha + \frac{1}{2} \int_{\mathbb{R}_+} u^2 p(du) \right) x_1 f''_{1,1}(x) \\ &\quad + \left(b + \beta x_1 + \int_{\mathbb{R}_+} u n(du) \right) f'_1(x), \end{aligned}$$

for $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}$ and $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R})$. Note that in fact X is the identically zero process. Finally, Theorem 9.30 in Li [27] yields the assertion. \square

3. Least squares estimator for a critical two-dimensional affine diffusion process

In this section continuous time stochastic processes will be written as $(\xi_t)_{t \in \mathbb{R}_+}$ instead of $(\xi(t))_{t \in \mathbb{R}_+}$. Let $(\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in \mathbb{R}_+}, \mathbb{P})$ be a filtered probability space satisfying the usual conditions, i.e., $(\Omega, \mathcal{F}, \mathbb{P})$ is complete, the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ is right-continuous and \mathcal{F}_0 contains all the \mathbb{P} -null sets in \mathcal{F} . Let $(W_t)_{t \in \mathbb{R}_+}$ and $(B_t)_{t \in \mathbb{R}_+}$ be independent standard $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Wiener processes. Let us consider the following two-dimensional diffusion process given by the SDE

$$\begin{cases} dY_t = (a - bY_t) dt + \sqrt{Y_t} dW_t, \\ dX_t = (m - \theta X_t) dt + \sqrt{Y_t} dB_t, \end{cases} \quad t \in \mathbb{R}_+, \quad (3.1)$$

where $a \in \mathbb{R}_{++}$ and $b, \theta, m \in \mathbb{R}$.

3.1. Preparations and (sub)(super)criticality

The next proposition is about the existence and uniqueness of a strong solution of the SDE (3.1).

Proposition 3.1. *Let (η, ζ) be a random vector independent of $(W_t, B_t)_{t \in \mathbb{R}_+}$ satisfying $\mathbb{P}(\eta \geq 0) = 1$. Then, for all $a \in \mathbb{R}_{++}$ and $b, m, \theta \in \mathbb{R}$, there is a (pathwise) unique strong solution $(Y_t, X_t)_{t \in \mathbb{R}_+}$ of the SDE (3.1) such that $\mathbb{P}((Y_0, X_0) = (\eta, \zeta)) = 1$ and $\mathbb{P}(Y_t \geq 0 \text{ for all } t \in \mathbb{R}_+) = 1$. Further, for all $0 \leq s \leq t$,*

$$Y_t = e^{-b(t-s)} \left(Y_s + a \int_s^t e^{-b(s-u)} du + \int_s^t e^{-b(s-u)} \sqrt{Y_u} dW_u \right), \quad (3.2)$$

and

$$X_t = e^{-\theta(t-s)} \left(X_s + m \int_s^t e^{-\theta(s-u)} du + \int_s^t e^{-\theta(s-u)} \sqrt{Y_u} dB_u \right). \quad (3.3)$$

Proof. By Ikeda and Watanabe [21, Example 8.2, page 221], there is a pathwise unique non-negative strong solution $(Y_t)_{t \in \mathbb{R}_+}$ of the first equation in (3.1) with any initial value η satisfying $\mathbb{P}(\eta \geq 0) = 1$. Next, by applications of the Itô's formula to the processes $(e^{bt} Y_t)_{t \in \mathbb{R}_+}$ and $(e^{\theta t} X_t)_{t \in \mathbb{R}_+}$, respectively, we have

$$\begin{aligned} d(e^{bt} Y_t) &= b e^{bt} Y_t dt + e^{bt} dY_t = b e^{bt} Y_t dt + e^{bt} ((a - bY_t) dt + \sqrt{Y_t} dW_t) \\ &= a e^{bt} dt + e^{bt} \sqrt{Y_t} dW_t, \quad t \in \mathbb{R}_+, \end{aligned}$$

and

$$\begin{aligned} d(e^{\theta t} X_t) &= \theta e^{\theta t} X_t dt + e^{\theta t} dX_t = \theta e^{\theta t} X_t dt + e^{\theta t} ((m - \theta X_t) dt + \sqrt{Y_t} dB_t) \\ &= m e^{\theta t} dt + e^{\theta t} \sqrt{Y_t} dB_t, \quad t \in \mathbb{R}_+, \end{aligned}$$

which imply (3.2) and (3.3) in case of $s = 0$. If $0 \leq s \leq t$, then, by

$$e^{bt} Y_t = e^{bs} Y_s + a \int_s^t e^{bu} du + \int_s^t e^{bu} \sqrt{Y_u} dB_u,$$

and

$$e^{\theta t} X_t = e^{\theta s} X_s + m \int_s^t e^{\theta u} du + \int_s^t e^{\theta u} \sqrt{Y_u} dB_u,$$

we have (3.2) and (3.3). Finally, we note that the existence of a pathwise unique strong solution $(Y_t, X_t)_{t \in \mathbb{R}_+}$ of the SDE (3.1) with $\mathbb{P}(Y_t \geq 0 \text{ for all } t \in \mathbb{R}_+) = 1$ follows also by a general result of Dawson and Li [10, Theorem 6.2]. \square

Note that it is the assumption $a \in \mathbb{R}_{++}$ that ensures $\mathbb{P}(Y_t \geq 0, \forall t \in \mathbb{R}_+) = 1$.

Next we present a result about the first moment of $(Y_t, X_t)_{t \in \mathbb{R}_+}$.

Proposition 3.2. *Let $(Y_t, X_t)_{t \in \mathbb{R}_+}$ be a strong solution of the SDE (3.1) satisfying $\mathbb{P}(Y_0 \geq 0) = 1$, $\mathbb{E}(Y_0) < \infty$, and $\mathbb{E}(X_0) < \infty$. Then*

$$\begin{bmatrix} \mathbb{E}(Y_t) \\ \mathbb{E}(X_t) \end{bmatrix} = \begin{bmatrix} e^{-bt} & 0 \\ 0 & e^{-\theta t} \end{bmatrix} \begin{bmatrix} \mathbb{E}(Y_0) \\ \mathbb{E}(X_0) \end{bmatrix} + \begin{bmatrix} \int_0^t e^{-bs} ds & 0 \\ 0 & \int_0^t e^{-\theta s} ds \end{bmatrix} \begin{bmatrix} a \\ m \end{bmatrix}, \quad t \in \mathbb{R}_+,$$

Proof. By Proposition 3.1, we have

$$\begin{aligned} Y_t &= e^{-bt} \left(Y_0 + a \int_0^t e^{bu} du + \int_0^t e^{bu} \sqrt{Y_u} dW_u \right), \quad t \in \mathbb{R}_+, \\ X_t &= e^{-\theta t} \left(X_0 + m \int_0^t e^{\theta u} du + \int_0^t e^{\theta u} \sqrt{Y_u} dB_u \right), \quad t \in \mathbb{R}_+, \end{aligned}$$

and so, taking expectations of both sides,

$$\begin{aligned} \mathbb{E}(Y_t) &= e^{-bt} \mathbb{E}(Y_0) + a e^{-bt} \int_0^t e^{bu} du = e^{-bt} \mathbb{E}(Y_0) + a \int_0^t e^{-bu} du, \quad t \in \mathbb{R}_+, \\ \mathbb{E}(X_t) &= e^{-\theta t} \mathbb{E}(X_0) + m e^{-\theta t} \int_0^t e^{\theta u} du = e^{-\theta t} \mathbb{E}(X_0) + m \int_0^t e^{-\theta u} du, \quad t \in \mathbb{R}_+, \end{aligned}$$

where we used that the processes

$$\left(\int_0^t e^{bu} \sqrt{Y_u} dW_u \right)_{t \in \mathbb{R}_+} \quad \text{and} \quad \left(\int_0^t e^{\theta u} \sqrt{Y_u} dB_u \right)_{t \in \mathbb{R}_+}$$

are martingales which can be checked as follows. First we check that they are local martingales with respect to the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$. Let us define the increasing sequence of stopping times by $\delta_n := \inf\{t \geq 0 : Y_t \geq n\}$, $n \in \mathbb{N}$. Since Y has continuous trajectories almost surely, we have $\mathbb{P}(\lim_{n \rightarrow \infty} \delta_n = \infty) = 1$. Using $(\delta_n)_{n \in \mathbb{N}}$ as a localizing sequence, we have

$$\mathbb{E} \left(\int_0^{t \wedge \delta_n} e^{2bu} Y_u du \right) \leq nt \max(1, e^{2bt}), \quad t \in \mathbb{R}_+, n \in \mathbb{N}.$$

The local martingale property of $(\int_0^t e^{bu} \sqrt{Y_u} dW_u)_{t \in \mathbb{R}_+}$ follows by Ikeda and Watanabe [21, page 57]. Hence, using (3.2) and that $a \in \mathbb{R}_{++}$, we find that

$$\mathbb{E}(e^{b(t \wedge \delta_n)} Y_{t \wedge \delta_n}) = \mathbb{E}(Y_0) + a \mathbb{E} \left(\int_0^{t \wedge \delta_n} e^{bu} du \right) \leq \mathbb{E}(Y_0) + at \max(1, e^{bt})$$

for all $t \in \mathbb{R}_+$ and $n \in \mathbb{N}$, and then, by Fatou's lemma,

$$\mathbb{E}(e^{bt} Y_t) \leq \liminf_{n \rightarrow \infty} \mathbb{E}(e^{b(t \wedge \delta_n)} Y_{t \wedge \delta_n}) \leq \mathbb{E}(Y_0) + at \max(1, e^{bt}), \quad t \in \mathbb{R}_+. \quad (3.4)$$

Next, we can deduce that $(\int_0^t e^{bu} \sqrt{Y_u} dW_u)_{t \in \mathbb{R}_+}$ is indeed a martingale. First, we note that a local martingale M is a square integrable martingale if $E([M, M]_t) < \infty$ for all $t \in \mathbb{R}_+$, where $([M, M]_t)_{t \in \mathbb{R}_+}$ denotes the quadratic variation process of M , see, e.g., Corollary 3 on page 73 in Protter [32]. Here the quadratic variation process of $(\int_0^t e^{bu} \sqrt{Y_u} dW_u)_{t \in \mathbb{R}_+}$ takes the form

$$E \left(\int_0^t e^{2bu} Y_u du \right) < \infty, \quad t \in \mathbb{R}_+,$$

where, for the inequality, we used Fubini's theorem, (3.4) and our assumption $\mathbb{E}(Y_0) < \infty$. Replacing b by θ , we have the desired martingale property of $(\int_0^t e^{\theta u} \sqrt{Y_u} dB_u)_{t \in \mathbb{R}_+}$, too. \square

Next we show that the process $(Y_t, X_t)_{t \in \mathbb{R}_+}$ given by the SDE (3.1) is an affine process.

Proposition 3.3. *Let $(Y_t, X_t)_{t \in \mathbb{R}_+}$ be a strong solution of the SDE (3.1) satisfying $\mathbb{P}(Y_0 \geq 0) = 1$. Then $(Y_t, X_t)_{t \in \mathbb{R}_+}$ is an affine process with infinitesimal generator*

$$\begin{aligned} (\mathcal{A}_{(Y,X)} f)(x) &= (a - bx_1) f'_1(x) + (m - \theta x_2) f'_2(x) \\ &\quad + \frac{1}{2} x_1 (f''_{1,1}(x) + f''_{2,2}(x)) \end{aligned} \quad (3.5)$$

for $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}$ and $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R})$.

Proof. For calculating the infinitesimal generator of $(Y_t, X_t)_{t \in \mathbb{R}_+}$, without loss of generality, we may suppose that $\mathbb{P}((Y_0, X_0) = (y_0, x_0)) = 1$, where $(y_0, x_0) \in \mathbb{R}_+ \times \mathbb{R}$. By Itô's formula, for all real-valued functions $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R})$ we have

$$\begin{aligned} f(Y_t, X_t) &= f(y_0, x_0) + \int_0^t f'_1(Y_s, X_s) \sqrt{Y_s} dW_s + \int_0^t f'_2(Y_s, X_s) \sqrt{Y_s} dB_s \\ &\quad + \int_0^t f'_1(Y_s, X_s) (a - bY_s) ds + \int_0^t f'_2(Y_s, X_s) (m - \theta X_s) ds \\ &\quad + \frac{1}{2} \left(\int_0^t f''_{1,1}(Y_s, X_s) Y_s ds + \int_0^t f''_{2,2}(Y_s, X_s) Y_s ds \right) \\ &= f(y_0, x_0) + \int_0^t (\mathcal{A}_{(Y,X)} f)(Y_s, X_s) ds + M_t(f), \quad t \in \mathbb{R}_+, \end{aligned}$$

where

$$M_t(f) := \int_0^t f'_1(Y_s, X_s)\sqrt{Y_s} dW_s + \int_0^t f'_2(Y_s, X_s)\sqrt{Y_s} dB_s, \quad t \in \mathbb{R}_+,$$

and $\mathcal{A}_{(Y,X)}f$ is given by (3.5). Here $(M_t(f))_{t \in \mathbb{R}_+}$ is a square integrable martingale with respect to the filtration $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$, since

$$\begin{aligned} \int_0^t \mathbb{E}((f'_1(Y_s, X_s))^2 Y_s) ds &\leq C_1 \int_0^t \mathbb{E}(Y_s) ds < \infty, & t \in \mathbb{R}_+, \\ \int_0^t \mathbb{E}((f'_2(Y_s, X_s))^2 Y_s) ds &\leq C_2 \int_0^t \mathbb{E}(Y_s) ds < \infty, & t \in \mathbb{R}_+, \end{aligned}$$

with some constants $C_1 > 0$ and $C_2 > 0$, where the finiteness of the integrals follow by Proposition 3.2. Finally, if $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R})$ is complex valued, then, by decomposing f into real and imaginary parts, one can argue in the same way as above. \square

By Proposition 3.3, the process $(Y_t, X_t)_{t \in \mathbb{R}_+}$ given by (3.1) is a two-dimensional affine process with admissible parameters

$$\left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} a \\ m \end{bmatrix}, \begin{bmatrix} -b & 0 \\ 0 & -\theta \end{bmatrix}, 0, 0 \right).$$

In what follows we define and study criticality of the affine process given by the SDE (3.1).

Definition 3.1. Let $(Y_t, X_t)_{t \in \mathbb{R}_+}$ be an affine diffusion process given by the SDE (3.1) satisfying $\mathbb{P}(Y_0 \geq 0) = 1$. We call $(Y_t, X_t)_{t \in \mathbb{R}_+}$ subcritical, critical or supercritical if the spectral radius of the matrix

$$\begin{pmatrix} e^{-bt} & 0 \\ 0 & e^{-\theta t} \end{pmatrix}$$

is less than 1, equal to 1 or greater than 1, respectively.

Note that, since the spectral radius of the matrix given in Definition 3.1 is $\max(e^{-bt}, e^{-\theta t})$, the affine process given in Definition 3.1 is

$$\begin{aligned} \text{subcritical} & \quad \text{if } b > 0 \text{ and } \theta > 0, \\ \text{critical} & \quad \text{if } b = 0, \theta \geq 0 \text{ or } b \geq 0, \theta = 0, \\ \text{supercritical} & \quad \text{if } b < 0 \text{ or } \theta < 0. \end{aligned}$$

Definition 3.1 of criticality is in accordance with the corresponding definition for one-dimensional continuous state branching processes, see, e.g., Li [27, page 58].

In this section we will always suppose that

$$\begin{aligned} \text{Condition (C):} \quad & (b, \theta) = (0, 0), \mathbb{P}(Y_0 \geq 0) = 1, \\ & \mathbb{E}(Y_0) < \infty, \text{ and } \mathbb{E}(X_0^2) < \infty. \end{aligned}$$

For some explanations why we study only this special case, see Remarks 3.1, 3.2 and 3.3. In the next sections under Condition (C) we will study asymptotic behaviour of least squares estimator of θ and (θ, m) , respectively. Before doing so we recall some critical models both in discrete and continuous time.

In general, parameter estimation for critical models has a long history. A common feature of the estimators for parameters of critical models is that one may prove weak limit theorems for them by using norming factors that are usually different from the norming factors for the subcritical and supercritical models. Further, it may happen that one has to use different norming factors for two different critical cases.

We recall some discrete time critical models. If $(\xi_k)_{k \in \mathbb{Z}_+}$ is an AR(1) process, i.e., $\xi_k = \rho \xi_{k-1} + \zeta_k$, $k \in \mathbb{N}$, with $\xi_0 = 0$ and an i.i.d. sequence $(\zeta_k)_{k \in \mathbb{N}}$ having mean 0 and positive variance, then the (ordinary) least squares estimator of the so-called stability parameter ρ based on the sample ξ_1, \dots, ξ_n takes the form

$$\tilde{\rho}_n = \frac{\sum_{k=1}^n \xi_{k-1} \xi_k}{\sum_{k=1}^n \xi_k^2}, \quad n \in \mathbb{N},$$

see, e.g., Hamilton [16, 17.4.2]. In the critical case, i.e., when $\rho = 1$, by Hamilton [16, 17.4.7],

$$n(\tilde{\rho}_n - 1) \xrightarrow{\mathcal{L}} \frac{\int_0^1 \mathcal{W}_t d\mathcal{W}_t}{\int_0^1 \mathcal{W}_t^2 dt} \quad \text{as } n \rightarrow \infty,$$

where $(\mathcal{W}_t)_{t \in \mathbb{R}_+}$ is a standard Wiener process and $\xrightarrow{\mathcal{L}}$ denotes convergence in distribution. Here $n(\tilde{\rho}_n - 1)$ is known as the Dickey-Fuller statistics. We emphasize that the asymptotic behaviour of $\tilde{\rho}_n$ is completely different in the subcritical ($|\rho| < 1$) and supercritical ($|\rho| > 1$) cases, where it is asymptotically normal and asymptotically Cauchy, respectively, see, e.g., Mann and Wald [29], Anderson [2] and White [36].

For continuous time critical models, we recall that Huang et al. [20, Theorem 2.4] studied asymptotic behaviour of weighted conditional least squares estimator of the drift parameters for discretely observed continuous time critical branching processes with immigration given by

$$\begin{aligned} \tilde{Y}_t &= \tilde{Y}_0 + \int_0^t (a + b\tilde{Y}_s) ds + \sigma \int_0^t \sqrt{\tilde{Y}_s} d\mathcal{W}_s + \int_0^t \int_{[0, \infty)} \xi \mathcal{N}_0(ds, d\xi) \\ &\quad + \int_0^t \int_0^{\tilde{Y}_{s-}} \int_{[0, \infty)} \xi (\mathcal{N}_1(ds, du, d\xi) - ds du p(d\xi)), \quad t \in \mathbb{R}_+, \end{aligned}$$

where $\tilde{Y}_0 \geq 0$, $a \geq 0$, $b \in \mathbb{R}$, $\sigma \geq 0$, \mathcal{W} is a standard Wiener process, $\mathcal{N}_0(ds, d\xi)$ is a Poisson random measure on $(0, \infty) \times [0, \infty)$ with intensity $ds n(d\xi)$, $\mathcal{N}_1(ds, du, d\xi)$ is a Poisson random measure on $(0, \infty) \times (0, \infty) \times [0, \infty)$ with intensity $ds du p(d\xi)$ such that the σ -finite measures n and p are supported by $(0, \infty)$ and

$$\int_0^\infty \xi n(d\xi) + \int_0^\infty \xi \wedge \xi^2 p(d\xi) < \infty.$$

Our technique differs from that of Huang et al. [20] and for completeness we note that the limit distribution and some parts of the proof of their Theorem 2.4 suffer from some misprints. Furthermore, Hu and Long [19] studied the problem of parameter estimation for critical mean-reverting α -stable motions

$$d\tilde{X}_t = (m - \theta\tilde{X}_t) dt + dZ_t, \quad t \in \mathbb{R}_+,$$

where Z is an α -stable Lévy motion with $\alpha \in (0, 2)$ observed at discrete instants. A least squares estimator is obtained and its asymptotics is discussed in the singular case $(m, \theta) = (0, 0)$. We note that the forms of the limit distributions of least squares estimators for critical two-dimensional affine diffusion processes in our Theorems 3.1 and 3.2 are the same as that of the limit distributions in Theorems 3.2 and 4.1 in Hu and Long [19], respectively. We also recall that Hu and Long [18] considered the problem of parameter estimation not only for critical mean-reverting α -stable motions, but also for some subcritical ones ($m = 0$ and $\theta > 0$) by proving limit theorems for the least squares estimators that are completely different from the ones in the critical case. Huang et al. [20] investigated the asymptotic behaviour of weighted conditional least squares estimator of the drift parameters not only for critical continuous time branching processes with immigration, but also for subcritical and supercritical ones.

Using our scaling Theorem 2.2 we can only handle a special critical affine diffusion model given by (1.1) (for a more detailed discussion, see Remark 3.2). The other critical and non-critical cases are under investigation but different techniques are needed.

3.2. Least squares estimator of θ when m is known

The least squares estimator (LSE) of θ based on the observations $X_i, i = 0, 1, \dots, n$, can be obtained by solving the extremum problem

$$\tilde{\theta}_n^{\text{LSE}} := \arg \min_{\theta \in \mathbb{R}} \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1}))^2.$$

This definition of LSE of θ can be considered as the counterpart of the one given in Hu and Long [18, formula (1.2)] for generalized Ornstein-Uhlenbeck processes driven by α -stable motions, see also Hu and Long [19, formulas (3.1) and (4.1)]. For a mathematical motivation of the definition of the LSE of θ , see later on Remark 3.4. With the notation $f(\theta) := \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1}))^2$, $\theta \in \mathbb{R}$, the equation $f'(\theta) = 0$ takes the form:

$$2 \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1})) X_{i-1} = 0.$$

Hence

$$\left(\sum_{i=1}^n X_{i-1}^2 \right) \theta = - \sum_{i=1}^n (X_i - X_{i-1} - m) X_{i-1},$$

i.e.,

$$\begin{aligned}\tilde{\theta}_n^{\text{LSE}} &= -\frac{\sum_{i=1}^n (X_i - X_{i-1} - m)X_{i-1}}{\sum_{i=1}^n X_{i-1}^2} \\ &= -\frac{\sum_{i=1}^n (X_i - X_{i-1})X_{i-1} - (\sum_{i=1}^n X_{i-1})m}{\sum_{i=1}^n X_{i-1}^2}\end{aligned}\quad (3.6)$$

provided that $\sum_{i=1}^n X_{i-1}^2 > 0$. Since $f''(\theta) = 2\sum_{i=1}^n X_{i-1}^2$, $\theta \in \mathbb{R}$, we have $\tilde{\theta}_n^{\text{LSE}}$ is indeed the solution of the extremum problem provided that $\sum_{i=1}^n X_{i-1}^2 > 0$.

Theorem 3.1. *Let us assume that Condition (C) holds. Then $\mathbb{P}(\sum_{i=1}^n X_{i-1}^2 > 0) = 1$ for all $n \geq 2$, and there exists a unique LSE $\tilde{\theta}_n^{\text{LSE}}$ which has the form given in (3.6). Further,*

$$n\tilde{\theta}_n^{\text{LSE}} \xrightarrow{\mathcal{L}} -\frac{\int_0^1 \mathcal{X}_t d\mathcal{X}_t - m \int_0^1 \mathcal{X}_t dt}{\int_0^1 \mathcal{X}_t^2 dt} \quad \text{as } n \rightarrow \infty, \quad (3.7)$$

where $(\mathcal{X}_t)_{t \in \mathbb{R}_+}$ is the second coordinate of a two-dimensional affine process $(\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+}$ given by the unique strong solution of the SDE

$$\begin{cases} d\mathcal{Y}_t = a dt + \sqrt{\mathcal{Y}_t} d\mathcal{W}_t, \\ d\mathcal{X}_t = m dt + \sqrt{\mathcal{Y}_t} d\mathcal{B}_t, \end{cases} \quad t \in \mathbb{R}_+, \quad (3.8)$$

with initial value $(\mathcal{Y}_0, \mathcal{X}_0) = (0, 0)$, where $(\mathcal{W}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{B}_t)_{t \in \mathbb{R}_+}$ are independent standard Wiener processes.

Remark 3.1. (i) The limit distributions in Theorem 3.1 have the same forms as those of the limit distributions in Theorem 3.2 in Hu and Long [19].

(ii) The limit distribution of $n\tilde{\theta}_n^{\text{LSE}}$ as $n \rightarrow \infty$ in Theorem 3.1 can be written also in the form

$$-\frac{\int_0^1 \mathcal{X}_t d(\mathcal{X}_t - mt)}{\int_0^1 \mathcal{X}_t^2 dt} = -\frac{\int_0^1 \mathcal{X}_t \sqrt{\mathcal{Y}_t} d\mathcal{B}_t}{\int_0^1 \mathcal{X}_t^2 dt}.$$

(iii) By Proposition 3.3, the affine process $(\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+}$ given in Theorem 3.1 has infinitesimal generator

$$(\mathcal{A}_{(\mathcal{Y}, \mathcal{X})} f)(x) = \frac{1}{2}x_1 f''_{1,1}(x) + \frac{1}{2}x_1 f''_{2,2}(x) + a f'_1(x) + m f'_2(x)$$

where $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}$ and $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R})$.

(iv) Under the Condition (C), by Theorem 3.1 and Slutsky's lemma, we get $\tilde{\theta}_n^{\text{LSE}}$ converges stochastically to the parameter $\theta = 0$ as $n \rightarrow \infty$. \square

Proof of Theorem 3.1. By (3.3), we have

$$X_t = X_0 + mt + \int_0^t \sqrt{Y_s} dB_s, \quad t \in \mathbb{R}_+.$$

Hence for all $t \in \mathbb{R}_{++}$, the conditional distribution of X_t given X_0 and $(Y_s)_{s \in [0,t]}$ is a normal distribution with mean $X_0 + mt$ and with variance $\int_0^t Y_s ds$. Here the variance $\int_0^t Y_s ds$ is positive almost surely for all $t \in \mathbb{R}_{++}$. Indeed, let $A_t := \{\omega \in \Omega : s \mapsto Y_s(\omega) \text{ is continuous on } [0, t]\}$. Then $\mathbb{P}(A_t) = 1$, and, since $\mathbb{P}(Y_0 \geq 0) = 1$, for all $\omega \in A_t$, $\int_0^t Y_s(\omega) ds = 0$ if and only if $Y_s(\omega) = 0$ for all $s \in [0, t]$. By (3.1), we have

$$Y_s = Y_0 + as + \int_0^s \sqrt{Y_u} dW_u, \quad s \in \mathbb{R}_+.$$

The stochastic integral on the right hand side can be approximated as

$$\sup_{s \in [0,t]} \left| \sum_{i=1}^{\lfloor ns \rfloor} \sqrt{Y_{(i-1)/n}} (W_{i/n} - W_{(i-1)/n}) - \int_0^s Y_u dW_u \right| \xrightarrow{\mathbb{P}} 0 \quad \text{as } n \rightarrow \infty$$

for all $t \in \mathbb{R}_+$, by Jacod and Shiryaev [22, Theorem I.4.44]. Hence there exists a sequence $(n_k)_{k \in \mathbb{N}}$ of positive integers such that

$$\sup_{s \in [0,t]} \left| \sum_{i=1}^{\lfloor n_k t \rfloor} \sqrt{Y_{(i-1)/n_k}} (W_{i/n_k} - W_{(i-1)/n_k}) - \int_0^s Y_u dW_u \right| \xrightarrow{\text{a.s.}} 0 \quad \text{as } k \rightarrow \infty$$

for all $t \in \mathbb{R}_+$. Consequently, with the notation

$$\tilde{A}_t := \left\{ \omega \in \Omega : \int_0^t Y_s(\omega) ds = 0 \right\},$$

we have

$$\begin{aligned} \tilde{A}_t \cap A_t &\subset \left\{ \tilde{A}_t \cap \left\{ \int_0^s Y_u dW_u = 0 \text{ for all } s \in [0, t] \right\} \right\} \\ &\subset \left\{ \tilde{A}_t \cap \{Y_s = Y_0 + as \text{ for all } s \in [0, t]\} \right\} \\ &= \left\{ \tilde{A}_t \cap \{Y_0 s + as^2/2 = 0 \text{ for all } s \in [0, t]\} \right\} \\ &= \left\{ \tilde{A}_t \cap \{Y_0 = -as/2 \text{ for all } s \in [0, t]\} \right\} = \emptyset, \end{aligned}$$

implying $\mathbb{P}(\int_0^t Y_s ds = 0) = 0$, and hence $\mathbb{P}(\int_0^t Y_s ds > 0) = 1$. It yields that

$$\mathbb{P}(X_t = 0) = \mathbb{E} \left(\mathbb{P}(X_t = 0 \mid X_0, (Y_s)_{s \in [0,t]}) \right) = 0, \quad t \in \mathbb{R}_{++}, \quad (3.9)$$

and hence $\mathbb{P}(\sum_{i=1}^n X_{i-1}^2 > 0) = 1$ for all $n \geq 2$.

Now we turn to prove (3.7). By Itô's formula, we have $d(\mathcal{X}_t^2) = 2\mathcal{X}_t d\mathcal{X}_t + \mathcal{Y}_t dt$, $t \in \mathbb{R}_+$, and hence, using also $\mathcal{X}_0 = 0$, we have

$$\int_0^1 \mathcal{X}_s d\mathcal{X}_s = \frac{1}{2} \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_s ds \right). \tag{3.10}$$

For the process $(X_t)_{t \in \mathbb{R}_+}$, a discrete version

$$\sum_{i=1}^n (X_i - X_{i-1})X_{i-1} = \frac{1}{2} \left(X_n^2 - X_0^2 - \sum_{i=1}^n (X_i - X_{i-1})^2 \right) \tag{3.11}$$

of the identity (3.10) can be easily checked. The aim of the following discussion is to prove

$$\begin{aligned} & \left(\frac{1}{n^2} \sum_{i=1}^n X_{i-1}, \frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2, \frac{1}{n} X_n, \frac{1}{n} X_0, \frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})^2 \right) \\ & \xrightarrow{\mathcal{L}} \left(\int_0^1 \mathcal{X}_t dt, \int_0^1 \mathcal{X}_t^2 dt, \mathcal{X}_1, 0, \int_0^1 \mathcal{Y}_t dt \right) \quad \text{as } n \rightarrow \infty. \end{aligned} \tag{3.12}$$

Let us consider the unique strong solution of the SDE

$$\begin{cases} d\tilde{Y}_t = a dt + \sqrt{\tilde{Y}_t} dW_t, \\ d\tilde{X}_t = m dt + \sqrt{\tilde{Y}_t} dB_t, \end{cases} \quad t \in \mathbb{R}_+, \tag{3.13}$$

with initial value $(\tilde{Y}_0, \tilde{X}_0) = (0, 0)$, where $(W_t)_{t \in \mathbb{R}_+}$ and $(B_t)_{t \in \mathbb{R}_+}$ are the independent standard $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -Wiener processes appearing in the SDE (3.1). First note that $(\tilde{Y}_t, \tilde{X}_t)_{t \in \mathbb{R}_+} \stackrel{\mathcal{L}}{=} (\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+}$, and, by Proposition 3.3, it is an affine process having admissible parameters

$$\left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} a \\ m \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, 0, 0 \right),$$

and condition (2.3) is trivially fulfilled. Hence, by part (ii) of Remark 2.7, we have

$$\left(n^{-1}\tilde{Y}_{nt}, n^{-1}\tilde{X}_{nt} \right)_{t \in \mathbb{R}_+} \stackrel{\mathcal{L}}{=} (\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+} \quad \text{for all } n \in \mathbb{N}. \tag{3.14}$$

Consequently, for all $n \in \mathbb{N}$, we have

$$\begin{aligned} & \left(\frac{1}{n^2} \sum_{i=1}^n \tilde{X}_{i-1}, \frac{1}{n^3} \sum_{i=1}^n \tilde{X}_{i-1}^2, \frac{1}{n} \tilde{X}_n, \frac{1}{n} \tilde{X}_0, \frac{1}{n^2} \sum_{i=1}^n (\tilde{X}_i - \tilde{X}_{i-1})^2 \right) \\ & \stackrel{\mathcal{L}}{=} (A_n, B_n, C_n, D_n, E_n), \end{aligned}$$

where

$$\begin{aligned}
 A_n &:= \frac{1}{n} \sum_{i=1}^n \mathcal{X}_{(i-1)/n} \xrightarrow{\text{a.s.}} \int_0^1 \mathcal{X}_t dt, & \text{as } n \rightarrow \infty, \\
 B_n &:= \frac{1}{n} \sum_{i=1}^n \mathcal{X}_{(i-1)/n}^2 \xrightarrow{\text{a.s.}} \int_0^1 \mathcal{X}_t^2 dt, & \text{as } n \rightarrow \infty, \\
 C_n &:= \mathcal{X}_1, \\
 D_n &:= \mathcal{X}_0, \\
 E_n &:= \sum_{i=1}^n (\mathcal{X}_{i/n} - \mathcal{X}_{(i-1)/n})^2 \xrightarrow{\mathbb{P}} \int_0^1 \mathcal{Y}_t dt, & \text{as } n \rightarrow \infty. \tag{3.15}
 \end{aligned}$$

Here the first two convergences are consequences of the definition of the Riemann integral using also that $(\mathcal{X}_t)_{t \in \mathbb{R}_+}$ has continuous sample paths almost surely. The convergence (3.15) can be checked as follows. With the notations of Jacod and Shiryaev [22], $(\tau_n := (\frac{i}{n} \wedge 1)_{i \in \mathbb{N}})_{n \in \mathbb{N}}$ is a Riemann sequence of (adapted) subdivisions and hence, by Jacod and Shiryaev [22, Theorem I.4.47], the sequence of processes

$$\left(\sum_{i=1}^n \left(\mathcal{X} \left(\frac{i}{n} \wedge 1 \wedge t \right) - \mathcal{X} \left(\frac{i-1}{n} \wedge 1 \wedge t \right) \right)^2 \right)_{t \in \mathbb{R}_+}, \quad n \in \mathbb{N},$$

converges to the quadratic variation process of \mathcal{X} in probability, uniformly on every compact interval. Especially, with $t = 1$, using also the SDE (3.8), we have (3.15). Hence, in order to prove (3.12), it suffices to show convergences

$$\frac{1}{n^2} \sum_{i=0}^{n-1} X_i - \frac{1}{n^2} \sum_{i=0}^{n-1} \tilde{X}_i \xrightarrow{\mathbb{P}} 0, \tag{3.16}$$

$$\frac{1}{n^3} \sum_{i=0}^{n-1} X_i^2 - \frac{1}{n^3} \sum_{i=0}^{n-1} \tilde{X}_i^2 \xrightarrow{\mathbb{P}} 0, \tag{3.17}$$

$$\frac{1}{n} X_n - \frac{1}{n} \tilde{X}_n \xrightarrow{\mathbb{P}} 0, \tag{3.18}$$

$$\frac{1}{n} X_0 - \frac{1}{n} \tilde{X}_0 \xrightarrow{\mathbb{P}} 0, \tag{3.19}$$

$$\frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})^2 - \frac{1}{n^2} \sum_{i=1}^n (\tilde{X}_i - \tilde{X}_{i-1})^2 \xrightarrow{\mathbb{P}} 0, \tag{3.20}$$

as $n \rightarrow \infty$. Indeed, one can refer to Slutsky's lemma using also that for any random vectors $U_n, V_n, n \in \mathbb{N}, U, V$ such that $U_n \xrightarrow{\mathbb{P}} U$ and $V_n \xrightarrow{\mathbb{P}} V$ as $n \rightarrow \infty$, we have $(U_n, V_n) \xrightarrow{\mathbb{P}} (U, V)$ as $n \rightarrow \infty$, see, e.g., van der Vaart [33, Theorem 2.7, part (vi)].

The convergence (3.19) is trivial. Next we show

$$\mathbb{E}(|Y_t - \tilde{Y}_t|) \leq \mathbb{E}(Y_0), \quad t \in \mathbb{R}_+. \quad (3.21)$$

By (3.1) and (3.13), we have

$$Y_t - \tilde{Y}_t = Y_0 + \int_0^t (\sqrt{Y_s} - \sqrt{\tilde{Y}_s}) dW_s, \quad t \in \mathbb{R}_+.$$

For each $n \in \mathbb{N}$, there exists an even and twice continuously differentiable function $\psi_n : \mathbb{R} \rightarrow \mathbb{R}_+$ with $|\psi_n(x)| \leq |x|$, $|\psi'_n(x)| \leq 1$, $\psi_n(x) \uparrow |x|$ as $n \rightarrow \infty$ for all $x \in \mathbb{R}$, and

$$\psi''_n(x-y)(\sqrt{x} - \sqrt{y})^2 \leq \frac{2(\sqrt{x} - \sqrt{y})^2}{n|x-y|} \leq \frac{2}{n}$$

for all $n \in \mathbb{N}$ and $x, y \in \mathbb{R}_+$, see, e.g., in Karatzas and Shreve [24, Proof of Proposition 5.2.13]. By Itô's formula,

$$\begin{aligned} \psi_n(Y_t - \tilde{Y}_t) &= \psi_n(Y_0) + \frac{1}{2} \int_0^t \psi''_n(Y_s - \tilde{Y}_s) \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right)^2 ds \\ &\quad + \int_0^t \psi'_n(Y_s - \tilde{Y}_s) \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right) dW_s \end{aligned} \quad (3.22)$$

for all $t \in \mathbb{R}_+$ and $n \in \mathbb{N}$. The last term is an $(\mathcal{F}_t)_{t \in \mathbb{R}_+}$ -martingale, since

$$\begin{aligned} \mathbb{E} \left(\int_0^t |\psi'_n(Y_s - \tilde{Y}_s)| \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right)^2 ds \right) &\leq \mathbb{E} \left(\int_0^t |Y_s - \tilde{Y}_s| ds \right) \\ &\leq \int_0^t (\mathbb{E}(Y_s) + \mathbb{E}(\tilde{Y}_s)) ds < \infty, \end{aligned}$$

where the last inequality follows by Lemma 3.2. Thus the expectation of the last term on the right hand side of (3.22) is zero, whereas the expectation of the second integral is bounded by $2t/n$. We conclude

$$\mathbb{E}(\psi_n(Y_t - \tilde{Y}_t)) \leq \mathbb{E}(\psi_n(Y_0)) + \frac{t}{n}, \quad t \in \mathbb{R}_+, \quad n \in \mathbb{N}.$$

By monotone convergence theorem, we have

$$\begin{aligned} \mathbb{E}(|Y_t - \tilde{Y}_t|) &= \mathbb{E} \left(\lim_{n \rightarrow \infty} \psi_n(Y_t - \tilde{Y}_t) \right) = \lim_{n \rightarrow \infty} \mathbb{E}(\psi_n(Y_t - \tilde{Y}_t)) \\ &\leq \liminf_{n \rightarrow \infty} \left(\mathbb{E}(\psi_n(Y_0)) + \frac{t}{n} \right) = \lim_{n \rightarrow \infty} \mathbb{E}(\psi_n(Y_0)) = \mathbb{E} \left(\lim_{n \rightarrow \infty} \psi_n(Y_0) \right) = \mathbb{E}(|Y_0|), \end{aligned}$$

which yields (3.21).

Next, we derive

$$\mathbb{E}(|X_t - \tilde{X}_t|) \leq \mathbb{E}(|X_0|) + \sqrt{t \mathbb{E}(Y_0)}, \quad t \in \mathbb{R}_+. \quad (3.23)$$

Again by (3.1) and (3.13), we have

$$X_t - \tilde{X}_t = X_0 + \int_0^t \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right) dB_s, \quad t \in \mathbb{R}_+, \quad (3.24)$$

hence

$$\begin{aligned} \mathbb{E}(|X_t - \tilde{X}_t|) &\leq \mathbb{E}(|X_0|) + \sqrt{\mathbb{E} \left(\left(\int_0^t (\sqrt{Y_s} - \sqrt{\tilde{Y}_s}) dB_s \right)^2 \right)} \\ &= \mathbb{E}(|X_0|) + \sqrt{\mathbb{E} \left(\int_0^t (\sqrt{Y_s} - \sqrt{\tilde{Y}_s})^2 ds \right)} \\ &\leq \mathbb{E}(|X_0|) + \sqrt{\mathbb{E} \left(\int_0^t |Y_s - \tilde{Y}_s| ds \right)} \\ &= \mathbb{E}(|X_0|) + \sqrt{\int_0^t \mathbb{E}(|Y_s - \tilde{Y}_s|) ds}, \end{aligned}$$

which yields (3.23) by (3.21).

By (3.23), we have

$$\mathbb{E} \left(\left| \frac{1}{n} X_n - \frac{1}{n} \tilde{X}_n \right| \right) \leq \frac{1}{n} \left(\mathbb{E}(|X_0|) + \sqrt{n \mathbb{E}(Y_0)} \right) \rightarrow 0 \quad \text{as } n \rightarrow \infty,$$

hence we obtain (3.18). In a similar way,

$$\mathbb{E} \left(\left| \frac{1}{n^2} \sum_{i=0}^{n-1} X_i - \frac{1}{n^2} \sum_{i=0}^{n-1} \tilde{X}_i \right| \right) \leq \frac{1}{n^2} \sum_{i=0}^{n-1} \left(\mathbb{E}(|X_0|) + \sqrt{i \mathbb{E}(Y_0)} \right) \rightarrow 0$$

as $n \rightarrow \infty$, hence we obtain (3.16).

We also have

$$\mathbb{E} \left((X_t - \tilde{X}_t)^2 \right) \leq 2 \mathbb{E}(X_0^2) + 2t \mathbb{E}(Y_0), \quad t \in \mathbb{R}_+. \quad (3.25)$$

Indeed, by (3.24), using Minkowski inequality, we have

$$\begin{aligned} \sqrt{\mathbb{E} \left((X_t - \tilde{X}_t)^2 \right)} &\leq \sqrt{\mathbb{E}(X_0^2)} + \sqrt{\mathbb{E} \left(\left(\int_0^t (\sqrt{Y_s} - \sqrt{\tilde{Y}_s}) dB_s \right)^2 \right)} \\ &= \sqrt{\mathbb{E}(X_0^2)} + \sqrt{\mathbb{E} \left(\int_0^t (\sqrt{Y_s} - \sqrt{\tilde{Y}_s})^2 ds \right)} \\ &\leq \sqrt{\mathbb{E}(X_0^2)} + \sqrt{\mathbb{E} \left(\int_0^t |Y_s - \tilde{Y}_s| ds \right)} \\ &= \sqrt{\mathbb{E}(X_0^2)} + \sqrt{\int_0^t \mathbb{E}(|Y_s - \tilde{Y}_s|) ds} \leq \sqrt{\mathbb{E}(X_0^2)} + \sqrt{t \mathbb{E}(Y_0)} \end{aligned}$$

by (3.21), which yields (3.25). In a similar way,

$$\mathbb{E}(X_t^2) \leq 3 \mathbb{E}(X_0^2) + 3m^2t^2 + 3t \mathbb{E}(Y_0) + 3at^2/2, \quad t \in \mathbb{R}_+. \tag{3.26}$$

since, by (3.1) and (3.2),

$$\begin{aligned} \sqrt{\mathbb{E}(X_t^2)} &\leq \sqrt{\mathbb{E}(X_0^2)} + |m|t + \sqrt{\mathbb{E}\left(\left(\int_0^t \sqrt{Y_s} dB_s\right)^2\right)} \\ &= \sqrt{\mathbb{E}(X_0^2)} + |m|t + \sqrt{\mathbb{E}\left(\int_0^t Y_s ds\right)} \\ &= \sqrt{\mathbb{E}(X_0^2)} + |m|t + \sqrt{\int_0^t (\mathbb{E}(Y_0) + as) ds} \\ &= \sqrt{\mathbb{E}(X_0^2)} + |m|t + \sqrt{t \mathbb{E}(Y_0) + at^2/2} \end{aligned}$$

for $t \in \mathbb{R}_+$, which yields (3.26). Clearly, (3.26) implies also $\mathbb{E}(\tilde{X}_t^2) \leq 3m^2t^2 + 3at^2/2$ for all $t \in \mathbb{R}_+$, and hence, together with (3.25) and (3.26), we conclude

$$\begin{aligned} \mathbb{E}\left(\left|\frac{1}{n^3} \sum_{i=0}^{n-1} X_i^2 - \frac{1}{n^3} \sum_{i=0}^{n-1} \tilde{X}_i^2\right|\right) &\leq \frac{1}{n^3} \sum_{i=0}^{n-1} \mathbb{E}\left(|(X_i - \tilde{X}_i)(X_i + \tilde{X}_i)|\right) \\ &\leq \frac{1}{n^3} \sum_{i=0}^{n-1} \sqrt{\mathbb{E}((X_i - \tilde{X}_i)^2) \mathbb{E}((X_i + \tilde{X}_i)^2)} \\ &\leq \frac{1}{n^3} \sum_{i=0}^{n-1} \sqrt{2 \mathbb{E}((X_i - \tilde{X}_i)^2) (\mathbb{E}(X_i^2) + \mathbb{E}(\tilde{X}_i^2))} \\ &\leq \frac{1}{n^3} \sum_{i=0}^{n-1} \sqrt{12(\mathbb{E}(X_0^2) + i \mathbb{E}(Y_0)) (\mathbb{E}(X_0^2) + (\mathbb{E}(Y_0)i + (2m^2 + a)i^2))} \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$, hence we obtain (3.17).

Next, we show (3.20). Again by (3.1) and (3.13), we have

$$X_i - X_{i-1} = m + \int_{i-1}^i \sqrt{Y_s} dB_s, \quad \tilde{X}_i - \tilde{X}_{i-1} = m + \int_{i-1}^i \sqrt{\tilde{Y}_s} dB_s, \quad i \in \mathbb{N},$$

and hence

$$\begin{aligned} &\frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})^2 - \frac{1}{n^2} \sum_{i=1}^n (\tilde{X}_i - \tilde{X}_{i-1})^2 \\ &= \frac{2m}{n^2} \int_0^n \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s}\right) dB_s \\ &\quad + \frac{1}{n^2} \sum_{i=1}^n \left[\left(\int_{i-1}^i \sqrt{Y_s} dB_s\right)^2 - \left(\int_{i-1}^i \sqrt{\tilde{Y}_s} dB_s\right)^2\right] \\ &=: 2mR_n + S_n. \end{aligned}$$

Here, by (3.21),

$$\begin{aligned} \mathbb{E}(R_n^2) &= \frac{1}{n^4} \mathbb{E} \left(\left(\int_0^n \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right) dB_s \right)^2 \right) \\ &= \frac{1}{n^4} \mathbb{E} \left(\int_0^n \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right)^2 ds \right) \leq \frac{1}{n^4} \mathbb{E} \left(\int_0^n |Y_s - \tilde{Y}_s| ds \right) \\ &= \frac{1}{n^4} \int_0^n \mathbb{E}(|Y_s - \tilde{Y}_s|) ds \leq \frac{\mathbb{E}(Y_0)}{n^3} \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned}$$

hence $R_n \xrightarrow{\mathbb{P}} 0$ as $n \rightarrow \infty$. Further, by (3.21),

$$\begin{aligned} \mathbb{E}(|S_n|) &= \mathbb{E} \left(\left| \frac{1}{n^2} \sum_{i=1}^n \int_{i-1}^i \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right) dB_s \int_{i-1}^i \left(\sqrt{Y_s} + \sqrt{\tilde{Y}_s} \right) dB_s \right| \right) \\ &\leq \frac{1}{n^2} \sum_{i=1}^n \mathbb{E} \left(\left| \int_{i-1}^i \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right) dB_s \int_{i-1}^i \left(\sqrt{Y_s} + \sqrt{\tilde{Y}_s} \right) dB_s \right| \right) \\ &\leq \frac{1}{n^2} \sum_{i=1}^n \sqrt{\mathbb{E} \left(\left(\int_{i-1}^i \left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right) dB_s \right)^2 \right) \mathbb{E} \left(\left(\int_{i-1}^i \left(\sqrt{Y_s} + \sqrt{\tilde{Y}_s} \right) dB_s \right)^2 \right)} \\ &= \frac{1}{n^2} \sum_{i=1}^n \sqrt{\int_{i-1}^i \mathbb{E} \left(\left(\sqrt{Y_s} - \sqrt{\tilde{Y}_s} \right)^2 \right) ds \int_{i-1}^i \mathbb{E} \left(\left(\sqrt{Y_s} + \sqrt{\tilde{Y}_s} \right)^2 \right) ds} \\ &\leq \frac{1}{n^2} \sum_{i=1}^n \sqrt{\int_{i-1}^i \mathbb{E}(|Y_s - \tilde{Y}_s|) ds \int_{i-1}^i 2(\mathbb{E}(Y_s) + \mathbb{E}(\tilde{Y}_s)) ds} \\ &\leq \frac{1}{n^2} \sum_{i=1}^n \sqrt{\mathbb{E}(Y_0) \int_{i-1}^i 2(\mathbb{E}(Y_0) + 2as) ds} \\ &= \frac{1}{n^2} \sum_{i=1}^n \sqrt{2\mathbb{E}(Y_0)(\mathbb{E}(Y_0) + (2i-1)a)} \rightarrow 0 \quad \text{as } n \rightarrow \infty, \end{aligned}$$

thus $S_n \xrightarrow{\mathbb{P}} 0$ as $n \rightarrow \infty$, and we obtain (3.20), and hence (3.12).

Finally, by (3.12) and the continuous mapping theorem, and using that

$$n\tilde{\theta}_n^{\text{LSE}} = \frac{\frac{m}{n^2} \sum_{i=1}^n X_{i-1} - \frac{1}{2n^2} X_n^2 + \frac{1}{2n^2} X_0^2 + \frac{1}{2n^2} \sum_{i=1}^n (X_i - X_{i-1})^2}{\frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2},$$

we have the assertion. Indeed, the function $g : \mathbb{R}^5 \rightarrow \mathbb{R}$, defined by

$$g(x, y, z, u, v) := \begin{cases} \frac{mx - (z^2 - u^2 - v)/2}{y} & \text{if } y \neq 0, \\ 0 & \text{if } y = 0, \end{cases}$$

is continuous on the set $\{(x, y, z, u, v) \in \mathbb{R}^5 : y \neq 0\}$, and the limit distribution in (3.12) is concentrated on this set since $\mathbb{P}(\int_0^1 \mathcal{X}_t^2 dt > 0) = 1$. Indeed, if

$\mathbb{P}(\int_0^1 \mathcal{X}_t^2 dt = 0) > 0$ held, then, by the almost sure continuity of the sample paths of \mathcal{X} , we would have $\mathbb{P}(\mathcal{X}_t = 0, \forall t \in [0, 1]) > 0$. Hence on the event $\{\omega \in \Omega : \mathcal{X}_t(\omega) = 0, \forall t \in [0, 1]\}$, the quadratic variation of \mathcal{X} would be identically zero. Since $d\mathcal{X}_t = m dt + \sqrt{\mathcal{Y}_t} d\mathcal{B}_t$, $t \in \mathbb{R}_+$, the quadratic variation of \mathcal{X} is the process $(\int_0^t \mathcal{Y}_u du)_{t \in \mathbb{R}_+}$, and then we would have

$$\int_0^t \mathcal{Y}_u du = 0 \quad \text{for all } t \in [0, 1]$$

on the event $\{\omega \in \Omega : \mathcal{X}_t(\omega) = 0, \forall t \in [0, 1]\}$. This yields us to a contradiction similarly as at the beginning of the proof due to that $a \in \mathbb{R}_{++}$ and $d\mathcal{Y}_t = a dt + \sqrt{\mathcal{W}_t} d\mathcal{W}_t$, $t \in \mathbb{R}_+$. Hence the continuous mapping theorem (see, e.g., Theorem 2.3 in van der Vaart [33]) yields

$$g\left(\frac{1}{n^2} \sum_{i=1}^n X_{i-1}, \frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2, \frac{1}{n} X_n, \frac{1}{n} X_0, \frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})^2\right) \\ \xrightarrow{\mathcal{L}} g\left(\int_0^1 \mathcal{X}_t dt, \int_0^1 \mathcal{X}_t^2 dt, \mathcal{X}_1, 0, \int_0^1 \mathcal{Y}_t dt\right) \quad \text{as } n \rightarrow \infty.$$

Since

$$\mathbb{P}\left(n\tilde{\theta}_n^{\text{LSE}} = g\left(\frac{1}{n^2} \sum_{i=1}^n X_{i-1}, \frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2, \frac{1}{n} X_n, \frac{1}{n} X_0, \frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})^2\right)\right) \\ \geq \mathbb{P}\left(\sum_{i=1}^n X_{i-1}^2 > 0\right) = 1$$

for all $n \geq 2$, the assertion follows using (3.10) and that if $\xi_n, \eta_n, n \in \mathbb{N}$, and ξ are random variables such that $\xi_n \xrightarrow{\mathcal{L}} \xi$ as $n \rightarrow \infty$ and $\lim_{n \rightarrow \infty} \mathbb{P}(\xi_n = \eta_n) = 1$, then $\eta_n \xrightarrow{\mathcal{L}} \xi$ as $n \rightarrow \infty$, see, e.g., Barczy et al. [3, Lemma 3.1]. \square

Remark 3.2. If the affine diffusion process given by the SDE (3.1) is critical but $(b, \theta) \neq (0, 0)$ (i.e., $b = 0, \theta > 0$ or $b > 0, \theta = 0$), then the asymptotic behaviour of the LSE $\tilde{\theta}_n^{\text{LSE}}$ cannot be studied using Theorem 2.2 since its condition $\lim_{\theta \rightarrow \infty} \theta \beta^{(\theta)} = \beta$ is not satisfied. \square

3.3. Least squares estimator of (θ, m)

The LSE of (θ, m) based on the observations $X_i, i = 0, 1, \dots, n$, can be obtained by solving the extremum problem

$$(\hat{\theta}_n^{\text{LSE}}, \hat{m}_n^{\text{LSE}}) := \arg \min_{(\theta, m) \in \mathbb{R}^2} \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1}))^2.$$

We need to solve the following system of equations with respect to (θ, m) :

$$\begin{aligned} 2 \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1})) X_{i-1} &= 0, \\ 2 \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1})) &= 0, \end{aligned}$$

which can be written also in the form

$$\begin{bmatrix} \sum_{i=1}^n X_{i-1}^2 & -\sum_{i=1}^n X_{i-1} \\ -\sum_{i=1}^n X_{i-1} & n \end{bmatrix} \begin{bmatrix} \theta \\ m \end{bmatrix} = \begin{bmatrix} -\sum_{i=1}^n (X_i - X_{i-1}) X_{i-1} \\ \sum_{i=1}^n (X_i - X_{i-1}) \end{bmatrix}.$$

Then one can check that

$$\hat{\theta}_n^{\text{LSE}} = -\frac{n \sum_{i=1}^n (X_i - X_{i-1}) X_{i-1} - \sum_{i=1}^n X_{i-1} \sum_{i=1}^n (X_i - X_{i-1})}{n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2}, \quad (3.27)$$

and

$$\hat{m}_n^{\text{LSE}} = \frac{\sum_{i=1}^n X_{i-1}^2 \sum_{i=1}^n (X_i - X_{i-1}) - \sum_{i=1}^n X_{i-1} \sum_{i=1}^n (X_i - X_{i-1}) X_{i-1}}{n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2} \quad (3.28)$$

provided that $n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2 > 0$. Since the matrix

$$\begin{bmatrix} 2 \sum_{i=1}^n X_{i-1}^2 & -2 \sum_{i=1}^n X_{i-1} \\ -2 \sum_{i=1}^n X_{i-1} & 2n \end{bmatrix}$$

which consists of the second order partial derivatives of the function $\mathbb{R}^2 \ni (\theta, m) \mapsto \sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1}))^2$ is positive definite provided that $n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2 > 0$, we have $(\hat{\theta}_n^{\text{LSE}}, \hat{m}_n^{\text{LSE}})$ is indeed the solution of the extremum problem provided that $n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2 > 0$.

Theorem 3.2. *Let us assume that Condition (C) holds. Then*

$$\mathbb{P} \left(n \sum_{i=1}^n X_{i-1}^2 - \left(\sum_{i=1}^n X_{i-1} \right)^2 > 0 \right) = 1 \quad \text{for all } n \geq 2, \quad (3.29)$$

and there exists a unique LSE $(\hat{\theta}_n^{\text{LSE}}, \hat{m}_n^{\text{LSE}})$ which has the form given in (3.27) and (3.28). Further,

$$n \hat{\theta}_n^{\text{LSE}} \xrightarrow{\mathcal{L}} -\frac{\int_0^1 \mathcal{X}_t d\mathcal{X}_t - \mathcal{X}_1 \int_0^1 \mathcal{X}_t dt}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt \right)^2} \quad \text{as } n \rightarrow \infty,$$

and

$$\widehat{m}_n^{\text{LSE}} \xrightarrow{\mathcal{L}} \frac{\mathcal{X}_1 \int_0^1 \mathcal{X}_t^2 dt - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d\mathcal{X}_t}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt \right)^2} \quad \text{as } n \rightarrow \infty,$$

where $(\mathcal{X}_t)_{t \in \mathbb{R}_+}$ is the second coordinate of a two-dimensional affine process $(\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+}$ given by the unique strong solution of the SDE

$$\begin{cases} d\mathcal{Y}_t = a dt + \sqrt{\mathcal{Y}_t} d\mathcal{W}_t, \\ d\mathcal{X}_t = m dt + \sqrt{\mathcal{Y}_t} d\mathcal{B}_t, \end{cases} \quad t \in \mathbb{R}_+,$$

with initial value $(\mathcal{Y}_0, \mathcal{X}_0) = (0, 0)$, where $(\mathcal{W}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{B}_t)_{t \in \mathbb{R}_+}$ are independent standard Wiener processes.

Remark 3.3. (i) The limit distributions in Theorem 3.2 have the same forms as those of the limit distributions in Theorem 4.1 in Hu and Long [19].

(ii) By Proposition 3.3, the affine process $(\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+}$ given in Theorem 3.2 has infinitesimal generator

$$(\mathcal{A}_{(\mathcal{Y}, \mathcal{X})} f)(x) = \frac{1}{2} x_1 f''_{1,1}(x) + \frac{1}{2} x_1 f''_{2,2}(x) + a f'_1(x) + m f'_2(x),$$

where $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}$ and $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R})$.

(iii) Under the Condition (C), by Theorem 3.2 and Slutsky's lemma, we get $\widehat{\theta}_n^{\text{LSE}}$ converges stochastically to the parameter $\theta = 0$ as $n \rightarrow \infty$, and one can show that $\widehat{m}_n^{\text{LSE}}$ does not converge stochastically to the parameter m as $n \rightarrow \infty$, see Appendix B. \square

Proof of Theorem 3.2. By an easy calculation,

$$n \sum_{i=1}^n X_{i-1}^2 - \left(\sum_{i=1}^n X_{i-1} \right)^2 = n \sum_{i=1}^n \left(X_{i-1} - \frac{1}{n} \sum_{j=1}^n X_{j-1} \right)^2 \geq 0,$$

and equality holds if and only if

$$X_{i-1} = \frac{1}{n} \sum_{j=1}^n X_{j-1}, \quad i = 1, \dots, n \quad \iff \quad X_0 = X_1 = \dots = X_{n-1}.$$

By (3.3), for all $n \geq 2$,

$$\begin{aligned} \mathbb{P}(X_0 = X_1 = \dots = X_{n-1}) &\leq \mathbb{P}(X_0 = X_1) = \mathbb{P}\left(\int_0^1 \sqrt{Y_s} dB_s = m\right) \\ &= \mathbb{E}\left(\mathbb{P}\left(\int_0^1 \sqrt{Y_s} dB_s = m \mid (Y_s)_{s \in [0,1]}\right)\right) = 0, \end{aligned}$$

where we used that the conditional distribution of $\int_0^1 \sqrt{Y_s} dB_s$ given $(Y_s)_{s \in [0,1]}$ is a normal distribution with mean 0 and with variance $\int_0^1 Y_s ds$. Here the variance $\int_0^1 Y_s ds$ is positive almost surely, see the proof of Theorem 3.1. This yields (3.29).

By (3.27) and (3.28), we have

$$\begin{aligned} n\widehat{\theta}_n^{\text{LSE}} &= -\frac{\frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})X_{i-1} - \frac{1}{n^2} \sum_{i=1}^n X_{i-1} \frac{1}{n} (X_n - X_0)}{\frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2 - \left(\frac{1}{n^2} \sum_{i=1}^n X_{i-1}\right)^2}, \\ \widehat{m}_n^{\text{LSE}} &= \frac{\frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2 \frac{1}{n} (X_n - X_0) - \frac{1}{n^2} \sum_{i=1}^n X_{i-1} \frac{1}{n^2} \sum_{i=1}^n (X_i - X_{i-1})}{\frac{1}{n^3} \sum_{i=1}^n X_{i-1}^2 - \left(\frac{1}{n^2} \sum_{i=1}^n X_{i-1}\right)^2}, \end{aligned}$$

and using (3.11) and (3.12), as in the proof of Theorem 3.1, the continuous mapping theorem yields the assertion. We only remark that

$$\mathbb{P}\left(\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2 > 0\right) = 1. \tag{3.30}$$

Indeed,

$$\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2 = \int_0^1 \left(\mathcal{X}_t - \int_0^1 \mathcal{X}_s ds\right)^2 \geq 0,$$

and equality holds if and only if

$$\mathcal{X}_t = \int_0^1 \mathcal{X}_s ds \quad \text{a.e. } t \in [0, 1].$$

Since \mathcal{X} has continuous sample paths almost surely,

$$\mathbb{P}\left(\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2 = 0\right) > 0 \tag{3.31}$$

holds if and only if

$$\mathbb{P}\left(\mathcal{X}_t = \int_0^1 \mathcal{X}_s ds, \forall t \in [0, 1]\right) > 0.$$

Hence, since $\mathcal{X}_0 = 0$, we have (3.31) holds if and only if $\mathbb{P}(\mathcal{X}_t = 0, \forall t \in [0, 1]) > 0$, which is a contradiction due to our assumption $a \in \mathbb{R}_{++}$ (for more details, see the proof of Theorem 3.1). \square

3.4. Conditional least squares estimator of (θ, m)

For all $t \in \mathbb{R}_+$, let $\mathcal{F}_t^{(Y, X)}$ be the σ -algebra generated by $(Y_s, X_s)_{s \in [0, t]}$. The conditional least squares estimator (CLSE) of (θ, m) based on the observations $X_i, i = 0, 1, \dots, n$, can be obtained by solving the extremum problem

$$(\widehat{\theta}_n^{\text{CLSE}}, \widehat{m}_n^{\text{CLSE}}) := \arg \min_{(\theta, m) \in \mathbb{R}^2} \sum_{i=1}^n \left(X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1}^{(Y, X)})\right)^2.$$

By (3.3), for all $(y_0, x_0) \in \mathbb{R}_+ \times \mathbb{R}$, we have

$$\mathbb{E}(X_t | (Y_0, X_0) = (y_0, x_0)) = e^{-\theta t} x_0 + m \int_0^t e^{-\theta(t-u)} du, \quad t \geq 0,$$

where we used that $(\int_0^t e^{\theta u} \sqrt{Y_u} dB_u)_{t \in \mathbb{R}_+}$ is a martingale (which follows by the proof of Proposition 3.2). Using that $(Y_t, X_t)_{t \in \mathbb{R}_+}$ is a time-homogeneous Markov process, we have

$$\mathbb{E}(X_t | \mathcal{F}_s^{(Y, X)}) = \mathbb{E}(X_t | (Y_s, X_s)) = e^{-\theta(t-s)} X_s + m \int_s^t e^{-\theta(t-u)} du$$

for $0 \leq s \leq t$. Then

$$\begin{aligned} X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1}^{(Y, X)}) &= X_i - e^{-\theta} X_{i-1} - m \int_{i-1}^i e^{-\theta(i-u)} du \\ &= X_i - e^{-\theta} X_{i-1} - m \int_0^1 e^{-\theta v} dv \\ &= X_i - \gamma X_{i-1} - \delta, \quad i = 1, \dots, n, \end{aligned}$$

where

$$\gamma := e^{-\theta} \quad \text{and} \quad \delta := m \int_0^1 e^{-\theta v} dv = \begin{cases} m \frac{1-e^{-\theta}}{\theta} & \text{if } \theta \neq 0, \\ m & \text{if } \theta = 0. \end{cases}$$

Hence for all $n \in \mathbb{N}$,

$$\begin{aligned} \widehat{\gamma}_n^{\text{CLSE}} &= e^{-\widehat{\theta}_n^{\text{CLSE}}}, \\ \widehat{\delta}_n^{\text{CLSE}} &= \widehat{m}_n^{\text{CLSE}} \int_0^1 e^{-\widehat{\theta}_n^{\text{CLSE}} v} dv, \end{aligned} \tag{3.32}$$

where $(\widehat{\gamma}_n^{\text{CLSE}}, \widehat{\delta}_n^{\text{CLSE}})$ is a CLSE of (γ, δ) based on the observations X_i , $i = 0, 1, \dots, n$, which can be obtained by solving the extremum problem

$$(\widehat{\gamma}_n^{\text{CLSE}}, \widehat{\delta}_n^{\text{CLSE}}) := \arg \min_{(\gamma, \delta) \in \mathbb{R}^2} \sum_{i=1}^n (X_i - \gamma X_{i-1} - \delta)^2. \tag{3.33}$$

Indeed, the function $A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$,

$$\mathbb{R}^2 \ni (\theta', m') \mapsto A(\theta', m') := \begin{bmatrix} \gamma' \\ \delta' \end{bmatrix} =: \begin{bmatrix} e^{-\theta'} \\ m' \int_0^1 e^{-\theta' v} dv \end{bmatrix} \in \mathbb{R}_+ \times \mathbb{R}$$

is bijective and measurable, and then there is a bijection between the set of CLSEs of the parameters (θ, m) and the set of CLSEs of the parameters

$A(\theta, m)$. This follows easily, since for all $n \in \mathbb{N}$, $(x_0, x_1, \dots, x_n) \in \mathbb{R}^{n+1}$ and $(\gamma', \delta') \in \mathbb{R}_+ \times \mathbb{R}$,

$$\begin{aligned} \sum_{i=1}^n (x_i - \gamma' x_{i-1} - \delta')^2 &= \sum_{i=1}^n \left(x_i - \begin{bmatrix} \gamma' \\ \delta' \end{bmatrix}^\top \begin{bmatrix} x_{i-1} \\ 1 \end{bmatrix} \right)^2 \\ &= \sum_{i=1}^n \left(x_i - (A(\theta', m'))^\top \begin{bmatrix} x_{i-1} \\ 1 \end{bmatrix} \right)^2, \end{aligned}$$

hence $(\widehat{\theta}_n^{\text{CLSE}}, \widehat{m}_n^{\text{CLSE}})$ is a CLSE of (θ, m) if and only if $A(\widehat{\theta}_n^{\text{CLSE}}, \widehat{m}_n^{\text{CLSE}})$ is a CLSE of $A(\theta, m)$.

For the extremum problem (3.33), we need to solve the following system of equations with respect to (γ, δ) :

$$\begin{aligned} 2 \sum_{i=1}^n (X_i - \gamma X_{i-1} - \delta) X_{i-1} &= 0, \\ 2 \sum_{i=1}^n (X_i - \gamma X_{i-1} - \delta) &= 0, \end{aligned}$$

which can be written also in the form

$$\begin{bmatrix} \sum_{i=1}^n X_{i-1}^2 & \sum_{i=1}^n X_{i-1} \\ \sum_{i=1}^n X_{i-1} & n \end{bmatrix} \begin{bmatrix} \gamma \\ \delta \end{bmatrix} = \begin{bmatrix} \sum_{i=1}^n X_{i-1} X_i \\ \sum_{i=1}^n X_i \end{bmatrix}.$$

Then

$$\widehat{\gamma}_n^{\text{CLSE}} = \frac{n \sum_{i=1}^n X_{i-1} X_i - \sum_{i=1}^n X_{i-1} \sum_{i=1}^n X_i}{n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2}, \tag{3.34}$$

and

$$\widehat{\delta}_n^{\text{CLSE}} = \frac{\sum_{i=1}^n X_{i-1}^2 \sum_{i=1}^n X_i - \sum_{i=1}^n X_{i-1} \sum_{i=1}^n X_{i-1} X_i}{n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2}, \tag{3.35}$$

provided that $n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2 \neq 0$. Since the matrix

$$\begin{bmatrix} 2 \sum_{i=1}^n X_{i-1}^2 & 2 \sum_{i=1}^n X_{i-1} \\ 2 \sum_{i=1}^n X_{i-1} & 2n \end{bmatrix}$$

consisting of the second order partial derivatives of the function $\mathbb{R}^2 \ni (\gamma, \delta) \mapsto \sum_{i=1}^n (X_i - \gamma X_{i-1} - \delta)^2$ is positive definite provided that $n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2 > 0$, we have $(\widehat{\gamma}_n^{\text{CLSE}}, \widehat{\delta}_n^{\text{CLSE}})$ is indeed the solution of the extremum problem provided that $n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2 > 0$.

Remark 3.4. Using the definition of CLSE of (θ, m) we give a mathematical motivation of the definition of the LSE $\widetilde{\theta}_n$ of θ introduced in Section 3.2. Note that if $\theta = 0$, then

$$X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1}^{(Y, X)}) = X_i - X_{i-1} - m, \quad i = 1, \dots, n.$$

If $\theta \neq 0$, then, by Taylor's theorem, $1 - e^{-\theta} = e^{-\tau\theta}$ with some $\tau = \tau(\theta) \in [0, 1]$, and hence

$$\begin{aligned} X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1}^{(Y,X)}) &= X_i - e^{-\theta} X_{i-1} - m \int_0^1 e^{-\theta v} dv \\ &= X_i - X_{i-1} + e^{-\tau\theta} \theta X_{i-1} - m e^{-\tau\theta} \end{aligned}$$

for $i = 1, \dots, n-1$. Hence for small values of θ one can approximate $X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1}^{(Y,X)})$ by $X_i - X_{i-1} + \theta X_{i-1} - m = X_i - X_{i-1} - (m - \theta X_{i-1})$, $i = 1, \dots, n$. Based on this, for small values of θ , in the definition of the LSE of θ , the sum $\sum_{i=1}^n (X_i - X_{i-1} - (m - \theta X_{i-1}))^2$ can be considered as an approximation of the sum $\sum_{i=1}^n (X_i - \mathbb{E}(X_i | \mathcal{F}_{i-1}^{(Y,X)}))^2$ in the definition of the CLSE of (θ, m) . \square

Theorem 3.3. *Let us assume that Condition (C) holds. Then*

$$\mathbb{P} \left(n \sum_{i=1}^n X_{i-1}^2 - \left(\sum_{i=1}^n X_{i-1} \right)^2 > 0 \right) = 1 \quad \text{for all } n \geq 2, \quad (3.36)$$

and there exists a unique CLSE $(\hat{\theta}_n^{\text{CLSE}}, \hat{m}_n^{\text{CLSE}})$ which has the form given in (3.32). Further,

$$n \hat{\theta}_n^{\text{CLSE}} \xrightarrow{\mathcal{L}} - \frac{\int_0^1 \mathcal{X}_t d\mathcal{X}_t - \mathcal{X}_1 \int_0^1 \mathcal{X}_t dt}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt \right)^2} \quad \text{as } n \rightarrow \infty, \quad (3.37)$$

and

$$\hat{m}_n^{\text{CLSE}} \xrightarrow{\mathcal{L}} \frac{\mathcal{X}_1 \int_0^1 \mathcal{X}_t^2 dt - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d\mathcal{X}_t}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt \right)^2} \quad \text{as } n \rightarrow \infty, \quad (3.38)$$

where $(\mathcal{X}_t)_{t \in \mathbb{R}_+}$ is the second coordinate of a two-dimensional affine process $(\mathcal{Y}_t, \mathcal{X}_t)_{t \in \mathbb{R}_+}$ given by the unique strong solution of the SDE

$$\begin{cases} d\mathcal{Y}_t = a dt + \sqrt{\mathcal{Y}_t} d\mathcal{W}_t, \\ d\mathcal{X}_t = m dt + \sqrt{\mathcal{Y}_t} d\mathcal{B}_t, \end{cases} \quad t \in \mathbb{R}_+,$$

with initial value $(\mathcal{Y}_0, \mathcal{X}_0) = (0, 0)$, where $(\mathcal{W}_t)_{t \in \mathbb{R}_+}$ and $(\mathcal{B}_t)_{t \in \mathbb{R}_+}$ are independent standard Wiener processes.

Proof. By the proof of Theorem 3.2, we have (3.36). By (3.34) and (3.35), for all $n \geq 2$ we have

$$\hat{\gamma}_n^{\text{CLSE}} - 1 = \frac{n \sum_{i=1}^n (X_i - X_{i-1}) X_{i-1} - X_n \sum_{i=1}^n X_{i-1}}{n \sum_{i=1}^n X_{i-1}^2 - \left(\sum_{i=1}^n X_{i-1} \right)^2},$$

and

$$\widehat{\delta}_n^{\text{CLSE}} = \frac{X_n \sum_{i=1}^n X_{i-1}^2 - \sum_{i=1}^n X_{i-1} \sum_{i=1}^n (X_i - X_{i-1}) X_{i-1}}{n \sum_{i=1}^n X_{i-1}^2 - (\sum_{i=1}^n X_{i-1})^2}.$$

Using (3.11) and (3.12), the continuous mapping theorem, by the same technique as in the proof of Theorem 3.2, we get

$$n(\widehat{\gamma}_n^{\text{CLSE}} - 1) \xrightarrow{\mathcal{L}} \frac{\int_0^1 \mathcal{X}_t d\mathcal{X}_t - \mathcal{X}_1 \int_0^1 \mathcal{X}_t dt}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2} \quad \text{as } n \rightarrow \infty, \quad (3.39)$$

and

$$\widehat{\delta}_n^{\text{CLSE}} \xrightarrow{\mathcal{L}} \frac{\mathcal{X}_1 \int_0^1 \mathcal{X}_t^2 dt - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d\mathcal{X}_t}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2} \quad \text{as } n \rightarrow \infty. \quad (3.40)$$

By Slutsky's lemma, we also have $\widehat{\gamma}_n^{\text{CLSE}} \xrightarrow{\mathbb{P}} 1$ as $n \rightarrow \infty$. Hence, by Taylor's theorem using also that $\widehat{\gamma}_n^{\text{CLSE}} > 0$, $n \in \mathbb{N}$ (due to its definition given in (3.32)), we have

$$\widehat{\theta}_n^{\text{CLSE}} = -\log(\widehat{\gamma}_n^{\text{CLSE}}) = -\log(\widehat{\gamma}_n^{\text{CLSE}}) - \log(1) = -\frac{1}{\xi_n}(\widehat{\gamma}_n^{\text{CLSE}} - 1), \quad (3.41)$$

where ξ_n is in the interval with endpoints 1 and $\widehat{\gamma}_n^{\text{CLSE}}$. Since $\widehat{\gamma}_n^{\text{CLSE}} \xrightarrow{\mathbb{P}} 1$ as $n \rightarrow \infty$, we have $\xi_n \xrightarrow{\mathbb{P}} 1$ as $n \rightarrow \infty$, and hence using the decomposition

$$n\widehat{\theta}_n^{\text{CLSE}} = -\frac{1}{\xi_n}n(\widehat{\gamma}_n^{\text{CLSE}} - 1), \quad n \in \mathbb{N},$$

Slutsky's lemma and (3.39), we get (3.37).

Next we turn to prove (3.38). For this, by (3.32), (3.40) and by Slutsky's lemma, it is enough to check that

$$\int_0^1 e^{-\widehat{\theta}_n^{\text{CLSE}} v} dv \xrightarrow{\mathbb{P}} 1 \quad \text{as } n \rightarrow \infty.$$

Since

$$\int_0^1 e^{-\widehat{\theta}_n^{\text{CLSE}} v} dv = \begin{cases} \frac{1 - e^{-\widehat{\theta}_n^{\text{CLSE}}}}{\widehat{\theta}_n^{\text{CLSE}}} & \text{if } \widehat{\theta}_n^{\text{CLSE}} \neq 0, \\ 1 & \text{if } \widehat{\theta}_n^{\text{CLSE}} = 0, \end{cases}$$

by (3.41), for all $\varepsilon > 0$ we have

$$\begin{aligned} & \mathbb{P} \left(\left| \int_0^1 e^{-\widehat{\theta}_n^{\text{CLSE}} v} dv - 1 \right| \geq \varepsilon \right) \\ &= \mathbb{P} \left(\left| \frac{1 - \widehat{\gamma}_n^{\text{CLSE}}}{\widehat{\theta}_n^{\text{CLSE}}} - 1 \right| \geq \varepsilon \mid \widehat{\theta}_n^{\text{CLSE}} \neq 0 \right) \mathbb{P}(\widehat{\theta}_n^{\text{CLSE}} \neq 0) \\ & \quad + \mathbb{P} \left(|1 - 1| \geq \varepsilon \mid \widehat{\theta}_n^{\text{CLSE}} = 0 \right) \mathbb{P}(\widehat{\theta}_n^{\text{CLSE}} = 0) \\ &= \mathbb{P} \left(|\xi_n - 1| \geq \varepsilon \mid \widehat{\theta}_n^{\text{CLSE}} \neq 0 \right) \mathbb{P}(\widehat{\theta}_n^{\text{CLSE}} \neq 0) \xrightarrow{\mathbb{P}} 0, \end{aligned}$$

since $\xi_n \xrightarrow{\mathbb{P}} 0$ as $n \rightarrow \infty$. \square

Remark 3.5. (i) We do not consider the CLSE of θ supposing that m is known since the corresponding extremum problem is rather complicated, and from statistical point of view it has less importance.

(ii) Under the Condition (C), by Theorem 3.3 and Slutsky's lemma, we get $\widehat{\theta}_n^{\text{CLSE}}$ converges stochastically to the parameter $\theta = 0$ as $n \rightarrow \infty$, and one can show that $\widehat{m}_n^{\text{CLSE}}$ does not converge stochastically to the parameter m as $n \rightarrow \infty$, see Appendix B. \square

Appendix

Appendix A: The integrals in (2.2)

We check that the integrals in (2.2) are well-defined, i.e., elements of \mathbb{C} , under the conditions (v) and (vi) of Definition 2.2. For this, by decomposing a complex-valued function to real and imaginary parts, it is enough to verify that

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'(x), \xi \rangle) x_1 \mu(d\xi) < \infty, \quad (\text{A.1})$$

$$\int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'_{(2)}(x), \xi_2 \rangle) m(d\xi) < \infty \quad (\text{A.2})$$

for all $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and real-valued $f \in \mathcal{C}_c^2(\mathbb{R}_+ \times \mathbb{R}^d)$.

First we check (A.1). If $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in \mathbb{R}_+ \times \mathbb{R}^d$ with $\|\xi\| \leq 1$, then, by (2.11) with $\theta = 1$,

$$|f(x + \xi) - f(x) - \langle f'(x), \xi \rangle| = \frac{1}{2} |\langle f''(x + \tau\xi), \xi \rangle| \leq \frac{1}{2} \|f''\|_\infty \|\xi\|^2,$$

where $\tau = \tau(x, \xi) \in [0, 1]$. If $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in \mathbb{R}_+ \times \mathbb{R}^d$ with $\|\xi\| > 1$, then, by (2.9) with $\theta = 1$,

$$\begin{aligned} |f(x + \xi) - f(x) - \langle f'(x), \xi \rangle| &\leq |f(x + \xi) - f(x)| + |\langle f'(x), \xi \rangle| \\ &= |\langle f'(x + \tau\xi), \xi \rangle| + |\langle f'(x), \xi \rangle| \\ &\leq 2\|f'\|_\infty \|\xi\|, \end{aligned}$$

where $\tau = \tau(x, \xi) \in [0, 1]$. Hence

$$\begin{aligned} & \left| \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'(x), \xi \rangle) x_1 \mu(d\xi) \right| \\ & \leq \int_{\{\xi \in \mathbb{R}_+ \times \mathbb{R}^d : \|\xi\| \leq 1\}} |(f(x + \xi) - f(x) - \langle f'(x), \xi \rangle) x_1| \mu(d\xi) \\ & \quad + \int_{\{\xi \in \mathbb{R}_+ \times \mathbb{R}^d : \|\xi\| > 1\}} |(f(x + \xi) - f(x) - \langle f'(x), \xi \rangle) x_1| \mu(d\xi) \\ & \leq \frac{1}{2} \|f''\|_\infty x_1 \int_{\{\xi \in \mathbb{R}_+ \times \mathbb{R}^d : \|\xi\| \leq 1\}} \|\xi\|^2 \mu(d\xi) \\ & \quad + 2 \|f'\|_\infty x_1 \int_{\{\xi \in \mathbb{R}_+ \times \mathbb{R}^d : \|\xi\| > 1\}} \|\xi\| \mu(d\xi) < \infty, \end{aligned}$$

where the last inequality follows by assumption (vi) of Definition 2.2.

Next we check (A.2). If $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in \mathbb{R}_+ \times \mathbb{R}^d$ with $\|\xi_2\| \leq 1$, then, by (2.11) with $\theta = 1$,

$$\begin{aligned} |f(x + \xi) - f(x) - \langle f'_{(2)}(x), \xi_2 \rangle| &= |f(x + \xi) - f(x) - \langle f'(x), \xi \rangle + \langle f'_1(x), \xi_1 \rangle| \\ &\leq \frac{1}{2} |\langle f''(x + \tau\xi), \xi \rangle| + |\langle f'_1(x), \xi_1 \rangle| \\ &\leq \frac{1}{2} \|f''\|_\infty \|\xi\|^2 + \|f'_1\|_\infty \xi_1, \end{aligned}$$

where $\tau = \tau(x, \xi) \in [0, 1]$. If $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\xi \in \mathbb{R}_+ \times \mathbb{R}^d$ with $\|\xi_2\| > 1$, then, by (2.9) with $\theta = 1$,

$$\begin{aligned} |f(x + \xi) - f(x) - \langle f'_{(2)}(x), \xi_2 \rangle| &= |\langle f'(x + \tau\xi), \xi \rangle - \langle f'_{(2)}(x), \xi_2 \rangle| \\ &= |\langle f'_1(x + \tau\xi), \xi_1 \rangle + \langle f'_{(2)}(x + \tau\xi), \xi_2 \rangle - \langle f'_{(2)}(x), \xi_2 \rangle| \\ &\leq \|f'_1\|_\infty \xi_1 + |\langle f'_{(2)}(x + \tau\xi) - f'_{(2)}(x), \xi_2 \rangle| \\ &\leq \|f'_1\|_\infty \xi_1 + 2 \|f'_{(2)}\| \|\xi_2\|, \end{aligned}$$

where $\tau = \tau(x, \xi) \in [0, 1]$. Using assumption (v) of Definition 2.2, the finiteness of the integral in (A.2) follows as for the integral in (A.1).

Having proved that the integrals in (2.2) are well-defined, we check that under the conditions (v) and (vi) of Definition 2.2, one can rewrite (2.12) in Duffie et al. [11] into the form (2.2), by changing the 2-nd, ..., (1 + d)-th coordinates of $b \in \mathbb{R}_+ \times \mathbb{R}^d$ and the first column of $\beta \in \mathbb{R}^{(1+d) \times (1+d)}$, respectively. More precisely, with the notations

$$\chi(\xi) := (\chi_1(\xi), \dots, \chi_{1+d}(\xi)) \quad \text{and} \quad \chi_{(2)}(\xi) := (\chi_2(\xi), \dots, \chi_{1+d}(\xi))$$

for $\xi \in \mathbb{R}^{1+d}$, where

$$\chi_k(\xi) := \begin{cases} (1 \wedge |\xi_k|) \frac{\xi_k}{|\xi_k|} & \text{if } \xi_k \neq 0, \\ 0 & \text{if } \xi_k = 0, \end{cases} \quad k = 1, \dots, 1 + d,$$

for all $x = (x_1, x_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $f \in C_c^2(\mathbb{R}_+ \times \mathbb{R}^d)$, we have

$$\begin{aligned} (\mathcal{A}f)(x) &= \sum_{i,j=1}^{1+d} (a_{i,j} + \alpha_{i,j}x_1) f''_{i,j}(x) + \langle f'(x), b + \beta x \rangle \\ &\quad + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'_{(2)}(x), \xi_2 \rangle) m(d\xi) \\ &\quad + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'(x), \xi \rangle) x_1 \mu(d\xi) \\ &= \sum_{i,j=1}^{1+d} (a_{i,j} + \alpha_{i,j}x_1) f''_{i,j}(x) + \langle f'(x), \tilde{b} + \tilde{\beta} x \rangle \\ &\quad + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'_{(2)}(x), \chi_{(2)}(\xi) \rangle) m(d\xi) \\ &\quad + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (f(x + \xi) - f(x) - \langle f'(x), \chi(\xi) \rangle) x_1 \mu(d\xi), \end{aligned}$$

where $\tilde{b} = (\tilde{b}_1, \tilde{b}_2) \in \mathbb{R}_+ \times \mathbb{R}^d$ and $\tilde{\beta} = (\tilde{\beta}_{i,j})_{i,j=1}^{1+d} \in \mathbb{R}^{(1+d) \times (1+d)}$ with

$$\begin{aligned} \tilde{b}_1 &:= b_1, \\ \tilde{b}_2 &:= b_2 + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (\chi_{(2)}(\xi) - \xi_2) m(d\xi), \\ (\tilde{\beta}_{i,1})_{i=1}^{1+d} &:= (\beta_{i,1})_{i=1}^{1+d} + \int_{\mathbb{R}_+ \times \mathbb{R}^d} (\chi(\xi) - \xi) \mu(d\xi), \\ (\tilde{\beta}_{i,j})_{i=1, \dots, 1+d}^{j=2, \dots, 1+d} &:= (\beta_{i,1})_{i=1, \dots, 1+d}^{j=2, \dots, 1+d}. \end{aligned}$$

Note also that there is another way for checking that the integrals in (2.2) are well-defined under the conditions (v) and (vi) of Definition 2.2. Namely, using that the integrals in (2.12) in Duffie et al. [11] are well-defined, the assertion follows since

$$\begin{aligned} \left\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} (\chi_{(2)}(\xi) - \xi_2) m(d\xi) \right\| &= \left(\sum_{i=2}^{1+d} \left(\int_{\mathbb{R}_+ \times \mathbb{R}^d} (\chi_i(\xi) - \xi_i) m(d\xi) \right)^2 \right)^{1/2} \\ &= \left(\sum_{i=2}^{1+d} \left(\int_{\mathbb{R}_+ \times \mathbb{R}^d} |\xi_i| \mathbf{1}_{\{|\xi_i| > 1\}} m(d\xi) \right)^2 \right)^{1/2} \\ &\leq \sqrt{d} \int_{\mathbb{R}_+ \times \mathbb{R}^d} \|\xi\| \mathbf{1}_{\{\|\xi\| > 1\}} m(d\xi) < \infty, \end{aligned}$$

and similarly

$$\left\| \int_{\mathbb{R}_+ \times \mathbb{R}^d} (\chi(\xi) - \xi) \mu(d\xi) \right\| < \infty.$$

Appendix B: On consistency properties of the LSE and CLSE of (θ, m)

Let us suppose that Condition (C) holds. Using Slutsky’s lemma and Theorems 3.2 and 3.3 we get $\widehat{\theta}_n^{\text{LSE}}$ and $\widehat{\theta}_n^{\text{CLSE}}$ converge stochastically to the parameter $\theta = 0$ as $n \rightarrow \infty$, respectively, and in what follows we show that $\widehat{m}_n^{\text{LSE}}$ and $\widehat{m}_n^{\text{CLSE}}$ do not converge stochastically to the parameter m as $n \rightarrow \infty$, respectively. For this it is enough to check that the weak limits of $\widehat{m}_n^{\text{LSE}}$ and $\widehat{m}_n^{\text{CLSE}}$ given in Theorems 3.2 and 3.3 do not equal to m almost surely, respectively. Since the weak limits in question are the same, we can give a common proof. First note that

$$\begin{aligned} & \frac{\mathcal{X}_1 \int_0^1 \mathcal{X}_t^2 dt - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d\mathcal{X}_t}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2} - m \\ &= \frac{(\mathcal{X}_1 - m) \int_0^1 \mathcal{X}_t^2 dt - \int_0^1 \mathcal{X}_t dt \left(\int_0^1 \mathcal{X}_t d\mathcal{X}_t - m \int_0^1 \mathcal{X}_t dt\right)}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2} \\ &= \frac{\int_0^1 \mathcal{X}_t^2 dt \int_0^1 d[\mathcal{X}_t - mt] - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d[\mathcal{X}_t - mt]}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2}, \end{aligned}$$

and hence

$$\frac{\mathcal{X}_1 \int_0^1 \mathcal{X}_t^2 dt - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d\mathcal{X}_t}{\int_0^1 \mathcal{X}_t^2 dt - \left(\int_0^1 \mathcal{X}_t dt\right)^2} \stackrel{\text{a.s.}}{=} m$$

if and only if

$$J := \int_0^1 \mathcal{X}_t^2 dt \int_0^1 d[\mathcal{X}_t - mt] - \int_0^1 \mathcal{X}_t dt \int_0^1 \mathcal{X}_t d[\mathcal{X}_t - mt] \stackrel{\text{a.s.}}{=} 0,$$

where $\stackrel{\text{a.s.}}{=}$ denotes equality almost surely. Here J can be written in the form

$$J = \int_0^1 \mathcal{X}_s^2 \left(\int_0^1 d[\mathcal{X}_t - mt]\right) ds - \int_0^1 \mathcal{X}_s \left(\int_0^1 \mathcal{X}_t d[\mathcal{X}_t - mt]\right) ds,$$

and hence $\mathbb{E}(J)$ takes the following form

$$\int_0^1 \mathbb{E} \left(\mathcal{X}_s^2 \int_0^1 d[\mathcal{X}_t - mt] \right) ds - \int_0^1 \mathbb{E} \left(\mathcal{X}_s \int_0^1 \mathcal{X}_t d[\mathcal{X}_t - mt] \right) ds.$$

Here $(\mathcal{X}_t - mt)_{t \in \mathbb{R}_+} = \left(\int_0^t \sqrt{\mathcal{Y}_u} d\mathcal{B}_u\right)_{t \in \mathbb{R}_+}$ is a square integrable martingale (see the proof of Proposition 3.2) and $\mathcal{X}_s = ms + \int_0^s \sqrt{\mathcal{Y}_u} d\mathcal{B}_u$ for $s \in \mathbb{R}_+$, thus,

for all $s \in [0, 1]$, we have

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_s^2 \int_0^1 d[\mathcal{X}_t - mt] \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= m^2 s^2 \mathbb{E} \left(\int_0^1 \sqrt{\mathcal{Y}_t} d\mathcal{B}_t \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 2ms \mathbb{E} \left(\int_0^s \sqrt{\mathcal{Y}_u} d\mathcal{B}_u \int_0^1 \sqrt{\mathcal{Y}_t} d\mathcal{B}_t \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ & \quad + \mathbb{E} \left(\left(\int_0^s \sqrt{\mathcal{Y}_u} d\mathcal{B}_u \right)^2 \int_0^1 \sqrt{\mathcal{Y}_t} d\mathcal{B}_t \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= 2ms \int_0^s \mathcal{Y}_u du, \end{aligned}$$

where $\mathcal{F}_1^{\mathcal{Y}}$ denotes the σ -algebra generated by $(\mathcal{Y}_u)_{u \in [0, 1]}$. For the last equality above, we used that conditionally on the σ -algebra $\mathcal{F}_1^{\mathcal{Y}}$, the stochastic process $(\int_0^t \sqrt{\mathcal{Y}_u} d\mathcal{B}_u)_{t \in [0, 1]}$ is a Gauss process with mean function identically 0 and with covariance function

$$\mathbb{E} \left(\int_0^s \sqrt{\mathcal{Y}_u} d\mathcal{B}_u \int_0^t \sqrt{\mathcal{Y}_u} d\mathcal{B}_u \middle| \mathcal{F}_1^{\mathcal{Y}} \right) = \int_0^{s \wedge t} \mathcal{Y}_u du, \quad s, t \in [0, 1],$$

(for the mean and covariation function, see Karatzas and Shreve [24, formulas (3.2.21) and (3.2.23)]), and we also used that the third moment of a centered normally distributed random variable is 0. Similarly, for all $s \in [0, 1]$, we have

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_s \int_0^1 \mathcal{X}_t d[\mathcal{X}_t - mt] \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= ms \mathbb{E} \left(\int_0^1 \mathcal{X}_t \sqrt{\mathcal{Y}_t} d\mathcal{B}_t \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + \mathbb{E} \left(\int_0^s \sqrt{\mathcal{Y}_u} d\mathcal{B}_u \int_0^1 \mathcal{X}_t \sqrt{\mathcal{Y}_t} d\mathcal{B}_t \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \mathbb{E} \left(\int_0^s \mathcal{X}_u \mathcal{Y}_u du \middle| \mathcal{F}_1^{\mathcal{Y}} \right) = \int_0^s \mathbb{E}(\mathcal{X}_u \mathcal{Y}_u | \mathcal{F}_1^{\mathcal{Y}}) du = m \int_0^s u \mathcal{Y}_u du. \end{aligned}$$

Thus

$$\begin{aligned} \mathbb{E} \left(\mathcal{X}_s^2 \int_0^1 d[\mathcal{X}_t - mt] \right) &= 2ms \int_0^s \mathbb{E}(\mathcal{Y}_u) du, \\ \mathbb{E} \left(\mathcal{X}_s \int_0^1 \mathcal{X}_t d[\mathcal{X}_t - mt] \right) &= m \int_0^s u \mathbb{E}(\mathcal{Y}_u) du. \end{aligned}$$

Consequently,

$$\begin{aligned} \mathbb{E}(J) &= 2m \int_0^1 s \left(\int_0^s \mathbb{E}(\mathcal{Y}_u) du \right) ds - m \int_0^1 \left(\int_0^s u \mathbb{E}(\mathcal{Y}_u) du \right) ds \\ &= 2m \int_0^1 \left(\int_u^1 s ds \right) \mathbb{E}(\mathcal{Y}_u) ds - m \int_0^1 \left(\int_u^1 ds \right) u \mathbb{E}(\mathcal{Y}_u) du \end{aligned}$$

$$= m \int_0^1 (1-u) \mathbb{E}(\mathcal{Y}_u) du = ma \int_0^1 (1-u)u du.$$

Hence if $m \neq 0$, then $\mathbb{E}(J) \neq 0$, which clearly yields that $J \stackrel{\text{a.s.}}{=} 0$ is impossible.

If $m = 0$, then $\mathbb{E}(J) = 0$, and hence for proving $\mathbb{P}(J = 0) < 1$, it is enough to show that $\mathbb{E}(J^2) > 0$. Now J can be written in the form $J = J_1 - J_2$ with

$$J_1 := \int_0^1 \mathcal{X}_s^2 ds \int_0^1 d\mathcal{X}_t, \quad J_2 := \int_0^1 \mathcal{X}_s ds \int_0^1 \mathcal{X}_t d\mathcal{X}_t.$$

Clearly, $\mathbb{E}(J^2) = \mathbb{E}(J_1^2) - 2\mathbb{E}(J_1 J_2) + \mathbb{E}(J_2^2)$. Here $J_1 J_2$ can be written in the form

$$J_1 J_2 = \int_0^1 \int_0^1 \mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \left(\int_0^1 d\mathcal{X}_t \right) \left(\int_0^1 \mathcal{X}_t d\mathcal{X}_t \right) ds_1 ds_2,$$

hence

$$\mathbb{E}(J_1 J_2) = \int_0^1 \int_0^1 \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \left(\int_0^1 d\mathcal{X}_t \right) \left(\int_0^1 \mathcal{X}_t d\mathcal{X}_t \right) \right) ds_1 ds_2.$$

If $s_1, s_2 \in [0, 1]$, then, by (3.10),

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \left(\int_0^1 d\mathcal{X}_t \right) \left(\int_0^1 \mathcal{X}_t d\mathcal{X}_t \right) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \frac{1}{2} \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1 \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_s ds \right) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \frac{1}{2} \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) - \frac{1}{2} \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1 \int_0^1 \mathcal{Y}_s ds \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \frac{1}{2} \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) - \frac{1}{2} \left(\int_0^1 \mathcal{Y}_s ds \right) \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1 \middle| \mathcal{F}_1^{\mathcal{Y}} \right). \end{aligned}$$

Similarly to the proof of Proposition 3.1 one can check that conditionally of $\mathcal{F}_1^{\mathcal{Y}}$, the process $(\mathcal{X}_t)_{t \in [0,1]}$ is a centered Gauss process (especially having independent increments). Hence if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \mathbb{E} \left(\mathcal{X}_{s_1}^6 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 4 \mathbb{E} \left(\mathcal{X}_{s_1}^3 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + \mathbb{E} \left(\mathcal{X}_{s_1}^3 (\mathcal{X}_1 - \mathcal{X}_{s_2})^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &+ 4 \mathbb{E} \left(\mathcal{X}_{s_1}^5 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1}) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^5 (\mathcal{X}_1 - \mathcal{X}_{s_2}) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &+ 6 \mathbb{E} \left(\mathcal{X}_{s_1}^4 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^4 (\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &+ 6 \mathbb{E} \left(\mathcal{X}_{s_1}^3 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})(\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \end{aligned}$$

$$\begin{aligned}
& + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^3 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 (\mathcal{X}_1 - \mathcal{X}_{s_2}) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& + 9 \mathbb{E} \left(\mathcal{X}_{s_1}^4 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1}) (\mathcal{X}_1 - \mathcal{X}_{s_2}) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& + \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1}) (\mathcal{X}_1 - \mathcal{X}_{s_2})^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 (\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^3 (\mathcal{X}_1 - \mathcal{X}_{s_2}) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& + 6 \mathbb{E} \left(\mathcal{X}_{s_1}^3 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 (\mathcal{X}_1 - \mathcal{X}_{s_2}) \middle| \mathcal{F}_1^{\mathcal{Y}} \right).
\end{aligned}$$

Using that \mathcal{X} has independent increments and that the odd moments of a centered normally distributed random variable are 0, if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then

$$\begin{aligned}
& \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1^3 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& = \mathbb{E} \left(\mathcal{X}_{s_1}^6 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 6 \mathbb{E} \left(\mathcal{X}_{s_1}^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& \quad + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& \quad + \mathbb{E} \left(\mathcal{X}_{s_1}^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& \quad + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& = 15 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^3 + 18 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \\
& \quad + 9 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right) + 3 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right)^2 \\
& \quad + 3 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right).
\end{aligned}$$

One can also check that if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then

$$\begin{aligned}
\mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \mathcal{X}_1 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) & = \mathbb{E} \left(\mathcal{X}_{s_1}^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\
& = 3 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 + \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right).
\end{aligned}$$

Hence if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \left(\int_0^1 d\mathcal{X}_t \right) \left(\int_0^1 \mathcal{X}_t d\mathcal{X}_t \right) \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= 6 \left(\int_0^{s_1} \mathcal{Y}_u du \right)^3 + 7 \left(\int_0^{s_1} \mathcal{Y}_u du \right)^2 \left(\int_{s_1}^{s_2} \mathcal{Y}_u du \right) \\ &+ 3 \left(\int_0^{s_1} \mathcal{Y}_u du \right)^2 \left(\int_{s_2}^1 \mathcal{Y}_u du \right) + \left(\int_0^{s_1} \mathcal{Y}_u du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u du \right)^2 \\ &+ \left(\int_0^{s_1} \mathcal{Y}_u du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u du \right) \left(\int_{s_2}^1 \mathcal{Y}_u du \right). \end{aligned}$$

Similar expression hold in case of $s_1, s_2 \in [0, 1]$ with $s_1 \geq s_2$, we have to change s_1 by s_2 .

Moreover, J_1^2 can be written in the form

$$J_1^2 = \int_0^1 \int_0^1 \mathcal{X}_{s_1}^2 \mathcal{X}_{s_2}^2 \left(\int_0^1 d\mathcal{X}_t \right)^2 ds_1 ds_2,$$

hence

$$\mathbb{E}(J_1^2) = \int_0^1 \int_0^1 \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2}^2 \left(\int_0^1 d\mathcal{X}_t \right)^2 \right) ds_1 ds_2.$$

If $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2}^2 \left(\int_0^1 d\mathcal{X}_t \right)^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2}^2 \mathcal{X}_1^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \mathbb{E} \left(\mathcal{X}_{s_1}^6 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + \mathbb{E} \left(\mathcal{X}_{s_1}^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &+ 6 \mathbb{E} \left(\mathcal{X}_{s_1}^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &+ \mathbb{E} \left(\mathcal{X}_{s_1}^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &+ \mathbb{E} \left(\mathcal{X}_{s_1}^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \mathbb{E} \left((\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= 15 \left(\int_0^{s_1} \mathcal{Y}_u du \right)^3 + 3 \left(\int_0^{s_1} \mathcal{Y}_u du \right)^2 \left(\int_{s_2}^1 \mathcal{Y}_u du \right) \\ &+ 18 \left(\int_0^{s_1} \mathcal{Y}_u du \right)^2 \left(\int_{s_1}^{s_2} \mathcal{Y}_u du \right) + 3 \left(\int_0^{s_1} \mathcal{Y}_u du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u du \right)^2 \\ &+ \left(\int_0^{s_1} \mathcal{Y}_u du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u du \right) \left(\int_{s_2}^1 \mathcal{Y}_u du \right). \end{aligned}$$

Similar expression hold in case of $s_1, s_2 \in [0, 1]$ with $s_1 \geq s_2$, we have to change s_1 by s_2 .

Furthermore, using (3.10), J_2^2 can be written in the form

$$J_2^2 = \frac{1}{4} \int_0^1 \int_0^1 \mathcal{X}_{s_1} \mathcal{X}_{s_2} \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_u \, du \right)^2 \, ds_1 \, ds_2,$$

hence

$$\mathbb{E}(J_2^2) = \frac{1}{4} \int_0^1 \int_0^1 \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_u \, du \right)^2 \right) \, ds_1 \, ds_2.$$

Here if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then we have

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_u \, du \right)^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \mathcal{X}_1^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) - 2 \left(\int_0^1 \mathcal{Y}_u \, du \right) \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \mathcal{X}_1^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ & \quad + \left(\int_0^1 \mathcal{Y}_u \, du \right)^2 \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \middle| \mathcal{F}_1^{\mathcal{Y}} \right), \end{aligned}$$

where, using the arguments as above, one can check that

$$\begin{aligned} & \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \mathcal{X}_1^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= \mathbb{E} \left(\mathcal{X}_{s_1}^6 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 5 \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ & \quad + \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_1 - \mathcal{X}_{s_2})^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 10 \mathbb{E} \left(\mathcal{X}_{s_1}^4 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ & \quad + 6 \mathbb{E} \left(\mathcal{X}_{s_1}^4 (\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 18 \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 (\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ &= 15 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^3 + 15 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right)^2 \\ & \quad + 3 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right)^2 + 30 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \\ & \quad + 18 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right) \\ & \quad + 18 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right), \end{aligned}$$

and

$$\begin{aligned} \mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \mathcal{X}_1^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) &= \mathbb{E} \left(\mathcal{X}_{s_1}^4 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) + 3 \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_{s_2} - \mathcal{X}_{s_1})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \\ & \quad + \mathbb{E} \left(\mathcal{X}_{s_1}^2 (\mathcal{X}_1 - \mathcal{X}_{s_2})^2 \middle| \mathcal{F}_1^{\mathcal{Y}} \right) \end{aligned}$$

$$\begin{aligned}
 &= 3 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 + 3 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \\
 &\quad + \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right),
 \end{aligned}$$

and

$$\mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \mid \mathcal{F}_1^{\mathcal{Y}} \right) = \mathbb{E} \left(\mathcal{X}_{s_1}^2 \mid \mathcal{F}_1^{\mathcal{Y}} \right) = \int_0^{s_1} \mathcal{Y}_u \, du.$$

Hence, by an easy calculation, if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then

$$\begin{aligned}
 &\mathbb{E} \left(\mathcal{X}_{s_1} \mathcal{X}_{s_2} \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_u \, du \right)^2 \mid \mathcal{F}_1^{\mathcal{Y}} \right) \\
 &= 10 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^3 + 10 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right)^2 \\
 &\quad + 20 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \\
 &\quad + 12 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right) \\
 &\quad + 2 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right)^2 \\
 &\quad + 12 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right).
 \end{aligned}$$

Similar expression hold in case of $s_1, s_2 \in [0, 1]$ with $s_1 \geq s_2$, we have to change s_1 by s_2 .

Hence if $s_1, s_2 \in [0, 1]$ with $s_1 \leq s_2$, then we have

$$\begin{aligned}
 &\mathbb{E} \left(\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2}^2 \left(\int_0^1 d\mathcal{X}_t \right)^2 - 2\mathcal{X}_{s_1}^2 \mathcal{X}_{s_2} \left(\int_0^1 d\mathcal{X}_t \right) \left(\int_0^1 \mathcal{X}_t \, d\mathcal{X}_t \right) \right. \\
 &\quad \left. + \frac{1}{4} \mathcal{X}_{s_1} \mathcal{X}_{s_2} \left(\mathcal{X}_1^2 - \int_0^1 \mathcal{Y}_u \, du \right)^2 \mid \mathcal{F}_1^{\mathcal{Y}} \right) \\
 &= \frac{11}{2} \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^3 + 9 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \\
 &\quad + \frac{7}{2} \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right)^2 + \frac{1}{2} \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right)^2 \\
 &\quad + 2 \left(\int_0^{s_1} \mathcal{Y}_u \, du \right) \left(\int_{s_1}^{s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_2}^1 \mathcal{Y}_u \, du \right),
 \end{aligned}$$

and a similar expression hold in case of $s_1, s_2 \in [0, 1]$ with $s_1 \geq s_2$, we have to change s_1 by s_2 . Then $\mathbb{E}(J^2) = \mathbb{E}(J_1^2) - 2\mathbb{E}(J_1 J_2) + \mathbb{E}(J_2^2)$ takes the form

$$\begin{aligned} & \int_0^1 \int_0^1 \mathbb{E} \left(\frac{11}{2} \left(\int_0^{s_1 \wedge s_2} \mathcal{Y}_u \, du \right)^3 + 9 \left(\int_0^{s_1 \wedge s_2} \mathcal{Y}_u \, du \right)^2 \left(\int_{s_1 \wedge s_2}^{s_1 \vee s_2} \mathcal{Y}_u \, du \right) \right. \\ & + \frac{7}{2} \left(\int_0^{s_1 \wedge s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_1 \wedge s_2}^{s_1 \vee s_2} \mathcal{Y}_u \, du \right)^2 + \frac{1}{2} \left(\int_0^{s_1 \wedge s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_1 \vee s_2}^1 \mathcal{Y}_u \, du \right)^2 \\ & \left. + 2 \left(\int_0^{s_1 \wedge s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_1 \wedge s_2}^{s_1 \vee s_2} \mathcal{Y}_u \, du \right) \left(\int_{s_1 \vee s_2}^1 \mathcal{Y}_u \, du \right) \right) ds_1 ds_2 > 0, \end{aligned}$$

where for the last inequality we used that

$$\mathbb{E} \left(\left(\int_0^{s_1 \wedge s_2} \mathcal{Y}_u \, du \right)^i \left(\int_{s_1 \wedge s_2}^{s_1 \vee s_2} \mathcal{Y}_u \, du \right)^j \left(\int_{s_1 \vee s_2}^1 \mathcal{Y}_u \, du \right)^k \right) > 0$$

for $i, j, k \in \{0, 1, 2, 3\}$, which follows by $a \in \mathbb{R}_{++}$ and that $\mathbb{P}(Y_t \geq 0 \text{ for all } t \in \mathbb{R}_+) = 1$. Consequently, we conclude $\mathbb{E}(J^2) > 0$, which clearly yields that $J \stackrel{\text{a.s.}}{=} 0$ is impossible.

References

- [1] ANDERSEN, L. B. G. and PITERBARG, V. V. (2007). Moment explosions in stochastic volatility models. *Finance and Stochastics* **11**(1) 29–50. [MR2284011](#)
- [2] ANDERSON, T. W. (1959). On asymptotic distributions of estimates of parameters of stochastic difference equations. *The Annals of Mathematical Statistics* **30** 676–687. [MR0107347](#)
- [3] BARCZY, M., ISPÁNY, M. and PAP, G. (2010). Asymptotic behavior of CLS estimators of autoregressive parameter for nonprimitive unstable INAR(2) models. Available on the ArXiv: <http://arxiv.org/abs/1006.4641>
- [4] BARNDORFF-NIELSEN, O. and SHEPHARD, N. (2001). Non-Gaussian Ornstein-Uhlenbeck-based models and some of their uses in financial economics. *Journal of the Royal Statistical Society, Series B* **63** 167–241. [MR1841412](#)
- [5] BEN ALAYA, M. and KEBAIER, M. (2012). Parameter estimation for the square root diffusions: ergodic and nonergodic cases. *Stochastic Models* **28**(4) 609–634.
- [6] BEN ALAYA, M. and KEBAIER, M. (2012). Asymptotic behavior of the maximum likelihood estimator for ergodic and nonergodic square-root diffusions. Available on the HAL: <http://hal.archives-ouvertes.fr/hal-00640053>
- [7] CARR, P. and WU, L. (2003). The finite moment log stable process and option pricing. *The Journal of Finance* **58** 753–777.

- [8] CHEN, H. and JOSLIN, S. (2012). Generalized transform analysis of affine processes and applications in finance. *Review of Financial Studies* **25**(7) 2225–2256.
- [9] CUCHIERO, C., FILIPOVIĆ, D. and TEICHMANN, J. (2010). Affine Models. Contribution to *Encyclopedia of Quantitative Finance*. John Wiley & Sons, Ltd.
- [10] DAWSON, D. A. and LI, Z. (2006). Skew convolution semigroups and affine processes. *The Annals of Probability* **34**(3) 1103–1142. [MR2243880](#)
- [11] DUFFIE, D., FILIPOVIĆ, D. and SCHACHERMAYER, W. (2003). Affine processes and applications in finance. *Annals of Applied Probability* **13** 984–1053. [MR1994043](#)
- [12] ETHIER, S. N. and KURTZ, T. G. (1986). *Markov Processes: Characterization and Convergence*. Wiley. [MR0838085](#)
- [13] FILIPOVIĆ, D. and MAYERHOFER, E. (2009). Affine diffusion processes: theory and applications. In Albrecher, Hansjörg (ed.) et al., *Advanced Financial Modelling, Radon Series on Computational and Applied Mathematics* **8** 125–164, Berlin, Walter de Gruyter. [MR2648460](#)
- [14] FRIZ, P. K. and KELLER-RESSEL, M. (2010). Moment Explosions in Stochastic Volatility Models. Contribution to *Encyclopedia of Quantitative Finance*. John Wiley & Sons, Ltd.
- [15] GLASSERMAN, P. and KIM, K.-K. (2010). Moment explosions and stationary distributions in affine diffusion models. *Mathematical Finance* **20**(1) 1–33. [MR2599675](#)
- [16] HAMILTON, J. D. (1994). *Time Series Analysis*. Princeton University Press, Princeton. [MR1278033](#)
- [17] HESTON, S. (1993). A closed-form solution for options with stochastic volatilities with applications to bond and currency options. *The Review of Financial Studies* **6** 327–343.
- [18] HU, Y. and LONG, H. (2009). Least squares estimator for Ornstein–Uhlenbeck processes driven by α -stable motions. *Stochastic Processes and their Applications* **119** 2465–2480. [MR2532208](#)
- [19] HU, Y. and LONG, H. (2009). On the singularity of least squares estimator for mean-reverting α -stable motions. *Acta Mathematica Scientia* **29B**(3) 599–608. [MR2514363](#)
- [20] HUANG, J., MA, C. and ZHU, C. (2011). Estimation for discretely observed continuous state branching processes with immigration. *Statistics and Probability Letters* **81** 1104–1111. [MR2803751](#)
- [21] IKEDA, N. and WATANABE, S. (1981). *Stochastic Differential Equations and Diffusion Processes*. North-Holland Publishing Company. [MR0637061](#)
- [22] JACOD, J. and SHIRYAEV, A. N. (2003). *Limit Theorems for Stochastic Processes*, 2nd ed. Springer-Verlag, Berlin. [MR1943877](#)
- [23] JENA, R. P., KIM, K.-K. and XING, H. (2012). Long-term and blow-up behaviors of exponential moments in multi-dimensional affine diffusions. *Stochastic Processes and their Applications* **122**(8) 2961–2993. [MR2931348](#)
- [24] KARATZAS, I. and SHREVE, S.E. (1991). *Brownian Motion and Stochastic Calculus, 2nd edition*. Springer-Verlag. [MR1121940](#)

- [25] KELLER-RESSEL, M., SCHACHERMAYER, W. and TEICHMANN, J. (2011). Affine processes are regular. *Probability Theory and Related Fields* **151** 591–611. [MR2851694](#)
- [26] KELLER-RESSEL, M., SCHACHERMAYER, W. and TEICHMANN, J. (2012). Regularity of affine processes on general state spaces. Available on the ArXiv: <http://arxiv.org/abs/1105.0632v2>
- [27] LI, Z. (2011). *Measure-Valued Branching Markov Processes*. Springer-Verlag, Heidelberg. [MR2760602](#)
- [28] MA, C. (2009). A limit theorem of two-type Galton–Watson branching processes with immigration. *Statistics and Probability Letters* **79** 1710–1716. [MR2547941](#)
- [29] MANN, H. B. and WALD, A. (1943). On the statistical treatment of linear stochastic difference equations. *Econometrica* **11**(3/4) 173–220. [MR0009291](#)
- [30] OVERBECK, L. and RYDÉN, T. (1997). Estimation in the Cox-Ingersoll-Ross model. *Econometric Theory* **13**(3) 430–461. [MR1455180](#)
- [31] OVERBECK, L. (1998). Estimation for continuous branching processes. *Scandinavian Journal of Statistics* **25**(1) 111–126. [MR1614256](#)
- [32] PROTTER, P. (2004). *Stochastic Integration and Differential Equations, 2nd edition*. Springer-Verlag. [MR2020294](#)
- [33] VAN DER VAART, A. W. (1998). *Asymptotic Statistics*. Cambridge University Press, Cambridge. [MR1652247](#)
- [34] WEI, C. Z. and WINNICKI, J. (1989). Some asymptotic properties of branching processes with immigration. *Stochastic Processes and their Applications* **31** 261–282. [MR0998117](#)
- [35] WEI, C. Z. and WINNICKI, J. (1990). Estimation of the means in the branching processes with immigration. *The Annals of Mathematical Statistics* **18**(4) 1757–1773. [MR1074433](#)
- [36] WHITE, J. S. (1958). The limiting distribution of the serial correlation coefficient in the explosive case. *The Annals of Mathematical Statistics* **29** 1188–1197. [MR0100952](#)