

Averaged vs. quenched large deviations and entropy for random walk in a dynamic random environment*

Firas Rassoul-Agha[†] Timo Seppäläinen[‡] Atilla Yilmaz[§]

Abstract

We consider random walk with bounded jumps on a hypercubic lattice of arbitrary dimension in a dynamic random environment. The environment is temporally independent and spatially translation invariant. We study the rate functions of the level-3 averaged and quenched large deviation principles from the point of view of the particle. In the averaged case the rate function is a specific relative entropy, while in the quenched case it is a Donsker-Varadhan type relative entropy for Markov processes. We relate these entropies to each other and seek to identify the minimizers of the level-3 to level-1 contractions in both settings. Motivation for this work comes from variational descriptions of the quenched free energy of directed polymer models where the same Markov process entropy appears.

Keywords: random walk; dynamic random environment; large deviations; averaged; quenched; empirical process; Donsker-Varadhan relative entropy; specific relative entropy; Doob h -transform; nonstationary process.

AMS MSC 2010: 60K37; 60F10; 82C41; 82C44.

Submitted to EJP on July 24, 2016, final version accepted on June 1, 2017.

Supersedes arXiv:1607.07000v1.

1 Introduction

After surveying the background of the present work, this introductory section describes the random walk in a dynamic random environment (RWDRE) model and then some general notions such as large deviation principles and the point of view of the particle. The section concludes with an overview of the rest of the paper.

*F. Rassoul-Agha was partially supported by National Science Foundation grant DMS-1407574 and by Simons Foundation grant 306576. T. Seppäläinen was partially supported by National Science Foundation grants DMS-1306777 and DMS-1602486, by Simons Foundation grant 338287, and by the Wisconsin Alumni Research Foundation. A. Yilmaz was partially supported by European Union FP7 Marie Curie Career Integration Grant no. 322078.

[†]Department of Mathematics, University of Utah, 155 South 1400 East, Salt Lake City, UT 84109, USA.
E-mail: firas@math.utah.edu, <http://www.math.utah.edu/~firas>

[‡]Department of Mathematics, University of Wisconsin-Madison, 480 Lincoln Dr., Madison, WI 53706, USA.
E-mail: seppalai@math.wisc.edu, <http://www.math.wisc.edu/~seppalai>

[§]Department of Mathematics, Koç University, Rumelifeneri Yolu, Sarıyer, Istanbul 34450, Turkey.
E-mail: atillayilmaz@ku.edu.tr, <http://home.ku.edu.tr/~atillayilmaz>

1.1 Background

This paper studies an entropy function for Markov processes that appears in random medium models. We give here some background motivation. A much-studied model is the random path in a random potential model, also called the polymer model. The random environment ω comes from a probability space $(\Omega, \mathfrak{G}, \mathbb{P})$ with an ergodic group action $\{T_x\}_{x \in \mathbb{Z}^d}$. The random path is a classical random walk $(X_k)_{k \geq 0}$ on \mathbb{Z}^d whose law is independent of the environment. The potential $V(\omega, z)$ is a function of ω and a step z of the random walk. A key quantity is the limiting *quenched free energy*

$$g(V) = \lim_{n \rightarrow \infty} n^{-1} \log E_0^{RW} \left[e^{\sum_{k=0}^{n-1} V(T_{X_k} \omega, X_{k+1} - X_k)} \right] \tag{1.1}$$

where E_0^{RW} is the expectation of the random walk and ω is fixed. The limit exists for \mathbb{P} -almost every ω under hypotheses on the moments of V and the degree of mixing of \mathbb{P} .

The limit $g(V)$ can be calculated only in a handful of exactly solvable models that exist only in 1 + 1 dimension. More generally, properties of $g(V)$ have remained an insurmountable problem. This question includes the positive temperature version of the question of understanding limit shapes of stochastic growth models such as first- and last-passage percolation. The latter question has also remained insurmountable since the origins of the subject over 50 years ago, except for a few exactly solvable models in 1 + 1 dimension. For surveys of models of type (1.1), see [10, 16].

Our article [43] introduced two variational formulas for $g(V)$. Let $p(z)$ be the jump kernel of the underlying random walk. The first formula

$$g(V) = \inf_F \mathbb{P}\text{-ess sup}_{\omega} \log \sum_z p(z) e^{V(\omega, z) + F(\omega, z)} \tag{1.2}$$

expresses $g(V)$ as an infimum over the $L^1(\mathbb{P})$ closure of gradients $F(\omega, z) = f(T_z \omega) - f(\omega)$, which we called the space of *cocycles*. Since this formula is not the topic of the present paper, we refer to [24, 43, 44] for precise definitions.

The second formula gives $g(V)$ as the dual of an entropy adapted to the point of view of the particle:

$$g(V) = \sup \{ E^\mu[V] - H(\mu) : \mu_\Omega \ll \mathbb{P}, E^\mu[V^-] < \infty \}. \tag{1.3}$$

The supremum is over probability measures μ on $\Omega \times \{\text{steps}\}$ with a natural invariance property and with a \mathbb{P} -absolutely continuous Ω -marginal μ_Ω . The entropy is given by

$$H(\mu) = \int_\Omega \sum_z \mu(d\omega, z) \log \frac{\mu(z | \omega)}{p(z)}. \tag{1.4}$$

Formula (1.3) was proved in [43], and this formulation is Theorem 7.5 in [24].

Article [24] extended these formulas from positive to zero temperature, that is, to last-passage percolation models. The goal is to shed light on $g(V)$ and limit shapes through the variational formulas. The relationship between formulas (1.2) and (1.3) is well understood presently only for directed polymers in weak disorder (Examples 3.7 and 7.7 in [24]) and in periodic environments (Section 8 in [24]).

Here is a brief overview of the current state of the study of these formulas. The cocycle variational formula (1.2) has been studied in several subsequent papers while the entropy formula (1.3) has received no serious attention before the present paper. [44] shows that (1.2) always has a minimizer and uses the minimizer(s) to characterize weak and strong disorder of directed polymers. [27] proves the existence of Busemann functions for the exactly solvable 1+1 dimensional log-gamma polymer and shows that these provide minimizing cocycles for (1.2) and also a limiting polymer measure

for infinite paths. [25, 26] construct the minimizing cocycles for the 2-dimensional corner growth model with general i.i.d. weights and use these to investigate Busemann functions, geodesics and the competition interface. These notions have become central in the field of random medium models over the last twenty years, beginning with the work of Newman in the early 1990s on the geodesics of first-passage percolation [34].

In the current paper we begin the study of the entropy (1.4). This entropy is the level-2 projection of an entropy that appears in the rate function of a level-3 quenched large deviation principle (LDP) for RWDRE. (See (2.2) and Theorem 2.2 in Section 2.) In fact, it was shown in [43] that the rate function of the induced LDP at level-2 (via contraction) is the convex dual of the map $h \mapsto g(h + V) - g(V)$, with V as in (1.8). We study the entropy in this large deviations context. In particular, we consider its relation to the entropy that serves as the rate function for a level-3 averaged LDP.

The *point-to-point* version of the quenched free energy (1.1) is

$$g(V, \xi) = \lim_{n \rightarrow \infty} n^{-1} \log E_0^{RW} \left[e^{\sum_{k=0}^{n-1} V(T_{X_k} \omega, X_{k+1} - X_k)}, X_n = [n\xi] \right] \tag{1.5}$$

defined for ξ in the convex hull of the support of the kernel $p(z)$, and where $[n\xi]$ is a lattice point that approximates $n\xi$ under certain constraints (see [41]) and is reachable from the origin in n steps. The entropy variational formula now takes the form

$$g(V, \xi) = \sup \{ E^\mu[V] - H(\mu) : \mu_\Omega \ll \mathbb{P}, E^\mu[V^-] < \infty, E^\mu[Z_1] = \xi \} \tag{1.6}$$

where Z_1 is the step variable under distribution μ . Formula (1.6) was proved in [41] for a directed walk in an i.i.d. environment and a local potential $V \in L^{d+\varepsilon}(\mathbb{P})$ for $\varepsilon > 0$. This formulation is Theorem 7.6 in [24].

Minimizing entropy under a mean step condition $E^\mu[Z_1] = \xi$ as in (1.6) is also done in the level-3 to level-1 contraction in large deviation theory. For this reason the main focus of the present paper is to study these contractions, both averaged and quenched. The averaged contraction can be understood completely. Then we seek to characterize when the averaged and quenched contractions lead to the same level-1 rate function and have the same minimizers.

The quenched rate function is hard to study. It begins with an entropy of a familiar type. But this entropy is corrected in a singular manner to account for the environment distribution \mathbb{P} , and then regularized again to be lower semicontinuous. The opaqueness of the l.s.c. regularization makes it difficult to analyze examples. By simplifying the situation so that the environment varies only temporally we can describe fully also the quenched contraction. We discover that the connection between the averaged and quenched rate functions can break down rather spectacularly. This part of the paper illuminates earlier large deviation work by Comets [8] and one of the authors [2, 47] that appears in the equilibrium statistical mechanics of disordered Gibbs measures.

The present paper studies only random walk in a dynamic random environment while connections to polymer models are left for future work. Our results in Section 3 begin with the level-3 averaged LDP from the point of view of the particle and the existence of the relevant limiting specific relative entropy. After understanding the contraction from the level-3 to level-1 averaged LDP we turn to study the quenched rate functions.

1.2 The model

Consider the d -dimensional hypercubic lattice \mathbb{Z}^d with an arbitrary $d \in \mathbb{N} = \{1, 2, \dots\}$. Fix a finite $\mathcal{R} \subset \mathbb{Z}^d$ with at least two elements and let

$$\mathcal{P} = \{q : \mathcal{R} \rightarrow [0, 1] : \sum_{z \in \mathcal{R}} q(z) = 1\} \tag{1.7}$$

denote the set of probability measures on \mathcal{R} . Elements of $\Omega = \mathcal{P}^{\mathbb{Z} \times \mathbb{Z}^d}$ are called space-time environments and they are of the form $\omega = (\omega_{i,x})_{(i,x) \in \mathbb{Z} \times \mathbb{Z}^d}$. Each $\omega \in \Omega$ defines a time-inhomogeneous discrete-time Markov chain $(X_i)_{i \geq 0}$ on \mathbb{Z}^d for which $X_0 = 0$ and the transition probability from state x to y at time i is

$$\pi_{i,i+1}(x, y | \omega) = \begin{cases} \omega_{i,x}(y - x) & \text{if } y - x \in \mathcal{R}, \\ 0 & \text{otherwise.} \end{cases}$$

If ω is randomly sampled from a probability distribution \mathbb{P} on (Ω, \mathfrak{S}) rather than being deterministic, then $(X_i)_{i \geq 0}$ is a random walk (RW) in a dynamic (or space-time) random environment, which we abbreviate as RWDRE. Here, \mathfrak{S} is the Borel σ -algebra with respect to (w.r.t.) the product topology on Ω .

RWDRE induces a probability measure $P_0(d\omega, dz) = \mathbb{P}(d\omega)P_0^\omega(dz)$ on the space $\Omega_{\mathbb{N}} = \Omega \times \mathcal{R}^{\mathbb{N}}$ of environments and walks (starting at the origin). Here, $\mathbf{z} = (z_i)_{i \geq 1} \in \mathcal{R}^{\mathbb{N}}$ is a sequence of steps, and P_0^ω is the quenched path measure defined by

$$P_0^\omega(z_1, \dots, z_n) = \prod_{i=0}^{n-1} \pi_{i,i+1}(x_i, x_{i+1} | \omega), \quad n \geq 1, x_0 = 0 \text{ and } x_{i+1} = x_i + z_{i+1}.$$

The marginal of P_0 on $\mathcal{R}^{\mathbb{N}}$ is called the averaged path measure and also denoted by P_0 whenever no confusion occurs. \mathbb{E}, E_0 and E_0^ω stand for expectation under \mathbb{P}, P_0 and P_0^ω , respectively. In general, we will write $E^\mu[f]$ or $\langle f, \mu \rangle$ for the integral of a function f against a probability measure μ .

Denote the entire spatial environment at a given time $i \in \mathbb{Z}$ by $\bar{\omega}_i = (\omega_{i,x} : x \in \mathbb{Z}^d)$. Let $(T_y^s)_{y \in \mathbb{Z}^d}$ be the group of spatial translations, defined by $(T_y^s \bar{\omega}_i)_x = \omega_{i,x+y}$ for $x, y \in \mathbb{Z}^d$. Throughout the article, we will make the following underlying assumptions.

- *Temporal independence:* $(\bar{\omega}_i)_{i \in \mathbb{Z}}$ are independent and identically distributed (i.i.d.) under \mathbb{P} with a common distribution \mathbb{P}_s on $\mathcal{P}^{\mathbb{Z}^d}$, i.e., $\mathbb{P} = (\mathbb{P}_s)^{\otimes \mathbb{Z}}$. (The subscript of \mathbb{P}_s stands for “spatial”.)
- *Spatial translation invariance:* \mathbb{P}_s is invariant under $(T_y^s)_{y \in \mathbb{Z}^d}$.

These two conditions are of course satisfied when $(\omega_{i,x})_{(i,x) \in \mathbb{Z} \times \mathbb{Z}^d}$ are i.i.d. However, restricting to that special case would not change the statements or the proofs of our results in this paper (except those in Section 3.7 on spatially constant environments). Moreover, it should be relatively straightforward to adapt our results to various discrete-time continuous-space models (such as RWDRE on \mathbb{R}^d considered in [4, 29]) where spatial independence is not applicable. Note in particular that we do not assume ergodicity under spatial translations.

For the averaged results, the only condition we impose on the one-step range \mathcal{R} of the walk is $2 \leq |\mathcal{R}| < \infty$. ($|\mathcal{R}|$ is the number of elements in the set \mathcal{R} . The case $|\mathcal{R}| = 1$ is trivial.) We will assume without loss of generality that $\mathbb{P}(\omega_{0,0}(z) > 0) > 0$ for $z \in \mathcal{R}$. (Otherwise, we can replace \mathcal{R} by $\{z \in \mathcal{R} : \mathbb{P}(\omega_{0,0}(z) > 0) > 0\}$.) Our quenched results will require various ellipticity conditions which we will indicate as needed in their statements. These ellipticity conditions will be compared in Remark 3.13.

As the name suggests, RWDRE is a variant of the much-studied random walk in a random environment (RWRE) model (see [53] for a survey). In fact, $(i, X_i)_{i \geq 0}$ can be viewed as a directed RWRE on \mathbb{Z}^{d+1} because its component in the direction of $(1, 0, \dots, 0)$ is strictly increasing. This directedness simplifies certain aspects of the analysis of the model. Most notably, RWDRE under the averaged measure P_0 is a classical RW on \mathbb{Z}^d with transition probabilities $\hat{q}(z) = \mathbb{E}[\omega_{0,0}(z)] > 0$. In particular, the strong law of large numbers (LLN) and Donsker’s invariance principle (IP) hold for the averaged walk. Since

any P_0 -almost sure statement holds P_0^ω -almost surely for \mathbb{P} -a.e. ω , there is no need for a separate strong LLN for the quenched walk. On the other hand, an averaged IP does not a priori imply a quenched one. Nevertheless, for the i.i.d. case, there is an IP under P_0^ω for \mathbb{P} -a.e. ω [38]. In stark contrast to these limit theorems, for (undirected) RWRE the validity of even the LLN is an open problem in the i.i.d. case when $d \geq 3$. See [3, 6] for two of the best sufficient conditions in the literature.

Finally, RWDRE is a special case of the random path in a random potential model (on \mathbb{Z}^{d+1}) that we described in Section 1.1 with

$$V(T_{i,x}\omega, z) = \log \left(\frac{\omega_{i,x}(z)}{p(z)} \right) \tag{1.8}$$

for every $\omega \in \Omega$, $i \in \mathbb{Z}$, $x \in \mathbb{Z}^d$ and $z \in \mathcal{R}$. The limiting quenched free energy $g(V)$ for this choice of potential is trivially zero. However, it is not $g(V)$ but the map $h \mapsto g(h+V) - g(V)$ and certain variants of it that play a central role in describing the large deviation behavior of these models (see [43, Theorem 3.1] for details).

1.3 Large deviation principles, the point of view of the particle, and empirical measures

Recall that a sequence $(Q_n)_{n \geq 1}$ of Borel probability measures on a topological space \mathbb{X} is said to satisfy a large deviation principle (LDP) with (exponential scale n and) rate function $I : \mathbb{X} \rightarrow [0, \infty]$ if I is lower semicontinuous, and for any measurable set G ,

$$-\inf_{x \in G^\circ} I(x) \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \log Q_n(G) \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log Q_n(G) \leq -\inf_{x \in \overline{G}} I(x).$$

G° is the topological interior of G and \overline{G} its topological closure. See [14, 15, 42] for general background regarding large deviations.

In the context of RWDRE, the LDP for $(P_0(X_n/n \in \cdot))_{n \geq 1}$ is nothing but Cramér’s theorem for classical multidimensional RW (see, e.g., [42, Chapter 4]), with rate function $I_{1,a} : \mathbb{R}^d \rightarrow [0, \infty]$ given by

$$I_{1,a}(\xi) = \sup_{\rho \in \mathbb{R}^d} \{ \langle \rho, \xi \rangle - \log \phi_a(\rho) \} = (\log \phi_a)^*(\xi), \tag{1.9}$$

the convex conjugate of the logarithm of the moment generating function

$$\phi_a(\rho) = \sum_{z \in \mathcal{R}} \hat{q}(z) e^{\langle \rho, z \rangle}, \tag{1.10}$$

where $\langle \cdot, \cdot \rangle$ denotes inner product. This is an averaged LDP, hence the subscript a . (The other subscript of $I_{1,a}$ stands for level-1 which is explained two paragraphs below.) Establishing the analogous quenched LDP for $(P_0^\omega(X_n/n \in \cdot))_{n \geq 1}$ and identifying the rate function is more arduous. It involves considering certain empirical measures from the point of view (POV) of the particle which we introduce next.

Define space-time translations $(T_{j,y})_{(j,y) \in \mathbb{Z} \times \mathbb{Z}^d}$ on Ω by $(T_{j,y}\omega)_{i,x} = \omega_{i+j,x+y}$. Then, $(T_{i,X_i}\omega)_{i \geq 0}$ is a discrete-time Markov chain taking values in Ω , and its transition probability from state ω to state ω' is given by

$$\bar{\pi}(\omega' | \omega) = \sum_{z \in \mathcal{R} : T_{1,z}\omega = \omega'} \pi_{0,1}(0, z | \omega).$$

Every limit theorem about this so-called *environment Markov chain* implies a corresponding limit theorem for the walk. This general and robust approach was first introduced in

the context of diffusion processes with random coefficients [35] and was later successfully adapted to RWRE (see for example [31, 30, 13, 36, 39, 50]).

In light of the previous paragraph, the large deviation behavior of RWDRE can be analyzed via various statistics of either the walk itself or the environment Markov chain. Among these statistics, the empirical velocity X_n/n is the coarsest one, and hence its large deviation analysis is referred to as level-1. Finer statistics are provided by the occupation measure

$$L_n = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{T_i, X_i \omega}$$

which records the environments seen from the POV of the particle. The pair-empirical measure

$$L_n^2 = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{T_i, X_i \omega, Z_{i+1}}$$

goes one step further by essentially keeping track of the pairs of consecutive environments that the particle sees. (In the Markov chain literature, the pair-empirical measure typically refers to $\frac{1}{n} \sum_{i=0}^{n-1} \delta_{T_i, X_i \omega, T_{i+1}, X_{i+1} \omega}$ which is measurable w.r.t. our choice of L_n^2 .) Pairs can be replaced with ℓ -tuples for any $\ell \geq 2$ to define more detailed empirical measures. Large deviations of each of these empirical measures are called level-2. Finally, level-3 involves the so-called empirical process

$$L_n^\infty = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{T_i, X_i \omega, \theta^i \mathbf{Z}}. \tag{1.11}$$

Here and throughout, $\mathbf{Z} = (Z_i)_{i \geq 1}$ denotes the sequence of steps $Z_i = X_i - X_{i-1}$ of the random path $(X_i)_{i \geq 0}$, and θ is the forward shift on sequences, i.e., $(\theta \mathbf{Z})_j = Z_{j+1}$ for every $j \in \mathbb{N}$. Under the topology of weak convergence of measures, the empirical process contains precisely the same information as all of the empirical measures for ℓ -tuples combined. Level-1,2,3 large deviations for Markov processes were established (under certain conditions) in a series of papers by Donsker and Varadhan [18, 19, 20]. The level terminology was introduced later in [22].

1.4 Further notation for steps, environments and σ -algebras

Throughout, for any bi-infinite sequence $\bar{z} = (\dots, z_{-2}, z_{-1}, z_0, z_1, z_2, \dots) \in \mathcal{R}^{\mathbb{Z}}$ of steps and any pair of indices $-\infty < i \leq j < \infty$, we write

$$z_{i,j} = (z_i, z_{i+1}, \dots, z_j), \quad z_{i,\infty} = (z_i, z_{i+1}, z_{i+2}, \dots) \quad \text{and} \quad z_{-\infty,j} = (\dots, z_{j-2}, z_{j-1}, z_j).$$

We also use $\mathbf{z} = z_{1,\infty}$ and $\bar{z} = z_{-\infty,\infty}$. Similarly, for any environment $\omega = (\bar{\omega}_i)_{i \in \mathbb{Z}}$ and any pair of indices $-\infty < k \leq \ell < \infty$,

$$\bar{\omega}_{k,\ell} = (\bar{\omega}_k, \bar{\omega}_{k+1}, \dots, \bar{\omega}_\ell), \quad \bar{\omega}_{k,\infty} = (\bar{\omega}_k, \bar{\omega}_{k+1}, \bar{\omega}_{k+2}, \dots) \quad \text{and} \quad \bar{\omega}_{-\infty,\ell} = (\dots, \bar{\omega}_{\ell-2}, \bar{\omega}_{\ell-1}, \bar{\omega}_\ell).$$

We use this notation to introduce the σ -algebras

$$\mathcal{A}_{k,\ell}^{i,j} = \sigma\{\bar{\omega}_{k,\ell-1}, Z_{i+1,j}\} \quad \text{and} \quad \mathfrak{S}_{k,\ell} = \sigma\{\bar{\omega}_{k,\ell-1}\}$$

on appropriate spaces, for $-\infty \leq i < j \leq \infty$ and $-\infty \leq k < \ell \leq \infty$. The reason for the indexing convention is that the distribution of step Z_{n+1} is part of environment $\bar{\omega}_n$. Note also that $\mathcal{A}_{-\infty,\infty}^{0,\infty}$ and $\mathcal{A}_{-\infty,\infty}^{-\infty,\infty}$ are the Borel σ -algebras (w.r.t. the product topology) on $\Omega_{\mathbb{N}} = \Omega \times \mathcal{R}^{\mathbb{N}}$ and $\Omega_{\mathbb{Z}} = \Omega \times \mathcal{R}^{\mathbb{Z}}$, respectively. For any σ -algebra \mathcal{F} , the space of bounded and \mathcal{F} -measurable functions is denoted by $b\mathcal{F}$.

1.5 Content and organization of the article

Section 2 reviews previous results on large deviations for RWDRE. The new results are in Section 3. The paper is organized so that the results of Section 3. n are proved in Section 3 + n . Section 3 concludes with remarks and open problems. The following list summarizes the results (with proofs in the indicated sections):

- (i) level-3 averaged LDP for the joint environment-path Markov chain (Section 4);
- (ii) analysis of the averaged contraction from level-3 to level-1 (Section 5);
- (iii) alternative formula for the level-3 quenched rate function (Section 6);
- (iv) relationship of level-3 averaged and quenched rate functions (Section 7);
- (v) characterizations of the equality of level-1 rate functions (Section 8);
- (vi) minimizers of quenched contractions from level-3 to level-1 (Section 9);
- (vii) spatially constant environments (Section 10).

2 Summary of previous results on large deviations

Recall from the Introduction that the level-1 averaged LDP, i.e., the LDP for the sequence $(P_0(\frac{X_n}{n} \in \cdot))_{n \geq 1}$, is simply the multidimensional Cramér theorem with the rate function $I_{1,a}$ given in (1.9), whereas the statement and the proof of its quenched counterpart is relatively technical. In fact, it is more convenient to first present the level-3 quenched LDP for the environment Markov chain, and we will proceed in this order.

Let S denote the temporal shift operator from the POV of the particle. It acts on $\Omega_{\mathbb{N}} = \Omega \times \mathcal{R}^{\mathbb{N}}$ via $S(\omega, \mathbf{z}) = (T_{1,z_1}\omega, \theta\mathbf{z})$, and on $\Omega_{\mathbb{Z}} = \Omega \times \mathcal{R}^{\mathbb{Z}}$ via $S(\omega, \bar{\mathbf{z}}) = (T_{1,z_1}\omega, \theta\bar{\mathbf{z}})$. On $\Omega_{\mathbb{Z}}$ S is invertible. We can write $S^k(\omega, \bar{\mathbf{z}}) = (T_{k,x_k}\omega, \theta^k\bar{\mathbf{z}})$ for all $k \in \mathbb{Z}$, with this convention: bi-infinite paths x , through the origin and sequences $\bar{\mathbf{z}} \in \mathcal{R}^{\mathbb{Z}}$ are bijectively associated to each other by

$$x_0 = 0, \quad x_k = - \sum_{i=k+1}^0 z_i \quad \text{and} \quad x_\ell = \sum_{i=1}^{\ell} z_i \quad \text{for} \quad k < 0 < \ell. \tag{2.1}$$

Remark 2.1. The empirical process L_n^∞ defined in (1.11) satisfies

$$\int f dL_n^\infty = \frac{1}{n} \sum_{i=0}^{n-1} f \circ S^i$$

for every $f \in b\mathcal{A}_{-\infty, \infty}^{0, \infty}$. In particular,

$$\left| \int (f \circ S) dL_n^\infty - \int f dL_n^\infty \right| \leq \frac{2\|f\|_\infty}{n}.$$

Thus, L_n^∞ is an asymptotically S -invariant element of $\mathcal{M}_1(\Omega_{\mathbb{N}})$ for \mathbb{P} -a.e. ω and every realization of $\mathbf{Z} \in \mathcal{R}^{\mathbb{N}}$.

For any S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$, let

- (i) $\bar{\mu}$ be the unique S -invariant extension of μ to $\Omega_{\mathbb{Z}}$,
- (ii) $\bar{\mu}_-$ the restriction of $\bar{\mu}$ to $\mathcal{A}_{-\infty, \infty}^{-\infty, 0}$, and
- (iii) $\pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty, 0}) = \bar{\mu}(Z_1 = z | \mathcal{A}_{-\infty, \infty}^{-\infty, 0})(\omega, z_{-\infty, 0})$ for every $z \in \mathcal{R}$.

Define $\bar{\mu}_- \times \pi$ and $\bar{\mu}_- \times \pi^{\bar{\mu}}$ on $\mathcal{A}_{-\infty, \infty}^{-\infty, 1}$ by

$$\begin{aligned} (\bar{\mu}_- \times \pi)(d\omega, dz_{-\infty, 1}) &= \bar{\mu}_-(d\omega, dz_{-\infty, 0})\pi_{0,1}(0, z_1 | \omega)c_{\mathcal{R}}(z_1) \quad \text{and} \\ (\bar{\mu}_- \times \pi^{\bar{\mu}})(d\omega, dz_{-\infty, 1}) &= \bar{\mu}_-(d\omega, dz_{-\infty, 0})\pi_{0,1}^{\bar{\mu}}(0, z_1 | \omega, z_{-\infty, 0})c_{\mathcal{R}}(z_1), \end{aligned}$$

respectively. Here, $c_{\mathcal{R}} = \sum_{z \in \mathcal{R}} \delta_z$ is the counting measure on \mathcal{R} . Note that $\bar{\mu}_- \times \pi^{\bar{\mu}}$ is simply the restriction of $\bar{\mu}$ to $\mathcal{A}_{-\infty, \infty}^{-\infty, 1}$. Let $H_q(\mu)$ denote the entropy of $\bar{\mu}_- \times \pi^{\bar{\mu}}$ relative to $\bar{\mu}_- \times \pi$ on $\mathcal{A}_{-\infty, \infty}^{-\infty, 1}$, i.e.,

$$\begin{aligned} H_q(\mu) &= H_{\mathcal{A}_{-\infty, \infty}^{-\infty, 1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \\ &= \int \bar{\mu}_-(d\omega, dz_{-\infty, 0}) \sum_{z \in \mathcal{R}} \pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty, 0}) \log \left(\frac{\pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty, 0})}{\pi_{0,1}(0, z | \omega)} \right). \end{aligned} \tag{2.2}$$

Projecting this entropy to $\mathcal{A}_{-\infty, \infty}^{0, 1}$ and replacing $\pi_{0,1}(0, z | \omega)$ with a constant jump kernel $p(z)$ gives the entropy (1.4) discussed in the Introduction.

The rate function of the level-3 quenched LDP is obtained via the following modification of H_q . For any $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$, denote its Ω -marginal by μ_{Ω} , and set

$$H_{q, \mathbb{P}}^S(\mu) = \begin{cases} H_q(\mu) & \text{if } \mu \text{ is } S\text{-invariant and } \mu_{\Omega} \ll \mathbb{P}, \\ \infty & \text{otherwise.} \end{cases} \tag{2.3}$$

$H_{q, \mathbb{P}}^S$ is convex but not lower semicontinuous, and the double convex conjugate $(H_{q, \mathbb{P}}^S)^{**}$ of $H_{q, \mathbb{P}}^S$ gives its lower semicontinuous regularization (see [42, Theorem 4.17]).

Theorem 2.2 (Level-3 quenched LDP). *Assume*

$$\exists p > d + 1 \text{ such that } \mathbb{E}[|\log \omega_{0,0}(z)|^p] < \infty \text{ for every } z \in \mathcal{R}. \tag{2.4}$$

Then, for \mathbb{P} -a.e. ω , the sequence $(P_0^{\omega}(L_n^{\infty} \in \cdot))_{n \geq 1}$ satisfies an LDP with rate function $I_{3,q} : \mathcal{M}_1(\Omega_{\mathbb{N}}) \rightarrow [0, \infty]$ given by

$$I_{3,q}(\mu) = (H_{q, \mathbb{P}}^S)^{**}(\mu).$$

This result is a special case of the level-3 quenched LDP we established in [43] for a class of models including both directed and undirected RWRE with a rather general but technical condition on the environment measure. We show in Proposition A.2 in Appendix A that this technical condition holds in our current setting under the ellipticity assumption (2.4).

Since the empirical velocity

$$\frac{X_n}{n} = E^{L_n^{\infty}}[Z_1] = \int z_1 L_n^{\infty}(d\omega, dz)$$

is a bounded and continuous function of the empirical process, the level-1 quenched LDP follows immediately from Theorem 2.2 via the contraction principle (see, e.g., [42, Chapter 3]).

Corollary 2.3 (Level-1 quenched LDP). *Assume (2.4). Then, for \mathbb{P} -a.e. ω , the sequence $(P_0^{\omega}(\frac{X_n}{n} \in \cdot))_{n \geq 1}$ satisfies an LDP with rate function $I_{1,q} : \mathbb{R}^d \rightarrow [0, \infty]$ given by*

$$I_{1,q}(\xi) = \inf\{I_{3,q}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), E^{\mu}[Z_1] = \xi\}. \tag{2.5}$$

After the appearance of [43], the level-1 quenched LDP was established in [7] using an alternative method involving the subadditive ergodic theorem, under the stronger assumption of

$$\text{uniform ellipticity: } \exists c > 0 \text{ such that } \mathbb{P}(\omega_{0,0}(z) \geq c) = 1 \text{ for every } z \in \mathcal{R}. \tag{2.6}$$

Originally developed in [48] for undirected RWRE, this second method is less technical and it avoids empirical measures, but does not give a formula for the rate function $I_{1,q}$.

Let $\mathcal{D} := \text{conv}(\mathcal{R})$ denote the convex hull of \mathcal{R} , and $\xi^* := \sum_{z \in \mathcal{R}} \hat{q}(z)z$ stand for the LLN velocity of the walk. The following proposition lists some elementary facts regarding the level-1 averaged and quenched rate functions. We provide its proof in Appendix B for the sake of completeness.

Proposition 2.4. *Assume (2.4). Then, the following hold.*

- (a) $I_{1,a}$ and $I_{1,q}$ are convex and continuous on \mathcal{D} .
- (b) $I_{1,a}(\xi) \leq I_{1,q}(\xi) \leq \max\{\mathbb{E}[|\log \omega_{0,0}(z)|] : z \in \mathcal{R}\} < \infty$ for every $\xi \in \mathcal{D}$.
- (c) $I_{1,a}(\xi) = 0$ iff $I_{1,q}(\xi) = 0$ iff $\xi = \xi^*$.
- (d) $I_{1,a}(z) < I_{1,q}(z)$ for every $z \in \mathcal{R}$ that is an extreme point of \mathcal{D} (unless $\omega_{0,0}(z)$ is deterministic).

Under additional assumptions, the following further results have been obtained regarding the comparison of the level-1 averaged and quenched rate functions in relation with the spatial dimension.

Theorem 2.5. *Assume that*

$$\begin{aligned}
 &(\omega_{i,x})_{(i,x) \in \mathbb{Z} \times \mathbb{Z}^d} \text{ are i.i.d.,} \\
 &\text{the environment is uniformly elliptic (see (2.6)), and} \\
 &\text{the walk is nearest-neighbor, i.e., } \mathcal{R} = U := \{\pm e_1, \dots, \pm e_d\}.
 \end{aligned} \tag{2.7}$$

Then, the following hold at the indicated spatial dimensions.

- (a) ($d = 1$) $I_{1,a}(\xi) < I_{1,q}(\xi)$ for all $\xi \in \mathcal{D} \setminus \{\xi^*\}$, see [52, Theorem 1.5].
- (b) ($d = 2$) $I_{1,a}(\xi) < I_{1,q}(\xi)$ for all $\xi \in \mathcal{D}$ in a punctured neighborhood of ξ^* , see [52, Theorem 1.6].
- (c) ($d \geq 3$) $I_{1,a}(\xi) = I_{1,q}(\xi)$ for all $\xi \in \mathcal{D}$ in a neighborhood of ξ^* , see [49, Theorem 2].

Examining the proofs given in the references reveals that the last two conditions in (2.7) can be replaced with somewhat weaker versions. However, the spatial independence of the environment is crucial to the proofs and cannot be relaxed much.

There are other previous results on large deviations for RWDRE such as the ones in [49] regarding the analysis of the averaged and quenched contractions from level-3 to level-1, but we prefer to mention them in later parts of this paper because they will be either covered by our new results or used in the proofs.

Our temporal independence assumption excludes various concrete models such as RW on particle systems. Level-1,2,3 quenched LDPs for such models (which satisfy uniform ellipticity (2.6)) are covered in [43], but averaged LDPs are open in general. See [1] for level-1 averaged and quenched LDPs for RW on one-dimensional shift-invariant attractive spin-flip systems. Article [28] proves a level-1 LDP for a variant of the RWDRE model where there is mutual influence between the particle and the environment. Finally, for previous results on large deviations for RWRE and closely related models, see [51, Section 2], [37] and [43, Section 1.3], and the references therein.

3 Results

3.1 Level-3 averaged LDP

For any S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$, the specific relative entropy

$$h(\mu | P_0) = \lim_{\ell \rightarrow \infty} \frac{1}{\ell} H_{0,\ell}(\mu | P_0) = \sup_{0 < \ell < \infty} \frac{1}{\ell} H_{0,\ell}(\mu | P_0) \tag{3.1}$$

exists, where

$$H_{k,\ell}(\mu | P_0) := H_{\mathcal{A}_{k,\ell}^{k,\ell}}(\mu | P_0) = \sup_{f \in b\mathcal{A}_{k,\ell}^{k,\ell}} \{E^\mu[f] - \log E_0[e^f]\} \tag{3.2}$$

is the entropy of μ relative to P_0 on $\mathcal{A}_{k,\ell}^{k,\ell}$. The existence of the limit and the identity in (3.1) follow from superadditivity and the independence built into P_0 , and will be justified in Section 4.

Our first result in this paper is the averaged counterpart of Theorem 2.2. Note that it requires only the temporal independence and spatial translation invariance conditions which we assume throughout the paper (see Section 1.2).

Theorem 3.1 (Level-3 averaged LDP). *The sequence $(P_0(L_n^\infty \in \cdot))_{n \geq 1}$ satisfies an LDP with rate function*

$$I_{3,a} : \mathcal{M}_1(\Omega_{\mathbb{N}}) \rightarrow [0, \infty]$$

given by

$$I_{3,a}(\mu) = \begin{cases} h(\mu | P_0) & \text{if } \mu \text{ is } S\text{-invariant,} \\ \infty & \text{otherwise.} \end{cases} \tag{3.3}$$

Remark 3.2. The appearance of S -invariance in (2.3) and (3.3) is natural, as observed in Remark 2.1. Every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ arises in the following way. Consider $\Omega_{\mathbb{Z}}$ as the product space $(\mathcal{P}^{\mathbb{Z}^d} \times \mathcal{R})^{\mathbb{Z}}$ with generic variable $(\omega, \bar{z}) = (\bar{\omega}_i, z_{i+1})_{i \in \mathbb{Z}}$ and temporal shift mapping $(\tau(\omega, \bar{z}))_i = (\bar{\omega}_{i+1}, z_{i+2})$. Let ν be a τ -invariant probability measure on $\Omega_{\mathbb{Z}}$. Recalling (2.1), let $\bar{\mu} \in \mathcal{M}_1(\Omega_{\mathbb{Z}})$ be the distribution of the sequence $(T_{-x_i}^s \bar{\omega}_i, z_{i+1})_{i \in \mathbb{Z}}$ under ν , and finally let μ be the marginal of $\bar{\mu}$ on $\Omega_{\mathbb{N}}$ obtained by dropping the nonpositive steps $z_{-\infty,0}$.

P_0 is not S -invariant (on $\mathcal{A}_{-\infty,\infty}^{0,\infty}$), but there is a unique S -invariant probability measure P_0^∞ on $\Omega_{\mathbb{Z}}$ that agrees with P_0 on $\mathcal{A}_{0,\infty}^{0,\infty}$ (see Lemmas 4.2 and 4.3). The LDP of Theorem 3.1 is valid also for the distributions $(P_0^\infty(L_n^\infty \in \cdot))_{n \geq 1}$ and will in fact be proved first for these.

Similar to Corollary 2.3, the contraction principle gives the following (infinite-dimensional) variational formula for the level-1 averaged rate function:

$$I_{1,a}(\xi) = \inf \{ I_{3,a}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), E^\mu[Z_1] = \xi \}. \tag{3.4}$$

Since (1.9) is a much simpler formula than (3.4), the significance of the latter lies not in providing a numerical value for $I_{1,a}(\xi)$, but in the questions it raises regarding the minimizer(s) of this variational formula, which we pursue next.

3.2 Minimizer of the averaged contraction

Recall from (1.9) that the level-1 averaged rate function $I_{1,a}$ is the convex conjugate of the logarithm of the moment generating function ϕ_a defined in (1.10). We have not assumed that \mathcal{D} has nonempty interior. Consequently $I_{1,a}$ is not necessarily differentiable, and instead of its gradient we have to work with the set-valued subdifferential $\partial I_{1,a}(\xi)$. Facts from convex analysis and proofs of some of the claims below are collected in Appendix C.

Let $\xi \in \text{ri}(\mathcal{D})$, the relative interior of \mathcal{D} . By basic convex analysis, every $\rho \in \partial I_{1,a}(\xi)$ maximizes in (1.9), that is,

$$I_{1,a}(\xi) = \langle \rho, \xi \rangle - \log \phi_a(\rho).$$

$I_{1,a}$ is differentiable at ξ if and only if $\partial I_{1,a}(\xi)$ is a singleton if and only if $\dim(\mathcal{D}) = d$. In general $\partial I_{1,a}(\xi)$ is a nonempty affine subset of \mathbb{R}^d parallel to the orthogonal complement of the affine hull of \mathcal{R} . From this last point it follows that any $\rho \in \partial I_{1,a}(\xi)$ can be used

below to define a measure $\mu^\xi \in \mathcal{M}_1(\Omega_{\mathbb{N}})$: for $-\infty < k \leq 0 < \ell < \infty$ and a test function $f \in b\mathcal{A}_{k,\ell}^{0,\ell}$,

$$\int f(\omega, \mathbf{z}) \mu^\xi(d\omega, d\mathbf{z}) := E_0[e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}(\omega, \mathbf{Z})]. \tag{3.5}$$

Proposition 5.1 in Section 5 provides basic properties of μ^ξ , beginning with its well-definedness.

The second result in this paper identifies μ^ξ as the unique minimizer of the averaged contraction from level-3 to level-1.

Theorem 3.3. *For every $\xi \in \text{ri}(\mathcal{D})$, μ^ξ is the unique minimizer of the variational formula (3.4) of the averaged contraction from level-3 to level-1.*

Measure μ^ξ was introduced in [49, Definition 1] with different notation and under the stronger assumptions in (2.7). Theorem 3.3 follows from an adaptation of [49, Theorem 1] which roughly says that, conditioned on $\{X_n/n \approx \xi\}$, the empirical process L_n^∞ converges to μ^ξ under P_0 . See Proposition 5.3 for the precise statement.

Next we start analyzing the structure of the averaged contraction minimizer $\mu^\xi \in \mathcal{M}_1(\Omega_{\mathbb{N}})$. First of all, μ^ξ is S -invariant (see Proposition 5.1(a)). Using the notation introduced in Section 2, let $\bar{\mu}^\xi$ be the unique S -invariant extension of μ^ξ to $\Omega_{\mathbb{Z}}$, and

$$\pi_{0,1}^{\bar{\mu}^\xi}(0, z | \omega, z_{-\infty,0}) = \bar{\mu}^\xi(Z_1 = z | \mathcal{A}_{-\infty,\infty}^{-\infty,0})(\omega, z_{-\infty,0})$$

for $z \in \mathcal{R}$.

Proposition 3.4. *For every $\xi \in \text{ri}(\mathcal{D})$, $j \geq 0$, and $z \in \mathcal{R}$,*

$$\bar{\mu}^\xi(Z_{j+1} = z | \mathcal{A}_{-\infty,\infty}^{-\infty,j})(\omega, z_{-\infty,j}) = \mu^\xi(Z_1 = z | \mathfrak{S}_{0,\infty})(T_{j,x_j}\omega). \tag{3.6}$$

Hence, the quenched walk under $\bar{\mu}^\xi$ is Markovian, and its transition kernel

$$\pi_{0,1}^\xi(0, z | \omega) := \mu^\xi(Z_1 = z | \mathfrak{S}_{0,\infty})(\omega) = \pi_{0,1}^{\bar{\mu}^\xi}(0, z | \omega, z_{-\infty,0}) \tag{3.7}$$

is $\mathfrak{S}_{0,\infty}$ -measurable.

We denote the Ω -marginal of μ^ξ by μ_Ω^ξ . The proof of Proposition 3.4 in Section 5 shows that, by martingale convergence, the transition kernel in (3.7) is given by

$$\pi_{0,1}^\xi(0, z | \omega) = \lim_{n \rightarrow \infty} \frac{E_0^\omega[e^{\langle \rho, X_n \rangle}, Z_1 = z]}{E_0^\omega[e^{\langle \rho, X_n \rangle}]}, \quad \mu_\Omega^\xi\text{-a.s.}, \tag{3.8}$$

for any $\rho \in \partial I_{1,a}(\xi)$. The following result provides a characterization of the absolute continuity of μ_Ω^ξ in terms of a structural representation of $\pi_{0,1}^\xi$ involving a Doob h -transform.

Theorem 3.5. *For every $\xi \in \text{ri}(\mathcal{D})$, consider the following statements.*

(i) *There exists a function $u \in L^1(\Omega, \mathfrak{S}_{0,\infty}, \mathbb{P})$ such that $\mathbb{P}(u > 0) = 1$ and*

$$\pi_{0,1}^\xi(0, z | \omega) = \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle} u(T_{1,z}\omega)}{\phi_a(\rho) u(\omega)} \tag{3.9}$$

for every $\rho \in \partial I_{1,a}(\xi)$.

(ii) $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$.

Then, (i) \implies (ii). Conversely, if

$$\exists z' \in \mathcal{R} \text{ such that } \mathbb{P}(\omega_{0,0}(z') > 0) = 1, \tag{3.10}$$

then (ii) \implies (i). Furthermore, whenever (i) holds, u is equal (up to a multiplicative constant) to $\left. \frac{d\mu_\Omega^\xi}{d\mathbb{P}} \right|_{\mathfrak{S}_{0,\infty}}$.

The proof that (i) implies (ii) in Theorem 3.5 is adapted from that of [44, Lemma 4.1] which is concerned with disorder regimes of directed random walks in random potentials. The other implication follows from (3.8) under the mild ellipticity condition (3.10) which ensures that μ_Ω^ξ and \mathbb{P} are in fact mutually absolutely continuous on $\mathfrak{S}_{0,\infty}$. For closely related results on directed polymers and ballistic (undirected) RWRE, see [12, Proposition 3.1] and [51, Theorem 3.3], respectively.

Remark 3.6. When we choose ξ to be the LLN velocity $\xi^* = \sum_{z \in \mathcal{R}} \hat{q}(z)z$, we can take $\rho = 0$ because $\nabla \log \phi_a(0) = \xi^*$ which is equivalent to $0 \in \partial I_{1,a}(\xi^*)$ (see (C.2) in Appendix C). Then (3.8) shows that $\pi_{0,1}^{\xi^*}(0, z | \omega) = \pi_{0,1}(0, z | \omega)$, the original kernel, and in (3.9) we can take $u \equiv 1$. Thus $\mu_\Omega^{\xi^*} = \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, which is also evident directly from the definition of μ^{ξ^*} in (3.5).

When $d \geq 3$ and the conditions in (2.7) hold, it was shown by one of the authors [49, Theorem 4] that statements (i) and (ii) in Theorem 3.5 are true not only at $\xi = \xi^*$ but also for ξ sufficiently close to ξ^* , and in this case $u \in L^2(\Omega, \mathfrak{S}_{0,\infty}, \mathbb{P})$.

3.3 Modified variational formulas for the quenched rate functions

Recall from (2.3) that the formula given in Theorem 2.2 for the level-3 quenched rate function $I_{3,q}$ involves absolute continuity w.r.t. \mathbb{P} (on \mathfrak{S}). This formula is valid for a general class of RWRE models. However, in the case of RWDRE, as we have seen in Proposition 3.4 and Theorem 3.5, the relevant σ -algebra is $\mathfrak{S}_{0,\infty}$. Therefore, we next provide appropriately modified formulas for $I_{3,q}$ and $I_{1,q}$ which will be central to some of our subsequent results. Define

$$H_{q,\mathbb{P}}^{S,+}(\mu) = \begin{cases} H_q(\mu) & \text{if } \mu \text{ is } S\text{-invariant and } \mu_\Omega \ll \mathbb{P} \text{ on } \mathfrak{S}_{0,\infty}, \\ \infty & \text{otherwise.} \end{cases} \tag{3.11}$$

Theorem 3.7. Assume (2.4). Then, for every $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$,

$$I_{3,q}(\mu) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu). \tag{3.12}$$

Corollary 3.8. Assume (2.4). Then, for every $\xi \in \text{ri}(\mathcal{D})$,

$$I_{1,q}(\xi) = \inf\{(H_{q,\mathbb{P}}^{S,+})^{**}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), E^\mu[Z_1] = \xi\} \tag{3.13}$$

$$= \inf\{H_q(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), E^\mu[Z_1] = \xi, \mu \text{ is } S\text{-inv.}, \mu_\Omega \ll \mathbb{P} \text{ on } \mathfrak{S}_{0,\infty}\}. \tag{3.14}$$

Example 3.9. The need for Theorem 3.7 is justified by the fact that $H_{q,\mathbb{P}}^S(\mu) = H_{q,\mathbb{P}}^{S,+}(\mu)$ does not hold in general. The following counterexample is adapted from [5]. Assume (2.7) and the following extra condition on the law of the environment:

$$\mathbb{P}(\omega_{0,0}(z) > \omega_{0,0}(z') \text{ for every } z' \in U \setminus \{z\}) = \frac{1}{2d}$$

for every $z \in U$. Consider a new transition kernel π' defined by

$$\pi'_{0,1}(0, z | \omega) = \begin{cases} 1 & \text{if } \omega_{0,0}(z) > \omega_{0,0}(z') \text{ for every } z' \in U \setminus \{z\}, \\ 0 & \text{otherwise.} \end{cases}$$

For \mathbb{P} -a.e. ω , the quenched walk under this new kernel is deterministic, the law of the environment Markov chain $(T_{i,X_i}\omega)_{i \geq 0}$ converges weakly to a π' -invariant probability measure \mathbb{Q} on Ω (see [5, Proposition 1.4]), $\mathbb{Q} = \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, but $\mathbb{Q} \perp \mathbb{P}$ on \mathfrak{S} (see [5,

Proposition 1.5]). Define an S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ by setting $\pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty,0}) = \pi_{0,1}'(0, z | \omega)$ and $\mu_{\Omega} = \mathbb{Q}$. Then, $H_{q,\mathbb{P}}^S(\mu) = \infty$, but (2.4) ensures that

$$H_{q,\mathbb{P}}^{S,+}(\mu) = H_q(\mu) = \mathbb{E} \left[\sum_{z \in U} \pi_{0,1}'(0, z | \omega) \log \left(\frac{\pi_{0,1}'(0, z | \omega)}{\pi_{0,1}(0, z | \omega)} \right) \right] < \infty.$$

3.4 Decomposing the level-3 averaged rate function

The level-3 averaged and quenched LDPs hold with rate functions $I_{3,a}$ and $I_{3,q}$ given in (3.3) and (3.12), respectively. Note that $I_{3,a}(\mu) \leq I_{3,q}(\mu)$ for every $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$. This follows from Jensen's inequality applied to the convex conjugates of the rate functions, and is shown in Corollary 3.11 for the sake of completeness. How are these two rate functions related beyond this basic inequality? The following theorem provides a partial answer. Additional remarks follow in Section 3.8.

Theorem 3.10. *For every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$,*

$$h(\mu | P_0) = h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega} | \mathbb{P}) + H_q(\mu), \tag{3.15}$$

where

$$h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega} | \mathbb{P}) = \lim_{n \rightarrow \infty} \frac{1}{n} H_{\mathfrak{S}_{0,n}}(\mu_{\Omega} | \mathbb{P}).$$

Theorem 3.10 is an application of the chain rule for relative entropy (see [17, Lemma 4.4.7]). It does not require any ellipticity condition. $H_{\mathfrak{S}_{0,n}}(\mu_{\Omega} | \mathbb{P})$ is the entropy of μ_{Ω} relative to \mathbb{P} on $\mathfrak{S}_{0,n}$, and $h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega} | \mathbb{P})$ is the specific relative entropy whose existence is shown in the proof of Theorem 3.10.

Corollary 3.11. *Assume (2.4). Then, for every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$,*

$$\begin{aligned} H_q(\mu) &\leq I_{3,a}(\mu) = h(\mu | P_0) = h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega} | \mathbb{P}) + H_q(\mu) \\ &\leq I_{3,q}(\mu) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu) \leq H_{q,\mathbb{P}}^{S,+}(\mu). \end{aligned}$$

The outermost inequality $H_q(\mu) \leq H_{q,\mathbb{P}}^{S,+}(\mu)$ in Corollary 3.11 holds by definition (3.11), yet it is still important. Indeed, if $H_q(\mu) = H_{q,\mathbb{P}}^{S,+}(\mu)$, then all of the inner inequalities are equalities, too. We will use this simple observation below in the proof of Corollary 3.14.

3.5 Equality of the averaged and quenched rate functions

Proposition 2.4 and Theorem 2.5 summarized what is known about the equality of $I_{1,a}(\xi)$ and $I_{1,q}(\xi)$. The following result complements this picture by providing three characterizations of $I_{1,a}(\xi) = I_{1,q}(\xi)$, each of which involve $\mu^{\xi} \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ (defined in (3.5) for $\xi \in \text{ri}(\mathcal{D})$) or its Ω -marginal μ_{Ω}^{ξ} .

Theorem 3.12. *Assume (2.4). For every $\xi \in \text{ri}(\mathcal{D})$, consider the following statements.*

- (i) $I_{1,a}(\xi) = I_{1,q}(\xi)$.
- (ii) $I_{1,q}(\xi) = H_q(\mu^{\xi})$.
- (iii) $(H_{q,\mathbb{P}}^{S,+})^{**}(\mu^{\xi}) = H_q(\mu^{\xi})$.
- (iv) $h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega}^{\xi} | \mathbb{P}) = 0$.

Then, (i) \iff (ii) \iff (iii) \implies (iv). Moreover, if

$$\exists \delta > 0 \text{ such that } \mathbb{E}[\omega_{0,0}(z)^{-\delta}] < \infty \text{ for every } z \in \mathcal{R}, \tag{3.16}$$

then (iv) \implies (i) and hence all four statements are equivalent.

Remark 3.13. The ellipticity conditions that appear in the statements of our results are related as follows:

$$(3.16) \implies (2.4) \implies (3.10).$$

They are all strictly weaker than uniform ellipticity (2.6).

Regarding the equality of the level-3 averaged and quenched rate functions, the following result provides a sufficient condition. It is also noteworthy that under the stronger condition of uniform ellipticity, the entropy $H_{\mathfrak{S}_{0,n}}(\mu_\Omega | \mathbb{P})$ can grow at most sublinearly for the absolutely continuous marginals of S -invariant measures.

Corollary 3.14. Assume (2.4). Then, for every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$,

$$I_{3,a}(\mu) = I_{3,q}(\mu) = H_q(\mu).$$

Furthermore, if we strengthen (2.4) to uniform ellipticity (2.6), then every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$ satisfies $h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega | \mathbb{P}) = 0$.

3.6 Minimizers of the quenched contractions

Recall from Theorem 3.3 that, for every $\xi \in \text{ri}(\mathcal{D})$, μ^ξ is the unique minimizer of the averaged contraction (3.4) from level-3 to level-1. Finding the minimizers of the quenched contractions (3.13) and (3.14) is more difficult in general. The following result treats the case where the level-1 rate functions are equal.

Theorem 3.15. Assume (2.4). For every $\xi \in \text{ri}(\mathcal{D})$:

(a) if $I_{1,a}(\xi) = I_{1,q}(\xi)$, then

$$I_{1,a}(\xi) = I_{1,q}(\xi) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu^\xi) = H_q(\mu^\xi), \tag{3.17}$$

and μ^ξ is the unique minimizer of the quenched contraction (3.13);

(b) if $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, then (3.17) holds, and μ^ξ is the unique minimizer of the quenched contractions (3.13) and (3.14).

Remark 3.16. Theorem 3.15(b) is not vacuous or trivial (see Remark 3.6). A similar result (regarding level-2 to level-1 contractions for ξ close to ξ^*) was previously obtained for certain ballistic (undirected) RWREs on \mathbb{Z}^d with $d \geq 4$ (see [51, Theorem 3.9]).

The lower semicontinuity of $(H_{q,\mathbb{P}}^{S,+})^{**}$ and the compactness of the set $\{\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}) : E^\mu[Z_1] = \xi\}$ ensure that the quenched contraction (3.13) always has a minimizer. On the other hand, there is currently no general existence result for minimizers of the quenched contraction (3.14). See Section 3.8 for further remarks.

3.7 Spatially constant environments

We illustrate our results in a simplified setting where the spatial variation of the environment is removed. The quenched process \mathbf{Z} is now a process of independent but not identically distributed variables. LDPs for such processes were originally established in [2, 8, 47], motivated in part by their application to the equilibrium statistical mechanics of disordered lattice systems such as the Ising or Curie-Weiss models with random fields or coupling constants. (Some of these large deviation results have been reproduced in Chapter 15 of the textbook [42].) The novelty we provide here is the identification of the averaged and quenched contraction minimizers. We find that many properties such as equality of averaged and quenched rate functions and minimizers fail.

Take a Borel probability measure λ on \mathcal{P} (defined in (1.7)). Let $(\bar{q}_i)_{i \in \mathbb{Z}}$ be sampled from $\mathcal{P}^{\mathbb{Z}}$ according to $\lambda^{\otimes \mathbb{Z}}$. Define $\omega \in \Omega = \mathcal{P}^{\mathbb{Z} \times \mathbb{Z}^d}$ by setting

$$\omega_{i,x} = \bar{q}_i \text{ for every } i \in \mathbb{Z} \text{ and } x \in \mathbb{Z}^d. \tag{3.18}$$

This induces a probability measure \mathbb{P} on (Ω, \mathfrak{S}) . Environments under \mathbb{P} are temporally i.i.d. and spatially constant. Hence, \mathbb{P} is invariant but not ergodic under the spatial translations $(T_y^s)_{y \in \mathbb{Z}^d}$.

For $\rho \in \mathbb{R}^d$ and $\omega \in \Omega$, define

$$W(\rho, \omega) = E_0^\omega[e^{\langle \rho, Z_1 \rangle}].$$

Observe that $\mathbb{E}[W(\rho, \omega)] = \phi_a(\rho)$. For the sake of eliminating trivial cases where the environment is effectively deterministic, we assume that

$$\mathbb{P}(W(\rho, \omega) = \phi_a(\rho)) < 1 \text{ unless } \phi_a(\rho) = e^{\langle \rho, \xi^* \rangle}. \tag{3.19}$$

The condition $\phi_a(\rho) = e^{\langle \rho, \xi^* \rangle}$ is the same as $\rho \in \partial I_{1,a}(\xi^*)$ (Proposition C.3 in Appendix C).

We start our study by giving a simple formula for the level-1 quenched rate function and showing that it is not equal to the averaged one at any atypical velocity.

Proposition 3.17. *Assume (2.4) and (3.18). Then, for every $\xi \in \mathcal{D}$,*

$$I_{1,q}(\xi) = \sup_{\rho \in \mathbb{R}^d} \{ \langle \rho, \xi \rangle - \mathbb{E}[\log W(\rho, \omega)] \} \geq \sup_{\rho \in \mathbb{R}^d} \{ \langle \rho, \xi \rangle - \log \mathbb{E}[W(\rho, \omega)] \} = I_{1,a}(\xi). \tag{3.20}$$

If $\xi \in \text{ri}(\mathcal{D}) \setminus \{ \xi^* \}$ and (3.19) holds, then the inequality in (3.20) is strict.

Remark 3.18. In Proposition 3.17, we assume (2.4) in order to apply Corollary 2.3. In fact, when the environment is spatially constant, a weaker ellipticity condition is sufficient for the level-1 quenched LDP, but we do not pursue such technical improvements here.

Next we present the structure of the unique minimizer μ^ξ (defined in (3.5)) of the averaged contraction (3.4) (see Theorem 3.3). For $\rho \in \mathbb{R}^d$ and $\omega \in \Omega$ let

$$u_1(\rho, \omega) = \frac{W(\rho, \omega)}{\phi_a(\rho)}.$$

Proposition 3.19. *Assume (3.18). Then, for every $\xi \in \text{ri}(\mathcal{D})$, the pairs $(\bar{\omega}_i, Z_{i+1})_{i \geq 0}$ are i.i.d. under μ^ξ . The Ω -marginal μ_{Ω}^ξ and the Markov transition kernel $\pi_{0,1}^\xi$ of μ^ξ are given by*

$$\left. \frac{d\mu_{\Omega}^\xi}{d\mathbb{P}} \right|_{\mathfrak{S}_{0,n}}(\omega) = \prod_{i=0}^{n-1} u_1(\rho, T_{i,0}\omega) \quad \text{and} \quad \pi_{0,1}^\xi(0, z | \omega) = \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle}}{W(\rho, \omega)},$$

for any $\rho \in \partial I_{1,a}(\xi)$. If $\xi \neq \xi^*$ and (3.19) holds, then $\mu_{\Omega}^\xi \not\ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, and $\pi_{0,1}^\xi$ is not obtained from $\pi_{0,1}$ via a Doob h -transform as in (3.9).

The simultaneous lack of absolute continuity and Doob h -transform are consistent with Theorem 3.5. Now that we have simple formulas for μ_{Ω}^ξ and $\pi_{0,1}^\xi$, we can compute $h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega}^\xi | \mathbb{P})$ and $H_q(\mu^\xi)$.

Proposition 3.20. *Assume (3.18). Then, for every $\xi \in \text{ri}(\mathcal{D})$ and $\rho \in \partial I_{1,a}(\xi)$,*

$$h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega}^\xi | \mathbb{P}) = \mathbb{E}[u_1(\rho, \omega) \log u_1(\rho, \omega)] \quad \text{and} \\ H_q(\mu^\xi) = \langle \rho, \xi \rangle - \log \phi_a(\rho) - \mathbb{E}[u_1(\rho, \omega) \log u_1(\rho, \omega)].$$

If $\xi \neq \xi^*$ and (3.19) holds, then $h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega}^\xi | \mathbb{P}) > 0$.

Proposition 3.20 implies that

$$h_{\mathfrak{S}_{0,\infty}}(\mu_{\Omega}^\xi | \mathbb{P}) + H_q(\mu^\xi) = \langle \rho, \xi \rangle - \log \phi_a(\rho) = I_{1,a}(\xi) = h(\mu^\xi | P_0)$$

for every $\xi \in \text{ri}(\mathcal{D})$ and $\rho \in \partial I_{1,a}(\xi)$, which is consistent with Theorem 3.10.

Since the environments are spatially constant, under the quenched conditioning on ω the Ω -marginal of the empirical process L_n^∞ of (1.11) is a deterministic measure that converges to \mathbb{P} . Consequently the quenched rate must blow up at measures with the “wrong” Ω -marginal. This was observed in [8, Theorem III.1] and [47, Theorem 3.4].

Proposition 3.21. *Assume (2.4) and (3.18). For every $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$, if $\mu_\Omega \neq \mathbb{P}$, then $I_{3,q}(\mu) = \infty$. Consequently, if (3.19) holds, then*

$$(H_{q,\mathbb{P}}^{S,+})^{**}(\mu^\xi) = H_{q,\mathbb{P}}^{S,+}(\mu^\xi) = \infty$$

for every $\xi \in \text{ri}(\mathcal{D}) \setminus \{\xi^*\}$.

If (2.4), (3.18) and (3.19) hold, then it follows from Propositions 3.17, 3.20 and 3.21 that all four statements in Theorem 3.12 are false for every $\xi \in \text{ri}(\mathcal{D}) \setminus \{\xi^*\}$, which is consistent with their equivalence.

Proposition 3.21 shows in a striking way how the alteration of the entropy H_q can completely remove the averaged minimizers μ^ξ from the effective domain of the quenched rate function. In particular, μ^ξ cannot be a minimizer of the quenched contractions (3.13) or (3.14). Our final result identifies the minimizer(s) of these quenched contractions.

For $\xi \in \text{ri}(\mathcal{D})$ define $\nu^\xi \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ by setting

$$\nu^\xi(d\omega, dz_{1,n}) = \mathbb{P}(d\omega) \prod_{i=0}^{n-1} \left\{ \pi_{0,1}(0, z_{i+1} \mid T_{i,0}\omega) \frac{e^{\langle \rho, z_{i+1} \rangle}}{W(\rho, T_{i,0}\omega)} \right\} \tag{3.21}$$

for every $n \in \mathbb{N}$, where $\rho \in \partial I_{1,q}(\xi)$.

Proposition 3.22. *Assume (2.4) and (3.18). Then, for every $\xi \in \text{ri}(\mathcal{D})$:*

- (a) ν^ξ is well-defined and S -invariant;
- (b) $E^{\nu^\xi}[Z_1] = \xi$;
- (c) $(H_{q,\mathbb{P}}^{S,+})^{**}(\nu^\xi) = H_q(\nu^\xi) = I_{1,q}(\xi)$; and
- (d) ν^ξ is the unique minimizer of the variational formulas (3.13) and (3.14) of the quenched contractions from level-3 to level-1.

The Ω -marginal of ν^ξ is $\nu_\Omega^\xi = \mathbb{P}$, which is consistent with Proposition 3.21. The transition kernels of μ^ξ (see Proposition 3.19) and ν^ξ are both of the form $\pi_{0,1}(0, z \mid \omega) \frac{e^{\langle \rho, z \rangle}}{W(\rho, \omega)}$, but defined using $\rho \in \partial I_{1,a}(\xi)$ and $\rho \in \partial I_{1,q}(\xi)$, respectively.

3.8 Additional remarks and open problems

3.8.1 Minimizers of the contractions

For every $\xi \in \text{ri}(\mathcal{D})$, Theorem 3.3 identifies μ^ξ as the unique minimizer of the averaged contraction (3.4) from level-3 to level-1. When $I_{1,a}(\xi) = I_{1,q}(\xi)$, Theorem 3.15 says that μ^ξ is also the unique minimizer of the quenched contraction (3.13). Moreover, if $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$ (see Remark 3.6 for examples), then μ^ξ is the unique minimizer of the quenched contraction (3.14), too. In the latter case, Theorem 3.5 gives a representation for the Markov transition kernel $\pi_{0,1}^\xi$ of the quenched walk under μ^ξ via a Doob h -transform.

When $I_{1,a}(\xi) < I_{1,q}(\xi)$, identifying the minimizers (if any) of the quenched contractions (3.13) and (3.14) or saying anything about their structure is an open problem in general. Note that (3.13) always has a minimizer (see Remark 3.16). In contrast, we expect that (3.14) has no minimizers when the environment $(\omega_{i,x})_{(i,x) \in \mathbb{Z} \times \mathbb{Z}^d}$ is i.i.d., but this is yet to be shown. On the other hand, in the case of spatially constant environments, Proposition 3.22 provides the unique minimizer of both of these quenched contractions.

In a recent article [44], we obtained results on the existence and identification of minimizers of variational formula (1.2) for the quenched free energy of random paths

in random potentials and the counterpart of this variational formula for the so-called annealed free energy

$$\log \mathbb{E} \left[E_0^{RW} \left[e^{V(\cdot, Z_1)} \right] \right]$$

in the directed i.i.d. case. These results cover the averaged and quenched logarithmic moment generating functions

$$\log \phi_a(\rho) = \log E_0[e^{\langle \rho, Z_1 \rangle}] \quad \text{and} \quad \Lambda_{1,q}(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega[e^{\langle \rho, X_n \rangle}] \quad (3.22)$$

for RWDRE. (In general, the term averaged refers to first dividing a quenched expectation by the quenched partition function and then taking the \mathbb{P} -expectation of the quotient, whereas annealed refers to first taking the \mathbb{P} -expectation of a quenched expectation and then dividing by the \mathbb{P} -expectation of the quenched partition function. However, there is no difference between them in the RWDRE setting since $E_0^{RW} [e^{V(\omega, Z_1)}] = 1$ for every $\omega \in \Omega$ by (1.8).) The functions in (3.22) are the convex conjugates of $I_{1,a}$ and $I_{1,q}$, respectively, by Varadhan’s lemma. In future work, we hope to combine these previous results with the current ones and thereby deepen our understanding of the large deviation behavior of RWDRE.

3.8.2 Connecting the rate functions

How the averaged and quenched rate functions are related to each other is an important question in the study of processes in random environments. For example, at level-1, obtaining an expression for $I_{1,a}$ in terms of $I_{1,q}$ would provide us with valuable information regarding how the path and the environment conspire towards the realization of atypical velocities. This question is answered with variational formulas in [9] for one-dimensional nearest-neighbor classical RWRE under the i.i.d. environment assumption and in [23] for the exactly solvable corner growth model with random parameters. It is an open problem for example for RWRE in higher dimensions or under more general conditions.

In the context of RWDRE, Theorem 3.10 provides a partial answer to the aforementioned question at level-3 since it connects $I_{3,a}(\mu) = h(\mu | P_0)$ with $I_{3,q}(\mu) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu)$ only indirectly via $H_q(\mu)$. This reduces the original question to understanding the variational expression $(H_{q,\mathbb{P}}^{S,+})^{**}(\mu) - H_q(\mu)$, which is one of our goals for future work. So far, we know that this difference is nonnegative (see Corollary 3.11), and equal to zero at μ^ξ if and only if $I_{1,a}(\xi) = I_{1,q}(\xi)$ (see Theorem 3.12).

3.8.3 Equality of the rate functions

When $I_{1,a}(\xi) = I_{1,q}(\xi)$ at an atypical velocity $\xi \in \text{ri}(\mathcal{D}) \setminus \{\xi^*\}$, the walk is solely responsible (in the exponential scale) for the occurrence of the rare event $\{X_n/n \approx \xi\}$ under the joint measure P_0 . Theorem 3.12 makes this precise by the statement $h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega^\xi | \mathbb{P}) = 0$.

Theorem 2.5 lists the previous results regarding the equality of the level-1 rate functions. The decisive statement for $d = 1$ is believed to be true also for $d = 2$. In contrast, recalling Proposition 2.4 (a,d), both $\mathcal{C} = \{\xi \in \mathcal{D} : I_{1,a}(\xi) = I_{1,q}(\xi)\}$ and $\mathcal{D} \setminus \mathcal{C}$ have nonempty interiors when $d \geq 3$. Hence, there is a phase transition at the boundary of \mathcal{C} , and we would like to analyze the structure of μ^ξ when $\xi \in \partial\mathcal{C}$. The characterizations in Theorem 3.12 can potentially shed light on this problem.

Theorem 3.15(b) provides a sufficient condition for $I_{1,a}(\xi) = I_{1,q}(\xi)$, namely $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$. Whether this condition is also necessary for $I_{1,a}(\xi) = I_{1,q}(\xi)$ is an important open problem which is related to the existence of the critical (i.e., strong but not very strong) disorder regime for directed polymers. See [44, Section 1.3] for details including

the general definitions of the weak, strong and very strong disorder regimes. When specialized to the RWDRE setting and notation, the environment is said to manifest

- (i) weak disorder if $\mu_{\Omega}^{\xi} \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$ (see Remark 3.6 for examples),
- (ii) strong disorder if $\mu_{\Omega}^{\xi} \not\ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, and
- (iii) very strong disorder if $\left. \frac{d\mu_{\Omega}^{\xi}}{d\mathbb{P}} \right|_{\mathfrak{S}_{0,n}}$ decays exponentially to zero as $n \rightarrow \infty$ (equivalently $I_{1,a}(\xi) < I_{1,q}(\xi)$).

Here, ξ is a multidimensional analog of inverse temperature (since $\rho \in \partial I_{1,a}(\xi)$ appears in the exponent inside the expectation on the RHS of (3.5)), with the LLN velocity ξ^* corresponding to infinite temperature. It is tempting to connect this problem of critical disorder with the previous one regarding the structure of μ^{ξ} at the boundary of \mathcal{C} , but we refrain from proposing any conjectures.

4 Level-3 averaged LDP from the point of view of the particle

We start this section with an important point regarding the relative entropies $H_{k,\ell}(\mu | P_0)$ defined in (3.2). We will refer to this point below in the proof of Theorem 3.1.

Remark 4.1. It is not necessarily the case that $H_{k,\ell}(\mu | P_0) = H_{0,\ell-k}(\mu | P_0)$ holds for S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ and $0 < k < \ell$. This is because the distribution of $(\bar{\omega}_i, Z_{i+1})$ under P_0 changes with i . Here is an example: The simplest S -invariant probability measure on $\Omega_{\mathbb{N}}$ is of the product type

$$\mu(d\omega, dz) = \bigotimes_{i \in \mathbb{Z}} \nu(d\bar{\omega}_i) \otimes \bigotimes_{j \in \mathbb{N}} \alpha(z_j).$$

Take $\nu(d\bar{\omega}_i) = \mathbb{P}_s(d\bar{\omega}_i)$ and $\alpha = \delta_z$ for some fixed $z \in \mathcal{R}$. Then,

$$H_{i,i+1}(\mu | P_0) = -\mathbb{E}[\log \sum_x P_0(X_i = x) \pi_{i,i+1}(x, x + z | \omega)].$$

Proof of Theorem 3.1. We will transform the problem into a level-3 LDP for an i.i.d. sequence on the space $\Omega_{\mathbb{Z}} = \Omega \times \mathcal{R}^{\mathbb{Z}}$. First, we define the measure on $\Omega_{\mathbb{Z}}$ that will give the desired i.i.d. sequence.

Recall the definition (2.1) $x_i = -\sum_{j=i+1}^0 z_j$ of the backward path $(x_i)_{i \leq 0}$. For $n \in \mathbb{N}$, let

$$\varphi_{-n}^{\omega}(z_{-n+1,0}) = \prod_{i=-n}^{-1} \pi_{i,i+1}(x_i, x_{i+1} | \omega) = \prod_{i=-n}^{-1} \pi_{0,1}(0, z_{i+1} | T_{i,x_i} \omega).$$

Note that $\varphi_{-n}^{\omega}(z_{-n+1,0})$ is not a probability distribution on vectors $z_{-n+1,0}$ because it does not sum up to one. Set

$$f_n(\omega) = \sum_{z_{-n+1,0} \in \mathcal{R}^n} \varphi_{-n}^{\omega}(z_{-n+1,0}) = \sum_{x \in \mathbb{Z}^d} P_0^{T^{-n,-x}\omega}(X_n = x). \tag{4.1}$$

On the σ -algebra $\mathcal{A}_{-n,\infty}^{-n,\infty}$, we define a measure $\tilde{P}^{(-n)}$ by setting

$$\tilde{P}^{(-n)}(d\bar{\omega}_{-n,\infty}, dz_{-n+1,m}) = \mathbb{P}(d\bar{\omega}_{-n,\infty}) \varphi_{-n}^{\omega}(z_{-n+1,0}) P_0^{\omega}(z_{1,m}) \prod_{i=-n+1}^m c_{\mathcal{R}}(z_i)$$

for every $m \in \mathbb{N}$. Here, $c_{\mathcal{R}}$ denotes the counting measure on \mathcal{R} .

Lemma 4.2. $(\tilde{P}^{(-n)})_{n \in \mathbb{N}}$ are consistent probability measures and hence they induce a probability measure P_0^∞ on $\Omega_{\mathbb{Z}}$.

Proof. For every $m, n \in \mathbb{N}$ and every test function $f \in b\mathcal{A}_{-n, \infty}^{-n, m}$,

$$\begin{aligned} & \int \tilde{P}^{(-n-1)}(d\bar{\omega}_{-n-1, \infty}, dz_{-n, m}) f(\bar{\omega}_{-n, \infty}, z_{-n+1, m}) \\ &= \int \mathbb{P}(d\bar{\omega}_{-n-1, \infty}) \sum_{z_{-n, m}} \varphi_{-n-1}^\omega(z_{-n, 0}) P_0^\omega(z_{1, m}) f(\bar{\omega}_{-n, \infty}, z_{-n+1, m}) \\ &= \int \mathbb{P}(d\bar{\omega}_{-n, \infty}) \sum_{z_{-n+1, m}} \varphi_{-n}^\omega(z_{-n+1, 0}) P_0^\omega(z_{1, m}) f(\bar{\omega}_{-n, \infty}, z_{-n+1, m}) \\ & \quad \times \int \mathbb{P}_s(d\bar{\omega}_{-n-1}) \sum_{z_{-n}} \pi_{0,1}(0, z_{-n} | T_{-n-1, x_{-n}-z_{-n}} \omega) \quad (4.2) \\ &= \int \mathbb{P}(d\bar{\omega}_{-n, \infty}) \sum_{z_{-n+1, m}} \varphi_{-n}^\omega(z_{-n+1, 0}) P_0^\omega(z_{1, m}) f(\bar{\omega}_{-n, \infty}, z_{-n+1, m}) \\ & \quad \times \int \mathbb{P}_s(d\bar{\omega}_{-n-1}) \sum_{z_{-n}} \pi_{0,1}(0, z_{-n} | T_{-n-1, x_{-n}} \omega) \quad (4.3) \\ &= \int \tilde{P}^{(-n)}(d\bar{\omega}_{-n, \infty}, dz_{-n+1, m}) f(\bar{\omega}_{-n, \infty}, z_{-n+1, m}). \end{aligned}$$

Here, (4.2) uses the temporal independence of the environment, whereas (4.3) follows from exchanging the order of the last integral and sum, recalling the spatial translation invariance assumption, and restoring the order of the last integral and sum. Taking $f \equiv 1$, we similarly get

$$\begin{aligned} & \int \tilde{P}^{(-n)}(d\bar{\omega}_{-n, \infty}, dz_{-n+1, m}) = \int \tilde{P}^{(-1)}(d\bar{\omega}_{-1, \infty}, dz_{0, m}) \\ &= \int \mathbb{P}(d\bar{\omega}_{0, \infty}) \sum_{z_{1, m}} P_0^\omega(z_{1, m}) \times \int \mathbb{P}_s(d\bar{\omega}_{-1}) \sum_{z_0} \pi_{0,1}(0, z_0 | T_{-1, 0} \omega) = 1. \end{aligned}$$

We conclude that $(\tilde{P}^{(-n)})_{n \in \mathbb{N}}$ are consistent probability measures. □

Recall that $(T_y^s)_{y \in \mathbb{Z}^d}$ are spatial translations defined by $(T_y^s \bar{\omega}_j)_x = \omega_{j, x+y}$ for $j \in \mathbb{Z}$ and $x, y \in \mathbb{Z}^d$. We use these translations to introduce the so-called slab variables

$$\mathbf{s}_j = (T_{x_j}^s \bar{\omega}_j, z_{j+1}), \quad j \in \mathbb{Z}. \quad (4.4)$$

This choice of terminology comes from viewing RWDRE in \mathbb{Z}^d as a directed RWRE in \mathbb{Z}^{d+1} . Note that \mathbf{s}_j is centered at the point x_j on the path. In this sense, the slab variables are adapted to the POV of the particle. Equivalently, they satisfy $\mathbf{s}_j = \mathbf{s}_0 \circ S^j$ for $j \in \mathbb{Z}$. For any pair of indices $-\infty < k \leq \ell < \infty$, we write $\mathbf{s}_{k, \ell} = (\mathbf{s}_k, \mathbf{s}_{k+1}, \dots, \mathbf{s}_\ell)$.

Lemma 4.3. P_0^∞ is S -invariant and the slab variables $(\mathbf{s}_j)_{j \in \mathbb{Z}}$ are i.i.d. under P_0^∞ .

Proof. This is an immediate consequence of the following induction steps. Let E_0^∞ stand for expectation under P_0^∞ . For $-n < 0 < m$,

$$\begin{aligned} E_0^\infty[f(\mathbf{s}_{-n, m-1})g(\mathbf{s}_m)] &= \sum_{z_{-n+1, m+1}} \mathbb{E} \left[\varphi_{-n}^\omega(z_{-n+1, 0}) P_0^\omega(z_{1, m}) f((T_{x_j}^s \bar{\omega}_j, z_{j+1})_{-n \leq j \leq m-1}) \right. \\ & \quad \left. \times \pi_{0,1}(0, z_{m+1} | T_{m, x_m} \omega) g(T_{x_m}^s \bar{\omega}_m, z_{m+1}) \right] \end{aligned}$$

$$\begin{aligned}
 &= \sum_{z_{-n+1}, m+1} \mathbb{E} \left[\varphi_{-n}^\omega(z_{-n+1}, 0) P_0^\omega(z_{1,m}) f((T_{x_j}^s \bar{\omega}_j, z_{j+1})_{-n \leq j \leq m-1}) \right] \\
 &\quad \times \mathbb{E} \left[\pi_{0,1}(0, z_{m+1} \mid T_{m, x_m} \omega) g(T_{x_m}^s \bar{\omega}_m, z_{m+1}) \right] \\
 &= E_0^\infty [f(\mathbf{s}_{-n, m-1})] E_0 [g(\bar{\omega}_0, Z_1)]
 \end{aligned}$$

by temporal independence and spatial translation invariance, where f and g are test functions on appropriate spaces. Similarly,

$$\begin{aligned}
 E_0^\infty [g(\mathbf{s}_{-n}) f(\mathbf{s}_{-n+1, m})] &= \sum_{z_{-n+1}, m+1} \mathbb{E} \left[\pi_{0,1}(0, z_{-n+1} \mid T_{-n, x_{-n}} \omega) g(T_{x_{-n}}^s \bar{\omega}_{-n}, z_{-n+1}) \right. \\
 &\quad \left. \times \varphi_{-n+1}^\omega(z_{-n+2}, 0) P_0^\omega(z_{1, m+1}) f((T_{x_j}^s \bar{\omega}_j, z_{j+1})_{-n+1 \leq j \leq m}) \right] \\
 &= \sum_{z_{-n+1}, m+1} \mathbb{E} \left[\pi_{0,1}(0, z_{-n+1} \mid T_{-n, x_{-n}} \omega) g(T_{x_{-n}}^s \bar{\omega}_{-n}, z_{-n+1}) \right] \\
 &\quad \times \mathbb{E} \left[\varphi_{-n+1}^\omega(z_{-n+2}, 0) P_0^\omega(z_{1, m+1}) f((T_{x_j}^s \bar{\omega}_j, z_{j+1})_{-n+1 \leq j \leq m}) \right] \\
 &= \sum_z \mathbb{E} [\pi_{0,1}(0, z \mid \omega) g(\bar{\omega}_0, z)] \\
 &\quad \times \sum_{z_{-n+2}, m+1} \mathbb{E} \left[\varphi_{-n+1}^\omega(z_{-n+2}, 0) P_0^\omega(z_{1, m+1}) f((T_{x_j}^s \bar{\omega}_j, z_{j+1})_{-n+1 \leq j \leq m}) \right] \\
 &= E_0 [g(\bar{\omega}_0, Z_1)] E_0^\infty [f(\mathbf{s}_{-n+1, m})]. \quad \square
 \end{aligned}$$

Denote the full sequence of slab variables by $\bar{\mathbf{s}} = (\mathbf{s}_j)_{j \in \mathbb{Z}}$ and let τ be the temporal shift on these sequences, defined by $(\tau \bar{\mathbf{s}})_j = \mathbf{s}_{j+1}$. With this notation, the empirical process induced by the slab variables is

$$L_n^{slab} = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{\tau^i \bar{\mathbf{s}}}.$$

The LDP for L_n^{slab} under P_0^∞ is an instance of the well-known Donsker-Varadhan level-3 LDP for sequences of i.i.d. random variables taking values in Polish spaces, see, e.g., [42, Chapter 6]. We state this result as Proposition 4.4 below, after some preparation for providing a formula for the corresponding rate function.

We can glue together the environment components of the slab variables to form an $\omega' \in \Omega$ with $\bar{\omega}'_j = T_{x_j}^s \bar{\omega}_j$ for $j \in \mathbb{Z}$, and thereby identify the space of slab sequences with $\Omega_{\mathbb{Z}}$. (For the sake of convenience, we will write ω instead of ω' .) This identification already factors in the POV of the particle, and the shift τ acts on (environment, path) pairs simply by

$$(\tau(\bar{\omega}, z_{\cdot+1}))_j = (\bar{\omega}_{j+1}, z_{j+2}).$$

In other words, the sequence $\bar{\mathbf{s}}$ can be thought of as a bijective map on $\Omega_{\mathbb{Z}}$. It induces a τ -invariant distribution $P_0^\infty \circ \bar{\mathbf{s}}^{-1}$ on this space. Since the σ -algebras $\mathcal{A}_{k,\ell}^{k,\ell}$ are now regarded as being generated by the i.i.d. slab variables \mathbf{s}_j , the problem with shifting relative entropy (cf. Remark 4.1) disappears. For τ -invariant probability measures Q on $\Omega_{\mathbb{Z}}$, the specific relative entropy

$$h(Q \mid P_0^\infty \circ \bar{\mathbf{s}}^{-1}) = \lim_{\ell \rightarrow \infty} \frac{H_{0,\ell}(Q \mid P_0^\infty \circ \bar{\mathbf{s}}^{-1})}{\ell} = \sup_{-\infty < k < \ell < \infty} \frac{H_{k,\ell}(Q \mid P_0^\infty \circ \bar{\mathbf{s}}^{-1})}{\ell - k} \quad (4.5)$$

exists by a standard superadditivity argument (see [42, Theorem 6.7]).

Proposition 4.4. *The sequence $(P_0^\infty(L_n^{slab} \in \cdot))_{n \geq 1}$ satisfies an LDP with rate function $I^{slab} : \mathcal{M}_1(\Omega_{\mathbb{Z}}) \rightarrow [0, \infty]$ given by*

$$I^{slab}(Q) = \begin{cases} h(Q | P_0^\infty \circ \bar{s}^{-1}) & \text{if } Q \text{ is } \tau\text{-invariant,} \\ \infty & \text{otherwise.} \end{cases}$$

Next, we transform this LDP into one for the empirical measures

$$L_n^{\mathbb{Z}} = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{T_{i,x_i \omega, \theta^i \bar{\mathbf{z}}}} = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{S^i(\omega, \bar{\mathbf{z}})}.$$

The inverse of the map $(\omega, \bar{\mathbf{z}}) \mapsto \bar{s}$ is $\gamma : \Omega_{\mathbb{Z}} \rightarrow \Omega_{\mathbb{Z}}$ that acts on slabs via

$$(\gamma(\bar{\omega}, z_{\cdot+1}))_j = (T_{-x_j}^s \bar{\omega}_j, z_{j+1}), \quad j \in \mathbb{Z}.$$

Note that

$$S \circ \gamma = \gamma \circ \tau$$

and $L_n^{\mathbb{Z}} = L_n^{slab} \circ \gamma^{-1}$. A probability measure Q on $\Omega_{\mathbb{Z}}$ is S -invariant iff $Q' = Q \circ \gamma$ is τ -invariant. Apply the contraction principle to the LDP in Proposition 4.4 with the map $Q' = Q \circ \gamma \mapsto Q$ on $\mathcal{M}_1(\Omega_{\mathbb{Z}})$. This gives an LDP for $(P_0^\infty(L_n^{\mathbb{Z}} \in \cdot))_{n \geq 1}$ with rate function

$$I^{\mathbb{Z}}(Q) = \begin{cases} h(Q \circ \gamma | P_0^\infty \circ \gamma) & \text{if } Q \text{ is } S\text{-invariant,} \\ \infty & \text{otherwise.} \end{cases}$$

Since γ acts bijectively on any collection of adjacent slabs that includes the zeroth slab and hence preserves $\mathcal{A}_{k,\ell}^{k,\ell}$ -measurability for $k \leq 0 < \ell$,

$$H_{k,\ell}(Q \circ \gamma | P_0^\infty \circ \gamma) = H_{k,\ell}(Q | P_0^\infty).$$

Thus, using (4.5), we can define a specific relative entropy for S -invariant Q by restricting the intervals $[k, \ell)$ to include 0:

$$h(Q | P_0^\infty) = \lim_{\ell \rightarrow \infty} \frac{H_{0,\ell}(Q | P_0^\infty)}{\ell} = \sup_{-\infty < k \leq 0 < \ell < \infty} \frac{H_{k,\ell}(Q | P_0^\infty)}{\ell - k} = h(Q \circ \gamma | P_0^\infty \circ \gamma). \quad (4.6)$$

The statement of the LDP we have established is simplified as follows.

Proposition 4.5. *The sequence $(P_0^\infty(L_n^{\mathbb{Z}} \in \cdot))_{n \geq 1}$ satisfies an LDP with rate function $I^{\mathbb{Z}} : \mathcal{M}_1(\Omega_{\mathbb{Z}}) \rightarrow [0, \infty]$ given by*

$$I^{\mathbb{Z}}(Q) = \begin{cases} h(Q | P_0^\infty) & \text{if } Q \text{ is } S\text{-invariant,} \\ \infty & \text{otherwise.} \end{cases}$$

As the last step, we transform this LDP into the one we want. Denote the natural $\Omega_{\mathbb{Z}} \rightarrow \Omega_{\mathbb{N}}$ projection by $\Phi(\omega, \bar{\mathbf{z}}) = (\omega, \mathbf{z})$. We can think of the empirical process L_n^∞ (introduced in (1.11)) as a function from $\Omega_{\mathbb{Z}}$ into $\mathcal{M}_1(\Omega_{\mathbb{N}})$ by replacing L_n^∞ with $L_n^\infty \circ \Phi$. We drop the projection from the notation since the coordinates of $\theta^i \mathbf{z} = z_{i+1,\infty}$ are defined on $\Omega_{\mathbb{Z}}$ as well as on $\Omega_{\mathbb{N}}$. The contraction principle gives an LDP for $(P_0^\infty(L_n^\infty \in \cdot))_{n \geq 1}$ with rate function $I_{3,a}^\infty : \mathcal{M}_1(\Omega_{\mathbb{N}}) \rightarrow [0, \infty]$ defined by

$$I_{3,a}^\infty(\mu) = \begin{cases} h(\bar{\mu} | P_0^\infty) & \text{if } \mu \text{ is } S\text{-invariant,} \\ \infty & \text{otherwise.} \end{cases}$$

Recall that, for any S -invariant μ on $\Omega_{\mathbb{N}}$, $\bar{\mu}$ denotes the unique S -invariant extension of μ to $\Omega_{\mathbb{Z}}$. By the S -invariance of both $\bar{\mu}$ and P_0^∞ , the entropies can be shifted to nonnegative levels so that

$$H_{k,\ell}(\bar{\mu} | P_0^\infty) = H_{0,\ell-k}(\bar{\mu} | P_0^\infty) = H_{0,\ell-k}(\mu | P_0) \quad \text{for } k \leq 0 < \ell.$$

The first equality above follows from the observation that $f \mapsto f \circ S^{-k}$ and $g \mapsto g \circ S^k$ are bijections between $b\mathcal{A}_{k,\ell}^{k,\ell}$ and $b\mathcal{A}_{0,\ell-k}^{0,\ell-k}$ for $k \leq 0 < \ell$. (However, this is not the case when $0 < k < \ell$, cf. Remark 4.1.) The second equality is valid because $\bar{\mu}$ and μ (resp. P_0^∞ and P_0) agree at nonnegative times. Comparing (3.1) and (4.6), we conclude that $I_{3,a}^\infty$ is equal to the level-3 averaged rate function $I_{3,a}$ defined in (3.3).

It remains to transfer the LDP from $(P_0^\infty(L_n^\infty \in \cdot))_{n \geq 1}$ to $(P_0(L_n^\infty \in \cdot))_{n \geq 1}$ separately for lower and upper bounds. This works easily because (i) weak topology on $\mathcal{M}_1(\Omega_{\mathbb{N}})$ is determined by finite-dimensional distributions, (ii) the dependence of L_n^∞ on environments at negative times vanishes as $n \rightarrow \infty$, and (iii) the measures P_0 and P_0^∞ agree on environments and steps at nonnegative times. We leave the routine details to the reader. This completes the proof of Theorem 3.1. \square

5 Minimizer of the averaged contraction

We start by listing some properties of the measure μ^ξ which was introduced in (3.5) for $\xi \in \text{ri}(\mathcal{D})$.

Proposition 5.1. *For every $\xi \in \text{ri}(\mathcal{D})$:*

- (a) μ^ξ is well-defined and S -invariant;
- (b) the slab variables $(s_\ell)_{\ell \geq 0}$ are i.i.d. under μ^ξ ;
- (c) $E^{\mu^\xi}[Z_1] = \xi$;
- (d) $H_{0,\ell}(\mu^\xi | P_0) = \ell I_{1,a}(\xi)$ for every $\ell \in \mathbb{N}$; and
- (e) $I_{3,a}(\mu^\xi) = h(\mu^\xi | P_0) = I_{1,a}(\xi)$.

Remark 5.2. At the LLN velocity ξ^* , the RHS of (3.5) gets simplified since $0 \in \partial I_{1,a}(\xi^*)$, and we deduce that $\mu^{\xi^*} = P_0$ on $\mathcal{A}_{0,\infty}^{0,\infty}$. (They are not equal on $\mathcal{A}_{-\infty,\infty}^{0,\infty}$, cf. Remark 3.2.) Moreover, the S -invariant extension $\bar{\mu}^{\xi^*} \in \mathcal{M}_1(\Omega_{\mathbb{Z}})$ of μ^{ξ^*} is equal to the measure P_0^∞ which was defined in Lemma 4.2. Therefore, Proposition 5.1(a,b) generalize Lemma 4.3.

Proof of Proposition 5.1. Fix an arbitrary $\xi \in \text{ri}(\mathcal{D})$.

- (a) We prove in Theorem C.2(b) from Appendix C that $\langle \rho, z \rangle - \log \phi_a(\rho) = \langle \rho', z \rangle - \log \phi_a(\rho')$ for every $\rho, \rho' \in \partial I_{1,a}(\xi)$ and $z \in \mathcal{R}$. Therefore, the RHS of (3.5) is well-defined. In order to conclude that μ^ξ is well-defined, it remains to show (for $-\infty < k \leq 0 < \ell < \infty$ and $f \in b\mathcal{A}_{k,\ell}^{0,\ell}$) that the RHS of (3.5) does not change if we replace (i) k by $k - 1$ or (ii) ℓ by $\ell + 1$.

- (i) Since $f \circ S^{-k+1} \in b\mathcal{A}_{1,\ell-k+1}^{-k+1,\ell-k+1}$,

$$\begin{aligned} & E_0[e^{\langle \rho, X_{\ell-k+1} \rangle - (\ell-k+1) \log \phi_a(\rho)} f \circ S^{-k+1}] \\ &= \sum_{z \in \mathcal{R}} \mathbb{E} \left[\pi_{0,1}(0, z | \omega) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} E_0^{T_1, z, \omega} [e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}] \right] \\ &= \sum_{z \in \mathcal{R}} \hat{q}(z) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} E_0[e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}] \\ &= E_0[e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}] \end{aligned}$$

by temporal independence and spatial translation invariance.

(ii) Similarly, since $f \circ S^{-k} \in b\mathcal{A}_{0,\ell-k}^{-k,\ell-k}$,

$$\begin{aligned} & E_0[e^{\langle \rho, X_{\ell-k+1} \rangle - (\ell-k+1) \log \phi_a(\rho)} f \circ S^{-k}] \\ &= \sum_x \mathbb{E} \left[E_0^\omega [e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}, X_{\ell-k} = x] \right. \\ &\quad \left. \times \sum_{z \in \mathcal{R}} \pi_{0,1}(0, z | T_{\ell-k,x}\omega) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} \right] \\ &= \sum_x E_0[e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}, X_{\ell-k} = x] \sum_{z \in \mathcal{R}} \hat{q}(z) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} \\ &= E_0[e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}]. \end{aligned}$$

Finally, regarding the S -invariance of μ^ξ , observe that $f \circ S \in b\mathcal{A}_{k+1,\ell+1}^{1,\ell+1} \subset b\mathcal{A}_{k+1,\ell+1}^{0,\ell+1}$ and

$$\begin{aligned} \int f \circ S d\mu^\xi &= E_0[e^{\langle \rho, X_{(\ell+1)-(k+1)} \rangle - ((\ell+1)-(k+1)) \log \phi_a(\rho)} (f \circ S) \circ S^{-(k+1)}] \\ &= E_0[e^{\langle \rho, X_{\ell-k} \rangle - (\ell-k) \log \phi_a(\rho)} f \circ S^{-k}] = \int f d\mu^\xi. \end{aligned}$$

(b) For every $\ell \in \mathbb{N}$,

$$\begin{aligned} E^{\mu^\xi} [f(\mathbf{s}_{0,\ell-1})g(\mathbf{s}_\ell)] &= E_0[e^{\langle \rho, X_{\ell+1} \rangle - (\ell+1) \log \phi_a(\rho)} f(\mathbf{s}_{0,\ell-1})g(\mathbf{s}_\ell)] \\ &= \sum_x \mathbb{E} \left[E_0^\omega [e^{\langle \rho, X_\ell \rangle - \ell \log \phi_a(\rho)} f(\mathbf{s}_{0,\ell-1}), X_\ell = x] \right. \\ &\quad \left. \times \sum_{z \in \mathcal{R}} \pi_{0,1}(0, z | T_{\ell,x}\omega) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} g(T_x^s \bar{\omega}_\ell, z) \right] \\ &= \sum_x E_0[e^{\langle \rho, X_\ell \rangle - \ell \log \phi_a(\rho)} f(\mathbf{s}_{0,\ell-1}), X_\ell = x] \\ &\quad \times \mathbb{E} \left[\sum_{z \in \mathcal{R}} \pi_{0,1}(0, z | T_{\ell,x}\omega) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} g(T_x^s \bar{\omega}_\ell, z) \right] \\ &= \sum_x E_0[e^{\langle \rho, X_\ell \rangle - \ell \log \phi_a(\rho)} f(\mathbf{s}_{0,\ell-1}), X_\ell = x] \\ &\quad \times \mathbb{E} \left[\sum_{z \in \mathcal{R}} \pi_{0,1}(0, z | \omega) e^{\langle \rho, z \rangle - \log \phi_a(\rho)} g(\bar{\omega}_0, z) \right] \\ &= E^{\mu^\xi} [f(\mathbf{s}_{0,\ell-1})] E^{\mu^\xi} [g(\mathbf{s}_0)], \end{aligned}$$

where f (resp. g) is a test function on ℓ (resp. 1) slab variable(s).

(c) $E^{\mu^\xi} [Z_1] = E_0[e^{\langle \rho, Z_1 \rangle - \log \phi_a(\rho)} Z_1] = \nabla \log \phi_a(\rho) = \xi$ (see (C.2) in Appendix C for the last equality.)

(d) For every $\ell \in \mathbb{N}$,

$$\begin{aligned} H_{0,\ell}(\mu^\xi | P_0) &= \ell H_{0,1}(\mu^\xi | P_0) = \ell E_0[e^{\langle \rho, Z_1 \rangle - \log \phi_a(\rho)} (\langle \rho, Z_1 \rangle - \log \phi_a(\rho))] \\ &= \ell(\langle \rho, \xi \rangle - \log \phi_a(\rho)) = \ell I_{1,a}(\xi), \end{aligned}$$

where the first and third equalities use (b) and (c), respectively. (See (C.2) in Appendix C for the last equality.)

(e) This is immediate from (a) and (d). □

The main ingredient in the proof of Theorem 3.3 is the following result.

Proposition 5.3. For every $\xi \in \text{ri}(\mathcal{D})$, $\varepsilon > 0$, $\ell \in \mathbb{N}$ and $f \in b\mathcal{A}_{-\ell, \ell}^{0, \ell}$,

$$\limsup_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0 \left(\left| \int f dL_n^\infty - \int f d\mu^\xi \right| > \varepsilon \mid \left| \frac{X_n}{n} - \xi \right| < \delta \right) < 0.$$

This result is essentially [49, Theorem 1] which is stated there under the assumptions in (2.7) which are more stringent than our current assumptions. For the sake of completeness and convenience, we provide below a streamlined adaptation of the proof to our setting.

Proof of Proposition 5.3. Let $g = f - \int f d\mu^\xi$. Then,

$$\int f dL_n^\infty - \int f d\mu^\xi = \int g dL_n^\infty =: \langle g, L_n^\infty \rangle.$$

By a standard change-of-measure argument and the level-1 averaged LDP, we see that for any $s > 0$,

$$\begin{aligned} & \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0(\langle g, L_n^\infty \rangle > \varepsilon \mid \left| \frac{X_n}{n} - \xi \right| < \delta) \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0(\langle g, L_n^\infty \rangle > \varepsilon, \left| \frac{X_n}{n} - \xi \right| < \delta) - \liminf_{n \rightarrow \infty} \frac{1}{n} \log P_0(\left| \frac{X_n}{n} - \xi \right| < \delta) \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{\langle \rho, X_n \rangle}, \langle g, L_n^\infty \rangle > \varepsilon, \left| \frac{X_n}{n} - \xi \right| < \delta] - \langle \rho, \xi \rangle + I_{1,a}(\xi) + |\rho|\delta \\ & \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}, \langle g, L_n^\infty \rangle > \varepsilon] + |\rho|\delta \end{aligned} \tag{5.1}$$

$$\leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + ns \langle g, L_n^\infty \rangle}] - s\varepsilon + |\rho|\delta, \tag{5.2}$$

where (5.1) and (5.2) follow from (C.2) in Appendix C and the exponential Chebyshev inequality, respectively. Let $G_j = g \circ S^j$ for $0 \leq j \leq n - 1$ and note that

$$\begin{aligned} E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + ns \langle g, L_n^\infty \rangle}] &= E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + s \sum_{j=0}^{n-1} G_j}] \\ &\leq \prod_{i=0}^{2\ell-1} E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + 2\ell s(G_i + G_{2\ell+i} + G_{4\ell+i} + \dots)}]^{1/2\ell} \end{aligned} \tag{5.3}$$

holds by Hölder’s inequality under $e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)} dP_0$.

For $0 \leq i \leq 2\ell - 1$, let c be the largest integer such that $2c\ell + i \leq n - 1$. For $n \geq 4\ell$,

$$\begin{aligned} & E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + 2\ell s(G_i + G_{2\ell+i} + \dots + G_{2(c-1)\ell+i} + G_{2c\ell+i})}] \\ &= \sum_x \mathbb{E} \left[E_0^\omega [e^{\langle \rho, X_{(2c-1)\ell+i} \rangle - ((2c-1)\ell+i) \log \phi_a(\rho) + 2\ell s(G_i + G_{2\ell+i} + \dots + G_{2(c-1)\ell+i}), X_{(2c-1)\ell+i} = x}] \right. \\ & \quad \times E_0^{T(2c-1)\ell+i, x^\omega} [e^{\langle \rho, X_{n-(2c-1)\ell-i} \rangle - (n-(2c-1)\ell-i) \log \phi_a(\rho) + 2\ell s G_\ell}] \end{aligned} \tag{5.4}$$

$$\begin{aligned} &= \sum_x E_0 [e^{\langle \rho, X_{(2c-1)\ell+i} \rangle - ((2c-1)\ell+i) \log \phi_a(\rho) + 2\ell s(G_i + G_{2\ell+i} + \dots + G_{2(c-1)\ell+i}), X_{(2c-1)\ell+i} = x}] \\ & \quad \times E_0 [e^{\langle \rho, X_{n-(2c-1)\ell-i} \rangle - (n-(2c-1)\ell-i) \log \phi_a(\rho) + 2\ell s G_\ell}] \end{aligned} \tag{5.5}$$

$$\begin{aligned} &= E_0 [e^{\langle \rho, X_{(2c-1)\ell+i} \rangle - ((2c-1)\ell+i) \log \phi_a(\rho) + 2\ell s(G_i + G_{2\ell+i} + \dots + G_{2(c-1)\ell+i})}] \\ & \quad \times E_0 [e^{\langle \rho, X_{n-(2c-1)\ell-i} \rangle - (n-(2c-1)\ell-i) \log \phi_a(\rho) + 2\ell s G_\ell}] \\ &= \dots = E_0 [e^{\langle \rho, X_{\ell+i} \rangle - (\ell+i) \log \phi_a(\rho) + 2\ell s G_i} \left(E_0 [e^{\langle \rho, X_{2\ell} \rangle - 2\ell \log \phi_a(\rho) + 2\ell s G_\ell}] \right)^{c-1}] \\ & \quad \times E_0 [e^{\langle \rho, X_{n-(2c-1)\ell-i} \rangle - (n-(2c-1)\ell-i) \log \phi_a(\rho) + 2\ell s G_\ell}]. \end{aligned} \tag{5.6}$$

Some explanation is in order: Since $g \in b\mathcal{A}_{-\ell, \ell}^{0, \ell}$, each term in the sum in (5.4) is the \mathbb{P} -expectation of the product of two random variables that are measurable w.r.t. $\mathfrak{G}_{-\infty, (2c-1)\ell+i}$ and $\mathfrak{G}_{(2c-1)\ell+i, \infty}$, respectively. Hence, (5.5) follows from the temporal independence and spatial translation invariance of the environment. The repeated application of this argument gives (5.6).

The boundedness of f (and, hence, of g) allows us to control the first and last expectations in (5.6). Therefore,

$$\begin{aligned} & \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + 2\ell s(G_i + G_{2\ell+i} + \dots + G_{2(c-1)\ell+i} + G_{2c\ell+i})}] \\ &= \frac{1}{2\ell} \log E_0[e^{\langle \rho, X_{2\ell} \rangle - 2\ell \log \phi_a(\rho) + 2\ell s G_\ell}]. \end{aligned}$$

Recalling (5.3), we deduce the following inequality:

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho) + ns \langle g, L_n^\infty \rangle}] \leq \frac{1}{2\ell} \log E_0[e^{\langle \rho, X_{2\ell} \rangle - 2\ell \log \phi_a(\rho) + 2\ell s G_\ell}] =: \zeta(s).$$

Note that $\zeta(0) = 0$ and

$$\zeta'(0) = E_0[e^{\langle \rho, X_{2\ell} \rangle - 2\ell \log \phi_a(\rho)} G_\ell] = \int g d\mu^\xi = \int f d\mu^\xi - \int f d\mu^\xi = 0.$$

Therefore, $\zeta(s) = o(s)$ as $s \rightarrow 0$, and it follows from (5.2) that

$$\limsup_{\delta \rightarrow 0} \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0(\langle g, L_n^\infty \rangle > \varepsilon \mid \left| \frac{X_n}{n} - \xi \right| < \delta) < 0.$$

Combining this inequality with the analogous one for $-f$ (and, hence, $-g$), we obtain the desired result. \square

Proof of Theorem 3.3. We checked in Proposition 5.1(e) that μ^ξ is a minimizer of (3.4). It remains to rule out other minimizers. For every $\nu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\nu \neq \mu^\xi$ and $E^\nu[Z_1] = \xi$, there exist $\varepsilon > 0$, $\ell \in \mathbb{N}$ and $f \in b\mathcal{A}_{-\ell, \ell}^{0, \ell}$ such that

$$\nu \in \{ \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}) : \left| \int f d\mu - \int f d\mu^\xi \right| > \varepsilon, |E^\mu[Z_1] - \xi| < \delta \}$$

for every $\delta > 0$, which is an open set. Therefore,

$$\begin{aligned} -I_{3,a}(\nu) &\leq \liminf_{n \rightarrow \infty} \frac{1}{n} \log P_0\left(\left| \int f dL_n^\infty - \int f d\mu^\xi \right| > \varepsilon, \left| \frac{X_n}{n} - \xi \right| < \delta\right) \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0\left(\left| \int f dL_n^\infty - \int f d\mu^\xi \right| > \varepsilon \mid \left| \frac{X_n}{n} - \xi \right| < \delta\right) \\ &\quad + \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0\left(\left| \frac{X_n}{n} - \xi \right| \leq \delta\right) \\ &\leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0\left(\left| \int f dL_n^\infty - \int f d\mu^\xi \right| > \varepsilon \mid \left| \frac{X_n}{n} - \xi \right| < \delta\right) \\ &\quad - \inf\{I_{1,a}(\xi') : |\xi' - \xi| \leq \delta\} \end{aligned}$$

by the level-3 and level-1 averaged LDPs. Taking limit superior as $\delta \rightarrow 0$ gives $-I_{3,a}(\nu) < -I_{1,a}(\xi)$ by Proposition 5.3. Thus ν cannot be a minimizer. \square

Proof of Proposition 3.4. Fix $\xi \in \text{ri}(\mathcal{D})$ and $\rho \in \partial I_{1,a}(\xi)$. Let

$$\alpha_n(\omega, z) = \frac{E_0^\omega[e^{\langle \rho, X_n \rangle}, Z_1 = z]}{E_0^\omega[e^{\langle \rho, X_n \rangle}]}.$$

Let $m, n \geq 1$ and $j \geq 0$. The calculation below shows that

$$\alpha_n(T_{j,x_j}\omega, z) = \bar{\mu}^\xi(Z_{j+1} = z | \mathcal{A}_{-m,j+n}^{-m,j})(\omega, z_{-m+1,j}). \tag{5.7}$$

Take a test function $f \in b\mathcal{A}_{-m,j+n}^{-m,j}$. Then, by S -invariance of $\bar{\mu}^\xi$, the definition (3.5) of μ^ξ , and two uses of the Markov property of the quenched walk,

$$\begin{aligned} & \int f(\omega, Z_{-m+1,j}) \alpha_n(T_{j,x_j}\omega, z) d\bar{\mu}^\xi = \int f(T_{m,X_m}\omega, Z_{1,m+j}) \alpha_n(T_{m+j,X_{m+j}}\omega, z) d\mu^\xi \\ & = E_0[f(T_{m,X_m}\omega, Z_{1,m+j}) \alpha_n(T_{m+j,X_{m+j}}\omega, z) e^{\langle \rho, X_{m+j+n} \rangle - (m+j+n) \log \phi_a(\rho)}] \\ & = E_0[f(T_{m,X_m}\omega, Z_{1,m+j}) e^{\langle \rho, X_{m+j} \rangle - (m+j) \log \phi_a(\rho)} \\ & \quad \times \alpha_n(T_{m+j,X_{m+j}}\omega, z) E_0^{T_{m+j,X_{m+j}}\omega}[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}]] \\ & = E_0[f(T_{m,X_m}\omega, Z_{1,m+j}) e^{\langle \rho, X_{m+j} \rangle - (m+j) \log \phi_a(\rho)} E_0^{T_{m+j,X_{m+j}}\omega}[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}, Z_1 = z]] \\ & = E_0[f(T_{m,X_m}\omega, Z_{1,m+j}) \mathbb{1}\{Z_{m+j+1} = z\} e^{\langle \rho, X_{m+j+n} \rangle - (m+j+n) \log \phi_a(\rho)}] \\ & = \int f(T_{m,X_m}\omega, Z_{1,m+j}) \mathbb{1}\{Z_{m+j+1} = z\} d\mu^\xi = \int f(\omega, Z_{-m+1,j}) \mathbb{1}\{Z_{j+1} = z\} d\bar{\mu}^\xi. \end{aligned}$$

This verifies (5.7). Let $m \rightarrow \infty$ in (5.7). Martingale convergence yields

$$\alpha_n(T_{j,x_j}\omega, z) = \bar{\mu}^\xi(Z_{j+1} = z | \mathcal{A}_{-\infty,j+n}^{-\infty,j})(\omega, z_{-\infty,j}).$$

For the case $j = 0$, by the $\mathfrak{S}_{0,n}$ -measurability of $\alpha_n(\cdot, z)$,

$$\begin{aligned} \alpha_n(\omega, z) & = \bar{\mu}^\xi(Z_1 = z | \mathcal{A}_{-\infty,n}^{-\infty,0})(\omega, z_{-\infty,0}) \\ & = \mu^\xi(Z_1 = z | \mathfrak{S}_{0,n})(\omega). \end{aligned}$$

In the last expression above $\bar{\mu}^\xi$ can be replaced with μ^ξ since the statement does not involve the backward path. Combining the last two displays gives, for $j \geq 0$ and $n \geq 1$,

$$\bar{\mu}^\xi(Z_{j+1} = z | \mathcal{A}_{-\infty,j+n}^{-\infty,j})(\omega, z_{-\infty,j}) = \mu^\xi(Z_1 = z | \mathfrak{S}_{0,n})(T_{j,x_j}\omega). \tag{5.8}$$

As $n \rightarrow \infty$ martingale convergence yields (3.6). The remainder of Proposition 3.4 follows from this. \square

For every $\xi \in \text{ri}(\mathcal{D})$,

$$\left. \frac{d\mu_\Omega^\xi}{d\mathbb{P}} \right|_{\mathfrak{S}_{0,n}}(\omega) = E_0^\omega[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}] =: u_n(\rho, \omega) \tag{5.9}$$

is a positive martingale on $(\Omega, \mathfrak{S}_{0,\infty}, \mathbb{P})$. Throughout the paper, we will sometimes suppress ρ and simply write u_n or $u_n(\omega)$ whenever it does not lead to any confusion.

Proof of Theorem 3.5. (i) \implies (ii): Summing both sides of (3.9) over $z \in \mathcal{R}$, we see that $u \in L^1(\Omega, \mathfrak{S}_{0,\infty}, \mathbb{P})$ satisfies

$$u(\omega) = \sum_{z \in \mathcal{R}} \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle}}{\phi_a(\rho)} u(T_{1,z}\omega) = \sum_{z \in \mathcal{R}} E_0^\omega[e^{\langle \rho, Z_1 \rangle - \log \phi_a(\rho)}, Z_1 = z] u(T_{1,z}\omega).$$

Iterating this identity $n \geq 1$ times, we deduce that

$$u(\omega) = \sum_x u_n(\rho, \omega, x) u(T_{n,x}\omega), \tag{5.10}$$

where

$$u_n(\rho, \omega, x) = E_0^\omega [e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}, X_n = x]. \tag{5.11}$$

Taking the conditional expectation of both sides of (5.10), we get

$$\mathbb{E}[u | \mathfrak{G}_{0,n}](\omega) = \sum_x u_n(\rho, \omega, x) \mathbb{E}[u \circ T_{n,x}] = \mathbb{E}[u] u_n(\rho, \omega)$$

by temporal independence and spatial translation invariance. Since $\mathbb{P}(u > 0) = 1$, we have $\mathbb{E}[u] > 0$. Therefore, as $n \rightarrow \infty$,

$$u_n = \frac{\mathbb{E}[u | \mathfrak{G}_{0,n}]}{\mathbb{E}[u]} \rightarrow \frac{u}{\mathbb{E}[u]}$$

\mathbb{P} -a.s. and in $L^1(\Omega, \mathfrak{G}_{0,\infty}, \mathbb{P})$ (see [21, Theorem 5.5.6]). We conclude that $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{G}_{0,\infty}$, and

$$\left. \frac{d\mu_\Omega^\xi}{d\mathbb{P}} \right|_{\mathfrak{G}_{0,\infty}} = \frac{u}{\mathbb{E}[u]}.$$

(ii) \implies (i): Let $u = \left. \frac{d\mu_\Omega^\xi}{d\mathbb{P}} \right|_{\mathfrak{G}_{0,\infty}}$. Note that $\mathbb{E}[u] = 1$ and hence $\mathbb{P}(u = 0) < 1$. We will first show that (3.10) implies $\mathbb{P}(u = 0) = 0$.

By martingale convergence, $u_n \rightarrow u$ \mathbb{P} -a.s. It follows immediately from the Markov property and the definition in (5.11) that

$$u_{m+n}(\rho, \omega) = \sum_x u_n(\rho, \omega, x) u_m(\rho, T_{n,x}\omega)$$

for every $m, n \in \mathbb{N}$. Sending $m \rightarrow \infty$, we deduce (5.10). In particular,

$$u(\omega) \geq u_n(\rho, \omega, nz) u(T_{n,nz}\omega)$$

for every $z \in \mathcal{R}$. If (3.10) holds, then $\exists z' \in \mathcal{R}$ such that $u_n(\rho, \omega, nz') > 0$ for every $n \in \mathbb{N}$. Therefore,

$$\{\omega : u(\omega) = 0\} \subset \bigcap_{n=1}^\infty \{\omega : u(T_{n,nz'}\omega) = 0\}. \tag{5.12}$$

By our temporal independence and spatial translation invariance assumptions, the Ω -valued process $(T_{n,nz'}\omega)_{n \geq 1}$ is stationary and ergodic under \mathbb{P} . Since $\mathbb{P}(u = 0) < 1$, we apply the ergodic theorem and deduce that, for \mathbb{P} -a.e. ω , there exists an $n \in \mathbb{N}$ such that $u(T_{n,nz'}\omega) > 0$, i.e., the RHS of (5.12) is a \mathbb{P} -null set. Consequently, $\mathbb{P}(u = 0) = 0$.

Finally, we derive (3.9): For \mathbb{P} -a.e. ω and every $z \in \mathcal{R}$,

$$\pi_{0,1}^\xi(0, z | \omega) = \mu^\xi(Z_1 = z | \mathfrak{G}_{0,\infty})(\omega) = \lim_{n \rightarrow \infty} \mu^\xi(Z_1 = z | \mathfrak{G}_{0,n})(\omega) \tag{5.13}$$

$$\begin{aligned} &= \lim_{n \rightarrow \infty} \frac{E_0^\omega [e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}, Z_1 = z]}{E_0^\omega [e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}]} \\ &= \lim_{n \rightarrow \infty} \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle} E_0^{T_{1,z}\omega} [e^{\langle \rho, X_{n-1} \rangle - (n-1) \log \phi_a(\rho)}]}{\phi_a(\rho) E_0^\omega [e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}]} \\ &= \lim_{n \rightarrow \infty} \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle} u_{n-1}(T_{1,z}\omega)}{\phi_a(\rho) u_n(\omega)} \tag{5.14} \\ &= \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle} u(T_{1,z}\omega)}{\phi_a(\rho) u(\omega)}. \end{aligned}$$

Note that the second equality in (5.13) follows from martingale convergence under μ_Ω^ξ which is mutually absolutely continuous with \mathbb{P} since $\mathbb{P}(u > 0) = 1$. \square

6 Modified variational formulas for the quenched rate functions

Since $\bigcup_{j \in \mathbb{N}} \mathcal{A}_{-j, \infty}^{0,j}$ is an algebra that generates $\mathcal{A}_{-\infty, \infty}^{0, \infty}$ on $\Omega_{\mathbb{N}}$, we can fix a sequence $(f_j)_{j \in \mathbb{N}}$ of test functions $f_j \in b\mathcal{A}_{-j, \infty}^{0,j}$ that separate $\mathcal{M}_1(\Omega_{\mathbb{N}})$ and satisfy $\|f_j\|_{\infty} = 1$. For every $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ and $\ell \in \mathbb{N}$, the set

$$G_{\mu, \ell} = \{ \nu \in \mathcal{M}_1(\Omega_{\mathbb{N}}) : |\langle f_j, \mu \rangle - \langle f_j, \nu \rangle| < \ell^{-1} \text{ for } 1 \leq j \leq \ell \}$$

is a weakly open neighborhood of μ . Note that $\bigcap_{\ell \in \mathbb{N}} G_{\mu, \ell} = \bigcap_{\ell \in \mathbb{N}} \overline{G}_{\mu, \ell} = \{ \mu \}$. The following result gives the lower bound in Theorem 3.7.

Theorem 6.1. *Assume that*

$$\mathbb{E}[|\log \omega_{0,0}(z)|] < \infty \text{ for every } z \in \mathcal{R}. \tag{6.1}$$

Then, for every $\ell \in \mathbb{N}$ and every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\mu_{\Omega} \ll \mathbb{P}$ on $\mathfrak{S}_{0, \infty}$,

$$\liminf_{n \rightarrow \infty} \frac{1}{n} \log P_0^{\omega}(L_n^{\infty} \in G_{\mu, \ell}) \geq -H_q(\mu). \tag{6.2}$$

Proof. The proof uses a strategy involving a change-of-measure, Jensen’s inequality, and the ergodic theorem, which is standard for obtaining LDP lower bounds for Markov chains, and has been successfully carried out in the context of (undirected) RWRE (see [46, 50, 40] for the level-1,2,3 quenched LDPs). In fact, keeping future applications in mind, the level-3 quenched LDP lower bound was derived in [40, Section 4] in full detail and without using the assumption that the walk is undirected. In particular, the lower bound of the LDP in Theorem 2.2 is covered by [40, Section 4], which readily implies that (6.2) holds for every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\mu_{\Omega} \ll \mathbb{P}$ (on \mathfrak{S}). Therefore, to prove Theorem 6.1, we need to replace \mathfrak{S} with $\mathfrak{S}_{0, \infty}$. Since the walk is directed in time, this modification requires only two minor changes in the proofs in [40, Section 4]. Below we go over the whole argument for the sake of completeness, point out the two differences, and provide references for further details.

Step 1. For every $\ell \in \mathbb{N}$, denote the marginal of P_0 on $\Omega_{\ell} = \Omega \times \mathcal{R}^{\ell}$ by $P_0^{(\ell)}$. Let $\bar{\eta}_i = (T_{i, X_i} \omega, Z_{i+1, i+\ell})$, $i \geq 0$. Then, under $P_0^{\omega}(\cdot | Z_{1, \ell} = z_{1, \ell})$, $(\bar{\eta}_i)_{i \geq 0}$ is a Markov chain with state space Ω_{ℓ} and transition kernel

$$\pi^{(\ell)}(S_z^+ \eta | \eta) = \pi_{0,1}(0, z | T_{\ell, x_{\ell}} \omega).$$

Here and throughout, $\eta = (\omega, z_{1, \ell}) \in \Omega_{\ell}$ and $S_z^+ : \Omega_{\ell} \rightarrow \Omega_{\ell} : (\omega, z_{1, \ell}) \mapsto (T_{1, z_1} \omega, z_{2, \ell}, z)$ for $z \in \mathcal{R}$.

For every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$, let μ_{Ω} and $\mu_{\Omega_{\ell}}$ denote the marginals of μ on Ω and Ω_{ℓ} , respectively, and define

$$\pi_{\mu}^{(\ell)}(S_z^+ \eta | \eta) = \mu(Z_{\ell+1} = z | \bar{\eta}_0 = \eta)$$

which can be viewed as the transition kernel of a Markov chain with state space Ω_{ℓ} . Since μ is S -invariant, $\mu_{\Omega_{\ell}}$ is an invariant measure for $\pi_{\mu}^{(\ell)}$. Moreover, if $\mu_{\Omega} \ll \mathbb{P}$ on $\mathfrak{S}_{0, \infty}$ and

$$\pi_{\mu}^{(\ell)}(S_z^+ \eta | \eta) > 0 \text{ for } \mu_{\Omega_{\ell}}\text{-a.e. } \eta \text{ and every } z \in \mathcal{R}, \tag{6.3}$$

then $\mu_{\Omega_{\ell}}$ is an ergodic invariant measure for $\pi_{\mu}^{(\ell)}$ and mutually absolutely continuous with $P_0^{(\ell)}$ on $\mathcal{A}_{0, \infty}^{0, \ell}$. This follows from a minor modification of [40, Lemma 4.1] (see Remark 6.2 below).

Step 2. Let P_η (resp. P_η^μ) stand for the law of the Markov chain $(\bar{\eta}_i)_{i \geq 0}$ with initial state $\bar{\eta}_0 = \eta$ and transition kernel $\pi^{(\ell)}$ (resp. $\pi_\mu^{(\ell)}$). Observe that

$$\begin{aligned} P_0^\omega(L_n^\infty \in G_{\mu,\ell}) &= \sum_{z_{1,\ell} \in \mathcal{R}^\ell} P_0^\omega(Z_{1,\ell} = z_{1,\ell}) P_0^\omega(L_n^\infty \in G_{\mu,\ell} \mid Z_{1,\ell} = z_{1,\ell}) \\ &= \sum_{z_{1,\ell} \in \mathcal{R}^\ell} P_0^\omega(Z_{1,\ell} = z_{1,\ell}) P_\eta(L_n^\ell \in G_{\mu,\ell}^{(\ell)}) \\ &\geq \sum_{z_{1,\ell} \in \mathcal{R}^\ell} P_0^\omega(Z_{1,\ell} = z_{1,\ell}) P_\eta(\tilde{L}_n^\ell \in \tilde{G}_{\mu,\ell}^{(\ell)}) \end{aligned} \tag{6.4}$$

for $n \geq 4\ell^2$, where

$$\begin{aligned} L_n^\ell &= \frac{1}{n} \sum_{i=0}^{n-1} \delta_{T_i, X_i, \omega, Z_{i+1, i+\ell}} = \frac{1}{n} \sum_{i=0}^{n-1} \delta_{\bar{\eta}_i} \in \mathcal{M}_1(\Omega_\ell), \quad \tilde{L}_n^\ell = \frac{1}{n-\ell} \sum_{i=\ell}^{n-1} \delta_{\bar{\eta}_i} \in \mathcal{M}_1(\Omega_\ell), \\ G_{\mu,\ell}^{(\ell)} &= \{ \nu \in \mathcal{M}_1(\Omega_\ell) : |\langle f_j, \mu_{\Omega_\ell} \rangle - \langle f_j, \nu \rangle| < \ell^{-1} \text{ for } 1 \leq j \leq \ell \} \quad \text{and} \\ \tilde{G}_{\mu,\ell}^{(\ell)} &= \{ \nu \in \mathcal{M}_1(\Omega_\ell) : |\langle f_j, \mu_{\Omega_\ell} \rangle - \langle f_j, \nu \rangle| < (2\ell)^{-1} \text{ for } 1 \leq j \leq \ell \}. \end{aligned}$$

The inequality in (6.4) is needed in our case because $(\omega, z_{1,\ell}) = \eta \mapsto P_\eta(\tilde{L}_n^\ell \in \tilde{G}_{\mu,\ell}^{(\ell)})$ is $\mathcal{A}_{0,\infty}^{0,\ell}$ -measurable whereas $\eta \mapsto P_\eta(L_n^\ell \in G_{\mu,\ell}^{(\ell)})$ is not.

Step 3. If $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$ and (6.3) holds, then for $P_0^{(\ell)}$ -a.e. η , we change the measure from P_η to P_η^μ , apply Jensen's inequality (with the logarithm function), send $n \rightarrow \infty$, use the ergodicity of μ_{Ω_ℓ} for $\pi_\mu^{(\ell)}$, and thereby deduce that

$$\begin{aligned} \liminf_{n \rightarrow \infty} \frac{1}{n} \log P_\eta(\tilde{L}_n^\ell \in \tilde{G}_{\mu,\ell}^{(\ell)}) &\geq \lim_{n \rightarrow \infty} \frac{1}{n} \log P_\eta^\mu(\tilde{L}_n^\ell \in \tilde{G}_{\mu,\ell}^{(\ell)}) - H(\mu_{\Omega_\ell} \times \pi_\mu^{(\ell)} \mid \mu_{\Omega_\ell} \times \pi^{(\ell)}) \\ &= -H_{\mathcal{A}_{-\infty,\infty}^{-\ell,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} \mid \bar{\mu}_- \times \pi) \geq -H_{\mathcal{A}_{-\infty,\infty}^{-\infty,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} \mid \bar{\mu}_- \times \pi) = -H_q(\mu). \end{aligned}$$

For further details regarding this step, see [40, Lemma 4.2]. The desired bound (6.2) now follows from (6.4).

Step 4. If $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$ but (6.3) fails to hold, we introduce a $\hat{\mu} \in \mathcal{M}_1(\Omega_\mathbb{N})$ of the form $\hat{\mu}(d\omega, d\mathbf{z}) = \mathbb{P}(d\omega) \otimes p^{\otimes \mathbb{N}}(d\mathbf{z})$ for some (deterministic) $p \in \mathcal{P}$ such that $p(z) > 0$ for every $z \in \mathcal{R}$. Note that

$$H_q(\hat{\mu}) = \mathbb{E} \left[\sum_{z \in \mathcal{R}} p(z) \log \left(\frac{p(z)}{\omega_{0,0}(z)} \right) \right] < \infty \tag{6.5}$$

by (6.1). (In fact, this is the only point in the proof where (6.1) is fully used.) We replace μ with $\mu_\epsilon = (1 - \epsilon)\mu + \epsilon\hat{\mu}$ which is an element of the open set $G_{\mu,\ell}$ for sufficiently small $\epsilon > 0$. Since $\hat{\mu}$ is S -invariant, its marginal $\hat{\mu}_{\Omega_\ell}$ on Ω_ℓ is an invariant measure for the transition kernel

$$\pi_{\hat{\mu}}^{(\ell)}(S_z^+ \eta \mid \eta) = \hat{\mu}(Z_{\ell+1} = z \mid \bar{\eta}_0 = \eta) = p(z) > 0.$$

The Ω_ℓ -marginal μ_ϵ^ℓ of μ_ϵ is an invariant measure for a transition kernel $\pi_{\mu_\epsilon}^{(\ell)}$ (suitably defined as a combination of $\pi_\mu^{(\ell)}$ and $\pi_{\hat{\mu}}^{(\ell)}$) which satisfies the analog of (6.3). For further details regarding this step, see [40, Proof of the lower bound in Theorem 3.1, page 224]. Therefore, μ_ϵ^ℓ and $P_0^{(\ell)}$ are mutually absolutely continuous on $\mathcal{A}_{0,\infty}^{0,\ell}$, and for $P_0^{(\ell)}$ -a.e. η ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log P_\eta(\tilde{L}_n^\ell \in \tilde{G}_{\mu,\ell}^{(\ell)}) \geq -H_q(\mu_\epsilon) \geq -(1 - \epsilon)H_q(\mu) - \epsilon H_q(\hat{\mu}).$$

The last inequality follows from the convexity of the relative entropy H_q . Finally, we send ϵ to 0, and recall (6.4) to deduce (6.2) as in Step 3. \square

Remark 6.2. In Step 1 of the proof of Theorem 6.1, we cited [40, Lemma 4.1] which assumes that $\mu_\Omega \ll \mathbb{P}$ (on \mathfrak{S}), which is equivalent to $\mu_{\Omega_\ell} \ll P_0^{(\ell)}$ on $\mathcal{A}_{-\infty, \infty}^{0, \ell}$ since $\mathbb{P}(\omega_{0,0}(z) > 0) = 1$ for every $z \in \mathcal{R}$ by (6.1). In that paper, the mutual absolute continuity of μ_{Ω_ℓ} and $P_0^{(\ell)}$ is established by showing that $f = \frac{d\mu_{\Omega_\ell}}{dP_0^{(\ell)}}$ satisfies

$$\mathbb{1}_{\{f(\omega, z_{1,\ell}) > 0\}} \leq \mathbb{1}_{\{f(T_{\ell+1, x_\ell+z}\omega, \tilde{z}_{1,\ell}) > 0\}}$$

for \mathbb{P} -a.e. ω , every $z_{1,\ell}, \tilde{z}_{1,\ell} \in \mathcal{R}^\ell$ and $z \in \mathcal{R}$, and then using the ergodicity of \mathbb{P} under $(T_{1,z})_{z \in \mathcal{R}}$ to argue that $P_0^{(\ell)}(f > 0) = 1$. Using this, the ergodicity of μ_{Ω_ℓ} for $\pi_\mu^{(\ell)}$ follows from a similar argument.

If $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0, \infty}$, then we replace $f = \frac{d\mu_{\Omega_\ell}}{dP_0^{(\ell)}}$ with $g = \frac{d\mu_{\Omega_\ell}}{dP_0^{(\ell)}} \Big|_{\mathcal{A}_{0, \infty}^{0, \ell}}$ in the proof of [40, Lemma 4.1]. This modification causes no complications since the negative environment levels $\bar{\omega}_{-\infty, -1}$ do not play any role. For example, the function $(\omega, z_{1,\ell}) \mapsto g(T_{\ell+1, x_\ell+z}\omega, \tilde{z}_{1,\ell})$ is measurable w.r.t. $\mathcal{A}_{\ell+1, \infty}^{0, \ell} \subset \mathcal{A}_{0, \infty}^{0, \ell}$ for every $\tilde{z}_{1,\ell} \in \mathcal{R}^\ell$ and $z \in \mathcal{R}$.

We are now ready to verify the modified variational formula for the level-3 quenched rate function.

Proof of Theorem 3.7. It follows immediately from the definitions in (2.3) and (3.11) that $H_{q, \mathbb{P}}^{S,+}(\mu) \leq H_{q, \mathbb{P}}^S(\mu)$ for every $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$. Therefore,

$$(H_{q, \mathbb{P}}^{S,+})^{**}(\mu) \leq (H_{q, \mathbb{P}}^S)^{**}(\mu) = I_{3,q}(\mu). \tag{6.6}$$

On the other hand, Theorem 6.1 and the upper bound in the level-3 quenched LDP (Theorem 2.2) give

$$-H_{q, \mathbb{P}}^{S,+}(\mu) \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \log P_0^\omega(L_n^\infty \in G_{\mu, \ell}) \leq \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0^\omega(L_n^\infty \in G_{\mu, \ell}) \leq - \inf_{\nu \in \bar{G}_{\mu, \ell}} I_{3,q}(\nu)$$

for every $\ell \in \mathbb{N}$. Sending $\ell \rightarrow \infty$, we get

$$I_{3,q}(\mu) \leq H_{q, \mathbb{P}}^{S,+}(\mu)$$

since $I_{3,q}$ is lower semicontinuous and $\bigcap_{\ell \in \mathbb{N}} \bar{G}_{\mu, \ell} = \{\mu\}$, and then deduce that

$$I_{3,q}(\mu) = (I_{3,q})^{**}(\mu) \leq (H_{q, \mathbb{P}}^{S,+})^{**}(\mu). \tag{6.7}$$

Finally, we put (6.6) and (6.7) together to obtain the desired equality (3.12). \square

Proof of Corollary 3.8. For every $\xi \in \mathcal{D}$, the variational formula

$$I_{1,q}(\xi) = \inf\{(H_{q, \mathbb{P}}^{S,+})^{**}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), E^\mu[Z_1] = \xi\}$$

follows immediately from Theorem 3.7 by the contraction principle. Define

$$\tilde{I}_{1,q}(\xi) = \inf\{H_{q, \mathbb{P}}^{S,+}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), E^\mu[Z_1] = \xi\}$$

which is equal to the RHS of (3.14). $\tilde{I}_{1,q}(\xi) < \infty$ because we can choose p to have mean ξ in the measure $\hat{\mu}$ in (6.5). Since $H_{q, \mathbb{P}}^{S,+}$ is convex (which readily follows from the convexity of H_q), $\tilde{I}_{1,q}$ is convex on \mathcal{D} and hence continuous on $\text{ri}(\mathcal{D})$. For every $\xi \in \text{ri}(\mathcal{D})$,

$$\begin{aligned} I_{1,q}(\xi) &= \liminf_{\delta \rightarrow 0} \{I_{1,q}(\xi') : \xi' \in \mathcal{D}, |\xi' - \xi| < \delta\} \\ &= \liminf_{\delta \rightarrow 0} \{(H_{q, \mathbb{P}}^{S,+})^{**}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), |E^\mu[Z_1] - \xi| < \delta\} \\ &= \liminf_{\delta \rightarrow 0} \{H_{q, \mathbb{P}}^{S,+}(\mu) : \mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}), |E^\mu[Z_1] - \xi| < \delta\} \\ &= \liminf_{\delta \rightarrow 0} \{\tilde{I}_{1,q}(\xi') : \xi' \in \mathcal{D}, |\xi' - \xi| < \delta\} = \tilde{I}_{1,q}(\xi) \end{aligned}$$

by the fact that $(H_{q, \mathbb{P}}^{S,+})^{**}$ is the lower semicontinuous regularization of $H_{q, \mathbb{P}}^{S,+}$ (see [42, Theorem 4.17]) and $\{\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}) : |E^\mu[Z_1] - \xi| < \delta\}$ is an open set. \square

7 Decomposing the level-3 averaged rate function

Proof of Theorem 3.10. Observe that, for $1 \leq k \leq n$,

$$P_0(Z_k = z | \mathcal{A}_{0,n}^{0,k-1})(\bar{\omega}_{0,n-1}, z_{1,k-1}) = \pi_{k-1,k}(x_{k-1}, x_{k-1} + z | \omega) = \pi_{0,1}(0, z | T_{k-1,x_{k-1}}\omega).$$

If $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ is S -invariant, then

$$\begin{aligned} H_{0,n}(\mu | P_0) &= H_{\mathcal{A}_{0,n}^{0,n}}(\mu | P_0) \\ &= H_{\mathcal{A}_{0,n}^{0,n-1}}(\mu | P_0) + \int H(\mu(Z_n = \cdot | \mathcal{A}_{0,n}^{0,n-1}) | \pi_{0,1}(0, \cdot | T_{n-1,x_{n-1}}\omega)) \mu(d\bar{\omega}_{0,n-1}, dz_{1,n-1}) \\ &= H_{\mathcal{A}_{0,n}^{0,n-1}}(\mu | P_0) + \int H(\bar{\mu}(Z_1 = \cdot | \mathcal{A}_{-n+1,1}^{-n+1,0}) | \pi_{0,1}(0, \cdot | \omega)) \bar{\mu}(d\bar{\omega}_{-n+1,0}, dz_{-n+2,0}) \\ &= H_{\mathcal{A}_{0,n}^{0,n-1}}(\mu | P_0) + H_{\mathcal{A}_{-n+1,1}^{-n+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \end{aligned}$$

by the chain rule for relative entropy. We can apply the chain rule repeatedly and thereby successively remove all the z -coordinates from the first relative entropy on the RHS. The general step is, for $1 \leq k \leq n - 1$,

$$\begin{aligned} H_{0,n}(\mu | P_0) &= H_{\mathcal{A}_{0,n}^{0,k}}(\mu | P_0) + \sum_{j=k+1}^n H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \\ &= H_{\mathcal{A}_{0,n}^{0,k-1}}(\mu | P_0) + \int H(\mu(Z_k = \cdot | \mathcal{A}_{0,n}^{0,k-1}) | \pi_{0,1}(0, \cdot | T_{k-1,x_{k-1}}\omega)) \mu(d\bar{\omega}_{0,n-1}, dz_{1,k-1}) \\ &\quad + \sum_{j=k+1}^n H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \\ &= H_{\mathcal{A}_{0,n}^{0,k-1}}(\mu | P_0) + \int H(\bar{\mu}(Z_1 = \cdot | \mathcal{A}_{-k+1,n-k+1}^{-k+1,0}) | \pi_{0,1}(0, \cdot | \omega)) \bar{\mu}(d\bar{\omega}_{-k+1,n-k}, dz_{-k+2,0}) \\ &\quad + \sum_{j=k+1}^n H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \\ &= H_{\mathcal{A}_{0,n}^{0,k-1}}(\mu | P_0) + \sum_{j=k}^n H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi). \end{aligned}$$

When all z -coordinates have been removed, we end up with this identity:

$$H_{0,n}(\mu | P_0) = H_{\mathfrak{S}_{0,n}}(\mu_{\Omega} | \mathbb{P}) + \sum_{j=1}^n H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi). \tag{7.1}$$

Lemma 7.1. *If $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ is S -invariant, then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) = H_{\mathcal{A}_{-\infty,\infty}^{-\infty,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) = H_q(\mu). \tag{7.2}$$

Proof. The relative entropy on the RHS of (7.2) is an upper bound on each term in the sum on the LHS. On the other hand, if simultaneously $j \nearrow \infty$ and $n - j \nearrow \infty$, then

$$H_{\mathcal{A}_{-j+1,n-j+1}^{-j+1,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \rightarrow H_{\mathcal{A}_{-\infty,\infty}^{-\infty,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi),$$

which implies the desired result. □

Continuing with the proof of Theorem 3.10, we have seen in Section 4 that the specific relative entropy

$$h(\mu | P_0) = \lim_{n \rightarrow \infty} \frac{1}{n} H_{0,n}(\mu | P_0)$$

exists. In combination with (7.1) and Lemma 7.1, this implies that the limit

$$h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega | \mathbb{P}) = \lim_{n \rightarrow \infty} \frac{1}{n} H_{\mathfrak{S}_{0,n}}(\mu_\Omega | \mathbb{P})$$

exists, and satisfies

$$h(\mu | P_0) = h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega | \mathbb{P}) + H_q(\mu). \quad \square$$

Proof of Corollary 3.11. For every S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$,

$$I_{3,a}(\mu) = h(\mu | P_0) = h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega | \mathbb{P}) + H_q(\mu) \geq H_q(\mu)$$

by Theorems 3.1 and 3.10. Moreover, if (2.4) holds, then from Theorem 3.7 and a basic property of the double convex conjugate (see [42, Proposition 4.10]),

$$I_{3,q}(\mu) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu) \leq H_{q,\mathbb{P}}^{S,+}(\mu).$$

It remains to show that $I_{3,a}(\mu) \leq I_{3,q}(\mu)$. Define

$$\Lambda_{3,a}(f) = \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{n\langle f, L_n^\infty \rangle}] \quad \text{and} \quad \Lambda_{3,q}(f) = \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega[e^{n\langle f, L_n^\infty \rangle}] \quad (7.3)$$

for every continuous $f \in b\mathcal{A}_{\infty,\infty}^{0,\infty}$ and \mathbb{P} -a.e. ω . By Varadhan's lemma (see, e.g., [42, Section 3.2]) these limits exist and are convex conjugates of the rate functions:

$$\Lambda_{3,a}(f) = (I_{3,a})^*(f) \quad \text{and} \quad \Lambda_{3,q}(f) = (I_{3,q})^*(f).$$

Then $I_{3,a}(\mu) \leq I_{3,q}(\mu)$ follows from

$$\begin{aligned} \Lambda_{3,q}(f) &= \mathbb{E} \left[\lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega[e^{n\langle f, L_n^\infty \rangle}] \right] = \lim_{n \rightarrow \infty} \mathbb{E} \left[\frac{1}{n} \log E_0^\omega[e^{n\langle f, L_n^\infty \rangle}] \right] \\ &\leq \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0[e^{n\langle f, L_n^\infty \rangle}] = \Lambda_{3,a}(f). \end{aligned} \quad \square$$

8 Equality of the averaged and quenched rate functions

Throughout this section, we assume (2.4) which ensures that the quenched LDPs hold. Again by Varadhan's lemma the limit

$$\Lambda_{1,q}(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega[e^{\langle \rho, X_n \rangle}] \quad (8.1)$$

exists for every $\rho \in \mathbb{R}^d$ and \mathbb{P} -a.e. ω , and satisfies

$$\Lambda_{1,q}(\rho) = (I_{1,q})^*(\rho) = \sup_{\xi \in \mathcal{D}} \{ \langle \rho, \xi \rangle - I_{1,q}(\xi) \}. \quad (8.2)$$

We have seen in Proposition 2.4(b) (and its proof in Appendix B) that $I_{1,a}(\xi) \leq I_{1,q}(\xi)$ and $\Lambda_{1,q}(\rho) \leq \log \phi_a(\rho)$ for every $\xi \in \mathcal{D}$ and $\rho \in \mathbb{R}^d$.

Lemma 8.1. For every $\xi \in \text{ri}(\mathcal{D})$ and $\rho \in \partial I_{1,a}(\xi)$,

$$I_{1,a}(\xi) < I_{1,q}(\xi) \quad \text{if and only if} \quad \Lambda_{1,q}(\rho) < \log \phi_a(\rho).$$

Proof. For every $\xi \in \text{ri}(\mathcal{D})$ and $\rho \in \partial I_{1,a}(\xi)$, if $\Lambda_{1,q}(\rho) < \log \phi_a(\rho)$, then

$$I_{1,a}(\xi) = \langle \rho, \xi \rangle - \log \phi_a(\rho) < \langle \rho, \xi \rangle - \Lambda_{1,q}(\rho) \leq \sup_{\rho' \in \mathbb{R}^d} \{ \langle \rho', \xi \rangle - \Lambda_{1,q}(\rho') \} = I_{1,q}(\xi).$$

Here, the first equality is shown in (C.2) from Appendix C, and the last equality follows from the convexity of $I_{1,q}$ (see Proposition 2.4(a)).

Conversely, if $I_{1,a}(\xi) < I_{1,q}(\xi)$, then the continuity of $I_{1,q}$ on \mathcal{D} (see Proposition 2.4(a)) implies that

$$\begin{aligned} \Lambda_{1,q}(\rho) &= \sup_{\xi' \in \mathcal{D}} \{ \langle \rho, \xi' \rangle - I_{1,q}(\xi') \} = \langle \rho, \xi'' \rangle - I_{1,q}(\xi'') \\ &\leq \langle \rho, \xi'' \rangle - I_{1,a}(\xi'') \leq \langle \rho, \xi \rangle - I_{1,a}(\xi) = \log \phi_a(\rho) \end{aligned} \tag{8.3}$$

for some $\xi'' \in \mathcal{D}$. If $\xi'' = \xi$, then the first inequality in (8.3) is strict; if $\xi'' \neq \xi$, then the second inequality in (8.3) is strict by (C.2). \square

Recall $u_n(\rho, \omega, x)$ from definition (5.11). When ρ is understood we can drop it from the notation. The next theorem is adapted from [11, Theorem 3.3] which is concerned with upper bounds for the free energy of directed polymers in random environments.

Theorem 8.2. For every $\rho \in \mathbb{R}^d$,

$$\Lambda_{1,q}(\rho) - \log \phi_a(\rho) \leq \inf \left\{ \frac{1}{tm} \log \mathbb{E} \left[\sum_x u_m(\rho, \cdot, x)^t \right] : t \in (0, 1), m \in \mathbb{N} \right\}.$$

Proof. It follows from the definition of $u_n(\omega) = u_n(\rho, \omega)$ in (5.9) and the Markov property of the quenched walk that

$$u_n(\omega) = \sum_{x \in \mathbb{Z}^d} u_n(\omega, x) \quad \text{and} \quad u_{n_1+n_2}(\omega) = \sum_{x_1, x_2} u_{n_1}(\omega, x_1) u_{n_2}(T_{n_1, x_1} \omega, x_2 - x_1).$$

For every $\rho \in \mathbb{R}^d$, $t \in (0, 1)$, and $m, n \geq 1$,

$$\begin{aligned} \mathbb{E} \left[\frac{1}{n} \log u_{nm} \right] &= \mathbb{E} \left[\frac{1}{tn} \log (u_{nm})^t \right] \\ &= \mathbb{E} \left[\frac{1}{tn} \log \left(\sum_{x_1, \dots, x_n} u_m(\cdot, x_1) u_m(T_{m, x_1} \cdot, x_2 - x_1) \cdots u_m(T_{(n-1)m, x_{n-1}} \cdot, x_n - x_{n-1}) \right)^t \right] \\ &\leq \mathbb{E} \left[\frac{1}{tn} \log \left(\sum_{x_1, \dots, x_n} u_m(\cdot, x_1)^t u_m(T_{m, x_1} \cdot, x_2 - x_1)^t \cdots u_m(T_{(n-1)m, x_{n-1}} \cdot, x_n - x_{n-1})^t \right) \right] \\ &\leq \frac{1}{tn} \log \mathbb{E} \left[\sum_{x_1, \dots, x_n} u_m(\cdot, x_1)^t u_m(T_{m, x_1} \cdot, x_2 - x_1)^t \cdots u_m(T_{(n-1)m, x_{n-1}} \cdot, x_n - x_{n-1})^t \right] \\ &= \frac{1}{tn} \log \left(\mathbb{E} \left[\sum_x u_m(\cdot, x)^t \right] \right)^n = \frac{1}{t} \log \mathbb{E} \left[\sum_x u_m(\cdot, x)^t \right] \end{aligned}$$

by the temporal independence and spatial translation invariance assumptions. Sending $n \rightarrow \infty$ and using the bounded convergence theorem, we get

$$m(\Lambda_{1,q}(\rho) - \log \phi_a(\rho)) = \lim_{n \rightarrow \infty} \frac{1}{n} \log u_{nm} = \lim_{n \rightarrow \infty} \mathbb{E} \left[\frac{1}{n} \log u_{nm} \right] \leq \frac{1}{t} \log \mathbb{E} \left[\sum_x u_m(\cdot, x)^t \right]. \quad \square$$

Corollary 8.3. For every $\xi \in \text{ri}(\mathcal{D})$ and $\rho \in \partial I_{1,a}(\xi)$, if there exist $t \in (0, 1)$ and $m \in \mathbb{N}$ such that

$$f(t) := \mathbb{E} \left[\sum_x u_m(\rho, \cdot, x)^t \right] < 1,$$

then $I_{1,a}(\xi) < I_{1,q}(\xi)$.

Proof. This follows immediately from Lemma 8.1 and Theorem 8.2. □

We need two additional lemmas before giving the proof of Theorem 3.12. In fact, the second one is part of Theorem 3.15, but we state and prove it separately here to avoid circular reasoning (we will later use Theorem 3.12 in the proof of Theorem 3.15).

Lemma 8.4. *Let μ and λ be probability measures with finite relative entropy given by*

$$H(\mu | \lambda) = \sup_g \{E^\mu[g] - \log E^\lambda[e^g]\}$$

with supremum over bounded measurable functions g . Then for any event A ,

$$\lambda(A) \geq \exp\{-\mu(A)^{-1}(H(\mu | \lambda) + \log 2)\}. \tag{8.4}$$

Proof. Assume $\mu(A) > 0$ for otherwise the inequality is trivially true. Then also $\lambda(A) > 0$ because finite entropy implies $\mu \ll \lambda$. Take $g = (-\log \lambda(A)) \cdot \mathbb{1}_A$ in the variational formula. □

Lemma 8.5. *Assume (2.4). If $I_{1,a}(\xi) = I_{1,q}(\xi)$, then*

$$I_{1,q}(\xi) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu^\xi),$$

and μ^ξ is the unique minimizer of the quenched contraction (3.13).

Proof. If $I_{1,a}(\xi) = I_{1,q}(\xi)$, then for every $\nu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\nu \neq \mu^\xi$ and $E^\nu[Z_1] = \xi$,

$$I_{1,q}(\xi) = I_{1,a}(\xi) < I_{3,a}(\nu) \leq I_{3,q}(\nu) = (H_{q,\mathbb{P}}^{S,+})^{**}(\nu)$$

by Theorem 3.3 and Corollary 3.11. Hence, ν is not a minimizer of the quenched contraction (3.13). However, the compactness of $\{\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}}) : E^\mu[Z_1] = \xi\}$ and the lower semicontinuity of $(H_{q,\mathbb{P}}^{S,+})^{**}$ guarantee that there is a minimizer. This implies the desired result. □

Proof of Theorem 3.12. $\neg(iv) \implies \neg(i)$: Observe that

$$\begin{aligned} H_{\mathfrak{S}_{0,n}}(\mu_\Omega | \mathbb{P}) &= \mathbb{E}[u_n \log u_n] = \sum_x \mathbb{E}[u_n(\cdot, x) \log u_n(\cdot, x)] - \mathbb{E}\left[u_n \sum_x \frac{u_n(\cdot, x)}{u_n} \log \frac{u_n(\cdot, x)}{u_n}\right] \\ &\leq \sum_x \mathbb{E}[u_n(\cdot, x) \log u_n(\cdot, x)] - \mathbb{E}[u_n] \log \left(\frac{1}{(cn)^d}\right) \\ &= \sum_x \mathbb{E}[u_n(\cdot, x) \log u_n(\cdot, x)] + d \log(cn). \end{aligned} \tag{8.5}$$

Here, we used the following facts: the entropy $-\sum p_i \log p_i$ of a discrete probability distribution with a finite support is dominated by that of the uniform distribution (with the same support); and $(cn)^d$ is a crude upper bound for the number of distinct endpoints of paths of length n started at the origin (with steps in \mathcal{R}).

If $h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega^\xi | \mathbb{P}) > 0$, then $H_{\mathfrak{S}_{0,n}}(\mu_\Omega^\xi | \mathbb{P})$ grows linearly in n . From (8.5) we take the weak consequence that there exists an $m \geq 1$ such that $\sum_x \mathbb{E}[u_m(\cdot, x) \log u_m(\cdot, x)] > 0$. The desired result $I_{1,a}(\xi) < I_{1,q}(\xi)$ follows from Corollary 8.3 which is applicable since $f(1) = 1$ and

$$f'(1) = \sum_x \mathbb{E}[u_m(\cdot, x) \log u_m(\cdot, x)] > 0.$$

$(i) \iff (ii)$: If (i) is true, then so is (iv) by the previous part. Therefore,

$$I_{1,q}(\xi) = I_{1,a}(\xi) = I_{3,a}(\mu^\xi) = h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega^\xi | \mathbb{P}) + H_q(\mu^\xi) = H_q(\mu^\xi)$$

by Proposition 5.1(e) and Corollary 3.11, and hence (ii) is true. Conversely, if (ii) is true, then

$$I_{1,q}(\xi) = H_q(\mu^\xi) \leq I_{3,a}(\mu^\xi) = I_{1,a}(\xi) \leq I_{1,q}(\xi)$$

by Corollary 3.11, Proposition 5.1(e) and Proposition 2.4(b), and hence (i) is true.

(i) \iff (iii) : If (i) is true, then so is (iv) by the first part. Therefore,

$$(H_{q,\mathbb{P}}^{S,+})^{**}(\mu^\xi) = I_{1,q}(\xi) = I_{1,a}(\xi) = I_{3,a}(\mu^\xi) = H_q(\mu^\xi) \leq (H_{q,\mathbb{P}}^{S,+})^{**}(\mu^\xi)$$

by Lemma 8.5, Proposition 5.1(e) and Corollary 3.11, and hence (iii) is true. Conversely, if (iii) is true, then

$$I_{3,q}(\mu^\xi) = I_{3,a}(\mu^\xi) = I_{1,a}(\xi) \leq I_{1,q}(\xi) \leq I_{3,q}(\mu^\xi)$$

by Corollary 3.11, Proposition 5.1(c,e), Proposition 2.4(b) and (2.5), and hence (i) is true.

$\neg(i) \implies \neg(iv)$: Assume (3.16). Theorem D.1 in Appendix D gives the concentration inequality

$$\mathbb{P}(|\log u_n - \mathbb{E}[\log u_n]| \geq n\varepsilon) \leq 2 \exp(-cn)$$

with a constant $c = c(\varepsilon) > 0$. If $I_{1,a}(\xi) < I_{1,q}(\xi)$, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}[\log u_n] = \Lambda_{1,q}(\rho) - \log \phi_a(\rho) < 0$$

by Lemma 8.1, with $\rho \in \partial I_{1,a}(\xi)$. Therefore, there is a $\delta > 0$ such that for large enough n ,

$$\mathbb{P}(u_n \geq \frac{1}{2}) = \mathbb{P}(\log u_n \geq -\log 2) \leq \mathbb{P}(\log u_n \geq -n\delta) \leq 2 \exp(-cn). \quad (8.6)$$

On the other hand,

$$\mu_\Omega^\xi(u_n \geq \frac{1}{2}) = 1 - \mu_\Omega^\xi(u_n < \frac{1}{2}) = 1 - \mathbb{E}[u_n \mathbf{1}_{\{u_n < \frac{1}{2}\}}] \geq \frac{1}{2}.$$

Applying Lemma 8.4 with $\mu = \mu_\Omega^\xi$, $\lambda = \mathbb{P}$ and $A = \{u_n \geq \frac{1}{2}\}$ on $\mathfrak{S}_{0,n}$, we see that (8.4) becomes

$$\mathbb{P}(u_n \geq \frac{1}{2}) \geq \exp\{-2(\mathbb{E}[u_n \log u_n] + \log 2)\}. \quad (8.7)$$

Combining (8.6)–(8.7) shows that $\mathbb{E}[u_n \log u_n]$ grows linearly in n , which contradicts $h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega^\xi | \mathbb{P}) = 0$. \square

Proof of Corollary 3.14. If $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ is S -invariant and $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, then $H_{q,\mathbb{P}}^{S,+}(\mu) = H_q(\mu)$ by (3.11). Therefore, $I_{3,a}(\mu) = I_{3,q}(\mu) = H_q(\mu)$ by Corollary 3.11. In fact, the second equality follows directly from the lower semicontinuity of H_q :

$$H_{q,\mathbb{P}}^{S,+}(\mu) = H_q(\mu) \leq (H_{q,\mathbb{P}}^{S,+})^{**}(\mu) \leq H_{q,\mathbb{P}}^{S,+}(\mu).$$

Under uniform ellipticity (2.6), if μ is S -invariant and $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, then

$$H_q(\mu) \leq \sum_{z \in \mathcal{R}} E^{\mu_\Omega} [|\log \omega_{0,0}(z)|] \leq |\mathcal{R}| |\log c| < \infty,$$

and $h(\mu | P_0) = I_{3,a}(\mu) = H_q(\mu)$ can be canceled from (3.15) to give $h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega | \mathbb{P}) = 0$. \square

9 Minimizers of the quenched contractions

Proof of Theorem 3.15. Fix an arbitrary $\xi \in \text{ri}(\mathcal{D})$. If $I_{1,a}(\xi) = I_{1,q}(\xi)$, then (3.17) follows immediately from Theorem 3.12, and we have shown in Lemma 8.5 that μ^ξ is the unique minimizer of the quenched contraction (3.13). This concludes the proof of part (a).

If $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, then recall from the proof of Theorem 3.5 that $u_n(\omega) = E_0^\omega[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}]$ converges to $u(\omega) = \left. \frac{d\mu_\Omega^\xi}{d\mathbb{P}} \right|_{\mathfrak{S}_{0,\infty}}(\omega)$ for \mathbb{P} -a.e. ω , and $\mathbb{P}(u > 0) = 1$. Therefore,

$$\Lambda_{1,q}(\rho) - \log \phi_a(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} \log u_n(\omega) = \lim_{n \rightarrow \infty} \frac{1}{n} \log u(\omega) = 0.$$

We deduce from Lemma 8.1 that $I_{1,a}(\xi) = I_{1,q}(\xi)$, and part (a) is applicable. Since μ^ξ is S -invariant and $E^{\mu^\xi}[Z_1] = \xi$ (see Proposition 5.1(a,c)), it is a minimizer of the quenched contraction (3.14).

It remains to show that (3.14) has no minimizers other than μ^ξ . To this end, consider any S -invariant $\nu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $\nu \neq \mu^\xi$, $E^\nu[Z_1] = \xi$, and $\nu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$. Observe that

$$I_{1,q}(\xi) < (H_{q,\mathbb{P}}^{S,+})^{**}(\nu) = H_q(\nu)$$

by part (a) and Corollary 3.14. This concludes the proof of part (b). □

10 Spatially constant environments

Proof of Proposition 3.17. The quenched walk X_n is now a sum of independent steps $Z_i \sim \bar{q}_{i-1}$, and so, by the strong LLN, for \mathbb{P} -a.e. ω ,

$$\Lambda_{1,q}(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega[e^{\langle \rho, X_n \rangle}] = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log W(\rho, T_{i,0}\omega) = \mathbb{E}[\log W(\rho, \omega)].$$

Therefore, the first equality in (3.20) follows from (8.2) and the convexity of $I_{1,q}$ (see Proposition 2.4(a)), and the rest from Jensen’s inequality and (1.9).

Assume (3.19). Let $\xi \in \text{ri}(\mathcal{D}) \setminus \{\xi^*\}$ and $\rho \in \partial I_{1,a}(\xi)$. Then $\rho \notin \partial I_{1,a}(\xi^*)$ by (C.2) from Appendix C. Consequently by Proposition C.3 the inequality in (3.19) holds and gives

$$\Lambda_{1,q}(\rho) = \mathbb{E}[\log W(\rho, \omega)] < \log \mathbb{E}[W(\rho, \omega)] = \log \phi_a(\rho) \tag{10.1}$$

by Jensen’s inequality. This implies $I_{1,a}(\xi) < I_{1,q}(\xi)$ by Lemma 8.1. □

Proof of Proposition 3.19. Recall from Proposition 5.1(b) that the slab variables $(s_i)_{i \geq 0}$ (defined in (4.4)) are i.i.d. under μ^ξ for every $\xi \in \text{ri}(\mathcal{D})$. Since the environments are spatially constant, the slab variables are simply $(\bar{\omega}_i, Z_{i+1})_{i \geq 0}$.

By definition (5.9), for every $\rho \in \partial I_{1,a}(\xi)$,

$$\begin{aligned} \left. \frac{d\mu_\Omega^\xi}{d\mathbb{P}} \right|_{\mathfrak{S}_{0,n}}(\omega) &= u_n(\rho, \omega) = E_0^\omega[e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}] \\ &= \prod_{i=0}^{n-1} \frac{W(\rho, T_{i,0}\omega)}{\phi_a(\rho)} = \prod_{i=0}^{n-1} u_1(\rho, T_{i,0}\omega). \end{aligned} \tag{10.2}$$

Therefore, by the limit in (5.14),

$$\begin{aligned} \pi_{0,1}^\xi(0, z | \omega) &= \lim_{n \rightarrow \infty} \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle} u_{n-1}(\rho, T_{1,0}\omega)}{\phi_a(\rho) u_n(\rho, \omega)} \\ &= \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle}}{\phi_a(\rho)} \frac{1}{u_1(\rho, \omega)} = \pi_{0,1}(0, z | \omega) \frac{e^{\langle \rho, z \rangle}}{W(\rho, \omega)}. \end{aligned} \tag{10.3}$$

If $\xi \neq \xi^*$ and (3.19) holds, then for \mathbb{P} -a.e. ω ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log u_n(\rho, \omega) = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log u_1(\rho, T_{i,0}\omega) = \mathbb{E}[\log u_1(\rho, \omega)] < \log \mathbb{E}[u_1(\rho, \omega)] = 0$$

by the strong LLN and Jensen's inequality. In particular, $u_n(\rho, \omega) \rightarrow 0$ as $n \rightarrow \infty$, and $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$.

Under the same assumptions, if $\pi_{0,1}^\xi$ satisfied (3.9) for some $u \in L^1(\Omega, \mathfrak{S}_{0,\infty}, \mathbb{P})$ such that $\mathbb{E}[u] = 1$ and $\mathbb{P}(u > 0) = 1$, then comparison with (10.3) gives

$$u(\omega) = u_1(\rho, \omega)u(T_{1,0}\omega).$$

Iterating this identity, we get

$$u(\omega) = \prod_{i=0}^{n-1} u_1(\rho, T_{i,0}\omega)u(T_{n,0}\omega) = u_n(\rho, \omega)u(T_{n,0}\omega).$$

Therefore, for \mathbb{P} -a.e. ω , $\mathbb{E}[u | \mathfrak{S}_{0,n}](\omega) = u_n(\rho, \omega) \rightarrow u(\omega) > 0$ as $n \rightarrow \infty$, which is a contradiction. \square

Proof of Proposition 3.20. Equality $h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega^\xi | \mathbb{P}) = \mathbb{E}[u_1(\rho, \omega) \log u_1(\rho, \omega)]$ comes from (10.2).

Substitute the second-last formula of (10.3) into (2.2) and use the independence of $(\bar{\omega}_i, Z_{i+1})_{i \geq 0}$ under μ^ξ :

$$\begin{aligned} H_q(\mu^\xi) &= E^{\mu^\xi} \left[\log \left(\frac{\pi_{0,1}^\xi(0, Z_1 | \omega)}{\pi_{0,1}(0, Z_1 | \omega)} \right) \right] = E^{\mu^\xi} \left[\log \left(\frac{e^{\langle \rho, Z_1 \rangle - \log \phi_a(\rho)}}{u_1(\rho, \omega)} \right) \right] \\ &= E^{\mu^\xi} [\langle \rho, Z_1 \rangle] - \log \phi_a(\rho) - E^{\mu^\xi} [\log u_1(\rho, \omega)] \\ &= \langle \rho, \xi \rangle - \log \phi_a(\rho) - \mathbb{E}[u_1(\rho, \omega) \log u_1(\rho, \omega)]. \end{aligned}$$

The last equality used Proposition 5.1(c).

If $\xi \neq \xi^*$ and (3.19) holds, then $\mathbb{P}(u_1(\rho, \omega) = 1) < 1$ for every $\rho \in \partial I_{1,a}(\xi)$ while $\mathbb{E}[u_1(\rho, \omega)] = 1$. Strict convexity of $u \mapsto u \log u$ gives

$$h_{\mathfrak{S}_{0,\infty}}(\mu_\Omega^\xi | \mathbb{P}) = \mathbb{E}[u_1(\rho, \omega) \log u_1(\rho, \omega)] > 0. \quad \square$$

Proof of Proposition 3.21. Under the quenched measure P_0^ω , the Ω -marginal of L_n^∞ is now a deterministic measure $n^{-1} \sum_{i=0}^{n-1} \delta_{T_{i,X_i}\omega} = n^{-1} \sum_{i=0}^{n-1} \delta_{T_{i,0}\omega}$ that converges weakly to \mathbb{P} , for \mathbb{P} -a.e. ω . Hence the rate $I_{3,q}(\mu)$ must be infinite if $\mu_\Omega \neq \mathbb{P}$.

By Proposition 3.19, if $\xi \in \text{ri}(\mathcal{D}) \setminus \{\xi^*\}$ and (3.19) holds, then $\mu_\Omega^\xi \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$. Therefore, $H_{q,\mathbb{P}}^{S,+}(\mu^\xi) = \infty$ by definition (see (3.11)), and $I_{3,q}(\mu^\xi) = (H_{q,\mathbb{P}}^{S,+})^{**}(\mu^\xi) = \infty$ by (3.12) and the paragraph above. \square

Proof of Proposition 3.22. Fix an arbitrary $\xi \in \text{ri}(\mathcal{D})$.

- (a) We prove in Theorem C.4(b) in Appendix C that $\langle \rho, z \rangle - \log W(\rho, \omega) = \langle \rho', z \rangle - \log W(\rho', \omega)$ for every $\rho, \rho' \in \partial I_{1,q}(\xi)$, $z \in \mathcal{R}$ and \mathbb{P} -a.e. ω . Therefore, the RHS of (3.21) is well-defined, and so is ν^ξ by consistency. Taking the POV of the particle, ν^ξ induces a Markov chain on Ω with transition kernel $\bar{\pi}^{\nu^\xi}(\omega' | \omega) = \mathbb{1}_{\{T_{1,0}\omega\}}(\omega')$, for which the Ω -marginal $\nu_\Omega^\xi = \mathbb{P}$ of ν^ξ is an invariant measure. Therefore, ν^ξ is S -invariant.

(b) Recall (C.3) and (C.5) from Appendix C and observe that

$$\begin{aligned} E^{\nu^\xi}[Z_1] &= \mathbb{E} \left[\sum_{z \in \mathcal{R}} \bar{q}_0(z) \frac{e^{\langle \rho, z \rangle} z}{W(\rho, \omega)} \right] = \mathbb{E} \left[\frac{E_0^\omega[e^{\langle \rho, Z_1 \rangle} Z_1]}{E_0^\omega[e^{\langle \rho, Z_1 \rangle}]} \right] \\ &= \mathbb{E}[\nabla \log W(\rho, \omega)] = \nabla \Lambda_{1,q}(\rho) = \xi. \end{aligned}$$

(c) $(H_{q,\mathbb{P}}^{S,+})^{**}(\nu^\xi) = H_q(\nu^\xi)$ by Corollary 3.14 since ν^ξ is S -invariant and $\nu_\Omega^\xi = \mathbb{P}$. Similar to the proof of Proposition 3.20,

$$\begin{aligned} H_q(\nu^\xi) &= E^{\nu^\xi} \left[\log \left(\frac{\pi_{0,1}^{\bar{\nu}^\xi}(0, Z_1 | \omega)}{\pi_{0,1}(0, Z_1 | \omega)} \right) \right] = E^{\nu^\xi} \left[\log \left(\frac{e^{\langle \rho, Z_1 \rangle}}{W(\rho, \omega)} \right) \right] \\ &= E^{\nu^\xi}[\langle \rho, Z_1 \rangle] - E^{\nu_\Omega^\xi}[\log W(\rho, \omega)] = \langle \rho, \xi \rangle - \mathbb{E}[\log W(\rho, \omega)] = I_{1,q}(\xi), \end{aligned}$$

where $\rho \in \partial I_{1,q}(\xi)$, and the fourth equality uses part (b). See (C.3) in Appendix C for the last equality.

(d) We know from part (c) that ν^ξ is a minimizer of (3.13) and (3.14). Take any S -invariant $\mu \in \mathcal{M}_1(\Omega_{\mathbb{N}})$ such that $E^\mu[Z_1] = \xi$.

(i) If μ is a minimizer of (3.13), then $\mu_\Omega = \mathbb{P}$ by Proposition 3.21, and hence $(H_{q,\mathbb{P}}^{S,+})^{**}(\mu) = H_q(\mu)$ by Corollary 3.14.

$$\begin{aligned} I_{1,q}(\xi) &= H_q(\mu) = H_{\mathcal{A}_{-\infty,\infty}^{-\infty,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi) \\ &= \int \bar{\mu}_-(d\omega, dz_{-\infty,0}) \sum_{z \in \mathcal{R}} \pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty,0}) \log \left(\frac{\pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty,0})}{\pi_{0,1}(0, z | \omega)} \right) \\ &= \int \bar{\mu}_-(d\omega, dz_{-\infty,0}) \sum_{z \in \mathcal{R}} \pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty,0}) \log \left(\frac{\pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty,0})}{\pi_{0,1}^{\bar{\nu}^\xi}(0, z | \omega)} \right) \\ &\quad + E^\mu[\langle \rho, Z_1 \rangle] - E^{\mu_\Omega}[\log W(\rho, \omega)] \\ &= H_{\mathcal{A}_{-\infty,\infty}^{-\infty,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi^{\bar{\nu}^\xi}) + \langle \rho, \xi \rangle - \mathbb{E}[\log W(\rho, \omega)] \\ &= H_{\mathcal{A}_{-\infty,\infty}^{-\infty,1}}(\bar{\mu}_- \times \pi^{\bar{\mu}} | \bar{\mu}_- \times \pi^{\bar{\nu}^\xi}) + I_{1,q}(\xi). \end{aligned}$$

Therefore, $\pi_{0,1}^{\bar{\mu}}(0, z | \omega, z_{-\infty,0}) = \pi_{0,1}^{\bar{\nu}^\xi}(0, z | \omega)$ for $\bar{\mu}_-$ -a.e. $(\omega, z_{-\infty,0})$ and $z \in \mathcal{R}$. Since $\mu_\Omega = \nu_\Omega^\xi = \mathbb{P}$, we conclude that $\mu = \nu^\xi$.

(ii) If μ is a minimizer of (3.14), then $\mu_\Omega \ll \mathbb{P}$ on $\mathfrak{S}_{0,\infty}$, therefore $(H_{q,\mathbb{P}}^{S,+})^{**}(\mu) = H_q(\mu)$ by Corollary 3.14, which implies that μ is a minimizer of (3.13), and the previous part is applicable. \square

Appendices

A Sufficient condition for the level-3 quenched LDP

The following definition is adapted from [43, Section 2] to our specific space-time setting and notation. Let $c = \max\{|z|_1 : z \in \mathcal{R}\}$. Here and below, $|\cdot|_1$ denotes the ℓ_1 -norm.

Definition A.1. A function $g : \Omega \rightarrow \mathbb{R}$ is said to be in class \mathcal{L} if $g \in L^1(\Omega, \mathfrak{S}, \mathbb{P})$ and

$$\limsup_{\varepsilon \rightarrow 0} \limsup_{n \rightarrow \infty} \max \left\{ \frac{1}{n} \sum_{0 \leq j \leq \varepsilon n} |g \circ T_{i+j, x+jz}| : 0 \leq i \leq n, |x|_1 \leq ci \right\} = 0 \quad \mathbb{P}\text{-a.s.} \quad (\text{A.1})$$

for every $z \in \mathcal{R}$.

The level-3 quenched LDP we have established in [43, Sections 3& 4] covers RWDRE subject to the following conditions: (i) \mathbb{P} is stationary and ergodic under the family of shifts $(T_{1,z})_{z \in \mathcal{R}}$; and (ii) the function

$$\omega \mapsto \log \pi_{0,1}(0, z | \omega) = \log \omega_{0,0}(z)$$

is in class \mathcal{L} for every $z \in \mathcal{R}$. The first condition is satisfied thanks to the temporal independence of the environment. (In fact, \mathbb{P} is stationary and ergodic under $T_{1,z}$ for each $z \in \mathcal{R}$.) Therefore, to prove Theorem 2.2 (under the ellipticity assumption (2.4)), it suffices to show the following result.

Proposition A.2. *If a Borel measurable function $g_0 : \mathcal{P}^{\mathbb{Z}^d} \rightarrow \mathbb{R}$ satisfies*

$$\int |g_0(\bar{\omega}_0)|^p d\mathbb{P}_s(\bar{\omega}_0) < \infty \tag{A.2}$$

for some $p > d + 1$, then $g : \Omega \rightarrow \mathbb{R}$ defined by $\omega \mapsto g(\omega) := g_0(\bar{\omega}_0)$ is in class \mathcal{L} .

Proof. Since constant functions are in class \mathcal{L} , we can assume without loss of generality that $\mathbb{E}[g] = 0$. It suffices to show a modified version of (A.1), namely,

$$\limsup_{n \rightarrow \infty} \max_{(i,x) \in A_n^\varepsilon} \frac{1}{n} \sum_{0 \leq j \leq 2\varepsilon n} |g \circ T_{i+j,x+jz}| = 0 \quad \text{P-a.s.} \tag{A.3}$$

for every $\varepsilon > 0$, where A_n^ε is a thinned out subset of $\{(i, x) \in \mathbb{Z} \times \mathbb{Z}^d : 0 \leq i \leq n, |x|_1 \leq c_i\}$ of size $|A_n^\varepsilon| \leq C_1 n^d \varepsilon^{-1}$ with some constant $C_1 = C_1(c, d)$.

For each $(i, x) \in A_n^\varepsilon$, the summands in $\sum_{0 \leq j \leq 2\varepsilon n} |g \circ T_{i+j,x+jz}|$ are i.i.d. Therefore, for every $\delta > 0$,

$$\mathbb{P} \left(\sum_{0 \leq j \leq 2\varepsilon n} |g \circ T_{i+j,x+jz}| \geq n\delta \right) \leq C_2(n\delta)^{-p} \varepsilon n$$

by (A.2) and the Fuk-Nagaev inequality (see [33, Corollary 1.8]), where $C_2 = C_2(p)$ is some constant and n is sufficiently large (depending on p, δ, ε). Hence,

$$\mathbb{P} \left(\max_{(i,x) \in A_n^\varepsilon} \sum_{0 \leq j \leq 2\varepsilon n} |g \circ T_{i+j,x+jz}| \geq n\delta \right) \leq C_1 n^d \varepsilon^{-1} C_2 (n\delta)^{-p} \varepsilon n = C_1 C_2 \delta^{-p} n^{d+1-p}$$

by a union bound.

Consider the subsequence $n_m = m^\gamma$ with some $\gamma > (p - d - 1)^{-1}$. Then,

$$\sum_{m=1}^{\infty} C_1 C_2 \delta^{-p} (n_m)^{d+1-p} < \infty$$

and

$$\limsup_{m \rightarrow \infty} \max_{(i,x) \in A_{n_m}^\varepsilon} \frac{1}{n_m} \sum_{0 \leq j \leq 2\varepsilon n_m} |g \circ T_{i+j,x+jz}| \leq \delta \quad \text{P-a.s.}$$

by the Borel-Cantelli lemma. This bound generalizes to the full sequence, too, since $\lim_{m \rightarrow \infty} \frac{n_{m+1}}{n_m} = 1$. Finally, sending $\delta \rightarrow 0$ implies (A.3). \square

B Elementary facts regarding the level-1 rate functions

Proof of Proposition 2.4. (a) $I_{1,a} = (\log \phi_a)^*$ and $I_{3,q} = (H_{q,\mathbb{P}}^S)^{**}$ are convex conjugates and hence convex. $I_{1,q}$ is defined in (2.5) via contraction, and therefore it is convex, too.

Since the rate functions $I_{1,a}$ and $I_{1,q}$ are lower semicontinuous on their domain \mathcal{D} , they are in fact continuous on \mathcal{D} , see [45, Theorem 10.2].

(b) Recall from (8.1) that

$$\Lambda_{1,q}(\rho) := \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega [e^{\langle \rho, X_n \rangle}]$$

for every $\rho \in \mathbb{R}^d$. Varadhan’s lemma gives $\Lambda_{1,q}(\rho) = (I_{1,q})^*(\rho)$. Observe that

$$\Lambda_{1,q}(\rho) = \mathbb{E} \left[\lim_{n \rightarrow \infty} \frac{1}{n} \log E_0^\omega [e^{\langle \rho, X_n \rangle}] \right] \leq \lim_{n \rightarrow \infty} \frac{1}{n} \log E_0 [e^{\langle \rho, X_n \rangle}] = \log \phi_a(\rho)$$

by the bounded convergence theorem and Jensen’s inequality. Thus, for every $\xi \in \mathcal{D}$,

$$\begin{aligned} I_{1,a}(\xi) &= (\log \phi_a)^*(\xi) = \sup_{\rho \in \mathbb{R}^d} \{ \langle \rho, \xi \rangle - \log \phi_a(\rho) \} \\ &\leq \sup_{\rho \in \mathbb{R}^d} \{ \langle \rho, \xi \rangle - \Lambda_{1,q}(\rho) \} = (\Lambda_{1,q})^*(\xi) = I_{1,q}(\xi) \end{aligned}$$

by (1.9) and the convexity of $I_{1,q}$.

For every $z \in \mathcal{R}$, the level-1 quenched LDP upper bound gives

$$\begin{aligned} -I_{1,q}(z) &\geq \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0^\omega (X_n = nz) \geq \limsup_{n \rightarrow \infty} \frac{1}{n} \log \prod_{i=0}^{n-1} \pi_{i,i+1}(iz, (i+1)z | \omega) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log \pi_{0,1}(0, z | T_{i,iz}\omega) = \mathbb{E}[\log \omega_{0,0}(z)]. \end{aligned}$$

The desired bound follows from (2.4) and the convexity of $I_{1,q}$.

(c) $I_{1,a}(\xi^*) = I_{1,q}(\xi^*) = 0$ by the LLN. $\log \phi_a$ is analytic on \mathbb{R}^d and hence $I_{1,a} = (\log \phi_a)^*$ is strictly convex on $\text{ri}(\mathcal{D})$. Therefore, $0 < I_{1,a}(\xi) \leq I_{1,q}(\xi)$ for every $\xi \neq \xi^*$ by part (b), which proves the desired implications.

(d) For every $z \in \mathcal{R}$, the level-1 averaged LDP upper bound gives

$$-I_{1,a}(z) \geq \limsup_{n \rightarrow \infty} \frac{1}{n} \log P_0(X_n = nz) \geq \lim_{n \rightarrow \infty} \frac{1}{n} \log (\hat{q}(z))^n = \log \hat{q}(z) = \log \mathbb{E}[\omega_{0,0}(z)].$$

If z is an extremal point of \mathcal{D} , then for every $\varepsilon > 0$,

$$-I_{1,q}(z) \leq \liminf_{n \rightarrow \infty} \frac{1}{n} \log P_0^\omega \left(\left| \frac{X_n}{n} - z \right| < \varepsilon \right) \leq (1 - c\varepsilon) \mathbb{E}[\log \omega_{0,0}(z)] + O(\varepsilon).$$

Here, the first inequality is an instance of the level-1 quenched LDP lower bound. The second inequality follows from three observations: (i) the event $\{|\frac{X_n}{n} - z| < \varepsilon\}$ consists of $e^{nO(\varepsilon)}$ paths, (ii) each path contains at least $(1 - c\varepsilon)n$ many z -steps for some constant $c = c(\mathcal{R})$, and (iii) the probabilities of these z -steps are i.i.d. by assumption. Sending $\varepsilon \rightarrow 0$, we deduce that

$$-I_{1,q}(z) \leq \mathbb{E}[\log \omega_{0,0}(z)] < \log \mathbb{E}[\omega_{0,0}(z)] \leq -I_{1,a}(z)$$

by Jensen’s inequality (unless $\omega_{0,0}(z)$ is deterministic). □

C Subdifferentials of the level-1 rate functions

The convex hull and the affine hull of the finite set $\mathcal{R} \subset \mathbb{Z}^d$ are defined as

$$\begin{aligned} \mathcal{D} = \text{conv}(\mathcal{R}) &= \left\{ \sum_{z \in \mathcal{R}} \lambda(z)z : \lambda(z) \in [0, 1] \text{ for every } z \in \mathcal{R}, \sum_{z \in \mathcal{R}} \lambda(z) = 1 \right\} \text{ and} \\ M = \text{aff}(\mathcal{R}) &= \left\{ \sum_{z \in \mathcal{R}} \lambda(z)z : \lambda(z) \in \mathbb{R} \text{ for every } z \in \mathcal{R}, \sum_{z \in \mathcal{R}} \lambda(z) = 1 \right\}, \end{aligned}$$

respectively. The relative interior $\text{ri}(\mathcal{D})$ is the interior of \mathcal{D} in the relative topology of M .

Recall from Appendix B that the functions $I_{1,a}$ and $\log \phi_a$ (resp. $I_{1,q}$ and $\Lambda_{1,q}$) are convex conjugates of each other. The subdifferential $\partial I_{1,a}(\xi)$ of $I_{1,a}$ at $\xi \in \mathcal{D}$ is defined as

$$\partial I_{1,a}(\xi) = \{\rho \in \mathbb{R}^d : I_{1,a}(\xi') \geq I_{1,a}(\xi) + \langle \rho, \xi' - \xi \rangle \text{ for every } \xi' \in \mathcal{D}\}. \quad (\text{C.1})$$

$\partial I_{1,q}(\xi)$, $\partial \Lambda_{1,q}(\rho)$ and $\partial \log \phi_a(\rho)$ are defined similarly. Note that $\log \phi_a$ is a smooth function, therefore

$$\partial \log \phi_a(\rho) = \{\nabla \log \phi_a(\rho)\}$$

at every $\rho \in \mathbb{R}^d$ (see [45, Theorem 25.1]).

Theorem C.1. *If $\xi \in \text{ri}(\mathcal{D})$, then $\partial I_{1,a}(\xi)$ and $\partial I_{1,q}(\xi)$ are nonempty and convex. For every $\rho \in \mathbb{R}^d$,*

$$\rho \in \partial I_{1,a}(\xi) \iff I_{1,a}(\xi) + \log \phi_a(\rho) = \langle \rho, \xi \rangle \iff \xi = \nabla \log \phi_a(\rho) \quad (\text{C.2})$$

and

$$\rho \in \partial I_{1,q}(\xi) \iff I_{1,q}(\xi) + \Lambda_{1,q}(\rho) = \langle \rho, \xi \rangle \iff \xi \in \partial \Lambda_{1,q}(\rho). \quad (\text{C.3})$$

Proof. These statements are special instances of [45, Theorems 23.4 and 23.5]. (Note that convexity is clear from the definition of subdifferentials.) \square

There is a unique linear subspace L of \mathbb{R}^d , given by $L := M - M = \{\xi - \xi' : \xi, \xi' \in M\}$, that is parallel to M , i.e., $M = \xi + L$ for every $\xi \in M$ (see [45, Theorem 1.2]). Set $\dim(\mathcal{D}) = \dim(L)$, where \dim denotes dimension. Let L^\perp be the orthogonal complement of L in \mathbb{R}^d .

Theorem C.2. *For every $\xi \in \text{ri}(\mathcal{D})$:*

- (a) $\partial I_{1,a}(\xi)$ is an affine set that is parallel to L^\perp , i.e., $\partial I_{1,a}(\xi) = \rho + L^\perp$ for every $\rho \in \partial I_{1,a}(\xi)$.
- (b) $\langle \rho, z \rangle - \log \phi_a(\rho) = \langle \rho', z \rangle - \log \phi_a(\rho')$ for every $\rho, \rho' \in \partial I_{1,a}(\xi)$ and $z \in \mathcal{R}$.
- (c) $\dim(\mathcal{D}) + \dim(\partial I_{1,a}(\xi)) = d$.
- (d) $I_{1,a}$ is differentiable at ξ if and only if $\dim(\mathcal{D}) = d$.

Proof.

- (a) That $\rho + L^\perp \subset \partial I_{1,a}(\xi)$ for any $\rho \in \partial I_{1,a}(\xi)$ follows immediately from definition (C.1) because $\langle \rho', \xi' - \xi \rangle = 0$ for all $\xi, \xi' \in \mathcal{D}$ and $\rho' \in L^\perp$.

Conversely, suppose $\rho' \notin L^\perp$. Then $\langle \rho', z \rangle$ is not constant over $z \in \mathcal{R}$, and

$$\langle \rho', \mathbf{J}(\nabla \log \phi_a)(\rho) \rho' \rangle = E_0[e^{\langle \rho, Z_1 \rangle - \log \phi_a(\rho)} \langle \rho', Z_1 \rangle^2] - E_0[e^{\langle \rho, Z_1 \rangle - \log \phi_a(\rho)} \langle \rho', Z_1 \rangle]^2 > 0$$

by Jensen's inequality. Here, \mathbf{J} denotes the Jacobian and $\mathbf{J}(\nabla \log \phi_a)(\rho)$ is the Hessian matrix of $\log \phi_a$ at ρ . Therefore, $\nabla \log \phi_a(\rho + \epsilon \rho') \neq \xi$ for sufficiently small $\epsilon > 0$, and hence $\rho + \epsilon \rho' \notin \partial I_{1,a}(\xi)$ by (C.2). Since $\partial I_{1,a}(\xi)$ is convex, we deduce that $\rho + \rho' \notin \partial I_{1,a}(\xi)$.

- (b) If $\rho, \rho' \in \partial I_{1,a}(\xi)$, then $\rho - \rho' \in L^\perp$ by part (a), and so $\langle \rho - \rho', z' \rangle$ is constant over $z' \in \mathcal{R}$. Consequently, for any particular $z \in \mathcal{R}$,

$$\begin{aligned} \langle \rho, z \rangle - \log \phi_a(\rho) &= \langle \rho - \rho' + \rho', z \rangle - \log \sum_{z' \in \mathcal{R}} \hat{q}(z') e^{\langle \rho - \rho' + \rho', z' \rangle} \\ &= \langle \rho', z \rangle - \log \sum_{z' \in \mathcal{R}} \hat{q}(z') e^{\langle \rho', z' \rangle} = \langle \rho', z \rangle - \log \phi_a(\rho'). \end{aligned} \quad (\text{C.4})$$

- (c) $\dim(\mathcal{D}) + \dim(\partial I_{1,a}(\xi)) = \dim(L) + \dim(L^\perp) = d$ by part (a).

(d) This follows from part (c) and [45, Theorem 25.1]. □

The next proposition states some properties of $\partial I_{1,a}(\xi^*)$ where $\xi^* = \sum_{z \in \mathcal{R}} \hat{q}(z)z$ is the LLN velocity. It is used in conjunction with assumption (3.19) for results on spatially constant environments.

Proposition C.3. *For every $\rho \in \mathbb{R}^d$, the following are equivalent:*

- (i) $\phi_a(\rho) = e^{\langle \rho, \xi^* \rangle}$;
- (ii) $\rho \in \partial I_{1,a}(\xi^*)$;
- (iii) $\langle \rho, z \rangle = \langle \rho, \xi^* \rangle$ for every $z \in \mathcal{R}$.

Proof. For every $\rho \in \mathbb{R}^d$,

$$\log \phi_a(\rho) = \log \sum_{z \in \mathcal{R}} \hat{q}(z) e^{\langle \rho, z \rangle} \geq \sum_{z \in \mathcal{R}} \hat{q}(z) \langle \rho, z \rangle = \langle \rho, \xi^* \rangle$$

by Jensen's inequality, and equality holds if and only if $\langle \rho, z \rangle$ is constant over $z \in \mathcal{R}$. This proves the equivalence of (i) and (iii).

Observe that $\nabla \log \phi_a(0) = \sum_{z \in \mathcal{R}} \hat{q}(z)z = \xi^*$. Therefore, $0 \in \partial I_{1,a}(\xi^*)$ by (C.2), and $\partial I_{1,a}(\xi^*) = L^\perp$ by Theorem C.2. The equivalence of (ii) and (iii) now follows since $\{z - \xi^* : z \in \mathcal{R}\}$ spans L . □

When the environment is spatially constant, recall from (10.1) that

$$\Lambda_{1,q}(\rho) = \mathbb{E}[\log W(\rho, \omega)].$$

In particular, it is a smooth function and

$$\partial \Lambda_{1,q}(\rho) = \{\nabla \Lambda_{1,q}(\rho)\} \tag{C.5}$$

at every $\rho \in \mathbb{R}^d$. In this case, the following quenched version of Theorem C.2 holds, with the same proof.

Theorem C.4. *Assume (2.4) and (3.18). Then, for every $\xi \in \text{ri}(\mathcal{D})$:*

- (a) $\partial I_{1,q}(\xi)$ is an affine set that is parallel to L^\perp , i.e., $\partial I_{1,q}(\xi) = \rho + L^\perp$ for every $\rho \in \partial I_{1,q}(\xi)$.
- (b) $\langle \rho, z \rangle - \log W(\rho, \omega) = \langle \rho', z \rangle - \log W(\rho', \omega)$ for all $\rho, \rho' \in \partial I_{1,q}(\xi)$, $z \in \mathcal{R}$ and \mathbb{P} -a.e. ω .
- (c) $\dim(\mathcal{D}) + \dim(\partial I_{1,q}(\xi)) = d$.
- (d) $I_{1,q}$ is differentiable at ξ if and only if $\dim(\mathcal{D}) = d$.

D A concentration inequality

Consider a random walk on \mathbb{Z}^d starting at the origin whose steps are independent and uniformly distributed on \mathcal{R} . Denote the corresponding path measure (resp. expectation) by \hat{P}_0 (resp. \hat{E}_0). For any $\rho \in \mathbb{R}^d$, define a function $\eta : \Omega \times \mathcal{R} \rightarrow \mathbb{R}$ by

$$\eta(\omega, z) = \langle \rho, z \rangle + \log(|\mathcal{R}| \pi_{0,1}(0, z | \omega)).$$

With this notation,

$$u_n(\omega) = E_0^\omega [e^{\langle \rho, X_n \rangle - n \log \phi_a(\rho)}] = \hat{E}_0 [e^{\sum_{i=0}^{n-1} (\eta(T_i, X_i, \omega, Z_{i+1}) - \log \phi_a(\rho))}].$$

This representation enables one to study RWDRE via techniques developed in the context of directed polymers (see [10] for a survey). For instance, the following result is an adaptation of a concentration inequality by Liu and Watbled for the quenched free energy of directed polymers (see [32, Section 6]).

Theorem D.1. Assume (3.16). Then, for every $\rho \in \mathbb{R}^d$ and $\varepsilon > 0$, $\exists c = c(\rho, \varepsilon) > 0$ such that

$$\mathbb{P}(|\log u_n - \mathbb{E}[\log u_n]| \geq n\varepsilon) \leq 2 \exp(-cn).$$

Proof. We can write $\log u_n - \mathbb{E}[\log u_n]$ as a sum of $(\mathfrak{S}_{0,i+1})_{0 \leq i \leq n-1}$ martingale differences:

$$\log u_n - \mathbb{E}[\log u_n] = \sum_{i=0}^{n-1} V_{n,i}, \quad \text{with} \quad V_{n,i} = \mathbb{E}_{i+1}[\log u_n] - \mathbb{E}_i[\log u_n],$$

where $\mathbb{E}_i[\dots]$ is shorthand for $\mathbb{E}[\dots | \mathfrak{S}_{0,i}]$.

Lemma D.2. For every $0 \leq i \leq n - 1$ and $t \in \mathbb{R}$,

$$\mathbb{E}_i[\exp(tV_{n,i})] \leq K(t) := \begin{cases} (\sup_{z \in \mathcal{R}} \mathbb{E}[e^{-|t|\eta(0,0,z)}]) (\sup_{z \in \mathcal{R}} \mathbb{E}[e^{\eta(0,0,z)}]^{|t|}) & \text{if } |t| < 1; \\ (\sup_{z \in \mathcal{R}} \mathbb{E}[e^{-|t|\eta(0,0,z)}]) (\sup_{z \in \mathcal{R}} \mathbb{E}[e^{|t|\eta(0,0,z)}]) & \text{if } |t| \geq 1. \end{cases}$$

Proof. Set

$$e_{n,i} = \exp \left(\sum_{0 \leq j \leq n-1, j \neq i} (\eta(T_j, X_j \omega, Z_{j+1}) - \log \phi_a(\rho)) \right), \quad u_{n,i} = \hat{E}_0[e_{n,i}].$$

Since $\mathbb{E}_{i+1}[\log u_{n,i}] = \mathbb{E}_i[\log u_{n,i}]$, we have

$$V_{n,i} = \mathbb{E}_{i+1} \left[\log \frac{u_n}{u_{n,i}} \right] - \mathbb{E}_i \left[\log \frac{u_n}{u_{n,i}} \right]. \tag{D.1}$$

For every $x \in \mathbb{Z}^d$ and $z \in \mathcal{R}$, define

$$\bar{\eta}(i, x, z) = \exp(\eta(T_{i,x} \omega, z) - \log \phi_a(\rho)), \quad \alpha(i, x, z) = \frac{\hat{E}_0[e_{n,i} \mathbb{1}_{\{X_i=x, Z_{i+1}=z\}}]}{u_{n,i}}.$$

Then,

$$\sum_{x \in \mathbb{Z}^d} \sum_{z \in \mathcal{R}} \alpha(i, x, z) = 1 \quad \text{and} \quad \frac{u_n}{u_{n,i}} = \sum_{x \in \mathbb{Z}^d} \sum_{z \in \mathcal{R}} \alpha(i, x, z) \bar{\eta}(i, x, z).$$

By (D.1), Jensen's inequality and the fact that $\mathfrak{S}_{0,i} \subset \mathfrak{S}_{0,i+1}$, we get

$$\begin{aligned} \mathbb{E}_i[\exp(tV_{n,i})] &= \exp \left(-t \mathbb{E}_i \left[\log \frac{u_n}{u_{n,i}} \right] \right) \mathbb{E}_i \left[\exp \left(t \mathbb{E}_{i+1} \left[\log \frac{u_n}{u_{n,i}} \right] \right) \right] \\ &\leq \mathbb{E}_i \left[\left(\frac{u_n}{u_{n,i}} \right)^{-t} \right] \mathbb{E}_i \left[\left(\frac{u_n}{u_{n,i}} \right)^t \right]. \end{aligned} \tag{D.2}$$

If $t < 0$ or $t \geq 1$, then the function $u \rightarrow u^t$ is convex; therefore Jensen's inequality gives

$$\left(\frac{u_n}{u_{n,i}} \right)^t = \left(\sum_{x \in \mathbb{Z}^d} \sum_{z \in \mathcal{R}} \alpha(i, x, z) \bar{\eta}(i, x, z) \right)^t \leq \sum_{x \in \mathbb{Z}^d} \sum_{z \in \mathcal{R}} \alpha(i, x, z) (\bar{\eta}(i, x, z))^t.$$

For every $x \in \mathbb{Z}^d$ and $z \in \mathcal{R}$, the random variables $\bar{\eta}(i, x, z)$ and $\alpha(i, x, z)$ are measurable w.r.t. $\sigma\{\omega_i\}$ and $\sigma\{\omega_j : 0 \leq j \leq n - 1, j \neq i\}$, respectively. Since these two σ -algebras are independent and the latter one contains $\mathfrak{S}_{0,i}$, we get

$$\mathbb{E}_i[\alpha(i, x, z) (\bar{\eta}(i, x, z))^t] = \mathbb{E}_i[\alpha(i, x, z)] \mathbb{E}[(\bar{\eta}(i, x, z))^t] = \mathbb{E}_i[\alpha(i, x, z)] \frac{\mathbb{E}[e^{t\eta(0,0,z)}]}{(\phi_a(\rho))^t}.$$

Hence, for $t < 0$ or $t \geq 1$,

$$\mathbb{E}_i \left[\left(\frac{u_n}{u_{n,i}} \right)^t \right] \leq \sup_{z \in \mathcal{R}} \frac{\mathbb{E}[e^{t\eta(0,0,z)}]}{(\phi_a(\rho))^t}.$$

If $t \in (0, 1)$, then the function $u \rightarrow u^t$ is concave; therefore Jensen's inequality gives

$$\mathbb{E}_i \left[\left(\frac{u_n}{u_{n,i}} \right)^t \right] \leq \left(\mathbb{E}_i \left[\frac{u_n}{u_{n,i}} \right] \right)^t \leq \sup_{z \in \mathcal{R}} \frac{(\mathbb{E}[e^{\eta(0,0,z)}])^t}{(\phi_a(\rho))^t}.$$

The desired result follows from plugging these bounds in (D.2). □

Continuing with the proof of Theorem D.1, recall the ellipticity assumption (3.16). Lemma D.2 implies

$$\mathbb{E}_i[\exp(\delta|V_{n,i}|)] \leq \mathbb{E}_i[\exp(\delta V_{n,i})] + \mathbb{E}_i[\exp(-\delta V_{n,i})] \leq 2K(\delta).$$

Since $\mathbb{E}[e^{-\delta\eta(\cdot,z)}] = (|\mathcal{R}|e^{\langle \rho, z \rangle})^{-\delta} \mathbb{E}[\omega_{0,0}(z)^{-\delta}]$, we deduce that $K(\delta) < \infty$. A suitable generalization of the Azuma-Hoeffding inequality (see [32, Theorem 2.1]) gives

$$\mathbb{E} \left[e^{\delta t(\log u_n - \mathbb{E}[\log u_n])} \right] \leq \exp \left(\frac{2nK(\delta)t^2}{1-t} \right)$$

for every $t \in (0, 1)$. Therefore,

$$\mathbb{P}(|\log u_n - \mathbb{E}[\log u_n]| \geq n\varepsilon) \leq \exp \left(-n\varepsilon\delta t + \frac{2nK(\delta)t^2}{1-t} \right)$$

by the exponential Chebyshev inequality. The desired result is obtained by optimizing over $t \in (0, 1)$. □

References

- [1] L. Avena, F. den Hollander, and F. Redig, *Large deviation principle for one-dimensional random walk in dynamic random environment: attractive spin-flips and simple symmetric exclusion*, Markov Process. Related Fields **16** (2010), no. 1, 139–168. MR-2664339
- [2] J. R. Baxter, N. C. Jain, and T. O. Seppäläinen, *Large deviations for nonstationary arrays and sequences*, Illinois J. Math. **37** (1993), no. 2, 302–328. MR-1208824
- [3] N. Berger, A. Drewitz, and A. F. Ramírez, *Effective polynomial ballisticity conditions for random walk in random environment*, Comm. Pure Appl. Math. **67** (2014), no. 12, 1947–1973. MR-3272364
- [4] C. Boldrighini, R. A. Minlos, and A. Pellegrinotti, *Discrete-time random motion in a continuous random medium*, Stochastic Process. Appl. **119** (2009), no. 10, 3285–3299. MR-2568274
- [5] E. Bolthausen and A.-S. Sznitman, *On the static and dynamic points of view for certain random walks in random environment*, Methods Appl. Anal. **9** (2002), no. 3, 345–375, Special issue dedicated to Daniel W. Stroock and Srinivasa S. R. Varadhan on the occasion of their 60th birthday. MR-2023130
- [6] E. Bolthausen, A.-S. Sznitman, and O. Zeitouni, *Cut points and diffusive random walks in random environment*, Ann. Inst. H. Poincaré Probab. Statist. **39** (2003), no. 3, 527–555. MR-1978990
- [7] D. Campos, A. Drewitz, A. F. Ramírez, F. Rassoul-Agha, and T. Seppäläinen, *Level 1 quenched large deviation principle for random walk in dynamic random environment*, Bull. Inst. Math. Acad. Sin. (N.S.) **8** (2013), no. 1, 1–29. MR-3097414
- [8] F. Comets, *Large deviation estimates for a conditional probability distribution. Applications to random interaction Gibbs measures*, Probab. Theory Related Fields **80** (1989), no. 3, 407–432. MR-0976534

- [9] F. Comets, N. Gantert, and O. Zeitouni, *Quenched, annealed and functional large deviations for one-dimensional random walk in random environment*, *Probab. Theory Related Fields* **118** (2000), no. 1, 65–114. MR-1785454
- [10] F. Comets, T. Shiga, and N. Yoshida, *Probabilistic analysis of directed polymers in a random environment: a review*, *Stochastic analysis on large scale interacting systems*, *Adv. Stud. Pure Math.*, vol. 39, Math. Soc. Japan, Tokyo, 2004, pp. 115–142. MR-2073332
- [11] F. Comets and V. Vargas, *Majorizing multiplicative cascades for directed polymers in random media*, *ALEA Lat. Am. J. Probab. Math. Stat.* **2** (2006), 267–277. MR-2249671
- [12] F. Comets and N. Yoshida, *Directed polymers in random environment are diffusive at weak disorder*, *Ann. Probab.* **34** (2006), no. 5, 1746–1770. MR-2271480
- [13] A. De Masi, P. A. Ferrari, S. Goldstein, and W. D. Wick, *An invariance principle for reversible Markov processes. Applications to random motions in random environments*, *J. Statist. Phys.* **55** (1989), no. 3–4, 787–855. MR-1003538
- [14] A. Dembo and O. Zeitouni, *Large deviations techniques and applications*, *Stochastic Modelling and Applied Probability*, vol. 38, Springer-Verlag, Berlin, 2010, Corrected reprint of the second (1998) edition. MR-2571413
- [15] F. den Hollander, *Large deviations*, *Fields Institute Monographs*, vol. 14, American Mathematical Society, Providence, RI, 2000. MR-1739680
- [16] F. den Hollander, *Random polymers*, *Lecture Notes in Mathematics*, vol. 1974, Springer-Verlag, Berlin, 2009, Lectures from the 37th Probability Summer School held in Saint-Flour, 2007. MR-2504175
- [17] J.-D. Deuschel and D. W. Stroock, *Large deviations*, *Pure and Applied Mathematics*, vol. 137, Academic Press, Inc., Boston, MA, 1989. MR-0997938
- [18] M. D. Donsker and S. R. S. Varadhan, *Asymptotic evaluation of certain Markov process expectations for large time. I. II*, *Comm. Pure Appl. Math.* **28** (1975), 1–47; *ibid.* **28** (1975), 279–301. MR-0386024
- [19] M. D. Donsker and S. R. S. Varadhan, *Asymptotic evaluation of certain Markov process expectations for large time. III*, *Comm. Pure Appl. Math.* **29** (1976), no. 4, 389–461. MR-0428471
- [20] M. D. Donsker and S. R. S. Varadhan, *Asymptotic evaluation of certain Markov process expectations for large time. IV*, *Comm. Pure Appl. Math.* **36** (1983), no. 2, 183–212. MR-0690656
- [21] R. Durrett, *Probability: theory and examples*, fourth ed., *Cambridge Series in Statistical and Probabilistic Mathematics*, Cambridge University Press, Cambridge, 2010. MR-2722836
- [22] R. S. Ellis, *Entropy, large deviations, and statistical mechanics*, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 271, Springer-Verlag, New York, 1985. MR-0793553
- [23] E. Emrah and C. Janjigian, *Large deviations for some corner growth models with inhomogeneity*, To appear in *Markov Process. Related Fields*, available at arXiv:1509.02234.
- [24] N. Georgiou, F. Rassoul-Agha, and T. Seppäläinen, *Variational formulas and cocycle solutions for directed polymer and percolation models*, *Comm. Math. Phys.* **346** (2016), no. 2, 741–779. MR-3535900
- [25] N. Georgiou, F. Rassoul-Agha, and T. Seppäläinen, *Stationary cocycles and Busemann functions for the corner growth model*, To appear in *Probab. Theory Related Fields*, available at arXiv:1510.00859.
- [26] N. Georgiou, F. Rassoul-Agha, and T. Seppäläinen, *Geodesics and the competition interface for the corner growth model*, To appear in *Probab. Theory Related Fields*, available at arXiv:1510.00860.
- [27] N. Georgiou, F. Rassoul-Agha, T. Seppäläinen, and A. Yilmaz, *Ratios of partition functions for the log-gamma polymer*, *Ann. Probab.* **43** (2015), no. 5, 2282–2331. MR-3395462
- [28] I. Ignatiouk-Robert, *Large deviations for a random walk in dynamical random environment*, *Ann. Inst. H. Poincaré Probab. Statist.* **34** (1998), no. 5, 601–636. MR-1641666

- [29] M. Joseph and F. Rassoul-Agha, *Almost sure invariance principle for continuous-space random walk in dynamic random environment*, ALEA Lat. Am. J. Probab. Math. Stat. **8** (2011), 43–57. MR-2748407
- [30] S. M. Kozlov, *The averaging method and walks in inhomogeneous environments*, Uspekhi Mat. Nauk **40** (1985), no. 2(242), 61–120, 238. MR-0786087
- [31] G. F. Lawler, *Weak convergence of a random walk in a random environment*, Comm. Math. Phys. **87** (1982/83), no. 1, 81–87. MR-0680649
- [32] Q. Liu and F. Watbled, *Exponential inequalities for martingales and asymptotic properties of the free energy of directed polymers in a random environment*, Stochastic Process. Appl. **119** (2009), no. 10, 3101–3132. MR-2568267
- [33] S. V. Nagaev, *Large deviations of sums of independent random variables*, Ann. Probab. **7** (1979), no. 5, 745–789. MR-0542129
- [34] C. M. Newman, *A surface view of first-passage percolation*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Zürich, 1994) (Basel), Birkhäuser, 1995, pp. 1017–1023. MR-1404001
- [35] G. C. Papanicolaou and S. R. S. Varadhan, *Boundary value problems with rapidly oscillating random coefficients*, Random fields, Vol. I, II (Esztergom, 1979), Colloq. Math. Soc. János Bolyai, vol. 27, North-Holland, Amsterdam-New York, 1981, pp. 835–873. MR-0712714
- [36] F. Rassoul-Agha, *The point of view of the particle on the law of large numbers for random walks in a mixing random environment*, Ann. Probab. **31** (2003), no. 3, 1441–1463. MR-1989439
- [37] F. Rassoul-Agha, *Large deviations for random walks in a mixing random environment and other (non-Markov) random walks*, Comm. Pure Appl. Math. **57** (2004), no. 9, 1178–1196. MR-2059678
- [38] F. Rassoul-Agha and T. Seppäläinen, *An almost sure invariance principle for random walks in a space-time random environment*, Probab. Theory Related Fields **133** (2005), no. 3, 299–314. MR-2198014
- [39] F. Rassoul-Agha and T. Seppäläinen, *Almost sure functional central limit theorem for ballistic random walk in random environment*, Ann. Inst. Henri Poincaré Probab. Stat. **45** (2009), no. 2, 373–420. MR-2521407
- [40] F. Rassoul-Agha and T. Seppäläinen, *Process-level quenched large deviations for random walk in random environment*, Ann. Inst. Henri Poincaré Probab. Stat. **47** (2011), no. 1, 214–242. MR-2779403
- [41] F. Rassoul-Agha and T. Seppäläinen, *Quenched point-to-point free energy for random walks in random potentials*, Probab. Theory Related Fields **158** (2014), no. 3–4, 711–750. MR-3176363
- [42] F. Rassoul-Agha and T. Seppäläinen, *A course on large deviations with an introduction to Gibbs measures*, Graduate Studies in Mathematics, vol. 162, American Mathematical Society, Providence, RI, 2015. MR-3309619
- [43] F. Rassoul-Agha, T. Seppäläinen, and A. Yilmaz, *Quenched free energy and large deviations for random walks in random potentials*, Comm. Pure Appl. Math. **66** (2013), no. 2, 202–244. MR-2999296
- [44] F. Rassoul-Agha, T. Seppäläinen, and A. Yilmaz, *Variational formulas and disorder regimes of random walks in random potentials*, Bernoulli **23** (2017), no. 1, 405–431. MR-3556777
- [45] R. T. Rockafellar, *Convex analysis*, Princeton Mathematical Series, No. 28, Princeton University Press, Princeton, N.J., 1970. MR-0274683
- [46] J. M. Rosenbluth, *Quenched large deviation for multidimensional random walk in random environment: A variational formula*, ProQuest LLC, Ann Arbor, MI, 2006, Thesis (Ph.D.)–New York University. MR-2708406
- [47] T. Seppäläinen, *Large deviations for lattice systems. I. Parametrized independent fields*, Probab. Theory Related Fields **96** (1993), no. 2, 241–260. MR-1227034
- [48] S. R. S. Varadhan, *Large deviations for random walks in a random environment*, Comm. Pure Appl. Math. **56** (2003), no. 8, 1222–1245, Dedicated to the memory of Jürgen K. Moser. MR-1989232

- [49] A. Yilmaz, *Large deviations for random walk in a space-time product environment*, Ann. Probab. **37** (2009), no. 1, 189–205. MR-2489163
- [50] A. Yilmaz, *Quenched large deviations for random walk in a random environment*, Comm. Pure Appl. Math. **62** (2009), no. 8, 1033–1075. MR-2531552
- [51] A. Yilmaz, *Harmonic functions, h -transform and large deviations for random walks in random environments in dimensions four and higher*, Ann. Probab. **39** (2011), no. 2, 471–506. MR-2789504
- [52] A. Yilmaz and O. Zeitouni, *Differing averaged and quenched large deviations for random walks in random environments in dimensions two and three*, Comm. Math. Phys. **300** (2010), no. 1, 243–271. MR-2725188
- [53] O. Zeitouni, *Random walks in random environment*, Lectures on probability theory and statistics, Lecture Notes in Math., vol. 1837, Springer, Berlin, 2004, pp. 189–312. MR-2071631

Electronic Journal of Probability

Electronic Communications in Probability

Advantages of publishing in EJP-ECP

- Very high standards
- Free for authors, free for readers
- Quick publication (no backlog)
- Secure publication (LOCKSS¹)
- Easy interface (EJMS²)

Economical model of EJP-ECP

- Non profit, sponsored by IMS³, BS⁴, ProjectEuclid⁵
- Purely electronic

Help keep the journal free and vigorous

- Donate to the IMS open access fund⁶ (click here to donate!)
- Submit your best articles to EJP-ECP
- Choose EJP-ECP over for-profit journals

¹LOCKSS: Lots of Copies Keep Stuff Safe <http://www.lockss.org/>

²EJMS: Electronic Journal Management System <http://www.vtex.lt/en/ejms.html>

³IMS: Institute of Mathematical Statistics <http://www.imstat.org/>

⁴BS: Bernoulli Society <http://www.bernoulli-society.org/>

⁵Project Euclid: <https://projecteuclid.org/>

⁶IMS Open Access Fund: <http://www.imstat.org/publications/open.htm>