

Fractional Brownian motion with zero Hurst parameter: a rough volatility viewpoint

Eyal Neuman* Mathieu Rosenbaum†

Abstract

Rough volatility models are becoming increasingly popular in quantitative finance. In this framework, one considers that the behavior of the log-volatility process of a financial asset is close to that of a fractional Brownian motion with Hurst parameter around 0.1. Motivated by this, we wish to define a natural and relevant limit for the fractional Brownian motion when H goes to zero. We show that once properly normalized, the fractional Brownian motion converges to a Gaussian random distribution which is very close to a log-correlated random field.

Keywords: fractional Brownian motion; log-correlated random field; rough volatility; multifractal processes.

AMS MSC 2010: Primary 60G22; 60G15; 60G57, Secondary 60G18; 28A80.

Submitted to ECP on November 1, 2017, final version accepted on July 26, 2018.

1 Introduction

The fractional Brownian motion (fBm for short) is a very popular modeling object in many fields such as hydrology, see for example [29], telecommunications and network traffic, see [23, 28] among others and finance, see the seminal paper [11]. A fBm $(B_t^H)_{t \in \mathbb{R}}$ with Hurst parameter $H \in (0, 1)$ is a zero-mean Gaussian process with covariance kernel given by

$$\mathbb{E}[B_t^H B_s^H] = \frac{1}{2} (|t|^{2H} + |s|^{2H} - |t - s|^{2H}).$$

It has stationary increments and is self-similar with parameter H , that is $(B_{at}^H)_{t \in \mathbb{R}}$ has the same law as $(a^H B_t^H)_{t \in \mathbb{R}}$ for any $a > 0$. Furthermore, sample paths of fBm have almost surely Hölder regularity $H - \varepsilon$ for any $\varepsilon > 0$.

One of the main probabilistic features which motivates the use of fBm in the applications mentioned above, is the long memory property of the increments when $H > 1/2$. This means that for $H > 1/2$, we have

$$\sum_{i=1}^{+\infty} \text{Cov}[(B_{i+1}^H - B_i^H), B_1^H] = +\infty.$$

Thus the auto-covariance function of the fBm increments decays slowly, which is interesting when modeling persistent phenomena.

*Imperial College London, Department of Mathematics. E-mail: e.neumann@imperial.ac.uk

†École Polytechnique, CMAP. E-mail: mathieu.rosenbaum@polytechnique.edu

However, recently, a new paradigm has been introduced in [20] for the use of fBm in finance. Indeed, a careful analysis of financial time series suggests that the log-volatility process, that is the intensity of the price variations of an asset, actually behaves like a fBm with Hurst parameter of order 0.1. Hence various approaches using a fBm with small Hurst parameter have been introduced for volatility modeling. These models are referred to as *rough volatility models*, see [4, 5, 6, 8, 13, 14, 16, 21] for more details and practical applications.

Such small estimated values for H (between 0.05 and 0.2) have been found when studying the volatility process of thousands of assets, see [7]. Consequently, a natural question is the behavior of the fBm in the limiting case when H is sent to zero. Of course, putting directly $H = 0$ in the covariance function does not lead to a relevant process. Thus, in this work, we wish to build a suitable sequence of normalized fBms and describe its limit as H goes to zero. This will lead us to a possible definition of the fractional Brownian motion for $H = 0$. Note that several authors already defined some fractional Brownian motion for $H = 0$, see in particular [19]. This is usually done through a regularization procedure. Our approach here is quite simple and probably more natural from the financial viewpoint we have in mind. Instead of regularizing the process, we choose to normalize it in order to get a non-degenerate limit. Our normalized sequence of processes $(X^H)_{H \in (0,1)}$ is defined through

$$X_t^H = \frac{B_t^H - \frac{1}{t} \int_0^t B_u^H du}{\sqrt{H}}, \quad t \in \mathbb{R},$$

where $X_0^H = 0$. Subtracting the integral in the numerator and dividing by \sqrt{H} enables us to get a non-trivial limit for our sequence as H goes to 0.

Our main result is the convergence of X^H , seen as a random element in the space of tempered distributions, towards an approximately *log-correlated Gaussian field*. Denote by \mathcal{S} the real Schwartz space, that is the set of real-valued functions on \mathbb{R} whose derivatives of all orders exist and decay faster than any polynomial at infinity. We write \mathcal{S}' for the dual of \mathcal{S} , that is the space of tempered distributions. We also define the subspace \mathcal{S}_0 of the real Schwartz space, consisting of functions ϕ from \mathcal{S} with $\int_{\mathbb{R}} \phi(s) ds = 0$, and its topological dual \mathcal{S}'/\mathbb{R} . A log-correlated Gaussian field (LGF for short) $X \in \mathcal{S}'/\mathbb{R}$, is a centered Gaussian field whose covariance kernel satisfies

$$\mathbb{E}[\langle X, \phi_1 \rangle \langle X, \phi_2 \rangle] = \int_{\mathbb{R}} \int_{\mathbb{R}} \log \frac{1}{|t-s|} \phi_1(t) \phi_2(s) dt ds,$$

for any $\phi_1, \phi_2 \in \mathcal{S}_0$, see [12] for an overview on LGF. We show in this paper that the limit of X^H as H goes to zero is "almost" a log-correlated Gaussian field, see Section 2 for an accurate result.

LGFs are closely related to some multifractal processes pioneered by Mandelbrot (see for example [26]), and further developed in [1, 2, 3, 9], among others. A process $(Y_t)_{t \geq 0}$ is said to be multifractal if for a range of values of q , we have for some $T > 0$

$$\mathbb{E}[|Y_{t+\ell} - Y_t|^q] \sim C(q) \ell^{\zeta(q)}, \quad \text{for } 0 < \ell \leq T,$$

where $C(q) > 0$ is a constant and $\zeta(\cdot)$ is a non-linear concave function. In particular, the multifractal random walk model for the log-price of an asset in [1] satisfies such property. It is defined as $Y_t = B_{M([0,t])}$, where B is a Brownian motion and

$$M(t) = \lim_{l \rightarrow 0} \sigma^2 \int_0^t e^{w_l(u)} du, \quad \text{a.s.},$$

with $\sigma^2 > 0$ and w_l a Gaussian process such that for some $\lambda^2 > 0$ and $T > 0$

$$\text{Cov}[w_l(t), w_l(t')] = \lambda^2 \log(T/|t - t'|), \text{ for } l < |t - t'| \leq T,$$

see [2] for details. Hence we see that M formally corresponds to a measure of the form $\exp(X_t)dt$, where X is a LGF. For the precise definition of such measures, see [22] and the generalizations on Gaussian multiplicative chaos in [30, 31, 32] and the references therein. Finally note that LGFs and more generally the associated theory of Gaussian multiplicative chaos have extensive use in other fields than finance, such as turbulence, see [10, 18], disordered systems, see [17, 25] and Liouville quantum gravity, see [30, 31].

In the following section we introduce our main theorem, that is an accurate statement about the convergence of the normalized fBm towards a LGF as H goes to zero. We also discuss the multifractal properties of the limiting LGF in the same section. The proof of our theorem can be found in Section 3.

2 Convergence of the fBM towards a LGF

2.1 Main result

We define the weak convergence of elements in \mathcal{S}' as in Proposition 12.2 in [24]. We say that X^H converges weakly to X as H tends to 0 if for any $\phi \in \mathcal{S}$ we have

$$\langle X^H, \phi \rangle \rightarrow \langle X, \phi \rangle,$$

in law, as H tends to 0. The main result of our paper is the following.

Theorem 2.1. *The sequence $\{X_t^H\}_{t \in \mathbb{R}}$ converges weakly as H tends to zero towards a centered Gaussian field X satisfying for any $\phi_1, \phi_2 \in \mathcal{S}$*

$$\mathbb{E}[\langle X, \phi_1 \rangle \langle X, \phi_2 \rangle] = \int_{\mathbb{R}} \int_{\mathbb{R}} K(t, s) \phi_1(t) \phi_2(s) dt ds,$$

where for $-\infty < s, t < \infty$, $s \neq t$ and $s, t \neq 0$

$$K(t, s) = \log \frac{1}{|t - s|} + g(t, s),$$

with

$$g(t, s) = \frac{1}{t} \int_0^t \log |s - u| du + \frac{1}{s} \int_0^s \log |t - u| du - \frac{1}{ts} \int_0^t \int_0^s \log |u - v| dudv.$$

We see that when $t, s > \delta$ for some $\delta > 0$, then $g(t, s)$ is a bounded continuous function. Hence the covariance kernel exhibits the same type of singularity as that of a LGF. Consequently, in our framework, the limit when H goes to zero of a normalized version of the fBm is "almost" a LGF.

2.2 Multifractal properties

In [30], the authors study the case of centered Gaussian fields on any domain $D \subset \mathbb{R}$ with covariance kernel satisfying

$$K(x, y) = \log_+ \frac{1}{|x - y|} + f(x, y), \quad x, y \in D, \tag{2.1}$$

where $\log_+(x) := \max\{0, \log x\}$ and $f(x, y)$ is a bounded continuous function. Thus, if we restrict X to the domain $[\delta, 1]$ for some fixed $\delta > 0$, then X is included in the framework

of [30]. In particular, their results on the multifractal spectrum of Gaussian fields apply on this restricted domain.

Motivated by the preceding paragraph, we first fix $\delta > 0$ and define an approximate volatility measure ξ_γ^H by

$$\xi_\gamma^H(dt) = e^{\gamma X_t^H - \frac{\gamma^2}{2} \mathbb{E}[(X_t^H)^2]} dt, \quad \delta \leq t \leq 1,$$

for some constant $\gamma > 0$. Here we assume that $\xi_\gamma^H(\cdot)$ vanishes on $[\delta, 1]^c$. From the result of Theorem 2.1 we deduce the following corollary. In what follows, convergence in the L^1 norm stands for the usual convergence of random variables in L^1 .

Corollary 2.2. *For $\gamma < \sqrt{2}$, $\{\xi_\gamma^H\}_{H \in (0,1)}$ converges as H approaches zero to a random measure ξ_γ in the following sense,*

$$\int_{\mathbb{R}} \phi(t) \xi_\gamma^H(dt) \xrightarrow{L^1} \int_{\mathbb{R}} \phi(t) \xi_\gamma(dt), \quad \text{for all } \phi \in \mathcal{S}.$$

Moreover, the limiting measure ξ_γ is the so-called Gaussian multiplicative chaos.

The proof Corollary 2.2 is given in Section 3. For the definition and properties of Gaussian multiplicative chaos we refer to a survey paper by Rhodes and Vargas [30]. We will briefly explain some of the properties of ξ_γ from the theory of Gaussian multiplicative chaos.

We first describe the behavior of the moments of ξ_γ . Note that for $t \in [\delta, 1]$, K is of the form (2.1). From Proposition 2.5 in [31], it follows that for all $t \in (\delta, 1)$ and $q \in (-\infty, 2/\gamma^2)$, there exists $C(t, q) > 0$ such that

$$\mathbb{E}[\xi_\gamma(B(t, r))^q] \sim C(t, q)r^{\zeta(q)}, \quad \text{as } r \rightarrow 0,$$

where $\zeta(q) = (1 + \gamma^2/2)q - \gamma^2 q^2/2$. Hence we do obtain a multifractal scaling as described in the introduction.

Finally we consider a quantity closely related to the function ζ above: the spectrum of singularities of ξ_γ . For any $0 < \gamma < \sqrt{2}$ and $0 < r < \sqrt{2}/\gamma$, we define

$$G_{\gamma,r} = \left\{ x \in (\delta, 1); \lim_{\varepsilon \rightarrow 0} \frac{\log \xi_\gamma(B(x, \varepsilon))}{\log \varepsilon} = 1 + \left(\frac{1}{2} - r\right)\gamma^2 \right\}.$$

The set $G_{\gamma,r}$ somehow corresponds to the points x where the Hölder regularity of ξ_γ is equal to $1 + (1/2 - r)\gamma^2$. Let $\dim_H(A)$ denote the Hausdorff dimension of a set A . Theorem 2.6 in [31] states that

$$\dim_H(G_{\gamma,r}) = 1 - \frac{\gamma^2 r^2}{2}.$$

In particular, we remark that

$$\dim_H(G_{\gamma,r}) = \inf_{p \in \mathbb{R}} \left\{ p \left(1 + \left(\frac{1}{2} - r\right)\gamma^2 \right) - \zeta(p) + 1 \right\}.$$

This equality means that the Frisch-Parisi conjecture relating the scaling exponents of a process to its spectrum of singularities holds in our case, see [15] and [30] for more details on the multifractal formalism.

The rest of this paper is devoted to the proofs of Theorem 2.1 and Corollary 2.2.

3 Proof of Theorem 2.1 and Corollary 2.2.

For $t, s \in \mathbb{R}$, let $K_H(t, s) = \mathbb{E}[X_t^H X_s^H]$. We start the proof with the following important lemma.

Lemma 3.1. For any non-zero $-\infty < s, t < \infty$ with $s \neq t$, we have

$$\lim_{H \rightarrow 0} K_H(t, s) = K(t, s). \tag{3.1}$$

Proof. We write $K_H(t, s) = I_1^H(t, s) + I_2^H(t, s) + I_3^H(t, s) + I_4^H(t, s)$, where

$$\begin{aligned} I_1^H(t, s) &= -\frac{1}{2H} |t-s|^{2H}, \quad I_2^H(t, s) = \frac{1}{2H} \frac{1}{t} \int_0^t |s-u|^{2H} du, \\ I_3^H(t, s) &= \frac{1}{2H} \frac{1}{s} \int_0^s |t-u|^{2H} du, \quad I_4^H(t, s) = -\frac{1}{2H} \frac{1}{st} \int_0^t \int_0^s |u-v|^{2H} dudv \end{aligned} \tag{3.2}$$

and $K(t, s) = I_1^0(t, s) + I_2^0(t, s) + I_3^0(t, s) + I_4^0(t, s)$, where

$$\begin{aligned} I_1^0(t, s) &= \log \frac{1}{|t-s|}, \quad I_2^0(t, s) = \frac{1}{t} \int_0^t \log |s-u| du, \\ I_3^0(t, s) &= \frac{1}{s} \int_0^s \log |t-u| du, \quad I_4^0(t, s) = -\frac{1}{st} \int_0^t \int_0^s \log |u-v| dudv. \end{aligned}$$

We have

$$K_H(t, s) = K_H(s, t) \text{ and } K(t, s) = K(s, t),$$

so we can assume that $-\infty < s < t < \infty$.

Note that for any $s \neq t$,

$$I_1^H(t, s) + \frac{1}{2H} = -\frac{1}{2H} (e^{2H \log |t-s|} - 1)$$

and therefore

$$\lim_{H \rightarrow 0} I_1^H(t, s) + \frac{1}{2H} = -\log |t-s| = I_1^0(t, s).$$

Next we deal with I_i^H , $i = 2, \dots, 4$. We consider several cases.

Case 1: Assume that $0 < s < t$. We easily get

$$I_2^H(t, s) + I_3^H(t, s) = \frac{1}{2H(2H+1)ts} (t^{2H+2} + s^{2H+2} - (t-s)^{2H+2}).$$

For $I_4^H(t, s)$, note that

$$\int_0^t |u-v|^{2H} du = \frac{1}{2H+1} v^{2H+1} + \frac{1}{2H+1} (t-v)^{2H+1}.$$

Hence we obtain

$$\int_0^s \int_0^t |u-v|^{2H} dudv = \frac{1}{(2H+1)(2H+2)} (t^{2H+2} - (t-s)^{2H+2} + s^{2H+2})$$

and therefore

$$I_4^H(t, s) = -\frac{1}{2H(2H+1)(2H+2)ts} (t^{2H+2} + s^{2H+2} - (t-s)^{2H+2}).$$

Define

$$I_{2,4}^H(t, s) = \sum_{i=2}^4 I_i^H(t, s).$$

It follows that

$$I_{2,4}^H(t, s) = \frac{1}{2H(2H+2)ts} (t^{2H+2} + s^{2H+2} - (t-s)^{2H+2}).$$

Thus we have

$$\begin{aligned} & I_{2,4}^H(t, s) - \frac{1}{H(2H+2)} \\ &= \frac{1}{2H(2H+2)ts} (t^{2H+2} + s^{2H+2} - (t-s)^{2H+2}) - \frac{1}{2H(2H+2)ts} (t^2 + s^2 - (t-s)^2) \\ &= \frac{1}{2H(2H+2)ts} (t^2(t^{2H} - 1) + s^2(s^{2H} - 1) - (t-s)^2((t-s)^{2H} - 1)). \end{aligned}$$

Consequently,

$$\lim_{H \rightarrow 0} I_{2,4}^H(t, s) - \frac{1}{H(2H+2)} = \frac{1}{2ts} (t^2 \log t + s^2 \log s - (t-s)^2 \log(t-s))$$

and

$$\lim_{H \rightarrow 0} I_{2,4}^H(t, s) - \frac{1}{2H} = \frac{1}{2ts} (t^2 \log t + s^2 \log s - (t-s)^2 \log(t-s)) - \frac{1}{2}.$$

We also easily get that for $0 < s < t$

$$I_2^0(t, s) + I_3^0(t, s) = \frac{1}{st} (t^2 \log t + s^2 \log s - (t-s)^2 \log(t-s)) - 2.$$

Furthermore,

$$\int_0^t \log |u-v| du = v \log v + (t-v) \log(t-v) - t.$$

Thus we obtain

$$\begin{aligned} \int_0^s \int_0^t \log |u-v| du dv &= -st + \frac{s^2}{4} (2 \log s - 1) + \frac{t^2}{4} (2 \log t - 1) \\ &\quad - \frac{(t-s)^2}{4} (2 \log(t-s) - 1) \\ &= \frac{1}{2} (t^2 \log t + s^2 \log s - (s-t)^2 \log(s-t) - 3st) \end{aligned}$$

and we deduce

$$I_4^0(t, s) = -\frac{1}{2ts} (t^2 \log t + s^2 \log s - (s-t)^2 \log(s-t)) + \frac{3}{2}.$$

Now define

$$I_{2,4}^0(t, s) = \sum_{i=2}^4 I_i^0(t, s).$$

We have

$$I_{2,4}^0(t, s) = \frac{1}{2st} (t^2 \log t + s^2 \log s - (t-s)^2 \log(t-s)) - \frac{1}{2}.$$

Finally we obtain

$$\lim_{H \rightarrow 0} I_{2,4}^H(t, s) - \frac{1}{2H} = I_{2,4}^0(t, s),$$

from which (3.1) readily follows for $0 < s < t < \infty$.

Case 2: Assume that $-\infty < s < t < 0$. We use that

$$K_H(t, s) = K_H(-t, -s) = K_H(-s, -t) \text{ and } K(t, s) = K(-t, -s) = K(-s, -t)$$

to deduce that (3.1) follows from the proof of Case 1.

Case 3: Assume that $s < 0 < t$. We write $s = -u$. Repeating the same steps as in Case 1, we get

$$\begin{aligned} & I_{2,4}^H(t, -u) - \frac{1}{H(2H + 2)} \\ &= \frac{1}{2H(2H + 2)tu} \left((t + u)^2((t + u)^{2H} - 1) - t^2(t^{2H} - 1) - u^2(u^{2H} - 1) \right), \end{aligned}$$

and

$$I_{2,4}^0(t, -u) = \frac{1}{2ut} \left((t + u)^2 \log(t + u) - t^2 \log t - u^2 \log u \right) - \frac{1}{2}.$$

It follows that

$$\lim_{H \rightarrow 0} I_{2,4}^H(t, -u) - \frac{1}{2H} = I_{2,4}^0(t, -u)$$

and we get (3.1). □

Let $\phi_1, \phi_2 \in \mathcal{S}$. Since X^H and X are centered Gaussians taking values in \mathcal{S}' , to prove Theorem 2.1, it is enough to show that

$$\lim_{H \rightarrow 0} \mathbb{E}[\langle X^H, \phi_1 \rangle \langle X^H, \phi_2 \rangle] = \int_{\mathbb{R}} \int_{\mathbb{R}} K(t, s) \phi_1(t) \phi_2(s) dt ds.$$

Furthermore, for any $H \in (0, 1)$ we have

$$\mathbb{E}[\langle X^H, \phi_1 \rangle \langle X^H, \phi_2 \rangle] = \int_{\mathbb{R}} \int_{\mathbb{R}} K_H(t, s) \phi_1(t) \phi_2(s) ds dt.$$

Hence Theorem 2.1 immediately follows from the next lemma.

Lemma 3.2. For any $\phi_1, \phi_2 \in \mathcal{S}$,

$$\lim_{H \rightarrow 0} \int_{\mathbb{R}} \int_{\mathbb{R}} K_H(t, s) \phi_1(t) \phi_2(s) ds dt = \int_{\mathbb{R}} \int_{\mathbb{R}} K(t, s) \phi_1(t) \phi_2(s) ds dt. \tag{3.3}$$

Proof. First note that for $x \geq 0$, $1 - e^{-x} \leq x$. Therefore for any $0 < |t - s| \leq 1$,

$$\frac{|1 - |t - s|^{2H}|}{2H} = \frac{1}{2H} (1 - e^{2H \log |t - s|}) \leq -\log |t - s|.$$

Moreover,

$$\begin{aligned} - \int \int_{|t-s| \leq 1} \log |t - s| |\phi_1(t)| |\phi_2(s)| dt ds &\leq -\|\phi_2\|_{\infty} \int_{\mathbb{R}} |\phi_1(t)| \int_{|v| \leq 1} \log |v| dv dt \\ &= -2\|\phi_2\|_{\infty} \int_{\mathbb{R}} |\phi_1(t)| \int_0^1 \log(v) dv dt \\ &\leq C \|\phi_1\|_1 \|\phi_2\|_{\infty}. \end{aligned}$$

Hence, from dominated convergence, it follows that

$$\lim_{H \rightarrow 0} \int \int_{|t-s| \leq 1} \frac{1 - |t-s|^{2H}}{2H} \phi_1(t) \phi_2(s) dt ds = - \int \int_{|t-s| \leq 1} \log |t-s| \phi_1(t) \phi_2(s) dt ds.$$

Let $f(x) = x^{2H}/(2H)$, where $x > 0$ and $H \in (0, 1/2)$. Since f is concave, for any $x \geq 1$, we have $0 \leq f(x) - f(1) \leq f'(1)(x - 1)$. Therefore for any $|t - s| \geq 1$,

$$\frac{||t - s|^{2H} - 1|}{2H} \leq |t - s| - 1 \leq 2(|t| + |s|).$$

Since

$$2 \int \int_{|t-s| > 1} (|t| + |s|) |\phi_1(t)| |\phi_2(s)| dt ds < \infty,$$

we get from dominated convergence

$$\lim_{H \rightarrow 0} \int \int_{|t-s| \geq 1} \frac{1 - |t-s|^{2H}}{2H} \phi_1(t) \phi_2(s) dt ds = - \int \int_{|t-s| \geq 1} \log |t-s| \phi_1(t) \phi_2(s) dt ds.$$

It remains to show that

$$\lim_{H \rightarrow 0} \int_{\mathbb{R}} \int_{\mathbb{R}} (I_{2,4}^H(t, s) - \frac{1}{2H}) \phi_1(t) \phi_2(s) dt ds = \int_{\mathbb{R}} \int_{\mathbb{R}} I_{2,4}^0(t, s) \phi_1(t) \phi_2(s) dt ds.$$

We again consider several cases.

Case 1: Assume that $0 < s < t$. We obviously have

$$\frac{s^2}{2H(2H + 2)ts} |s^{2H} - 1| \leq \frac{1}{2H(2H + 2)} |s^{2H} - 1|.$$

Moreover, from a Taylor expansion, we get $t^{2H} - (t-s)^{2H} \leq 2H(t-s)^{2H-1}s$ and therefore

$$\begin{aligned} & \frac{1}{2Hts} |t^2(t^{2H} - 1) - (t-s)^2((t-s)^{2H} - 1)| \\ & \leq \frac{1}{2Hts} (|t^2(t^{2H} - 1) - (t-s)^2(t^{2H} - 1)| \\ & \quad + |(t-s)^2(t^{2H} - 1) - (t-s)^2((t-s)^{2H} - 1)|) \\ & \leq \frac{1}{2Hts} (|t^{2H} - 1| |s^2 - 2st| + 2H(t-s)^2(t-s)^{2H-1}s) \\ & \leq \frac{1}{H} |t^{2H} - 1| + (t-s)^{2H}. \end{aligned}$$

It follows that for any $0 < s < t$,

$$\left| I_{2,4}^H(t, s) - \frac{1}{H(2H + 2)} \right| \leq \frac{1}{2H} |s^{2H} - 1| + \frac{1}{2H} |t^{2H} - 1| + (t-s)^{2H} := L_1^H(t, s). \tag{3.4}$$

Case 1bis: Assume that $0 < t < s$. Since $I_{2,4}^H(t, s) = I_{2,4}^H(s, t)$ we get

$$\left| I_{2,4}^H(t, s) - \frac{1}{H(2H + 2)} \right| \leq L_1^H(s, t).$$

Case 2: Assume that $-\infty < -t, -s < 0$. Since $I_{2,4}^H(-t, -s) = I_{2,4}^H(t, s)$, we have

$$\left| I_{2,4}^H(-t, -s) - \frac{1}{H(2H + 2)} \right| \leq L_1^H(t, s).$$

Case 3: Assume that $-s < 0 < t$ and $s \leq t$. Recall that when $-s < 0 < t$,

$$\begin{aligned} & I_{2,4}^H(t, -s) - \frac{1}{H(2H+2)} \\ &= \frac{1}{2H(2H+2)ts} ((t+s)^2((t+s)^{2H}-1) - t^2(t^{2H}-1) - s^2(s^{2H}-1)). \end{aligned}$$

Since $s < t$ we have

$$\left| \frac{s^2}{2H(2H+2)ts} (s^{2H}-1) \right| \leq \frac{1}{2H(2H+2)} |s^{2H}-1|.$$

Using a Taylor expansion, we get $(t+s)^{2H} - t^{2H} \leq 2Ht^{2H-1}s$ and therefore

$$\begin{aligned} & \frac{1}{2Hts} |(t+s)^2((t+s)^{2H}-1) - t^2(t^{2H}-1)| \\ & \leq \frac{1}{2Hts} (|(t+s)^2((t+s)^{2H}-1) - (t+s)^2(t^{2H}-1)| \\ & \quad + |(t+s)^2(t^{2H}-1) - t^2(t^{2H}-1)|) \\ & \leq \frac{1}{2Hts} (2H(t+s)^2t^{2H-1}s + |t^{2H}-1||s^2+2st|) \\ & \leq (t+s)^{2H} + \frac{3}{2H}|t^{2H}-1|. \end{aligned}$$

It follows that

$$\left| I_{2,4}^H(t, -s) - \frac{1}{H(2H+2)} \right| \leq \frac{1}{2H}|s^{2H}-1| + \frac{1}{H}|t^{2H}-1| + (t+s)^{2H} := L_2^H(t, s).$$

Case 4: Assume that $-s < 0 < t$ and $t < s$. Repeating the same lines as in Case 3 but exchanging the roles of t and s we get

$$\left| I_{2,4}^H(t, -s) - \frac{1}{H(2H+2)} \right| \leq L_2^H(s, t).$$

Case 5: Assume that $-t < 0 < s$ and $t < s$. Since $I_{2,4}^H(-t, s) = I_{2,4}^H(s, -t)$, it follows from Case 3 that

$$\left| I_{2,4}^H(-t, s) - \frac{1}{H(2H+2)} \right| \leq L_2^H(s, t).$$

Case 6: Assume that $-t < 0 < s$ and $s < t$. Using again $I_{2,4}^H(-t, s) = I_{2,4}^H(s, -t)$, we get from Case 4 that

$$\left| I_{2,4}^H(-t, s) - \frac{1}{H(2H+2)} \right| \leq L_2^H(t, s).$$

Finally, we clearly have

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \left(\sup_{H \in (0, 1/2)} (L_1^H(s, t) + L_1^H(t, s) + L_2^H(s, t) + L_2^H(t, s)) \right) \phi_1(t) \phi_2(s) dt ds < \infty.$$

Thus we conclude the proof using the dominated convergence theorem. \square

Before we prove Corollary 2.2, we recall an approximation of $\log_+(1/|x|)$, from Example 2.2 in [30]. For any $x \in \mathbb{R}$ we define the cone $\mathcal{C}(x)$ in $\mathbb{R} \times \mathbb{R}_+$:

$$\mathcal{C}(x) = \left\{ (y, t) \in \mathbb{R} \times \mathbb{R}_+; |x-y| \leq \frac{t \wedge 1}{2} \right\}.$$

A direct computation gives

$$f(x) = \int_{\mathcal{C}(0) \cap \mathcal{C}(x)} \frac{dy dt}{t^2} = \log_+ \frac{1}{|x|}.$$

We approximate f by the following functions,

$$\tilde{f}_\varepsilon(x) := \int_{\mathcal{C}(0) \cap \mathcal{C}(x); \varepsilon < t < \infty} \frac{dy dt}{t^2}, \quad \text{for } \varepsilon > 0. \tag{3.5}$$

Proof of Corollary 2.2 From Theorem 25 in [33] we obtain that Corollary 2.2 follows from (3.3), provided we show that $\{\xi^H\}_{H \in (0,1)}$ are uniformly integrable.

First note that all ξ^H , $0 < H < 1$, can be constructed on the same probability space, as a convolution of the same Brownian motion with different kernels which depend on H (see [27]). From (3.2) and (3.4) we have for $\delta \leq s, t \leq 1$,

$$\begin{aligned} K_H(s, t) &\leq \frac{1}{2H} |1 - |t - s|^{2H}| + \frac{1}{2H} |s^{2H} - 1| + \frac{1}{2H} |t^{2H} - 1| + (t - s)^{2H} + C \\ &\leq \frac{1}{2H} |1 - |t - s|^{2H}| + C(\delta). \end{aligned}$$

Using $1 - e^{-x} \leq x$ for $x > 0$ we get for all $\delta \leq s, t \leq 1$,

$$K_H(s, t) \leq -\log |t - s| + C(\delta).$$

On the other hand for $\delta \leq s, t \leq 1$ we have

$$K_H(s, t) \leq \frac{1}{2H} + C(\delta).$$

It follows that

$$\begin{aligned} K_H(s, t) &\leq \left(\frac{1}{2H}\right) \wedge \log \frac{1}{|t - s|} + C(\delta) \\ &= \log \frac{1}{|t - s| \vee e^{-\frac{1}{2H}}} + C(\delta). \end{aligned}$$

Let $\varepsilon \in (0, 1)$ be arbitrary small. Let \tilde{X}^ε be a centred Gaussian random field with the covariance kernel $\tilde{f}_\varepsilon + C(\delta)$, where \tilde{f}_ε is given in (3.5), i.e.

$$\tilde{K}_\varepsilon(t, s) := \mathbb{E}[\tilde{X}_t^\varepsilon \tilde{X}_s^\varepsilon] = \tilde{f}_\varepsilon(t - s) + C(\delta), \quad -\infty < t, s < \infty.$$

Since $\varepsilon \in (0, 1)$, we get from (3.5) for $0 < x \leq 1$,

$$\begin{aligned} \tilde{f}_\varepsilon(x) &= \int_{\mathcal{C}(0) \cap \mathcal{C}(x); \varepsilon < r < \infty} \frac{dy dr}{r^2} \\ &= \int_{x \vee \varepsilon}^\infty \frac{dr}{r^2} \int_{x - (r \wedge 1)/2}^{(r \wedge 1)/2} dy \\ &= -\log(x \vee \varepsilon) - \frac{x}{x \vee \varepsilon} + 1. \end{aligned}$$

Note that \tilde{f}_ε is an even function and therefore,

$$\tilde{K}_\varepsilon(t, s) \geq \log \frac{1}{|t - s| \vee \varepsilon} + C(\delta), \quad 0 < t, s \leq 1.$$

We define the following of measure

$$\tilde{\xi}_\gamma^\varepsilon(dt) = e^{\gamma\tilde{X}_t^\varepsilon - \frac{\gamma^2}{2}\mathbb{E}[(\tilde{X}_t^\varepsilon)^2]} dt, \quad \delta \leq t \leq 1.$$

Recall that $\gamma^2 < 2$ by our assumption. Then from the proof of Proposition 3.5 in [32] it follows that there exists $p = p(\gamma) > 1$, such that

$$\sup_{\varepsilon > 0} \mathbb{E}[(\tilde{\xi}_\gamma^\varepsilon([0, 1]))^p] < \infty.$$

By choosing $H = H(\varepsilon) = (-2 \log \varepsilon)^{-1}$, we note that $K_H(s, t) \leq \tilde{K}_\varepsilon(s, t)$, and from the comparison principal with $F(x) = x^p$ (see Corollary A.2 in [32]), we get

$$\sup_{H \in (0, 1)} \mathbb{E}[\xi_\gamma^H([\delta, 1])^p] < \infty.$$

We therefore conclude that $\{\xi^H\}_{H \in (0, 1)}$ are uniformly integrable. □

References

- [1] E. Bacry, J. Delour, and J.F. Muzy. Multifractal random walk. *Physical Review E*, 64(2):026103, 2001.
- [2] E. Bacry and J.F. Muzy. Log-infinitely divisible multifractal processes. *Communications in Mathematical Physics*, 236(3):449–475, Jun 2003. MR-2021198
- [3] J. Barral and B.B. Mandelbrot. Multifractal products of cylindrical pulses. *Probability Theory and Related Fields*, 124(3):409–430, Nov 2002. MR-1939653
- [4] C. Bayer, P.K. Friz, P. Gassiat, J. Martin, and B. Stemper. A regularity structure for rough volatility. *arXiv preprint arXiv:1710.07481*, 2017.
- [5] C. Bayer, P.K. Friz, and J. Gatheral. Pricing under rough volatility. *Quantitative Finance*, 16(6):887–904, 2016. MR-3494612
- [6] C. Bayer, P.K. Friz, A. Gulisashvili, B. Horvath, and B. Stemper. Short-time near-the-money skew in rough fractional volatility models. *arXiv preprint arXiv:1703.05132*, 2017.
- [7] M. Bennedsen, A. Lunde, and M.S. Pakkanen. Decoupling the short-and long-term behavior of stochastic volatility. *arXiv preprint arXiv:1610.00332*, 2016.
- [8] M. Bennedsen, A. Lunde, and M.S. Pakkanen. Hybrid scheme for Brownian semistationary processes. *Finance and Stochastics*, 21(4):931–965, 2017. MR-3723378
- [9] L. E. Calvet and A.J. Fisher. How to forecast long-run volatility: Regime switching and the estimation of multifractal processes. *Journal of Financial Econometrics*, 2(1):49–83, 2004.
- [10] L. Chevillard, R. Robert, and V. Vargas. A stochastic representation of the local structure of turbulence. *EPL (Europhysics Letters)*, 89(5):54002, 2010.
- [11] F. Comte and E. Renault. Long memory in continuous-time stochastic volatility models. *Mathematical Finance*, 8(4):291–323, 1998. MR-1645101
- [12] B. Duplantier, R. Rhodes, S. Sheffield, and V. Vargas. Log-correlated Gaussian fields: an overview. *arXiv preprint arXiv:1407.5605*, 2014.
- [13] O. El Euch and M. Rosenbaum. The characteristic function of rough Heston models. *Mathematical Finance*, to appear, 2016.
- [14] M. Forde and H. Zhang. Asymptotics for rough stochastic volatility and Lévy models. *preprint available at <http://www.mth.kcl.ac.uk/fordem>*, 2015.
- [15] U. Frisch and G. Parisi. On the singularity structure of fully developed turbulence. *Turbulence and Predictability in Geophysical Fluid Dynamics and Climate Dynamics*, pages 84–88, 1985. MR-1024082
- [16] M. Fukasawa. Short-time at-the-money skew and rough fractional volatility. *Quantitative Finance*, 17(2):189–198, 2017. MR-3592946

- [17] Y. V. Fyodorov and J.P. Bouchaud. Freezing and extreme-value statistics in a random energy model with logarithmically correlated potential. *Journal of Physics A: Mathematical and Theoretical*, 41(37):372001, 2008. MR-2430565
- [18] Y. V. Fyodorov, P. Le Doussal, and A. Rosso. Freezing transition in decaying burgers turbulence and random matrix dualities. *EPL (Europhysics Letters)*, 90(6):60004, 2010.
- [19] Y. V. Fyodorov, B. A. Khoruzhenko, and N. J. Simm. Fractional Brownian motion with hurst index $H = 0$ and the Gaussian unitary ensemble. *The Annals of Probability*, 44(4):2980–3031, 2016. MR-3531684
- [20] J. Gatheral, T. Jaisson, and M. Rosenbaum. Volatility is rough. *arXiv preprint arXiv:1410.3394*, 2014. MR-3805308
- [21] A. Jacquier, M.S. Pakkanen, and H. Stone. Pathwise large deviations for the rough Bergomi model. *arXiv preprint arXiv:1706.05291*, 2017.
- [22] J.P. Kahane. *Sur le chaos multiplicatif*. Prépublications mathématiques d’Orsay. Département de mathématique, 1985. MR-0829798
- [23] W.E. Leland, M.S. Taqqu, W. Willinger, and D.V. Wilson. On the self-similar nature of ethernet traffic (extended version). *IEEE/ACM Transactions on networking*, 2(1):1–15, 1994.
- [24] A. Lodhia, S. Sheffield, X. Sun, and S. S. Watson. Fractional Gaussian fields: a survey. *arXiv preprint arXiv:1407.5598v2*, 2016. MR-3466837
- [25] T. Madaule, R. Rhodes, and V. Vargas. Glassy phase and freezing of log-correlated Gaussian potentials. *The Annals of Applied Probability*, 26(2):643–690, 2016. MR-3476621
- [26] B.B. Mandelbrot, A.J. Fisher, and L. Calvet. A multifractal model of asset returns. Cowles Foundation Discussion Papers 1164, Cowles Foundation for Research in Economics, Yale University, September 1997.
- [27] B.B. Mandelbrot, J.W. Van Ness. Fractional Brownian Motions, Fractional Noises and Applications. *SIAM Rev.*, 10(4):422–437, 1968. MR-0242239
- [28] T. Mikosch, S. Resnick, H. Rootzén, and A. Stegeman. Is network traffic approximated by stable Lévy motion or fractional Brownian motion? *The Annals of Applied Probability*, 12(1):23–68, 2002. MR-1890056
- [29] F.J. Molz, H.H. Liu, and J. Szulga. Fractional Brownian motion and fractional Gaussian noise in subsurface hydrology: A review, presentation of fundamental properties, and extensions. *Water Resources Research*, 33(10):2273–2286, 1997.
- [30] R. Rhodes and V. Vargas. Gaussian multiplicative chaos and applications: a review. *Probability Surveys*, 11:315–392, 2014. MR-3274356
- [31] R. Rhodes and V. Vargas. Lecture notes on Gaussian multiplicative chaos and Liouville quantum gravity. *arXiv preprint arXiv:1602.07323*, 2016. MR-3274356
- [32] R. Robert and V. Vargas. Gaussian multiplicative chaos revisited. *The Annals of Probability*, 38(2):605–631, 2010. MR-2642887
- [33] A. Shamov. On Gaussian multiplicative chaos. *Journal of Functional Analysis*, 270(9):3224–3261, 2016. MR-3475456

Acknowledgments. We are very grateful to Vincent Vargas, an anonymous referee and the associate editor whose numerous useful comments enabled us to significantly improve this paper.