

# Nonparametric and high-dimensional functional graphical models\*

Eftychia Solea and Holger Dette

*CREST and ENSAI, Rennes, France,  
Ruhr-Universität Bochum, Germany*

*e-mail: [eftychia.solea@ensai.fr](mailto:eftychia.solea@ensai.fr); [holger.dette@rub.de](mailto:holger.dette@rub.de)*

**Abstract:** We consider the problem of constructing nonparametric undirected graphical models for high-dimensional functional data. Most existing statistical methods in this context assume either a Gaussian distribution on the vertices or linear conditional means. In this article, we provide a more flexible model which relaxes the linearity assumption by replacing it by an arbitrary additive form. The use of functional principal components offers an estimation strategy that uses a group lasso penalty to estimate the relevant edges of the graph. We establish statistical guarantees for the resulting estimators, which can be used to prove consistency if the dimension and the number of functional principal components diverge to infinity with the sample size. We also investigate the empirical performance of our method through simulation studies and a real data application

**MSC2020 subject classifications:** Primary 60K35, 60K35; secondary 60K35.

**Keywords and phrases:** Additive models, conditional independence, functional data, lasso, undirected graphical models.

Received October 2021.

## Contents

1	Introduction . . . . .	6176
2	Additive functional graphical models . . . . .	6178
3	Estimation and computation . . . . .	6181
4	Statistical guarantees . . . . .	6184
5	Finite sample properties . . . . .	6188
5.1	Simulated data . . . . .	6188
5.1.1	Scenario 1: nonlinear models . . . . .	6190
5.1.2	Scenario 2: linear model . . . . .	6190
6	Real data application . . . . .	6191
7	Conclusions . . . . .	6192
A	Auxiliary results . . . . .	6193
A.1	Rates of convergence for sample design matrices . . . . .	6201
A.2	Proof of Theorem 4.1 . . . . .	6205

---

arXiv: [2103.10568](https://arxiv.org/abs/2103.10568)

\*This work has been supported in part by the Collaborative Research Center “Statistical modeling of nonlinear dynamic processes” (SFB 823, Teilprojekt A1,C1) of the German Research Foundation (DFG).

A.3 Proof of Proposition A.2 . . . . .	6205
A.4 Proof of Proposition A.2 . . . . .	6227
Acknowledgments . . . . .	6228
References . . . . .	6228

## 1. Introduction

In recent years, there has been a large amount of work on estimating undirected graphical models that describe the conditional dependencies among the components of a  $p$ -dimensional random vector  $X = (X^1, \dots, X^p)^\top$ . Mathematically, let  $\mathbf{V} = \{1, \dots, p\}$ , and  $\mathbf{E}$  denote a subset of  $\{(i, j) \in \mathbf{V} \times \mathbf{V} : i \neq j\}$ , which satisfies  $(i, j) \in \mathbf{E}$  if and only if  $(j, i) \in \mathbf{E}$ . The pair  $\mathbf{G} = (\mathbf{V}, \mathbf{E})$  constitutes an undirected graph, with  $\mathbf{V}$  representing the set of vertices and  $\mathbf{E}$  the set of edges. The vector  $X$  follows a graphical model if

$$(i, j) \notin \mathbf{E} \quad \Leftrightarrow \quad X^i \perp\!\!\!\perp X^j | X^{-\{i, j\}}, \quad (1.1)$$

where  $X^{-\{i, j\}}$  represents the vector  $X$  with its  $i$ th and  $j$ th components removed, and for random elements  $A$ ,  $B$ , and  $C$ ,  $A \perp\!\!\!\perp B | C$  means that  $A$  and  $B$  are conditionally independent given  $C$ . The goal is to estimate the edge set  $\mathbf{E}$  based on a random sample from  $X$ .

However, many recent applications, particularly medical applications, involve multivariate functional data, such as electroencephalogram (EEG) and functional magnetic resonance imaging (fMRI) data, where each sampling unit is modelled as a realization of a  $p$ -dimensional stochastic process  $X^1(t), \dots, X^p(t)$ , where  $t \in T$  and  $T$  is a closed subset of  $\mathbb{R}$ . The central goal is to depict the conditional dependence structure among these  $p$  random functions by *functional graphical models (FGM)*, which have become very popular in recent years. In this paper, we are interested in estimating a nonparametric and high-dimensional undirected graphical model for multivariate functional data.

Our motivation is brain network connectivity based on EEG in neuroscience research. EEG measures brain activities by placing electrodes on various locations on the subject's scalp and measuring voltage values across time. This results in a vector of interdependent random functions. One of the interests in network connectivity analysis is to characterize the inter-dependence of the random functions by an undirected graph, where the nodes corresponds to the electrodes on the subject's scalp and the edges represent conditional dependencies among the electrodes.

In contrast to the finite dimensional case, less literature can be found on graphical models for multivariate functional data. [26] proposed the Functional Gaussian Graphical Model (FGGM) assuming that  $X$  is a multivariate Gaussian random process. Roughly speaking, they used a truncated Karhunen-Loève expansion, say of order  $m_n$ , to reduce the infinite dimensional problem to a  $pm_n$ -dimensional problem for the principal component scores. The conditional independencies of the graph define a block sparsity structure, such that the zero blocks of the precision matrix of the scores can be used to identify the

edge set using a group lasso penalty. They called this method functional glasso, or simply fglasso, and the authors showed that, when  $m_n$  approaches infinity, consistent estimation of the edge set is possible. The FGGM is the extension of the Gaussian graphical model of [38] to the functional setting. [40] proposes a FGGM based on a new notion of separability for the covariance operator of multivariate functional data, termed partial separability. [43] considered a neighbourhood selection approach to estimate the FGGM, which is an extension of the popular work of [21] to the functional setting. [45] proposed a Bayesian framework for the analysis of FGGM, [42] developed a method that estimates the difference between two functional graphical models and [27] considered a dynamic functional graphical model, where the dependency structure among the Gaussian random functions is allowed to change over time. Finally, to relax the Gaussian assumption [16] replace conditional independence with the concept of functional additive conditional independence and [31] proposed a copula transformation on the principal scores, which is an extension of the Gaussian copula model of [18, 17] and [36] from the multivariate setting to the functional setting.

In this paper, we introduce an alternative approach to relax the Gaussian assumption in the functional graphical model. Our research is motivated by the fact that in many applications the relation between the functional principal scores is rarely linear as implied by the assumption of the FGGM. To illustrate this observation, we consider an EEG dataset that consists of two groups of subjects: 77 subjects in the alcoholic group, and 45 in the control group [41, 13]. For each subject, an EEG activity was recorded at 256 time points over a one second time interval using 64 electrodes placed on the subject's scalp. The goal is to construct a FGM to characterize brain network connectivity for the two groups of subjects based on the functional data collected by the electrodes. In Figure 1, we display the pairwise scatterplots between channels using the first two principal components for the random functions of the two groups. Clearly, this figure indicates that the conditional relationships among the scores corresponding to different vertices of the graph are nonlinear. Therefore, the Gaussian assumption of the FGGM is difficult to justify for the analysis of this type of data.

As an alternative, we propose a new nonparametric FGM that allows the conditional relationships among the principal scores to take an additive structure. Our approach uses the traditional probabilistic concept of conditional independence, and then applies the additive structure to the scores of the Karhunen-Loève expansion of each random function. We approximate each nonparametric additive component by a linear combination of B-splines basis functions. This enables us to estimate the edge set of the graph by imposing the group lasso penalty on a matrix formed by the coefficients in the spline approximation. We derive statistical guarantees for the resulting estimates, which can be used to prove consistency if the dimension  $p$  and the number of scores diverge to infinity with the sample size. This provides a useful methodology for general nonparametric analysis of high-dimensional functional graphical models.

The remainder of the article is organized as follows. Section 2 describes the




FIG 1. Pairwise scatterplots for the alcoholic group between channels F7 and FP2 (left) and for the control group between channels O1 and X (right) with locally-weighted polynomial regression (blue line).

methodology and proposes the nonparametric functional graphical model. Section 3 presents the estimation procedure. In Section 4 we study the theoretical properties of the resulting estimator. In Section 5 we conduct simulation studies to evaluate the finite sample properties of the proposed methodology, and in Section 6 we apply the new model to the motivating EGG dataset. We conclude with some final remarks in Section 7, while all proofs of the theoretical results are deferred to the Appendix.

## 2. Additive functional graphical models

We first provide a formal definition of an *additive function-on-function regression model* which will be used to define the functional graphical model considered in this paper. We begin by introducing some basic concepts from functional data analysis.

Throughout this paper,  $\mathcal{L}^2([0, 1])$  denotes the space of all square-integrable functions defined on the interval  $[0, 1] \subset \mathbb{R}$ . We denote by  $\langle f, g \rangle = \int_{[0,1]} f(t)g(t)dt$  the common inner product in  $\mathcal{L}^2([0, 1])$  and by  $\|f\| = \langle f, f \rangle^{1/2}$  the corresponding norm. Let  $X = (X^1, \dots, X^p)$  denote a  $p$ -dimensional random element with mean 0 whose  $i$ th component  $X^i$  is an element of  $\mathcal{L}^2([0, 1])$  such that  $E\|X^i\|^2 < \infty$ . For each  $X^i$ , we define the corresponding covariance operator

$$\Sigma_{X^i X^i}(f)(t) = \int_0^1 f(s)\sigma_{X^i X^i}(s, t)ds, \quad f \in \mathcal{L}^2([0, 1]), \quad (2.1)$$

where  $\sigma_{X^i X^i}(s, t) = \text{cov}(X^i(s), X^i(t)) = E(X^i(s)X^i(t))$  is the covariance function of the random element  $X^i$ . The operator  $\Sigma_{X^i X^i}$  is a compact Hilbert-Schmidt operator [see, for example, 11], and there exists a spectral decomposi-

tion of the covariance function of the form

$$\sigma_{X^i X^i}(s, t) = \sum_{r=1}^{\infty} \lambda_r^i \phi_r^i(s) \phi_r^i(t), \tag{2.2}$$

where  $\lambda_1^i \geq \lambda_2^i \geq \dots \geq 0$  are the eigenvalues and  $\{\phi_k^i\}_{k \in \mathbb{N}}$  are orthonormal eigenfunctions satisfying

$$\int_0^1 \sigma_{X^i X^i}(s, t) \phi_r^i(s) ds = \lambda_r^i \phi_r^i(t).$$

Consequently, each  $X^i \in \mathcal{L}^2([0, 1])$  can be represented by its Karhunen-Loève expansion

$$X^i = \sum_{r=1}^{\infty} \sqrt{\lambda_r^i} \xi_r^i \phi_r^i \quad i = 1, \dots, p, \tag{2.3}$$

where the random variables  $\xi_r^i = \langle X^i, \phi_r^i \rangle / \sqrt{\lambda_r^i}$  are called the *functional principal component scores* and satisfy  $E(\xi_r^i) = 0$ ,  $\text{var}(\xi_r^i) = 1$ ,  $E(\xi_q^i \xi_r^i) = 0$  for  $q \neq r$  for  $i = 1, \dots, p$ .

We next give our formal definition of the functional graphical model. Let  $G = (V, E)$  be an undirected graph, where  $V$  denotes the finite set  $\{1, \dots, p\}$ , and  $E$  denotes a subset of  $\{(i, j) \in V \times V : i \neq j\}$ , which satisfies  $(i, j) \in E$  if and only if  $(j, i) \in E$ .

**Definition 2.1.** A vector of random functions  $X = (X^1, \dots, X^p) \in \mathcal{L}^2([0, 1]) \times \dots \times \mathcal{L}^2([0, 1])$  is said to follow a functional graphical model with respect to an undirected graph  $G = (V, E)$  if and only if

$$X^i \perp\!\!\!\perp X^j | X^{-\{i,j\}}, \quad \forall (i, j) \notin E.$$

**Example 2.1.** [26] assumed that  $X = (X^1, \dots, X^p)$  is a  $p$ -dimensional Gaussian process on  $\mathcal{L}^2([0, 1]) \times \dots \times \mathcal{L}^2([0, 1])$  and define a Functional Gaussian Graphical model (FGGM) by the condition

$$(i, j) \notin E \quad \Leftrightarrow \quad \text{cov}[X^i(s), X^j(t) | X^{-\{i,j\}}] = 0 \quad \forall s, t \in [0, 1]. \tag{2.4}$$

Next, they proposed to approximate each  $X^i$  by the first  $m_n$  coefficients from the Karhunen-Loève expansion (2.3). Thus, for each  $X^i$ , one obtains a  $pm_n$ -dimensional Gaussian random vector  $\xi^\top = ((\xi^1)^\top, \dots, (\xi^p)^\top)$  of scores, where  $\xi^i = (\xi_1^i, \dots, \xi_{m_n}^i)^\top$  is the vector of the first  $m_n$  functional principal component scores in the Karhunen-Loève expansion (2.3) of each  $X^i$ . Based on this finite  $m_n$ -dimensional representation, the FGGM can be represented as a conditional multivariate linear regression model with respect to the scores. Indeed, each  $\xi_q^i$  can be expressed as

$$\xi_q^i = \sum_{j \neq i}^p \sum_{r=1}^{m_n} B_{qr}^{ij} \xi_r^j + \epsilon_q^i, \quad i \in V, q = 1, \dots, m_n, \tag{2.5}$$

such that  $(\epsilon_q^i)_{1 \leq q \leq m_n}$  is uncorrelated with  $(\xi_r^j)_{1 \leq r \leq m_n}$ ,  $i \neq j$  if and only if

$$B_n^{ij} = (B_{qr}^{ij})_{1 \leq q, r \leq m_n} = -(\Theta_n^{ii})^{-1} \Theta_n^{ij}, \quad (i, j) \in \mathbb{V} \times \mathbb{V}, i \neq j,$$

where  $\Theta_n^{ij} \in \mathbb{R}^{m_n \times m_n}$  is the  $(i, j)$ th element of the block precision matrix  $\Theta_n = (\Theta_n^{ij})_{1 \leq i, j \leq p}$  of the  $pm_n$ -dimensional vector  $\xi$ . Hence, under the Gaussian assumption the conditional relationships between nodes  $i$  and  $j$  are linear, and the network structure of the FGGM can also be recovered by the sparsity structure of the regression coefficient matrix  $B_n^{ij}$ . They used group-lasso penalized maximum likelihood estimation to address the blockwise sparsity of  $\Theta_n$ , and showed that  $\Theta_n$  is a consistent estimate of the set  $\mathbf{E}$ , when  $p$  and  $m_n$  approach infinity with increasing sample size.

We use a generalization of the representation (2.5) to give a formal definition of the additive function-on-function model for multivariate functional data.

**Definition 2.2.** Consider a vector of random functions  $X = (X^1, \dots, X^p) \in \mathcal{L}^2([0, 1]) \times \dots \times \mathcal{L}^2([0, 1])$  and suppose that each  $X^i$  has a Karhunen-Loève expansion of the form (2.3). The vector  $X$  follows the function-on-function additive model if for each pair  $(i, j) \in \mathbb{V} \times \mathbb{V}$  there exists a sequence of smooth functions  $f_{qr}^{ij} = \{f_{qr}^{ij} : q, r \in \mathbb{N}\}$  defined on  $\mathbb{R}$  with  $E[f_{qr}^{ij}(\xi_r^j)] = 0$ ,  $q, r \in \mathbb{N}$ , such that

$$E[\xi_q^i | \{\xi_r^j, j \neq i\}] = \sum_{j \neq i} \sum_{r=1}^{\infty} f_{qr}^{ij}(\xi_r^j). \quad (2.6)$$

Similar to the functional additive regression model of [9], model (2.6) relaxes the linearity assumption in FGGM by imposing an additive structure on the scores in the Karhunen-Loève expansion, giving rise to a more flexible model than the FGGM. By definition, the scores  $\xi_r^i$  are uncorrelated, but we also require them to be independent in the following discussion as also postulated in [9]. Furthermore, we assume that they take values in a closed and bounded interval  $[-1, 1]$ . For example, this can be achieved by taking a monotone transformation  $\Psi : \mathbb{R} \rightarrow [-1, 1]$ , such as  $\Psi(x) = -1 + \frac{2 \arctan(x) + 0.5\pi}{\pi}$ . [see 46, 34].

We now define a new nonparametric functional graphical model which we call the Additive Functional Graphical Model (AFGM).

**Definition 2.3.** Suppose  $X = (X^1, \dots, X^p) \in \mathcal{L}^2([0, 1]) \times \dots \times \mathcal{L}^2([0, 1])$  is associated with a functional graphical model  $\mathbf{G} = (\mathbb{V}, \mathbf{E})$ . If  $X$  is additionally a function-on-function additive model of the form (2.6), then we say that  $X$  follows an additive functional graphical model, and write this statement as  $X \sim \text{AFGM}(\mathbf{G})$ .

The definition implies that the independence structure of  $X$  can be recovered by the sparse structure of the additive components  $f_{qr}^{ij}$  in the representation (2.6). In most cases, the sequence of eigenvalues decreases fast, and hence without much loss of generality, we will assume a finite  $m_n$ -dimensional model,

where  $(m_n)_{n \in \mathbb{N}}$  is a sequence of positive real numbers converging to infinity with increasing sample size. Specifically,

$$\xi_q^i = \sum_{j \neq i} \sum_{r=1}^{m_n} f_{qr}^{ij}(\xi_r^j) + \epsilon_q^i, \quad i = 1, \dots, p, q = 1, \dots, m_n, \quad (2.7)$$

such that the error terms  $\epsilon_q^i = \xi_q^i - \sum_{j \neq i} \sum_{r=1}^{m_n} f_{qr}^{ij}(\xi_r^j)$  have 0 mean, they are mutually independent, and independent of the predictors  $\xi_r^j, j \neq i$  if and only if conditional independence among  $X^i$  can be recovered from the additive functions  $f_{qr}^{ij}$ , since in this case the conditional mean of each  $\xi_q^i$  in the graph only depends on the values of its neighbours. In the Gaussian model,  $f_{qr}^{ij}(\xi_r^j) = B_{qr}^{ij} \xi_r^j$ , and model (2.7) reduces to the FGGM (2.5).

Thus, our goal is to estimate the edge set

$$E_n = \{(i, j) \in V \times V : i \neq j, f_{qr}^{ij} \neq 0 \text{ for some } q, r = 1, \dots, m_n\}. \quad (2.8)$$

Note that we aim to recover the edge set when each  $X^i$  is exactly a finite sum of  $m_n$  terms rather than an infinite sum. This allows us to work with finite dimensional graphs that is  $E = E_n$ . Our theoretical results in Section 4 show that the edge set can be identified with probability converging to 1 as  $m_n \rightarrow \infty$ ,  $p \rightarrow \infty$  and  $n \rightarrow \infty$ .

**Remark 2.1.**

- (a) Note that it is not necessary to fix the sign of the eigenfunctions in the definition of scores  $\xi_r^i = \langle X^i, \phi_r^i \rangle / \sqrt{\lambda_r}$  used in the representation (2.6) or (2.7), because a sign change can always be compensated by choosing the function  $f_{qr}^{ij}(-x)$  instead of  $f_{qr}^{ij}(x)$ .
- (b) Model (2.7) can be regarded as the nonparametric and additive version of the FGGM (2.5), and the generalization of the model of [33] to the functional setting, where they propose a semi-parametric method which allows the conditional means of the random variables to take on an arbitrary additive structure.

**3. Estimation and computation**

In this section, we develop an estimation procedure for fully observed functional data to estimate the scores  $\xi^i$  for each  $X^i$ , which is used afterwards for the estimation of the edge set  $E_n$ .

To be precise, let  $X_1, \dots, X_n$  be an independent sample from  $X$ , such that for each  $u = 1, \dots, n$ ,  $X_u = (X_u^1, \dots, X_u^p)$  is a vector in  $\mathcal{L}^2([0, 1]) \times \dots \times \mathcal{L}^2([0, 1])$ . Then, for each  $i = 1, \dots, p$ , the covariance operator  $\Sigma_{X^i X^i}$  can be estimated by

$$\hat{\Sigma}_{X^i X^i}(f)(t) = \int_{[0,1]} f(s) \hat{\sigma}_{X^i X^i}(s, t) ds, \quad f \in \mathcal{L}^2([0, 1]),$$

where

$$\hat{\sigma}_{X^i X^i}(s, t) = \frac{1}{n} \sum_{u=1}^n X_u^i(s) X_u^i(t),$$

is the common estimator of the covariance function (note that the  $X^i$  are centered). Let  $\hat{\lambda}_r^i$  and  $\hat{\phi}_r^i$  be the sample eigenvalues and eigenfunctions obtained by solving the equation

$$\int_0^1 \hat{\sigma}_{X^i X^i}(s, t) \phi_r^i(s) ds = \lambda_r^i \phi_r^i(t), \quad r = 1, \dots, m_n,$$

subject to the constraints  $\langle \phi_q^i, \phi_r^i \rangle = 0$ , for  $q \neq r, q, r = 1, \dots, m_n$  and  $\|\phi_r^i\| = 1$ . Then, the estimated scores  $\hat{\xi}_{ur}^i$  are given by

$$\hat{\xi}_{ur}^i = (\hat{\lambda}_r^i)^{-1/2} \langle X_u^i, \hat{\phi}_r^i \rangle, \quad u = 1, \dots, n, r = 1, \dots, m_n, i \in \mathbb{V}.$$

For each  $i \in \mathbb{V}$ , let  $\hat{\xi}_u^i = (\hat{\xi}_{u1}^i, \dots, \hat{\xi}_{um_n}^i)^\top$  be the  $m_n$ -dimensional vector of the estimated scaled scores corresponding to the observation  $X_u$ ,  $u = 1, \dots, n$ . Following [12] and [8], we use B-spline functions to approximate the additive components  $f_{qr}^{ij}$  in model (2.7). To be precise, let  $-1 = \tau_0 < \tau_1 < \dots < \tau_{L_n} < \tau_{L_n+1} = 1$  be an equidistant partition of the interval  $[-1, 1]$  into  $L_n + 1$  subintervals  $I_b = [\tau_b, \tau_{b+1}), b = 0, \dots, L_n - 1$ , and  $I_{L_n} = [\tau_{L_n}, \tau_{L_n+1}]$ .

For the number of knots we make the following assumption. Define  $S_{\ell L_n}$  as the space of polynomial splines of degree  $\ell \geq 1$  consisting of functions  $s$  satisfying: (a) the restriction of  $s$  to the interval  $I_b$  is a polynomial of degree  $\ell$  for  $1 \leq b \leq L_n$ ; (b) for  $\ell \geq 2$  and  $1 \leq \ell' \leq \ell - 2$ ,  $s$  is a  $\ell'$  times continuously differentiable on the interval  $[-1, 1]$ . Then, there exists a basis of normalized B-splines functions  $(h_k)_{1 \leq k \leq k_n}$  for the space  $S_{\ell L_n}$ , where  $k_n = L_n + \ell + 1$ , such that every function  $s \in S_{\ell L_n}$  can be represented as

$$s(x) = \sum_{k=1}^{k_n} \beta_k h_k(x)$$

(see [30]). Under some smoothness conditions, the additive functions  $f_{qr}^{ij}$  can be represented by linear combinations of B-splines functions

$$f_{qr}^{ij}(x) \approx \tilde{f}_{qr}^{ij}(x) = \sum_{k=1}^{k_n} h_k(x) \beta_{qrk}^{ij}, \quad q, r = 1, 2, \dots, \quad (3.1)$$

where the sequence of positive real numbers,  $(k_n)_{n \in \mathbb{N}}$ , diverges to infinity as  $n \rightarrow \infty$  (note that this can always be achieved by increasing the number of knots in the partition). Hence, the corresponding function  $f_{qr}^{ij}$  will be zero approximately if and only if  $\|\beta_{qr}^{ij}\|_2^2 = 0$ , where  $\|\cdot\|_2$  denotes the Euclidean norm of the  $k_n$ -dimensional vector  $\beta_{qr}^{ij} = (\beta_{qr1}^{ij}, \dots, \beta_{qrk_n}^{ij})^\top$ ,  $q, r = 1, \dots, m_n$ . Thus, to encourage sparsity we propose to minimize the criterion

$$PL_i(\beta, \hat{\xi}) = \frac{1}{2n} \sum_{q=1}^{m_n} \sum_{u=1}^n \left( \hat{\xi}_{uq}^i - \sum_{j \neq i}^p \sum_{r=1}^{m_n} h^\top(\hat{\xi}_{ur}^j) \beta_{qr}^{ij} \right)^2 + \lambda_n \sum_{j \neq i}^p \left\{ \sum_{q=1}^{m_n} \sum_{r=1}^{m_n} \|\beta_{qr}^{ij}\|_2^2 \right\}^{1/2},$$



subject to the constraint

$$\sum_{u=1}^n h^\top(\hat{\xi}_{ur}^j) \beta_{qr}^{ij} = 0, \quad q, r = 1, \dots, m_n, j \in \mathcal{V}, \quad (3.2)$$

where  $h^\top(x) = (h_1(x), \dots, h_{k_n}(x))$  is the  $k_n$ -dimensional vector of the B-splines basis functions and  $\lambda_n \geq 0$  is a tuning parameter. Essentially, this is a group lasso penalty that enforces all regression coefficients  $\beta_{qr1}^{ij}, \dots, \beta_{qrk_n}^{ij}$  to either be all 0 or all nonzero. Note that the centering constraint (3.2) accounts for the fact that the function  $f_{qr}^{ij}$  in model (2.7) satisfies  $E(f_{qr}^{ij}(\xi_r^j)) = 0$  for  $q, r = 1, \dots, m_n$ .

This problem can be converted to an unconstrained optimisation problem by centering the basis functions. More precisely, defining

$$\tilde{h}_{nk}(\hat{\xi}_{ur}^j) = h_k(\hat{\xi}_{ur}^j) - \frac{1}{n} \sum_{u=1}^n h_k(\hat{\xi}_{ur}^j), \quad k = 1, \dots, k_n, r = 1, \dots, m_n, j \in \mathcal{V}, \quad (3.3)$$

we consider the unconstrained optimization problem

$$\widehat{PL}_i(\beta, \hat{\xi}) = \frac{1}{2n} \sum_{q=1}^{m_n} \sum_{u=1}^n \left( \hat{\xi}_{uq}^i - \sum_{j \neq i}^p \sum_{r=1}^{m_n} \tilde{h}_n^\top(\hat{\xi}_{ur}^j) \beta_{qr}^{ij} \right)^2 + \lambda_n \sum_{j \neq i}^p \left\{ \sum_{q=1}^{m_n} \sum_{r=1}^{m_n} \|\beta_{qr}^{ij}\|_2^2 \right\}^{1/2}, \quad (3.4)$$

where

$$\tilde{h}_n(\hat{\xi}_{ur}^j) = (\tilde{h}_{n1}(\hat{\xi}_{ur}^j), \dots, \tilde{h}_{nk_n}(\hat{\xi}_{ur}^j))^\top, \quad (3.5)$$

is the  $k_n$ -dimensional vector of the centered B-splines evaluated at the estimated scores.

Now let  $\hat{\xi}^i = (\hat{\xi}_{ur}^i)_{1 \leq u \leq n, 1 \leq r \leq m_n}$  be the  $n \times m_n$  matrix of the estimated scores, and define

$$\tilde{\mathbf{H}}_n^\top(\hat{\xi}^{-i}) = (\tilde{H}_n(\hat{\xi}^1), \dots, \tilde{H}_n(\hat{\xi}^{i-1}), \tilde{H}_n(\hat{\xi}^{i+1}), \dots, \tilde{H}_n(\hat{\xi}^p)) \in \mathbb{R}^{n \times (p-1)k_n m_n} \quad (3.6)$$

as the vector of matrices  $\tilde{H}_n(\hat{\xi}^j) = (\tilde{h}_n^\top(\hat{\xi}_{ur}^j))_{1 \leq u \leq n, 1 \leq r \leq m_n} \in \mathbb{R}^{n \times k_n m_n}$ . Similarly, let

$$B^i = (B^{ij}, j \in \mathcal{V} \setminus \{i\}) \in \mathbb{R}^{(p-1)k_n m_n \times m_n},$$

be the vector of matrices  $B^{ij} = (\beta_{qr}^{ij})_{1 \leq q \leq m_n, 1 \leq r \leq m_n} \in \mathbb{R}^{k_n m_n \times m_n}, j \neq i$ . Then, following some algebraic manipulations, the objective function in (3.4) can be rewritten as

$$\widehat{PL}_i(B, \hat{\xi}) = \frac{1}{2n} \|\hat{\xi}^i - \tilde{\mathbf{H}}_n^\top(\hat{\xi}^{-i}) B^i\|_F^2 + \lambda_n \sum_{j \neq i}^p \|B^{ij}\|_F, \quad (3.7)$$

where  $\|\cdot\|_F$  denotes the Frobenius norm. Finally, we define  $\hat{B}_n^i$  as the solution of

$$\hat{B}_n^i = \operatorname{argmin}\{\widehat{PL}_i(B, \hat{\xi}) : B \in \mathbb{R}^{(p-1)k_n m_n \times m_n}\},$$

and we propose to estimate the set  $E_n$  in (2.8) by

$$\hat{E}_n = \{(i, j) \in \mathcal{V} \times \mathcal{V} : i \neq j, \|\hat{B}_n^{ij}\|_{\mathbb{F}} \neq 0 \text{ and } \|\hat{B}_n^{ji}\|_{\mathbb{F}} \neq 0\}. \quad (3.8)$$

Following [21], we can also estimate the graph by

$$\hat{\hat{E}}_n = \{(i, j) \in \mathcal{V} \times \mathcal{V} : i \neq j, \|\hat{B}_n^{ij}\|_{\mathbb{F}} \neq 0 \text{ or } \|\hat{B}_n^{ji}\|_{\mathbb{F}} \neq 0\}. \quad (3.9)$$

As shown in the Corollary 4.1 in Section 4, the differences between these estimated graphs vanish asymptotically. See also [21] and [43]. We summarize the algorithm below.

- (a) Implement FPCA to obtain the estimated scores  $\hat{\xi}_{ur}^i$  of each observation  $X_u^i$  and then transform the scores into the range  $[-1, 1]$  using a monotone transformation. Choose  $m_n$  such that at least 90% of the total variation is explained.
- (b) For a given  $\lambda_n$  and for each  $i \in \mathcal{V}$ , solve the optimisation problem (3.7) using, for example, the Fast Iterative Shrinkage Thresholding Algorithm (See [4] for more details) to find a sparse estimate of  $B^i$ .
- (c) Declare that there is an edge between node  $i$  and node  $j$  if and only if  $\|\hat{B}_n^{ij}\|_{\mathbb{F}}^2$  and  $\|\hat{B}_n^{ji}\|_{\mathbb{F}}^2$  are not zero.

#### 4. Statistical guarantees

In this section we study the theoretical properties of the proposed estimator of the graph structure of the AFGM, where we allow the number of nodes  $p$  to diverge to infinity with increasing sample size. We also assume that the functions are fully observed. A particular technical challenge in deriving the asymptotic theory consists in the fact that the additive structure is applied to the unobserved variables  $\xi_{ur}^i$ , and the estimator  $\hat{B}_n^i$  obtained from minimizing (3.7) is based on the estimated scores. Thus, the error in these estimated coefficients must be taken into account for the analysis of the procedure.

We begin by introducing some notation. For any two positive sequences of real numbers  $(a_n)_{n \in \mathbb{N}}$  and  $(b_n)_{n \in \mathbb{N}}$ , we write  $a_n \lesssim b_n$  if  $a_n \leq K_1 b_n$  for some constant  $0 < K_1 < \infty$  which does not depend on  $n$ . We use the notation  $a_n \asymp b_n$  representing the property  $A \leq \inf_n |a_n/b_n| \leq \sup_n |a_n/b_n| \leq B$ , for positive constants  $A$  and  $B$ . Moreover, given a matrix  $A = (a_{ij})_{1 \leq i \leq M_1, 1 \leq j \leq M_2} \in \mathbb{R}^{M_1 \times M_2}$ , we use  $\|A\|_{\mathbb{F}}$  for the Frobenius norm and  $\|A\|_2$  for the operator norm. Finally, for any two symmetric matrices  $A$  and  $B$ , we use the notation  $A \preceq B$  to denote the property that the matrix  $B - A$  is nonnegative definite.

We define the neighbourhood of each node  $i \in \mathcal{V}$  by

$$\mathcal{N}_n^i = \{j \in \mathcal{V} \setminus \{i\} : f_{qr}^{ij} \neq 0, \text{ for some } 1 \leq q, r \leq m_n\},$$

where  $(f_{qr}^{ij})_{1 \leq q, r \leq m_n}$  are the true population additive functions in the additive regression

$$\xi_{uq}^i = \sum_{j \in \mathcal{N}_n^i} \sum_{r=1}^{m_n} f_{qr}^{ij}(\xi_{ur}^j) + \epsilon_{uq}^i, \quad q = 1, \dots, m_n, u = 1, \dots, m_n, i \in \mathcal{V}, \quad (4.1)$$

where  $\epsilon_{uq}^i = \xi_{uq}^i - E[\xi_{uq}^i | \{\xi_{ur}^j, j \neq i; r = 1, \dots, m_n\}]$ . Thus, the best predictor of  $\xi_{uq}^i$  is an additive function of the scores in the set of neighbours  $N_n^i$  of the node  $i$  only. Then, the edge set  $E_n$  defined in (2.8) can be rewritten as

$$E_n = \{(i, j) \in V \times V : i \neq j, i \in N_n^j \text{ and } j \in N_n^i\}.$$

Let  $\beta_{qr}^{*ij} = (\beta_{qr1}^{*ij}, \dots, \beta_{qrk_n}^{*ij})^\top$  be the  $k_n$ -dimensional vector of the true spline coefficients in

$$\xi_{uq}^i = \sum_{j \in N_n^i} \sum_{r=1}^{m_n} h^\top(\xi_{ur}^j) \beta_{qr}^{*ij} + w_{uq}^i + \epsilon_{uq}^i, \quad q = 1, \dots, m_n, u = 1, \dots, n, i \in V,$$

where  $h^\top(\xi_{ur}^j) = (h_1(\xi_{ur}^j), \dots, h_{k_n}(\xi_{ur}^j)) \in \mathbb{R}^{k_n}$  is the vector of the B-splines evaluated at the unobserved scaled scores  $\xi_{ur}^j$  and  $w_{uq}^i$  is the truncation error

$$w_{uq}^i = \sum_{j \in N_n^i} \sum_{r=1}^{m_n} (f_{qr}^{ij}(\xi_{ur}^j) - h^\top(\xi_{ur}^j) \beta_{qr}^{*ij}), \quad q = 1, \dots, m_n, u = 1, \dots, n, i \in V.$$

In Proposition A.1 in the Appendix, we show that this truncation error is uniformly small. Let

$$B_n^{*i} = (B_{m_n k_n}^{*ij}, j \in V \setminus \{i\}) \in \mathbb{R}^{(p-1)k_n m_n \times m_n},$$

where

$$B_{m_n k_n}^{*ij} = \{\beta_{qrk}^{*ij} : 1 \leq q, r \leq m_n, 1 \leq k \leq k_n\}, \tag{4.2}$$

and define

$$B_n^{*N_n^i} = (B_{m_n k_n}^{*ij}, j \in N_n^i) \in \mathbb{R}^{n^i k_n m_n \times m_n}. \tag{4.3}$$

We wish to determine the 0 blocks of the truncated  $B_n^{*i}$  matrix. Recalling that  $\hat{B}_n^i = (\hat{B}_n^{ij}, j \in V \setminus \{i\}) \in \mathbb{R}^{(p-1)k_n m_n \times m_n}$  is the solution to the minimization problem (3.7). Then, the estimated neighbourhood for each node  $i \in V$  is

$$\hat{N}_n^i = \{j \in V \setminus \{i\} : \|\hat{B}_n^{ij}\|_F \neq 0\},$$

which yields the estimated edge set

$$\hat{E}_n = \{(i, j) \in V \times V : i \in \hat{N}_n^j \text{ and } j \in \hat{N}_n^i\}.$$

Finally, to derive our theoretical results, we need to introduce the following non-random matrices. Let  $\tilde{h}(\xi_r^j) = (\tilde{h}_1(\xi_r^j), \dots, \tilde{h}_{k_n}(\xi_r^j))^\top \in \mathbb{R}^{k_n}$  be the vector of the centered  $k_n$  B-splines evaluated at the unobserved scaled scores  $\xi_r^j$ ,  $r = 1, \dots, m_n, j \in V$ , and define the  $1 \times k_n m_n$  and  $1 \times n^i k_n m_n$  vectors

$$\tilde{H}(\xi^j) = (\tilde{h}^\top(\xi_r^j))_{1 \leq r \leq m_n},$$

$$\tilde{\mathbf{H}}^\top(\xi^{\mathbf{N}_n^i}) = (\tilde{H}(\xi^j), j \in \mathbf{N}_n^i), \quad (4.4)$$

where  $1 \leq n^i \leq p$  is the cardinality of the set  $\mathbf{N}_n^i$ . We introduce the following matrices

$$\Sigma_{\mathbf{N}_n^i \mathbf{N}_n^i}^* = E\left(\tilde{\mathbf{H}}(\xi^{\mathbf{N}_n^i})\tilde{\mathbf{H}}^\top(\xi^{\mathbf{N}_n^i})\right) \in \mathbb{R}^{n^i k_n m_n \times n^i k_n m_n} \quad (4.5)$$

and

$$\Sigma_{\xi^j \mathbf{N}_n^i}^* = E\left(\tilde{H}^\top(\xi^j)\tilde{\mathbf{H}}^\top(\xi^{\mathbf{N}_n^i})\right) \in \mathbb{R}^{k_n m_n \times n^i k_n m_n}. \quad (4.6)$$

For the statement of our theoretical results we require several assumptions. Assumption 4.1 is a similar assumption as made by [26] and refers to the eigen-system of the covariance operator defined in (2.1).

**Assumption 4.1.**

(i) There exist positive constants  $d_0, d_1$  and  $d_2$  such that

$$d_0 r^{-\beta} \leq \lambda_r^i \leq d_1 r^{-\beta}, \quad \lambda_r^i - \lambda_{r+1}^i \geq d_2^{-1} r^{-1-\beta} \quad \text{for } r \geq 1,$$

for some  $\beta > 1$ . We assume that  $X^i$  is  $m_n$ -dimensional; that is,  $\lambda_{m_n}^i \neq 0$  and  $\lambda_{m_n+1}^i = 0$ .

(ii) The number of principal component scores  $m_n$  satisfies  $m_n \asymp n^\alpha$  for some constant  $\alpha \in [0, \frac{1}{2+3\beta})$ .

(iii) The eigenfunctions  $\phi_r^i$  of the covariance operator defined in (2.2) are continuous and satisfy

$$\max_{j \in \mathbf{V}} \sup_{s \in [0,1]} \sup_{r \in \mathbf{N}} |\phi_r^j(s)| \leq C < \infty.$$

The next two conditions refer to the smoothness of the functions  $f_{qr}^{ij}$  in model (2.6). To be precise, let  $\kappa$  be a nonnegative integer and let  $\rho \in (0, 1]$ . We define  $\mathcal{F}_{\kappa, \rho}$  as the Hölder space of functions  $f : [-1, 1] \rightarrow \mathbb{R}$  whose  $\kappa$ th derivative exists and satisfies a Lipschitz condition of order  $\rho$ , and additionally satisfies the condition

$$\|f\|_\infty = \sup_{x \in [-1,1]} |f(x)| \leq F, \quad (4.7)$$

for some  $F > 0$ .

**Assumption 4.2.** Let  $d = \kappa + \rho > 0.5$  and assume  $f_{qr}^{ij} \in \mathcal{F}_{\kappa, \rho}$  and  $E[f_{qr}^{ij}(\xi_{ur}^j)] = 0$ , for all  $q, r = 1, \dots, m_n$  and  $(i, j) \in \mathbf{V} \times \mathbf{V}$ .

**Assumption 4.3.** The joint density function, say  $p^j$ , of the random vector  $\xi^j = (\xi_1^j, \dots, \xi_{m_n}^j)^\top$  is bounded away from zero and infinity on  $[-1, 1]^{m_n}$  for every  $j = 1, \dots, p$ .

In order to derive graph estimation consistency, we make the following assumption about the errors  $\epsilon_1^i, \dots, \epsilon_{m_n}^i$  in model (2.7). A similar condition was also postulated by [33] for joint additive models in the multivariate setting.

**Assumption 4.4.** *There exists a constant  $C > 0$  such that  $P(|\epsilon_q^i| > x) \leq 2 \exp(-Cx^2)$  for all  $x \geq 0$  and  $q = 1, \dots, m_n, i \in \mathcal{V}$ .*

**Assumption 4.5.**  *$k_n = O(n^\nu)$  for some  $\nu > 0$ , where  $\nu \geq \frac{\alpha(2+3\beta)}{2d-4}$  if  $d > 2$ .*

**Assumption 4.6** (Bounded eigenspectrum). *The minimum eigenvalue  $\Lambda_{\min}(\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^*)$  of the matrix  $\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^*$  defined in (4.5) satisfies for each  $i \in \mathcal{V}$  and all  $n \in \mathbb{N}$ ,*

$$\Lambda_{\min}(\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^*) > C_{\min},$$

for some constant  $C_{\min} > 0$ .

**Assumption 4.7** (Irrepresentable condition). *There exists a constant  $0 < \eta \leq 1$  such that for all  $i \in \mathcal{V}$ ,*

$$\max_{j \notin \mathbb{N}_n^i} \|\Sigma_{\xi^j \mathbb{N}_n^i}^* (\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^*)^{-1}\|_{\mathbb{F}} \leq \frac{1 - \eta}{\sqrt{n^i}}. \tag{4.8}$$

Assumption 4.6 states that the minimum eigenvalue of the population matrix  $\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}$  is bounded away from 0. Assumption 4.7 is the classical irrepresentable condition which is necessary and sufficient to show model selection consistency of the group lasso [39, 1]. According to [22] if the irrepresentable condition is relaxed, the lasso selects the correct non-zero coefficients but it may select some additional zero components. [28] and [23] considered similar assumptions for sparse additive and for high-dimensional multivariate regression models respectively. We now state our main theoretical result for the estimator  $\hat{\mathbf{N}}_n^i$  of the neighbourhood corresponding to the node  $i \in \mathcal{V}$ .

**Theorem 4.1.** *Suppose that Assumptions 4.1 - 4.7 are satisfied and the regularization parameter  $\lambda_n$  satisfies for all  $i$ ,*

$$\frac{n^i m_n^{3/2}}{k_n^d \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}} \lesssim \lambda_n \lesssim (n^i)^{-3/2} (b_n^{*i})^3 \left( \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \right)^{-2}, \tag{4.9}$$

where  $b_n^{*i} = \min_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}$ . Then,

$$P(\hat{\mathbf{N}}_n^i \neq \mathbf{N}_n^i) \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(pm_n k_n)\right),$$

where  $C_1 > 0$ .

The proof of Theorem 4.1 is complicated and is given in the Appendix. A major difficulty consists in the fact that the objective function (3.7) is based on the estimated scores and one has to establish concentration bounds in the estimation of the sample design matrix  $\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n$  using the estimated scores, rather than the true scores (stated as Theorem A.1 in the Appendix).

Using the union bound of probability and Theorem 4.1 we obtain the following result that states that the true edge set  $\mathbf{E}_n$  is equal to its estimate  $\hat{\mathbf{E}}_n$  with probability tending to 1.

**Corollary 4.1.** *If the assumptions of Theorem 4.1 are satisfied, we have for a positive constant  $C_1 > 0$*

$$P(\hat{E}_n \neq E_n) \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \min_{i=1}^p \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 3 \log(pm_n k_n)\right),$$

where  $\hat{E}_n$  is obtained by (3.8) or (3.9).

**Remark 4.1.**

(a) Under the assumptions of Theorem 4.1, it follows that

$$P(\hat{E}_n \neq E_n) \rightarrow 0,$$

if  $n \rightarrow \infty, p \rightarrow \infty$  and

$$\frac{n^{1-\alpha(2+3\beta)} (\lambda_n \min_{i=1}^p b_n^{*i})^2}{n^i m_n^2 k_n^4} \rightarrow \infty \quad \text{and} \quad \frac{n^i m_n^2 k_n^4 \log(pm_n k_n)}{\lambda_n^2} = o\left(\min_{i=1}^p b_n^{*i}\right)^2. \quad (4.10)$$

Note that the graph consistency result is achieved when  $p$  is much larger than the degree of each node  $i \in V$ ,  $n_i$ . For example, if  $p = O(n^\theta)$ ,  $\theta > 0$ ,  $\max_{i \in V} n^i = O(n^\gamma)$  with  $0 \leq \gamma < 1$ ,  $m_n = O(n^\alpha)$  with  $\alpha \in [0, \frac{1}{2+3\beta})$  and  $k_n = O(n^\nu)$ , then, conditions (4.10) reduce to

$$\frac{\log(pm_n k_n)}{n^{1-(\alpha(4+3\beta)+\gamma+4\nu)} \lambda_n} = o\left(\min_{i=1}^p b_n^{*i}\right).$$

(b) In the case of scalar data ( $\alpha = 0$ ), the conditions of Theorem 4.1 will be implied by

$$\frac{n^i}{k_n^d} \lesssim \lambda_n b_n^{*i}, \quad \lambda_n (n^i)^{3/2} \lesssim (b_n^{*i})^3 \quad \text{and} \quad \sqrt{\frac{n^i k_n^4 \log(pk_n)}{n \lambda_n^2}} = o(b_n^{*i}),$$

which are similar to the assumptions made in [28] and [33] for the analysis of scalar data by sparse additive models.

## 5. Finite sample properties

### 5.1. Simulated data

In this section we investigate the finite sample performance of the proposed model (AFGM) by means of a simulation study. We also compare the new methodology with the functional additive precision operator (FAPO) of [16] and the FGM of [26], where we consider two scenarios: nonlinear dependence and linear dependence.

It is not obvious how to construct flexible joint distributions for continuous random variables. However, joint distributions can be induced via structural

equation models with respect to a directed acyclic graph (DAG) [24], where each variable is expressed as a function of some other variables as well as some noise. Here, we consider additive noise models with additive non-linear differentiable functions and Gaussian noise [5]. These are special cases of structural equation models. In particular, given a DAG,  $G = (V, E)$ , we generate non-Gaussian scores by following the structural equations,

$$\xi_q^i = \sum_{j \in \text{pa}(i)} \sum_{r=1}^5 f_{qr}^{ij}(\xi_r^j) + \tilde{\epsilon}_q^j, \quad i = 1, \dots, p, q = 1, \dots, 5, \quad (5.1)$$

where  $\text{pa}(i)$  is the set of parents of node  $i$  in the associated DAG  $G$ ,  $\tilde{\epsilon}_q^j$  are Gaussian distributed random variables with 0 mean and  $f_{qr}^{ij}$  are non-linear and smooth functions. Next, we moralized the directed graph by connecting with an undirected edge all parent nodes and dropping the orientations of the directed edges. This gives us the undirected conditional independence graph of the distribution [6, 19]. Our goal is to recover the undirected conditional independence graph.

Then, we generate the multivariate functional data as

$$X_u^i(t_s) = \sum_{j \in \text{pa}_i} \sum_{q=1}^5 \sum_{r=1}^5 f_{qr}^{ij}(\xi_{ur}^j) \phi_q(t_s) + \epsilon_{us}^i, \quad u = 1, \dots, n, \quad (5.2)$$

where  $\phi_1^i, \dots, \phi_5^i$  are the first 5 functions of the orthonormal Fourier basis, and the errors  $\epsilon_{us}^i$  form an i.i.d. sample from a  $\mathcal{N}(0, 0.5^2)$  distribution.

In all simulation experiments in this section, this data has been smoothed to obtain continuous functions  $X_u^i$  using 10 B-spline basis functions of order 4; that is, piecewise polynomials of degree 3. For simplicity we assume  $f_{qr}^{ij}(x) = f(x)$  for all  $q, r = 1, \dots, 5$  and for all  $(j, i) \in E$ . In all examples, we center  $f(\xi_{ur}^j)$  to have 0 mean, and we generated  $n = 100$  functions observed at 100 equally spaced time points  $0 = t_1, \dots, t_{100} = 1$ . We consider directed acyclic graphs with  $p = 100$  nodes so that 1% of pairs of vertices are randomly selected as edges. We choose  $m_n$  functional principal components scores so that at least 90% of the total variation is explained. Furthermore, we approximate each additive function using B-splines of order 4. For simplicity we choose the same spline functions for all  $j = 1, \dots, p$  and for all  $r = 1, \dots, m_n$ . For the choice of  $k_n$ , we follow [20] and take  $k_n = 4 + \lceil \sqrt{n} \rceil$ .

For each scenario, we produce the average ROC curves (over 50 replications) for a range of 50 tuning parameters for the 3 functional graphical models estimators. To draw the curves, we compute for different regularization parameters  $\lambda$  the positive rate (sensitivity) and false positive rate (1-specificity) which are defined as

$$\text{TP} = \frac{\sum_{1 \leq j < i \leq p} I\{(i, j) \in E_n, (i, j) \in \hat{E}_n\}}{\sum_{1 \leq j < i \leq p} I\{(i, j) \in E_n\}}, \quad \text{FP} = \frac{\sum_{1 \leq j < i \leq p} I\{(i, j) \notin E_n, (i, j) \in \hat{E}_n\}}{\sum_{1 \leq j < i \leq p} I\{(i, j) \notin E_n\}}.$$

### 5.1.1. Scenario 1: nonlinear models

We use the following nonlinear models, where the linearity assumption (2.5) does not hold. We first consider the following model, Model I, used in [46]

$$\text{Model I: } f(x) = 1.4 + 3x - \frac{1}{2} + \sin(2\pi(x - \frac{1}{2})) + 8(x - \frac{1}{3})^2 - \frac{8}{9}.$$

For the choice of scores in (5.1), we simulate  $\xi_{u_r}^i$  independently from the uniform distribution  $U[-1/2, 1/2]$  for all  $r = 1, \dots, 5, i \in \mathbf{V}, u = 1, \dots, n$ . Furthermore, the errors  $\tilde{\epsilon}_{u_q}^i$  in (5.1) form an i.i.d. sample from the normal distribution with variance 1, that is  $\mathcal{N}(0, 1)$ .

The second example was considered in [20]

$$\text{Model II: } f(x) = -\sin(2x) + x^2 - 25/12 + x + \exp(-x) - 2/5 \cdot \sinh(5/2).$$

The scores  $\xi_{u_r}^i$  were simulated independently from the uniform distribution  $U[-2.5, 2.5]$  for all  $r = 1, \dots, 5, i \in \mathbf{V}, u = 1, \dots, n$ , and the errors  $\tilde{\epsilon}_{u_q}^i$  in (5.1) were simulated from the normal distribution,  $\mathcal{N}(0, 1)$ .

For each  $i = 1, \dots, 100$  and  $q = 1, \dots, 5$ , we define the overall signal-to-noise (SNR) ratio as

$$\text{SNR}_q^i = \frac{E|\sum_{j \in \text{Pa}_i} \sum_{r=1}^5 f_{qr}^{ij}(\xi_r^j)|}{\sqrt{\text{var}(\tilde{\epsilon}_q^i)}}.$$

We have calculated these SNRs by Monte Carlo simulation. Note also that the true graph is very sparse. The maximum number of parents is 3, where 28 nodes in the graph have 1 parent, 8 nodes have two parents and 2 nodes have 3 parents. In Model I, for nodes with one parent, the integral in the numerator of the SNR,  $\int_{-1}^1 \dots \int_{-1}^1 \frac{1}{32} |\sum_{r=1}^5 f(\xi_r^j)| d\xi_1^j \dots d\xi_5^j$ , is approximately equal to 17.8, giving a SNR ratio of 25.3. For nodes with 2 and 3 parents the SNR is computed as 50.44 and 75.7 respectively. Similarly, for Model II, the SNR for nodes with 1 parent is 6.20, whereas the SNR for nodes with 2 and 3 parents is 8.75 and 10.70 respectively.

The left and middle panel of Figure 2 show the averaged ROC curves over 40 replications corresponding to the two models. In first two lines of Table 1, we report the means and standard deviations (in parentheses) of the associated area-under-curve (AUC) values. An AUC close to 1 means a better performance for the estimator. We observe from the plots in Figure 2 and from Table 1, that for the AFGM estimator the areas under the ROC are substantially larger than for the FGGM, indicating that our new method AFGM dominates the FGGM. Similarly, the AFGM performs better than FAPO, indicating the benefit of a sparse and high-dimensional scheme.

### 5.1.2. Scenario 2: linear model

Next, we consider a model, where the linearity assumption is satisfied, to see how much efficiency might be lost by employing a nonparametric model under






FIG 2. ROC curves ((AFGM (—), FAPO (---), FGGM (···)) for Model I (left), Model II (middle) and Model III (right).

TABLE 1  
Means and standard errors (in parentheses) for AUC for models I and II.

$p$	Models	Methods		
		AFGM	FAPO	FGGM
100	I	0.73 (0.02)	0.67 (0.02)	0.59 (0.01)
	II	0.76 (0.01)	0.70 (0.02)	0.69 (0.02)
	III	0.89 (0.01)	0.82 (0.01)	0.90 (0.01)

the Gaussian assumption. The model is generated by (5.1) and (5.2) with

$$\text{Model III: } f(x) = x$$

The scores  $\xi_{u,v}^i$  were simulated independently from the standard Gaussian distribution. To implement the AFGM we truncate the scores such that they are located in the interval  $[-1, 1]$ . The SNR for nodes with 1 parent is approximately equal to 0.78, whereas the SNR for nodes with 2 and 3 parents is 1.08 and 1.32 respectively. The right panel in Figure 2 presents the averaged ROC curves for Model III, whereas the third line in Table 1 reports the means and standard deviations of AUC. We can see that under the linearity assumption the AFGM is comparable with the FGGM and both methods show an improvement compared to FAPO.

### 6. Real data application

In this section we apply the new method to the EEG data set available at the UCI Machine Learning Repository. The data involve 77 subjects in the alcoholic group and 45 subjects in the control group. Each subject was exposed to a stimulus while brain activities were recorded from the 64 electrodes placed on the subject’s scalp, over a one-second period in which 256 time points were sampled. See [41] and [13] for more backgrounds of this data. The goal is to characterize functional connectivity among the 64 nodes for the two groups, based on the functional data collected from the electrodes.

We choose  $k_n = 4 + \lceil \sqrt{n} \rceil$  B-spline functions of order 4 and number of scores equal to  $m_n = 5$ . Since our goal is to capture outstanding differences in brain




FIG 3. Estimated brain networks by AFGM (upper panels), FAPO (middle panels) and FGM (lower panels) for the alcoholic group (left), the control group (middle) and differential brain networks (right)

connectivity between the alcoholic group and control group, we take the tuning constant  $\lambda_n$  to be such that 5% of the  $\binom{64}{2}$  pairs of vertices are retained as edges.

Figure 3 shows the estimated brain networks constructed by the three methods for the alcoholic group (left) and the control group (middle). The right plots in Figure 3 represent the differential brain networks, where the red lines indicate the edges that are in the alcoholic network but not in the control network, and the blue lines indicate the edges that are in the control network but not in the alcoholic network. We observe that the brain networks have different patterns for the two groups. For example, we observe for all methods, that there is increased functional connectivity in the left frontal area for the alcoholic group relative to the control.

## 7. Conclusions

In this paper, we utilize the idea of generalized additive models to develop a new nonparametric graphical model for multivariate functional data which does not require the assumption of a Gaussian distribution. The conditional relation-

ships among the principal scores in the Karhunen-Loève expansion of a random function are allowed to take an arbitrary additive rather than a linear form as imposed by the assumption of Gaussianity. The additive functions are then approximated by linear combinations of B-splines. This approximation allows us to develop a group lasso algorithm to estimate the graph that encourages block-wise sparsity to a matrix formed by the coefficients in the spline approximation. We establish consistency of the procedure while both the number of principal components and the number of nodes diverge to infinity with increasing sample size. By simulation study and an analysis of a data example, we demonstrate the applicability of the new methodology.

The proposed model and methodology suggests many directions for future research. First, the asymptotic results here are developed under the framework where the order  $m_n$  in the expansion (2.7) tends to infinity with the sample size, and it is of interest if similar statistical guarantees can be obtained in model (2.6) with the infinite representation. Second, the theory is developed under the assumption that the random functions are fully observed. Therefore, an interesting and important question for future research is the extension of the methodology to smooth functions that are observed on a dense time grid such that the covariance operators and the functions are consistently estimated. Another important direction is the consideration of the sparse setting, where the functions are observed on a relatively small number of time points and are contaminated with noise. In this case, alternative approaches such as [37], [35] and [25] might be useful and will be further investigated in the future.

**Appendix A: Auxiliary results**

In this section we state some auxiliary results, which will be used in the proof of the Theorem 4.1. The next Lemma provides a concentration inequality for the norm  $\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}}$ , where  $\|\cdot\|_{\text{HS}}$  denotes the Hilbert-Schmidt norm. It can be proved by similar arguments as given in the proof of Lemma 6 in [26] observing the independence of the random variables  $\xi_{ur}^i$ . The details are omitted for the sake of brevity.

**Lemma A.1.** *Suppose that Assumption 4.1 is satisfied. Then, there exists a constant  $C_1$  such that for all  $0 < \epsilon \leq C_1$  and for each  $i = 1, \dots, p$*

$$P\left(\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} \geq \epsilon\right) \lesssim \exp(-C_1 n \epsilon^2).$$

Let  $\xi^i = (\xi_{ur}^i)_{\substack{1 \leq r \leq m_n \\ 1 \leq u \leq n}} \in \mathbb{R}^{n \times m_n}$  be the matrix of unobserved scores, and define

$$\tilde{h}_n(\xi_{ur}^j) = (\tilde{h}_{n1}(\xi_{ur}^j), \dots, \tilde{h}_{nk_n}(\xi_{ur}^j))^{\top}$$

as the vector of the centered  $k_n$  B-splines functions evaluated at the score  $\xi_{ur}^j$ , where

$$\tilde{h}_{nk}(\xi_{ur}^j) = h_k(\xi_{ur}^j) - \frac{1}{n} \sum_{u=1}^n h_k(\xi_{ur}^j), \quad k = 1, \dots, k_n. \tag{A.1}$$

We also assume the constraint  $\sum_{u=1}^n \sum_{k=1}^{k_n} h_k(\xi_{ur}^j) \beta_{qrk}^{ij} = 0$ ,  $q, r = 1, \dots, m_n, j \in \mathbb{V}$ . Note that this corresponds to (3.3), where the estimated scores  $\hat{\xi}_{ur}^j$  have been replaced by the unobserved scores  $\xi_{ur}^j$ . Let

$$\tilde{H}_n(\xi^j) = (\tilde{h}_n^\top(\xi_{ur}^j))_{1 \leq u \leq n, 1 \leq r \leq m_n} \in \mathbb{R}^{n \times k_n m_n}, \quad (\text{A.2})$$

$$\tilde{\mathbf{H}}_n^\top(\xi^{\mathbb{N}_n^i}) = (\tilde{H}_n(\xi^j), j \in \mathbb{N}_n^i) \in \mathbb{R}^{n \times n^i k_n m_n}, \quad (\text{A.3})$$

and define

$$\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n = \frac{1}{n} \tilde{\mathbf{H}}_n(\xi^{\mathbb{N}_n^i}) \tilde{\mathbf{H}}_n^\top(\xi^{\mathbb{N}_n^i}) \in \mathbb{R}^{n^i k_n m_n \times n^i k_n m_n}, \quad (\text{A.4})$$

which is the sample analog of the matrix  $\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^*$  defined in (4.5). Similarly, let  $\hat{\xi}^i = (\hat{\xi}_{ur}^i)_{1 \leq u \leq n, 1 \leq r \leq m_n}$  be the  $n \times m_n$  matrix of the estimated scores, and

$$\tilde{\mathbf{H}}_n^\top(\hat{\xi}^{\mathbb{N}_n^i}) = (\tilde{H}_n(\hat{\xi}^j), j \in \mathbb{N}_n^i) \in \mathbb{R}^{n \times n^i k_n m_n}, \quad (\text{A.5})$$

where

$$\tilde{H}_n(\hat{\xi}^j) = (\tilde{h}_n^\top(\hat{\xi}_{ur}^j))_{1 \leq u \leq n, 1 \leq r \leq m_n} \in \mathbb{R}^{n \times k_n m_n}, \quad (\text{A.6})$$

and  $\tilde{h}_n(\hat{\xi}_{ur}^j)$  is defined in (3.5). Then,

$$\hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n = \frac{1}{n} \tilde{\mathbf{H}}_n(\hat{\xi}^{\mathbb{N}_n^i}) \tilde{\mathbf{H}}_n^\top(\hat{\xi}^{\mathbb{N}_n^i}) \in \mathbb{R}^{n^i k_n m_n \times n^i k_n m_n} \quad (\text{A.7})$$

is the estimated version of the sample design matrix  $\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n$  in (A.4). The next result provides tail bounds for all entries of the matrix  $\hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n$ .

**Theorem A.1.** *Suppose that Assumption 4.1 holds. Then, there exists a positive constant  $C_1$  such that for any  $\delta > 0$  satisfying  $0 < \delta \leq C_1$  and for all  $(i, j) \in \mathbb{V} \times \mathbb{V}$ ,  $i \neq j$ ,  $r, q = 1, \dots, m_n$  and  $k, \ell = 1, \dots, k_n$ , we have*

$$\begin{aligned} P\left(\left|\frac{1}{n} \sum_{u=1}^n \left(\tilde{h}_{nk}(\hat{\xi}_{ur}^i) \tilde{h}_{n\ell}(\hat{\xi}_{uq}^j) - \tilde{h}_{nk}(\xi_{ur}^i) \tilde{h}_{n\ell}(\xi_{uq}^j)\right)\right| \geq \delta\right) \\ \lesssim \exp\left(-C_1 n^{1-\alpha(2+3\beta)} k_n^{-2} \delta^2\right). \end{aligned}$$

*Proof.* First, we have

$$\left|\sum_{u=1}^n \left(\tilde{h}_{nk}(\hat{\xi}_{ur}^i) \tilde{h}_{n\ell}(\hat{\xi}_{uq}^j) - \tilde{h}_{nk}(\xi_{ur}^i) \tilde{h}_{n\ell}(\xi_{uq}^j)\right)\right| \leq T_1 + T_2,$$

where the terms  $T_1$  and  $T_2$  are defined as

$$T_1 = \left|\sum_{u=1}^n \tilde{h}_{n\ell}(\xi_{uq}^j) \left(\tilde{h}_{nk}(\hat{\xi}_{ur}^i) - \tilde{h}_{nk}(\xi_{ur}^i)\right)\right|,$$

$$T_2 = \left| \sum_{u=1}^n \tilde{h}_{nk}(\hat{\xi}_{ur}^i) \left( \tilde{h}_{n\ell}(\hat{\xi}_{uq}^j) - \tilde{h}_{n\ell}(\xi_{uq}^j) \right) \right|.$$

Consequently, for any  $\delta > 0$ ,

$$\begin{aligned} P\left( \left| \sum_{u=1}^n \left( \tilde{h}_{nk}(\hat{\xi}_{ur}^i) \tilde{h}_{n\ell}(\hat{\xi}_{uq}^j) - \tilde{h}_{nk}(\xi_{ur}^i) \tilde{h}_{n\ell}(\xi_{uq}^j) \right) \right| \geq 2n\delta \right) \\ \leq P(T_1 \geq n\delta) + P(T_2 \geq n\delta), \end{aligned} \tag{A.8}$$

and therefore it is sufficient to derive inequalities for the two probabilities on the right-hand side of (A.8).

(a) We start with the probability  $P(T_1 \geq n\delta)$ . By the definition of  $\tilde{h}_{nk}(\hat{\xi}_{ur}^i)$  and  $\tilde{h}_{nk}(\xi_{ur}^i)$  in (3.3) and (A.1), respectively, and some elementary calculations, we obtain for any  $\delta > 0$

$$P(T_1 \geq n\delta) \leq P\left(T_{11} \geq \frac{n\delta}{2}\right) + P\left(T_{12} \geq \frac{n\delta}{2}\right),$$

where

$$\begin{aligned} T_{11} &= \left| \sum_{u=1}^n h_\ell(\xi_{uq}^j) (h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i)) \right|, \\ T_{12} &= \left| n^{-1} \sum_{v=1}^n h_\ell(\xi_{vq}^j) \sum_{u=1}^n (h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i)) \right|. \end{aligned}$$

We now derive a concentration inequality for  $T_{11}$ . By Cauchy-Schwarz inequality and using the fact that  $|h_\ell(\xi_{uq}^j)| \leq 1$  we have

$$P\left(T_{11} \geq \frac{n\delta}{2}\right) \leq P\left(\sum_{u=1}^n |h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i)|^2 \geq \frac{n\delta^2}{4}\right),$$

and Taylor's expansion gives

$$\left| h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i) \right| = \left| \frac{\partial h_k(\xi^*)}{\partial \xi_{ur}^i} \left( \hat{\xi}_{ur}^i - \xi_{ur}^i \right) \right|, \tag{A.9}$$

where  $\xi^*$  lies in the line segment between  $\hat{\xi}_{ur}^i$  and  $\xi_{ur}^i$ . From the derivative formula of the B-splines (see [7], Ch.10) there exists a constant  $M > 0$ , independent of  $n$ , such that for all  $k = 1, \dots, k_n$  and  $x \in [-1, 1]$ ,

$$\left| \frac{\partial h_k(x)}{\partial x} \right| \leq ML_n, \tag{A.10}$$

where  $L_n$  is the number of knots. As a result, using (A.9) and (A.10) we obtain

$$P(T_{11} \geq n\delta) \leq P\left(\sum_{u=1}^n |h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i)|^2 \geq \frac{n\delta^2}{4}\right)$$

$$\leq P\left(\sum_{u=1}^n |\hat{\xi}_{ur}^i - \xi_{ur}^i|^2 \geq \frac{n\delta^2}{4M^2L_n^2}\right).$$

Recall that  $\hat{\xi}_{ur}^i = (\hat{\lambda}_r^i)^{-1/2} \langle X_u^i, \hat{\phi}_r^i \rangle$  and  $\xi_{ur}^i = (\lambda_r^i)^{-1/2} \langle X_u^i, \phi_r^i \rangle$ . Then,

$$\begin{aligned} \hat{\xi}_{ur}^i - \xi_{ur}^i &= (\hat{\lambda}_r^i)^{-1/2} \langle X_u^i, \hat{\phi}_r^i \rangle - (\lambda_r^i)^{-1/2} \langle X_u^i, \phi_r^i \rangle \\ &\leq ((\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}) \langle X_u^i, \hat{\phi}_r^i \rangle + (\lambda_r^i)^{-1/2} \langle X_u^i, \hat{\phi}_r^i - \phi_r^i \rangle. \end{aligned}$$

By the Cauchy-Schwarz inequality and using the fact that  $\|\hat{\phi}_r^i\| = 1$  we obtain,

$$\begin{aligned} |\hat{\xi}_{ur}^i - \xi_{ur}^i| &\leq |(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}| \|X_u^i\| + (\lambda_r^i)^{-1/2} \|X_u^i\| \|\hat{\phi}_r^i - \phi_r^i\| \\ &\leq |(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}| \|X_u^i\| + (\lambda_r^i)^{-1/2} d_r^i \|X_u^i\| \|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}}, \end{aligned}$$

where we have used the inequality  $\|\hat{\phi}_r^i - \phi_r^i\| \leq d_r^i \|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}}$  [see Lemma 4.3 in 2] and assume w.l.o.g. that  $\hat{\phi}_r^i$  can be chosen to satisfy  $\text{sgn}(\langle \hat{\phi}_r^i, \phi_r^i \rangle) = 1$  (see the discussion in Remark 2.1). Using the inequality  $(a+b)^2 \leq 2(a^2+b^2)$ ,  $a, b \in \mathbb{R}$ , this implies

$$\begin{aligned} P(T_{11} \geq n\delta) &\leq P\left(|(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}|^2 \sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n\delta^2}{16M^2L_n^2}\right) \\ &\quad + P\left((\lambda_r^i)^{-1} (d_r^i)^2 \|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}}^2 \sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n\delta^2}{16M^2L_n^2}\right). \end{aligned} \tag{A.11}$$

We now consider the first term at the right-hand side of (A.11). Observe that  $\sum_{u=1}^n \|X_u^i\|^2 = \sum_{r=1}^{\infty} \lambda_r^i \sum_{u=1}^n (\xi_{ur}^i)^2$ , and recall that  $\xi_{ur}^i \in [-1, 1]$  with  $E(\xi_{ur}^i) = 1$ ,  $E((\xi_{ur}^i)^2) = 1$ . Thus  $|(\xi_{ur}^i)^2 - 1| \leq 2$ , which implies that for each  $r = 1, \dots, m_n$ ,  $\sum_{u=1}^n ((\xi_{ur}^i)^2 - 1)$  is a sub-Gaussian random variable with parameter proxy  $\sigma^2 = 4n$ . Consequently, we obtain by Theorem 2.1 of [3]

$$E\left\{\sum_{u=1}^n ((\xi_{ur}^i)^2 - 1)\right\}^{2k} \leq k!(16n)^k, \quad k \geq 1.$$

Using the convexity of the function  $x \mapsto x^{2k}$  and Jensen's inequality, it follows

$$\begin{aligned} E\left|\sum_{u=1}^n (\|X_u^i\|^2 - E\|X_u^i\|^2)\right|^{2k} &= E\left\{\sum_{r=1}^{\infty} \lambda_r^i \sum_{u=1}^n ((\xi_{ur}^i)^2 - 1)\right\}^{2k} \\ &\leq \sum_{r=1}^{\infty} \lambda_r^i E\left\{\sum_{u=1}^n ((\xi_{ur}^i)^2 - 1)\right\}^{2k} \left(\sum_{r=1}^{\infty} \lambda_r^i\right)^{2k-1} \\ &\leq k!(16\lambda_0^2 n)^k, \quad k \geq 1, \end{aligned}$$

where  $\lambda_0 = \sup_{i \leq p} \sum_{r=1}^{\infty} \lambda_r^i < \infty$  (due to Assumption 4.1). Hence, from Theorem 2.1 of [3], we obtain for all  $\epsilon > 0$ ,

$$P\left(\sum_{u=1}^n (\|X_u^i\|^2 - E\|X_u^i\|^2) \geq \epsilon\right) \leq \exp\left(-\frac{\epsilon^2}{128\lambda_0^2 n}\right).$$

Furthermore,  $E(\sum_{u=1}^n \|X_u^i\|^2) \leq n\lambda_0$ . Thus, for all  $\epsilon/2 \geq n\lambda_0$ ,

$$P\left(\sum_{u=1}^n \|X_u^i\|^2 \geq \epsilon\right) \leq P\left(\sum_{u=1}^n (\|X_u^i\|^2 - E\|X_u^i\|^2) \geq \epsilon\right) \leq \exp\left(-\frac{\epsilon^2}{128\lambda_0^2 n}\right). \quad (\text{A.12})$$

Now, we obtain for the first probability on the right-hand side of (A.11)

$$\begin{aligned} & P\left(|(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}|^2 \sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n\delta^2}{16M^2 L_n}\right) \\ & \leq P\left(|(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}| \geq \frac{\delta}{2M^{1/2} L_n}\right) + P\left(\sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n}{4M}\right). \quad (\text{A.13}) \end{aligned}$$

Define the event  $\Omega_{m_n}^i = \{\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} < 2^{-1}\delta_{m_n}^i\}$ , where  $\delta_{m_n}^i = \min_{1 \leq r \leq m_n} \{\lambda_r^i - \lambda_{r+1}^i\}$ . Assumption 4.1(i) implies  $\delta_{m_n}^i \geq d_2^{-1} m_n^{-(1+\beta)}$  leading to

$$P((\Omega_{m_n}^i)^c) \leq P(\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} \geq 2^{-1} d_2^{-1} m_n^{-(1+\beta)}) \lesssim \exp(-C_1 n m_n^{-2(1+\beta)}), \quad (\text{A.14})$$

for some  $C_1 > 0$ , where we have used Lemma A.1 with  $\delta = 2^{-1} d_2^{-1} m_n^{-(1+\beta)}$ . Furthermore, from Lemma 4.43 of [2] we have on the event  $\Omega_{m_n}^i$

$$\sup_{r \geq 1} |\hat{\lambda}_r^i - \lambda_r^i| \leq \|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} \leq 2^{-1} \delta_{m_n}^i \leq 2^{-1} \lambda_{m_n}^i.$$

This implies  $\hat{\lambda}_r^i \geq \frac{\lambda_r^i}{2}$ ,  $\hat{\lambda}_r^i \leq 2\lambda_r^i$  and

$$|(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}| \leq \frac{(\hat{\lambda}_r^i)^{-1} |\hat{\lambda}_r^i - \lambda_r^i| (\lambda_r^i)^{-1}}{(\hat{\lambda}_r^i)^{-1/2} + (\lambda_r^i)^{-1/2}} \leq 2(\lambda_r^i)^{-3/2} |\hat{\lambda}_r^i - \lambda_r^i|.$$

This together with (A.14) imply that

$$\begin{aligned} & P\left(|(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}| \geq \frac{\delta}{2M^{1/2} L_n}\right) \\ & \leq P\left(|(\hat{\lambda}_r^i)^{-1/2} - (\lambda_r^i)^{-1/2}| \geq \frac{\delta}{2M^{1/2} L_n} \cap \Omega_{m_n}^i\right) + P((\Omega_{m_n}^i)^c) \\ & \leq P\left(|\hat{\lambda}_r^i - \lambda_r^i| \geq \frac{\delta(\lambda_r^i)^{3/2}}{4M^{1/2} L_n}\right) + P((\Omega_{m_n}^i)^c) \\ & \leq P\left(\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} \geq \frac{\delta(\lambda_r^i)^{3/2}}{4M^{1/2} L_n}\right) + P((\Omega_{m_n}^i)^c), \end{aligned}$$

where we used Lemma 4.43 of [2] for the third inequality. Therefore, from this, Lemma A.1, (A.12) with  $\epsilon = \frac{n}{4M}$  and the fact that  $d_0 r^{-\beta} \leq \lambda_r^i$ , the right-hand side of (A.13) can be upper-bounded by

$$P\left(\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} \geq \frac{\delta(\lambda_r^i)^{3/2}}{4M^{1/2} L_n}\right) + P\left(\sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n}{4M}\right) + P((\Omega_{m_n}^i)^c) \quad (\text{A.15})$$

$$\lesssim \exp(-C_1 n L_n^{-2} m_n^{-3\beta} \delta^2) + \exp(-C_2 n) + \exp(-C_3 n m_n^{-2(\beta+1)}),$$

for all  $0 < \delta L_n^{-1} m_n^{-3\beta/2} \leq C_1$  and some positive constants  $C_1, C_2$  and  $C_3$ .

For the second term on the right-hand side of the inequality (A.11) we use Lemma A.1, (A.12) and the fact that  $(d_r^i)^{-1} \geq \frac{d_2}{2\sqrt{2}} m_n^{-1-\beta}$ , to obtain,

$$\begin{aligned} & P\left((\lambda_r^i)^{-1} (d_r^i)^2 \|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}}^2 \sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n\delta^2}{16M^2 L_n^2}\right) \\ & \leq P\left(\|\hat{\Sigma}_{X^i X^i} - \Sigma_{X^i X^i}\|_{\text{HS}} \geq \frac{\delta(\lambda_r^i)^{1/2} (d_r^i)^{-1}}{2M^{1/2} L_n}\right) + P\left(\sum_{u=1}^n \|X_u^i\|^2 \geq \frac{n}{4M}\right) \\ & \lesssim \exp(-C_4 n) + \exp(-C_5 n L_n^{-2} m_n^{-(2+3\beta)} \delta^2), \end{aligned} \tag{A.16}$$

for  $0 < \delta L_n^{-1} m_n^{-(2+3\beta)/2} \leq C_5$  and  $C_4 > 0, C_5 > 0$ .

Combining (A.15) and (A.16) we obtain for all  $0 < \delta L_n^{-1} m_n^{-3\beta/2} \leq C_3$  the inequality

$$P(T_{11} \geq n\delta) \lesssim \exp(-C_1 n) + \exp(-C_2 n m_n^{-2(\beta+1)}) + \exp(-C_3 n L_n^{-2} m_n^{-(2+3\beta)} \delta^2). \tag{A.17}$$

We now consider the probability  $P(T_{12} \geq n\delta)$ . Using the fact that  $|h_\ell(\xi_{uq}^j)| \leq 1$  and Taylor's expansion (A.9) yield

$$P(T_{12} \geq n\delta) \leq P\left(\left|\sum_{u=1}^n (\hat{\xi}_{ur}^i - \xi_{ur}^i)\right| \geq \frac{n\delta}{2ML_n}\right) \leq P\left(\sum_{u=1}^n |\hat{\xi}_{ur}^i - \xi_{ur}^i|^2 \geq \frac{n\delta^2}{4M^2 L_n^2}\right),$$

where we have used the Cauchy-Schwarz inequality. Therefore, by similar arguments as used in the derivation of the bound for  $P(T_{11} \geq n\delta)$ , there exist positive constants  $C_4, C_5$  and  $C_6$  such that for all  $0 < \delta L_n^{-1} m_n^{-3\beta/2} \leq C_6$

$$P(T_{12} \geq n\delta) \lesssim \exp(-C_4 n) + \exp(-C_5 n m_n^{-2(\beta+1)}) + \exp(-C_6 n L_n^{-2} m_n^{-(2+3\beta)} \delta^2). \tag{A.18}$$

Combining (A.17) and (A.18) and choosing suitable constants, we obtain for  $0 < \delta L_n^{-1} m_n^{-3\beta/2} \leq C_1$ ,

$$P(T_1 \geq n\delta) \lesssim \exp(-C_1 n L_n^{-2} m_n^{-(2+3\beta)} \delta^2). \tag{A.19}$$

(b) To derive a bound for the term  $P(T_2 \geq n\delta)$  in (A.8) we use the decomposition

$$\begin{aligned} P(T_2 \geq n\delta) & \leq P\left(\left|\sum_{u=1}^n \left(\tilde{h}_{nk}(\hat{\xi}_{ur}^i) - \tilde{h}_{nk}(\xi_{ur}^i)\right) \left(\tilde{h}_{n\ell}(\hat{\xi}_{uq}^j) - \tilde{h}_{n\ell}(\xi_{uq}^j)\right)\right| \geq \frac{n\delta}{2}\right) \\ & \quad + P\left(\left|\sum_{u=1}^n \tilde{h}_k(\xi_{ur}^i) \left(\tilde{h}_{n\ell}(\hat{\xi}_{uq}^j) - \tilde{h}_{n\ell}(\xi_{uq}^j)\right)\right| \geq \frac{n\delta}{2}\right) \end{aligned}$$



$$=P\left(T_{21} \geq \frac{n\delta}{2}\right) + P\left(T_{22} \geq \frac{n\delta}{2}\right),$$

where the last inequality defines the terms  $T_{21}$  and  $T_{22}$  in an obvious manner. For the second term, we obtain by the same arguments as used to estimate  $P(T_1 \geq n\delta)$  for any  $0 < \delta L_n^{-1} m_n^{-3\beta/2} \leq C_2$ ,

$$P\left(T_{22} \geq \frac{n\delta}{2}\right) \lesssim \exp(-C_2 n L_n^{-2} m_n^{-(2+3\beta)} \delta^2).$$

For the first term, we use the definition of the centred B-splines in (3.3) and (A.1) to obtain for any  $\delta > 0$

$$P\left(T_{21} \geq \frac{n\delta}{2}\right) \leq P\left(T_{211} \geq \frac{n\delta}{4}\right) + P\left(T_{212} \geq \frac{n\delta}{4}\right),$$

where

$$T_{211} = \left| \sum_{u=1}^n \left( h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i) \right) \left( h_\ell(\hat{\xi}_{uq}^j) - h_\ell(\xi_{uq}^j) \right) \right|,$$

$$T_{212} = n^{-1} \left| \sum_{u=1}^n \left( h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i) \right) \right| \left| \sum_{u=1}^n \left( h_\ell(\hat{\xi}_{uq}^j) - h_\ell(\xi_{uq}^j) \right) \right|.$$

To derive a concentration bound for the first term, we use (A.9), (A.10) and the Cauchy-Schwarz inequality to obtain

$$P\left(T_{211} \geq \frac{n\delta}{4}\right) \leq P\left(\sum_{u=1}^n |\hat{\xi}_{ur}^i - \xi_{ur}^i| |\hat{\xi}_{uq}^j - \xi_{uq}^j| \geq \frac{n\delta}{4M^2 L_n^2}\right)$$

$$\leq 2P\left(\sum_{u=1}^n |\hat{\xi}_{ur}^i - \xi_{ur}^i|^2 \geq \frac{n\delta}{4M^2 L_n^2}\right) \lesssim \exp(-C_3 n L_n^{-2} m_n^{-(2+3\beta)} \delta),$$

for a positive constant  $C_3$  such that for  $0 < \delta^{1/2} L_n^{-1} m_n^{-3\beta/2} \leq C_3$ . Here the last inequality follows by the same arguments as used for the bound of  $P(T_{11} \geq n\delta)$ . Finally, for the term  $P(T_{212} \geq \frac{n\delta}{4})$  we have

$$P\left(T_{212} \geq \frac{n\delta}{4}\right) = P\left(\left| \sum_{u=1}^n \left( h_k(\hat{\xi}_{ur}^i) - h_k(\xi_{ur}^i) \right) \right| \left| \sum_{u=1}^n \left( h_\ell(\hat{\xi}_{uq}^j) - h_\ell(\xi_{uq}^j) \right) \right| \geq \frac{n^2 \delta}{4}\right).$$

Thus, we can apply similar techniques as used in the derivation of the bound (A.17) with  $\frac{n\delta}{2}$  replaced by  $\frac{n\delta^{1/2}}{2}$  leading to

$$P\left(T_{212} \geq \frac{n\delta}{4}\right) \lesssim \exp(-C_4 n L_n^{-2} m_n^{-(2+3\beta)} \delta),$$

for  $0 < \delta^{1/2} L_n^{-1} m_n^{-3\beta/2} \leq C_4$ . Combining these results with (A.8) and (A.19) and using the fact that  $m_n \asymp n^\alpha$  and  $k_n > L_n$ , we obtain the assertion of the Theorem.  $\square$

Theorem A.1 states that the elements of the matrix  $\hat{\Sigma}_{N_n^i N_n^i} - \Sigma_{N_n^i N_n^i}^n$  exhibit exponential-type probability tails. We also observe that the decay rate  $\beta$  of the eigenvalues appears in the tail behaviour. A similar condition of exponential tails is imposed on the elements of the sample covariance matrix of scalar and functional Gaussian data for the analysis of high-dimensional Gaussian graphical models [see, for example, 29, 26].

**Proposition A.1.** *Suppose that Assumptions 4.2, 4.3 and condition (4.7) are satisfied. Then, there exist functions  $\tilde{f}_{nqr}^{ij} = \sum_{k=1}^{k_n} \beta_{qrk}^{*ij} \tilde{h}_{nk}$  and positive constants  $c_1, C_1$ , such that*

$$P(\Omega^c) \leq 2 \exp\left(-C_1 \frac{nk_n^{-2d}}{n^i m_n^2} + \log(n^i m_n^2)\right),$$

where

$$\Omega = \left\{ \max_{j \in N_n^i} \max_{1 \leq q, r \leq m_n} \frac{1}{\sqrt{n}} \|\mathbf{f}_{qr}^{ij} - \tilde{\mathbf{f}}_{nqr}^{ij}\|_2 < c_1 k_n^{-d} \right\}, \quad (\text{A.20})$$

and  $\mathbf{f}_{qr}^{ij} = (f_{qr}^{ij}(\xi_{1r}^j), \dots, f_{qr}^{ij}(\xi_{nr}^j))^\top$ ,  $\tilde{\mathbf{f}}_{nqr}^{ij} = (\tilde{f}_{nqr}^{ij}(\xi_{1r}^j), \dots, \tilde{f}_{nqr}^{ij}(\xi_{nr}^j))^\top$ .

*Proof.* By Assumptions 4.2 and 4.3 for any  $f_{qr}^{ij} \in \mathcal{F}_{\kappa, \rho}$  there exists a B-spline  $g_{qr}^{ij} = \sum_{k=1}^{k_n} \beta_{qrk}^{*ij} h_k \in \mathcal{S}_{\ell L_n}$  and a positive constant  $c_1$  such that

$$\|f_{qr}^{ij} - g_{qr}^{ij}\|_\infty \leq c_1 k_n^{-d},$$

(see Lemma 5 in [32]). Let  $\tilde{f}_{nqr}^{ij}(\xi_{ur}^j) = g_{qr}^{ij}(\xi_{ur}^j) - \frac{1}{n} \sum_{u=1}^n g_{qr}^{ij}(\xi_{ur}^j)$ . Then recalling the notation (A.14) we have  $\tilde{f}_{nqr}^{ij}(\xi_{ur}^j) = \sum_{k=1}^{k_n} \beta_{qrk}^{*ij} \tilde{h}_{nk}(\xi_{ur}^j)$ , and we obtain

$$\begin{aligned} \frac{1}{n} \|\mathbf{f}_{qr}^{ij} - \tilde{\mathbf{f}}_{nqr}^{ij}\|_2^2 &\leq 2c_1^2 k_n^{-2d} + 2 \left( \frac{1}{n} \sum_{u=1}^n g_{qr}^{ij}(\xi_{ur}^j) \right)^2 \\ &\leq 2c_1^2 k_n^{-2d} + 4 \left( \frac{1}{n} \sum_{u=1}^n (g_{qr}^{ij}(\xi_{ur}^j) - f_{qr}^{ij}(\xi_{ur}^j)) \right)^2 + 4 \left( \frac{1}{n} \sum_{u=1}^n f_{qr}^{ij}(\xi_{ur}^j) \right)^2 \\ &\leq 6c_1^2 k_n^{-2d} + 4 \left( \frac{1}{n} \sum_{u=1}^n f_{qr}^{ij}(\xi_{ur}^j) \right)^2. \end{aligned}$$

From this, condition (4.7), Hoeffding's inequality and the union bound, it follows with an appropriate constant  $c_2 > 0$

$$\begin{aligned} &P\left( \max_{j \in N_n^i} \max_{1 \leq q, r \leq m_n} \frac{1}{n} \|f_{qr}^{ij}(\xi_r^j) - \tilde{f}_{nqr}^{ij}(\xi_r^j)\|_2^2 \geq c_1^2 k_n^{-2d} \right) \\ &\leq P\left( \max_{j \in N_n^i} \max_{1 \leq q, r \leq m_n} \left| \frac{1}{n} \sum_{u=1}^n f_{qr}^{ij}(\xi_{ur}^j) \right| \geq c_2 k_n^{-d} \right) \\ &\leq 2 \exp\left(-\frac{nk_n^{-2d}}{2M^2 n^i m_n^2} + \log(n^i m_n^2)\right), \end{aligned}$$

which completes the proof.  $\square$

### A.1. Rates of convergence for sample design matrices

In this section we show that if Assumptions 4.6 and 4.7 hold, then with high probability, the assumptions hold also for the corresponding *sample matrices*

$$\begin{aligned}\Sigma_{N_n^i N_n^i}^n &= \frac{1}{n} \tilde{\mathbf{H}}_n(\xi^{N_n^i}) \tilde{\mathbf{H}}_n^\top(\xi^{N_n^i}) \in \mathbb{R}^{n^i k_n m_n \times n^i k_n m_n}, \\ \Sigma_{\xi^j N_n^i}^n &= \frac{1}{n} \tilde{H}_n^\top(\xi^j) \tilde{\mathbf{H}}_n^\top(\xi^{N_n^i}) \in \mathbb{R}^{k_n m_n \times k_n m_n n^i},\end{aligned}\quad (\text{A.21})$$

where  $\tilde{H}_n(\xi^j)$  and  $\tilde{\mathbf{H}}_n(\xi^{N_n^i})$  are defined in (A.2) and (A.3), respectively. Note that the matrices in (A.21) are based on the unobserved scores and are the sample analogs of the matrices  $\Sigma_{N_n^i N_n^i}^*$  and  $\Sigma_{\xi^j N_n^i}^*$  in (4.5) and (4.6), respectively.

**Lemma A.2.** *Suppose that Assumption 4.6 holds. Then, there exists a constant  $C_1 > 0$  such that for any  $\delta > 0$ ,*

$$P\left(\left\|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\right\|_F \geq \delta\right) \leq 2 \exp\left(-C_1 \frac{n\delta^2}{(n^i m_n k_n)^2} + 2 \log(n^i m_n k_n)\right). \quad (\text{A.22})$$

$$P\left(\Lambda_{\min}(\Sigma_{N_n^i N_n^i}^n) \leq C_{\min} - \delta\right) \leq 2 \exp\left(-C_1 \frac{n\delta^2}{(n^i m_n k_n)^2} + 2 \log(n^i m_n k_n)\right). \quad (\text{A.23})$$

*Proof.* Weyl's Lemma yields

$$\Lambda_{\min}(\Sigma_{N_n^i N_n^i}^*) - \Lambda_{\min}(\Sigma_{N_n^i N_n^i}^n) \leq \|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\|_2 \leq \|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\|_F,$$

and by Assumption 4.6 we have,

$$P\left(\Lambda_{\min}(\Sigma_{N_n^i N_n^i}^n) \leq C_{\min} - \delta\right) \leq P\left(\|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\|_F \geq \delta\right). \quad (\text{A.24})$$

By definition the  $n^i m_n k_n \times n^i m_n k_n$  matrix  $\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*$  contains elements of the form

$$W_{k\ell, r q}^{j_1, j_2} = \frac{1}{n} \sum_{u=1}^n \tilde{h}_{nk}(\xi_{uq}^{j_1}) \tilde{h}_{n\ell}(\xi_{ur}^{j_2}) - E(\tilde{h}_k(\xi_q^{j_1}) \tilde{h}_\ell(\xi_r^{j_2})),$$

which can be rewritten (recalling the notation (A.1)) as  $A_1 - A_2$ , where

$$\begin{aligned}A_1 &= \frac{1}{n} \sum_{u=1}^n h_k(\xi_{uq}^{j_1}) h_\ell(\xi_{ur}^{j_2}) - E(h_k(\xi_q^{j_1}) h_\ell(\xi_r^{j_2})), \\ A_2 &= \frac{1}{n^2} \sum_{u_1=1}^n \sum_{u_2=1}^n h_k(\xi_{u_1 q}^{j_1}) h_\ell(\xi_{u_2 r}^{j_2}) - E(h_k(\xi_q^{j_1})) E(h_\ell(\xi_r^{j_2})).\end{aligned}$$

Next, observe that the summands of  $A_1$  have expectation 0 and are bounded in absolute value by 2. Therefore, by Hoeffding's inequality, for any  $\epsilon > 0$ , we have  $P(|A_1| \geq \epsilon) \leq 2 \exp(-\frac{n\epsilon^2}{128})$ . Moreover, the term  $A_2$  can be written as  $\frac{n-1}{n}A_{21} + A_{22}$ , where

$$A_{21} = \frac{1}{n(n-1)} \sum_{u_1 \neq u_2}^n h_k(\xi_{u_1q}^{j_1}) h_\ell(\xi_{u_2r}^{j_2}) - E(h_k(\xi_{u_1q}^{j_1}) h_\ell(\xi_{u_2r}^{j_2})),$$

is a  $U$ -statistic and  $A_{22} = \frac{1}{n^2} \sum_{u_1=1}^n h_k(\xi_{u_1q}^{j_1}) h_\ell(\xi_{u_1r}^{j_2}) - E(h_k(\xi_{u_1q}^{j_1}) h_\ell(\xi_{u_1r}^{j_2}))$ . Consequently, by Hoeffding's inequality for  $U$ -statistics [10], for any  $\epsilon > 0$ ,  $P(|A_{21}| \geq \epsilon) \leq 2 \exp(-\frac{n\epsilon^2}{128})$ , and it is easy to see (due to the additional factor  $1/n$ ) that  $A_{22}$  satisfies an even stronger concentration inequality. Therefore, it follows that for any  $\epsilon > 0$

$$P(|W_{k\ell,rq}^{j_1,j_2}| \geq \epsilon) \leq 2 \exp(-C_1 n \epsilon^2), \quad (\text{A.25})$$

for some constant  $C_1 > 0$ . Thus, the union bound over the  $(n^i m_n k_n)^2$  indices and the choice of  $\epsilon = \frac{\delta}{n^i m_n k_n}$  in (A.25) yields (A.22). Finally, the assertion (A.23) follows from relation (A.24) at the beginning of the proof.  $\square$

The next Lemma guarantees that the matrices defined in (A.21) satisfy the irrepresentable condition in Assumption 4.7 with high probability.

**Lemma A.3.** *If Assumption 4.6 and 4.7 are satisfied for some  $0 < \eta \leq 1$ , then*

$$P\left(\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1}\|_{\mathbb{F}} \geq \frac{1-\eta}{\sqrt{n^i}}\right) \lesssim \exp\left(-C_1 \frac{n}{((n^i)^{5/4} m_n k_n)^2} + 2 \log(pm_n k_n)\right),$$

where  $C_1$  is a positive constant that depends only on  $C_{\min}$  and  $\eta$ .

*Proof.* First, we decompose

$$\begin{aligned} \max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1}\|_{\mathbb{F}} &\leq \max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1} - \Sigma_{\xi^j N_n^i}^* (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_{\mathbb{F}} \\ &\quad + \max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^* (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_{\mathbb{F}}. \end{aligned}$$

By Assumption 4.7 we have  $\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^* (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_{\mathbb{F}} \leq \frac{1-\eta}{\sqrt{n^i}}$  and it suffices to consider

$$P\left(\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1} - \Sigma_{\xi^j N_n^i}^* (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_{\mathbb{F}} \geq \frac{\eta}{2\sqrt{n^i}}\right).$$

For this purpose we use the decomposition  $\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1} - \Sigma_{\xi^j N_n^i}^* (\Sigma_{N_n^i N_n^i}^*)^{-1} = T_1^j + T_2^j + T_3^j$  where

$$T_1^j = \Sigma_{\xi^j N_n^i}^* \left( (\Sigma_{N_n^i N_n^i}^n)^{-1} - (\Sigma_{N_n^i N_n^i}^*)^{-1} \right), \quad T_2^j = \left( \Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^* \right) (\Sigma_{N_n^i N_n^i}^*)^{-1},$$

$$T_3^j = \left( \Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^* \right) \left( (\Sigma_{N_n^i N_n^i}^n)^{-1} - (\Sigma_{N_n^i N_n^i}^*)^{-1} \right),$$

and control the probabilities  $P(\max_{j \notin N_n^i} \|T_h^j\|_F \geq \frac{\eta}{6\sqrt{n^i}})$  separately.

(a) For the first term  $T_1^j$ , we use the identity  $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$  and obtain from Assumption 4.7

$$\begin{aligned} \max_{j \notin N_n^i} \|T_1^j\|_F &\leq \max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^* (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_F \|(\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*) (\Sigma_{N_n^i N_n^i}^n)^{-1}\|_F \\ &\leq \frac{(1-\eta)}{\sqrt{n^i}} \|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\|_F \|(\Sigma_{N_n^i N_n^i}^n)^{-1}\|_2. \end{aligned}$$

Thus, defining the event  $\mathcal{T} = \{\|(\Sigma_{N_n^i N_n^i}^n)^{-1}\|_2 \leq \frac{2}{C_{\min}}\}$  we obtain

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|T_1^j\|_F \geq \frac{\eta}{6\sqrt{n^i}}\right) &\leq P\left(\|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\|_F \|(\Sigma_{N_n^i N_n^i}^n)^{-1}\|_2 \geq \frac{\eta}{6(1-\eta)}\right) \\ &\leq P\left(\|\Sigma_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^*\|_F \geq \frac{\eta C_{\min}}{12(1-\eta)}\right) + P(\mathcal{T}^c) \\ &\leq 4 \exp\left(-C_1 \frac{n}{(n^i m_n k_n)^2} + 2 \log(n^i m_n k_n)\right), \end{aligned} \tag{A.26}$$

where we used Lemma A.2 with  $\delta = \frac{\eta C_{\min}}{12(1-\eta)}$  and  $\delta = \frac{C_{\min}}{2}$  for the last inequality.

(b) For the second term  $T_2^j$ , we have

$$\max_{j \notin N_n^i} \|T_2^j\|_F \leq \|(\Sigma_{N_n^i N_n^i}^*)^{-1}\|_2 \max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_F \leq C_{\min}^{-1} \max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_F,$$

where we used Assumption 4.6 in the second inequality. Thus,

$$P\left(\max_{j \notin N_n^i} \|T_2^j\|_F \geq \frac{\eta}{6\sqrt{n^i}}\right) \leq P\left(\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_F \geq \frac{\eta C_{\min}}{6\sqrt{n^i}}\right).$$

Now, using similar arguments as in the proof of (A.22) in Lemma A.2, we can show

$$P\left(\|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_F \geq \delta\right) \leq 2 \exp\left(-C_1 \frac{n\delta^2}{n^i m_n^2 k_n^2} + \log(n^i m_n^2 k_n^2)\right), \tag{A.27}$$

for some positive constant  $C_1 > 0$ . This bound with  $\delta = \frac{\eta C_{\min}}{6\sqrt{n^i}}$  and the union bound yield

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|T_2^j\|_F \geq \frac{\eta}{6\sqrt{n^i}}\right) &\leq (p - n^i) P\left(\|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_F \geq \frac{\eta C_{\min}}{6\sqrt{n^i}}\right) \\ &\leq 2 \exp\left(-C_1 \frac{n}{(n^i m_n k_n)^2} + \log(n^i m_n^2 k_n^2) + \log(p - n^i)\right). \end{aligned} \tag{A.28}$$

(c) For the third term  $T_3^j$ , we have

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|T_3^j\|_{\mathbb{F}} \geq \frac{\eta}{6\sqrt{n^i}}\right) &\leq P\left(\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_{\mathbb{F}} \geq \sqrt{\frac{\eta}{6\sqrt{n^i}}}\right) \\ &\quad + P\left(\|(\Sigma_{N_n^i N_n^i}^n)^{-1} - (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_{\mathbb{F}} \geq \sqrt{\frac{\eta}{6\sqrt{n^i}}}\right). \end{aligned} \quad (\text{A.29})$$

Using (A.27) with  $\delta = \sqrt{\frac{\eta}{6\sqrt{n^i}}}$  we obtain for the first term on the right-hand side of (A.29)

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^*\|_{\mathbb{F}} \geq \sqrt{\frac{\eta}{6\sqrt{n^i}}}\right) & \quad (\text{A.30}) \\ &\lesssim \exp\left(-C_1 \frac{n}{(n^i)^{3/2} m_n^2 k_n^2} + \log(n^i m_n^2 k_n^2) + \log((p - n^i) m_n^2 k_n^2)\right). \end{aligned}$$

To derive a bound for the second term in (A.29) we apply the same arguments as used for the term  $T_1^j$

$$\begin{aligned} P\left(\|(\Sigma_{N_n^i N_n^i}^n)^{-1} - (\Sigma_{N_n^i N_n^i}^*)^{-1}\|_{\mathbb{F}} \geq \sqrt{\frac{\eta}{6\sqrt{n^i}}}\right) & \\ \lesssim \exp\left(-C_1 \frac{n}{(n^i)^{5/2} m_n^2 k_n^2} 2 \log(n^i m_n k_n)\right) &+ \exp\left(-C_1 \frac{n}{(n^i m_n k_n)^2} + 2 \log(n^i m_n k_n)\right). \end{aligned} \quad (\text{A.31})$$

Thus, from (A.29), (A.30) and (A.31) we obtain

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|T_3^j\|_{\mathbb{F}} \geq \frac{\eta}{6\sqrt{n^i}}\right) & \quad (\text{A.32}) \\ &\lesssim \exp\left(-C_1 \frac{n}{(n^i)^{3/2} (m_n k_n)^2} + \log(n^i m_n^2 k_n^2) + \log((p - n^i) m_n^2 k_n^2)\right) \\ &+ 2 \exp\left(-C_1 \frac{n}{(n^i)^{5/2} m_n^2 k_n^2} + 2 \log(n^i m_n k_n)\right) & (\text{A.33}) \\ &+ 2 \exp\left(-C_1 \frac{n}{(n^i m_n k_n)^2} + 2 \log(n^i m_n k_n)\right). \end{aligned}$$

Putting together (A.26), (A.28) and (A.32) and using the fact  $\log n^i \leq \log(p - n^i) \leq \log p$  (since  $n^i \leq p$ ) we conclude

$$P\left(\max_{j \notin N_n^i} \|\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1}\|_{\mathbb{F}} \geq \frac{1 - \frac{\eta}{2}}{\sqrt{n^i}}\right) \leq 2 \exp\left(-C_1 \frac{n}{(n^i)^{5/2} m_n^2 k_n^2} + 2 \log(pm_n k_n)\right),$$

for some positive constant  $C_1$  that depends on  $C_{\min}$  and  $\eta$ . This completes the proof.  $\square$

**A.2. Proof of Theorem 4.1**

We begin establishing the model selection consistency given that Assumptions 4.6 and 4.7 are satisfied by the sample matrices defined in (A.21). In particular we define the event

$$\mathcal{N} = \{\Sigma_{N_n^i N_n^i}^n, \Sigma_{\xi^j N_n^i}^n \text{ satisfy Assumptions 4.6 and 4.7}\} \tag{A.34}$$

and state the following result, which is the essential step in the proof of Theorem 4.1 and will be proved in Section A.3 below.

**Proposition A.2.** *If the assumptions of Theorem 4.1 are satisfied. Then,*

$$P(\hat{N}_n^i \neq N_n^i \cap \mathcal{N}) \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right), \tag{A.35}$$

where  $C_1$  is a positive constant.

We have

$$P(\hat{N}_n^i \neq N_n^i) \leq P(\hat{N}_n^i \neq N_n^i \text{ and } \mathcal{N}) + P(\mathcal{N}^c),$$

where the first probability on the right hand side can be estimated by (A.35). Moreover, by Lemmas A.2 and A.3,

$$P(\mathcal{N}^c) \lesssim \exp\left(-C_1 \frac{n}{((n^i)^{5/4} m_n k_n)^2} + 2 \log(pm_n k_n)\right) + \exp\left(-C_1 \frac{n}{(n^i)^2 m_n k_n)^2} + 2 \log(pm_n k_n)\right),$$

and this proves Theorem 4.1.

**A.3. Proof of Proposition A.2**

We follow a similar strategy as in [1] and [14], who showed consistency of the group lasso in a reproducing kernel Hilbert space framework. First, we consider the following alternative form of the group lasso problem (3.7)

$$\widehat{PL}_i(B^i, \hat{\xi}) = \frac{1}{2n} \|\hat{\xi}^i - \tilde{\mathbf{H}}_n^T(\hat{\xi}^{-i})B^i\|_{\mathbb{F}}^2 + \frac{\lambda_n}{2} \left(\sum_{j \neq i}^p \|B^{ij}\|_{\mathbb{F}}\right)^2. \tag{A.36}$$

Because the function  $x \rightarrow x^2, x \geq 0$  is monotone, problem (A.36) leads to the same regularisation paths as problem (3.7) [see 1, , page 1187 for more details]. To derive the Karush-Kuhn-Tucker (KKT) conditions, we recall the notations (A.3), (A.4) and define the matrices

$$\hat{\Sigma}_{N_n^i \xi^j}^n = \frac{1}{n} \tilde{\mathbf{H}}_n(\hat{\xi}^{N_n^i}) \tilde{H}_n(\hat{\xi}^j) \in \mathbb{R}^{n^i k_n m_n \times k_n m_n}, \tag{A.37}$$

$$\hat{\Sigma}_{\xi^j \xi^i}^n = \frac{1}{n} \tilde{H}_n^T(\hat{\xi}^j) \hat{\xi}^i \in \mathbb{R}^{k_n m_n \times m_n} \quad (\text{A.38})$$

when  $j \neq i$  and

$$\hat{\Sigma}_{\mathbf{N}_n^i \xi^i}^n = \frac{1}{n} \tilde{\mathbf{H}}_n(\hat{\xi}^{\mathbf{N}_n^i}) \hat{\xi}^i \in \mathbb{R}^{n^i k_n m_n \times m_n}, \quad (\text{A.39})$$

where the matrices  $\tilde{H}_n(\hat{\xi}^j)$  and  $\tilde{\mathbf{H}}_n(\hat{\xi}^{\mathbf{N}_n^i})$  have been defined in (A.6) and (A.5) respectively. We also denote by  $\Sigma_{\mathbf{N}_n^i \xi^j}^n$ ,  $\Sigma_{\xi^j \xi^i}^n$ ,  $\Sigma_{\mathbf{N}_n^i \xi^i}^n$  the versions of (A.37), (A.38), (A.39) that use the true scores  $\xi_{u_r}^i$  instead of the estimated  $\hat{\xi}_{u_r}^i$  (see also equation (A.7)).

**Lemma A.4** (KKT conditions). *A matrix  $B^i = (B^{ij}, j \in \mathbb{V} \setminus \{i\}) \in \mathbb{R}^{(p-1)k_n m_n \times m_n}$  with support  $\mathbf{N}_n^i$  is optimal for problem (A.36) if and only if*

$$(\hat{\Sigma}_{\mathbf{N}_n^i \mathbf{N}_n^i}^n + \lambda_n \hat{D}_{\mathbf{N}_n^i}) B^{\mathbf{N}_n^i} - \hat{\Sigma}_{\mathbf{N}_n^i \xi^i}^n = 0, \quad \text{for all } j \in \mathbf{N}_n^i, \quad (\text{A.40a})$$

$$\|\hat{\Sigma}_{\xi^j \mathbf{N}_n^i}^n B^{\mathbf{N}_n^i} - \hat{\Sigma}_{\xi^j \xi^i}^n\|_{\mathbb{F}} \leq \lambda_n \sum_{j \neq i}^p \|B^{ij}\|_{\mathbb{F}}, \quad \text{for all } j \notin \mathbf{N}_n^i, \quad (\text{A.40b})$$

where  $\hat{\Sigma}_{\mathbf{N}_n^i \mathbf{N}_n^i}^n$  is defined in (A.7),  $B^{\mathbf{N}_n^i} = (B^{ij}, j \in \mathbf{N}_n^i) \in \mathbb{R}^{n^i k_n m_n \times m_n}$ ,  $B = (\beta_{qrk}^{ij} : 1 \leq q, r \leq m_n, 1 \leq k \leq k_n)$  and

$$\hat{D}_{\mathbf{N}_n^i} = \text{diag}((\hat{D}_{\mathbf{N}_n^i})_{jj} : j \in \hat{\mathbf{N}}_n^i)$$

is a block diagonal matrix with  $n^i$  elements  $(\hat{D}_{\mathbf{N}_n^i})_{jj} = \frac{\sum_{\ell \neq i}^p \|\hat{B}^{i\ell}\|_{\mathbb{F}}}{\|\hat{B}^{ij}\|_{\mathbb{F}}} I_{k_n m_n} \in \mathbb{R}^{k_n m_n \times k_n m_n}$ .

The idea of the proof is to first construct an estimator  $\hat{B}_{\mathbf{N}_n^i}^{\mathbf{N}_n^i}$  by minimizing the following restricted problem given the true support  $\mathbf{N}_n^i$ . That is,

$$\hat{B}_{\mathbf{N}_n^i}^{\mathbf{N}_n^i} = \text{argmin}\{\widehat{PL}_{\mathbf{N}_n^i}(B, \hat{\xi}) : B \in \mathbb{R}^{n^i k_n m_n \times m_n}\}, \quad (\text{A.41})$$

where

$$\widehat{PL}_{\mathbf{N}_n^i}(B, \hat{\xi}) = \frac{1}{2n} \|\hat{\xi}^i - \tilde{\mathbf{H}}_n^T(\hat{\xi}^{\mathbf{N}_n^i}) B\|_{\mathbb{F}}^2 + \frac{\lambda_n}{2} \left( \sum_{j \in \mathbf{N}_n^i}^p \|B^{ij}\|_{\mathbb{F}} \right)^2, \quad (\text{A.42})$$

(note that  $\widehat{PL}_{\mathbf{N}_n^i}(B, \hat{\xi})$  corresponds to the function (A.36), where we put  $B^{ij} = 0$  whenever  $j \notin \mathbf{N}_n^i$ ), and to show that the minimizer in (A.41) is “close” to the true matrix  $B_{\mathbf{N}_n^i}^{\mathbf{N}_n^i}$  defined in (4.3). To achieve this we use similar arguments as in [1] and construct another auxiliary estimator  $\tilde{B}_{\mathbf{N}_n^i}^{\mathbf{N}_n^i}$  that minimizes the restricted penalized function, where the group lasso penalty in (A.41) is replaced by an  $\ell_2$ -type penalty. More precisely,  $\tilde{B}_{\mathbf{N}_n^i}^{\mathbf{N}_n^i}$  is defined by

$$\tilde{B}_{\mathbf{N}_n^i}^{\mathbf{N}_n^i} = \text{argmin}\{\widetilde{PL}_{\mathbf{N}_n^i}(B, \hat{\xi}) : B \in \mathbb{R}^{n^i k_n m_n \times m_n}\}, \quad (\text{A.43})$$



where

$$\widetilde{PL}_{N_n^i}(B, \hat{\xi}) = \frac{1}{2n} \|\hat{\xi}^i - \tilde{\mathbf{H}}_n^\top(\hat{\xi}^{-i})B\|_F^2 + \frac{\lambda_n}{2} \left( \sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_F \right) \left( \sum_{j \in N_n^i} \frac{\|B^{ij}\|_F^2}{\|B_{m_n k_n}^{*ij}\|_F} \right).$$

We now proceed in the following steps:

- (a) In Proposition A.3 we show that the distance  $\|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_F$  is small with high probability.
- (b) In Proposition A.4 we show that  $\hat{B}_n^{N_n^i}$  is close to  $\tilde{B}_n^{N_n^i}$  with high probability.
- (c) In Proposition A.5 we use this result to derive a concentration bound for  $\|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_F$ .
- (d) We then construct the oracle minimiser  $(\hat{B}_n^{N_n^i}, \mathbf{0})$ , where  $\hat{B}_n^{N_n^i}$  is the minimiser of (A.41) and  $\mathbf{0}$  consists of  $(p - 1 - n^i)$  zero  $k_n m_n \times m_n$  matrices.
- (e) Finally, in Proposition A.6 we show that the oracle minimiser is optimal for the restricted problem (A.41) given the true support  $N_n^i$ ; that is, it satisfies (A.40b).

The minimisation problem (A.41) is convex; however, for  $p > n$ , it need not to be strictly convex, so that there may not be a unique solution. Nevertheless, the next lemma shows that the matrix  $\hat{\Sigma}_{N_n^i N_n^i}^n$  defined in (A.7) is strictly positive definite with high probability, and hence the objective function (A.41) is strictly convex, and thus  $\hat{B}_n^{N_n^i}$  is the unique optimal solution.

**Lemma A.5.** *There exists a constant  $C_1 > 0$  such that,*

$$P\left(\Lambda_{\min}(\hat{\Sigma}_{N_n^i N_n^i}^n) \geq \frac{C_{\min}}{4}\right) \gtrsim 1 - \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right).$$

*Proof.* By Weyl’s Lemma, we have  $\Lambda_{\min}(\Sigma_{N_n^i N_n^i}^n) \leq \Lambda_{\min}(\hat{\Sigma}_{N_n^i N_n^i}^n) + \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_2$ , and we get

$$P\left(\Lambda_{\min}(\hat{\Sigma}_{N_n^i N_n^i}^n) \leq \frac{C_{\min}}{4} \text{ and } \Lambda_{\min}(\Sigma_{N_n^i N_n^i}^n) > \frac{C_{\min}}{2}\right) \leq P\left(\|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_2 \geq \frac{C_{\min}}{4}\right).$$

Furthermore, using  $\delta^2 = \frac{1}{(n^i m_n k_n)^2} \frac{C_{\min}}{4}$  in Theorem A.1 with the union bound over the  $(n^i m_n k_n)^2$  index pairs of the matrix  $\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n$ , yields for some positive constant  $C_1$

$$P\left(\|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_F \geq \frac{C_{\min}}{4}\right) \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right).$$

The assertion now follows by the same arguments as given in the proof of Lemma A.2. □

**Proposition A.3.** *Suppose Assumptions 4.1-4.6 hold and the regularization parameter  $\lambda_n$  satisfies*

$$\frac{n^i m_n^{3/2}}{k_n^d} \lesssim \sqrt{\frac{2}{C_{\min}}} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}. \tag{A.44}$$

If  $\Lambda_{\min}(\hat{\Sigma}_{N_n^i N_n^i}^n) \geq \frac{C_{\min}}{4}$ , then there exists a constant  $c_2 \in (0, 1/2)$  such that, for any  $\delta > 0$  satisfying

$$\frac{2}{C_{\min}} \sqrt{n^i} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \leq c_2 \delta, \tag{A.45}$$

we have for the minimizer of (A.43)

$$P(\|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \geq \delta) \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} \delta^2}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right),$$

where  $B_n^{*N_n^i}$  is defined in (4.3) and the constant  $C_1$  satisfies  $0 < \delta \leq C_1$ .

Before we start with the proof we note that condition (A.44) refers to the spline approximation error from including only  $k_n$  terms and the second condition (A.45) represents the bias due to ridge penalisation.

For the proof we use similar arguments as given in the proof of Proposition 2 of [15]. The main change that we need to consider is the approximation error of the additive regression functions by splines. First, the minimizer  $\tilde{B}_n^{N_n^i}$  defined in (A.43) is of the form

$$\tilde{B}_n^{N_n^i} = (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \hat{\Sigma}_{N_n^i \xi^i}^n,$$

where  $D_{N_n^i}^*$  is a block diagonal matrix with  $(D_{N_n^i}^*)_{jj} = \sum_{\ell \neq i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}} / \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} I_{k_n m_n}$ ,  $j \in N_n^i$  as diagonal blocks, and the matrices  $\hat{\Sigma}_{N_n^i N_n^i}^n$  and  $\hat{\Sigma}_{N_n^i \xi^i}^n$  are defined in (A.7) and (A.37), respectively.

A simple calculation shows that

$$\|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \leq T_1 + T_2 + T_3,$$

where the terms  $T_1$ ,  $T_2$  and  $T_3$  are defined by

$$\begin{aligned} T_1 &= \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} (\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n)\|_{\mathbb{F}}, \\ T_2 &= \|\{(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} - (\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1}\} \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}}, \\ T_3 &= \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \Sigma_{N_n^i \xi^i}^n - B_n^{*N_n^i}\|_{\mathbb{F}}. \end{aligned}$$

Thus, for any  $\delta > 0$ ,  $P(\|\tilde{B}_n^i - B_n^{*N_n^i}\|_{\mathbb{F}} \geq 3\delta) \leq \sum_{i=1}^3 P(T_i \geq \delta)$ , and it is sufficient to derive bounds for the three probabilities corresponding to the random variables  $T_1$ ,  $T_2$  and  $T_3$ . Starting with  $T_1$  we have

$$T_1 \leq \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1}\|_2 \|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}} \leq \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}},$$

where we use the fact that

$$\|\hat{\Sigma}_{N_n^i N_n^i}^n\|_2 \geq \frac{C_{\min}}{4}$$

on the event  $\mathcal{N}$  and that  $(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \preceq (\hat{\Sigma}_{N_n^i N_n^i}^n)^{-1}$ . Therefore, using Lemma A.2, similar arguments as given in the proof of Theorem A.1 and applying the union bound over the  $n^i m_n^2 k_n$  pairs, we obtain

$$\begin{aligned} P(T_1 \geq \delta) &\leq P\left(\|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min} \delta}{4}\right) \\ &\lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} \delta^2}{n^i m_n^2 k_n^3} + \log(n^i m_n^2 k_n)\right), \end{aligned} \quad (\text{A.46})$$

for  $0 < \frac{\delta}{n^i m_n^2 k_n} \leq C_1$  with  $C_1 > 0$  depending on  $C_{\min}$ . To derive the bound for the probability  $P(T_2 \geq \delta)$  we use the identity  $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$  to obtain (on the event  $\mathcal{N}$ )

$$\begin{aligned} T_2 &\leq \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1}\|_2 \|(\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n)(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}} \\ &\leq \frac{4}{C_{\min}} \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \Sigma_{N_n^i \xi^i}^n\|_2. \end{aligned} \quad (\text{A.47})$$

Recall the relation (4.1) and let

$$w_{uq}^i = \sum_{j \in N_n^i} \sum_{r=1}^{m_n} (f_{qr}^{ij}(\xi_{ur}^j) - \tilde{f}_{qr}^{ij}(\xi_{ur}^j)), \quad q = 1, \dots, m_n, u = 1, \dots, n, \quad (\text{A.48})$$

where  $\tilde{f}_{qr}^{ij}$  denotes the function from Proposition A.1. Then, we can rewrite relation (4.1) in the form

$$\xi^i = \tilde{\mathbf{H}}_n^{\top}(\xi^{N_n^i}) B_n^{*N_n^i} + w^i + \epsilon^i \in \mathbb{R}^{n \times m_n},$$

where  $w^i = (w_{uq}^i)_{1 \leq u \leq n, 1 \leq q \leq m_n}$ ,  $\epsilon^i = (\epsilon_{uq}^i)_{1 \leq u \leq n, 1 \leq q \leq m_n} \in \mathbb{R}^{n \times m_n}$  and  $\tilde{\mathbf{H}}_n^{\top}(\xi^{N_n^i})$  is defined in (A.3). Furthermore, by multiplying from the left the above equation with  $\frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n}$  we obtain

$$\Sigma_{N_n^i \xi^i}^n = \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i} + \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i + \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i, \quad (\text{A.49})$$

$$\Sigma_{\xi^j \xi^i}^n = \Sigma_{\xi^j N_n^i}^n B_n^{*N_n^i} - \frac{\tilde{\mathbf{H}}_n^{\top}(\xi^j)}{n} w^i - \frac{\tilde{\mathbf{H}}_n^{\top}(\xi^j)}{n} \epsilon^i. \quad (\text{A.50})$$

where the matrix  $\Sigma_{N_n^i \xi^i}^n$  is defined in Section A.3 and  $\Sigma_{\xi^j N_n^i}^n = (\Sigma_{N_n^i \xi^j}^n)^\top$ . Using this representation and the triangle inequality we get

$$\begin{aligned} & \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \Sigma_{N_n^i \xi^i}^n\|_2 \\ & \leq \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i}\|_2 + \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i\|_2 \\ & \quad + \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i\|_2. \end{aligned}$$

As a result from this and (A.47), it follows that for all  $\delta > 0$  (on the event  $\mathcal{N}$ )

$$P(T_2 \geq \delta) \leq P\left(T_{21} \geq \frac{\delta}{3}\right) + P\left(T_{22} \geq \frac{\delta}{3}\right) + P\left(T_{23} \geq \frac{\delta}{3}\right), \quad (\text{A.51})$$

where

$$\begin{aligned} T_{21} &= \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i}\|_2, \\ T_{22} &= \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i\|_2, \\ T_{23} &= \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i\|_2. \end{aligned}$$

Next we derive upper bounds for each of the probabilities in (A.51). For  $T_{21}$  observe that

$$\begin{aligned} T_{21} &\leq \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \Sigma_{N_n^i N_n^i}^n\|_2 \|B_n^{*N_n^i}\|_2 \\ &\leq \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}}, \end{aligned}$$

where the second inequality uses the fact that  $\|B_n^{*N_n^i}\|_2 < \infty$  and that the norm  $\|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \Sigma_{N_n^i N_n^i}^n\|_2$  is bounded by one. Therefore, it follows from Theorem A.1 with  $\delta$  replaced by  $\frac{C_{\min} \delta}{6n^i m_n k_n}$  and the union bound over the  $(n^i m_n k_n)^2$  pairs that

$$\begin{aligned} P\left(T_{21} \geq \frac{\delta}{3}\right) &\leq P\left(\|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{\delta C_{\min}}{12}\right) \\ &\lesssim \exp\left(-C_2 \frac{n^{1-\alpha(2+3\beta)} \delta^2}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right), \end{aligned} \quad (\text{A.52})$$

for  $0 < \frac{\delta}{n^i m_n k_n} \leq C_2$ , for  $C_2 > 0$  depending on  $C_{\min}$ .

For the term  $T_{22}$  note that

$$\|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^{*i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}}\|_2 \leq \|(\Sigma_{N_n^i N_n^i}^n)^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}}\|_2 = \Lambda_{\min}(\Sigma_{N_n^i N_n^i}^n)^{-1/2}, \quad (\text{A.53})$$

where we used Lemma A.2 for the last inequality with  $\delta = \frac{C_{\min}}{2}$ . Thus, (on the event  $\mathcal{N}$ ) the term  $T_{22}$  can be bounded by

$$T_{22} \leq \frac{2^{5/2}}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \left\| \frac{w^i}{\sqrt{n}} \right\|_{\mathbb{F}}.$$

Recall the notation of  $w_{uq}^i$  in (A.48) and the definition of the event  $\Omega$  in Proposition A.1. Then, if the event  $\Omega$  holds, we have,

$$\begin{aligned} \left\| \frac{1}{\sqrt{n}} w^i \right\|_{\mathbb{F}}^2 &= \frac{1}{n} \sum_{u=1}^n \sum_{q=1}^{m_n} (w_{uq}^i)^2 = \frac{1}{n} \sum_{u=1}^n \sum_{q=1}^{m_n} \left( \sum_{j \in N^i} \sum_{r=1}^{m_n} (f_{qr}^{ij}(\xi_{ur}^j) - \tilde{f}_{nqr}^{ij}(\xi_{ur}^j)) \right)^2, \\ &\leq \frac{n^i m_n}{n} \sum_{j \in N^i} \sum_{q=1}^{m_n} \sum_{r=1}^{m_n} \sum_{u=1}^n (f_{qr}^{ij}(\xi_{ur}^j) - \tilde{f}_{nqr}^{ij}(\xi_{ur}^j))^2 \\ &\leq \frac{n^i m_n}{n} \sum_{j \in N^i} \sum_{q=1}^{m_n} \sum_{r=1}^{m_n} \max_{j \in N^i} \max_{1 \leq q, r \leq m_n} \|f_{qr}^{ij} - \tilde{f}_{qr}^{ij}\|_2 \leq c_1 (n^i)^2 m_n^3 k_n^{-2d}, \end{aligned} \tag{A.54}$$

and by assumption (A.44) it follows that on the event  $\Omega$

$$\left\| \frac{1}{\sqrt{n}} w^i \right\|_{\mathbb{F}} \leq \sqrt{c_1} \sqrt{\frac{2}{C_{\min}}} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}. \tag{A.55}$$

As a result,

$$\begin{aligned} P\left(T_{22} \geq \frac{\delta}{3}\right) &\leq P\left(\left(\frac{8}{C_{\min}^{3/2}}\right)^2 \sqrt{c_1} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{\delta}{3}\right) + P(\Omega^c) \\ &\lesssim \exp\left(-C_3 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^{-2} \delta^2}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_3 \frac{n}{n^i m_n^2 k_n^{2d}} + \log(n^i m_n^2)\right) \end{aligned} \tag{A.56}$$

for  $0 < \delta (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^{-1} \leq C_3$ , for  $C_3 > 0$  depending on  $C_{\min}$  and  $c_1$ , where we have used Theorem A.1 and Proposition A.1.

We next derive an upper bound for the probability corresponding to the term  $T_{23}$  in (A.51) noting that (on the event  $\mathcal{N}$ )

$$T_{23} \leq \frac{8}{C_{\min}^2} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \left\| \frac{\tilde{\mathbf{H}}_n(\xi_{N_n^i}^i)}{n} \epsilon^i \right\|_{\mathbb{F}}. \tag{A.57}$$

The  $(i, j)$  element of the matrix  $\frac{\tilde{\mathbf{H}}_n(\xi_{N_n^i}^i)}{n} \epsilon^i$  can be written as an i.i.d sum of the form  $\frac{1}{n} \sum_{u=1}^n \tilde{h}_{nk}(\xi_{ur}^j) \epsilon_{uq}^i$ . Thus, by Assumption 4.4 it follows that

$$P\left(\left| \frac{1}{n} \sum_{u=1}^n \tilde{h}_{nk}(\xi_{ur}^j) \epsilon_{uq}^i \right| \geq \epsilon\right) \leq 2 \exp(-C_5 n \epsilon^2),$$

for any  $\epsilon > 0$ . Therefore, by applying the union bound over the  $n^i m_n^2 k_n$  gives

$$P\left(\left\|\frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n}\epsilon^i\right\|_{\mathbb{F}} \geq \epsilon\right) \leq 2 \exp\left(-C_5 \frac{n\epsilon^2}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right). \quad (\text{A.58})$$

Using this inequality with  $\epsilon = C_{\min}/6$ , (A.52) and (A.57) gives

$$\begin{aligned} P\left(T_{23} \geq \frac{\delta}{3}\right) &\lesssim \exp\left(-C_4 \frac{n^{1-\alpha(2+3\beta)}\delta^2}{(n^i)^2 m_n^2 k_n^4} + 2\log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_5 \frac{n}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right), \end{aligned} \quad (\text{A.59})$$

for  $0 < \frac{\delta}{n^i m_n k_n} \leq C_4$ , for  $C_4 > 0$  depending on  $C_{\min}$ . Therefore from (A.52), (A.56) and (A.59) it follows that

$$\begin{aligned} P(T_2 \geq \delta) &\lesssim \exp\left(-C_2 \frac{n^{1-\alpha(2+3\beta)}\delta^2}{(n^i)^2 m_n^2 k_n^4} + 2\log(n^i m_n k_n)\right) \\ &\quad + 2 \exp\left(-C_3 \frac{n}{n^i m_n^2 k_n^{2d}} + \log(n^i m_n^2)\right), \end{aligned} \quad (\text{A.60})$$

for  $0 < \frac{\delta}{n^i m_n k_n} \leq C_2$ , where the first term dominates the second one because of Assumption 4.5.

Finally, we derive an upper bound for the probability involving  $T_3$ . Using representation (A.49) we obtain

$$T_3 \leq T_{31} + T_{32} + T_{33}, \quad (\text{A.61})$$

where

$$\begin{aligned} T_{31} &= \left\|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i} - B_n^{*N_n^i}\right\|_{\mathbb{F}}, \\ T_{32} &= \left\|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i\right\|_{\mathbb{F}}, \\ T_{33} &= \left\|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i\right\|_{\mathbb{F}}. \end{aligned}$$

For the first term  $T_{31}$  (on the event  $\mathcal{N}$ )

$$\begin{aligned} T_{31} &= \lambda_n \left\|(\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} D_{N_n^i}^* B_n^{*N_n^i}\right\|_{\mathbb{F}} \\ &\leq \frac{2}{C_{\min}} \lambda_n (n^i)^{1/2} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \leq \frac{\delta}{2(1 + \sqrt{c_1})}, \end{aligned}$$

where we used condition (A.45) with  $\sqrt{c_1} = \frac{1}{2c_2} - 1$  and the fact that  $\left\|\text{diag}\left(\frac{B_{m_n k_n}^{*ij}}{\|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}}\right) : j \in N_n^i\right\|_{\mathbb{F}} = (n^i)^{1/2}$ . Moreover, by applying the same arguments for deriving the bound of  $T_{22}$  and by using (A.53), conditions (A.44)

and (A.45) it follows that on the event  $\Omega$

$$T_{32} \leq \sqrt{c_1} \left( \frac{2}{C_{\min}} \right)^{1/2} \frac{n^i m_n^{3/2}}{k_n^d} \leq \sqrt{c_1} \frac{2}{C_{\min}} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \leq \frac{\sqrt{c_1}}{2(1 + \sqrt{c_1})} \delta.$$

For  $T_{33}$ , we have

$$T_{33} \leq \frac{2}{C_{\min}} \left\| \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i}) \epsilon^i}{n} \right\|_{\mathbb{F}}.$$

Therefore, inequalities (A.61) and (A.58) imply that for all  $\delta > 0$

$$\begin{aligned} P(T_3 \geq \delta) &\leq P\left(\frac{2}{C_{\min}} \left\| \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i}) \epsilon^i}{n} \right\|_{\mathbb{F}} \geq \frac{\delta}{2}\right) + P(\Omega^c) \\ &\leq 2 \exp\left(-C_3 \frac{n \delta^2}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right) + 2 \exp\left(-C_3 \frac{n}{n^i m_n^2 k_n^{2d}} + \log(n^i m_n^2)\right), \end{aligned} \tag{A.62}$$

for  $C_3 > 0$  depending on  $C_{\min}$ . Thus, by (A.46), (A.60) and (A.62), we have shown, for any  $\delta > 0$  such that  $0 < \delta \leq C_1$  and  $0 < \delta \leq C_2$

$$\begin{aligned} P(\|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \geq \delta) &\lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} \delta^2}{n^i m_n^2 k_n^3} + \log(n^i m_n^2 k_n)\right) \\ &\quad + \exp\left(-C_2 \frac{n^{1-\alpha(2+3\beta)} \delta^2}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_3 \frac{n \delta^2}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right). \end{aligned}$$

Since the second term dominates the first and the third, the assertion in Proposition A.3 follows.

The next proposition brings  $\hat{B}_n^{N_n^i} = (\hat{B}_n^{ij}, j \in N_n^i)$  close to  $\tilde{B}_n^{N_n^i} = (\tilde{B}_n^{ij}, j \in N_n^i)$ , from which we can establish the concentration inequality for  $\hat{B}_n^{N_n^i}$ .

**Proposition A.4.** *Let  $\hat{B}_n^{N_n^i}$  be the minimiser of (A.41) and  $\tilde{B}_n^{N_n^i}$  be the minimiser of (A.43). If  $\Lambda_{\min}(\hat{\Sigma}_{N_n^i N_n^i}^n) \geq \frac{C_{\min}}{4}$  then,*

$$\|\hat{B}_n^{N_n^i} - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} \leq \frac{10}{C_{\min}} \lambda_n n^i (b_n^{*i})^{-1} \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}},$$

where  $b_n^{*i} = \min_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}$ .

*Proof.* The idea of the proof is similar as in the proof of Proposition 3 in [15]. Consider the sphere  $\mathcal{S}_n(\delta_n) = \{B \in \mathbb{R}^{n^i k_n m_n \times m_n} : \|B - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} = \delta_n\}$ , where  $(\delta_n)_{n \in \mathbb{N}}$  is a positive sequence of real numbers. For  $\epsilon \in [0, 1]$  let

$$f(\epsilon) = \widehat{PL}_{N_n^i}(\tilde{B}_n^{N_n^i} + \epsilon A, \hat{\xi}^i),$$

where the function  $\widehat{PL}_{N_n^i}$  is defined in (A.42) and  $A = B - \tilde{B}_n^{N_n^i}$ . A straightforward calculation gives for the first and the second derivatives of the function  $f(\epsilon)$

$$\begin{aligned} \dot{F}(\epsilon) &= -\langle \hat{\Sigma}_{N_n^i \xi^i}^n, A \rangle_{\mathbb{F}} + \langle A, \hat{\Sigma}_{N_n^i N_n^i}^n (\tilde{B}_n^{N_n^i} + \epsilon A) \rangle_{\mathbb{F}} \\ &\quad + \lambda_n \sum_{j \in N_n^i} \|\tilde{B}_n^{ij} + \epsilon A^j\|_{\mathbb{F}} \sum_{k \in N_n^i} (\langle \tilde{B}_n^{ik} + \epsilon A^k, A^k \rangle_{\mathbb{F}} \|\tilde{B}_n^{ik} + \epsilon A^k\|_{\mathbb{F}}^{-1}). \\ \ddot{F}(\epsilon) &= \langle A, \hat{\Sigma}_{N_n^i N_n^i}^n A \rangle_{\mathbb{F}} + \lambda_n \left( \sum_{j \in N_n^i} \|\tilde{B}_n^{ij} + \epsilon A^j\|_{\mathbb{F}}^{-1} \langle \tilde{B}_n^{ij} + \epsilon A^j, A^j \rangle_{\mathbb{F}} \right)^2 \\ &\quad + \lambda_n \sum_{j \in N_n^i} \|\tilde{B}_n^{ij} + \epsilon A^j\|_{\mathbb{F}} \sum_{k \in N_n^i} \|A^k\|_{\mathbb{F}}^2 \|\tilde{B}_n^{ik} + \epsilon A^k\|_{\mathbb{F}}^{-1} \\ &\quad - \lambda_n \sum_{j \in N_n^i} \|\tilde{B}_n^{ij} + \epsilon A^j\|_{\mathbb{F}} \sum_{k \in N_n^i} (\langle \tilde{B}_n^{ik} + \epsilon A^k, A^k \rangle_{\mathbb{F}})^2 \|\tilde{B}_n^{ik} + \epsilon A^k\|_{\mathbb{F}}^{-3}. \end{aligned}$$

where  $\langle \cdot, \cdot \rangle_{\mathbb{F}}$  denotes the Frobenious inner product and  $\hat{\Sigma}_{N_n^i \xi^i}^n, \hat{\Sigma}_{N_n^i N_n^i}^n$  are defined in (A.39) and (A.7) respectively. By construction,  $f(0) = \widehat{PL}_{N_n^i}(\tilde{B}_n^{N_n^i}, \hat{\xi}^i)$ ,  $f(1) = \widehat{PL}_{N_n^i}(B, \hat{\xi}^i)$  and by Taylor's theorem, we have for some  $\epsilon \in (0, 1)$

$$\widehat{PL}_{N_n^i}(B, \hat{\xi}^i) - \widehat{PL}_{N_n^i}(\tilde{B}_n^{N_n^i}, \hat{\xi}^i) = f(1) - f(0) = \dot{f}(0) + \frac{\ddot{f}(\epsilon)}{2}. \quad (\text{A.63})$$

The Cauchy-Schwarz inequality yields for any  $B \in \mathcal{S}_n(\delta_n)$  and  $\epsilon \in [0, 1]$

$$\ddot{f}(\epsilon) \geq \langle A, \hat{\Sigma}_{N_n^i N_n^i}^n A \rangle_{\mathbb{F}} + \lambda_n \left( \sum_{j \in N_n^i} \|\tilde{B}_n^{ij} + \epsilon A^j\|_{\mathbb{F}}^{-1} \langle \tilde{B}_n^{ij} + \epsilon A^j, A^j \rangle_{\mathbb{F}} \right)^2 \geq \frac{C_{\min}}{4} \delta_n^2. \quad (\text{A.64})$$

On the other hand, by Lemma A7 in [14] it follows that

$$\begin{aligned} |\dot{f}(0)| &\leq \lambda_n \sum_{j \in N_n^i} \sum_{k \in N_n^i} \left[ \|\tilde{B}_n^{ij} - B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{ik} - B^k\|_{\mathbb{F}} \right. \\ &\quad \left. + \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|B_{m_n k_n}^{*ik}\|_{\mathbb{F}}^{-1} \|\tilde{B}_n^{ij} - B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{ik} - B^k\|_{\mathbb{F}} \right]. \end{aligned}$$

A further application of the Cauchy-Schwarz inequality gives

$$\begin{aligned} |\dot{f}(0)| &\leq \lambda_n \sqrt{n^i \sum_{j \in N_n^i} \|\tilde{B}_n^{ij} - B_{m_n k_n}^{*ij}\|_{\mathbb{F}}^2 n^i \sum_{k \in N_n^i} \|\tilde{B}_n^{ik} - B^k\|_{\mathbb{F}}^2} \\ &\quad + \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{ij} - B_{m_n k_n}^{*ij}\|_{\mathbb{F}} (b_n^{*i})^{-1} \sum_{k \in N_n^i} \|\tilde{B}_n^{ik} - B^k\|_{\mathbb{F}} \\ &= \lambda_n n^i \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B\|_{\mathbb{F}} \\ &\quad + \lambda_n (b_n^{*i})^{-1} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{ij} - B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \sum_{k \in N_n^i} \|\tilde{B}_n^{ik} - B^k\|_{\mathbb{F}} \end{aligned}$$



$$\begin{aligned} &\leq \lambda_n n^i \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B\|_{\mathbb{F}} \\ &+ \lambda_n \sqrt{n^i} (b_n^{*i})^{-1} \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B\|_{\mathbb{F}} \|B_n^{*N_n^i}\|_{\mathbb{F}}. \end{aligned}$$

Using the fact  $b_n^{*i} \leq \|B_n^{*N_n^i}\|_{\mathbb{F}} \leq \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}$  we obtain

$$|\dot{f}(0)| \leq 2\lambda_n n^i (b_n^{*i})^{-1} \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B\|_{\mathbb{F}}. \quad (\text{A.65})$$

Hence, combining (A.63), (A.64) and (A.65), we obtain

$$\begin{aligned} \widehat{PL}_{N_n^i}(B, \hat{\xi}^i) - \widehat{PL}_{N_n^i}(\tilde{B}_n^{N_n^i}, \hat{\xi}^i) &\geq -2\lambda_n n^i (b_n^{*i})^{-1} \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \\ &+ \frac{C_{\min}}{4} \delta_n^2. \end{aligned}$$

If we choose  $\delta_n^2 = \frac{10}{C_{\min}} \lambda_n n^i (b_n^{*i})^{-1} \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}}$ , it follows that

$$\widehat{PL}_{N_n^i}(B, \hat{\xi}^i) - \widehat{PL}_{N_n^i}(\tilde{B}_n^{N_n^i}, \hat{\xi}^i) > 0.$$

Since the function  $\widehat{PL}_{N_n^i}(B, \hat{\xi}^i)$  is convex, the minimizer  $\hat{B}_n^{N_n^i}$  of  $\widehat{PL}_{N_n^i}(B, \hat{\xi}^i)$  is going to be inside the sphere defined by  $\mathcal{S}_n(\delta_n)$ , that is,

$$\|\hat{B}_n^{N_n^i} - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} \leq \frac{10}{C_{\min}} \lambda_n n^i (b_n^{*i})^{-1} \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \|\tilde{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}}. \quad \square$$

Using the Propositions A.3 and A.4, we now can establish the concentration bounds for  $\|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}}$ .

**Proposition A.5.** *Suppose Assumptions of Proposition A.3 are satisfied and that  $\delta$  satisfies*

$$\frac{2}{C_{\min}} \lambda_n (n^i)^{3/2} \left( \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \right)^2 \leq c_2 b_n^{*i} \delta, \quad (\text{A.66})$$

for some constant  $c_2 > 0$ . Then,

$$\begin{aligned} &P\left(\|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \geq \delta\right) \\ &\lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^2 \delta^2}{(n^i)^4 m_n^2 k_n^4 \left(\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right)^2} + 2 \log(n^i m_n k_n)\right), \end{aligned}$$

where  $C_1 > 0$  such that  $0 < \delta \leq C_1$ .

*Proof.* By Proposition A.4 and the triangle inequality,

$$\begin{aligned} \|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} &\leq \|\hat{B}_n^{N_n^i} - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} + \|B_n^{*N_n^i} - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} \\ &\leq (b_n^{*i})^{-1} \|B_n^{*N_n^i} - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} \left( \frac{10}{C_{\min}} \lambda_n n^i \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} + b_n^{*i} \right) \\ &\lesssim n^i (b_n^{*i})^{-1} \|B_n^{*N_n^i} - \tilde{B}_n^{N_n^i}\|_{\mathbb{F}} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}, \end{aligned}$$

where we have used the fact that  $b_n^{*i} \leq \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}$  and  $\lambda_n \lesssim 1$ . The assertion now follows from Proposition A.3 with  $\delta$  replaced by  $b_n^{*i} \delta (n^i \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^{-1}$ .  $\square$

Let  $\hat{B}_n^{N_n^i}$  be the minimizer of the restricted problem (A.41). By construction, the estimator  $(\hat{B}_n^{N_n^i}, \mathbf{0})$  obtained from  $\hat{B}_n^{N_n^i}$  by adding blocks with 0 elements whenever  $j \notin N_n^i$ , satisfies the first KKT-condition (A.40a). To prove that  $(\hat{B}_n^{N_n^i}, \mathbf{0})$  is, with high probability, optimal for problem (A.41), it is therefore sufficient to show that the second KKT-condition (A.40b) is satisfied. This is the statement of the following proposition.

**Proposition A.6.** *The matrix  $(\hat{B}_n^{N_n^i}, \mathbf{0})$  satisfies (A.40b) with high probability, in the sense that*

$$\begin{aligned} P(\max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n \hat{B}_n^{N_n^i} - \hat{\Sigma}_{\xi^j \xi^i}^n\|_{\mathbb{F}} \geq \lambda_n \sum_{j \neq i}^p \|\hat{B}_n^{ij}\|_{\mathbb{F}}) \\ \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right), \end{aligned}$$

where  $C_1$  is a positive constant.

*Proof.* The idea of the proof is similar as in the proof of Proposition 4 in [15]. By the first optimality condition (A.40a), we have for all  $j \in N_n^i$ ,

$$\hat{B}_n^{N_n^i} = (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \hat{\Sigma}_{N_n^i \xi^i}^n, \quad (\text{A.67})$$

where  $\hat{D}_{N_n^i}$  is defined in Lemma A.4. Using (A.67) in the expression at the left-hand side of condition (A.40b) gives

$$\hat{\Sigma}_{\xi^j N_n^i}^n \hat{B}_n^{N_n^i} - \hat{\Sigma}_{\xi^j \xi^i}^n = \hat{\Sigma}_{\xi^j N_n^i}^n (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \hat{\Sigma}_{N_n^i \xi^i}^n - \hat{\Sigma}_{\xi^j \xi^i}^n = R_1^j + \dots + R_7^j,$$

where

$$\begin{aligned} R_1^j &= (\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n) (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \Sigma_{N_n^i \xi^i}^n \\ R_2^j &= \Sigma_{\xi^j N_n^i}^n (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} (\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n) \end{aligned}$$

$$\begin{aligned}
R_3^j &= \Sigma_{\xi^j N_n^i}^n \{ (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} - (\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \} \Sigma_{N_n^i \xi^i}^n \\
R_4^j &= (\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n) (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} (\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n) \\
R_5^j &= \Sigma_{\xi^j \xi^i}^n - \hat{\Sigma}_{\xi^j \xi^i}^n \\
R_6^j &= \Sigma_{\xi^j N_n^i}^n \{ (\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} - (\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \} \Sigma_{N_n^i \xi^i}^n \\
R_7^j &= \Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*)^{-1} \Sigma_{N_n^i \xi^i}^n - \Sigma_{\xi^j \xi^i}^n.
\end{aligned}$$

In the following we derive bounds for the probabilities

$$P\left(\max_{j \notin N_n^i} \|R_r^j\|_{\mathbb{F}} \geq \frac{\lambda_n}{7} \sum_{j \in N_n^i} \|\hat{B}_n^{ij}\|_{\mathbb{F}}\right) \quad (\text{A.68})$$

( $r = 1, \dots, 7$ ). For this purpose we proceed in two steps.

**Step 1:** First, we define the event,  $\mathcal{A}_0$ , that there exists a constant  $0 < c_0 < 1$ , such that

$$\mathcal{A}_0 = \left\{ \|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \leq c_0 (n^i)^{-1/2} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \right\}.$$

Then, by Proposition A.5 with  $\delta = c_0 (n^i)^{-1/2} \sum_{j \in N_n^i} \|B_{m_n k_n}^*\|_{\mathbb{F}}$  we have

$$\begin{aligned}
P(\mathcal{A}_0^c) &\lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^2}{(n^i)^5 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\
&\lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right),
\end{aligned} \quad (\text{A.69})$$

for some constant  $C_1 > 0$  where for the second inequality we used the fact  $\frac{2}{C_{\min}} \lambda_n (n^i)^2 \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \lesssim b_n^{*i}$  which follows from condition A.66 in Proposition A.5.

Now, on the event  $\mathcal{A}_0$ , it follows by the Cauchy-Schwarz inequality that

$$\left| \sum_{j \in N_n^i} \|\hat{B}_n^{ij}\|_{\mathbb{F}} - \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \right| \leq c_0 \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}.$$

Therefore, we have for  $r = 1, \dots, 7$

$$\begin{aligned}
P\left(\max_{j \notin N_n^i} \|R_r^j\|_{\mathbb{F}} \geq \frac{\lambda_n}{7} \sum_{j \in N_n^i} \|\hat{B}_n^{ij}\|_{\mathbb{F}}\right) &\leq P\left(\max_{j \notin N_n^i} \|R_r^j\|_{\mathbb{F}} \geq \frac{\lambda_n(1-c_0)}{7} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\
&\quad + P(\mathcal{A}_0^c),
\end{aligned}$$

where the second probability can be bounded by (A.69). Thus, in order to control the probabilities of the terms  $\max_{j \notin N_n^i} \|R_r^j\|_{\mathbb{F}} \geq \frac{\lambda_n}{7} \sum_{j \in N_n^i} \|\hat{B}_n^{ij}\|_{\mathbb{F}}$  it suffices to

derive the probabilities  $P(\max_{j \notin N_n^i} \|R_r^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})$  for all  $r = 1, \dots, 7$ , where  $c_1 = \frac{1-c_0}{7}$ .

**Step 2:** Term  $R_1^j$ . Substituting the representation (A.49) used in the proof of Proposition A.3 we can rewrite  $R_1^j$  as

$$R_1^j = (\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n)(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \left( \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i} + \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i + \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i \right),$$

and by the union bound it follows that

$$P\left(\max_{j \notin N_n^i} \|R_1^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \leq \sum_{r=1}^3 P\left(\max_{j \notin N_n^i} \|R_{1r}^j\|_{\mathbb{F}} \geq \frac{c_1 \lambda_n}{3} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right), \quad (\text{A.70})$$

where  $R_{11}^j, R_{12}^j, R_{13}^j$  are defined in an obvious manner satisfying

$$\begin{aligned} \|R_{11}^j\|_{\mathbb{F}} &\leq \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i}\|_2. \\ \|R_{12}^j\|_{\mathbb{F}} &\leq \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i\|_2. \\ \|R_{13}^j\|_{\mathbb{F}} &\leq \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i\|_2. \end{aligned}$$

Next we derive bounds for the probabilities on the right hand side of (A.70) starting with the term  $\|R_{11}^j\|_{\mathbb{F}}$ . Observing that

$$\begin{aligned} (\hat{\Sigma}_{N_n^i N_n^i}^n - \lambda_n \hat{D}_{N_n^i})^{-1} \Sigma_{N_n^i N_n^i}^n &= (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} (\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n) \\ &\quad + (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \hat{\Sigma}_{N_n^i N_n^i}^n. \end{aligned}$$

We have

$$\|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \Sigma_{N_n^i N_n^i}^n\|_2 \leq \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} (\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n)\|_2 \quad (\text{A.71})$$

$$\begin{aligned} &+ \|(\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \hat{\Sigma}_{N_n^i N_n^i}^n\|_2 \\ &\leq \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} + 1, \quad (\text{A.72}) \end{aligned}$$

where we used the fact that  $\hat{\Sigma}_{N_n^i N_n^i}^n \preceq \hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n I_{N_n^i} \preceq \hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i}$ . From this, it follows that

$$\max_{j \notin N_n^i} \|R_{11}^j\|_{\mathbb{F}} \lesssim \max_{j \notin N_n^i} \{\|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}}\} \left( \frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} + 1 \right),$$

where we use the fact that  $\|B_n^{*N_n^i}\|_2$  is bounded. Consider the event

$$\mathcal{A}_1 = \left\{ \max_{j \notin N_n^i} \{\|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}}\} \leq \frac{c_1 \lambda_n \sum_{l \in N_n^i} \|B_{m_n k_n}^{*il}\|_{\mathbb{F}}}{6} \right\}.$$

Then,

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|R_{11}^j\|_{\mathbb{F}} \geq \frac{c_1 \lambda_n}{3} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) &\leq P\left(\|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) + P(\mathcal{A}_1^c), \\ &\lesssim \exp\left(-C_2 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &+ \exp\left(-C_2 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{l \in N_n^i} \|B_{m_n k_n}^{*il}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + \log(n^i m_n^2 k_n^2) + \log((p-n^i) m_n^2 k_n^2)\right), \end{aligned}$$

where we have used Theorem A.1 and the union bound and  $C_2$  is a positive constant depending on  $C_{\min}$ .

For the term  $R_{12}^j$ , we use the same arguments as for the term  $T_2$  in the proof of Proposition A.3. Specifically, recall the definition of the event  $\Omega$  in (A.20) and the calculation in (A.53) and (A.54) to obtain on the event  $\Omega$

$$\begin{aligned} \max_{j \notin N_n^i} \|R_{12}^j\|_{\mathbb{F}} &\lesssim \max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \left(\frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} + 1\right) \\ &\quad \times \left(\frac{2}{C_{\min}}\right)^{3/2} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}, \end{aligned}$$

where we used condition (4.9) for the last inequality.

Then, conditioning on the event  $\mathcal{A}_2 = \{\max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \geq \frac{c_1}{6} (\frac{C_{\min}}{2})^{3/2}\}$ , we have by Proposition A.1

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|R_{12}^j\|_{\mathbb{F}} \geq \frac{c_1 \lambda_n}{3} \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) &\leq P\left(\|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) + P\left(\max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \geq \frac{c_1}{6} (\frac{C_{\min}}{2})^{3/2}\right) \\ &+ P(\Omega^c) \\ &\lesssim \exp\left(-C_3 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &+ \exp\left(-C_3 \frac{n^{1-\alpha(2+3\beta)}}{n^i m_n^2 k_n^4} + \log(n^i m_n^2 k_n^2) + \log((p-n^i) m_n^2 k_n^2)\right) \\ &+ \exp\left(-C_3 \frac{n}{n^i m_n^2 k_n^{2d}} + \log(n^i m_n^2)\right), \end{aligned}$$

for some  $C_3 > 0$  depending on  $C_{\min}$  and  $c_1$ . For the term  $R_{13}^j$ , we have

$$\max_{j \notin N_n^i} \|R_{13}^j\|_{\mathbb{F}} \leq \frac{4}{C_{\min}} \max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \left\| \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i \right\|_{\mathbb{F}},$$

which yields

$$\begin{aligned}
& P\left(\max_{j \notin N_n^i} \|R_{13}^j\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\
& \leq P\left(\max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n - \Sigma_{\xi^j N_n^i}^n\|_{\mathbb{F}} \geq \frac{c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}}{3}\right) + P\left(\left\|\frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n}\right\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) \\
& \lesssim \exp\left(-C_4 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + \log(n^i m_n^2 k_n^2) + \log((p-n^i) m_n^2 k_n^2)\right) \\
& \quad + \exp\left(-C_4 \frac{n}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right),
\end{aligned}$$

where we used Theorem A.1 and (A.58) for some  $C_4 > 0$  that depends on  $C_{\min}$  and  $c_1$ . Combining together the results for the terms  $\|R_{13}^j\|_{\mathbb{F}}$ ,  $\|R_{12}^j\|_{\mathbb{F}}$  and  $\|R_{13}^j\|_{\mathbb{F}}$  we conclude that

$$\begin{aligned}
& P\left(\max_{j \notin N_n^i} \|R_1^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\
& \lesssim \exp\left(-C_2 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{l \in N_n^i} \|B_{m_n k_n}^{*il}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + \log(p m_n^2 k_n^2)\right),
\end{aligned}$$

since  $p - n^i \leq p$  and  $n^i \leq p$ .

*Term  $R_2^j$ .* First we write

$$\begin{aligned}
\max_{j \notin N_n^i} \|R_2^j\|_{\mathbb{F}} & \leq \max_{j \notin N_n^i} \left\{ \|\Sigma_{\xi^j N_n^i}^n (\Sigma_{N_n^i N_n^i}^n)^{-1}\|_{\mathbb{F}} \right\} \|\Sigma_{N_n^i N_n^i}^n (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i}^n)^{-1}\|_2 \|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}} \\
& \leq \frac{1-\eta}{\sqrt{n^i}} \|\Sigma_{N_n^i N_n^i}^n (\hat{\Sigma}_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i}^n)^{-1}\|_2 \|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}} \\
& \leq \frac{1-\eta}{\sqrt{n^i}} \left(\frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} + 1\right) \|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}},
\end{aligned}$$

where we used (A.71) and the second inequality holds with high probability by Lemma A.3. As a result, it follows that

$$\begin{aligned}
& P\left(\max_{j \notin N_n^i} \{\|R_2^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\}\right) \\
& \leq P\left(\frac{1-\eta}{\sqrt{n^i}} \left(\frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} + 1\right) \|\hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\
& \leq P\left(\|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) + P(\mathcal{A}_2^b) \\
& \lesssim \exp\left(-C_3 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\
& \quad + \exp\left(-C_3 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{m_n^2 k_n^3} + \log(n^i m_n^2 k_n)\right),
\end{aligned}$$

where

$$\mathcal{A}_2 = \left\{ \left\| \hat{\Sigma}_{N_n^i \xi^i}^n - \Sigma_{N_n^i \xi^i}^n \right\|_{\mathbb{F}} \leq \frac{c_1}{2} \frac{\sqrt{n^i}}{1 - \eta} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \right\},$$

and we used Theorem A.1 and the union bound for the last inequality for  $C_3 > 0$  depending on  $C_{\min}$ ,  $\eta$  and  $c_1$ .

*Term  $R_3^j$ .* Using the identity  $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$  and following similar arguments used to obtain bounds for the terms  $R_1$  and  $R_2$ , we get (note that we are working on the event  $\mathcal{N}$ )

$$P\left(\max_{j \notin N_n^i} \|R_3^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \leq \sum_{r=1}^3 P\left(\|R_{3r}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right),$$

where,

$$\begin{aligned} \|R_{31}\|_{\mathbb{F}} &= \frac{1 - \eta}{\sqrt{n^i}} \left(1 + \frac{4}{C_{\min}} \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}}\right) \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \\ &\quad \times \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \Sigma_{N_n^i N_n^i}^n B_n^{*N_n^i}\|_2, \\ \|R_{32}\|_{\mathbb{F}} &= \frac{1 - \eta}{\sqrt{n^i}} \left(1 + \frac{4}{C_{\min}} \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}}\right) \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \\ &\quad \times \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \tilde{\mathbf{H}}_n(\xi^{N_n^i}) w^i\|_{\mathbb{F}}, \\ \|R_{33}\|_{\mathbb{F}} &= \frac{1 - \eta}{\sqrt{n^i}} \left(1 + \frac{4}{C_{\min}} \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}}\right) \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \\ &\quad \times \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \tilde{\mathbf{H}}_n(\xi^{N_n^i}) \epsilon^i\|_{\mathbb{F}}, \end{aligned}$$

(note that the terms are independent of  $j$ ). We next derive bounds for the probabilities  $P\left(\|R_{3r}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right)$  for  $r = 1, 2, 3$ . For the term  $\|R_{31}\|_{\mathbb{F}}$ , note that the third term is finite. Hence,

$$\|R_{31}\|_{\mathbb{F}} \lesssim \frac{1 - \eta}{\sqrt{n^i}} \|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \left(\frac{4}{C_{\min}} \|\hat{\Sigma}_{N_n^i N_n^i}^n - \Sigma_{N_n^i N_n^i}^n\|_{\mathbb{F}} + 1\right),$$

which implies (using the same arguments as before)

$$P\left(\max_{j \notin N_n^i} \|R_{31}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \tag{A.73}$$

$$\leq P\left(\|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{\sqrt{n^i} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} c_1}{1 - \eta} \frac{c_1}{6}\right) \tag{A.74}$$

$$+ P\left(\|\Sigma_{N_n^i N_n^i}^n - \hat{\Sigma}_{N_n^i N_n^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) \tag{A.75}$$

$$\begin{aligned} &\lesssim \exp\left(-C_4 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_4 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right), \end{aligned} \quad (\text{A.76})$$

for some positive constant  $C_4$  that depends on  $\eta$  and  $C_{\min}$ . Similarly, for the term  $\|R_{32}\|_{\mathbb{F}}$  have

$$\|R_{32}\|_{\mathbb{F}} \leq \frac{1-\eta}{\sqrt{n^i}} \|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} \left(\frac{4}{C_{\min}} \|\hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} + 1\right) \|\frac{w^i}{\sqrt{n}}\|_{\mathbb{F}}.$$

Hence, recalling the definition of the set  $\Omega$  in (A.20) and using conditions (4.9) and (A.55), Proposition A.1 and Theorem A.1, we obtain for some positive constant

$$\begin{aligned} &P\left(\|R_{32}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ &\leq P\left(\|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} \geq \frac{\sqrt{n^i} c_1}{12c_2(1-\eta)}\right) + P\left(\|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) + P(\Omega^c) \\ &\lesssim \exp\left(-C_5 \frac{n^{1-\alpha(2+3\beta)}}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) + \exp\left(-C_5 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp(-C_3 \frac{n}{n^i m_n^2 k_n^{2d}} + \log(n^i m_n^2)), \end{aligned}$$

for some positive constant  $C_5$  that depends on  $\eta$  and  $C_{\min}$ . For the term  $\|R_{33}\|_{\mathbb{F}}$ , we write

$$\|R_{33}\|_{\mathbb{F}} \leq \frac{1-\eta}{\sqrt{n^i}} \left(\frac{2}{C_{\min}}\right)^{1/2} \left(1 + \frac{4}{C_{\min}} \|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}}\right) \|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} \|\frac{\tilde{\mathbf{H}}_n(\xi^{\mathbb{N}_n^i})}{n} \epsilon^i\|_{\mathbb{F}},$$

and consider the event  $\mathcal{A}_3 = \{\|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} \leq \frac{c_1 \sqrt{n^i}}{1-\eta} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\}$ . Then

$$\begin{aligned} &P\left(\|R_{33}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ &\leq P\left(\|R_{33}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \text{ and } \mathcal{A}_3\right) + P(\mathcal{A}_3^c) \\ &\leq P\left(\|\frac{\tilde{\mathbf{H}}_n(\xi^{\mathbb{N}_n^i})}{n} \epsilon^i\|_{\mathbb{F}} \geq \frac{\sqrt{C_{\min}}}{6\sqrt{2}}\right) + P\left(\|\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n - \hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n\|_{\mathbb{F}} \geq \frac{C_{\min}}{4}\right) + P(\mathcal{A}_3^c) \\ &\lesssim \exp\left(-C_6 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_6 \frac{n^{1-\alpha(2+3\beta)}}{(n^i)^2 m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) + \exp\left(-C_6 \frac{n}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right), \end{aligned}$$



where we used (A.58) and Theorem A.1. Combining the results of  $\|R_{31}\|_{\mathbb{F}}$ ,  $\|R_{32}\|_{\mathbb{F}}$  and  $\|R_{33}\|_{\mathbb{F}}$  we conclude,

$$\begin{aligned} & P\left(\|R_3^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ & \lesssim \exp\left(-C_4 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right). \end{aligned}$$

*Term  $R_4^j$ .* We have

$$\begin{aligned} \max_{j \notin \mathbb{N}_n^i} \|R_4^j\|_{\mathbb{F}} & \leq \max_{j \notin \mathbb{N}_n^i} \|\hat{\Sigma}_{\xi^j \mathbb{N}_n^i}^n - \Sigma_{\xi^j \mathbb{N}_n^i}^n\|_{\mathbb{F}} \|(\hat{\Sigma}_{\mathbb{N}_n^i \mathbb{N}_n^i}^n + \lambda_n \hat{D}_{\mathbb{N}_n^i})^{-1}\|_2 \|\hat{\Sigma}_{\mathbb{N}_n^i \xi^i}^n - \Sigma_{\mathbb{N}_n^i \xi^i}^n\|_{\mathbb{F}} \\ & \leq \frac{4}{C_{\min}} \max_{j \notin \mathbb{N}_n^i} \|\hat{\Sigma}_{\xi^j \mathbb{N}_n^i}^n - \Sigma_{\xi^j \mathbb{N}_n^i}^n\|_{\mathbb{F}} \|\hat{\Sigma}_{\mathbb{N}_n^i \xi^i}^n - \Sigma_{\mathbb{N}_n^i \xi^i}^n\|_{\mathbb{F}}. \end{aligned}$$

Recall the definition of event  $\mathcal{A}_1 = \{\max_{j \notin \mathbb{N}_n^i} \|\hat{\Sigma}_{\xi^j \mathbb{N}_n^i}^n - \Sigma_{\xi^j \mathbb{N}_n^i}^n\|_{\mathbb{F}} \leq \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\}$ , then we have

$$\begin{aligned} & P\left(\max_{j \notin \mathbb{N}_n^i} \|R_4^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \leq P\left(\|\hat{\Sigma}_{\mathbb{N}_n^i \xi^i}^n - \Sigma_{\mathbb{N}_n^i \xi^i}^n\|_{\mathbb{F}} \geq c_1 \frac{C_{\min}}{4}\right) + P(\mathcal{A}_1^c) \\ & \lesssim \exp\left(-C_5 \frac{n^{1-\alpha(2+3\beta)}}{n^i m_n^2 k_n^3} + \log(n^i m_n^2 k_n)\right) \\ & + \exp\left(-C_5 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + \log(p m_n^2 k_n^2)\right), \end{aligned}$$

for some positive constant  $C_5$  depending on  $C_{\min}$ , where the estimates follow from Theorem A.1.

*Term  $R_5^j$ .* For the term  $R_5^j$  we obtain directly for some positive constant  $C_6$

$$\begin{aligned} & P\left(\max_{j \notin \mathbb{N}_n^i} \|R_5^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ & \leq (p - n^i) P\left(\|\Sigma_{\xi^j \xi^i}^n - \hat{\Sigma}_{\xi^j \xi^i}^n\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ & \lesssim \exp\left(-C_6 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{m_n^2 k_n^3} + \log(p m_n^2 k_n)\right). \end{aligned}$$

*Term  $R_6^j$ .* Using again the identity  $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$  and (A.49), we have

$$\begin{aligned} R_6^j & = \lambda_n \Sigma_{\xi^j \mathbb{N}_n^i}^n (\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n + \lambda_n D_{\mathbb{N}_n^i}^*)^{-1} (D_{\mathbb{N}_n^i}^* - \hat{D}_{\mathbb{N}_n^i}) \\ & \quad \times (\Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n + \lambda_n \hat{D}_{\mathbb{N}_n^i})^{-1} \left( \Sigma_{\mathbb{N}_n^i \mathbb{N}_n^i}^n B_n^{* \mathbb{N}_n^i} + \frac{\tilde{\mathbf{H}}_n(\xi^{\mathbb{N}_n^i})}{n} w^i + \frac{\tilde{\mathbf{H}}_n(\xi^{\mathbb{N}_n^i})}{n} \epsilon^i \right). \end{aligned}$$

Obviously, on the event  $\mathcal{N}$  we have

$$P\left(\max_{j \notin N_n^i} \|R_6^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \leq \sum_{r=1}^3 P\left(\|R_{6r}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right),$$

where

$$\begin{aligned} \|R_{61}\|_{\mathbb{F}} &= \frac{1-\eta}{\sqrt{n^i}} \lambda_n \|D_{N_n^i}^* - \hat{D}_{N_n^i}\|_2, \\ \|R_{62}\|_{\mathbb{F}} &= \frac{1-\eta}{\sqrt{n^i}} \lambda_n \|D_{N_n^i}^* - \hat{D}_{N_n^i}\|_2 \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i\|_{\mathbb{F}}, \\ \|R_{63}\|_{\mathbb{F}} &= \frac{1-\eta}{\sqrt{n^i}} \lambda_n \|D_{N_n^i}^* - \hat{D}_{N_n^i}\|_2 \|(\Sigma_{N_n^i N_n^i}^n + \lambda_n \hat{D}_{N_n^i})^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i\|_{\mathbb{F}}. \end{aligned}$$

Now, using the same computations as in the proof of Proposition 4 in [15], we obtain for the operator norm of the matrix  $(D_{N_n^i}^*)_{jj} - (\hat{D}_{N_n^i})_{jj}$

$$\begin{aligned} &\|(D_{N_n^i}^*)_{jj} - (\hat{D}_{N_n^i})_{jj}\|_2 \\ &= \left| \frac{\sum_{\ell \in N_n^i} \|\hat{B}_n^{i\ell}\|_{\mathbb{F}}}{\|\hat{B}_n^{ij}\|_{\mathbb{F}}} - \frac{\sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}}}{\|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}} \right| \\ &\leq \frac{\|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \sum_{\ell \in N_n^i} \left| \|\hat{B}_n^{i\ell}\|_{\mathbb{F}} - \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}} \right| + \left| \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} - \|\hat{B}_n^{ij}\|_{\mathbb{F}} \right| \sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}}}{\|\hat{B}_n^{ij}\|_{\mathbb{F}} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}} \\ &\leq \frac{\|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} \sum_{\ell \in N_n^i} \|\hat{B}_n^{i\ell} - B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}} + \|B_{m_n k_n}^{*ij} - \hat{B}_n^{ij}\|_{\mathbb{F}} \sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}}}{\|\hat{B}_n^{ij}\|_{\mathbb{F}} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}} \\ &\leq \frac{\sum_{\ell \in N_n^i} \|\hat{B}_n^{i\ell} - B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}} \sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}}}{\|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} (\|B_{m_n k_n}^{*ij}\|_{\mathbb{F}} - \|\hat{B}_n^{ij} - B_{m_n k_n}^{*ij}\|_{\mathbb{F}})}. \end{aligned}$$

On the event  $\mathcal{A}_4 = \{\|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \leq \frac{b_n^{*i}}{2}\}$ , we obtain

$$\begin{aligned} \|(D_{N_n^i}^*)_{jj} - (\hat{D}_{N_n^i})_{jj}\|_2 &\leq 2(b_n^{*i})^{-2} \sum_{\ell \in N_n^i} \|\hat{B}_n^{i\ell} - B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}} \sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}} \\ &\leq 2(b_n^{*i})^{-2} (n^i)^{1/2} \|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \sum_{\ell \in N_n^i} \|B_{m_n k_n}^{*i\ell}\|_{\mathbb{F}}, \end{aligned}$$

where we used the Cauchy-Schwarz inequality. Consequently,

$$P\left(\|R_{61}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \leq P\left(\|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_{\mathbb{F}} \geq \frac{c_1}{6} \frac{1}{1-\eta} (b_n^{*i})^2\right) + P(\mathcal{A}_4^c).$$

Now, we apply Proposition A.5 with  $\delta = \frac{c_1}{6} \frac{1}{1-\eta} (b_n^{*i})^2$  and  $\delta = \frac{b_n^{*i}}{2}$  to obtain

$$\begin{aligned} P\left(\|R_{61}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) &\lesssim \exp\left(-C_7 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^6}{(n^i)^4 m_n^2 k_n^4 (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_7 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^4}{(n^i)^4 m_n^2 k_n^4 (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2} + 2 \log(n^i m_n k_n)\right) \\ &\lesssim \exp\left(-C_7 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right), \end{aligned}$$

where we used  $\frac{2}{C_{\min}} \lambda_n (n^i)^{3/2} (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2 \leq c_2 (b_n^{*i})^s$  ( $s = 2, 3$ ) for the last inequality (see condition (A.66)) with  $C_7 > 0$  depending on  $\eta$ .

By the same arguments and using (A.53), (A.54) and Assumption 4.9, we can show the existence of a constant  $C_8 > 0$  such that

$$\begin{aligned} P\left(\|R_{62}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) &\lesssim \exp\left(-C_8 \frac{n^{1-\alpha(2+3\beta)} \lambda_n^{-2} (b_n^{*i})^6}{(n^i)^4 m_n^2 k_n^4 (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_8 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^4}{(n^i)^4 m_n^2 k_n^4 (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2} + 2 \log(n^i m_n k_n)\right) + P(\Omega^{\circ}) \\ &\lesssim \exp\left(-C_7 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) + P(\Omega^{\circ}), \end{aligned}$$

where the probability  $P(\Omega^{\circ})$  can be estimated by Proposition A.1 and is dominated by the first term because of Assumption 4.5. Similarly, using (A.58), we obtain

$$\begin{aligned} P\left(\|R_{63}\|_{\mathbb{F}} \geq \frac{c_1}{3} \lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) &\lesssim \exp\left(-C_9 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^6}{(n^i)^4 m_n^2 k_n^4 (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_9 \frac{n^{1-\alpha(2+3\beta)} (b_n^{*i})^4}{(n^i)^4 m_n^2 k_n^4 (\sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_9 \frac{n}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right) \\ &\lesssim \exp\left(-C_7 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in \mathbb{N}_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right) \\ &\quad + \exp\left(-C_9 \frac{n}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right). \end{aligned}$$

Combining the results for  $\|R_{61}\|_F$ ,  $\|R_{62}\|_F$  and  $\|R_{63}\|_F$ , we can conclude that

$$\begin{aligned} P\left(\max_{j \notin N^i} \|R_6^j\|_F \geq \frac{C_1}{3} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_F\right) \\ \lesssim \exp\left(-C_7 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_F)^2}{n^i m_n^2 k_n^4} + 2 \log(n^i m_n k_n)\right), \end{aligned}$$

for a positive constant  $C_7 > 0$  that depends on  $\eta$ .

*Term  $R_7^j$ .* Observing (A.49) and (A.50), the term  $R_7^j$  can be further decomposed as

$$\begin{aligned} R_7^j &= \sum_{\xi^j N_n^i} \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right)^{-1} \left( \sum_{N_n^i N_n^i}^n B_n^{*N_n^i} + \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} w^i + \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{n} \epsilon^i \right) \\ &\quad - \sum_{\xi^j N_n^i} B_n^{*N_n^i} - \frac{\tilde{H}_n^\top(\xi^j)}{n} w^i - \frac{\tilde{H}_n^\top(\xi^j)}{n} \epsilon^i \\ &= R_{71}^j + R_{72}^j + R_{73}^j, \end{aligned}$$

where

$$\begin{aligned} R_{71}^j &= \sum_{\xi^j N_n^i} \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right)^{-1} \sum_{N_n^i N_n^i}^n B_n^{*N_n^i} - \sum_{\xi^j N_n^i} B_n^{*N_n^i} \\ R_{72}^j &= \sum_{\xi^j N_n^i} \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right)^{-1} \frac{\tilde{\mathbf{H}}_n^\top(\xi^{N_n^i})}{n} w^i - \frac{\tilde{H}_n^\top(\xi^j)}{n} w^i \\ R_{73}^j &= \sum_{\xi^j N_n^i} \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right)^{-1} \frac{\tilde{\mathbf{H}}_n^\top(\xi^{N_n^i})}{n} \epsilon^i - \frac{\tilde{H}_n^\top(\xi^j)}{n} \epsilon^i. \end{aligned}$$

Turning to  $\|R_{71}^j\|_F$ , we have

$$\begin{aligned} \max_{j \notin N_n^i} \|R_{71}^j\|_F &\leq \max_{j \notin N_n^i} \left\| \sum_{\xi^j N_n^i} \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right)^{-1} \left( \sum_{N_n^i N_n^i}^n B_n^{*N_n^i} - \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right) B_n^{*N_n^i} \right) \right\|_F \\ &= \lambda_n \max_{j \notin N_n^i} \left\| \sum_{\xi^j N_n^i} \left( \sum_{N_n^i N_n^i}^n \right)^{-1} D_{N_n^i}^* B_n^{*N_n^i} \right\|_F \\ &\leq (1-\eta) \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_F, \end{aligned} \tag{A.77}$$

where we have used  $\sum_{N_n^i N_n^i}^n \preceq \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^*$  and Assumption 4.7.

For the term  $R_{72}^j$ , we first write

$$\begin{aligned} \max_{j \notin N_n^i} \|R_{72}^j\|_F &\leq \max_{j \notin N_n^i} \left\| \frac{\tilde{H}_n^\top(\xi^j)}{\sqrt{n}} \left( I - \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}} \left( \sum_{N_n^i N_n^i}^n + \lambda_n D_{N_n^i}^* \right)^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}} \right) \frac{w^i}{\sqrt{n}} \right\|_F \\ &\leq \max_{j \notin N_n^i} \left\| \frac{\tilde{H}_n^\top(\xi^j)}{\sqrt{n}} \right\|_2 \left\| \frac{w^i}{\sqrt{n}} \right\|_F \left\| I - \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}} \left( \sum_{N_n^i N_n^i}^n \right)^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}} \right\|_2. \end{aligned}$$

The matrix  $I - \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}} (\sum_{N_n^i}^n)^{-1} \frac{\tilde{\mathbf{H}}_n(\xi^{N_n^i})}{\sqrt{n}}$  defines a projection and therefore its operator norm is 1. Moreover, by the Lemma 6.2 in [44] we have  $\|\frac{\tilde{H}_n^\top(\xi^j)}{\sqrt{n}} \frac{\tilde{H}_n(\xi^j)}{\sqrt{n}}\|_2 \leq \frac{1}{k_n}$  for all  $j \in \mathcal{V}$ . Thus, observing (A.54), it follows that on the event  $\Omega$  defined in (A.20)

$$\left\| \frac{\tilde{H}_n^\top(\xi^j)}{\sqrt{n}} \right\|_2 \left\| \frac{w^i}{\sqrt{n}} \right\|_{\mathbb{F}} \lesssim \frac{n^i m_n^{3/2}}{k_n^{d+1/2}}.$$

By condition (4.9),

$$\max_{j \notin N_n^i} \|R_{72}^j\|_{\mathbb{F}} \leq \frac{n^i m_n^{3/2}}{k_n^{d+1/2}} \lesssim \sqrt{\frac{2}{C_{\min}}} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}.$$

Combining this inequality and (A.77) now yields

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|R_7^j\|_{\mathbb{F}} \geq c_1 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) &\leq P\left(\max_{j \notin N_n^i} \|R_{73}^j\|_{\mathbb{F}} \geq c_3 \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ &\leq P\left(\left\| \frac{\tilde{\mathbf{H}}_n^\top(\xi^{N_n^i})}{n} \epsilon^i \right\|_{\mathbb{F}} \geq \frac{c_3}{2} \frac{\sqrt{n^i}}{(1-\eta)} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ &\quad + (p - n^i) P\left(\left\| \frac{\tilde{H}_n^\top(\xi^j)}{n} \epsilon^i \right\|_{\mathbb{F}} \geq \frac{c_3}{2} \lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right) \\ &\lesssim \exp\left(-C_s \frac{n \left(\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right)^2}{n^i m_n^2 k_n} + \log(n^i m_n^2 k_n)\right) \\ &\quad + \exp\left(-C_s \frac{n \left(\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right)^2}{m_n^2 k_n} + \log((p - n^i) m_n^2 k_n)\right) + P(\Omega^c). \end{aligned}$$

where  $c_3 = c_1 - (1 - \eta) - (\frac{2}{C_{\min}})^{1/2}$ . Finally, combining the result (A.69) from Step 1 with the estimates for  $R_1, \dots, R_7$ , we conclude that

$$\begin{aligned} P\left(\max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^j N_n^i}^n \hat{B}_n^{N_n^i} - \hat{\Sigma}_{\xi^j \xi^i}^n\|_{\mathbb{F}} \geq \lambda_n \sum_{j \neq i}^p \|\hat{B}_n^{ij}\|_{\mathbb{F}}\right) \\ \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} \left(\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}}\right)^2}{n^i m_n^2 k_n^4} + 2 \log(pm_n k_n)\right), \end{aligned}$$

such that  $\lambda_n (n^i)^{3/2} \lesssim (b_n^{*i})^3 (\sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_{\mathbb{F}})^{-2}$ . □

#### A.4. Proof of Proposition A.2

First, note that the event  $\hat{N}_n^i = N_n^i$  holds if and only if

$$\|\hat{B}_n^{ij}\|_{\mathbb{F}} \neq 0 \quad \forall j \in N_n^i \quad \text{and} \quad \|\hat{B}_n^{ij}\|_{\mathbb{F}} = 0 \quad \forall j \notin N_n^i,$$

which is implied by the conditions

$$\begin{aligned} \|\hat{B}_n^{N_n^i} - B_n^{*N_n^i}\|_F &< b_n^{*i} = \min_{j \in N^i} \|B_{m_n k_n}^{*ij}\|_F, \\ \max_{j \notin N_n^i} \|\hat{\Sigma}_{\xi^{jN^i}}^n \hat{B}_n^{N_n^i} - \hat{\Sigma}_{\xi^j \xi^i}^n\|_F &\leq \lambda_n \sum_{j \in N^i} \|\hat{B}_n^{ij}\|_F. \end{aligned}$$

Thus, the Proposition A.5 (with  $\delta = b_n^{*i}$ , an obvious estimate of the probability using condition (A.66)) and Proposition A.6 we can conclude that

$$P(\hat{N}_n^i \neq N_n^i \text{ and } \mathcal{N}) \lesssim \exp\left(-C_1 \frac{n^{1-\alpha(2+3\beta)} (\lambda_n \sum_{j \in N_n^i} \|B_{m_n k_n}^{*ij}\|_F)^2}{n^i m_n^2 k_n^4} + 2\log(pm_n k_n)\right)$$

and this completes the proof of Proposition A.2.

### Acknowledgments

The authors would like to thank Martina Stein who typed parts of this manuscript with considerable technical expertise.

### References

- [1] Francis R Bach. Consistency of the group lasso and multiple kernel learning. *Journal of Machine Learning Research*, 9(Jun):1179–1225, 2008. [MR2417268](#)
- [2] Denis Bosq. *Linear Processes in Function Spaces: Theory and Applications*, volume 149. Springer Science & Business Media, 2012. [MR1783138](#)
- [3] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013. [MR3185193](#)
- [4] Stephen Boyd, Stephen P Boyd, and Lieven Vandenberghe. *Convex optimization*. Cambridge university press, 2004. [MR2061575](#)
- [5] Peter Bühlmann, Jonas Peters, and Jan Ernest. Cam: Causal additive models, high-dimensional order search and penalized regression. *The Annals of Statistics*, 42(6):2526–2556, 2014. [MR3277670](#)
- [6] Robert G Cowell, Philip Dawid, Steffen L Lauritzen, and David J Spiegelhalter. *Probabilistic networks and expert systems: Exact computational methods for Bayesian networks*. Springer Science & Business Media, 2007. [MR1697175](#)
- [7] Carl De Boor. *A Practical Guide to Splines*, volume 27. springer-verlag New York, 1978. [MR0507062](#)
- [8] Yingying Fan, Gareth M James, and Peter Radchenko. Functional additive regression. *The Annals of Statistics*, 43(5):2296–2325, 2015. [MR3396986](#)
- [9] Kyunghye Han, Hans-Georg Müller, Byeong U Park, et al. Smooth back-fitting for additive modeling with small errors-in-variables, with an application to additive functional regression for multiple predictor functions. *Bernoulli*, 24(2):1233–1265, 2018. [MR3706793](#)

- [10] Wassily Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963. [MR0144363](#)
- [11] Tailen Hsing and Randall Eubank. *Theoretical Foundations of Functional Data Analysis, with an Introduction to Linear Operators*. John Wiley & Sons, 2015. [MR3379106](#)
- [12] Jian Huang, Joel L Horowitz, and Fengrong Wei. Variable selection in non-parametric additive models. *Annals of Statistics*, 38(4):2282–2313, 2010. [MR2676890](#)
- [13] Lester Ingber. Statistical mechanics of neocortical interactions: Canonical momenta indicators of electroencephalography. *Physical Review E*, 55(4):4578, 1997. [MR1609439](#)
- [14] Kuang-Yao Lee, Bing Li, and Hongyu Zhao. Variable selection via additive conditional independence. *Journal of the Royal Statistical Society, Series B (Statistical Methodology)*, 78(5):1037–1055, 2016. [MR3557188](#)
- [15] Kuang-Yao Lee, Lexi Li, Bing Li, and Hongyu Zhao. Nonparametric functional graphical modeling through functional additive regression operator. *Working paper*, 2020.
- [16] Bing Li and Eftychia Solea. A nonparametric graphical model for functional data with application to brain networks based on fmri. *Journal of the American Statistical Association*, 113(524):1637–1655, 2018. [MR3902235](#)
- [17] Han Liu, Fang Han, Ming Yuan, John Lafferty, Larry Wasserman, et al. High-dimensional semiparametric gaussian copula graphical models. *The Annals of Statistics*, 40(4):2293–2326, 2012. [MR3059084](#)
- [18] Han Liu, John Lafferty, and Larry Wasserman. The nonparanormal: Semiparametric estimation of high dimensional undirected graphs. *Journal of Machine Learning Research*, 10:2295–2328, 2009. [MR2563983](#)
- [19] Po-Ling Loh and Peter Bühlmann. High-dimensional learning of linear causal networks via inverse covariance estimation. *The Journal of Machine Learning Research*, 15(1):3065–3105, 2014. [MR3277162](#)
- [20] Lukas Meier, Sara Van de Geer, Peter Bühlmann, et al. High-dimensional additive modeling. *Annals of Statistics*, 37(6B):3779–3821, 2009. [MR2572443](#)
- [21] Nicolai Meinshausen and Peter Bühlmann. High-dimensional graphs and variable selection with the lasso. *Annals of Statistics*, pages 1436–1462, 2006. [MR2278363](#)
- [22] Nicolai Meinshausen, Bin Yu, et al. Lasso-type recovery of sparse representations for high-dimensional data. *Annals of Statistics*, 37(1):246–270, 2009. [MR2488351](#)
- [23] Guillaume Obozinski, Martin J Wainwright, Michael I Jordan, et al. Support union recovery in high-dimensional multivariate regression. *Annals of Statistics*, 39(1):1–47, 2011. [MR2797839](#)
- [24] Judea Pearl. *Causality: Models, Reasoning and Inference*. Cambridge University Press, 2009. [MR2548166](#)
- [25] Justin Petrovich, Matthew Reimherr, and Carrie Daymont. Functional regression models with highly irregular designs. *arXiv preprint arXiv:1805*.

- [08518](#), 2018.
- [26] Xinghao Qiao, Shaojun Guo, and Gareth M James. Functional graphical models. *Journal of the American Statistical Association*, 114(525):211–222, 2019. [MR3941249](#)
  - [27] Xinghao Qiao, Cheng Qian, Gareth M James, and Shaojun Guo. Doubly functional graphical models in high dimensions. *Biometrika*, 107(2):415–431, 2020. [MR4108937](#)
  - [28] Pradeep Ravikumar, John Lafferty, Han Liu, and Larry Wasserman. Sparse additive models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 71(5):1009–1030, 2009. [MR2750255](#)
  - [29] Pradeep Ravikumar, Martin J Wainwright, Garvesh Raskutti, Bin Yu, et al. High-dimensional covariance estimation by minimizing  $\ell_1$ -penalized log-determinant divergence. *Electronic Journal of Statistics*, 5:935–980, 2011. [MR2836766](#)
  - [30] Larry Schumaker. *Spline Functions: Basic Theory*. Cambridge University Press, 2007. [MR2348176](#)
  - [31] Eftychia Solea and Bing Li. Copula gaussian graphical models for functional data. *Journal of the American Statistical Association*, pages 1–13, 2020. [MR4436312](#)
  - [32] Charles J Stone et al. Additive regression and other nonparametric models. *Annals of Statistics*, 13(2):689–705, 1985. [MR0790566](#)
  - [33] Arend Voorman, Ali Shojaie, and Daniela Witten. Graph estimation with joint additive models. *Biometrika*, 101(1):85–101, 2013. [MR3180659](#)
  - [34] Raymond KW Wong, Yehua Li, and Zhengyuan Zhu. Partially linear functional additive models for multivariate functional data. *Journal of the American Statistical Association*, 114(525):406–418, 2019. [MR3941264](#)
  - [35] Luo Xiao, Cai Li, William Checkley, and Ciprian Crainiceanu. Fast covariance estimation for sparse functional data. *Statistics and computing*, 28(3):511–522, 2018. [MR3761337](#)
  - [36] Lingzhou Xue, Hui Zou, et al. Regularized rank-based estimation of high-dimensional nonparanormal graphical models. *Annals of Statistics*, 40(5):2541–2571, 2012. [MR3097612](#)
  - [37] Fang Yao, Hans-Georg Müller, and Jane-Ling Wang. Functional data analysis for sparse longitudinal data. *Journal of the American Statistical Association*, 100(470):577–590, 2005. [MR2160561](#)
  - [38] M. Yuan and Y. Lin. Model selection and estimation in gaussian graphical model. *Biometrika*, 94:19–35, 2007. [MR2367824](#)
  - [39] Ming Yuan and Yi Lin. Model selection and estimation in regression with grouped variables. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 68(1):49–67, 2006. [MR2212574](#)
  - [40] Javier Zapata, Sang-Yun Oh, and Alexander Petersen. Partial separability and functional graphical models for multivariate gaussian processes. *arXiv preprint arXiv: 1910.03134*, 2019. [MR4472841](#)
  - [41] Xiao Lei Zhang, Henri Begleiter, Bernice Porjesz, Wenyu Wang, and Ann Litke. Event related potentials during object recognition tasks. *Brain Research Bulletin*, 38(6):531–538, 1995.



- [42] Boxin Zhao, Y Samuel Wang, and Mladen Kolar. Fudge: Functional differential graph estimation with fully and discretely observed curves. *arXiv preprint arXiv: 2003.05402*, 2020.
- [43] Boxin Zhao, Shengjun Zhai, Y Samuel Wang, and Mladen Kolar. High-dimensional functional graphical model structure learning via neighborhood selection approach. *arXiv preprint arXiv: 2105.02487*, 2021.
- [44] S Zhou, X Shen, DA Wolfe, et al. Local asymptotics for regression splines and confidence regions. *Annals of Statistics*, 26(5):1760–1782, 1998. [MR1673277](#)
- [45] Hongxiao Zhu, Nate Strawn, and David B Dunson. Bayesian graphical models for multivariate functional data. *Journal of Machine Learning Research*, 17(1):7157–7183, 2016. [MR3580357](#)
- [46] Hongxiao Zhu, Fang Yao, and Hao Helen Zhang. Structured functional additive regression in reproducing kernel hilbert spaces. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(3):581–603, 2014. [MR3210729](#)