

# WIGNER THEOREMS FOR RANDOM MATRICES WITH DEPENDENT ENTRIES: ENSEMBLES ASSO- CIATED TO SYMMETRIC SPACES AND SAMPLE CO- VARIANCE MATRICES

KATRIN HOFMANN-CREDNER

*Ruhr-Universität Bochum, Fakultät für Mathematik, NA 3/31, D-44780 Bochum, Germany*  
email: [katrin.credner@rub.de](mailto:katrin.credner@rub.de)

MICHAEL STOLZ<sup>1</sup>

*Ruhr-Universität Bochum, Fakultät für Mathematik, NA 3/31, D-44780 Bochum, Germany*  
email: [michael.stolz@rub.de](mailto:michael.stolz@rub.de)

*Submitted July 19, 2007, accepted in final form June 17, 2008*

AMS 2000 Subject classification: Primary 15A52, Secondary 82B44

Keywords: random matrices, symmetric spaces, semicircle law, Wigner, Marcenko-Pastur, Wishart, sample covariance matrices, dependent random variables, density of states, universality

## *Abstract*

It is a classical result of Wigner that for an hermitian matrix with independent entries on and above the diagonal, the mean empirical eigenvalue distribution converges weakly to the semicircle law as matrix size tends to infinity. In this paper, we prove analogs of Wigner's theorem for random matrices taken from all infinitesimal versions of classical symmetric spaces. This is a class of models which contains those studied by Wigner and Dyson, along with seven others arising in condensed matter physics. Like Wigner's, our results are universal in that they only depend on certain assumptions about the moments of the matrix entries, but not on the specifics of their distributions. What is more, we allow for a certain amount of dependence among the matrix entries, in the spirit of a recent generalization of Wigner's theorem, due to Schenker and Schulz-Baldes. As a byproduct, we obtain a universality result for sample covariance matrices with dependent entries.

## 1 Introduction

Classical physics-inspired random matrix theory is chiefly concerned with probability measures on what Freeman Dyson in 1962 called the “threefold way”, namely, the spaces of hermitian, real symmetric, and quaternion real matrices (or their respective exponentiated, compact versions). The rationale behind this focus is Dyson's proof that any hermitian matrix (thought

---

<sup>1</sup>RESEARCH OF BOTH AUTHORS SUPPORTED BY DEUTSCHE FORSCHUNGSGEMEINSCHAFT VIA SFB/TR 12

of as a truncated Hamiltonian of a quantum system) that commutes with a group of unitary symmetries and “time reversals” breaks down to these three constituents ([7]), which are, in structural terms, the tangent spaces to the Riemannian Symmetric Spaces (RSS) of class A, AI and AII.

During the last decade, theoretical condensed matter physicists have pointed out that matrix descriptions of systems such as mesoscopic normal-superconducting hybrid structures are outside the scope of Dyson’s theorem, and that the tangent spaces to all ten infinite series of classical RSS may (and do indeed) arise. The deeper reasons are explained in [1, Section 6.4], [2], and [11]. Some of this material is summarized in [8]. An alternative motivation for the study of random matrices associated to the classical RSS comes from analytic number theory, see [13]. Concrete matrix realizations of this “tenfold way” of “symmetry classes” are given in Section 2 below.

The task of developing random matrix theories for the full “tenfold way”, i.e., studying probability measures on all ten series of matrix spaces, has been taken up in [8], where the probability measures enjoy invariance properties that guarantee an explicit analytic expression for the joint eigenvalue density, and in [5, 6], where the focus is on the compact versions of the classical RSS, endowed with their natural invariant probability measure. In the present paper, we abandon invariance properties and turn to analogs of Wigner’s famous result of 1958 ([18]), stating that for a symmetric matrix with independent entries on and above the diagonal, the mean empirical eigenvalue distribution converges weakly to the semicircle law as matrix size tends to infinity. This is a universality result in the sense that it only depends on certain assumptions about the moments of the matrix entries, but not on the specifics of their distributions. In recent years, related universality results for Gaussian fluctuations in Wigner’s set-up have become available (see [4, 17] and the references therein), but we will restrict our attention to the law of large numbers level that was considered in Wigner’s fundamental paper.

Actually, our starting point is not the classical version of Wigner’s result, but a recent generalization, due to Schenker and Schulz-Baldes ([16]), allowing for a certain amount of dependence to hold among the matrix entries. Specifically, the authors consider the following set-up: For each  $n \in \mathbb{N}$  write  $\mathbb{I}_n := \{1, \dots, n\}$  and suppose that  $\mathbb{I}_n^2 = \mathbb{I}_n \times \mathbb{I}_n$  comes with an equivalence relation  $\sim_n$ . The entries of the matrix  $X_n = (\frac{1}{\sqrt{n}} a_n(p, q))_{p, q=1, \dots, n}$  are complex random variables, with  $a_n(p_1, q_1), \dots, a_n(p_j, q_j)$  independent whenever  $(p_1, q_1), \dots, (p_j, q_j)$  belong to  $j$  distinct equivalence classes of the relation  $\sim_n$ . Furthermore, it is required that  $a_n(p, q) = \overline{a_n(q, p)}$  for all  $n, p, q$ . In the case that all equivalence classes of  $\sim_n$  are of the form  $\{(p, q), (q, p)\}$ , one is back to hermitian matrices with independent entries on and above the diagonal, i.e. to the situation of Wigner’s theorem. If some equivalence classes are larger, then there is some leeway for violations of independence. In the framework of Schenker and Schulz-Baldes, the following conditions on  $\sim_n$  serve as a less restrictive substitute for independence:

$$(W1) \max_p \#\{(q, p', q') \in \mathbb{I}_n^3 : (p, q) \sim_n (p', q')\} = o(n^2)$$

$$(W2) \max_{p, q, p'} \#\{q' \in \mathbb{I}_n : (p, q) \sim_n (p', q')\} \leq B, \text{ where } B < \infty \text{ is a constant}$$

$$(W3) \#\{(p, q, p') \in \mathbb{I}_n^3 : (p, q) \sim_n (q, p') \text{ and } p \neq p'\} = o(n^2).$$

Apart from that, one requires that for all  $n, p, q$ ,  $a_n(p, q)$  is centered and

$$\mathbb{E}(a_n(p, q)\overline{a_n(p, q)}) = 1. \tag{1}$$

Furthermore, a uniform bound on the  $k$ -th moments is assumed:

$$\sup_n \max_{p,q=1,\dots,n} \mathbb{E}(|a_n(p,q)|^k) < \infty \text{ for all } k \in \mathbb{N}. \quad (2)$$

For an hermitian matrix  $M \in \mathbb{C}^{n \times n}$  with eigenvalues  $\lambda_1, \dots, \lambda_n$  write

$$L_n(M) := \frac{1}{n} \sum_{j=1}^n \delta_{\lambda_j} \quad (3)$$

for the empirical measure of the eigenvalues of  $M$ . Then the main theorem of [16] can be stated as follows:

**THEOREM 1.1.** *Under conditions (W1), (W2), (W3), (1), and (2), the measure  $\mathbb{E}(L_n(X_n))$ , i.e. the mean empirical eigenvalue distribution, converges weakly to the semicircle law with density*

$$\frac{1}{2\pi} 1_{[-2,2]}(x) \sqrt{4 - x^2}. \quad (4)$$

So one ends up with the same limit distribution as in the independent case. In fact, conditions (W1), (W2), (W3) arose from a close reading of Wigner's proof, in order to understand how much independence is really needed to arrive at the semicircle law. So the approach of Schenker and Schulz-Baldes is complementary to the route taken in recent work of Anderson and Zeitouni ([3]), where a different dependence structure leads to a limit which is not a semicircle.

To establish analogs of Theorem 1.1 for all ten symmetry classes, we proceed as follows: In Section 2 we give precise definitions of the matrix spaces in question and introduce some auxiliary notation for the combinatorics of moment calculations. In Section 3 we treat those classes for which  $\mathbb{E}L_n$  converges to the semicircle distribution and for which nothing more than a slight extension of Theorem 1.1 is needed. On the other hand, substantial work has to be done for the so-called "chiral" classes, which, despite their roots in physics, are related to sample covariance matrices and hence lead to some relative of the Marčenko-Pastur distribution as limit for  $\mathbb{E}L_n$ . The main step is to rework the combinatorics of moment convergence to the Marčenko-Pastur law in the spirit of Schenker and Schulz-Baldes, yielding a universality result for sample covariance matrices with dependent entries. This is the content of Section 4. In the final Section 5, this result is applied to the chiral classes.

## 2 Background and notation

We begin by listing the ten "symmetry classes", i.e. series of matrix spaces, from which our matrices are taken. In structural terms, these spaces are of the form  $i\mathfrak{g}$ , where  $\mathfrak{g}$  is the Lie algebra of a compact classical group, or  $i\mathfrak{p}$ , where  $\mathfrak{p}$  is the  $-1$  eigenspace of a Cartan involution of  $\mathfrak{g}$  (see [8] for details). The  $i$  factor is to make sure that the matrices are hermitian. The labels A, AI, etc. in the list are those of Lie theory, but no Lie theoretic properties will be needed in what follows.  $X^*$  denotes the conjugate transpose of a matrix  $X$ .

**Class A**

$$\mathbb{M}_n^A = \{X \in \mathbb{C}^{n \times n} : X \text{ hermitian}\}$$

**Class AI**

$$\mathbb{M}_n^{AI} = \{X \in \mathbb{R}^{n \times n} : X \text{ symmetric}\}$$

**Class AII**

$$\mathbb{M}_n^{\text{AII}} = \left\{ \begin{pmatrix} X_1 & X_2 \\ -\overline{X_2} & \overline{X_1} \end{pmatrix} : \begin{array}{l} X_i \in \mathbb{C}^{n \times n}, X_1 \text{ hermitian,} \\ X_2 \text{ skew symmetric} \end{array} \right\}$$

**Class AIII**

$$\mathbb{M}_n^{\text{AIII}} = \left\{ \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} : X \in \mathbb{C}^{s \times t} \right\}$$

**Class B/D**

$$\mathbb{M}_n^{\text{B/D}} = \{X \in (i\mathbb{R})^{n \times n} : X \text{ skew symmetric}\}$$

**Class BDI**

$$\mathbb{M}_n^{\text{BDI}} = \left\{ \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} : X \in (i\mathbb{R})^{s \times t} \right\}$$

**Class DIII**

$$\mathbb{M}_n^{\text{DIII}} = \left\{ \begin{pmatrix} X_1 & X_2 \\ X_2 & -X_1 \end{pmatrix} : X_i \in (i\mathbb{R})^{n \times n} \text{ skew symmetric} \right\}$$

**Class C**

$$\mathbb{M}_n^{\text{C}} = \left\{ \begin{pmatrix} X_1 & X_2 \\ \overline{X_2} & -\overline{X_1} \end{pmatrix} : \begin{array}{l} X_i \in \mathbb{C}^{n \times n}, X_1 \text{ hermitian,} \\ X_2 \text{ symmetric} \end{array} \right\}$$

**Class CI**

$$\mathbb{M}_n^{\text{CI}} = \left\{ \begin{pmatrix} X_1 & X_2 \\ X_2 & -X_1 \end{pmatrix} : X_i \in \mathbb{R}^{n \times n} \text{ symmetric} \right\}$$

**Class CII**

$$\mathbb{M}_n^{\text{CII}} = \left\{ \begin{pmatrix} 0 & X \\ X^* & 0 \end{pmatrix} : X \in \mathbb{H}^{s \times t} \right\},$$

where the space  $\mathbb{H}^{s \times t}$  of quaternionic matrices is embedded into  $\mathbb{C}^{2s \times 2t}$  as

$$\mathbb{H}^{s \times t} = \left\{ \begin{pmatrix} U & V \\ -\overline{V} & \overline{U} \end{pmatrix} : U, V \in \mathbb{C}^{s \times t} \right\}.$$

Generically, we write  $\mathcal{C}$  for any label A, AI, ..., CII. We have  $\mathbb{M}_n^{\mathcal{C}} \subset \mathbb{C}^{\delta n \times \delta n}$  with  $\delta = \delta_{\mathcal{C}} = 2$  if  $\mathcal{C} = \text{AII, DIII, C, CI, CII}$  and  $\delta = 1$  otherwise. Classes AIII, BDI and CII, the ‘‘chiral’’ classes in physics terminology (since they are related to Dirac operators, see [9], [11]), are special in that the shape of the subblocks depends on an extra parameter  $s$ , and it is clear that one will have to control the relative growth of  $s = s(n)$  and  $t(n) = n - s(n)$  as  $n \rightarrow \infty$ . In fact, their large  $n$  behaviour is quite different from that of the other classes. Consequently, in what follows, we will treat these classes separately. A, AI, and AII are the classical Wigner-Dyson classes (underlying GUE, GOE, and GSE, resp.). The remaining classes, arising via Bogoliubov-de Gennes mean field approximation (see [1, Sec. 6.4]), will be termed Bogoliubov-de Gennes (BdG) or superconductor classes.

Before getting down to business, let us review some combinatorial notation and facts that will be useful later on. Write  $\mathcal{P}(n)$  for the set of all partitions of  $\mathbb{I}_n = \{1, 2, \dots, n\}$ . If  $\mathbf{p} \in \mathcal{P}(n)$  has  $r$  blocks, write  $|\mathbf{p}| = r$ . Define  $\mathcal{P}^{(i)}(n) := \{\mathbf{p} \in \mathcal{P}(n) : |\mathbf{p}| = i\}$ . If each of the blocks of  $\mathbf{p}$  consists of exactly two elements, we say that  $\mathbf{p}$  is a pair partition and write  $\mathbf{p} \in \mathcal{P}_2(n)$ . For  $\mathbf{p} \in \mathcal{P}(n)$ , write  $\sim_{\mathbf{p}} \subseteq \mathbb{I}_n \times \mathbb{I}_n$  for the corresponding equivalence relation. Given  $\mathbf{p}, \mathbf{q} \in \mathcal{P}(n)$ ,

define  $\sim_{\mathbf{p}} \vee \sim_{\mathbf{q}} \subseteq \mathbb{I}_n \times \mathbb{I}_n$  as the smallest equivalence relation which contains  $\sim_{\mathbf{p}}$  and  $\sim_{\mathbf{q}}$ . The partition corresponding to  $\sim_{\mathbf{p}} \vee \sim_{\mathbf{q}}$  is denoted by  $\mathbf{p} \vee \mathbf{q}$ .

$\mathbf{p} \in \mathcal{P}(n)$  is called crossing, if there exist  $p_1 < q_1 < p_2 < q_2$  in  $\mathbb{I}_n$  such that  $p_1 \sim_{\mathbf{p}} p_2 \not\sim_{\mathbf{p}} q_1 \sim_{\mathbf{p}} q_2$ . Otherwise, it is called noncrossing. Write  $\text{NC}(n)$ ,  $\text{NC}^{(i)}(n)$  and  $\text{NC}_2(n)$  for the set of noncrossing partitions, noncrossing partitions with  $i$  blocks, and noncrossing pair partitions of  $\mathbb{I}_n$ , respectively. For sets  $\Omega, \Omega'$  write  $\mathcal{F}(\Omega, \Omega') := \{\varphi : \Omega \rightarrow \Omega'\}$  and  $\mathcal{F}(k, n) := \mathcal{F}(\mathbb{I}_k, \mathbb{I}_n)$ . For  $\mathbf{p} \in \mathcal{P}(k)$  write  $\mathcal{F}(\mathbf{p}, \Omega)$  for the set of all  $\varphi \in \mathcal{F}(\mathbb{I}_k, \Omega)$  which are constant on the blocks of  $\mathbf{p}$ . Finally, let us give names to two important special pair partitions:

$$\begin{aligned} \mathbf{m} &:= \{\{1, 2\}, \{3, 4\}, \dots, \{2k-1, 2k\}\} \in \mathcal{P}_2(2k) \quad \text{and} \\ \mathbf{n} &:= \{\{2, 3\}, \{4, 5\}, \dots, \{2k, 1\}\} \in \mathcal{P}_2(2k). \end{aligned}$$

It is well-known that  $\#\text{NC}_2(2k)$  equals the  $k$ th Catalan number  $C_k$ , which in turn equals the  $2k$ th moment of the semicircular distribution with density given in (4) (whose odd moments vanish). On the other hand, setting for  $\kappa > 0$

$$m_k := \sum_{i=1}^k \#(\text{NC}^{(i)}(k)) \kappa^i, \tag{5}$$

$(m_k)_{k \in \mathbb{N}}$  is the sequence of moments of the Marčenko-Pastur distribution with density

$$\max\{0, (1 - \kappa)\} \delta_0 + \frac{\sqrt{4\kappa - (x - 1 - \kappa)^2}}{2\pi x} 1_{[(1-\sqrt{\kappa})^2, (1+\sqrt{\kappa})^2]}(x).$$

A reference for these facts is [10] or [12].

### 3 Wigner-Dyson and Bogoliubov-de Gennes classes

These are the easy cases, because they basically reduce to Theorem 1.1. In fact, one may interpret the symmetries of a matrix from  $\mathbb{M}_n^{\mathcal{C}}$  as an equivalence relation on pairs of indices, recalling that in the set-up of Theorem 1.1, if index pairs  $(p, q), (p', q')$  are equivalent, then the corresponding random variables  $a_n(p, q), a_n(p', q')$  may be identical. Of course, for  $\mathcal{C} = \text{AII}, \text{DIII}, \text{C}, \text{CI}, \text{CII}$ , the symmetries of  $\mathbb{M}_n^{\mathcal{C}}$  must be realized as equivalence relations on  $\mathbb{I}_{2n} \times \mathbb{I}_{2n}$ , so one obtains the desired theorem on random elements of  $\mathbb{M}_n^{\mathcal{C}}$  by passage to a subsequence in Theorem 1.1. A more serious caveat is the following: Some of the blocks which make up matrices from  $\mathbb{M}_n^{\mathcal{C}}$  are skew symmetric, so their diagonal elements vanish, contradicting condition (1). While we will see that this problem can be circumvented in the cases at hand, the full blocks of zeroes in the chiral cases make it impossible to apply Theorem 1.1 for them as well.

To make the set-up for this section precise, let  $\mathcal{C}$  be a Wigner-Dyson or BdG class, write  $\delta = \delta^{\mathcal{C}}$  as in Section 2,  $\mathbb{J}_{\delta n}^{\mathcal{C}} := \{(p, q) \in \mathbb{I}_{\delta n}^2 : \text{pr}_{p,q}(\mathbb{M}_n^{\mathcal{C}}) \neq 0\}$ , where  $\text{pr}_{p,q}$  projects each element of  $\mathbb{M}_n^{\mathcal{C}}$  onto its  $(p, q)$ -entry. Consider an equivalence relation  $\sim_{\delta n}$  on  $\mathbb{I}_{\delta n}^2$  and a random matrix  $X_{\delta n} = (\frac{1}{\sqrt{\delta n}} a_{\delta n}(p, q))_{p,q=1,\dots,\delta n}$  such that the centered complex random variables  $a_{\delta n}(p_1, q_1), \dots, a_{\delta n}(p_j, q_j)$  are independent whenever  $(p_1, q_1), \dots, (p_j, q_j)$  belong to  $j$  distinct equivalence classes of the relation  $\sim_{\delta n}$ . Assume that conditions (W1), (W2), (W3) hold, with  $\mathbb{I}_{\delta n}$  in the place of  $\mathbb{I}_n$ , and that the moment condition (2) is satisfied. As to (1), it is required that it holds for all  $(p, q) \in \mathbb{J}_{\delta n}^{\mathcal{C}}$ . All realizations of the matrix  $X_{\delta n}$  are supposed to be elements of  $\mathbb{M}_n^{\mathcal{C}}$ . It is straightforward to verify that this assumption is compatible with conditions (W1),

(W2), (W3). So we may take the symmetries of the matrices for granted, and have some leeway for extra dependence between the matrix entries. Under these conditions, there holds

**THEOREM 3.1.** *If  $\mathcal{C}$  is a Wigner-Dyson or Bogolioubov-de Gennes class,  $\mathbb{E}(L_{\delta_n}(X_{\delta_n}))$  converges weakly to the semicircle law.*

It only remains to address the complication that random elements of  $\mathbb{M}_n$  may have up to  $4n$  entries which are identically zero. To see that the effect of this complication is asymptotically negligible, recall from the proof of Theorem 1.1 in [16] that the  $k$ -th moment of  $\mathbb{E}L_n$  vanishes if  $k$  is odd and is asymptotically equivalent to

$$\frac{1}{n^{l+1}} \sum_{\mathfrak{p} \in \text{NC}_2(2l)} \#\mathcal{S}_n(\mathfrak{p})$$

if  $k = 2l$  is even. Here the set  $\mathcal{S}_n(\mathfrak{p})$  consists of all pairs  $(\varphi, \psi)$ ,  $\varphi, \psi \in \mathcal{F}(k, n)$ , with the following properties:

- (i)  $\psi(j) = \varphi(j + 1)$  for all  $j \in \mathbb{I}_k$ , where  $k + 1$  is cyclically identified with 1.
- (ii)

$$\begin{aligned} (\varphi(i), \psi(i)) &= (\psi(j), \varphi(j)) \quad \text{if } i \sim_{\mathfrak{p}} j \quad \text{and} \\ (\varphi(i), \psi(i)) &\not\sim_n (\varphi(j), \psi(j)) \quad \text{if } i \not\sim_{\mathfrak{p}} j. \end{aligned}$$

For  $n \in \mathbb{N}$  fix  $E_n \subset \mathbb{I}_n^2$  with  $\#E_n = o(n^2)$ . Actually, what we have in mind is that  $E_n$  contains the  $O(n)$  diagonal places of skew blocks. For  $\nu \in \mathbb{I}_k$  set

$$\mathcal{S}_n^{(\nu)}(\mathfrak{p}) := \{(\varphi, \psi) \in \mathcal{S}_n(\mathfrak{p}) : (\varphi(\nu), \psi(\nu)) \in E_n\}$$

and

$$\mathcal{S}_n(\mathfrak{p})' = \bigcup_{\nu \in \mathbb{I}_k} \mathcal{S}_n^{(\nu)}(\mathfrak{p}).$$

Then the following lemma makes it possible to neglect the effect of the zero entries on the diagonals of the blocks:

**LEMMA 3.2.** *For  $\mathfrak{p} \in \text{NC}_2(2l)$ ,  $\#\mathcal{S}_n(\mathfrak{p})' = o(n^{l+1})$ .*

*Proof.* Since  $\#\mathcal{S}_n(\mathfrak{p})' \leq \sum_{\nu=1}^{2l} \#\mathcal{S}_n^{(\nu)}(\mathfrak{p})$ , it suffices to show that for all  $\nu$  one has  $\#\mathcal{S}_n^{(\nu)}(\mathfrak{p}) = o(n^{l+1})$ . To this end, we construct an element  $(\varphi, \psi) \in \mathcal{S}_n^{(\nu)}(\mathfrak{p})$ , starting with  $(\varphi(\nu), \psi(\nu))$  and proceeding cyclically. By cyclically permuting the index set, we may assume that  $\nu = 1$ . For  $(\varphi(1), \psi(1))$  we have  $o(n^2)$  choices.  $\varphi(2)$  is then fixed by property (i). If  $1 \sim_{\mathfrak{p}} 2$ , then  $\psi(2)$  is fixed by (ii). Otherwise, we have at most  $n$  choices. Similarly, for  $j \geq 2$ , once we have chosen  $\psi(j - 1)$ ,  $\varphi(j)$  is fixed, and  $\psi(j)$  is either fixed or we have at most  $n$  choices for it, depending on whether  $i \sim_{\mathfrak{p}} j$  for some  $1 \leq i < j$  or not. The latter case occurs  $l - 1$  times. So  $\#\mathcal{S}_n^{(\nu)}(\mathfrak{p}) \leq o(n^2) \times n^{l-1} = o(n^{l+1})$ . □

### 4 Sample covariance matrices

In this section we prove a limit theorem for the mean empirical eigenvalue distribution of sample covariance matrices with dependence. By these we understand matrices of the form  $X^*X$ , where  $X$  admits a certain amount of dependence among its entries,  $X^*$  is the conjugate transpose of  $X$ , and the entries of  $X$  are not necessarily Gaussian, but subject to certain conditions on their moments. In our context, this is preparatory work for the study of the chiral classes in Section 5, but it is of interest for its own sake. In the case that  $X$  has independent entries, the result is well-known, and we will take the combinatorial proof of Oravecz and Petz ([15]) as starting-point for an analysis in the spirit of Schenker and Schulz-Baldes ([16]).

For  $n \in \mathbb{N}$  let  $s(n), t(n) \in \mathbb{N}$  and suppose that there exist  $\kappa, \mu \in ]0, \infty[$  such that  $\lim_n \frac{s(n)}{n} = \kappa$  and  $\lim_n \frac{t(n)}{n} = \mu$ . The classical case is  $t(n) = n$ , but we will need this more general set-up in Section 5. Consider an equivalence relation  $\sim_n$  on  $\mathbb{I}_{s(n)} \times \mathbb{I}_{t(n)}$  and a random matrix  $X_n = (\frac{1}{\sqrt{n}} a_n(p, q))_{p=1, \dots, s(n), q=1, \dots, t(n)}$  such that the complex random variables  $a_n(p_1, q_1), \dots, a_n(p_j, q_j)$  are independent whenever  $(p_1, q_1), \dots, (p_j, q_j)$  belong to  $j$  distinct equivalence classes of the relation  $\sim_n$ . We impose the following conditions on  $\sim_n$ :

- (MP1)  $\max_p \#\{(q, p', q') \in \mathbb{I}_{t(n)} \times \mathbb{I}_{s(n)} \times \mathbb{I}_{t(n)} : (p, q) \sim_n (p', q')\} = o(n^2)$ .
- (MP2)  $\max_{p, q, p'} \#\{q' \in \mathbb{I}_{t(n)} : (p, q) \sim_n (p', q')\} \vee \max_{p, q, q'} \#\{p' \in \mathbb{I}_{s(n)} : (p, q) \sim_n (p', q')\} \leq B$ , where  $B$  is a finite constant.
- (MP3)  $\#\{(p, q, q') \in \mathbb{I}_{s(n)} \times \mathbb{I}_{t(n)}^2 : (p, q) \sim_n (p, q') \text{ and } q \neq q'\} = o(n^2)$  and  $\#\{(p, p', q) \in \mathbb{I}_{s(n)}^2 \times \mathbb{I}_{t(n)} : (p, q) \sim_n (p', q) \text{ and } p \neq p'\} = o(n^2)$ .

We assume that (1) and (2) hold. Under these assumptions we will prove the following theorem.

**THEOREM 4.1.** *As  $n \rightarrow \infty$ ,  $\mathbb{E}(L_n(X_n^*X_n))$  converges weakly to a probability measure with  $k$ -th moment equal to*

$$\sum_{i=1}^k \#(\text{NC}^{(i)}(k)) \kappa^i \mu^{k-i+1}. \tag{6}$$

*If  $\mu = 1$ , this limit is the Marčenko-Pastur distribution.*

We are going prove Theorem 4.1 via the method of moments. So we fix  $k \in \mathbb{N}$  and show that

$$\int x^k \mathbb{E}(L_n(X_n^*X_n))(dx) = \mathbb{E} \int x^k L_n(X_n^*X_n)(dx) = \frac{1}{n} \mathbb{E} \text{Tr}((X_n^*X_n)^k) \tag{7}$$

converges to (6) as  $n \rightarrow \infty$ . To write the trace in (7) in an explicit way, we use the notation introduced in Section 2. In the course of the technical proofs it will be convenient to identify the index set  $\mathbb{I}_{2k}$  with the cyclic group  $\mathbb{Z}/2k\mathbb{Z}$ , i.e., to identify  $2k + 1$  with 1 and so on.

The starting point for all that follows is the observation that

$$\frac{1}{n} \text{Tr}((X_n^*X_n)^k) = \frac{1}{n^{k+1}} \sum_{\varphi \in \mathcal{F}(m, s(n))} \sum_{\psi \in \mathcal{F}(n, t(n))} \prod_{\nu=1}^k \overline{a_n(\varphi(2\nu - 1), \psi(2\nu - 1))} a_n(\varphi(2\nu), \psi(2\nu)), \tag{8}$$

with  $\mathbf{m}, \mathbf{n}$  as in Section 2. Let  $\mathbf{p} \in \mathcal{P}(2k)$ . We say that  $(\varphi, \psi) \in \mathcal{F}(\mathbf{m}, s(n)) \times \mathcal{F}(\mathbf{n}, t(n))$  is associated to  $\mathbf{p}$ , and write  $(\varphi, \psi) \in S_n(\mathbf{p})$ , if for all  $i, j = 1, \dots, 2k$  there holds

$$i \sim_{\mathbf{p}} j \iff (\varphi(i), \psi(i)) \sim_n (\varphi(j), \psi(j)). \tag{9}$$

Writing

$$\Sigma_n(\mathbf{p}) := \sum_{(\varphi, \psi) \in S_n(\mathbf{p})} \prod_{\nu=1}^k \overline{a_n(\varphi(2\nu-1), \psi(2\nu-1))} a_n(\varphi(2\nu), \psi(2\nu)), \tag{10}$$

we obtain

$$\frac{1}{n} \text{Tr}((X_n^* X_n)^k) = \frac{1}{n^{k+1}} \sum_{\mathbf{p} \in \mathcal{P}(2k)} \Sigma_n(\mathbf{p}). \tag{11}$$

**LEMMA 4.2.** *For  $\mathbf{p} \in \mathcal{P}(2k)$  one has  $\#S_n(\mathbf{p}) \leq O(n^{|\mathbf{p}|+1})$ .*

*Proof.* We construct an element  $(\varphi, \psi) \in S_n(\mathbf{p})$ , proceeding from 1 to  $2k$  and giving rather coarse upper bounds on the number of choices in each step. We have  $s(n)$  choices for  $\varphi(1)$  and  $t(n)$  choices for  $\psi(1)$ . For  $\varphi$  to be constant on the blocks of  $\mathbf{m}$ , we must have  $\varphi(2) = \varphi(1)$ . If  $1 \sim_{\mathbf{p}} 2$ , then by (MP2) we have at most  $B$  choices for  $\psi(2)$ . Otherwise, we have at most  $t(n)$  choices. Note that  $\psi(2) = \psi(3)$ , since  $\psi$  is supposed to be constant on the blocks of  $\mathbf{n}$ . In the general case, for  $\nu = 2, 3, \dots, 2k-1$ , one of  $\varphi(\nu), \psi(\nu)$  is fixed, and for the other we have  $\leq B$  choices if  $\nu \sim_{\mathbf{p}} \nu'$  for some  $\nu' \in \mathbb{I}_{\nu-1}$  or at most  $s(n) \vee t(n)$  choices otherwise. This latter case occurs precisely  $|\mathbf{p}|-1$  times.  $(\varphi(2k), \psi(2k))$  is fixed by the requirement that  $\varphi(2k-1) = \varphi(2k)$  and  $\psi(2k) = \psi(1)$ . In total, we have at most  $s(n) t(n) (s(n) \vee t(n))^{|\mathbf{p}|-1} B^{2k-(|\mathbf{p}|+1)} = O(n^{|\mathbf{p}|+1})$  choices for  $(\varphi, \psi)$ .  $\square$

**LEMMA 4.3.** *For  $\mathbf{p} \in \mathcal{P}(2k)$ ,  $|\mathbb{E}\Sigma_n(\mathbf{p})| \leq \#S_n(\mathbf{p})c_k$ , with  $c_k$  independent of  $n$ .*

*Proof.* By Hölder's inequality and (2) one has

$$\begin{aligned} & \left| \mathbb{E} \prod_{\nu=1}^k \overline{a_n(\varphi(2\nu-1), \psi(2\nu-1))} a_n(\varphi(2\nu), \psi(2\nu)) \right| \\ & \leq \prod_{\nu=1}^{2k} \mathbb{E} \left( |a_n(\varphi(\nu), \psi(\nu))|^k \right)^{\frac{1}{k}} \leq c_k < \infty, \end{aligned}$$

with  $c_k$  independent of  $n, \varphi$  and  $\psi$ .  $\square$

**COROLLARY 4.4.**  $\mathbb{E}\Sigma_n(\mathbf{p}) = o(n^{k+1})$  unless  $\mathbf{p} \in \mathcal{P}_2(2k)$ .

*Proof.* If  $|\mathbf{p}| \leq k-1$ , then by Lemmata 4.2 and 4.3,  $|\mathbb{E}\Sigma_n(\mathbf{p})| \leq \#S_n(\mathbf{p}) c_k \leq O(n^k)c_k = o(n^{k+1})$ . If  $\mathbf{p}$  contains a block which consists of precisely one element  $\nu_0$ , say, then we have  $\mathbb{E}\Sigma_n(\mathbf{p}) = 0$ , because for any  $(\varphi, \psi) \in S_n(\mathbf{p})$  the random variable  $a_n(\varphi(\nu_0), \psi(\nu_0))$  is centered and by construction independent of  $\{a_n(\varphi(\nu), \psi(\nu)) : \nu \in \mathbb{I}_{2k} \setminus \{\nu_0\}\}$ . So  $\mathbb{E}\Sigma_n(\mathbf{p})$  vanishes if  $|\mathbf{p}| \geq k+1$ , or if  $|\mathbf{p}| = k$ , but  $\mathbf{p} \notin \mathcal{P}_2(2k)$ .  $\square$

The following lemma is a straightforward adaptation of a key step of [16] to the present context.

**LEMMA 4.5.** *If  $\mathbf{p} \in \mathcal{P}_2(2k)$  is crossing, then  $\#S_n(\mathbf{p}) = o(n^{k+1})$ .*

*Proof.* Suppose that  $\mathfrak{p}$  contains a nearest neighbour pair, i.e. a block of the form  $\{\nu, \nu + 1\}$ . Assume that  $\nu$  is odd.

If  $\psi(\nu) = \psi(\nu + 1)$ , then  $\psi(\nu - 1) = \psi(\nu + 2)$ . Writing  $\mathbb{J} = \mathbb{I}_{2k} \setminus \{\nu, \nu + 1\}$ , we see that  $(\varphi|_{\mathbb{J}}, \psi|_{\mathbb{J}}) \in S_n(\tilde{\mathfrak{p}})$ , where  $\tilde{\mathfrak{p}}$  is the partition of  $\mathbb{J}$  whose blocks are those of  $\mathfrak{p}$  except for  $\{\nu, \nu + 1\}$ . Given  $(\tilde{\varphi}, \tilde{\psi}) \in S_n(\tilde{\mathfrak{p}})$ , there are  $s(n)$  choices for  $\varphi(\nu) = \varphi(\nu + 1)$ , hence  $s(n)$  possible extensions to  $(\varphi, \psi) \in S_n(\mathfrak{p})$ , since  $\psi(\nu), \psi(\nu + 1)$  are determined by  $\tilde{\psi}$ .

If  $\psi(\nu) \neq \psi(\nu + 1)$ , then by (MP3) there are only  $o(n^2)$  choices for the triplet  $(\varphi(\nu) = \varphi(\nu + 1), \psi(\nu), \psi(\nu + 1))$ . As in the proof of Lemma 4.2, we see that there are at most  $O(n^{k-1})$  choices for the remaining values of  $\varphi, \psi$ , since  $\tilde{\mathfrak{p}}$  consists of  $k - 1$  pairs.

Since we may argue analogously for  $\nu$  even, we have shown that

$$\#S_n(\mathfrak{p}) \leq O(n) \#S_n(\tilde{\mathfrak{p}}) + o(n^{k+1}). \tag{12}$$

Since  $\mathfrak{p}$  was assumed to be crossing, iterating this argument yields

$$\#S_n(\mathfrak{p}) \leq O(n^r) \#S_n(\mathfrak{p}') + o(n^{k+1}), \tag{13}$$

where  $\mathfrak{p}'$  contains no nearest neighbour pair. Upon relabelling, we may regard  $\mathfrak{p}'$  as an element of  $\mathcal{P}_2(2(k - r))$ , where  $k - r \geq 2$ . Note that we may have  $\mathfrak{p} = \mathfrak{p}'$ , whence it is possible that  $r = 0$ .

Let  $\lambda$  be the minimal  $l > 0$  such that there exists  $\nu \in \mathbb{I}_{2(k-r)}$  with  $\{\nu, \nu + l\} \in \mathfrak{p}'$  (where addition takes place in  $\mathbb{Z}/2(k - r)\mathbb{Z}$ ). Observe that  $\lambda \geq 2$ . Now, if  $\nu$  has the property that  $\{\nu, \nu + \lambda\} \in \mathfrak{p}'$ , then all elements of  $\{\nu + 1, \dots, \nu + \lambda - 1\}$  are paired with partners outside  $\{\nu, \nu + 1, \dots, \nu + \lambda\}$ . We find an upper bound for  $\#S_n(\mathfrak{p}')$  as follows. By (MP1), we have  $s(n)$  choices for  $\varphi(\nu)$  and  $o(n^2)$  choices for the triplet  $(\psi(\nu), \varphi(\nu + \lambda), \psi(\nu + \lambda))$ . By construction, going from left to right through  $\nu + l$  ( $l = 1, 2, \dots, \lambda - 2$ ), either  $\varphi(\nu + l)$  is fixed and there are at most  $t(n)$  choices for  $\psi(\nu + l)$ , or  $\psi(\nu + l)$  is fixed and there are at most  $s(n)$  choices for  $\varphi(\nu + l)$ . According to whether  $\nu$  is even or odd, we must have either  $\varphi(\nu + \lambda - 1) = \varphi(\nu + \lambda)$  and  $\psi(\nu + \lambda - 1) = \psi(\nu + \lambda - 2)$  or  $\psi(\nu + \lambda - 1) = \psi(\nu + \lambda)$  and  $\varphi(\nu + \lambda - 1) = \varphi(\nu + \lambda - 2)$ . So we have at most  $O(n^{\lambda-1})o(n^2)$  choices for the restrictions of  $\varphi$  and  $\psi$  to  $\{\nu, \nu + 1, \dots, \nu + \lambda\}$ . Going cyclically through the complement of this set, starting with  $\nu + \lambda + 1$ , in each step one of  $\varphi(\nu + \lambda + l), \psi(\nu + \lambda + l)$  is fixed, and there are at most  $B$  resp.  $O(n)$  choices for the other, depending on whether  $\nu + \lambda + l$  is paired with one of the previously considered points or not. This latter case occurs exactly  $k - r - \lambda$  times.

Putting all this together with (13), we arrive at

$$\#S_n(\mathfrak{p}) \leq O(n^r) o(n^{\lambda+1}) O(n^{k-r-\lambda}) + o(n^{k+1}) = o(n^{k+1}).$$

□

**LEMMA 4.6.** *For all  $\mathfrak{p} \in \text{NC}_2(2k)$  there holds  $l_{\mathfrak{p}} := |\mathfrak{p} \vee \mathfrak{m}| + |\mathfrak{p} \vee \mathfrak{n}| = k + 1$ , with  $\vee$  as in Section 2.*

*Proof.* For  $k = 1$ , the only (noncrossing pair) partition  $\mathfrak{p} \in \mathcal{P}(2)$  is  $\{\{1, 2\}\}$ , which satisfies  $l_{\mathfrak{p}} = 2$ . Suppose that  $k \geq 2$  and that the claim is true for  $k - 1$ . Note that any  $\mathfrak{p} \in \text{NC}_2(2k)$  contains a block of the form  $\{\nu, \nu + 1\}$ . Now consider the partition  $\tilde{\mathfrak{p}}$  of  $\mathbb{I}_{2k} \setminus \{\nu, \nu + 1\}$ , whose blocks are those of  $\mathfrak{p}$  except for  $\{\nu, \nu + 1\}$ . Define  $\tilde{\mathfrak{m}}, \tilde{\mathfrak{n}} \in \mathbb{I}_{2k} \setminus \{\nu, \nu + 1\}$  in the obvious way. We claim that

$$l_{\tilde{\mathfrak{p}}} = l_{\mathfrak{p}} - 1. \tag{14}$$

To see this for  $\nu$  even, note that in this case  $\{\nu, \nu + 1\}$  is also a block of  $\mathbf{n}$ . This means that it is a block of  $\mathbf{p} \vee \mathbf{n}$ , hence  $|\tilde{\mathbf{p}} \vee \tilde{\mathbf{n}}| = |\mathbf{p} \vee \mathbf{n}| - 1$ . On the other hand,  $\mathbf{p} \vee \mathbf{m}$  has a block which contains  $\nu - 1, \nu, \nu + 1, \nu + 2$ . This block does not split on eliminating  $\{\nu, \nu + 1\}$ , since  $\{\nu - 1, \nu + 2\}$  is a block of  $\tilde{\mathbf{m}}$ . So we arrive at  $|\tilde{\mathbf{p}} \vee \tilde{\mathbf{m}}| = |\mathbf{p} \vee \mathbf{m}|$ . For  $\nu$  odd, the argument is analogous, yielding (14). Hence by induction,  $l_{\mathbf{p}} = l_{\tilde{\mathbf{p}}} + 1 = ((k - 1) + 1) + 1 = k + 1$ .  $\square$

The following is evident:

**LEMMA 4.7.** *For  $\mathbf{p} \in \text{NC}_2(2k)$ , each of the blocks of  $\mathbf{p}$  consists of exactly one odd and exactly one even number.*

**LEMMA 4.8.**  $\#\{\mathbf{p} \in \text{NC}_2(2k) : |\mathbf{p} \vee \mathbf{m}| = j\} = \#\text{NC}^{(j)}(k)$ .

*Proof.* We may identify  $\mathcal{P}(k)$  with  $\{\mathbf{p} \in \mathcal{P}(2k) : \text{any block of } \mathbf{p} \text{ is the union of blocks of } \mathbf{m}\}$ . So  $\mathbf{p} \mapsto \mathbf{p} \vee \mathbf{m}$  maps  $\mathcal{P}(2k)$  onto  $\mathcal{P}(k)$ . It is easy to see that if  $\mathbf{p} \vee \mathbf{m}$  is crossing, so is  $\mathbf{p}$ . In fact  $\text{NC}_2(2k)$  is mapped bijectively onto  $\text{NC}(k)$ . To see this, it suffices to show that  $\mathbf{p} \mapsto \mathbf{p} \vee \mathbf{m}$  is injective, as it is known that  $\#\text{NC}(k) = \#\text{NC}_2(2k)$  (see [14, Remark 9.5]). A block of  $\mathbf{p} \vee \mathbf{m}$  is of the form  $b_J = \bigcup_{\nu \in J} \{2\nu - 1, 2\nu\}$  for some  $J \subseteq \mathbb{I}_k$ . We have to show that there exists precisely one  $\tilde{\mathbf{p}} \in \text{NC}_2(b_J)$  such that  $\tilde{\mathbf{p}} \vee \{\{2\nu - 1, 2\nu\} : \nu \in J\} = \{b_J\}$ . Since our aim is to show that a pairing of the elements of  $b_J$  with certain properties is uniquely determined, the embedding of  $b_J$  into  $\mathbb{I}_{2k}$  is irrelevant, and we may assume that  $b_J = \mathbb{I}_{2r}$  for some  $r \leq k$ . Let us start by finding a partner for 1. By Lemma 4.7, the partner must be even,  $1 \sim_{\tilde{\mathbf{p}}} 2s$ , say. Assume that  $s < r$ . Since we wish to construct a noncrossing  $\tilde{\mathbf{p}}$ , no  $1 < \nu < 2s$  can be paired with any  $\nu' > 2s$ . On the other hand,  $2s \not\sim_{\mathbf{m}} 2s + 1$ , so  $2s$  and  $2s + 1$  lie in distinct blocks of  $\tilde{\mathbf{p}} \vee \mathbf{m}$ , contradicting the requirement that  $\tilde{\mathbf{p}} \vee \mathbf{m} = \{\mathbb{I}_{2r}\}$ . Consequently, we must have  $1 \sim_{\tilde{\mathbf{p}}} 2r$ . The partner of 2 must be odd. We claim that necessarily  $2 \sim_{\tilde{\mathbf{p}}} 3$ . Otherwise the partner of 2 is  $2s - 1$ ,  $s \geq 3$ . Then for  $\tilde{\mathbf{p}}$  to be noncrossing, it is necessary that  $\{3, 4, \dots, 2s - 2\}$  be a union of blocks of  $\tilde{\mathbf{p}}$ . But  $2 \not\sim_{\mathbf{m}} 3$  and  $2s - 2 \not\sim_{\mathbf{m}} 2s - 1$ . So  $\{3, 4, \dots, 2s - 2\}$  splits into at least two distinct blocks of  $\tilde{\mathbf{p}} \vee \mathbf{m}$ , violating our conditions on  $\tilde{\mathbf{p}}$ . Deleting  $\{2, 3\}$  and relabelling, we inductively see that  $\tilde{\mathbf{p}} = \{\{1, 2r\}, \{2, 3\}, \{4, 5\}, \dots, \{2r - 2, 2r - 1\}\}$ , hence is uniquely determined.  $\square$

Define

$$S_n^\vee(\mathbf{p}) := \{(\varphi, \psi) : \varphi \in \mathcal{F}(\mathbf{p} \vee \mathbf{m}, s(n)), \psi \in \mathcal{F}(\mathbf{p} \vee \mathbf{n}, t(n))\}$$

and observe that  $S_n^\vee(\mathbf{p}) \subset S_n(\mathbf{p})$ . Write

$$S_n^\wedge(\mathbf{p}) := S_n(\mathbf{p}) \setminus S_n^\vee(\mathbf{p})$$

and

$$\Sigma_n^\vee(\mathbf{p}) := \sum_{(\varphi, \psi) \in S_n^\vee(\mathbf{p})} \prod_{\nu=1}^k \overline{a_n(\varphi(2\nu - 1), \psi(2\nu - 1))} a_n(\varphi(2\nu), \psi(2\nu)).$$

**LEMMA 4.9.** *For  $\mathbf{p} \in \text{NC}_2(2k)$ ,  $\#S_n^\wedge(\mathbf{p}) = o(n^{k+1})$ .*

*Proof.* Since  $\mathbf{p}$  is noncrossing, we find  $\nu \in \mathbb{I}_{2k-1}$  such that  $\nu \sim_{\mathbf{p}} \nu + 1$ . Suppose that  $\nu$  is odd. Then for any  $(\varphi, \psi) \in S_n^\wedge(\mathbf{p})$  we have  $\varphi(\nu) = \varphi(\nu + 1)$ . If  $\psi(\nu) \neq \psi(\nu + 1)$ , then by condition (MP3) there are  $o(n^2)$  possibilities for the triplet  $(\varphi(\nu), \psi(\nu), \psi(\nu + 1))$ , and one sees as in Lemma 4.5 that there are  $O(n^{k-1})$  choices for  $(\varphi, \psi)$  on  $\mathbb{J} := \mathbb{I}_{2k} \setminus \{\nu, \nu + 1\}$ . If  $\psi(\nu) = \psi(\nu + 1)$ , then  $\psi(\nu - 1) = \psi(\nu + 2)$ . If  $\tilde{\mathbf{p}}$  is the partition of  $\mathbb{J}$  induced by  $\mathbf{p}$ , then  $(\varphi|_{\mathbb{J}}, \psi|_{\mathbb{J}}) \in S_n^\wedge(\tilde{\mathbf{p}})$ . In

this case, hence,  $\#S_n^\wedge(\mathbf{p}) \leq O(n) \#S_n^\wedge(\tilde{\mathbf{p}})$ . Putting both cases for  $\psi(\nu), \psi(\nu + 1)$  together, we obtain

$$\#S_n^\wedge(\mathbf{p}) \leq O(n) \#S_n^\wedge(\tilde{\mathbf{p}}) + o(n^{k+1}).$$

This was proven for  $\nu$  odd. In view of the symmetry of (MP3), the proof for  $\nu$  even is analogous, with the roles of  $\varphi$  and  $\psi$  interchanged. Since  $(\varphi, \psi) \in S_n^\wedge(\mathbf{p})$ , iteration of this process will finally lead to a nearest neighbour pair  $\nu \sim_{\mathbf{p}} \nu + 1$  such that  $\varphi(\nu) \neq \varphi(\nu + 1)$  or  $\psi(\nu) \neq \psi(\nu + 1)$ . So we end up with

$$\#S_n^\wedge(\mathbf{p}) \leq O(n^{k-1})o(n^2) + o(n^{k+1}) = o(n^{k+1}).$$

□

Putting the ingredients together, we have that

$$\frac{1}{n} \mathbb{E}(\text{Tr}((X^* X)^k)) = \frac{1}{n^{k+1}} \sum_{\mathbf{p} \in \mathcal{P}(2k)} \mathbb{E}\Sigma_n(\mathbf{p})$$

is by Corollary 4.4 and Lemmata 4.3, 4.5, and 4.6 asymptotically equivalent to

$$\frac{1}{n^{k+1}} \sum_{j=1}^k \sum_{\substack{\mathbf{p} \in \text{NC}_2(2k), \\ |\mathbf{p} \vee \mathbf{m}|=j}} \mathbb{E}\Sigma_n(\mathbf{p}). \tag{15}$$

By Lemmata 4.9 and 4.3 we may replace  $\mathbb{E}\Sigma_n(\mathbf{p})$  by  $\mathbb{E}\Sigma_n^\vee(\mathbf{p})$  in (15). Recall that for  $(\varphi, \psi) \in S_n^\vee(\mathbf{p})$ ,  $\varphi$  and  $\psi$  are *a fortiori* constant on the blocks of  $\mathbf{p}$ . Comparing Lemma 4.7 with (10), one sees that this implies that given  $(\varphi, \psi) \in S_n^\vee(\mathbf{p})$ , a block of  $\mathbf{p}$  corresponds to a matrix element and its complex conjugate. Invoking (1) and Lemma 4.6, we see that (15) is asymptotically equivalent to

$$\frac{1}{n^{k+1}} \sum_{j=1}^k \sum_{\substack{\mathbf{p} \in \text{NC}_2(2k), \\ |\mathbf{p} \vee \mathbf{m}|=j}} \sum_{(\varphi, \psi) \in S_n^\vee(\mathbf{p})} 1 = \frac{1}{n^{k+1}} \sum_{j=1}^k \sum_{\substack{\mathbf{p} \in \text{NC}_2(2k), \\ |\mathbf{p} \vee \mathbf{m}|=j}} s(n)^j t(n)^{k-j+1},$$

which tends, as  $n \rightarrow \infty$ , to

$$\sum_{j=1}^k \#\{\mathbf{p} \in \text{NC}_2(2k) : |\mathbf{p} \vee \mathbf{m}| = j\} \kappa^j \mu^{k-j+1} = \sum_{j=1}^k \#\text{NC}^{(j)}(k) \kappa^j \mu^{k-j+1},$$

where the last equality follows from Lemma 4.8.

## 5 The chiral classes

In this section we apply our result about sample covariance matrices to random elements of the spaces  $\mathbb{M}_n^{\mathcal{C}}$  from Section 2, where  $\mathcal{C} = \text{BDI}, \text{AIII}$  or  $\text{CII}$ . It is convenient to consider the AIII case first. It consists of matrices of the form

$$\mathcal{X}_n = \begin{pmatrix} 0 & X_n \\ X_n^* & 0 \end{pmatrix} \in \mathbb{C}^{n \times n},$$

with  $X_n \in \mathbb{C}^{s(n) \times t(n)}$ , hence  $s(n) + t(n) = n$ . We assume that  $\lim_{n \rightarrow \infty} \frac{s(n)}{n} = \kappa \in ]0, 1[$ , hence  $\lim_{n \rightarrow \infty} \frac{t(n)}{n} = 1 - \kappa =: \mu$ . Note that this framework is more restrictive than the one considered in Section 4. But these restrictions naturally arise if one considers  $X_n$  as a subblock of an element of  $\mathbb{M}_n^{\text{AIII}}$ , whence  $n$  is the natural parameter for asymptotics. Observe that

$$\text{Tr} \begin{pmatrix} 0 & X_n \\ X_n^* & 0 \end{pmatrix}^k = \begin{cases} 0 & \text{if } k \text{ odd} \\ 2 \text{Tr}((X_n^* X_n)^l) & \text{if } k = 2l \text{ even.} \end{cases} \tag{16}$$

Assuming that  $X_n$  satisfies conditions (MP1), (MP2), (MP3) of Section 4, Theorem 4.1 implies that as  $n \rightarrow \infty$ , the  $2l$ -th moment of  $EL_n(\mathcal{X}_n)$  converges to

$$2 \sum_{j=1}^l \# \text{NC}^{(j)}(l) \kappa^j (1 - \kappa)^{l-j+1}. \tag{17}$$

The special case where the entries of  $X_n$  take purely imaginary values yields the same limit for class BDI. Since the extra symmetries of the CII case are compatible with (MP1), (MP2) and (MP3), we obtain the same limit for this class, too.

In [8], in the special case of chiral random matrices with independent Gaussian entries, the empirical limit distribution  $\mu_{\text{ch},2}$  of the squared eigenvalues was determined as

$$|1 - 2\kappa| \delta_0 + 1_{[a,b]}(x) \frac{1}{\pi x} \sqrt{(x-a)(b-x)} \, dx, \tag{18}$$

where

$$a = 1 - 2\sqrt{\kappa(1 - \kappa)}, \quad b = 1 + 2\sqrt{\kappa(1 - \kappa)}. \tag{19}$$

Note that (18) differs from the corresponding formula in [8], since a different definition of  $L_n$  is used in that paper, and since (1) above imposes a condition on complex rather than real variances. Now, the elegant approach of Haagerup and Thorbjørnsen ([10]) to the moments of the Marčenko-Pastur distribution can be easily adapted to  $\mu_{\text{ch},2}$ , to the effect that the  $l$ -th moment of  $\mu_{\text{ch},2}$  is indeed given by (17). In fact, for  $l \geq 1$ ,

$$\begin{aligned} \int x^l \mu_{\text{ch},2}(dx) &= \frac{1}{\pi} \int_{1-2\sqrt{\kappa(1-\kappa)}}^{1+2\sqrt{\kappa(1-\kappa)}} x^{l-1} \sqrt{4\kappa(1-\kappa) - (x-1)^2} \, dx \\ &= \frac{4\kappa(1-\kappa)}{\pi} \int_{-\pi}^0 \sin^2 \theta (2\sqrt{\kappa(1-\kappa)} \cos \theta + 1)^{l-1} \, d\theta \\ &= \frac{2\kappa(1-\kappa)}{\pi} \int_{-\pi}^{\pi} \sin^2 \theta (2\sqrt{\kappa(1-\kappa)} \cos \theta + 1)^{l-1} \, d\theta. \end{aligned} \tag{20}$$

Setting  $g(\theta) = (\sqrt{\kappa} + \sqrt{1-\kappa} e^{i\theta})^{l-1}$  and observing that  $\sin^2 \theta = \frac{1}{2}(1 - \cos 2\theta)$ , we see that (20) can be written as

$$\frac{\kappa(1-\kappa)}{\pi} \int_{-\pi}^{\pi} \text{Re}(1 - e^{i2\theta}) |g(\theta)|^2 \, d\theta$$

or as

$$2\kappa(1-\kappa) \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} |g(\theta)|^2 \, d\theta - \text{Re} \left( \frac{1}{2\pi} \int_{-\pi}^{\pi} h(\theta) \overline{k(\theta)} \, d\theta \right) \right)$$

with  $h(\theta) = e^{i\theta}g(\theta)$ ,  $k(\theta) = e^{-i\theta}g(\theta)$ . Invoking Parseval's formula and elementary computations with binomial coefficients, we obtain

$$\begin{aligned} \int x^l \mu_{\text{ch},2}(dx) &= 2 \sum_{j=0}^{l-1} \left\{ \binom{l-1}{j}^2 - \binom{l-1}{j-1} \binom{l-1}{j+1} \right\} \kappa^{j+1} (1-\kappa)^{l-j} \\ &= \frac{2}{l} \sum_{j=1}^l \binom{l}{j} \binom{l}{j-1} \kappa^j (1-\kappa)^{l+1-j} \\ &= 2 \sum_{j=1}^l \#(\text{NC}^{(j)}(l)) \kappa^j (1-\kappa)^{l+1-j}. \end{aligned}$$

A reference for the last equality is [14, Cor. 9.13]. In view of (16), we have

**THEOREM 5.1.** *If  $\mathcal{C}$  is a chiral class, then  $\mathbb{E}(L_{\delta_n}(X_{\delta_n}))$  converges to a probability measure  $\mu_{\text{ch}}$  on  $\mathbb{R}$  given by*

$$|1 - 2\kappa|\delta_0 + 1_{[a,b]}(x^2) \frac{2}{\pi x} \sqrt{(x^2 - a)(b - x^2)} dx$$

with  $a, b$  as in (19) above.

## References

- [1] A. Altland and B. Simons, *Condensed matter field theory*, Cambridge UP, Cambridge, 2006.
- [2] A. Altland and M. Zirnbauer, *Nonstandard symmetry classes in mesoscopic normal/superconducting hybrid structures*, Physical Review B **55** (1997), no. 2, 1142–1161.
- [3] G. Anderson and O. Zeitouni, *A law of large numbers for finite-range dependent random matrices*, arXiv.org:math/0609364.
- [4] Z. D. Bai and J. Yao, *On the convergence of the spectral empirical process of Wigner matrices*, Bernoulli **11** (2005), 1059 – 1092. MR2189081
- [5] B. Collins and M. Stolz, *Borel theorems for random elements of the classical compact symmetric spaces*, Ann. of Probab. **36** (2008), 876 – 895
- [6] E. Dueñez, *Random matrix ensembles associated to compact symmetric spaces*, Comm. Math. Phys. **244** (2004), no. 1, 29–61. MR2029949
- [7] F. J. Dyson, *The threefold way. Algebraic structure of symmetry groups and ensembles in quantum mechanics*, J. Mathematical Phys. **3** (1962), 1199–1215. MR0177643
- [8] P. Eichelsbacher and M. Stolz, *Large deviations for random matrix ensembles in mesoscopic physics*, math.PR/0610811, to appear in Markov Process. Related Fields
- [9] P. Forrester, *Log-gases and random matrices*, book in progress, see <http://www.ms.unimelb.edu.au/~matpjf/matpjf.html>.
- [10] U. Haagerup and S. Thorbjørnsen, *Random matrices with complex Gaussian entries*, Expo. Math. **21** (2003), no. 4, 293–337. MR2022002

- 
- [11] P. Heinzner, A. Huckleberry, and M. R. Zirnbauer, *Symmetry classes of disordered fermions*, Commun. Math. Phys. **257** (2005), 725–771. MR2164950
  - [12] F. Hiai and D. Petz, *The semicircle law, free random variables and entropy*, Mathematical Surveys and Monographs, vol. 77, American Mathematical Society, Providence, RI, 2000. MR1746976
  - [13] N. M. Katz and P. Sarnak, *Zeroes of zeta functions and symmetry*, Bull. Amer. Math. Soc. **36** (1999), 1–26. MR1640151
  - [14] A. Nica and R. Speicher, *Lectures on the combinatorics of free probability*, London Mathematical Society Lecture Note Series, vol. 335, Cambridge University Press, Cambridge, 2006. MR2266879
  - [15] F. Oravecz and D. Petz, *On the eigenvalue distribution of some symmetric random matrices*, Acta Sci. Math. (Szeged) **63** (1997), no. 3-4, 383–395. MR1480488
  - [16] J. H. Schenker and H. Schulz-Baldes, *Semicircle law and freeness for random matrices with symmetries or correlations*, Math. Res. Lett. **12** (2005), no. 4, 531–542. MR2155229
  - [17] J. H. Schenker and H. Schulz-Baldes, *Gaussian fluctuations for random matrices with correlated entries*, Int. Math. Res. Not. 2007, Article ID rnm047, doi:10.1093/imrn/rnm047.
  - [18] E. P. Wigner, *On the distribution of the roots of certain symmetric matrices*, Ann. of Math. (2) **67** (1958), 325–327. MR0095527