

# The critical branching random walk in a random environment dies out

Olivier Garet\*      Régine Marchand†

## Abstract

We study the possibility for branching random walks in random environment (BRWRE) to survive. The particles perform simple symmetric random walks on the  $d$ -dimensional integer lattice, while at each time unit, they split into independent copies according to time-space i.i.d. offspring distributions. As noted by Comets and Yoshida, the BRWRE is naturally associated with the directed polymers in random environment (DPRE), for which the quantity  $\Psi$  called the free energy is well studied. Comets and Yoshida proved that there is no survival when  $\Psi < 0$  and that survival is possible when  $\Psi > 0$ . We proved here that, except for degenerate cases, the BRWRE always die when  $\Psi = 0$ . This solves a conjecture of Comets and Yoshida.

**Keywords:** branching random walk; random environment; survival; critical behavior; renormalization; block construction.

**AMS MSC 2010:** 60K35; 82B43.

Submitted to ECP on November 15, 2012, final version accepted on January 30, 2013.

## 1 Introduction

### 1.1 Branching random walks in random environment

Let us introduce the model. We write  $\mathbb{N} = \{0, 1, 2, \dots\}$  and  $\mathbb{N}^* = \{1, 2, \dots\}$ .

To each  $(t, x) \in \mathbb{N} \times \mathbb{Z}^d$ , we associate a distribution  $q_{t,x} = (q_{t,x}(k))_{k \in \mathbb{N}}$  on the integers; the family  $\mathbf{q} = (q_{t,x})_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$  is called an environment. We denote by  $\Lambda = \mathcal{P}(\mathbb{N})^{\mathbb{N} \times \mathbb{Z}^d}$  the space of environments, where  $\mathcal{P}(\mathbb{N})$  is the set of distributions on  $\mathbb{N}$ .

Given an environment  $\mathbf{q} = (q_{t,x})_{t \in \mathbb{N}, x \in \mathbb{Z}^d} \in \Lambda$ , we define the branching random walk in environment  $\mathbf{q}$  as the following dynamics:

- At time  $t = 0$ , there is one particle at the origin  $x = 0$ .
- Each particle, located at site  $x \in \mathbb{Z}^d$  at time  $t$ , jumps at time  $t + 1$  to one of the  $2d$  neighbors of  $x$  chosen uniformly; upon arrival, it dies and is replaced by  $k$  new particles with probability  $q_{t,x}(k)$ . The number of newborn particles is independent of the jump, and all these variables, indexed by the full population at time  $t$ , are independent.

Take a random  $\mathbf{q}$ , we obtain a branching random walk in random environment (BRWRE). We assume here that  $\mathbf{q} = (q_{t,x})_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$  is a  $\mathcal{P}(\mathbb{N})$ -valued i.i.d. sequence with common distribution  $\gamma$  and we denote by  $\mathbb{P}^\gamma$  the annealed law of the branching random walk in this random environment. We also note  $\mathbb{E}^\gamma$  for the expectation with respect to  $\mathbb{P}^\gamma$ .

---

\*Université de Lorraine, Institut Élie Cartan de Lorraine, CNRS UMR 7502, France.  
E-mail: Olivier.Garet, Regine.Marchand@univ-lorraine.fr

Since the environment not only depends on the sites but also on the time, we can be more specific and say that we work with a random space-time or random dynamic environment. This is the model studied in Yoshida [12], Hu and Yoshida [8], and also Comets and Yoshida [3]. A natural question is to characterize the laws  $\gamma$  that allow the branching random walk to survive.

For  $(t, x) \in \mathbb{N} \times \mathbb{Z}^d$  and fixed  $\mathbf{q}$ , we introduce the mean progeny at site  $(t, x)$ :

$$m_{t,x} = \sum_{k \in \mathbb{N}} k q_{t,x}(k). \tag{1.1}$$

From here on, we assume that

$$\mathbb{P}^\gamma(m_{0,0} + m_{0,0}^{-1}) < +\infty, \tag{1.2}$$

$$\mathbb{P}^\gamma(q_{0,0}(0) > 0) > 0 \quad \text{and} \quad \mathbb{P}^\gamma(q_{0,0}(0) + q_{0,0}(1) < 1) > 0. \tag{1.3}$$

Assumptions (1.3) are intended to avoid obvious survival and obvious extinction.

### 1.2 The Comets–Yoshida Theorem and the associated conjecture

The Comets–Yoshida Theorem [3] relates the survival of the BRWRE with a functional on an associated directed polymer in random environment as follows: define on a probability space  $(\Omega_S, \mathcal{F}_S, P_S)$  a simple symmetric random walk  $(S_t)_{t \geq 0}$  on  $\mathbb{Z}^d$  starting from  $S_0 = 0$ . The partition function of the directed polymer in random environment  $\mathbf{q}$  is given by

$$Z_t = \int \prod_{u=0}^{t-1} m_{u, S_u} \, dP_S,$$

with  $m_{t,x}$  as in (1.1). It is easy to see (e.g., [12, Lemma 1.4]) that  $Z_t$  is the expectation of the number of particles of the BRWRE living at time  $t$ , knowing the random environment  $\mathbf{q} = (q_{t,x})_{(t,x) \in \mathbb{N} \times \mathbb{Z}^d}$ . Note that (1.2), combined with the inequality  $|\log u| \leq u \vee u^{-1}$  for  $u > 0$ , implies that

$$\forall t \in \mathbb{N}^* \quad \mathbb{E}^\gamma(|\log Z_t|) < +\infty$$

**Proposition 1.1** (Comets–Yoshida). *The following limit exists*

$$\Psi(\gamma) \stackrel{\text{def.}}{=} \lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E}^\gamma(\log Z_t).$$

Moreover,

- If  $\Psi(\gamma) < 0$ , then  $\mathbb{P}^\gamma(\text{survival}) = 0$ .
- If  $\Psi(\gamma) > 0$  and  $\mathbb{E}^\gamma[-\log(1 - q_{0,0}(0))] < +\infty$ , then  $\mathbb{P}^\gamma(\text{survival} | \mathbf{q}) > 0$  for  $\mathbb{P}^\gamma$ -almost every  $\mathbf{q}$ .

The number  $\Psi$  is called the free energy of the polymer. The aim of this article is to prove the following theorem, stated as a conjecture by Comets and Yoshida [3]:

**Theorem 1.2.** *Assume that (1.2) and (1.3) are satisfied. Then*

$$(\Psi(\gamma) = 0) \implies (\mathbb{P}^\gamma(\text{survival}) = 0).$$

It is a well-known fact that growth processes often die out in the critical case: it obviously holds for the Galton–Watson tree, but it also holds for more complicated models. This has been proved for the contact process in dimension  $d$  by Bezuidenhout and Grimmett in a famous paper entitled *The critical contact process dies out* [1]. Recently,

Steif and Warfheimer [10] generalized this result to the case of the contact process in randomly evolving environment introduced by Broman [2]. We also proved in [5] that is true for a class of dependent oriented percolations on  $\mathbb{Z}^d$ .

Remark: In Comets and Yoshida [3], it is proved, provided that  $\Psi > 0$ , that  $\Psi$  is the asymptotic growth rate of the total population of the BRWRE when it survives. With Theorem 1.2, we get that the total population grows exponentially as soon as it survives. A similar result is proved for a class of growth models in random environment by Fukushima and Yoshida [4].

### 1.3 Strategy of the proof

Recall that  $\gamma$  is the common law of the i.i.d sequence  $(q_{t,x})_{t \geq 0, x \in \mathbb{Z}^d}$  of distributions on  $\mathbb{N}$ . For  $\rho \in (0, 1]$ , we define a perturbation  $\gamma^\rho$  of  $\gamma$ . Precisely, we define  $\gamma^\rho$  as the image of  $\gamma$  by the map:

$$q \mapsto \rho q + (1 - \rho)\delta_0,$$

where  $\delta_0 \in \mathcal{P}(\mathbb{N})$  is the Dirac mass on 0.

Under  $\gamma^\rho$ , the environment is then less favorable to births of particles than under  $\gamma$ : under  $\mathbb{P}^{\gamma^\rho}$ , each particle sitting at  $x$  at time  $t$  flips an extra coin to decide whether it tries to have children or not.

**Lemma 1.3.**  $\forall \rho \in (0, 1] \quad \forall \gamma \in (0, 1] \quad \Psi(\gamma^\rho) = \log \rho + \Psi(\gamma)$ .

*Proof.* Note  $\mathbb{Q}^\gamma = \gamma^{\otimes \mathbb{N} \times \mathbb{Z}^d}$ . Let  $q_{t,x}^\rho(k) = \rho q_{t,x}(k) + (1 - \rho)\delta_0(k)$ . Obviously, the law of the field  $(q_{t,x}^\rho)$  under  $\mathbb{Q}^\gamma$  is  $\mathbb{Q}^{\gamma^\rho}$ . If we define, for  $(t, x) \in \mathbb{N} \times \mathbb{Z}^d$ ,

$$m_{t,x}^\rho = \sum_{k \in \mathbb{N}} k q_{t,x}^\rho(k) \quad \text{and} \quad Z_t^\rho = \int \prod_{u=0}^{t-1} m_{u,S_u}^\rho \, dP_S,$$

it is easy to see that  $m_{t,x}^\rho = \rho m_{t,x}$ , that  $Z_t^\rho = \rho^t Z_t$ , and finally

$$\Psi(\gamma^\rho) = \lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E}^\gamma[\log Z_t^\rho] = \log \rho + \lim_{t \rightarrow \infty} \frac{1}{t} \mathbb{E}^\gamma[\log Z_t] = \log \rho + \Psi(\gamma).$$

□

To prove Theorem 1.2, we will use the strategy of Bezuidenhout and Grimmett, which consists in characterising survival by a local event. More precisely, for a fixed law  $\gamma$  allowing the branching random walk to survive, we will exhibit a local event  $A$ , that only depends on what happens in a finite box  $[0, T] \times [-L, L]^d$ , and a level  $p_0 < 1$  such that

$$(C_1) \quad \mathbb{P}^\gamma(A) > p_0$$

$$(C_2) \quad \text{For every law } \gamma' \text{ on } \mathcal{P}(\mathbb{N}), (\mathbb{P}^{\gamma'}(A) > p_0) \implies (\mathbb{P}^{\gamma'}(\text{survival}) > 0).$$

A coupling argument (whose proof is detailed in section 3.4) implies that

$$\mathbb{P}^{\gamma^\rho}(A) \geq \rho^{M(T+1)(2L+1)^d} \mathbb{P}^\gamma(A \cap B_M), \tag{1.4}$$

where  $B_M$  is the event: if the process starts with one individual at each site in  $[-L, L]^d$ , no site in  $[-L, L]^d$  contains more than  $M$  particles between time 0 and time  $T$ . Taking  $M$ , then  $\rho$  large enough, we get

$$(\mathbb{P}^\gamma(\text{survival}) > 0) \implies (\exists \rho \in (0, 1) \quad \mathbb{P}^{\gamma^\rho}(\text{survival}) > 0). \tag{1.5}$$

Consider now  $\gamma$  such that  $\mathbb{P}^\gamma(\text{survival}) > 0$ , take  $\rho$  given by (1.5) and assume  $\Psi(\gamma) = 0$ . Lemma 1.3 ensures  $\Psi(\gamma^\rho) < 0$  while  $\mathbb{P}^{\gamma^\rho}(\text{survival}) > 0$ , which is in contradiction with Proposition 1.1. Thus

$$(\mathbb{P}^\gamma(\text{survival}) > 0) \implies (\Psi(\gamma) > 0),$$

which proves Theorem 1.2.

We focus now on the Bezuidenhout–Grimmett construction for the BRWRE. First we need to give a more precise construction of the BRWRE. In particular, we will need a FKG inequality, which we prove in the next section.

## 2 Notations and FKG inequality

An initial configuration  $A = (A(x))_{x \in \mathbb{Z}^d}$  is an element of the set  $\mathbb{N}^{\mathbb{Z}^d}$  satisfying

$$|A| = \sum_{x \in \mathbb{Z}^d} A(x) < +\infty.$$

The number  $A(x)$  is the number of particles sitting on site  $x$  at time  $t = 0$ .

We encode the BRWRE starting from the initial configuration  $A$  by the random variables  $\eta_t^A = (\eta_t^A(x))_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$  which represent the number of particles living on site  $x$  at time  $t$ . Thus,

$$|\eta_t^A| = \sum_{x \in \mathbb{Z}^d} \eta_t^A(x)$$

stands for the total number of particles living at time  $t$ . To exploit the independence properties of the environment, we also need to consider the occupied sites at time  $t$ :

$$\underline{\eta}_t^A(x) = \mathbf{1}_{\{\eta_t^A(x) > 0\}} \quad \text{and} \quad \underline{\eta}_t^A = \{x \in \mathbb{Z}^d : \eta_t^A(x) > 0\}.$$

We denote by  $|\underline{\eta}_t^A|$  the number of occupied sites at time  $t$ . We define the lifetime of the branching random walk starting from  $A$

$$\tau^A = \min\{t \in \mathbb{N} : \eta_t^A = 0_{\mathbb{Z}^d}\} = \min\{t \in \mathbb{N} : \underline{\eta}_t^A = \emptyset\}.$$

We say that there is survival starting from  $A$  when  $\tau^A = +\infty$ .

In this article, the genealogy of the BRWRE is not central, and we choose a description in terms of particles systems. In the proof of the next lemma on FKG inequalities, we give a full construction of the BRWRE in the spirit of probabilistic cellular automata.

**Lemma 2.1.** *For any bounded non decreasing functions  $f, g : \mathbb{N}^{\mathbb{N} \times \mathbb{Z}^d} \rightarrow \mathbb{R}$ ,*

$$\begin{aligned} \mathbb{E}^\gamma [f(\eta_t^A) g(\eta_t^A)] &\geq \mathbb{E}^\gamma [f(\eta_t^A)] \mathbb{E}^\gamma [g(\eta_t^A)], \\ \mathbb{E}^\gamma [f(\underline{\eta}_t^A) g(\underline{\eta}_t^A)] &\geq \mathbb{E}^\gamma [f(\underline{\eta}_t^A)] \mathbb{E}^\gamma [g(\underline{\eta}_t^A)]. \end{aligned}$$

*The same inequalities also hold for any bounded non increasing functions  $f, g : \mathbb{N}^{\mathbb{N} \times \mathbb{Z}^d} \rightarrow \mathbb{R}$ .*

*Proof.* To build the BRWRE, we need a probability space on which live the following random variables:

- The  $(q_{t,x})_{(t,x) \in \mathbb{N} \times \mathbb{Z}^d}$  are i.i.d, take their values in  $\mathcal{P}(\mathbb{N})$ , and have  $\gamma$  as common law.
- The  $(U_{t,x,k})_{(t,x,k) \in \mathbb{N} \times \mathbb{Z}^d \times \mathbb{N}^*}$  are i.i.d and follow the uniform law on  $[0, 1]$ . With the environment  $\mathbf{q}$ , the random variable  $U_{t,x,k}$  is used to generate the progeny of the  $k$ -th particle sitting on site  $x$  at time  $t$ .

- The  $(D_{t,x,k})_{(t,x,k) \in \mathbb{N} \times \mathbb{Z}^d \times \mathbb{N}^*}$  are i.i.d and follow the uniform law on the set  $V = \{v \in \mathbb{Z}^d : \|v\|_1 = 1\}$ . The random variable  $D_{t,x,k}$  gives the displacement of the children of the  $k$ -th particle sitting on site  $x$  at time  $t$ .
- All these variables are independent.

So we can choose  $\Omega = \mathcal{P}(\mathbb{N})^{\mathbb{N} \times \mathbb{Z}^d} \times [0, 1]^{\mathbb{N} \times \mathbb{Z}^d \times \mathbb{N}^*} \times V^{\mathbb{N} \times \mathbb{Z}^d \times \mathbb{N}^*}$  and

$$\mathbb{P}^\gamma = \gamma^{\otimes \mathbb{N} \times \mathbb{Z}^d} \otimes \mathcal{U}([0, 1])^{\otimes \mathbb{N} \times \mathbb{Z}^d \times \mathbb{N}^*} \otimes \mathcal{U}(V)^{\otimes \mathbb{N} \times \mathbb{Z}^d \times \mathbb{N}^*}.$$

For  $t \geq 1$ , we denote by  $\theta_t$  the time translation operator acting on each coordinate of  $\Omega$ .

We define then the number  $E_{t,x}(v, p)$  of children born on site  $x$  at time  $t$  and moving to  $x + v$  when  $p$  particles live on  $x$  at time  $t$ :

$$E_{t,x}(v, p) = \sum_{k=1}^p \mathbb{1}_{\{D_{t,x,k}=v\}} \sum_{i=1}^{+\infty} \mathbb{1}_{\{\sum_{j=0}^i q_{t,x}(j) \leq U_{t,x,k}\}}.$$

An initial configuration  $A \in \mathbb{N}^{\mathbb{Z}^d}$  being given, we set  $\eta_0 = A$ , and

$$\eta_{t+1}^A(x) = \sum_{v \in V} E_{t,x-v}(v, \eta_t^A(x - e)).$$

We note  $\mathcal{F}_t = \sigma((q_{s,x}, U_{s,x,k}, D_{s,x,k})_{0 \leq s \leq t, x \in \mathbb{Z}^d, k \in \mathbb{N}^*})$ . Then,  $\eta_{t+1}$  is  $\mathcal{F}_t$ -measurable.

Note that

- for every  $\omega \in \Omega$  and  $v \in V$ ,  $p \mapsto E_{t,x}(v, p)(\omega)$  is non-decreasing.
- The coordinates of the random field  $(E_{t,x})_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$  are i.i.d. under  $\mathbb{P}^\gamma$ .

By induction on  $t \in \mathbb{N}$ , we thus see that  $A \mapsto \eta_t^A(x)$  is non-decreasing; also for fixed  $A$ , the  $(\eta_t^A(x))_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$  are non-decreasing with respect to the vectors  $(E_{t,x})_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$ . As these vectors are independent, the FKG inequality holds for the  $(\eta_t^A(x))_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$ . As the  $(\eta_t^A(x))_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$  are non-decreasing functions of the  $(\eta_t^A(x))_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$ , the FKG inequality also holds for the  $(\eta_t^A(x))_{t \in \mathbb{N}, x \in \mathbb{Z}^d}$ . Note that we use here the independent FKG inequality, also known as the Harris inequality (see Harris [7] or Grimmett [6]).  $\square$

### Truncating

To localize the events we consider, we also need truncated versions of the BRWRE: for  $B \subset \mathbb{Z}^d$ , and  $A \in \mathbb{N}^B$  such that  $|A| < +\infty$ , we set  ${}_B\eta_0^A = A$  and

$${}_B\eta_{t+1}^A(x) = \mathbb{1}_B(x) \sum_{v \in V} E_{t,x-v}(v, {}_B\eta_t^A(x - v)).$$

In words, we keep in  ${}_B\eta_t^A$  only particles living in  $B$ : other particles are simply discarded. Note that  ${}_B\eta_t^A(x)$  is obviously non-decreasing in  $A$  and non-decreasing in  $B$ . A straight adaptation of the previous proof ensures that the FKG inequalities of Lemma 2.1 still hold for the truncated BRWRE. Note that the process  $({}_B\eta_t^A)_{t \geq 0}$  is  $\sigma((q_{t,x}, U_{t,x,k}, D_{t,x,k})_{t \in \mathbb{N}, x \in B, k \in \mathbb{N}^*})$ -measurable.

We mainly work with restrictions on subsets  $B = \{-L, \dots, L\}^d$ . In that case, we simply note  ${}_L\eta_t^A$  instead of  $\{{}_B\eta_t^A\}_{B = \{-L, \dots, L\}^d}$ .

## 3 Construction of the block event

### Outline of proof

Recall that we are looking for a local event  $A$  which satisfies  $(C_1)$  and  $(C_2)$ . The idea is to find an event that expresses the fact that if the branching random walk occupies a sufficiently large area at a given place, it will presumably extend itself a bit further.

Set, for every integer  $n \geq 1$ ,

$$A_n = \left\{ x \in \mathbb{Z}^d : \|x\|_1 \leq n \text{ and } \sum_{i=1}^n x_i \equiv 0 \pmod{2} \right\}.$$

Note that when  $n$  is even,  $A_n$  is the maximal set that a branching random walk can reach at time  $n$ .

The next proposition ensures that starting from  $A_n$  with  $n$  large enough, the BRWRE restrained to a large box  $\{-2L - 2n, \dots, 2L + 2n\}^d \times \{0, \dots, 2T\}$  occupies with high probability a translated copy of  $A_n$ :

**Proposition 3.1.** *For every  $\varepsilon > 0$ , there exist positive integers  $n, L, T$ , with  $n \leq L$  such that*

$$\mathbb{P}^\gamma \left( \begin{array}{c} \exists x \in \{L + n, \dots, 2L + n\} \times \{0, \dots, 2L\}^{d-1}, t \in \{T, \dots, 2T\} \\ 2L+2n\underline{\eta}_t^{A_n} \supset x + A_n \end{array} \right) \geq 1 - \varepsilon.$$

Proposition 3.1 provides an event  $A$  that satisfies  $(C_1)$ . The fact that  $A$  also satisfies  $(C_2)$  follows from a quite standard construction of an embedded supercritical percolation of blocks. This construction does not rely on the specificities of the model. We can find in the literature many examples of similar block events. Most of these papers adapt the initial construction of the Bezuidenhout–Grimmett article [1]: *The critical contact process dies out*. Their proof is also exposed in the reference book by Liggett [9].

A complete description of these procedures can be found in Steif–Warfheimer [10] in the case of a contact process where the death rate depends on a dynamical environment or in Garet–Marchand [5] for a class of dependent oriented percolation.

### 3.1 Some properties of the surviving branching random walk

We fix  $\gamma$  such that the probability of survival  $\mathbb{P}^\gamma(\tau^{\{0\}} = +\infty)$  is positive. At first, we prove that survival implies the explosion of the number of particles.

**Lemma 3.2.** *For every finite initial configuration  $A$ ,*

$$\mathbb{P}^\gamma(\tau^A = +\infty, \lim_{t \rightarrow +\infty} |\eta_t^A| = +\infty) = \mathbb{P}^\gamma(\tau^A = +\infty).$$

*Proof.* Let  $A$  be a fixed finite initial configuration, and  $N$  be a fixed positive integer. By assumption (1.3), there exist  $\varepsilon_0, \alpha_0 > 0$  such that

$$\mathbb{P}^\gamma(q_{0,0}(0) > \varepsilon_0) > \alpha_0.$$

By blocking the progenies of all living particles at time  $s$ , we see that,

$$\forall s > 1 \quad \mathbb{P}^\gamma(\tau^A < +\infty \mid \mathcal{F}_{s-1}) \geq \varepsilon_0^{|\eta_s^A|} \alpha_0^{|\eta_s^A|} \geq (\varepsilon_0 \alpha_0)^{|\eta_s^A|}.$$

By the martingale convergence theorem,  $\lim_{s \rightarrow +\infty} \mathbb{P}^\gamma(\tau^A < +\infty \mid \mathcal{F}_{s-1}) = \mathbb{1}_{\{\tau^A < +\infty\}}$ . So on the event  $\{\tau^A = +\infty\}$ ,  $\lim_{s \rightarrow +\infty} |\eta_s^A| = +\infty$ .  $\square$

To exploit the independence properties of the environment, we need particles to sit on distinct sites; the fact that the number of particles at time  $t$  explodes when the BRWRE survives does not directly ensure that the number of occupied sites also explodes. It will be a by-product of our construction, but we are not able to prove it at this stage. So we need to work separately under the two complementary assumptions:

$$\begin{aligned} (H) \quad & \mathbb{P}^\gamma(q_{0,0}(0) = 1) > 0, \\ (H^c) \quad & \mathbb{P}^\gamma(q_{0,0}(\mathbb{N}^*) > 0) = 1. \end{aligned}$$

Under Assumption (H), we prove that survival implies the explosion of the number of occupied sites:

**Lemma 3.3.** *Assume (H) is fulfilled. For every finite initial configuration A,*

$$\mathbb{P}^\gamma(\tau^A = +\infty, \lim_{t \rightarrow +\infty} |\eta_t^A| = +\infty) = \mathbb{P}^\gamma(\tau^A = +\infty).$$

*Proof.* Let A be a fixed finite initial configuration, and N be a fixed positive integer. Under (H)

$$\varepsilon_1 = \mathbb{P}^\gamma(q_{0,0}(0) = 1) > 0.$$

By blocking the progenies of all occupied sites at time s, we see that,

$$\forall s \geq 1 \quad \mathbb{P}^\gamma(\tau^A < +\infty \mid \mathcal{F}_{s-1}) \geq \varepsilon_1^{|\eta_s^A|}.$$

By the martingale convergence theorem,  $\lim_{s \rightarrow +\infty} \mathbb{P}^\gamma(\tau^A < +\infty \mid \mathcal{F}_{s-1}) = \mathbb{1}_{\{\tau^A < +\infty\}}$ . So on the event  $\{\tau^A = +\infty\}$ ,  $\lim_{s \rightarrow +\infty} |\eta_s^A| = +\infty$ . □

Besides, under Assumption (H<sup>c</sup>), we prove that if we start with many particles sitting on the same site, there is a large probability that after some time many sites are occupied. We denote by (N) the initial configuration where N particles sit on 0. Formally, we have (N) = (Nδ<sub>0</sub>(x))<sub>x∈Z<sup>d</sup></sub>.

**Lemma 3.4.** *Assume (H<sup>c</sup>) is fulfilled. Let n be a fixed even integer.*

*Set B<sub>n</sub> = {0, ..., 2n} × {−n, ..., n}<sup>d−1</sup>. Then  $\lim_{N \rightarrow +\infty} \mathbb{P}^\gamma(ne_1 + A_n \subset B_n \eta_{2n}^{(N)}) = 1$ .*

*Proof.* Fix n ≥ 1 and α > 0. Under (H<sup>c</sup>),

$$\forall \varepsilon > 0 \quad \exists \eta > 0 \quad \mathbb{P}^\gamma(q_{0,0}(\mathbb{N} \setminus \{0\}) \geq \eta) \geq 1 - \varepsilon/2. \tag{3.1}$$

We say that the environment q is fertile in the box B<sub>n</sub> × {0, ..., 2n} if

$$\forall x \in B_n \quad \forall t \in \{0, \dots, 2n\} \quad q_{x,t}(\mathbb{N} \setminus \{0\}) \geq \eta;$$

we denote this event by F. Thus, with ε and η satisfying (3.1),

$$\mathbb{P}^\gamma(F) \geq (1 - \varepsilon)^{(4n+1)^d(2n+1)}.$$

We choose ε > 0 such that  $\mathbb{P}^\gamma(F) \geq 1 - \alpha/2$ , and take the corresponding η given by (3.1). Now, as  $\mathbb{P}^\gamma(ne_1 + A_n \subset \eta_{2n}^{(N)}) \geq \mathbb{P}^\gamma(ne_1 + A_n \subset \eta_{2n}^{(N)} \mid F)(1 - \alpha/2)$ , we restrict ourselves to environments in F.

For an environment q in F, each coordinate q<sub>x,t</sub>, x ∈ B<sub>n</sub>, t ∈ {0, ..., 2n} stochastically dominates the fixed law γ<sub>0</sub> = ηδ<sub>1</sub> + (1 - η)δ<sub>0</sub>. Thus by coupling, it is sufficient to prove that

$$\lim_{N \rightarrow +\infty} \mathbb{P}^{\gamma_0}(ne_1 + A_n \subset B_n \eta_{2n}^{(N)}) = 1. \tag{3.2}$$

Let now x ∈ ne<sub>1</sub> + A<sub>n</sub> be fixed. As ||x||<sub>1</sub> ≤ 2n,

$$\mathbb{P}^{\gamma_0}(x \in B_n \eta_{2n}^{(1)}) \geq \left(\frac{\eta}{2d}\right)^{2n} > 0.$$

As the environment is non-random, the behavior of distinct particles are independent under  $\mathbb{P}^{\gamma_0}$ . Thus,

$$\lim_{N \rightarrow +\infty} \mathbb{P}^{\gamma_0}(x \in B_n \eta_{2n}^{(N)}) \geq \lim_{N \rightarrow +\infty} 1 - \left(1 - \left(\frac{\eta}{2d}\right)^{2n}\right)^N = 1,$$

which proves (3.2); this ends the proof. □

A last lemma ensures that starting from  $A_n$  with  $n$  large, the survival probability is large:

**Lemma 3.5.**  $\lim_{n \rightarrow +\infty} \mathbb{P}^\gamma(\tau^{A_n} = +\infty) = 1.$

*Proof.* Denote

$$A_\infty = \{x \in \mathbb{Z}^d : \sum_{i=1}^n x_i \in 2\mathbb{Z}\}.$$

We know that  $\eta_t^{\{x\}} \subset \eta_t^{A_n}$  holds for each  $n \geq 1, t \geq 0$  and  $x \in A_n$ . It follows that  $\{\tau^{A_n} = +\infty\} \supset \bigcup_{x \in A_n} \{\tau^{\{x\}} = +\infty\}$ . So

$$\begin{aligned} \lim_{n \rightarrow +\infty} \mathbb{P}^\gamma(\tau^{A_n} = +\infty) &\geq \lim_{n \rightarrow +\infty} \mathbb{P}^\gamma\left(\bigcup_{x \in A_n} \{\tau^{\{x\}} = +\infty\}\right) \\ &= \mathbb{P}^\gamma\left(\bigcup_{x \in A_\infty} \{\tau^{\{x\}} = +\infty\}\right). \end{aligned}$$

The event  $\bigcup_{x \in A_\infty} \{\tau^{\{x\}} = +\infty\}$  is invariant under the spatial translation  $x \mapsto x + 2e_1$ , so by ergodicity, it is either null or full.

Since  $\mathbb{P}^\gamma\left(\bigcup_{x \in A_\infty} \{\tau^{\{x\}} = +\infty\}\right) \geq \mathbb{P}^\gamma(\tau^{\{0\}} = +\infty) > 0$ , it is equal to 1. □

**Lemma 3.6.** Recall that  $B_n = \{0, \dots, 2n\} \times \{-n, \dots, n\}^{d-1}$ . For every  $n \geq 1$ , there exists  $h > 0$  such that

$$\mathbb{P}^\gamma(A_n \subset \underline{n}\eta_h^{\{0\}}) > 0 \text{ and } \mathbb{P}^\gamma(ne_1 + A_n \subset B_n \underline{n}\eta_h^{\{0\}}) > 0.$$

*Proof.* By (1.3), there exists  $\eta > 0$  and  $\varepsilon > 0$  such that

$$\mathbb{P}^\gamma(q_{0,0}(0) + q_{0,0}(1) \leq 1 - \eta) \geq \varepsilon.$$

Let  $k$  be even with  $2^k \geq |A_n|$  and let  $h = k + 2n$ . Note

$$G = \{\forall(x, t) \in B_n \times \{0, \dots, k + 2n\} \quad q_{0,0}(0) + q_{0,0}(1) \leq 1 - \eta\}.$$

$\mathbb{P}^\gamma(G) \geq \varepsilon^{|B_n \times \{0, \dots, h\}|} > 0$ . On the event  $G$ , the environment allows each individual to have more than one daughter with probability  $\eta$ .

For an environment  $q$  in  $F$ , each coordinate  $q_{x,t}, x \in B_n, t \in \{0, \dots, h\}$  stochastically dominates the fixed law  $\gamma_1 = \eta\delta_2 + (1 - \eta)\delta_0$ , so for each event  $E$ , we have

$$\mathbb{P}^\gamma(G) \geq \varepsilon^{|B_n \times \{0, \dots, h\}|} \mathbb{P}^{\gamma_1}(G).$$

Thus, it is sufficient to prove that  $\mathbb{P}^{\gamma_1}(A_n \subset \underline{n}\eta_h^{\{0\}}) > 0$  and  $\mathbb{P}^{\gamma_1}(ne_1 + A_n \subset B_n \underline{n}\eta_h^{\{0\}}) > 0$ .

We only prove the second inequality; the first one is similar.

Under  $\mathbb{P}^{\gamma_1}$ , the probability that there is at least  $|A_n|$  particles at the origin at time  $k$  whose ancestors never left  $\{0; e_1\}$  is at least  $(\frac{\eta}{2d})^{2^{k+1}-1}$ . Then, there is a probability at least  $(\frac{\eta}{2d})^{2n|A_n|}$  that their progeny fills  $ne_1 + A_n$  in  $2n$  steps, with trajectories remaining inside  $\{-2n, \dots, 2n\}^d$ . Finally,

$$\mathbb{P}^{\gamma_1}(ne_1 + A_n \subset B_n \underline{n}\eta_h^{\{0\}}) \geq (\frac{\eta}{2d})^{2^{k+1}-1} (\frac{\eta}{2d})^{2n|A_n|} > 0.$$

The other proof is similar. □



**3.2 From survival to local events**

**Lemma 3.7.** *For every finite  $A \subset \mathbb{Z}^d$ , for every positive integer  $N$ ,*

$$\lim_{t \rightarrow +\infty} \lim_{L \rightarrow +\infty} \mathbb{P}(|_L \eta_t^A| \geq N) = \mathbb{P}(\tau^A = +\infty).$$

*Under the extra assumption (H),*

$$\lim_{t \rightarrow +\infty} \lim_{L \rightarrow +\infty} \mathbb{P}(|_L \underline{\eta}_t^A| \geq N) = \mathbb{P}(\tau^A = +\infty).$$

*Proof.* Let  $A$  be a fixed finite subset of  $\mathbb{Z}^d$ , and  $N$  be a fixed positive integer.

$$\forall t \in \mathbb{N} \quad \eta_t^A = \bigcup_{L \in \mathbb{N}} |_L \eta_t^A.$$

Indeed, the inclusion  $\supset$  follows from positivity, and if  $L \geq \|A\|_\infty + 2t + 1$ , then for every  $s \leq t$ ,  $\eta_s^A = |_L \eta_s^A$ . Thus

$$\{|\eta_t^A| \geq N\} = \bigcup_{L \in \mathbb{N}} \{|_L \eta_t^A| \geq N\}$$

and thus  $\lim_{L \rightarrow +\infty} \mathbb{P}(|_L \eta_t^A| \geq N) = \mathbb{P}(|\eta_t^A| \geq N)$ . But Lemma 3.2 and the dominated convergence theorem ensure that

$$\lim_{t \rightarrow +\infty} \mathbb{P}^\gamma(\tau^A = +\infty, |\eta_t^A| \geq N) = \mathbb{P}(\tau^A = +\infty).$$

In the same way,  $\lim_{L \rightarrow +\infty} \mathbb{P}(|_L \underline{\eta}_t^A| \geq N) = \mathbb{P}(|\underline{\eta}_t^A| \geq N)$ . Under Assumption (H), we use Lemma 3.3 to conclude. □

Then, using the FKG inequality with a classical square root trick, we can ensure that the truncated process at time  $t$  contains many points in a prescribed orthant of  $\mathbb{Z}^d$ :

**Lemma 3.8.** *For every positive integers  $n, N, t$ , for every integer  $L > n$ ,*

$$\begin{aligned} \mathbb{P}^\gamma \left( |{}_L \eta_t^{A_n} \cap \{0, \dots, L\}^d| \leq N \right)^{2^d} &\leq \mathbb{P}^\gamma \left( |{}_L \eta_t^{A_n}| \leq N 2^d \right), \\ \mathbb{P}^\gamma \left( |{}_L \underline{\eta}_t^{A_n} \cap \{0, \dots, L\}^d| \leq N \right)^{2^d} &\leq \mathbb{P}^\gamma \left( |{}_L \underline{\eta}_t^{A_n}| \leq N 2^d \right). \end{aligned}$$

*Proof.* By the symmetries of the model and of  $A_n$ , the law of the intersection of the process with a given orthant does not depend on the orthant. Applying the FKG inequality to the non increasing indicator functions of the corresponding events, we get

$$\mathbb{P}^\gamma \left( |{}_L \eta_t^{A_n} \cap \{0, \dots, L\}^d| \leq N \right)^{2^d} \leq \mathbb{P}^\gamma \left( \bigcap_{\varepsilon \in \{+1, -1\}^d} \left\{ \left| {}_L \eta_t^{A_n} \cap \prod_{i=1}^n \varepsilon_i \{0, \dots, L\} \right| \leq N \right\} \right),$$

which is smaller than  $\mathbb{P}^\gamma \left( |{}_L \eta_t^{A_n}| \leq N 2^d \right)$ . The other proof is similar. □

For every positive integers  $n, L, T$  with  $L > n$ , we define  $N^{A_n}(L, T)$  as the number of particles sitting on sites  $(x, t)$  such that  $t \in \{0, \dots, T\}$ ,  $\|x\|_\infty = L$ ,  $x \in |_L \eta_t^{A_n}$ : this is the number of particles sitting on a given lateral face of the large box. We also define  $\underline{N}^{A_n}(L, T)$  as the number of occupied sites  $(x, t)$  such that  $t \in \{0, \dots, T\}$ ,  $\|x\|_\infty = L$ ,  $x \in |_L \underline{\eta}_t^{A_n}$ : this is the number of occupied sites on a given lateral face of the large box. The next point is to ensure that when the BRWRE survives, it must occupy many points and on the top face and on the lateral faces of a large box:

**Lemma 3.9.** *For any positive integers  $M, N, n$ , for any increasing sequences of integers  $(T_j)_{j \geq 1}$  and  $(L_j)_{j \geq 1}$ ,*

$$\lim_{j \rightarrow +\infty} \mathbb{P}^\gamma \left( N^{A_n}(L_j, T_j) \leq M \right) \mathbb{P}^\gamma \left( |_{L_j} \eta_{T_j}^{A_n}| \leq N \right) \leq \mathbb{P}^\gamma \left( \tau^{A_n} < +\infty \right).$$

*Under the extra assumption (H),*

$$\lim_{j \rightarrow +\infty} \mathbb{P}^\gamma \left( \underline{N}^{A_n}(L_j, T_j) \leq M \right) \mathbb{P}^\gamma \left( |_{L_j} \underline{\eta}_{T_j}^{A_n}| \leq N \right) \leq \mathbb{P}^\gamma \left( \tau^{A_n} < +\infty \right).$$

*Proof.* For integers  $L, T$ , let  $\mathcal{F}_{L,T} = \sigma((q_{s,x}, U_{s,x,k}, D_{s,x,k})_{0 \leq s \leq T, x \in \{-L, \dots, L\}^d, k \in \mathbb{N}^*})$ . Set  $k = M + N$ . Let  $(T_j)_j$  and  $(L_j)_j$  be two increasing sequences of integers.

$$H_j = \{N^{A_n}(L_j, T_j) + |_{L_j} \eta_{T_j}^{A_n}| \leq k\} \quad \text{and} \quad G = \{\tau^{A_n} < +\infty\}.$$

We proceed as in Lemma 3.2. Since  $H_j \in \mathcal{F}_{L_j, T_j-1}$ , we have

$$\mathbb{P}^\gamma(G | \mathcal{F}_{L_j, T_j}) \geq \mathbb{1}_{H_j} (\varepsilon_0 \alpha_0)^k.$$

By the martingale convergence theorem,  $\mathbb{P}^\gamma(G | \mathcal{F}_{L_j, T_j})$  almost surely converges to  $\mathbb{1}_G$ , which implies that

$$\overline{\lim}_{j \rightarrow +\infty} H_j \subset G, \quad \text{and thus} \quad \overline{\lim}_{j \rightarrow +\infty} \mathbb{P}^\gamma(H_j) \leq \mathbb{P} \left( \overline{\lim}_{j \rightarrow +\infty} H_j \right) \leq \mathbb{P}^\gamma(G).$$

Using the FKG inequality once again, note that

$$\begin{aligned} \mathbb{P}^\gamma(H_j) &= \mathbb{P}^\gamma(N^{A_n}(L_j, T_j) + |_{L_j} \eta_{T_j}^{A_n}| \leq M + N) \\ &\geq \mathbb{P}^\gamma(N^{A_n}(L_j, T_j) \leq M) \mathbb{P}^\gamma(|_{L_j} \eta_{T_j}^{A_n}| \leq N), \end{aligned}$$

which ends the proof. The proof of the result under (H) is obtained by adapting the proof as Lemma 3.3 is adapted from Lemma 3.2.  $\square$

Exactly as in Lemma 3.8, the FKG inequality and the symmetries of the process allow to control the number of colonized points in a prescribed orthant of a lateral face of the box  $\{-L, \dots, L\}^d \times \{0, \dots, T\}$ .

For every positive integers  $L, T$  and every finite  $A \subset \mathbb{Z}^d$ , we define  $N_+^A(L, T)$  as the number of particles sitting on sites  $(x, t)$  such that  $t \in \{0, \dots, T\}$ ,  $x_1 = L$ ,  $x_i \geq 0$  for  $2 \leq i \leq d$ ,  $x \in_L \underline{\eta}_t^A$ , and  $\underline{N}_+^A(L, T)$  as the number of occupied sites  $(x, t)$  such that  $t \in \{0, \dots, T\}$ ,  $x_1 = L$ ,  $x_i \geq 0$  for  $2 \leq i \leq d$ ,  $x \in_L \underline{\eta}_t^A$ . Then

**Lemma 3.10.** *For every positive integers  $n, N, K, t$ , for every integer  $L \geq n$ ,*

$$\mathbb{P}^\gamma \left( N_+^{A_n}(L, T) \leq M \right)^{d2^d} \leq \mathbb{P}^\gamma \left( N^{A_n}(L, T) \leq Md2^d \right).$$

$$\mathbb{P}^\gamma \left( \underline{N}_+^{A_n}(L, T) \leq M \right)^{d2^d} \leq \mathbb{P}^\gamma \left( \underline{N}^{A_n}(L, T) \leq Md2^d \right).$$

*Proof.* This is the same square root trick as in the proof of Lemma 3.8.  $\square$

### 3.3 Proof of Proposition 3.1

With the previous lemmas in hand, we can now prove Proposition 3.1.

**3.3.1 First case: assume that (H) holds.**

In that case, we can work with the number of occupied sites.

Let  $\varepsilon > 0$  be fixed and  $0 < \delta < 1$  to be chosen later.

With Lemma 3.5, we first choose a positive even integer  $n$  such that

$$\mathbb{P}^\gamma (\tau^{A_n} = +\infty) > 1 - 2\delta^2. \tag{3.3}$$

We take then  $h > 0$  given by Lemma 3.6. Choose an integer  $N'$  such that

$$\left(1 - \mathbb{P}^\gamma \left( {}_n \underline{\eta}_h^0 \supset A_n \right)\right)^{N'} \leq \delta, \tag{3.4}$$

and then  $N$  such that every finite subset  $A$  of  $\mathbb{Z}^d$  with cardinal  $N$  contains a subset  $A'$  of  $N'$  points such that

$$\forall x, y \in A' \quad (x \neq y) \implies \|x - y\|_\infty \geq 2n + 1.$$

Since background random variables in disjoint areas are independent, as a consequence of (3.4), we get that for every subset  $A$  of  $\mathbb{Z}^d$ .

$$|A| \geq N \implies \mathbb{P}^\gamma (\exists x \in A \quad {}_{n+\|A\|_\infty} \underline{\eta}_h^x \supset x + A_n) \geq 1 - \delta \tag{3.5}$$

Similarly, there exists  $M > 0$  such that

$$|A| \geq M \implies \mathbb{P}^\gamma (\exists x \in A \quad {}_{2n+\|A\|_\infty} \underline{\eta}_h^x \supset x + ne_1 + A_n) \geq 1 - \delta \tag{3.6}$$

Since  $1 - 2\delta < 1 - 2\delta^2 < \mathbb{P}^\gamma (\tau^{A_n} = +\infty)$ , there exist by Lemma 3.7  $T_0 \in \mathbb{N}$  and a map  $L' : [T_0, +\infty) \rightarrow \mathbb{N}$  such that

$$\forall T \geq T_0 \quad \forall L \geq L'(T) \quad \mathbb{P}^\gamma \left( |{}_L \underline{\eta}_T^{A_n}| > 2^d N \right) \geq 1 - \delta.$$

Define  $L_0 = L'(T_0)$ , then for  $i \geq 0$

$$T_{i+1} = \inf \left\{ T > T_i; \mathbb{P}^\gamma \left( |{}_{L_i} \underline{\eta}_T^{A_n}| > 2^d N \right) < 1 - 2\delta \right\}.$$

Note that  $T_{i+1} < +\infty$  because with assumption (H) we have

$$\lim_{t \rightarrow +\infty} \mathbb{P}^\gamma \left( |{}_{L_i} \underline{\eta}_t^{A_n}| > 2^d N \right) = 0.$$

Then simply take  $L_{i+1} = \max(L'(T_{i+1}), L_i + 1)$ . Then, we have two increasing sequences  $(T_i)_{i \geq 1}$  and  $(L_i)_{i \geq 1}$  such that

$$\begin{aligned} \forall i \geq 1 \quad \mathbb{P}^\gamma \left( |{}_{L_i} \underline{\eta}_{T_i}^{A_n}| > 2^d N \right) &\geq 1 - 2\delta \\ \forall i \geq 1 \quad \mathbb{P}^\gamma \left( |{}_{L_i} \underline{\eta}_{T_{i+1}}^{A_n}| > 2^d N \right) &< 1 - 2\delta. \end{aligned}$$

With Lemma 3.9 and (3.3), there exists  $K$  such that

$$\mathbb{P}^\gamma \left( \underline{N}^{A_n}(L_K, T_K + 1) \leq d2^d M \right) \mathbb{P}^\gamma \left( |{}_{L_K} \underline{\eta}_{T_K+1}^{A_n}| \leq 2^d N \right) \leq 2\delta^2.$$

From now on, set  $L = L_K$  and  $T = T_K + 1$ . We get

$$\begin{aligned} \mathbb{P}^\gamma \left( |{}_L \underline{\eta}_{T-1}^{A_n}| > 2^d N \right) &\geq 1 - 2\delta, \\ \mathbb{P}^\gamma \left( \underline{N}^{A_n}(L, T) > d2^d M \right) &\geq 1 - \frac{2\delta^2}{1 - \mathbb{P}^\gamma \left( |{}_L \underline{\eta}_T^{A_n}| > 2^d N \right)} \geq 1 - \delta. \end{aligned}$$

With lemmas 3.8 and 3.10, we obtain

$$\mathbb{P}^\gamma \left( |\underline{L}\eta_{T-1}^{A_n} \cap \{0, \dots, L\}^d | > N \right) \geq 1 - (2\delta)^{2^{-d}}, \tag{3.7}$$

$$\mathbb{P}^\gamma \left( \underline{N}_+^{A_n}(L, T) > M \right) \geq 1 - \delta^{2^{-d}/d}. \tag{3.8}$$

Putting (3.5) and (3.7) together, we get

$$\mathbb{P}^\gamma \left( \exists x \in \{0, \dots, L\}^d \quad \underline{L}_{L+2n}\eta_{T-1+h}^{A_n} \supset x + A_n \right) \geq (1 - (2\delta)^{2^{-d}})(1 - \delta).$$

Now, we can use (3.8) and (3.6) starting from the translated  $x + A_n$  occupied at time  $T - 1 + h$ : it leads to

$$\mathbb{P}^\gamma \left( \begin{array}{l} \exists x \in \{L + n, \dots, 2L + n\} \times \{0, \dots, 2L\}^{d-1}, \\ \exists t \in \{T - 1 + 2h, \dots, 2T - 1 + 2h\} \\ \underline{2L+2n}\eta_t^{A_n} \supset x + A_n \end{array} \right) \geq (1 - (2\delta)^{2^{-d}})(1 - \delta)(1 - \delta^{2^{-d}/d})(1 - \delta).$$

By choosing  $\delta > 0$  small enough, the right-hand side can be made larger than  $1 - \varepsilon$ , and renaming  $T + h$  by  $T$  gives the result.

### 3.3.2 Second case: assume that $(H^c)$ holds.

We can not work with the number of occupied sites anymore, so we must work with the number of particles.

Let  $\varepsilon > 0$ . Let  $\delta \in (0, 1)$  be such that  $\delta \leq \varepsilon/4$  and

$$(1 - (2\delta)^{2^{-d}})(1 - \delta)(1 - \delta^{2^{-d}/d})(1 - \delta) \geq 1 - \varepsilon.$$

We must again split the proof into two complementary subcases.

$$(A_\delta) : \quad \forall L > 0 \quad \overline{\lim}_{T \rightarrow +\infty} \mathbb{P}^\gamma(|\underline{L}\eta_T^{A_n}| > 0) < 1 - 2\delta$$

$$(A_\delta^c) : \quad \exists L > 0 \quad \overline{\lim}_{T \rightarrow +\infty} \mathbb{P}^\gamma(|\underline{L}\eta_T^{A_n}| > 0) \geq 1 - 2\delta.$$

Suppose first that  $(H^c)$  and  $(A_\delta)$  hold.

Choose  $n, N, M$  exactly as in (3.3), (3.5) and (3.6). Under  $(H^c)$ , by Lemma 3.4, we can choose  $J$  large enough to have

$$\left( 1 - \mathbb{P}^\gamma \left( ne_1 + A_n \subset B_n \eta_{2n}^{(J)} \right) \right) \leq \delta. \tag{3.9}$$

Set  $R = \max\{N, M, J\}^2$ .

Using  $1 - 2\delta < 1 - 2\delta^2 < \mathbb{P}^\gamma(\tau^{A_n} = +\infty)$ ,  $(A_\delta)$  and Lemma 3.7 together, the same reasoning as in case  $(H)$  gives the existence of two increasing sequences  $(T_i)_{i \geq 1}$  and  $(L_i)_{i \geq 1}$  such that

$$\begin{aligned} \forall i \geq 1 \quad \mathbb{P}^\gamma \left( |\underline{L}_i \eta_{T_i}^{A_n}| > 2^d R \right) &\geq 1 - 2\delta \\ \forall i \geq 1 \quad \mathbb{P}^\gamma \left( |\underline{L}_i \eta_{T_{i+1}}^{A_n}| > 2^d R \right) &< 1 - 2\delta. \end{aligned}$$

By Lemma 3.9 and (3.3), there exists  $K$  such that

$$\mathbb{P}^\gamma \left( N^{A_n}(L_K, T_K + 1) \leq d2^d R \right) \mathbb{P}^\gamma \left( |\underline{L}_K \eta_{T_K+1}^{A_n}| \leq 2^d R \right) \leq 2\delta^2,$$

From now on, set  $L = L_K$  and  $T = T_K + 1$ . We get

$$\begin{aligned} \mathbb{P}^\gamma \left( |{}_L\eta_{T-1}^{A_n}| > 2^d R \right) &\geq 1 - 2\delta, \\ \mathbb{P}^\gamma \left( N^{A_n}(L, T) > d2^d R \right) &\geq 1 - \frac{2\delta^2}{1 - \mathbb{P}^\gamma \left( |{}_L\eta_{T-1}^{A_n}| > 2^d R \right)} \geq 1 - \delta. \end{aligned}$$

By lemmas 3.8 and 3.10, we obtain

$$\mathbb{P}^\gamma \left( |{}_L\eta_{T-1}^{A_n} \cap \{0, \dots, L\}^d| > R \right) \geq 1 - (2\delta)^{2^{-d}}, \tag{3.10}$$

$$\mathbb{P}^\gamma \left( N_+^{A_n}(L, T) > R \right) \geq 1 - \delta^{2^{-d}/d}. \tag{3.11}$$

But if  $|{}_L\eta_{T-1}^{A_n} \cap \{0, \dots, L\}^d| > R$ ,

- either  $|{}_L\eta_{T-1}^{A_n} \cap \{0, \dots, L\}^d| \geq \sqrt{R} \geq N$ ,
- or there is one  $x \in {}_L\eta_{T-1}^{A_n} \cap \{0, \dots, L\}^d$  such that  $\eta_{T-1}^{A_n}(x) \geq \sqrt{R} \geq J$ .

Using Equation (3.10), either the choice we made for  $N$  or the choice (3.9) we made for  $J$  leads to

$$\mathbb{P}^\gamma \left( \exists x \in \{0, \dots, L\}^d \quad {}_{L+2n}\eta_{T-1+h}^{A_n} \supset x + A_n \right) \geq (1 - (2\delta)^{2^{-d}})(1 - \delta).$$

Now, we can use (3.11) starting from the translated  $x + A_n$  occupied at time  $T - 1 + h$ . But if  $N_+^{A_n}(L, T) > R$ ,

- either the number of occupied sites on this face is larger than  $\sqrt{R} \geq M$ ,
- or one of the occupied sites contains more than  $\sqrt{R} \geq J$  particles.

Using Equation (3.11), either the choice we made for  $M$  or the choice (3.9) we made for  $J$  leads to

$$\begin{aligned} \mathbb{P}^\gamma \left( \begin{array}{l} \exists x \in \{L+n, \dots, 2L+n\} \times \{0, \dots, 2L\}^{d-1}, \\ \exists t \in \{T-1+2h, \dots, 2T-1+2h\} \\ {}_{2L+2n}\eta_t^{A_n} \supset x + A_n \end{array} \right) \\ \geq (1 - (2\delta)^{2^{-d}})(1 - \delta)(1 - \delta^{2^{-d}/d})(1 - \delta) \geq 1 - \varepsilon. \end{aligned}$$

Renaming  $T + h$  by  $T$  gives the result.

Suppose finally that  $(H^c)$  and  $(A_\delta^c)$  hold

In other words, there exists  $L$  such that

$$\lim_{T \rightarrow +\infty} \mathbb{P}^\gamma(|{}_L\eta_T^{A_n}| > 0) = \mathbb{P}^\gamma({}_L\tau^{A_n} = +\infty) \geq 1 - 2\delta. \tag{3.12}$$

We see that this is the most favorable case for the branching random walk, because it can survive even in a finite box. We will see it is also the easiest case to deal with.

We fix then a large  $L$  such that (3.12) is satisfied. There exists  $h > 0$  such that

$$\mathbb{P}^\gamma(\forall x \in \{-L, \dots, L\}^d, (L+n)e_1 + A_n \subset {}_{2L+2n}\eta_h^x) = r > 0.$$

For  $k \geq 0$ , set  $T_k = kh$  and

$$B_k = \{\forall x \in \{-L, \dots, L\}^d, (L+n)e_1 + A_n \subset {}_{2L+2n}\eta_h^x \circ \theta_{T_k}\}.$$

The  $B_k$ 's are clearly independent. Moreover

$$\{{}_L\tau^{A_n} = +\infty\} \cap B_k \subset \{(L+n)e_1 + A_n \subset {}_{2L+2n}\eta_{T_{k+1}}^{A_n}\}.$$

Thus, for any  $K > 1$ ,

$$\begin{aligned} & \mathbb{P}^\gamma(L\tau^{A_n} = +\infty, \forall t \in \{T_K, \dots, T_{2K}\} (L+n)e_1 + A_n \not\subset_{2L+2n}\underline{\eta}_t^{A_n}) \\ & \leq \mathbb{P}^\gamma(L\tau^{A_n} = +\infty, \bigcap_{k=K-1}^{2K-1} B_k^c) \\ & \leq \mathbb{P}^\gamma\left(\bigcap_{k=K-1}^{2K-1} B_k^c\right) = (1-r)^{K+1}. \end{aligned}$$

We set now  $2\delta = \varepsilon/2$ , and take  $K$  large enough to have  $(1-r)^{K+1} < \varepsilon/2$ , and

$$\begin{aligned} & \mathbb{P}^\gamma\left(\exists x \in \{L+n, \dots, 2L+n\} \times \{0, \dots, 2L\}^{d-1}, t \in \{T, \dots, 2T\} \right. \\ & \quad \left. \begin{array}{c} 2L+2n\underline{\eta}_t^{A_n} \supset x + A_n \end{array} \right) \\ & \geq \mathbb{P}^\gamma(L\tau^{A_n} = +\infty, \exists t \in \{T_K, \dots, T_{2K}\} (L+n)e_1 + A_n \subset_{2L+2n}\underline{\eta}_t^{A_n}) \\ & \geq \mathbb{P}^\gamma(L\tau^{A_n} = +\infty) - \mathbb{P}^\gamma(L\tau^{A_n} = +\infty, \forall t \in \{T_K, \dots, T_{2K}\} (L+n)e_1 + A_n \not\subset_{2L+2n}\underline{\eta}_t^{A_n}) \\ & \geq 1 - 2\delta - \varepsilon/2 = 1 - \varepsilon. \end{aligned}$$

One can see that we had to split the study into three cases and to make (not so) different proofs. This can seem odd, but we think it is unavoidable. Note for instance that there is a similar disjunction in Yao and Chen [11].

### 3.4 Appendix

We enlarge our probability space and introduce the coin variables  $(Y_{t,x,k})_{t \geq 0, x \in \mathbb{Z}^d, k \geq 1}$ , that are independent Bernoulli variables with parameter  $\rho \in [0, 1]$  and are independent of every variable introduced previously. We denote by  $\mathbb{P}$  the law on this new probability space.

Fix a finite subset  $B$  of  $\mathbb{Z}^d$ ,  $M \geq 1$  and note  $S_{B,M} = \{0, \dots, M\}^B$ .

Define now, as previously, for  $t \in \mathbb{N}, x \in \mathbb{Z}^d, v \in V, p \in \mathbb{N}$ :

$$E_{t,x}^\rho(v, p) = \sum_{k=1}^p Y_{t,x,k} \mathbb{1}_{\{D_{t,x,k}=v\}} \sum_{i=1}^{+\infty} \mathbb{1}_{\{\sum_{j=0}^i q_{t,x}(j) \leq U_{t,x,k}\}}.$$

Then for  $A \in S_{B,M}$ , we set  ${}_{B}\eta_0^{A,\rho} = A$  and

$${}_{B}\eta_{t+1}^{A,\rho}(x) = \mathbb{1}_B(x) \sum_{v \in V} E_{t,x-v}^\rho(v, {}_{B}\eta_t^{A,\rho}(x-v)).$$

Let  $T \geq 1$ . If  $A \in S_{B,M}$ , then it follows from a natural induction that

$$\forall t \in \{0, \dots, T\} \quad \forall x \in B \quad {}_{B}\eta_t^{A,\rho}(x) = {}_{B}\eta_t^A(x)$$

on the event

$$B_M \cap \bigcap_{0 \leq t \leq T; x \in B; 1 \leq k \leq M} \{Y_{t,x,k} = 1\}.$$

where  $B_M = \{\forall x \in B \quad \forall t \in \{0, \dots, T\} \quad {}_{B}\eta_t^A(x) \leq M\}$ . Denote by  $\mathcal{A}_{B,M,T}$  the  $\sigma$ -field generated by the variables  $({}_{B}\eta_t^A(x))_{t \in \{0, \dots, T\}, A \in S_{B,M}, x \in B}$ . Since the law of the  ${}_{B}\eta_t^{A,\rho}$ 's under  $\mathbb{P}$  is the law of the  ${}_{B}\eta_t^A$ 's under  $\mathbb{P}^{\gamma\rho}$ , the coupling gives for  $E \in \mathcal{A}_{B,M,T}$ .

$$\begin{aligned} \mathbb{P}^{\gamma\rho}(E) & \geq \mathbb{P}\left(E \cap B_M \cap \bigcap_{0 \leq t \leq T; x \in B; 1 \leq k \leq M} \{Y_{t,x,k} = 1\}\right) \\ & = \rho^{(T+1)|B|M} \mathbb{P}(E \cap B_M) = \rho^{(T+1)|B|M} \mathbb{P}^\gamma(E \cap B_M), \end{aligned}$$

which gives Inequality (1.4).

Since the event in Proposition 3.1 belongs to  $\mathcal{A}_{B,1,T} \subset \mathcal{A}_{B,M,T}$ , the proof is complete.

**Acknowledgments.** We would like to thank an anonymous referee who noticed some flaws in a previous version of the article.

**References**

- [1] Carol Bezuidenhout and Geoffrey Grimmett. The critical contact process dies out. *Ann. Probab.*, 18(4):1462–1482, 1990. MR-1071804
- [2] Erik I. Broman. Stochastic domination for a hidden Markov chain with applications to the contact process in a randomly evolving environment. *Ann. Probab.*, 35(6):2263–2293, 2007. MR-2353388
- [3] Francis Comets and Nobuo Yoshida. Branching random walks in space-time random environment: survival probability, global and local growth rates. *J. Theoret. Probab.*, 24(3):657–687, 2011. MR-2822477
- [4] R. Fukushima and N. Yoshida. On exponential growth for a certain class of linear systems. *ALEA Lat. Am. J. Probab. Math. Stat.*, 9(2):323–336, 2012.
- [5] Olivier Garet and Régine Marchand. Growth of a population of bacteria in a dynamical hostile environment, arXiv:1010.4618 *preprint*, 2012.
- [6] Geoffrey Grimmett. *Percolation*, volume 321 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 1999. MR-1707339
- [7] T. E. Harris. A lower bound for the critical probability in a certain percolation process. *Proc. Cambridge Philos. Soc.*, 56:13–20, 1960. MR-0115221
- [8] Yueyun Hu and Nobuo Yoshida. Localization for branching random walks in random environment. *Stochastic Process. Appl.*, 119(5):1632–1651, 2009. MR-2513122
- [9] Thomas M. Liggett. *Stochastic interacting systems: contact, voter and exclusion processes*, volume 324 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999. MR-1717346
- [10] Jeffrey E. Steif and Marcus Warfheimer. The critical contact process in a randomly evolving environment dies out. *ALEA Lat. Am. J. Probab. Math. Stat.*, 4:337–357, 2008. MR-2461788
- [11] Qiang Yao and Xinxing Chen. The complete convergence theorem holds for contact processes in a random environment on  $\mathbb{Z}^d \times \mathbb{Z}_+$ . *Stochastic Process. Appl.*, 122(9):3066–3100, 2012. MR-2946436
- [12] Nobuo Yoshida. Central limit theorem for branching random walks in random environment. *Ann. Appl. Probab.*, 18(4):1619–1635, 2008. MR-2434183