

ON UNIQUENESS OF SOLUTIONS FOR THE STOCHASTIC DIFFERENTIAL EQUATIONS OF NONLINEAR FILTERING¹

BY VLADIMIR M. LUCIC AND ANDREW J. HEUNIS

University of Waterloo

We study a nonlinear filtering problem in which the signal to be estimated is conditioned by the observations. The main results establish pathwise uniqueness for the unnormalized filter equation and uniqueness in law for the normalized and unnormalized filter equations.

1. Introduction. An early work on uniqueness for the stochastic differential equations of nonlinear filtering is that of Szpirglas [15]. The basic viewpoint adopted in [15] is to regard the measure-valued stochastic differential equations of nonlinear filtering as entities quite separate from the original nonlinear filtering problem, for which one can formulate the notions of solution (or weak solution), pathwise uniqueness and uniqueness in law, by essentially adapting these concepts from the theory of Itô stochastic differential equations (see Section IV.1 of [4] or Section IX.1 of [12]). With these notions at hand, it is then established in [15] that pathwise uniqueness and uniqueness in law hold for both the normalized (Fujisaki–Kallianpur–Kunita) and unnormalized (Duncan–Mortensen–Zakai) filter equations, in the case of a nonlinear filtering problem where the signal is a Markov process which is independent of the Wiener process in the observation equation and the sensor function in the observation equation is uniformly bounded.

Our goal is to look at uniqueness for the stochastic differential equations of nonlinear filtering from a point of view very similar to that of [15], but for a nonlinear filtering problem in which there is dependence of the signal on the observations. In fact, we shall look at the specific nonlinear filtering problem where the signal $\{X_t\}$ is an \mathbb{R}^d -valued process solving an equation of the form

$$(1.1) \quad dX_t = b(X_t) dt + B(X_t) dW_t + c(X_t) dV_t,$$

the \mathbb{R}^{d_1} -valued observation process $\{Y_t\}$ is defined by

$$(1.2) \quad Y_t = W_t + \int_0^t h(X_s) ds,$$

and $\{(W_t, V_t)\}$ is a standard $\mathbb{R}^{d_1+d_2}$ -valued Wiener process [precise conditions on the mappings $b(\cdot)$, $B(\cdot)$, $c(\cdot)$ and $h(\cdot)$ will be stated in Section 2]. The pair (1.1) and (1.2) represents a simple model of a signal and observation in

Received November 1999; revised March 2000.

¹Supported by MITACS/NSERC.

AMS 2000 subject classifications. Primary 60G35; secondary 60G44, 60G57.

Key words and phrases. Nonlinear filter equations, weak solutions, uniqueness.

which the signal $\{X_t\}$ depends on the Wiener process $\{W_t\}$ of the observation equation.

Motivated by [15], we shall regard the normalized and unnormalized filter equations for this nonlinear filtering problem as measure-valued stochastic differential equations, defined quite independently of the filtering problem, and will formulate the notions of weak solution, pathwise uniqueness and uniqueness in law for the filter equations. Our main result (see Theorem 2.21 to follow) establishes pathwise uniqueness for the unnormalized filter equation, together with uniqueness in law for the normalized and unnormalized filter equations, subject to reasonably general conditions on the mappings $b(\cdot)$, $B(\cdot)$ and $c(\cdot)$ in the signal equation (1.1), and a uniform boundedness condition on the sensor function $h(\cdot)$ in the observation equation (1.2). As will be seen from the discussion of Section 2 (see Remark 2.22) the elegant semigroup ideas used in [15] to establish pathwise uniqueness do not seem to extend to the filtering problem represented by (1.1) and (1.2), where the signal $\{X_t\}$ depends on the observation Wiener process $\{W_t\}$, and our approach necessarily involves a different method of proof.

In Section 2 we review the normalized and unnormalized filter equations for the nonlinear filtering problem given by (1.1) and (1.2), define weak solutions, pathwise uniqueness, and uniqueness in law for the filter equations and state the main result, namely Theorem 2.21. We also discuss the relationship of this result with other works on uniqueness for the nonlinear filter equations in Remarks 2.22, 2.23 and 2.24. Section 3 is devoted to the proof of the main result, while the proofs of various technical facts and lemmas needed in Section 3 are relegated to Sections 4, 5 and 6.

2. Stochastic differential equations of nonlinear filtering.

REMARK 2.1. For easy access we first summarize most of the basic notation which will be used in the sequel:

(i) For a metric space E , let $\mathcal{B}(E)$ denote the Borel σ -algebra on E , let $B(E)$ denote the set of all real-valued uniformly bounded Borel measurable mappings on E , and, for $\phi \in B(E)$, define the supremum norm by $\|\phi\| := \sup_{x \in E} |\phi(x)|$. Likewise, write $C(E)$ for the set of all real-valued continuous mappings on E , and write $\bar{C}(E)$ for the collection of all members of $C(E)$ which are uniformly bounded.

(ii) For a complete separable metric space E , let $\mathcal{M}^+(E)$ denote the space of all positive bounded measures on the measurable space $(E, \mathcal{B}(E))$, with the usual topology of weak (or narrow) convergence. Then $\mathcal{M}^+(E)$ is separable and metrically topologically complete, and Exercise 9.5.6 of [3] shows that a simple variant of the Prohorov metric turns the topological space $\mathcal{M}^+(E)$ into a complete separable metric space. Also, let $\mathcal{P}(E)$ denote the collection of all members of $\mathcal{M}^+(E)$ which are probability measures. For $\mu \in \mathcal{M}^+(E)$ and a $\mathcal{B}(E)$ -measurable and μ -integrable mapping ϕ from E into \mathbb{R} , write $\mu(\phi)$ or $\mu\phi$ for the integral $\int_E \phi d\mu$.

(iii) For a vector x in a finite-dimensional Euclidean space \mathbb{R}^q , write x^k for the k th scalar entry of x , $\forall k = 1, \dots, q$, and write $|x|$ for the Euclidean norm of x , namely $|x|^2 := \sum_{k=1}^q (x^k)^2$. Also, let $C^\infty(\mathbb{R}^q)$ denote the set of all infinitely smooth real-valued mappings on \mathbb{R}^q , and let $C_c^\infty(\mathbb{R}^q)$ be the collection of all members of $C^\infty(\mathbb{R}^q)$ with compact support. Finally, let $\widehat{C}(\mathbb{R}^q)$ denote the collection of all members of $\overline{C}(\mathbb{R}^q)$ which vanish at infinity.

(iv) For the positive integers q, r , let $\mathbb{R}^{q \times r}$ denote the set of q by r matrices with real entries. Likewise, let $\mathbb{S}_+^{q \times q}$ denote the collection of all members of $\mathbb{R}^{q \times q}$ which are symmetric nonnegative definite, and let $\mathbb{S}_{++}^{q \times q}$ denote the collection of all members of $\mathbb{S}_+^{q \times q}$ which are strictly positive definite.

(v) For the positive integer q , let \mathbb{R}^{q*} denote the compact metric space which is the one-point compactification of the Euclidean space \mathbb{R}^q .

Now consider a nonlinear filtering problem made up of the following basic elements:

E.1 A fixed interval of interest $[0, T]$, with $T \in (0, \infty)$.

E.2 A complete probability space (Ω, \mathcal{F}, P) carrying a filtration $\{\mathcal{F}_t, t \in [0, T]\}$ such that \mathcal{F}_0 includes all null events of (Ω, \mathcal{F}, P) . Defined on (Ω, \mathcal{F}, P) is an \mathbb{R}^d -valued continuous $\{\mathcal{F}_t\}$ -adapted process $\{X_t, t \in [0, T]\}$ and an $\mathbb{R}^{d_1+d_2}$ -valued $\{\mathcal{F}_t\}$ -Wiener process $\{(W_t, V_t), t \in [0, T]\}$ such that (1.1) holds, where $b: \mathbb{R}^d \rightarrow \mathbb{R}^d$, $B: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d_1}$ and $c: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d_2}$ are Borel-measurable and locally bounded functions (i.e., uniformly bounded over bounded subsets of \mathbb{R}^d).

E.3 An \mathbb{R}^{d_1} -valued *observation process* $\{Y_t, t \in [0, T]\}$ defined by (1.2), where $h: \mathbb{R}^d \rightarrow \mathbb{R}^{d_1}$ is Borel-measurable, with

$$(2.1) \quad \mathbb{E} \left[\sum_{k=1}^{d_1} \int_0^T |h^k(X_u)|^2 du \right] < \infty.$$

Define the *observation filtration* $\{\mathcal{F}_t^Y, t \in [0, T]\}$ by

$$(2.2) \quad \begin{aligned} \mathcal{F}_t^Y &:= \sigma\{Y_u, u \in [0, t]\} \vee \mathcal{N}(P) \quad \text{where } \mathcal{N}(P) \\ &:= \{N \in \mathcal{F}: P(N) = 0\}. \end{aligned}$$

From Lemma 1.1 of [10] there exists some $\mathcal{P}(\mathbb{R}^d)$ -valued $\{\mathcal{F}_{t+}^Y\}$ -optional process $\{\pi_t, t \in [0, T]\}$, called the *optimal filter*, which is defined on (Ω, \mathcal{F}, P) and satisfies

$$(2.3) \quad \pi_t \phi = \mathbb{E}[\phi(X_t) | \mathcal{F}_{t+}^Y] \quad \text{a.s.} \quad \forall t \in [0, T], \forall \phi \in B(\mathbb{R}^d).$$

From (2.1) and Jensen's inequality we see that

$$\mathbb{E} \left[\sum_{k=1}^{d_1} \int_0^T [\pi_u(|h^k|)]^2 du \right] < \infty,$$

and we can therefore define the \mathbb{R}^{d_1} -valued *innovations process* $\{I_t, t \in [0, T]\}$ by

$$(2.4) \quad I_t^k := Y_t^k - \int_0^t \pi_s h^k ds \quad \forall t \in [0, T], k = 1, \dots, d_1.$$

An important property of the innovations process is that $\{I_t, t \in [0, T]\}$ is an \mathbb{R}^{d_1} -valued $\{\mathcal{F}_{t+}^Y\}$ -Wiener process [see Theorem VI.8.4 of [13], observing that the filtration $\{\mathcal{G}_t\}$ on page 322 of [13] corresponds to our $\{\mathcal{F}_{t+}^Y\}$]. Since $\{I_t\}$ is continuous, it is necessarily $\{\mathcal{F}_t^Y\}$ -adapted, thus $\{I_t, t \in [0, T]\}$ is a $\{\mathcal{F}_t^Y\}$ -Wiener process. Now define $m: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times (d_1 + d_2)}$ by

$$m(x) := [B(x) \quad c(x)] \quad \forall x \in \mathbb{R}^d,$$

and put

$$(2.5a) \quad \mathcal{A}\phi(x) := \sum_{i=1}^d b^i(x) \partial_i \phi(x) + \frac{1}{2} \sum_{i,j=1}^d [m(x)m^T(x)]^{ij} \partial_i \partial_j \phi(x) \\ \forall x \in \mathbb{R}^d, \phi \in C^\infty(\mathbb{R}^d);$$

$$(2.5b) \quad \mathcal{B}_k \phi(x) := \sum_{i=1}^d B^{ik}(x) \partial_i \phi(x) \quad \forall x \in \mathbb{R}^d, \phi \in C^\infty(\mathbb{R}^d), k = 1, \dots, d_1.$$

For each $\phi \in C_c^\infty(\mathbb{R}^d)$ one sees from (1.1), (1.2) and Itô's formula that the process

$$(2.6) \quad M_t^\phi := \phi(X_t) - \int_0^t \mathcal{A}\phi(X_s) ds, \quad t \in [0, T],$$

is a square-integrable $\{\mathcal{F}_t\}$ -martingale with

$$(2.7) \quad \langle M^\phi, W^k \rangle_t = \int_0^t \mathcal{B}_k \phi(X_u) du, \quad t \in [0, T], k = 1, \dots, d_1.$$

This observation, together with Theorem VI.8.11 of [13], establishes the following.

THEOREM 2.2. *For the nonlinear filtering problem given by E.1, E.2 and E.3, the optimal filter $\{\pi_t, t \in [0, T]\}$ satisfies*

$$(2.8) \quad \pi_t \phi = \pi_0 \phi + \int_0^t \pi_s (\mathcal{A}\phi) ds + \int_0^t \sum_{k=1}^{d_1} [\pi_s (h^k \phi + \mathcal{B}_k \phi) \\ - (\pi_s h^k)(\pi_s \phi)] dI_s^k \quad \forall \phi \in C_c^\infty(\mathbb{R}^d).$$

The relation (2.8) is known variously as the Fujisaki–Kallianpur–Kunita equation, the Kushner–Stratonovich equation or the normalized filter equation.

REMARK 2.3. Since $C_c^\infty(\mathbb{R}^d)$ is dense in $\widehat{C}(\mathbb{R}^d)$, with respect to the supremum norm, it must be convergence determining (see Problem 3.11.11 of [3]). Now it follows from (2.8) that $\{\pi_t, t \in [0, T]\}$ is a continuous $\mathcal{P}(\mathbb{R}^d)$ -valued process, and therefore $\{\mathcal{F}_t^Y\}$ -adapted. Thus, we can replace \mathcal{F}_{t+}^Y in (2.3) by \mathcal{F}_t^Y .

The characterization of $\{\pi_t\}$ given by Theorem 2.2 becomes useful when some form of uniqueness is established for (2.8). The approach adopted here is suggested by [15], which in turn is motivated by the results of [16] on weak solutions, pathwise uniqueness, and uniqueness in law for Itô stochastic differential equations (see Section IV.1 of [4] or Section IX.1 of [12] for a comprehensive account of these ideas). Taking advantage of the fact that the innovations process $\{I_t\}$ which “drives” 2.8 is a standard $\{\mathcal{F}_t^Y\}$ -Wiener process, we can follow [15] and regard the normalized filter equation as an entity quite separate from the nonlinear filtering problem, namely as a probability-measure valued stochastic differential equation driven by a standard Wiener process, for which one can formulate the notions of weak solution, pathwise uniqueness and uniqueness in law as follows (compare [15], Definition III.1, V.1, V.2 and [1], Definition 9.1).

DEFINITION 2.4. The pair $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t, \tilde{I}_t)\}$ is a *weak solution* of the normalized filter equation when:

- (i) $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P})$ is a complete filtered probability space.
- (ii) $\{\tilde{I}_t, t \in [0, T]\}$ is an \mathbb{R}^{d_1} -valued $\{\tilde{\mathcal{F}}_t\}$ -Wiener process on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$.
- (iii) $\{\tilde{\pi}_t, t \in [0, T]\}$ is a $\mathcal{P}(\mathbb{R}^d)$ -valued continuous $\{\tilde{\mathcal{F}}_t\}$ -adapted process such that

$$(2.9) \quad \tilde{P} \left(\int_0^T \sum_{k=1}^{d_1} [\tilde{\pi}_s |h^k|]^2 ds < \infty \right) = 1,$$

and, for each $\phi \in C_c^\infty(\mathbb{R}^d)$, the following holds to within indistinguishability:

$$(2.10) \quad \begin{aligned} \tilde{\pi}_t \phi &= \tilde{\pi}_0 \phi + \int_0^t \tilde{\pi}_s (\mathcal{A} \phi) ds \\ &+ \sum_{k=1}^{d_1} \int_0^t \left[\tilde{\pi}_s (h^k \phi + \mathcal{B}_k \phi) - (\tilde{\pi}_s h^k)(\tilde{\pi}_s \phi) \right] d\tilde{I}_s^k, \quad t \in [0, T]. \end{aligned}$$

REMARK 2.5. The terminology that $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P})$ is a “complete filtered probability space” will always be understood to mean that $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ is a complete probability space carrying the filtration $\{\tilde{\mathcal{F}}_t, t \in [0, T]\}$, and $\tilde{\mathcal{F}}_0$ includes all P -null events in $\tilde{\mathcal{F}}$.

REMARK 2.6. In view of Definition 2.4, it follows that $\{(\Omega, \mathcal{F}, \{\mathcal{F}_t^Y\}, P), (\pi_t, I_t)\}$ for $\{\mathcal{F}_t^Y, t \in [0, T]\}$, $\{\pi_t, t \in [0, T]\}$ and $\{I_t, t \in [0, T]\}$ defined by (2.2), (2.3) and 2.4 is a weak solution of the normalized filter equation.

DEFINITION 2.7. The normalized filter equation has the property of *pathwise uniqueness* when the following holds: If $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t^1, \tilde{I}_t)\}$ and $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t^2, \tilde{I}_t)\}$ are weak solutions of the normalized filter equation such that $\tilde{P}(\tilde{\pi}_0^1 = \tilde{\pi}_0^2) = 1$, then

$$\tilde{P}(\tilde{\pi}_t^1 = \tilde{\pi}_t^2 \forall t \in [0, T]) = 1.$$

REMARK 2.8. For the next definition we shall need the following notation: if ξ is a measurable mapping from some probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ into a separable metric space E , then $\mathcal{L}_{\tilde{P}}(\xi)$ is the probability measure on the Borel σ -algebra $\mathcal{B}(E)$ defined by $\mathcal{L}_{\tilde{P}}(\xi)(\Gamma) := \tilde{P}\{\xi \in \Gamma\}$ for each $\Gamma \in \mathcal{B}(E)$.

DEFINITION 2.9. The normalized filter equation has the property of *uniqueness in joint law* when the following holds: If $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t, \tilde{I}_t)\}$ and $\{(\bar{\Omega}, \bar{\mathcal{F}}, \{\bar{\mathcal{F}}_t\}, \bar{P}), (\bar{\pi}_t, \bar{I}_t)\}$ are weak solutions of the normalized filter equation such that $\mathcal{L}_{\tilde{P}}(\tilde{\pi}_0) = \mathcal{L}_{\bar{P}}(\bar{\pi}_0)$, then the processes $\{(\tilde{\pi}_t, \tilde{I}_t), t \in [0, T]\}$ and $\{(\bar{\pi}_t, \bar{I}_t), t \in [0, T]\}$ have the same finite-dimensional distributions.

REMARK 2.10. Under certain conditions one can associate a simpler *unnormlized filter equation* with the normalized filter equation. For this purpose the following additional notation is useful: if $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P})$ is a complete filtered probability space, $\{\tilde{M}_t\}$ is a continuous $\{\tilde{\mathcal{F}}_t\}$ -semimartingale, and $\{\tilde{\gamma}_t\}$ is a locally bounded $\{\tilde{\mathcal{F}}_t\}$ -progressively measurable process, then $\tilde{\gamma} \cdot \tilde{M}$ denotes the stochastic integral of $\tilde{\gamma}$ with respect to \tilde{M} . Also, put

$$\mathcal{E}(\tilde{M})_t := \exp\left(\tilde{M}_t - \frac{1}{2}\langle \tilde{M} \rangle_t\right).$$

Now let $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t, \tilde{I}_t)\}$ be a weak solution of the normalized filter equation, and define

$$(2.11) \quad \tilde{Y}_t^k := \tilde{I}_t^k + \int_0^t \tilde{\pi}_s h^k ds \quad \forall t \in [0, T], \quad k = 1, \dots, d_1;$$

$$(2.12) \quad \tilde{\chi}_t := \mathcal{E}\left(-\sum_{k=1}^{d_1} (\tilde{\pi} h^k) \cdot \tilde{I}^k\right)_t \quad \forall t \in [0, T].$$

Since $\{\tilde{I}_t, t \in [0, T]\}$ is a $\{\tilde{\mathcal{F}}_t\}$ -Wiener process, it follows that $\{\tilde{\chi}_t, t \in [0, T]\}$ is a continuous strictly positive $\{\tilde{\mathcal{F}}_t\}$ -local martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$, and

$$(2.13) \quad \frac{1}{\tilde{\chi}_t} = \mathcal{E}\left(\sum_{k=1}^{d_1} (\tilde{\pi} h^k) \cdot \tilde{Y}^k\right)_t \quad \forall t \in [0, T].$$

Define the $\mathcal{M}^+(\mathbb{R}^d)$ -valued process $\{\tilde{\sigma}_t, t \in [0, T]\}$ on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ by

$$(2.14) \quad \tilde{\sigma}_t \phi := \frac{\tilde{\pi}_t \phi}{\tilde{\chi}_t} \quad \forall t \in [0, T], \quad \phi \in B(\mathbb{R}^d).$$

Hence

$$(2.15) \quad \tilde{\sigma}_t \phi := (\tilde{\pi}_t \phi) \mathcal{E} \left(\sum_{k=1}^{d_1} (\tilde{\pi} h^k) \cdot \tilde{Y}^k \right)_t \quad \forall t \in [0, T], \quad \phi \in B(\mathbb{R}^d),$$

and, in light of (2.9), we see that

$$\tilde{P} \left(\int_0^T \sum_{k=1}^{d_1} [\tilde{\sigma}_s |h^k \phi + \mathcal{B}_k \phi|^2] ds < \infty \right) = 1 \quad \forall \phi \in C_c^\infty(\mathbb{R}^d) \cup \{1\}.$$

Using Itô's formula and relation (2.10), we easily arrive at the *Duncan–Mortensen–Zakai equation* or *unnormalized filter equation*: for each $\phi \in C_c^\infty(\mathbb{R}^d) \cup \{1\}$ we have

$$(2.16) \quad \begin{aligned} \tilde{\sigma}_t \phi &= \tilde{\sigma}_0 \phi + \int_0^t \tilde{\sigma}_s(\mathcal{A} \phi) ds \\ &+ \sum_{k=1}^{d_1} \int_0^t \tilde{\sigma}_s(h^k \phi + \mathcal{B}_k \phi) d\tilde{Y}_s^k \quad \forall t \in [0, T]. \end{aligned}$$

REMARK 2.11. From Remark 2.3 and (2.15) we see that $t \rightarrow \tilde{\sigma}_t \phi: [0, T] \rightarrow \mathbb{R}$ is continuous for each bounded continuous $\phi: \mathbb{R}^d \rightarrow \mathbb{R}$, thus $\{\tilde{\sigma}_t\}$ is a *continuous* $\mathcal{M}^+(\mathbb{R}^d)$ -valued process which is $\{\mathcal{F}_t\}$ -adapted. Moreover, from (2.15), we see that the random element $\tilde{\sigma}_0$ takes values in $\mathcal{P}(\mathbb{R}^d)$, the set of *probability measures* on \mathbb{R}^d .

REMARK 2.12. If, in (2.15), we use the optimal filter $\{\pi_t\}$ in place of $\{\tilde{\pi}_t\}$ and the observation process $\{Y_t\}$ in place of $\{\tilde{Y}_t\}$ to get an $\mathcal{M}^+(\mathbb{R}^d)$ -valued and $\{\mathcal{F}_t^Y\}$ -adapted process $\{\sigma_t\}$, namely

$$(2.17) \quad \sigma_t \phi := (\pi_t \phi) \mathcal{E} \left(\sum_{k=1}^{d_1} (\pi h^k) \cdot Y^k \right)_t \quad \forall t \in [0, T], \quad \phi \in B(\mathbb{R}^d),$$

then $\{\sigma_t\}$ is called the *unnormalized optimal filter* for the filtering problem given by (1.1) and (1.2).

REMARK 2.13. In (2.16) the “driving process” $\{\tilde{Y}_t\}$ is the continuous $\{\tilde{\mathcal{F}}_t\}$ -semimartingale defined on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ by (2.11). Equation (2.16) becomes more tractable if we can replace \tilde{P} with some equivalent probability measure \tilde{Q} such that $\{\tilde{Y}_t, t \in [0, T]\}$ is an $\{\tilde{\mathcal{F}}_t\}$ -Wiener process with respect to \tilde{Q} . To this end, observe from (2.12) that $\{(\tilde{\chi}_t, \tilde{\mathcal{F}}_t), t \in [0, T]\}$ is a continuous local martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$, and that, *if it is a martingale*, then

$$(2.18) \quad \tilde{Q}(A) := E^{\tilde{P}}[\tilde{\chi}_T; A] \quad \forall A \in \tilde{\mathcal{F}},$$

defines a probability measure \tilde{Q} on $(\tilde{\Omega}, \tilde{\mathcal{F}})$ which is equivalent to the probability measure \tilde{P} , namely

$$(2.19) \quad \tilde{P} \equiv \tilde{Q} [\tilde{\mathcal{F}}].$$

From (2.11), (2.12) and the Girsanov theorem, it then follows that $\{(\tilde{Y}_t, \tilde{\mathcal{F}}_t), t \in [0, T]\}$ is a Wiener process on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{Q})$.

REMARK 2.14. A sufficient condition on the weak solution $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t, \tilde{I}_t)\}$ and sensor function $h(\cdot)$ which ensures $\{(\tilde{\chi}_t, \tilde{\mathcal{F}}_t), t \in [0, T]\}$ is a martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{P})$ is that

$$E^{\tilde{P}} \left[\exp \left(\frac{1}{2} \sum_{k=1}^{d_1} \int_0^T [\tilde{\pi}_s h^k]^2 ds \right) \right] < \infty$$

(see Corollary 3.5.13 of [5]). In particular, this condition always holds when $h^k \in B(\mathbb{R}^d)$, $k = 1, \dots, d_1$.

With the preceding discussion in mind, we next formulate the notion of weak solution of the unnormalized filter equation, pathwise uniqueness and uniqueness in law (compare with Definition IV.1 of [15]).

DEFINITION 2.15. A pair $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t, \tilde{Y}_t)\}$ is a *weak solution* of the unnormalized filter equation when:

- (i) $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q})$ is a complete filtered probability space.
- (ii) $\{\tilde{Y}_t, t \in [0, T]\}$ is an \mathbb{R}^{d_1} -valued $\{\tilde{\mathcal{F}}_t\}$ -Wiener process.
- (iii) $\{\tilde{\sigma}_t, t \in [0, T]\}$ is a $\mathcal{M}^+(\mathbb{R}^d)$ -valued continuous $\{\tilde{\mathcal{F}}_t\}$ -adapted process such that the random element $\tilde{\sigma}_0$ takes values in $\mathcal{P}(\mathbb{R}^d)$, and, for each $\phi \in C_c^\infty(\mathbb{R}^d) \cup \{1\}$, we have the following:

$$(2.20) \quad \tilde{Q} \left(\int_0^T \sum_{k=1}^{d_1} [\tilde{\sigma}_s |h^k \phi + \mathcal{B}_k \phi|^2] ds < \infty \right) = 1;$$

- (a)
- (b) The LHS and RHS of (2.15) are indistinguishable.

DEFINITION 2.16. The unnormalized filter equation has the property of *pathwise uniqueness* when the following holds: if $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^1, \tilde{Y}_t)\}$ and $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^2, \tilde{Y}_t)\}$ are weak solutions of the unnormalized filter equation such that $\tilde{Q}(\tilde{\sigma}_0^1 = \tilde{\sigma}_0^2) = 1$, then

$$\tilde{Q}(\tilde{\sigma}_t^1 = \tilde{\sigma}_t^2, \forall t \in [0, T]) = 1.$$

DEFINITION 2.17. The unnormalized filter equation has the property of *uniqueness in joint law* when the following holds: if $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t, \tilde{Y}_t)\}$ and $\{(\bar{\Omega}, \bar{\mathcal{F}}, \{\bar{\mathcal{F}}_t\}, \bar{Q}), (\bar{\sigma}_t, \bar{Y}_t)\}$ are weak solutions of the unnormalized filter equation such that $\mathcal{L}_{\tilde{Q}}(\tilde{\sigma}_0) = \mathcal{L}_{\bar{Q}}(\bar{\sigma}_0)$, then $\{(\tilde{\sigma}_t, \tilde{Y}_t), t \in [0, T]\}$ and $\{(\bar{\sigma}_t, \bar{Y}_t), t \in [0, T]\}$ have the same finite-dimensional distributions.

In this paper our goal is to establish pathwise uniqueness for the unnormalized filter equation and uniqueness in joint law for both the normalized and unnormalized filter equations. To this end we postulate the following conditions on the mappings $b(\cdot)$, $B(\cdot)$, $c(\cdot)$ in (1.1), and the mapping $h(\cdot)$ in (1.2):

CONDITION 2.18. The mapping $b: \mathbb{R}^d \rightarrow \mathbb{R}^d$ is Borel-measurable, and the mappings $B: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d_1}$ and $c: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d_2}$ are continuous. There exists a constant $C \in [0, \infty)$ such that

$$\max_{i, j, k} \{|b^i(x)|, |B^{ij}(x)|, |c^{ik}(x)|\} \leq C[1 + |x|] \quad \forall x \in \mathbb{R}^d.$$

CONDITION 2.19. The mapping $c: \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d_2}$ is such that the matrix $c(x)c^T(x)$ is strictly positive definite for every $x \in \mathbb{R}^d$.

CONDITION 2.20. The mapping $h: \mathbb{R}^d \rightarrow \mathbb{R}^{d_1}$ is Borel-measurable and uniformly bounded.

We can now state our main result.

THEOREM 2.21. *Suppose that Conditions 2.18, 2.19 and 2.20 hold for the nonlinear filtering problem given by E.1, E.2 and E.3. Then:*

- (i) *The unnormalized filter equation has the property of pathwise uniqueness.*
- (ii) *The normalized filter equation has the property of uniqueness in joint law.*
- (iii) *The unnormalized filter equation has the property of uniqueness in joint law.*

REMARK 2.22. Here [15] establishes pathwise uniqueness and uniqueness in law for the normalized and unnormalized filter equations corresponding to the following nonlinear filtering problem: the signal $\{X_t\}$ is a homogeneous Markov process with values in a complete separable metric space E and weak infinitesimal generator \mathcal{A} , the observation process is

$$Y_t := W_t + \int_0^t h(X_u) du, \quad t \in [0, T],$$

where $\{W_t\}$ is an \mathbb{R}^{d_1} -valued Wiener process *independent* of the Markov process $\{X_t\}$, and the sensor function $h: E \rightarrow \mathbb{R}^{d_1}$ is uniformly bounded and $\mathcal{B}(E)$ -measurable. In this context, by a weak solution of the unnormalized

filter equation is meant a pair $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t, \tilde{Y}_t)\}$ such that:

- (a) $(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q})$ is a complete filtered probability space.
- (b) $\{\tilde{Y}_t, t \in [0, T]\}$ is an \mathbb{R}^{d_1} -valued $\{\tilde{\mathcal{F}}_t\}$ -Wiener process.
- (c) $\{\tilde{\sigma}_t, t \in [0, T]\}$ is a $\mathcal{M}^+(E)$ -valued, cadlag (or r.c.l.l.) $\{\tilde{\mathcal{F}}_t\}$ -adapted process, the random element $\tilde{\sigma}_0$ takes values in $\mathcal{P}(E)$, and $\sup_{t \in [0, T]} \times E[|\tilde{\sigma}_t 1|^2] < \infty$.
- (d) For each $\phi \in \mathcal{D}(\mathcal{A})$ (the domain of the generator \mathcal{A}) one has to within indistinguishability that

$$(2.21) \quad \begin{aligned} \tilde{\sigma}_t \phi &= \tilde{\sigma}_0 \phi + \int_0^t \tilde{\sigma}_s(\mathcal{A} \phi) ds \\ &+ \sum_{k=1}^{d_1} \int_0^t \tilde{\sigma}_s(h^k \phi) d\tilde{Y}_s^k, \quad \forall t \in [0, T]. \end{aligned}$$

(See Definition IV.1 of [15].) The nice thing about (2.21) is that it includes reference to just one unbounded linear operator, namely the infinitesimal generator \mathcal{A} of the signal process, and the resolvent identity can be used to eliminate \mathcal{A} and rewrite (2.21) in the form

$$(2.22) \quad \tilde{\sigma}_t \phi = \tilde{\sigma}_0(P_t \phi) + \sum_{k=1}^{d_1} \int_0^t \tilde{\sigma}_s(h^k P_{t-s} \phi) d\tilde{Y}_s^k \quad \forall t \in [0, T],$$

where $\{P_t\}$ is the Borel semigroup with infinitesimal generator \mathcal{A} . There is complete equivalence between (2.21) and (2.23) in the sense that if the pair $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t, \tilde{Y}_t)\}$ is subject to (a), (b), (c), then (2.23) holds for each $\phi \in \mathcal{D}(\mathcal{A})$ if and only if (2.24) holds for each $\phi \in B(E)$ (see Théorème IV.1 of [15]). Consequently, it is enough to establish pathwise uniqueness for (2.24) in order to conclude pathwise uniqueness for the unnormalized filter equation. The advantage of (2.22) is that it involves only the *bounded* linear operators $\{P_t\}$, and this structure makes it possible to establish pathwise uniqueness for solutions of (2.22) by iterating a simple integral inequality (see Section V.2 of [15]). Comparing (2.21) with the unnormalized filter equation (2.16) for the nonlinear filtering problem defined by (1.1) and (1.2), we see that (2.16) includes two unbounded linear operators, namely the first-order differential operator \mathcal{B}_k which results from dependence of the signal $\{X_t\}$ on the Wiener process $\{W_t\}$ of the observation equation, as well as the second-order differential operator \mathcal{A} corresponding to the signal process $\{X_t\}$. In this case there seems to be no clear way of adapting the elegant semigroup ideas of [15] to remove both of these unbounded operators and get an equivalent equation involving just bounded linear operators. Accordingly, the approach that we shall use to establish Theorem 2.21(i) is different from that of [15] and relies on a uniqueness theorem for measure-valued evolution equations (see Theorem 3.6 to follow).

REMARK 2.23. Uniqueness for the normalized and unnormalized filter equations has also been studied by [1], [10] and [14] from a somewhat different

point of view than that taken by [15] and the present work. To see this in the context of the filtering problem given by (1.1) and (1.2), observe from Remark 2.12 that the unnormalized optimal filter $\{\sigma_t\}$ solves the Duncan–Mortensen–Zakai equation, namely for each $\phi \in C_c^\infty(\mathbb{R}^d) \cup \{1\}$ we have

$$(2.23) \quad \begin{aligned} \sigma_t \phi &= \pi_0 \phi + \int_0^t \sigma_s(\mathcal{A}\phi) ds \\ &+ \sum_{k=1}^{d_1} \int_0^t \sigma_s(h^k \phi + \mathcal{B}_k \phi) dY_s^k \quad \forall t \in [0, T]. \end{aligned}$$

With this in mind, the following question is natural: suppose that $\{\rho_t\}$ is some $\mathcal{M}^+(\mathbb{R}^d)$ -valued, cadlag, and $\{\mathcal{F}_t^Y\}$ -adapted process on (Ω, \mathcal{F}, P) , such that for each $\phi \in C_c^\infty(\mathbb{R}^d) \cup \{1\}$ we have

$$(2.24) \quad \begin{aligned} \rho_t \phi &= \pi_0 \phi + \int_0^t \rho_s(\mathcal{A}\phi) ds \\ &+ \sum_{k=1}^{d_1} \int_0^t \rho_s(h^k \phi + \mathcal{B}_k \phi) dY_s^k \quad \forall t \in [0, T]. \end{aligned}$$

Does it follow that $\{\sigma_t\}$ and $\{\rho_t\}$ are indistinguishable? The works of [1], Theorem 3.1, [10], Theorems 4.2 and 4.7 and [14], Theorem 3.1 provide conditions on the nonlinear filtering problem for which the answer is in the affirmative. Uniqueness in this sense is useful for the following reason: the observation process $\{Y_t\}$ is the random data that “drives” the unnormalized filter equation (2.23), and if we can “nonanticipatively” use the individual paths of $\{Y_t\}$ as data to compute a measure-valued process $\{\rho_t\}$ which satisfies (2.26) (e.g., by a numerical method) then uniqueness ensures that $\{\rho_t\}$ is in fact the desired unnormalized optimal filter $\{\sigma_t\}$. It should be noted that uniqueness in this sense can be established for much more general nonlinear filtering problems than that represented by the simple model (1.1) and (1.2). In fact, Theorem 3.1 of [1] deals with a filtering problem in which the signal process takes values in a complete separable metric space (not necessarily locally compact), the sensor function $h(\cdot)$ need not be uniformly bounded but only satisfy a finite-energy condition similar to (2.1), the dependence of the signal $\{X_t\}$ on the Wiener process $\{W_t\}$ is more general than that given by the explicit model (1.1), (1.2) (see (1.3) of [1]), and the joint signal–observation process $\{(X_t, Y_t)\}$ is the cadlag solution of a well-posed martingale problem.

The sense of pathwise uniqueness in the preceding paragraph is different from that established by Theorem 2.21(i), since the candidate solution $\{\rho_t\}$ of the filter equation (2.24) is postulated to be adapted specifically to the observation filtration $\{\mathcal{F}_t^Y\}$ (in fact, the arguments used in [1], [10] and [14] rely crucially on this restriction). In contrast, Theorem 2.21(i) establishes pathwise uniqueness in the more general sense of Definition 2.16, where the candidate solutions $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^1, \tilde{Y}_t)\}$ and $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^2, \tilde{Y}_t)\}$ are defined on an arbitrary filtered probability space, and there is no insistence that the measure-valued components $\{\tilde{\sigma}_t^1\}$ and $\{\tilde{\sigma}_t^2\}$ of the two solutions be

adapted to the self-filtration of $\{\tilde{Y}_t\}$. The usefulness of this latter notion of pathwise uniqueness is that, by an adaptation to the filter equations of the construction of [16], it leads to uniqueness in law for the normalized and unnormalized filter equations [see Theorem 2.21(ii) and (iii), and recall Definitions 2.9 and 2.17]. Uniqueness in law turns out to be essential for studying weak limits and approximations of the nonlinear filter equations by the method of martingale problems and weak convergence.

REMARK 2.24. Using the method of stochastic flows and backward equations, [8], Theorem 6.2.8, and [7] establish a form of pathwise uniqueness for the unnormalized filter equation. Rather restrictive boundedness and smoothness conditions on the coefficients of (1.1) and (1.2) seem necessary for this method to work. One should also note that the approach of [15], which is based on the equivalence of (2.21) and (2.22), finds its inspiration in an earlier study of pathwise uniqueness for the filter equations due to [6].

REMARK 2.25. A basic property of Itô stochastic differential equations due to [16] is that pathwise uniqueness implies uniqueness in joint law, so that pathwise uniqueness is the stronger of the two uniqueness properties. It is shown in [15] that the basic Yamada–Watanabe argument extends to the measure-valued filter equations, so that pathwise uniqueness is again the stronger property [this is how we will conclude (ii) and (iii) from (i) in Theorem 2.21]. Linearity of the unnormalized filter equation in fact implies the converse, so that for this equation the two uniqueness properties are actually equivalent.

THEOREM 2.26. *Suppose that Conditions 2.18, 2.19 and 2.20 hold for the nonlinear filtering problem given by E.1, E.2 and E.3. Then uniqueness in joint law implies pathwise uniqueness for the unnormalized filter equation.*

3. Proofs of Theorems 2.21 and 2.26. The terminology in the next remark will be useful for dealing with measure-valued evolution equations.

REMARK 3.1. Suppose that E is a complete separable metric space, and $\mathcal{D}: \mathcal{D}(\mathcal{D}) \rightarrow B(E)$ is a mapping with domain $\mathcal{D}(\mathcal{D}) \subset B(E)$. Then $\{\mu_t, t \in [0, \infty)\}$ is an $\mathcal{M}^+(E)$ -valued solution of the evolution equation for $(\mathcal{D}, \mathcal{D}(\mathcal{D}))$, when (i) $\mu_t \in \mathcal{M}^+(E), \forall t \in [0, \infty)$ and $\mu_0 \in \mathcal{P}(E)$; (ii) for each $\Gamma \in \mathcal{B}(E)$, the mapping $t \rightarrow \mu_t(\Gamma): [0, \infty) \rightarrow [0, \infty)$ is Borel-measurable; (iii) for each $f \in \mathcal{D}(\mathcal{D})$ we have $\int_0^t |\mu_s(\mathcal{D}f)| ds < \infty, \forall t \in [0, \infty)$, and

$$(3.1) \quad \mu_t f = \mu_0 f + \int_0^t \mu_s(\mathcal{D}f) ds \quad \forall t \in [0, \infty).$$

Moreover, $\{\mu_t, t \in [0, \infty)\}$ is a $\mathcal{P}(E)$ -valued solution of the evolution equation for $(\mathcal{D}, \mathcal{D}(\mathcal{D}))$ when it is an $\mathcal{M}^+(E)$ -valued solution with $\mu_t(E) = 1, \forall t \in [0, \infty)$. The evolution equation for $(\mathcal{D}, \mathcal{D}(\mathcal{D}))$ is said to have uniqueness in the class of $\mathcal{M}^+(E)$ -valued solutions over the interval $[0, \infty)$ when, for any two

such solutions $\{\mu_t^i, t \in [0, \infty)\}$, $i = 1, 2$, with $\mu_0^1 = \mu_0^2$, it follows that $\mu_t^1 = \mu_t^2$, $\forall t \in [0, \infty)$. The notion of uniqueness within the class of $\mathcal{P}(E)$ -valued solutions over the interval $[0, \infty)$ has an analogous formulation. Finally, the preceding terminology adapts in an obvious way to solutions $\{\mu_t, t \in [0, T]\}$ defined over the finite interval $[0, T]$.

PROOF OF THEOREM 2.21(i). We shall need the following result, the proof of which is given in Section 5.

FACT 3.2. Suppose that Conditions 2.18–2.20 hold and let $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}, \tilde{Y})\}$ be a weak solution of the unnormalized filter equation. Then, for every $\alpha \in (1, \infty)$ there exists a constant $\gamma(\alpha) \in [0, \infty)$ such that

$$(3.2) \quad \mathbf{E}^{\tilde{Q}} \left[\sup_{0 \leq s \leq T} |\tilde{\sigma}_s^1|^\alpha \right] \leq \gamma(\alpha).$$

Now fix two weak solutions $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^i, \tilde{Y}_t)\}$, $i = 1, 2$, of the unnormalized filter equation, such that

$$(3.3) \quad \tilde{Q}[\tilde{\sigma}_0^1 = \tilde{\sigma}_0^2] = 1,$$

and define product measures on $(\mathbb{R}^{2d}, \mathcal{B}(\mathbb{R}^{2d}))$ by

$$\mu_t^{12}(\cdot, \tilde{\omega}) := (\tilde{\sigma}_t^1 \times \tilde{\sigma}_t^2)(\cdot, \tilde{\omega}) \quad \forall (t, \tilde{\omega}) \in [0, T] \times \tilde{\Omega}.$$

A simple application of the Dynkin class theorem establishes the following.

FACT 3.3. For every $\Gamma \in \mathcal{B}(\mathbb{R}^{2d})$, the mapping $(t, \tilde{\omega}) \rightarrow \mu_t^{12}(\Gamma, \tilde{\omega}) : \tilde{\Omega} \times [0, T] \rightarrow [0, \infty)$ is measurable with respect to the $\{\tilde{\mathcal{F}}_t\}$ -progressive σ -algebra.

Also put

$$(3.4) \quad \nu_t^{12}(\Gamma) := \mathbf{E}^{\tilde{Q}}[\mu_t^{12}(\Gamma)] \quad \forall \Gamma \in \mathcal{B}(\mathbb{R}^{2d}), t \in [0, T].$$

It readily follows that ν_t^{12} defines a positive measure on $(\mathbb{R}^{2d}, \mathcal{B}(\mathbb{R}^{2d}))$ for every $t \in [0, T]$. By Fact 3.2,

$$(3.5) \quad \begin{aligned} \nu_t^{12}(\mathbb{R}^{2d}) &= \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 1)(\tilde{\sigma}_t^2 1)] \\ &\leq \left(\mathbf{E}^{\tilde{Q}} \left[\sup_{0 \leq s \leq T} |\tilde{\sigma}_s^1|^2 \right] \right)^{1/2} \left(\mathbf{E}^{\tilde{Q}} \left[\sup_{0 \leq s \leq T} |\tilde{\sigma}_s^2|^2 \right] \right)^{1/2} \\ &\leq \gamma(2) \quad \forall t \in [0, T]. \end{aligned}$$

This shows that ν_t^{12} is a positive measure on $(\mathbb{R}^{2d}, \mathcal{B}(\mathbb{R}^{2d}))$, uniformly bounded with respect to $t \in [0, T]$, while Fact 3.3 with Fubini's theorem shows that

the mapping $t \rightarrow \nu_t^{12}(\Gamma): [0, T] \rightarrow \mathbb{R}$ is Borel-measurable for each $\Gamma \in \mathcal{B}(\mathbb{R}^{2d})$. Next, define $\nu_t^{11}, \nu_t^{22} \in \mathcal{M}^+(\mathbb{R}^{2d})$, $t \in [0, T]$, analogously to ν_t^{12} by

$$(3.6) \quad \nu_t^{11}(\Gamma) := \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 \times \tilde{\sigma}_t^1)(\Gamma)] \quad \forall \Gamma \in \mathcal{B}(\mathbb{R}^{2d}), t \in [0, T], i = 1, 2.$$

In the same way as for ν_t^{12} , we see that ν_t^{11} are positive measures on $(\mathbb{R}^{2d}, \mathcal{B}(\mathbb{R}^{2d}))$, uniformly bounded with respect to $t \in [0, T]$, and the mappings $t \rightarrow \nu_t^{11}(\Gamma): [0, T] \rightarrow \mathbb{R}$ are Borel-measurable for each $\Gamma \in \mathcal{B}(\mathbb{R}^{2d})$, $i = 1, 2$.

REMARK 3.4. For mappings $f_1, f_2 \in B(\mathbb{R}^d)$ define the tensor product of f_1 with f_2 to be the mapping $f_1 \otimes f_2: \mathbb{R}^{2d} \rightarrow \mathbb{R}$ given by

$$f_1 \otimes f_2(x_1, x_2) := f_1(x_1)f_2(x_2) \quad \forall x_1, x_2 \in \mathbb{R}^d.$$

In view of (3.4) and (3.6), for each $f_1, f_2 \in B(\mathbb{R}^d)$ we have

$$(3.7) \quad \nu_t^{12}(f_1 \otimes f_2) = \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 f_1)(\tilde{\sigma}_t^2 f_2)],$$

$$(3.8) \quad \nu_t^{11}(f_1 \otimes f_2) = \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 f_1)(\tilde{\sigma}_t^1 f_2)], \quad i = 1, 2.$$

From (3.3), (3.4) and (3.6) we see that

$$(3.9) \quad \nu_0^{11}, \nu_0^{22} \text{ and } \nu_0^{12} \text{ are probability measures on } \mathcal{B}(\mathbb{R}^{2d}) \text{ and } \nu_0^{11} = \nu_0^{22} = \nu_0^{12}.$$

Using this fact, we shall establish

$$(3.10) \quad \nu_t^{11} = \nu_t^{22} = \nu_t^{12} \quad \forall t \in [0, T],$$

from which pathwise uniqueness follows. Indeed, if (3.10) holds, then for each $f \in B(\mathbb{R}^d)$ we have

$$\nu_t^{11}(f \otimes f) = \nu_t^{22}(f \otimes f) = \nu_t^{12}(f \otimes f) \quad \forall t \in [0, T],$$

and therefore from (3.7) and (3.8),

$$\begin{aligned} \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 f - \tilde{\sigma}_t^2 f)^2] &= \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 f)(\tilde{\sigma}_t^1 f)] - 2\mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 f)(\tilde{\sigma}_t^2 f)] + \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^2 f)(\tilde{\sigma}_t^2 f)] \\ &= \nu_t^{11}(f \otimes f) - 2\nu_t^{12}(f \otimes f) + \nu_t^{22}(f \otimes f) = 0. \end{aligned}$$

Thus, for each $t \in [0, T]$ and $f \in B(\mathbb{R}^d)$, we have

$$(3.11) \quad \tilde{Q}[\tilde{\sigma}_t^1 f = \tilde{\sigma}_t^2 f] = 1.$$

Now $\widehat{C}(\mathbb{R}^d)$ equipped with the supremum norm $\|\cdot\|$ is separable. Thus, from (3.11), for each $t \in [0, T]$ there is a \tilde{Q} -null event $N_t \in \tilde{\mathcal{F}}$ such that, for each $\tilde{\omega} \notin N_t$, we have

$$(3.12) \quad \tilde{\sigma}_t^1(\tilde{\omega})f = \tilde{\sigma}_t^2(\tilde{\omega})f \quad \forall f \in \widehat{C}(\mathbb{R}^d).$$

But $\widehat{C}(\mathbb{R}^d)$ separates bounded positive measures on $\mathcal{B}(\mathbb{R}^d)$ (see Problem 5.4.25 of [5]), thus (3.12) establishes $\tilde{Q}[\tilde{\sigma}_t^1 = \tilde{\sigma}_t^2] = 1$ for each $t \in [0, T]$. Now Theorem 2.21(i) follows from the fact that $\{\tilde{\sigma}_t^i, t \in [0, T]\}$ are continuous (recall Definition 2.15).

It therefore remains to establish (3.10) in order to prove Theorem 2.21(i). To this end, for each $x_1, x_2 \in \mathbb{R}^d$ define the $2d \times 2d$ matrix $\bar{a}(x_1, x_2)$, the $2d$ vector $\bar{b}(x_1, x_2)$, and the real number $\bar{h}(x_1, x_2)$ by

$$(3.13a) \quad \bar{a}(x_1, x_2) := \begin{bmatrix} cc^T(x_1) & 0 \\ 0 & cc^T(x_2) \end{bmatrix} + \begin{bmatrix} B(x_1) \\ B(x_2) \end{bmatrix} \begin{bmatrix} B^T(x_1) & B^T(x_2) \end{bmatrix},$$

$$(3.13b) \quad \bar{b}(x_1, x_2) := \begin{bmatrix} b(x_1) + B(x_1)h(x_2) \\ b(x_2) + B(x_2)h(x_1) \end{bmatrix},$$

$$(3.13c) \quad \bar{h}(x_1, x_2) := \sum_{k=1}^{d_1} h^k(x_1)h^k(x_2).$$

Observe that the matrix $\bar{a}(x_1, x_2)$ is symmetric and strictly positive definite (see Condition 2.19), and let $\bar{\mathcal{A}}$ be the second-order linear differential operator corresponding to the matrices \bar{a} and \bar{b} , namely

$$(3.14) \quad \bar{\mathcal{A}}\phi(x) := \sum_{i=1}^{2d} \bar{b}^i(x)\partial_i\phi(x) + \frac{1}{2} \sum_{i,j=1}^{2d} \bar{a}^{ij}(x)\partial_i\partial_j\phi(x) \\ \forall \phi \in C^\infty(\mathbb{R}^{2d}), \quad x \in \mathbb{R}^{2d}.$$

From (3.13a), (3.13b), Conditions 2.18 and 2.20, there is a constant $K \in [0, \infty)$ such that

$$(3.15) \quad \max_i |\bar{b}^i(x)| \leq K[1 + |x|], \quad \max_{i,j} |\bar{a}^{ij}(x)| \leq K[1 + |x|^2] \quad \forall x \in \mathbb{R}^{2d},$$

and the operator $\bar{\mathcal{A}}$ has the following property, which is established in Section 4.

LEMMA 3.5. *Suppose that Conditions 2.18–2.20 hold. Then $\{\nu_t^{11}, t \in [0, T]\}$, $\{\nu_t^{12}, t \in [0, T]\}$, and $\{\nu_t^{22}, t \in [0, T]\}$, given by (3.4) and (3.6), are $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solutions of the evolution equation for $(\bar{\mathcal{A}} + \bar{h}, \text{span} \times \{1, C_c^\infty(\mathbb{R}^{2d})\})$.*

It remains to show that the evolution equation for $(\bar{\mathcal{A}} + \bar{h}, \text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\})$ has uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solutions over the interval $[0, T]$, since this fact, along with (3.9) and Lemma 3.5, gives (3.10), as required to establish Theorem 3.6. To this end we need the following result on uniqueness of measure-valued solutions of the evolution equation corresponding to a multiplicatively perturbed linear second-order differential operator

on Euclidean space:

THEOREM 3.6. *Let \mathcal{L} be the linear second-order differential operator on the finite-dimensional Euclidean space \mathbb{R}^q defined by*

$$(3.16a) \quad \mathcal{D}(\mathcal{L}) := \text{span}\{1, C_c^\infty(\mathbb{R}^q)\},$$

$$(3.16b) \quad \mathcal{L}f(x) := \sum_i \beta^i(x) \partial_i f(x) + \frac{1}{2} \sum_{i,j} \alpha^{ij}(x) \partial_i \partial_j f(x) \quad \forall x \in \mathbb{R}^q, \forall f \in \mathcal{D}(\mathcal{L}),$$

where $\beta: \mathbb{R}^q \rightarrow \mathbb{R}^q$ is Borel measurable, $\alpha: \mathbb{R}^q \rightarrow \mathbb{S}_{++}^{q \times q}$ is continuous, and there exists a constant $K \in [0, \infty)$ such that

$$(3.17) \quad |\beta^i(x)| \leq K(1 + |x|), \quad |\alpha^{ij}(x)| \leq K(1 + |x|^2) \quad \forall x \in \mathbb{R}^q.$$

If $\lambda \in B(\mathbb{R}^q)$ then the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$ has uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions over the interval $[0, T]$.

To complete the proof of Theorem 2.21(i) we note from (3.13) that $\bar{a}(\cdot)$ is continuous on \mathbb{R}^{2d} , $\bar{b}(\cdot)$ is Borel-measurable on \mathbb{R}^{2d} and $\bar{h} \in B(\mathbb{R}^{2d})$. That the evolution equation for $(\bar{\mathcal{A}} + \bar{h}, \text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\})$ has uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solutions over the interval $[0, T]$ now follows from (3.15) and Theorem 3.6 with $q := 2d$, $\beta(\cdot) := \bar{b}(\cdot)$, $\alpha(\cdot) := \bar{a}(\cdot)$, and $\lambda(\cdot) := -\bar{h}(\cdot)$. \square

REMARK 3.7. When $\beta(\cdot)$ and $\lambda(\cdot)$ in Theorem 3.6 are *continuous* then \mathcal{L} is a linear operator on $\bar{C}(\mathbb{R}^q)$, and Theorem 3.6 is just a very special consequence of a general theorem of Bhatt and Karandikar (see Theorem 3.4 and Remark 1 of [2]) on uniqueness of measure-valued solutions of perturbed evolution equations. However, when $\beta(\cdot)$ and $\lambda(\cdot)$ are only Borel-measurable, then $\mathcal{L}f(\cdot)$ is not continuous for $f \in \mathcal{D}(\mathcal{L})$, and we cannot directly use the result of [2]. We prove Theorem 3.6 in Section 6.

REMARK 3.8. The proof just given for Theorem 2.21(i) relies on the special structure of the unnormalized filter equation (2.16) and does not appear to extend to the normalized filter equation (2.8). We have therefore not been able to establish pathwise uniqueness in the sense of Definition 2.7 under conditions comparable to those of Theorem 2.21.

PROOF OF THEOREM 2.21(ii). Let $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{P}), (\tilde{\pi}_t, \tilde{I}_t)\}$ and $\{(\bar{\Omega}, \bar{\mathcal{F}}, \{\bar{\mathcal{F}}_t\}, \bar{P}), (\bar{\pi}_t, \bar{I}_t)\}$ be two weak solutions of the normalized filter equation. By an argument similar to that used for Proposition IX.1.4 of [12], to establish uniqueness in joint law it is enough to show that the processes $\{(\tilde{\pi}_t, \tilde{I}_t)\}$ and $\{(\bar{\pi}_t, \bar{I}_t)\}$ are identically distributed when

$$(3.18) \quad \tilde{\pi}_0 = \bar{\pi}_0 = \mu \quad \text{for each } \mu \in \mathcal{P}(\mathbb{R}^d).$$

Thus suppose (3.18) holds for some $\mu \in \mathcal{P}(\mathbb{R}^d)$. Put

$$\tilde{\chi}_t := \mathcal{E} \left(- \sum_{k=1}^{d_1} (\tilde{\pi} h^k) \cdot \tilde{I}^k \right)_t \quad \text{and} \quad \bar{\chi}_t := \mathcal{E} \left(- \sum_{k=1}^{d_1} (\bar{\pi} h^k) \cdot \bar{I}^k \right)_t \quad \forall t \in [0, T]$$

and define the measures \tilde{Q} and \bar{Q} on the measurable spaces $(\tilde{\Omega}, \tilde{\mathcal{F}})$, and $(\bar{\Omega}, \bar{\mathcal{F}})$, respectively, by

$$(3.19) \quad \tilde{Q}(A) := \mathbb{E}^{\tilde{P}}[\tilde{\chi}_T; A] \quad \forall A \in \tilde{\mathcal{F}},$$

$$(3.20) \quad \bar{Q}(A) := \mathbb{E}^{\bar{P}}[\bar{\chi}_T; A] \quad \forall A \in \bar{\mathcal{F}}.$$

Then, with

$$(3.21) \quad \tilde{Y}_t := \tilde{I}_t + \sum_{k=1}^{d_1} \int_0^t \tilde{\pi}_u h^k du, \quad \bar{Y}_t := \bar{I}_t + \sum_{k=1}^{d_1} \int_0^t \bar{\pi}_u h^k du, \quad t \in [0, T]$$

and

$$(3.22) \quad \tilde{\sigma}_t := \tilde{\pi}_t / \tilde{\chi}_t, \quad \bar{\sigma}_t := \bar{\pi}_t / \bar{\chi}_t \quad \forall t \in [0, T],$$

we see, as in Remark 2.10 and Remark 2.13, that the pairs $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t, \tilde{Y}_t)\}$ and $\{(\bar{\Omega}, \bar{\mathcal{F}}, \{\bar{\mathcal{F}}_t\}, \bar{Q}), (\bar{\sigma}_t, \bar{Y}_t)\}$ are weak solutions of the unnormalized filter equation, with

$$\tilde{\sigma}_0 = \bar{\sigma}_0 = \mu.$$

For a complete separable metric space E , let $C_E[0, T]$ denote the complete separable metric space of all continuous mappings from $[0, T]$ into E with the usual metric giving uniform convergence over $[0, T]$. Define

$$\hat{\Omega} := C_{\mathcal{M}^+(\mathbb{R}^d)}[0, T] \times C_{\mathcal{M}^+(\mathbb{R}^d)}[0, T] \times C_{\mathbb{R}^{d_1}}[0, T],$$

which is a complete separable metric space with the usual product metric, and let $\hat{\omega} = (\omega^1, \omega^2, \omega^3)$ be a generic member of $\hat{\Omega}$. By the Yamada–Watanabe construction (see Theorem IV.1.1 of [4]), there exists $\hat{P} \in \mathcal{P}(\hat{\Omega})$ such that:

$$\text{YW.1 } \mathcal{L}_{\hat{P}}(\omega^1, \omega^3) = \mathcal{L}_{\tilde{Q}}(\tilde{\sigma}, \tilde{Y}).$$

$$\text{YW.2 } \mathcal{L}_{\hat{P}}(\omega^2, \omega^3) = \mathcal{L}_{\bar{Q}}(\bar{\sigma}, \bar{Y}).$$

YW.3 If $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{P})$ is the completion of $(\hat{\Omega}, \mathcal{B}(\hat{\Omega}), \hat{P})$, and $\hat{\mathcal{F}}_t$ is the augmentation of the σ -algebra $\sigma\{\hat{\omega}(s), s \in [0, t]\}$ with the null events of $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{P})$, $\forall t \in [0, T]$, then $\{\omega_t^3, t \in [0, T]\}$ is a $\{\hat{\mathcal{F}}_t\}$ -Wiener process on $(\hat{\Omega}, \hat{\mathcal{F}}, \hat{P})$.

From YW.1, YW.2 and YW.3, along with Exercise IV.5.16 of [12], it follows that $\{(\hat{\Omega}, \hat{\mathcal{F}}, \{\hat{\mathcal{F}}_t\}, \hat{P}), (\omega^1, \omega^3)\}$ and $\{(\hat{\Omega}, \hat{\mathcal{F}}, \{\hat{\mathcal{F}}_t\}, \hat{P}), (\omega^2, \omega^3)\}$ are weak solutions for the unnormalized filter equation with

$$\omega_0^1 = \omega_0^2 = \mu,$$

and hence, from Theorem 2.21(i),

$$(3.23) \quad \widehat{P}(\omega_t^1 = \omega_t^2 \quad \forall t \in [0, T]) = 1.$$

From (3.22) we see that

$$(3.24) \quad \tilde{\pi}_t \phi = (\tilde{\sigma}_t \phi)(\tilde{\sigma}_t 1) \quad \forall t \in [0, T], \quad \phi \in B(\mathbb{R}^d),$$

and so, from (3.21),

$$(3.25) \quad \tilde{I}_t^k = \tilde{Y}_t^k - \sum_{k=1}^{d_1} \int_0^t (\tilde{\sigma}_u 1)(\tilde{\sigma}_u h^k) du \quad \forall k = 1, 2, \dots, d_1, \quad t \in [0, T].$$

From (3.24) and (3.25) there exists a measurable mapping $\Phi: C_{\mathcal{M}+(\mathbb{R}^d)}[0, T] \times C_{\mathbb{R}^{d_1}}[0, T] \rightarrow C_{\mathcal{P}(\mathbb{R}^d)}[0, T] \times C_{\mathbb{R}^{d_1}}[0, T]$ such that

$$(3.26) \quad (\tilde{\pi}, \tilde{I}) = \Phi(\tilde{\sigma}, \tilde{Y}).$$

Now (3.24) and (3.25) continue to hold with “overbar” in place of “tilde,” and hence

$$(3.27) \quad (\bar{\pi}, \bar{I}) = \Phi(\bar{\sigma}, \bar{Y}).$$

Thus, for each $\Gamma \in \mathcal{B}(C_{\mathcal{P}(\mathbb{R}^d)}[0, T] \times C_{\mathbb{R}^{d_1}}[0, T])$, we see from (3.19), (3.26) and YW.1 that

$$(3.28) \quad \begin{aligned} \tilde{P}((\tilde{\pi}, \tilde{I}) \in \Gamma) &= \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_T 1)^{-1} I_\Gamma(\Phi(\tilde{\sigma}, \tilde{Y}))] \\ &= \mathbf{E}^{\hat{P}}[(\omega_T^1 1)^{-1} I_\Gamma(\Phi(\omega^1, \omega^3))], \end{aligned}$$

and, from (3.23), (3.27) and YW.2, we similarly have

$$(3.29) \quad \begin{aligned} \bar{P}((\bar{\pi}, \bar{I}) \in \Gamma) &= \mathbf{E}^{\bar{Q}}[(\bar{\sigma}_T 1)^{-1} I_\Gamma(\Phi(\bar{\sigma}, \bar{Y}))] \\ &= \mathbf{E}^{\hat{P}}[(\omega_T^2 1)^{-1} I_\Gamma(\Phi(\omega^2, \omega^3))]. \end{aligned}$$

Now (3.23), (3.28) and (3.29) show that $\tilde{P}((\tilde{\pi}, \tilde{I}) \in \Gamma) = \bar{P}((\bar{\pi}, \bar{I}) \in \Gamma)$, as required.

The Proof of Theorem 2.21(iii) is an obvious simplification of the proof of Theorem 2.21(ii) and is omitted.

PROOF OF THEOREM 2.26. Let $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^i, \tilde{Y}_t)\}$, $i = 1, 2$ be two weak solutions of the unnormalized filter equation. Define

$$\tilde{\sigma}_t^3(\cdot) := \frac{\tilde{\sigma}_t^1(\cdot) + \tilde{\sigma}_t^2(\cdot)}{2}, \quad t \in [0, T].$$

It then follows that $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^3, \tilde{Y}_t)\}$ is a weak solution of the unnormalized filter equation. Therefore, the postulated uniqueness in joint law together with Fact 3.2 implies that for an arbitrary $\phi \in C_c^\infty(\mathbb{R}^d) \cup \{1\}$ we have

$$2\mathbf{E}^{\tilde{Q}}\left[\left(\frac{\tilde{\sigma}_t^1 \phi + \tilde{\sigma}_t^2 \phi}{2}\right)^2\right] - \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^1 \phi)^2] - \mathbf{E}^{\tilde{Q}}[(\tilde{\sigma}_t^2 \phi)^2] = 0 \quad \forall t \in [0, T].$$

Rearranging this expression gives

$$(3.30) \quad \mathbb{E}^{\tilde{Q}} \left[(\tilde{\sigma}_t^1 \phi - \tilde{\sigma}_t^2 \phi)^2 \right] = 0 \quad \forall t \in [0, T].$$

Since $\widehat{C}(\mathbb{R}^d)$ is separable (in the supremum norm), it follows that $C_c^\infty(\mathbb{R}^d)$ is likewise separable, and therefore, by Problem 5.4.25 of [5], there is a countable determining set for $\mathcal{M}^+(\mathbb{R}^d)$ in $C_c^\infty(\mathbb{R}^d) \cup \{1\}$. Then (3.30) shows that $\{\tilde{\sigma}_t^1\}$ and $\{\tilde{\sigma}_t^2\}$ are modifications of each other, hence indistinguishable (since $\{\tilde{\sigma}_t^1\}$ and $\{\tilde{\sigma}_t^2\}$ are continuous). \square

4. Proof of Lemma 3.5. For arbitrary $f_1, f_2 \in C_c^\infty(\mathbb{R}^d)$ put

$$(4.1) \quad \begin{aligned} \tilde{\mathcal{A}}(f_1 \otimes f_2) &:= f_1 \otimes (\mathcal{A}f_2) + (\mathcal{A}f_1) \otimes f_2 \\ &+ \sum_{k=1}^{d_1} \left[(h^k f_1) \otimes (h^k f_2) + (h^k f_1) \otimes (\mathcal{B}_k f_2) \right. \\ &\left. + (\mathcal{B}_k f_1) \otimes (h^k f_2) + (\mathcal{B}_k f_1) \otimes (\mathcal{B}_k f_2) \right], \end{aligned}$$

where \mathcal{A} and \mathcal{B}_k are given by (2.7). We need the following lemma, the proof of which is given in Section 5.

LEMMA 4.1. *Suppose Conditions 2.18–2.20 hold, let $\{(\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}), (\tilde{\sigma}_t^i, \tilde{Y}_t)\}$, $i = 1, 2$, be weak solutions of the unnormalized filter equation, and define the $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued functions $\{v_t^{12}, t \in [0, T]\}$, $\{v_t^{11}, t \in [0, T]\}$, and $\{v_t^{22}, t \in [0, T]\}$, as in (3.4) and (3.6). Then, for each $f_1, f_2 \in C_c^\infty(\mathbb{R}^d)$, we have*

$$(4.2) \quad v_t^{12}(f_1 \otimes f_2) = v_0^{12}(f_1 \otimes f_2) + \int_0^t v_u^{12}(\tilde{\mathcal{A}}(f_1 \otimes f_2)) du \quad \forall t \in [0, T],$$

with identical relations for v_t^{11} and v_t^{22} in place of v_t^{12} .

By direct evaluation of the right-hand side of (4.1) it is easy to establish:

LEMMA 4.2. *For $\tilde{\mathcal{A}}$ and $\bar{\mathcal{A}}$ defined in (4.1) and (4.14), respectively, we have*

$$(4.3) \quad \tilde{\mathcal{A}}(f_1 \otimes f_2)(x) = \bar{\mathcal{A}}(f_1 \otimes f_2)(x) + \bar{h}(x)(f_1 \otimes f_2)(x) \quad \forall x \in \mathbb{R}^{2d},$$

for each $f_1, f_2 \in C_c^\infty(\mathbb{R}^d)$.

Define

$$(4.4) \quad \tilde{\mathcal{G}} := \text{span}\{f_1 \otimes f_2 : f_1, f_2 \in C_c^\infty(\mathbb{R}^d)\}.$$

Putting Lemmas 4.2 and 4.1 together, we see that the mappings $\{v_t^{12}, t \in [0, T]\}$, $\{v_t^{11}, t \in [0, T]\}$ and $\{v_t^{22}, t \in [0, T]\}$, defined at (3.4) and (3.6), are $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solutions of the evolution equation for $(\bar{\mathcal{A}} + \bar{h}, \tilde{\mathcal{G}})$; that is,

$$(4.5) \quad v_t^{12} f = v_0^{12} f + \int_0^t v_u^{12}(\bar{\mathcal{A}}f + \bar{h}f) du \quad \forall t \in [0, T], \forall f \in \tilde{\mathcal{G}},$$

with identical relations for ν_t^{11} and ν_t^{22} in place of ν_t^{12} . In order to prove Lemma 3.5, it remains to show that (4.5) holds not only for $f \in \tilde{\mathcal{D}}$, but for each f in the larger domain $\text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\}$. That is, it must be shown that the mappings $\{\nu_t^{12}, t \in [0, T]\}$, $\{\nu_t^{11}, t \in [0, T]\}$ and $\{\nu_t^{22}, t \in [0, T]\}$ are $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solutions of the evolution equation for $(\mathcal{A} + \bar{h}, \text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\})$. To this end we need the following result, whose proof is deferred to Section 5:

LEMMA 4.3. *Suppose Conditions 2.18–2.20 hold. Then the closure of the relation $\{(f, \mathcal{A}f): f \in \tilde{\mathcal{D}}\}$ in the supremum norm of $B(\mathbb{R}^{2d}) \times B(\mathbb{R}^{2d})$ contains the relation $\{(f, \mathcal{A}f): f \in C_c^\infty(\mathbb{R}^{2d})\}$.*

From Lemma 4.3 and the notions of bp-closedness and bp-closure of a relation (see foot of page 111 of [3]), we see that

$$(4.6) \quad \{(f, \mathcal{A}f): f \in C_c^\infty(\mathbb{R}^{2d})\} \subset \text{bp-closure} \{(f, \mathcal{A}f): f \in \tilde{\mathcal{D}}\}.$$

Now put

$$(4.7) \quad S^{12} := \left\{ (f, g) \in B(\mathbb{R}^{2d}) \times B(\mathbb{R}^{2d}): \nu_t^{12} f \right. \\ \left. = \nu_0^{12} f + \int_0^t \nu_s^{12} (g + \bar{h}f) ds \forall t \in [0, T] \right\},$$

and observe that S^{12} is a linear relation. By (3.5) we have

$$\sup_{0 \leq t \leq T} \nu_t^{12}(\mathbb{R}^{2d}) < \infty,$$

and therefore, since $\bar{h} \in B(\mathbb{R}^{2d})$, it follows from the dominated convergence theorem that the linear relation S^{12} is bp-closed in $B(\mathbb{R}^{2d}) \times B(\mathbb{R}^{2d})$. Since the $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued mapping $\{\nu_t^{12}, t \in [0, T]\}$ solves the evolution equation for $(\mathcal{A} + \bar{h}, \tilde{\mathcal{D}})$, we have $\{(f, \mathcal{A}f): f \in \tilde{\mathcal{D}}\} \subset S^{12}$, and therefore, from the bp-closedness of S^{12} and (4.6), we have

$$(4.8) \quad \{(f, \mathcal{A}f): f \in C_c^\infty(\mathbb{R}^{2d})\} \subset S^{12}.$$

Next, observe from (3.15) and Problem 4.11.12 of [3] that the operator $(\mathcal{A}, C_c^\infty(\mathbb{R}^{2d}))$ is conservative, and hence (see page 166 of [3]) we have

$$(4.9) \quad (1, 0) \in \text{bp-closure} \{(f, \mathcal{A}f): f \in C_c^\infty(\mathbb{R}^{2d})\}.$$

In light of (4.8), (4.9), and the bp-closedness of S^{12} , we then get

$$(4.10) \quad (1, 0) \in S^{12}.$$

Now (4.8), (4.10), and linearity of the relation S^{12} shows that

$$\{(f, \mathcal{A}f): f \in \text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\}\} \subset S^{12},$$

which, in view of (4.1) shows that $\{\nu_t^{12}, t \in [0, T]\}$ is an $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solution of the evolution equation for $(\mathcal{A} + \bar{h}, \text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\})$. Defining S^{11} as in (4.7), but with ν_t^{11} in place of ν_t^{12} , we can similarly show that

$\{\nu_t^{11}, t \in [0, T]\}$ is an $\mathcal{M}^+(\mathbb{R}^{2d})$ -valued solution of the evolution equation for $(\mathcal{A} + \bar{h}, \text{span}\{1, C_c^\infty(\mathbb{R}^{2d})\})$, and likewise for $\{\nu_t^{22}, t \in [0, T]\}$. \square

5. Proofs of technical results.

PROOF OF FACT 3.2. Fix some $\alpha \in (1, \infty)$. Since $\tilde{\sigma}_0$ takes values in $\mathcal{P}(\mathbb{R}^d)$, we see from (2.16) with $\phi \equiv 1$ that

$$\tilde{\sigma}_t \mathbf{1} = 1 + \int_0^t \sum_{k=1}^{d_1} \left(\frac{\tilde{\sigma}_s h^k}{\tilde{\sigma}_s \mathbf{1}} \right) (\tilde{\sigma}_s \mathbf{1}) d\tilde{Y}_s^k.$$

This gives (see Exercise IV.3.10(1) of [12])

$$(5.1) \quad \tilde{\sigma}_t \mathbf{1} = \mathcal{E} \left(\sum_{k=1}^{d_1} \left(\frac{\tilde{\sigma} h^k}{\tilde{\sigma} \mathbf{1}} \cdot \tilde{Y}^k \right) \right)_t,$$

and hence

$$(5.2) \quad \begin{aligned} |\tilde{\sigma}_t \mathbf{1}|^\alpha &= \tilde{M}_t \exp \left(\frac{\alpha(\alpha-1)}{2} \sum_{k=1}^{d_1} \int_0^t \left| \frac{\tilde{\sigma}_s h^k}{\tilde{\sigma}_s \mathbf{1}} \right|^2 ds \right) \\ &\leq \tilde{M}_t \exp \left(\frac{\alpha(\alpha-1)}{2} \|h\| T \right) \quad \forall t \in [0, T], \end{aligned}$$

for

$$\tilde{M}_t := \mathcal{E} \left(\alpha \sum_{k=1}^{d_1} \left(\frac{\tilde{\sigma} h^k}{\tilde{\sigma} \mathbf{1}} \cdot \tilde{Y}^k \right) \right)_t \quad \text{and} \quad \|h\| := \sup_{x \in \mathbb{R}^d} |h(x)|.$$

Now Condition 2.20 ensures that the processes $\{(\tilde{\sigma}_t h^k)/(\tilde{\sigma}_t \mathbf{1}), t \in [0, T]\}$ are uniformly bounded (by $\|h^k\|$), and therefore $\{(\tilde{M}_t, \tilde{\mathcal{F}}_t), t \in [0, T]\}$ is a continuous martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{Q})$, with $\tilde{M}_0 = 1$. Taking \tilde{Q} -expectations in (5.2) then gives

$$(5.3) \quad \mathbb{E}^{\tilde{Q}}[|\tilde{\sigma}_t \mathbf{1}|^\alpha] \leq \exp \left(\frac{\alpha(\alpha-1)}{2} \|h\| T \right) \quad \forall t \in [0, T].$$

Again, by (5.1) and uniform-boundedness of the processes $\{(\tilde{\sigma}_t h^k)/(\tilde{\sigma}_t \mathbf{1}), t \in [0, T]\}$ we see that $\{(\tilde{\sigma}_t \mathbf{1}, \tilde{\mathcal{F}}_t), t \in [0, T]\}$ is a continuous martingale on $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{Q})$, which, in light of (5.3), is L_α -bounded. Thus, by Doob's inequality, there is some $\gamma(\alpha) \in (0, \infty)$ such that (3.2) holds. \square

PROOF OF LEMMA 4.1. Fix $f_1, f_2 \in C_c^\infty(\mathbb{R}^d)$. Since $\{\tilde{\Omega}, \tilde{\mathcal{F}}, \{\tilde{\mathcal{F}}_t\}, \tilde{Q}\}$, $(\tilde{\sigma}_t^i, \tilde{Y}_t^i)$, $i = 1, 2$ are weak solutions of the unnormalized filter equations,

we have

$$(5.4) \quad \begin{aligned} \tilde{\sigma}_t^i f_i &= \tilde{\sigma}_0^i f_i + \int_0^t \tilde{\sigma}_u^i(\mathcal{A} f_i) du \\ &+ \sum_k \int_0^t \tilde{\sigma}_u^i(h^k f_i + \mathcal{B}_k f_i) d\tilde{Y}_u^k, \quad i = 1, 2. \end{aligned}$$

Expand the product of semimartingales $\{(\tilde{\sigma}_t^1 f_1)(\tilde{\sigma}_t^2 f_2)\}$ by Itô's formula, and note that each stochastic integral in this expansion is with respect to a \tilde{Q} -standard Wiener process $\{\tilde{Y}_t^k, \tilde{\mathcal{F}}_t\}$, $k = 1, 2, \dots, d_1$. Fact 3.2 ensures that these stochastic integrals are genuine \tilde{Q} -martingales, and therefore have \tilde{Q} -expectation identically zero. Upon taking \tilde{Q} -expectations on each side of the resulting expansion of $\{(\tilde{\sigma}_t^1 f_1)(\tilde{\sigma}_t^2 f_2)\}$ and using (4.3) and (3.7), we get (4.2). The corresponding identities for ν^{11} and ν^{22} are similarly obtained. \square

PROOF OF LEMMA 4.3. Fix arbitrary $\varepsilon \in (0, \infty)$ and $g \in C_c^\infty(\mathbb{R}^{2d})$. Put

$$B_R := \{x \in \mathbb{R}^{2d} : |x| \leq R\}, \quad R \in [0, \infty),$$

and fix R such that $\text{supp}(g) \subset B_R$. Also fix some $q \in C_c^\infty(\mathbb{R}^d)$ such that

$$(5.5a) \quad \|q\| \leq 1;$$

$$(5.5b) \quad q(z) = 1 \quad \forall z \in \mathbb{R}^d \quad \text{with } |z| \leq R;$$

$$(5.5c) \quad q(z) = 0 \quad \forall z \in \mathbb{R}^d \quad \text{with } |z| \geq R\sqrt{2}.$$

By Proposition 7.1 in Appendix 7 of [3], there exists a polynomial $p: \mathbb{R}^{2d} \rightarrow \mathbb{R}$ such that

$$(5.6a) \quad \max_{x \in B_{2R}} |g(x) - p(x)| < \varepsilon;$$

$$(5.6b) \quad \max_{x \in B_{2R}} |\partial_i g(x) - \partial_i p(x)| < \varepsilon \quad \forall i = 1, \dots, 2d;$$

$$(5.6c) \quad \max_{x \in B_{2R}} |\partial_i \partial_j g(x) - \partial_i \partial_j p(x)| < \varepsilon \quad \forall i, j = 1, \dots, 2d.$$

Since $g(x) = 0$ when $x \notin B_R$, we note from (5.6a) that

$$(5.7) \quad \sup_{x \in B_{2R} \setminus B_R} |p(x)| < \varepsilon.$$

For all $x \in \mathbb{R}^{2d}$, put $x := (x_1, x_2)$, $x_1, x_2 \in \mathbb{R}^d$, and define

$$\bar{q}(x) := q(x_1)q(x_2),$$

$$f(x) := \bar{q}(x) p(x).$$

Since $q \in C_c^\infty(\mathbb{R}^d)$ and $p(x)$ is a polynomial in $x = (x_1, x_2)$, it follows that $f \in \tilde{\mathcal{S}}$ [recall (4.4)]. From (5.5), we have $\bar{q}(x) = 0$ when $x \notin B_{2R}$ and $\bar{q}(x) = 1$

when $x \in B_R$. Thus

$$\begin{aligned} \|f - g\| &= \sup_{x \in \mathbb{R}^{2d}} |\bar{q}(x)p(x) - g(x)| = \max_{x \in B_{2R}} |\bar{q}(x)p(x) - g(x)| \\ &\leq \max_{x \in B_R} |p(x) - g(x)| + \sup_{x \in B_{2R} \setminus B_R} |\bar{q}(x)p(x) - g(x)|, \end{aligned}$$

hence (5.6) and (5.7) give

$$(5.8) \quad \|f - g\| \leq \varepsilon + \sup_{x \in B_{2R} \setminus B_R} |p(x)| \leq 2\varepsilon.$$

Next, consider $\|\mathcal{A}f - \mathcal{A}g\|$. From (3.14) we have

$$(5.9) \quad \mathcal{A}f(x) = \bar{q}(x)\mathcal{A}p(x) + p(x)\mathcal{A}\bar{q}(x) + (\nabla p(x))^T \bar{a}(x) \nabla \bar{q}(x).$$

By the choice of R we have $g(x) = 0$ and therefore $\mathcal{A}g(x) = 0$, $\forall x \notin B_R$. Moreover, from (5.5c), we have $\bar{q}(x) = 1$, and therefore $\nabla \bar{q}(x) = 0$ and $\mathcal{A}\bar{q}(x) = 0$, $\forall x \in B_R$. Similarly, $\bar{q}(x) = 0$, and therefore $\mathcal{A}\bar{q}(x) = 0$, $\forall x \notin B_{2R}$. Then, it follows from (5.75) that

$$\begin{aligned} \|\mathcal{A}f - \mathcal{A}g\| &= \sup_{x \in B_{2R}} |\bar{q}(x)\mathcal{A}p(x) + p(x)\mathcal{A}\bar{q}(x) \\ &\quad + (\nabla p(x))^T \bar{a}(x) \nabla \bar{q}(x) - \mathcal{A}g(x)| \\ &\leq \sup_{x \in B_R} |\bar{q}(x)\mathcal{A}p(x) + p(x)\mathcal{A}\bar{q}(x) \\ (5.10) \quad &\quad + (\nabla p(x))^T \bar{a}(x) \nabla \bar{q}(x) - \mathcal{A}g(x)| \\ &\quad + \sup_{x \in B_{2R} \setminus B_R} |\bar{q}(x)\mathcal{A}p(x) + p(x)\mathcal{A}\bar{q}(x) + (\nabla p(x))^T \bar{a}(x) \nabla \bar{q}(x)| \\ &= \sup_{x \in B_R} |\mathcal{A}p(x) - \mathcal{A}g(x)| + \sup_{x \in B_{2R} \setminus B_R} (|\bar{q}(x)\mathcal{A}p(x)| \\ &\quad + |p(x)\mathcal{A}\bar{q}(x)| + |(\nabla p(x))^T \bar{a}(x) \nabla \bar{q}(x)|). \end{aligned}$$

Since \bar{a} and \bar{b} are locally bounded, we have

$$C_1 := \sup_{x \in B_{2R}} \left(\sum_{i=1}^{2d} |\bar{b}_i(x)| + \frac{1}{2} \sum_{i,j=1}^{2d} |\bar{a}_{ij}(x)| \right) < \infty.$$

Also let

$$C_2 := \|\bar{q}\| + \sum_i \|\partial_i \bar{q}\| + \sum_{i,j} \|\partial_i \partial_j \bar{q}\| < \infty.$$

Then by (5.6),

$$(5.11) \quad \sup_{x \in B_R} |\mathcal{A}p(x) - \mathcal{A}g(x)| < C_1 \varepsilon.$$

Similarly, by (5.6) and the fact that $g(x) = 0$, $\forall x \in B_{2R} \setminus B_R$, we obtain

$$\left. \begin{aligned} |\partial_i p(x)| &< \varepsilon, \\ |\mathcal{A}p(x)| &< C_1 \varepsilon, \end{aligned} \right\} \quad \forall x \in B_{2R} \setminus B_R,$$

and hence, from these bounds and (5.7),

$$(5.12) \quad \sup_{x \in B_{2R} \setminus B_R} (|\bar{q}(x) \cdot \bar{\mathcal{A}}p(x)| + |p(x) \cdot \bar{\mathcal{A}}\bar{q}(x)| + |(\nabla p(x))^T \bar{a}(x) \nabla \bar{q}(x)|) \\ \leq \varepsilon C_1 + \varepsilon C_1 C_2 + \varepsilon C_1 C_2.$$

Now, upon combining (5.10), (5.11) and (5.12) we have

$$(5.13) \quad \|\bar{\mathcal{A}}g - \bar{\mathcal{A}}f\| \leq 2\varepsilon(C_1 + C_1 C_2),$$

and the result follows. \square

6. Proof of Theorem 3.6.

REMARK 6.1. In view of Remark 1 on page 345 of [2], with no loss of generality we suppose that $\lambda \in B(\mathbb{R}^q)$ is a nonnegative mapping, and that the constant K in (3.17) is such that $0 \leq \lambda(x) \leq K$, $\forall x \in \mathbb{R}^q$.

We first show that the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$ has uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions over the interval $[0, \infty)$. Thus, let $\{\mu_t, t \in [0, \infty)\}$ be such a solution, hence

$$(6.1) \quad \mu_t f = \mu_0 f + \int_0^t \mu_s (\mathcal{L}f - \lambda f) ds \quad \forall t \in [0, \infty) \forall f \in \mathcal{D}(\mathcal{L}).$$

Since $(1, 0) \in \mathcal{L}$, it follows from (6.1) that

$$\mu_t(\mathbb{R}^q) = \mu_0(\mathbb{R}^q) - \int_0^t \mu_s \lambda ds \quad \forall t \in [0, \infty)$$

and therefore $\mu_t(\mathbb{R}^q) \in [0, 1]$, $\forall t \in [0, \infty)$ [since μ_0 is a probability measure and, by Remark 6.1, $\lambda(\cdot)$ is nonnegative]; that is, each μ_t is a subprobability measure on $\mathcal{B}(\mathbb{R}^q)$.

REMARK 6.2. We are going to use Theorem 6.4 (which follows) to establish uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions of the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$. However, Theorem 6.4 provides uniqueness in the class of *probability measure-valued* solutions of an evolution equation (recall Remark 3.1), and we have seen that, for an $\mathcal{M}^+(E)$ -valued solution $\{\mu_t, t \in [0, \infty)\}$ of the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$, the μ_t are only *subprobability measures* on \mathbb{R}^q . We shall therefore use an idea of Bhatt and Karandikar (see page 344 of [2]), and add a “point at infinity” to get a one-point compactification \mathbb{R}^{q*} of \mathbb{R}^q . Then the subprobability measures μ_t on \mathbb{R}^q are extended to *probability measures* μ_t^* on \mathbb{R}^{q*} [see (6.2)], and it will be seen that the resulting function $\{\mu_t^*, t \in [0, \infty)\}$ is a $\mathcal{P}(\mathbb{R}^{q*})$ -valued solution of the evolution equation for an operator $\mathcal{L}^* \subset B(\mathbb{R}^{q*}) \times B(\mathbb{R}^{q*})$ which is an “extension” of the operator $(\mathcal{L} - \lambda) \subset B(\mathbb{R}^q) \times B(\mathbb{R}^q)$ (see (6.4)). Theorem 6.4 will then be used to establish uniqueness in the class of $\mathcal{P}(\mathbb{R}^{q*})$ -valued solutions of the evolution equation for \mathcal{L}^* , and this in turn will yield uniqueness in the class of the $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions of the evolution equation (6.1), as required.

We use Δ to denote the point at infinity in the one-point compactification \mathbb{R}^{q*} of \mathbb{R}^q . Also, members of $B(\mathbb{R}^{q*})$ and operators on $B(\mathbb{R}^{q*})$ will be superscripted with “*”, and, for $f^* \in B(\mathbb{R}^{q*})$, we write $f^*|_{\mathbb{R}^q}$ to denote the restriction of f^* to the domain \mathbb{R}^q .

Using the postulated $\mathcal{M}^+(\mathbb{R}^q)$ -valued solution $\{\mu_t, t \in [0, \infty)\}$ of the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$, define $\mu_t^* \in \mathcal{P}(\mathbb{R}^{q*}), \forall t \in [0, \infty)$, as follows:

$$(6.2) \quad \mu_t^*(\Gamma) := \mu_t(\Gamma \cap \mathbb{R}^q) + (1 - \mu_t(\mathbb{R}^q))I_\Gamma(\Delta) \quad \forall \Gamma \in \mathcal{B}(\mathbb{R}^{q*}).$$

Next, define the linear operator $\mathcal{L}^*: \mathcal{D}(\mathcal{L}^*) \subset C(\mathbb{R}^{q*}) \rightarrow B(\mathbb{R}^{q*})$ by

$$(6.3a) \quad \mathcal{D}(\mathcal{L}^*) := \{f^* \in C(\mathbb{R}^{q*}): f^*|_{\mathbb{R}^q} - f^*(\Delta) \in C_c^\infty(\mathbb{R}^q)\},$$

$$(6.3b) \quad \mathcal{L}^*f^*(x) := \mathcal{L}(f^*|_{\mathbb{R}^q} - f^*(\Delta))(x) \quad \forall x \in \mathbb{R}^q, \quad \forall f^* \in \mathcal{D}(\mathcal{L}^*),$$

$$(6.3c) \quad \mathcal{L}^*f^*(\Delta) := 0 \quad \forall f^* \in \mathcal{D}(\mathcal{L}^*).$$

Finally, define linear operator $\mathcal{S}^*: \mathcal{D}(\mathcal{S}^*) \subset C(\mathbb{R}^{q*}) \rightarrow B(\mathbb{R}^{q*})$ by

$$(6.4) \quad \begin{aligned} \mathcal{S}^*f^*(x) &:= \mathcal{L}^*f^*(x) - \lambda(x)(f^*(x) - f^*(\Delta)) \\ &\forall x \in \mathbb{R}^{q*} \quad \forall f^* \in \mathcal{D}(\mathcal{S}^*) := \mathcal{D}(\mathcal{L}^*). \end{aligned}$$

From (6.2) and (6.1) one easily checks that $\{\mu_t^*, t \in [0, \infty)\}$ solves the evolution equation for $(\mathcal{S}^*, \mathcal{D}(\mathcal{S}^*))$, namely

$$(6.5) \quad \mu_t^*f^* = \mu_0^*f^* + \int_0^t \mu_s^*(\mathcal{S}^*f^*) ds \quad \forall t \in [0, \infty), \quad \forall f^* \in \mathcal{D}(\mathcal{S}^*),$$

REMARK 6.3. We clearly have the following: if the evolution equation for $(\mathcal{S}^*, \mathcal{D}(\mathcal{S}^*))$ has uniqueness in the class of $\mathcal{P}(\mathbb{R}^{q*})$ -valued solutions over $[0, \infty)$, then the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$ must have uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions over $[0, \infty)$. It therefore remains to establish the former type of uniqueness, and for this purpose we shall use the following special case of Theorem 2.7(c) from [9].

THEOREM 6.4. *Suppose that F is a compact metric space, $\mathbb{A}^0: \mathcal{D}(\mathbb{A}^0) \subset \overline{C}(\mathbb{R}^{q*}) \rightarrow \overline{C}(\mathbb{R}^{q*} \times F)$ is a linear operator, and η is a transition function from \mathbb{R}^{q*} to F . Define*

$$(6.6) \quad \begin{aligned} \mathbb{A}_\eta^0 f^*(x) &:= \int_F \mathbb{A}^0 f^*(x, y) \eta(x, dy) \quad \forall x \in \mathbb{R}^{q*}, \\ &\forall f^* \in \mathcal{D}(\mathbb{A}_\eta^0) := \mathcal{D}(\mathbb{A}^0). \end{aligned}$$

Suppose also that (i) $\mathcal{D}(\mathbb{A}^0)$ is closed under multiplication and separates points, (ii) $\mathbb{A}_y^0 f^ \equiv \mathbb{A}^0 f^*(\cdot, y)$ is a pregenerator for each $y \in F$, and (iii) \mathbb{A}_η^0 satisfies the following separability hypothesis: there exists some countable $\{g_k^*\} \subset \mathcal{D}(\mathbb{A}_\eta^0)$ such that the graph of \mathbb{A}_η^0 is included within the bp-closure of the linear span of $\{(g_k^*, \mathbb{A}_\eta^0 g_k^*)\}$. With these conditions we have the following: if uniqueness holds*

for the martingale problem for \mathbb{A}_η^0 then the evolution equation for $(\mathbb{A}_\eta^0, \mathcal{D}(\mathbb{A}_\eta^0))$ has uniqueness in the class of $\mathcal{P}(\mathbb{R}^{q*})$ -valued solutions over the interval $[0, \infty)$.

Define the compact metric space

$$F := \{y = (y_1, y_2, y_3) \in \mathbb{S}_+^{q \times q} \times \mathbb{R}^q \times \mathbb{R} : |y_1^{ij}| \leq K, |y_2^i| \leq K, 0 \leq y_3 \leq K\},$$

and, motivated by Example 3.4 of [11], for each $y \in F$ define the linear operator \mathcal{L}_y on $\overline{C}(\mathbb{R}^q)$ by

$$(6.7) \quad \begin{aligned} \mathcal{L}_y f(x) &:= \sum_i (1 + |x|) y_2^i \partial_i f(x) + \frac{1}{2} \sum_{i,j} (1 + |x|^2) y_1^{ij} \partial_i \partial_j f(x) \\ &\quad \forall x \in \mathbb{R}^q, f \in \mathcal{D}(\mathcal{L}_y) := C_c^\infty(\mathbb{R}^q). \end{aligned}$$

Also, put $\mathcal{D}(\mathbb{A}^0) := \mathcal{D}(\mathcal{L}^*)$ (see [see (6.3)] and

$$(6.8a) \quad \begin{aligned} \mathbb{A}^0 f^*(x, y) &:= \mathcal{L}_y(f^*|_{\mathbb{R}^q} - f^*(\Delta))(x) - y_3(f^*(x) - f^*(\Delta)) \\ &\quad \forall f^* \in \mathcal{D}(\mathbb{A}^0), \quad \forall (x, y) \in \mathbb{R}^q \times F, \end{aligned}$$

$$(6.8b) \quad \mathbb{A}^0 f^*(\Delta, y) := 0 \quad \forall f^* \in \mathcal{D}(\mathbb{A}^0), \quad \forall y \in F.$$

This defines a linear mapping $\mathbb{A}^0: \mathcal{D}(\mathbb{A}^0) \subset \overline{C}(\mathbb{R}^{q*}) \rightarrow \overline{C}(\mathbb{R}^{q*} \times F)$. Next, fix an arbitrary $\bar{y} \in F$, and define a transition function η from \mathbb{R}^q to F by

$$(6.9a) \quad \eta(x, \Gamma) := \delta_{(a(x)/1+|x|^2, b(x)/1+|x|, \lambda(x))}(\Gamma), \quad x \in \mathbb{R}^q, \Gamma \in \mathcal{B}(F),$$

$$(6.9b) \quad \eta(\Delta, \Gamma) := \delta_{\bar{y}}(\Gamma), \quad \Gamma \in \mathcal{B}(F).$$

Putting together (6.4), (6.7), (6.8) and (6.9), we get

$$(6.10) \quad \mathcal{G}^* \equiv \mathbb{A}_\eta^0,$$

[where \mathbb{A}_η^0 is given by (6.6) in terms of the operator \mathbb{A}^0 in (6.8) and the transition function η in (6.9)]. We next check the conditions of Theorem 6.4 for \mathbb{A}^0 and η given by (6.8) and (6.9): (i) From (6.3) it follows that $\mathcal{D}(\mathbb{A}^0) := \mathcal{D}(\mathcal{L}^*)$ is closed under multiplication and separates points. (ii) Fix $y \in F$ and $v^* \in \mathcal{P}(\mathbb{R}^{q*})$, and define the linear operator \mathcal{L}_y^* on $C(\mathbb{R}^{q*})$ by $\mathcal{D}(\mathcal{L}_y^*) := \mathcal{D}(\mathcal{L}^*)$ and

$$\mathcal{L}_y^* f^*(x) := \mathcal{L}_y(f^*|_{\mathbb{R}^q} - f^*(\Delta))(x) \quad \forall x \in \mathbb{R}^q, \quad \mathcal{L}_y^* f^*(\Delta) := 0 \quad \forall f^* \in \mathcal{D}(\mathcal{L}_y^*).$$

Clearly \mathcal{L}_y given by (6.7) satisfies the positive maximum principle, thus Theorem 4.5.4 of and [3] gives existence of a solution of the $D_{\mathbb{R}^{q*}}[0, \infty)$ -martingale problem for (\mathcal{L}_y^*, v^*) .

Then, for the linear operator \mathbb{A}_y^0 on $C(\mathbb{R}^{q*})$ given by $\mathcal{D}(\mathbb{A}_y^0) := \mathcal{D}(\mathcal{L}_y^*)$ and

$$\mathbb{A}_y^0 f^*(x) := \mathcal{L}_y^* f^*(x) - y_3(f^*(x) - f^*(\Delta)) \quad \forall x \in \mathbb{R}^{q*}, \quad \forall f^* \in \mathcal{D}(\mathbb{A}_y^0),$$

it follows from Theorem 4.10.2 of [3] that there exists a solution of the $D_{\mathbb{R}^{q*}}[0, \infty)$ -martingale problem for $(\mathbb{A}_y^0, \delta_x)$, $\forall x \in \mathbb{R}^{q*}$, and therefore \mathbb{A}_y^0 is a pre-generator (see remark at foot of page 4 in [9]). (iii) From Remark 2.5 of [9] there is a sequence $\{g_k\} \subset C_c^\infty(\mathbb{R}^q)$ such that the graph of \mathcal{L} is

included within the bp-closure of the linear span of $\{(g_k, \mathcal{L}g_k)\}$. Now define $g_{k,q}^*(x) := g_k(x) + q, \forall x \in \mathbb{R}^q, g_{k,q}^*(\Delta) := q, k = 1, 2, \dots, q$ rational. From (6.4) it follows that the countable set $\{g_{k,q}^*\}$ is a subset of $\mathcal{D}(\mathcal{G}^*)$ and the graph of \mathcal{G}^* is contained in the bp-closure of the linear span of $\{(g_{k,q}^*, \mathcal{G}^*g_{k,q}^*)\}$. In view of (6.10) we have thus verified condition (iii) of Theorem 6.4. Finally, note that uniqueness holds for the martingale problem for \mathcal{G}^* . Indeed, observe from Theorem 8.1.7 of [3] that the martingale problem for \mathcal{L} is well posed, from which it easily follows that the $D_{\mathbb{R}^{q^*}}[0, \infty)$ -martingale problem for \mathcal{L}^* is well posed, and hence Theorem 4.10.2 of [3] shows that the $D_{\mathbb{R}^{q^*}}[0, \infty)$ -martingale problem for \mathcal{G}^* is well posed. Now it easily follows from Theorem 4.3.6 of [3] that uniqueness holds for the martingale problem for \mathcal{G}^* . We thus conclude from Theorem 6.4 and (6.10) that the evolution equation for $(\mathcal{G}^*, \mathcal{D}(\mathcal{G}^*))$ has uniqueness in the class of $\mathcal{P}(\mathbb{R}^{q^*})$ -valued solutions over $[0, \infty)$, and hence (see Remark 6.3) the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$ has uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions over $[0, \infty)$.

It finally remains to establish that the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$ has uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions over the finite interval $[0, T]$. Let $\{\mu_t, t \in [0, T]\}$ be such a solution. Extend μ from $[0, T]$ to $[0, \infty)$ by defining

$$\mu_t(\Gamma) := \int_{\mathbb{R}^q} E_x \left[I_\Gamma(\omega_t) \exp \left(\int_0^{t-T} -\lambda(\omega_s) ds \right) \right] \mu_T(dx)$$

$$\forall t \in [T, \infty), \quad \forall \Gamma \in \mathcal{B}(\mathbb{R}^q),$$

where $P_x \in \mathcal{P}(C_{\mathbb{R}^q}[0, \infty))$, $x \in \mathbb{R}^q$, is the probability measure on the space $C_{\mathbb{R}^q}[0, \infty)$ (of continuous functions from $[0, \infty)$ into \mathbb{R}^q) which solves the martingale problem for (\mathcal{L}, δ_x) , and ω denotes a generic element of $C_{\mathbb{R}^q}[0, \infty)$ (existence and uniqueness of P_x follow from Theorem 8.1.7 of [3]). It is easily checked that the mapping $\{\mu_t, t \in [0, \infty)\}$ is an $\mathcal{M}^+(\mathbb{R}^q)$ -valued solution of the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$, and therefore the evolution equation for $(\mathcal{L} - \lambda, \mathcal{D}(\mathcal{L}))$ must have uniqueness in the class of $\mathcal{M}^+(\mathbb{R}^q)$ -valued solutions over the finite interval $[0, T]$. \square

Acknowledgment. The authors thank Professor T. G. Kurtz for valuable comments on this work.

REFERENCES

- [1] BHATT, A. G., KALLIANPUR, G. and KARANDIKAR, R. L. (1995). Uniqueness and robustness of solution of measure-valued equations of nonlinear filtering. *Ann. Probab.* **23** 1895–1938.
- [2] BHATT, A. G. and KARANDIKAR, R. L. (1995). Evolution equations for Markov processes: application to the white-noise theory of filtering. *Appl. Math. Optim.* **31** 327–348.
- [3] ETHIER, S. N. and KURTZ, T. G. (1986). *Markov Processes: Characterization and Convergence*. Wiley, New York.
- [4] IKEDA, N. and WATANABE, S. (1989). *Stochastic Differential Equations and Diffusion Processes*. North-Holland, Amsterdam.

- [5] KARATZAS, I. and SHREVE, S. E. (1991). *Brownian Motion and Stochastic Calculus*. Springer, New York.
- [6] KUNITA, H. (1971). Asymptotic behavior of the nonlinear filtering errors of Markov processes. *J. Multivariate Anal.* **1** 365–393.
- [7] KUNITA, H. (1981). Stochastic partial differential equations connected with nonlinear filtering. *Lecture Notes in Math.* **972** 100–169. Springer, Berlin.
- [8] KUNITA, H. (1990). *Stochastic Flows and Stochastic Differential Equations*. Cambridge Univ. Press.
- [9] KURTZ, T. G. (1998). Martingale problems for conditional distributions of Markov processes. *Electron J. Probab.* **3** 1–29.
- [10] KURTZ, T. G. and OCONE, D. L. (1988). Unique characterization of conditional distributions in nonlinear filtering. *Ann. Probab.* **16** 80–107.
- [11] KURTZ, T. G. and STOCKBRIDGE, R. H. (1988). Existence of Markov controls and characterization of optimal Markov controls. *SIAM J. Control Optim.* **36** 609–653.
- [12] REVUZ, D. and YOR, M. (1994). *Continuous Martingales and Brownian Motion*. Springer, New York.
- [13] ROGERS, L. C. G. and WILLIAMS, D. (1987). *Diffusions, Markov Processes and Martingales 2: Itô Calculus*. Wiley, New York.
- [14] ROZOVSKII, B. L. (1991). A simple proof of uniqueness for Kushner and Zakai equations. In *Stochastic Analysis* (E. Mayer-Wolf, E. Merzbach and A. Schwartz, eds.) 449–458. Academic Press, New York.
- [15] SZPIRGLAS, J. (1998). Sur l'équivalence d'équations différentielles stochastiques à valeurs mesures intervenant dans le filtrage markovien non linéaire. *Ann. Inst. H. Poincaré Probab. Statist.* **14** 33–59.
- [16] YAMADA, T. and WATANABE, S. (1971). On the uniqueness of solutions of stochastic differential equations. *J. Math. Kyoto Univ.* **11** 155–167.

DEPARTMENT OF ELECTRICAL
AND COMPUTER ENGINEERING
UNIVERSITY OF WATERLOO
WATERLOO, ONTARIO N2L 3G1
CANADA
E-MAIL: lucic@kingcong.uwaterloo.ca
heunis@kingcong.uwaterloo.ca