

## The Usual Model Construction for NFU Preserves Information

M. Randall Holmes

**Abstract** The usual construction of models of NFU (New Foundations with urelements, introduced by Jensen) is due to Maurice Boffa. A Boffa model is obtained from a model of (a fragment of) Zermelo–Fraenkel with Choice (ZFC) with an automorphism which moves a rank: the domain of the Boffa model is a rank that is moved. “Most” elements of the domain of the Boffa model are urelements in terms of the interpreted NFU. The main result of this paper is that the restriction of the membership relation of the original model of set theory with automorphism to the domain of the Boffa model is first-order definable in the language of NFU. In particular, all information about the extensions in the original model of the urelements of the model of NFU is definable in terms of NFU. A corollary (answering a question of Thomas Forster) is that the urelements in a Boffa model are not homogeneous.

### 1 Introduction

This paper was originally motivated by a question of Thomas Forster about the urelements in a model of Jensen’s modification New Foundations with urelements (NFU) of Quine’s New Foundations (NF): Is it possible for them to be homogeneous? We answer this question in the negative for a familiar class of models (those obtained from a nonstandard model of a fragment of set theory with an automorphism in a way originally proposed by Maurice Boffa) by showing that the membership relation of the underlying model is first-order definable in terms of the interpreted NFU, which is not at all obvious and perhaps surprising.

Forster asked whether there are models of NFU in which the urelements are homogeneous. By this we mean that for any formula  $\varphi(a_1 \cdots a_n)$ , in which the  $a_i$ ’s are the only parameters,  $(\forall a_1 \cdots a_n \cdot (\forall b_1 \cdots b_n \cdot U(a_1) \wedge \cdots \wedge U(a_n) \wedge U(b_1) \wedge \cdots$

Received May 27, 2010; accepted December 12, 2011

2010 Mathematics Subject Classification: Primary 03E70

Keywords: New Foundations, NF, urelements, homogeneity, models with automorphisms

© 2012 by University of Notre Dame 10.1215/00294527-1722764

$\wedge U(b_n) \rightarrow \varphi(a_1 \cdots a_n) \leftrightarrow \varphi(b_1 \cdots b_n))$  holds, where  $U(x)$  means “ $x$  is an urelement.” Of course a model of NF would be such a model of NFU. (It is relatively easy to show that the urelements of any model of NFU are homogeneous with respect to stratified formulas, though we do not give details here: one uses the permutation techniques adapted to NF by Dana Scott in [16] and discussed in the context of NFU in Crabbé [2].)

The author’s (incorrect!) response was that of course there are such models, because all the urelements in the “usual” models of NFU (described in detail below) are indistinguishable in the suitable sense, because all information about the extensions of the urelements of the model is discarded in the construction.

But it turns out that this is not the case, and for a reason which perhaps has more interest than the answer to the original question (which remains unanswered, and may have some of the flavor of the unsolved problem of the consistency of NF). The usual construction of a model of NFU starts with a model of (some fragment of) ordinary set theory with an automorphism which moves a rank of the cumulative hierarchy. The domain of the model of NFU is a rank in this model moved by the automorphism. Most elements of this rank are treated as urelements in the model of NFU, and it *appears* that information is being discarded in this process. But it is not: the restriction to the domain of the model of NFU of the membership relation of the original model of set theory with automorphism turns out to be first-order definable in the model of NFU, which quite incidentally gives a negative answer to the question as to whether the urelements in the model of NFU are homogeneous.

## 2 NFU

We briefly describe the theory NFU, along with the related theories TST (simple type theory) and NF.

Simple type theory TST is a strongly streamlined version of the type theory of Russell and Whitehead [19]. One follows Ramsey [14] in eliminating the orders in [19]. One follows Wiener [20] in noting that since there is a definable ordered pair in typed set theory, one does not need relation types. Theories of this kind were apparently first proposed about 1930 (see Wang [18] for historical remarks).

TST is a first-order theory with sorts (called *types*) indexed by the natural numbers. The typing conditions for atomic formulas are briefly given by the templates  $x^n = y^n$ ,  $x^n \in y^{n+1}$ . The axioms of TST are the following.

**Extensionality** We have that

$$(\forall A^{n+1} B^{n+1} . A^{n+1} = B^{n+1} \leftrightarrow (\forall x^n . x^n \in A^{n+1} \leftrightarrow x^n \in B^{n+1}))$$

is an axiom for each  $n$ .

**Comprehension** For each formula  $\varphi$  in which  $A^{n+1}$  is not free, and each variable  $x^n$ ,

$$(\exists A^{n+1} . (\forall x^n . x^n \in A^{n+1} \leftrightarrow \varphi))$$

is an axiom.

Axioms of infinity and choice are usually adjoined, whose exact form need not be investigated at this point.

In [13], W. V. O. Quine proposed that the sorts of TST could be collapsed so as to obtain an unsorted first-order theory with equality and membership whose axioms

are exactly the axioms of TST with distinctions of type between variables ignored (in a way which does not introduce any identification between variables of different types). This theory is called NF (New Foundations, after the name of the paper [13]). It is traditional to present it in a way which does not depend on the definition of another theory.

**Extensionality** We have that

$$(\forall AB \cdot A = B \leftrightarrow (\forall x \cdot x \in A \leftrightarrow x \in B))$$

is an axiom.

**Definition 2.1** Let  $\varphi$  be a formula in the language of first-order logic with equality and membership. We say that a function  $\sigma$  from variables to natural numbers is a *stratification* of  $\varphi$  iff for each atomic subformula “ $x = y$ ” of  $\varphi$  we have  $\sigma(“x”) = \sigma(“y”)$  and for each atomic subformula “ $x \in y$ ” we have  $\sigma(“x”) + 1 = \sigma(“y”)$ . We say that  $\varphi$  is *stratified* iff there is a stratification of  $\varphi$ .

**Stratified comprehension** For each stratified formula  $\varphi$  in which  $A$  is not free, and each variable  $x$ ,

$$(\exists A \cdot (\forall x \cdot x \in A \leftrightarrow \varphi))$$

is an axiom.

It should be clear that this is an equivalent description of the theory, as the stratified formulas are exactly those which can be obtained from well-formed formulas of TST by dropping type distinctions without introducing identifications between variables.

In fact, the notion of stratification can be eliminated from the definition of the theory and with it the last reference to even relative notions of type: the axiom scheme of stratified comprehension is equivalent to the conjunction of finitely many of its instances. The usually referenced, though far from the nicest, presentation of this is in Hailperin [5]. A nicer presentation is found in the author’s [8], though this needs to be modified to eliminate references to a primitive ordered pair. (The way to do this is indicated there.)

The consistency of NF remains an open question. In 1969 R. B. Jensen [9] proved the consistency of NFU, which differs from NF only in its formulation of extensionality.

**Weak extensionality** We have that

$$(\forall AB \cdot (\exists y \cdot y \in A) \rightarrow A = B \leftrightarrow (\forall x \cdot x \in A \leftrightarrow x \in B))$$

is an axiom.

Notice that NFU has the same comprehension scheme as NF. The idea is that NFU allows the existence of many urelements without elements in addition to the empty set. Strictly speaking, the formulation above does not allow one to specify an empty set. An inessential and very convenient modification of the theory is to add the empty set as a primitive (which allows one further to define a sethood predicate). We present a full axiomatization of NFU with empty set as a primitive notion.

**Empty set** We have  $(\forall x \cdot x \notin \emptyset)$ .

**Definition 2.2** We have

$$\text{set}(x) \equiv_{\text{def}} x = \emptyset \vee (\exists y \cdot y \in x).$$

**Weak extensionality** We have that

$$(\forall AB \cdot \text{set}(A) \wedge \text{set}(B) \rightarrow A = B \leftrightarrow (\forall x \cdot x \in A \leftrightarrow x \in B))$$

is an axiom for each  $n$ .

**Stratified comprehension** For each stratified formula  $\varphi$  in which  $A$  is not free, and each variable  $x$ ,

$$(\exists A \cdot \text{set}(A) \wedge (\forall x \cdot x \in A \leftrightarrow \varphi))$$

is an axiom.

Specker showed in [17] that NF disproves choice (and so proves infinity). Jensen showed that NFU is consistent with infinity and choice, and further that NFU is consistent with the negation of infinity.

### 3 The Boffa Model Construction

We do not describe the original consistency proof of NFU due to Jensen. Instead, we give a related model construction due to Maurice Boffa [1]. This is appropriate as the aim of this paper is to prove a result about models of the sort described by Boffa. (The class of models we consider is actually slightly less general than the class of models discussed by Boffa.)

We use the term “Mac Lane set theory” for Zermelo set theory with separation restricted to bounded formulas. This system was proposed as a foundation for mathematics by Saunders Mac Lane in [10]. Let  $M$  be a model of Mac Lane set theory with the additional axiom that every set belongs to a rank  $V_\alpha$ . The additional axiom provides no essential additional strength (see Mathias [11]). We can further suppose, using the Ehrenfeucht–Mostowski theorem of model theory (see Ehrenfeucht and Mostowski [3]), that  $M$  has an external automorphism  $j$  that moves a “rank”  $V_\alpha$  (and of course its “ordinal index”  $\alpha$ , which is not a standard ordinal). We can further suppose without loss of generality that  $M \models \alpha > j(\alpha)$  (as we could replace  $j$  with  $j^{-1}$  to make this true in the worst case). We stipulate that the structure we are talking about be of the form  $\langle M, \in_M, j, \alpha \rangle$ , so that we can talk about the automorphism. The structure  $\langle M, \in_M, j, \alpha \rangle$  is a model of Mac Lane set theory + “every set belongs to a rank” with the following modifications:  $j$  and  $\alpha$  are added to the language, along with axioms asserting that  $\alpha$  is an infinite ordinal,  $V_\alpha$  exists, and  $j(\alpha) < \alpha$ , and with separation restricted to formulas in which  $j$  does not appear (which can be extended to formulas in which  $j$  appears in parameters).

We construct a Boffa model  $\langle B, \in_B, j \rangle$  of NFU (with an additional function symbol  $j$  which cannot appear in instances of stratified comprehension, except in parameters; we include it merely so that  $j$  makes sense in  $B$ -formulas) as follows:  $B = \{x \mid M \models x \in V_\alpha\}$ ;  $B$  is the collection of  $M$ -elements of  $V_\alpha$ ;  $x \in_B y$  is defined as  $j(x) \in_M y \wedge y \in_M V_{j(\alpha)+1}$ . Of course  $B \models x = j(y)$  iff  $M \models x = j(y) \wedge y \in V_\alpha$ . Notice that if  $M \models u \in V_\alpha - V_{\alpha+1}$ , then  $B \models (\forall x \cdot x \notin u)$ . (The empty set of  $M$  is also elementless in the model of NFU, and naturally taken to be the empty set of  $B$ .) The  $M$ -elements of  $V_\alpha - V_{\alpha+1}$  are the urelements of the model  $B$ . It should be evident that weak extensionality holds in  $B$ .

We outline the proof that stratified comprehension holds in  $B$  (following Forster [4, pp. 68–70]). Let  $\varphi$  be a stratified formula in the language of NFU. It translates directly into a formula  $\varphi_1$  in the language of  $M$  (by replacing each  $u = v$  with

$u = v$ , each  $\emptyset$  with  $\emptyset^M$ , and each  $u \in v$  with  $j(u) \in v \wedge v \in V_{j(\alpha)+1}$ . If  $M \models \text{“}\{x \in V_\alpha \mid \varphi_1\} \text{ exists,}”$  then we give the name  $A$  to the object such that  $M \models A = \{x \in V_\alpha \mid \varphi_1\}$ . We claim that if  $A$  exists, then we have  $B \models x \in j(A)$  iff  $\varphi: B \models x \in j(A)$  is equivalent to  $M \models j(x) \in j(A) \wedge j(A) \in V_{j(\alpha)+1}$ , which is equivalent to  $M \models j(x) \in j(A)$  because  $M \models j(A) \in V_{j(\alpha)+1}$  is true (as obviously  $M \models A \in V_{\alpha+1}$ ), and of course this is equivalent to  $M \models x \in A$ .

It remains to show that  $A$  exists. This does not follow from the separation axiom of  $M$  immediately, because the formula  $\varphi_1$  will usually contain what seem to be essential occurrences of  $j$ . We show how to convert  $\varphi_1$  to an equivalent formula in which occurrences of  $j$  are confined to parameters, from which it follows that  $A$  exists in  $M$ .

Let  $\sigma$  be a stratification of  $\varphi$ . Let  $N$  be a constant larger than any element of the range of  $\sigma$  (restricted to variables which occur in  $\varphi$ ). Replace each atomic formula  $u = v$  in  $\varphi_1$  with the equivalent  $j^{N-\sigma(u)}(u) = j^{N-\sigma(v)}(v)$ , noting that this is the same as  $j^{N-\sigma(u)}(u) = j^{N-\sigma(v)}(v)$ , because  $\varphi$  is stratified. Replace each atomic formula  $j(u) \in v$  in  $\varphi_1$  with the equivalent  $j^{N-\sigma(u)}(u) \in j^{N-\sigma(v)-1}(v)$ , noting that this is the same as  $j^{N-\sigma(u)}(u) \in j^{N-\sigma(v)}(v)$ , because  $\varphi$  is stratified. The remaining atomic formulas  $v \in V_{j(\alpha)+1}$  are replaced with  $j^{N-\sigma(v)}(v) \in j^{N-\sigma(v)}(V_{j(\alpha)+1})$ . The resulting formula we call  $\varphi_2$ . In  $\varphi_2$ , a variable  $u$  always occurs in the context  $j^{N-\sigma(u)}(u)$ , without any further applications of  $j$ . Now each bound variable  $u$  is restricted to  $V_\alpha$ , so we can replace all occurrences of  $j^{N-\sigma(u)}(u)$  with  $u$  while replacing the bound on the quantifier with  $j^{N-\sigma(u)}(V_\alpha)$ . We do this for each bound variable. We also replace  $j^{N-\sigma(x)}(x)$  with  $x$ . We call the resulting formula  $\varphi_3$ . Since  $j$  now appears only in parameters in  $A_0 = \{x \in j^{N-\sigma(x)}(V_\alpha) \mid \varphi_3\}$ , this set exists in  $M$ , and  $j^{\sigma(x)-N}(A_0)$  is clearly the desired set  $A$ . This completes the proof that the model  $B$  satisfies stratified comprehension.

It is straightforward to establish that  $B$  satisfies infinity and that  $B$  satisfies choice iff  $M$  does. If we modified the construction by providing that  $M \models \text{“}\alpha \text{ is a finite ordinal,}”$  the model  $B$  would satisfy NFU with the negation of the axiom of infinity.

#### 4 Recovering Information from a Boffa Model

If  $M \models A \in V_{\alpha+1}$ , then for each  $y$ ,  $M \models x \in A$  iff  $M \models j(x) \in j(A)$  iff  $B \models x \in j(A)$ : informally, we say that the  $M$ -set  $A$  is implemented by the  $B$ -set  $j(A)$ . An unordered pair  $\{a, b\}$  is implemented by  $\{j(a), j(b)\}$  (the second object being named here in the language of  $M$ ; in the language appropriate to  $B$ , it would be called  $\{a, b\}$ : the extremely precise statement of this is that for any  $u, a, b$ ,  $B \models u = \{a, b\}$  iff  $M \models u = \{j(a), j(b)\}$ ). The ordered pair  $\langle a, b \rangle$  in the sense of  $B$  is then  $\langle j^2(a), j^2(b) \rangle$  in the sense of  $M$ . Finally, if  $f$  is a function in  $M$ , we see that  $j^3(f)$  is the function with the same extension in  $B$ :  $M \models y = f(x)$  iff  $M \models \langle x, y \rangle \in f$  iff  $M \models \langle j^2(x), j^2(y) \rangle \in j^2(f)$  iff  $B \models \langle x, y \rangle \in j^3(f)$  (clearly  $M \models \langle j^2(x), j^2(y) \rangle \in V_{j(\alpha)+1}$ ). So we see that every  $M$ -function from  $V_\alpha$  to  $V_\alpha$  is implemented in  $B$  as a function from the universe to the universe (note that  $j(V_\alpha)$  is the universal set in the sense of  $B$ :  $M \models x \in V_\alpha$  implies  $M \models j(x) \in j(V_\alpha)$  which in turn implies  $B \models x \in j(V_\alpha)$ ). We denote the universal set by  $V$  and the collection of all sets by  $P(V)$  in  $B$ -formulas: the same objects are  $j(V_\alpha)$  and  $j^2(V_{\alpha+1})$  in  $M$ .

We introduce a nonstandard piece of notation common in  $NF$  studies (ultimately derived from [19]), because we will shortly use it.  $\iota(x)$  is a notation for  $\{x\}$ , and

so  $\iota^{\ast}A$  is notation for the collection of one-element subsets of  $A$  (the elementwise image of  $A$  under the singleton operation).

We introduce a specific function found in  $M$ , namely, the function  $\mathbf{S} = j^3(\{\{\{x\}, x\} \mid x \in j(V_\alpha)\})$ .  $M \models$  “the domain of  $j^{-3}(\mathbf{S})$  is  $\iota^{\ast}j(V_\alpha)$  and its range is  $j(V_\alpha)$ ,” so of course  $M \models j^{-3}(\mathbf{S}) : \iota^{\ast}j(V_\alpha) \rightarrow j(V_{\alpha+1})$ . An  $M$ -element of  $\iota^{\ast}j(V_\alpha)$  is of the form  $\{j(x)\}$ , which  $B$  sees as  $\{x\}$ , whereas  $M$ -elements of  $j(V_{\alpha+1})$  are exactly the sets in the sense of  $B$ :  $B \models \mathbf{S} : \iota^{\ast}V \rightarrow P(V)$ : that is,  $B$  says that  $\mathbf{S}$  is a function from the set of all singletons into the set of all sets. Further,  $M \models (\forall x \in V_\alpha \cdot j^{-3}(\mathbf{S})(\{j(x)\}) = j(x))$ : thus  $B \models (\forall x \cdot \mathbf{S}(\{x\}) = j(x))$ .  $\mathbf{S}$ , which is a set map, codes the external automorphism  $j$  in a certain sense ( $j$  is not a set function in either  $M$  or  $B$ ).

The function  $\mathbf{S}$  has a further property of considerable interest in  $B$ . Suppose  $M \models A \in j(V_{\alpha+1})$  (equivalently,  $B \models$  “ $A$  is a set”). We have  $B \models x \in A$  iff  $M \models j(x) \in A$  iff  $M \models j^2(x) \in j(A)$  iff  $B \models j(x) \in j(A)$  (certainly  $M \models j(A) \in V_{j(\alpha+1)}$ ) iff  $B \models \mathbf{S}(\{x\}) \in \mathbf{S}(\{A\})$ . Further, if  $B \models y \in j(A)$ , then  $M \models j(y) \in j(A)$ , so  $M \models y \in A$ , so  $M \models y \in V_{j(\alpha)}$ , so  $M \models y = j(x)$  for some  $x$  such that  $M \models x \in V_\alpha$ , so  $B \models y = \mathbf{S}(\{x\})$  for some such  $x$ . Thus  $B \models$  “for any set  $A$ ,  $\mathbf{S}(\{A\}) = \{\mathbf{S}(\{x\}) \mid x \in A\}$ .”

We summarize this statement in the format of an axiom to adjoin to NFU (as we did in Holmes [7] where we first introduced these ideas with a rather different application in view).

**Axiom of Endomorphism (to adjoin to NFU)** There is an injective function  $\mathbf{S} : \iota^{\ast}V \rightarrow P(V)$  such that for any set  $A$ ,  $\mathbf{S}(\{A\}) = \{\mathbf{S}(\{x\}) \mid x \in A\}$ .

We now observe that in the Boffa model  $B$  it is possible to define the restriction of the membership relation of  $M$  to  $V_\alpha$  in terms of the membership relation of  $B$  and the function  $\mathbf{S}$ :  $M \models x \in y$  iff  $M \models j(x) \in j(y)$  iff  $B \models x \in j(y)$  iff  $B \models x \in \mathbf{S}(\{y\})$ .

**Definition 4.1 (NFU + endomorphism)**  $x E y$  is defined as  $x \in \mathbf{S}(\{y\})$ . Notice that as the relative types of  $x$  and  $y$  are the same in the definition of  $x E y$ ,  $E$  is a set relation:  $\{\langle x, y \rangle \mid x E y\}$  exists by stratified comprehension:

$$B \models x E y \text{ iff } M \models x \in y.$$

The final claim which establishes our main result is that if there is a function  $\mathbf{S}$  which witnesses the truth of the Axiom of Endomorphism in a Boffa model, then there is exactly one such function. We can then define  $x \in^* y$  as the following: there is a function  $\mathbf{S}$  which witnesses the truth of the Axiom of Endomorphism, and  $x \in \mathbf{S}(\{y\})$ , a statement which can be expanded out into a sentence in the first-order language of NFU. Considerations shown above already show that (subject to our final claim) the relation  $\in^*$  will coincide with  $\in_M$  restricted to the domain of  $B$ .

Because the domain of  $B$  is a rank of the cumulative hierarchy in  $M$ , there is a notion of ordinal rank of elements of the domain (an element  $x$  has rank  $\beta$  iff  $\beta$  is the least ordinal such that  $x \subseteq V_\beta$ ). It is the case that the ordinal rank of a set  $A$  is the successor of the supremum of the set of ranks of elements of  $A$ . We briefly explain why ranks of the cumulative hierarchy are definable in the weak set theory we are using. Recall that this is Mac Lane set theory (Zermelo set theory with bounded separation) with the assumption that every set belongs to a rank, which we will formally state in the course of this development.

**Definition 4.2** A subhierarchy is a well-ordering  $\leq_H$  such that the  $\leq_H$ -least set is  $\emptyset$ , the  $\leq_H$ -immediate successor of any set  $x$  is  $P(x)$  if it exists, and the  $\leq_H$ -supremum of a subset  $A$  of the domain with no  $\leq_H$ -maximum is  $\bigcup A$ . Note that any subhierarchy agrees with the inclusion order on its domain.

**Theorem 4.3** *If  $\leq_G$  and  $\leq_H$  are subhierarchies, either they are equal or one is an initial segment of the other.*

The proof is omitted.

**Definition 4.4** A rank of the cumulative hierarchy is a set which is an element of some subhierarchy.

**Axiom of rank** Every set is a subset of some rank. (This is a more formal statement of the assumption made about our model  $M$  that every set belongs to some rank of the cumulative hierarchy.)

**Definition 4.5** The rank of a set  $A$  is the smallest rank in the inclusion order containing  $A$  as a subset.

**Definition 4.6** The (Scott) order type of a well-ordering  $W$  is the set of all well-orderings isomorphic to  $W$  and belonging to the smallest rank in the inclusion order containing well-orderings isomorphic to  $W$ . A (Scott) ordinal is a set which is the Scott order type of some well-ordering.

**Definition 4.7** If  $r$  is a rank, we say  $r = V_\alpha$  iff  $\alpha$  is the Scott order type of the inclusion order on proper subranks of  $r$ . It is not expected here that  $V_\alpha$  exists for every ordinal  $\alpha$ ; it is a consequence of Mac Lane set theory with the Axiom of Rank (and not a consequence of Zermelo or Mac Lane set theory by themselves) that there is an infinite rank, so  $V_\omega$  does exist. It can also be noted that if the rank  $V_\alpha$  exists, it follows that the usual von Neumann ordinal  $\alpha$  exists.

**Definition 4.8** The ordinal rank of a set  $x$  is the least  $\alpha$  such that  $x \subseteq V_\alpha$ .

The trick used here to define ordinals in Mac Lane set theory with the Axiom of Rank is due to Dana Scott in [15]. The same trick can be used to represent other sorts of isomorphism classes as sets. (A common application of Scott's trick is to define cardinals in ZF, where the usual von Neumann definition of cardinal number does not work). In Mac Lane or Zermelo set theory without rank there is no reasonable global way to represent cardinal or ordinal numbers: an extensive development of mathematics in a theory which is essentially Zermelo set theory with the Axiom of Rank, making extensive use of Scott's trick, is found in Potter [12]. The axiom of rank does not essentially strengthen Zermelo or Mac Lane set theory; this can be seen in [11].

Suppose that  $B$  satisfies the assertion that there is another map  $S^*$  such that  $S^* : {}^tV \rightarrow P(V)$  and for any set  $A$ ,  $S^*({}^tA) = \{S^*({}^t\{x\}) \mid x \in A\}$ . Define  $x E^* y$  as  $x \in S^*({}^t\{y\})$ .

Suppose  $S \neq S^*$ . Then there is a minimal ordinal  $\beta$  such that there is an element  $x$  of  $V_\alpha$  of ordinal rank  $\beta$  such that  $S(\{x\}) \neq S^*(\{x\})$ .

Now  $B \models S(\{\{x\}\}) = \{S(\{x\})\} \neq \{S^*(\{x\})\} = S^*(\{\{x\}\})$ , so if  $x$  is a counterexample, so is the object that the model  $B$  calls  $\{x\}$ , which is the object that  $M$  calls  $\{j(x)\}$ , which has ordinal rank  $j(\beta) + 1$ . It follows that  $\beta \leq j(\beta) + 1$ . In fact,

since  $\beta = j(\beta) + 1$  is impossible (consider the parity of the finite part of the ordinal  $\beta$ ), it follows that  $\beta < j(\beta) + 1$ , so  $\beta \leq j(\beta)$ .

Since  $\beta \leq j(\beta)$  and  $\beta < \alpha$ , it is clear that  $\beta \leq j(\beta) < j(\alpha)$ , from which it follows that a counterexample must be a subset of  $V_{j(\alpha)}$ , so an element of  $V_{j(\alpha)+1}$ , so a  $B$ -set.

If  $x$  is a counterexample of minimal ordinal rank  $\beta$  and  $B \models y \in x$ , then  $M \models j(y) \in x$ , so the ordinal rank of  $j(y)$  is less than the ordinal rank of  $x$ : if the ordinal rank of  $y$  is  $\gamma$ , we have  $j(\gamma) < \beta$ , so we have  $\gamma < j^{-1}(\beta) \leq \beta$ . It follows that for any  $y$  such that  $B \models y \in x$ , we have  $B \models \mathbf{S}(\{y\}) = \mathbf{S}^*(\{y\})$ .

Now since  $x$  is a  $B$ -set, we have  $B \models \mathbf{S}(\{x\}) = \{ \mathbf{S}(\{y\} \mid y \in x \} = \{ \mathbf{S}^*(\{y\} \mid y \in x \} = \mathbf{S}^*(\{x\})$ , which contradicts our initial assumptions about  $x$ . We conclude that there can be only one function witnessing the Axiom of Endomorphism in a Boffa model, from which it follows that the membership relation of the original model is definable in the Boffa model, which is what we set out to prove.

## 5 Further Remarks and Conclusions

It is worth noting that, while the notion of ordinal rank used in this argument is an  $M$ -notion, it is definable in  $B$ , and in fact we can formulate an assumption about  $B$  which is equivalent to the assertion that the universe of  $B$  is a rank of the cumulative hierarchy according to  $M$ . In any well-founded relation  $W$ , we can define in NFU as in ordinary set theory a notion of ordinal rank of elements of the domain of  $W$ . Further, we can say that an ordinal rank  $\beta$  in a relation  $W$  is *complete* if every subset of the collection of ordinals of rank  $\leq \beta$  is the  $W$ -preimage of some element of the domain of  $W$  of rank  $\leq \beta + 1$ . Because of the way  $B$  is constructed, an assertion satisfied by  $B$  is that  $E$  is a well-founded relation, and any ordinal rank in  $E$  is either complete or the entire domain of  $E$ . NFU with the Axiom of Endomorphism and this additional assertion proves that there is at most one function witnessing the truth of the Axiom of Endomorphism. It is much easier to formulate the argument as we have given it in terms of  $M$ 's notion of rank, as we avoid technicalities about the way ordinals are defined in NFU and peculiar properties of the ordinals of NFU.

It is further worth remarking that, while it was already well known (and is easily seen from the consistency proof for NFU given above) that NFU + infinity has precisely the consistency strength of Mac Lane set theory, the argument of this paper shows that there is a slight extension of NFU (NFU + Endomorphism + “ $E$  is a well-founded relation and any ordinal rank in  $E$  is either complete or the entire domain of  $E$ ”) which has a rather more intimate relationship of mutual interpretability with Mac Lane set theory (without infinity) enhanced with an automorphism of the universe which moves a rank: we have seen above that we can derive a model of the extension of NFU from the extension of Mac Lane which defines all the concepts of Mac Lane set theory as restricted to its domain. It can further be noted that from the model of NFU we can recover an entire model of Mac Lane, not just a rank: the elements of the interpreted Mac Lane are pairs  $\langle x, n \rangle$ , where  $x$  is a set and  $n$  is a natural number, with the intention that  $\langle x, n \rangle$  code  $j^{-n}(x)$ ; we do not give the (easy) full details of this development here. The model of Mac Lane that is recovered is in effect truncated at the supremum of the ranks indexed by  $j^{-n}(\alpha)$ 's; this is of course not an ordinal of  $M$ , and it is important here that we are working in Mac Lane rather than Zermelo set theory, as unbounded quantifiers would be a problem

in the latter context. There is another way to interpret Mac Lane set theory with an automorphism in NFU (or more accurately the theory of a rank in a model of Mac Lane moved downward by an automorphism): the objects of this interpretation are isomorphism classes of well-founded extensional relations with a top element. The methods used are derived from Hinnion [6], and full details can be found in [8].

It is quite striking that in the usual models of NFU, the apparently featureless urelements are thus seen to be far from featureless. A predicate of an urelement  $u$  which distinguishes urelements from one another is easily described:  $\emptyset E u$  is a very simple example. (Its full form in the language of NFU would be the expansion of “there is a function  $\mathbf{S}$  witnessing the truth of the Axiom of Endomorphism such that  $\emptyset \in \mathbf{S}(\{u\})$ .”) But this is not the main point of this result: the main point is that one has stronger interpretability of the ambient Mac Lane set theory in which a model of NFU is constructed by the Boffa procedure than one would expect.

### References

- [1] Boffa, M., “ZFJ and the consistency problem for NF,” pp. 102–6 in *Jahrbuch der Kurt Gödel Gesellschaft 1988*. [Zbl 0721.03035](#). [MR 1023332](#). [574](#)
- [2] Crabbé, M., “On the set of atoms,” *Logic Journal of the IGPL*, vol. 8 (2000), pp. 751–59. [Zbl 0966.03049](#). [MR 1830525](#). [572](#)
- [3] Ehrenfeucht, A., and A. Mostowski, “Models of axiomatic theories admitting automorphisms,” *Fundamenta Mathematicae*, vol. 43 (1956), pp. 50–68. [Zbl 0073.00704](#). [MR 0084456](#). [574](#)
- [4] Forster, T. E., *Set Theory with a Universal Set: Exploring an Untyped Universe*, 2nd ed., vol. 31 of *Oxford Logic Guides*, Oxford University Press, New York, 1995. [MR 1366833](#). [574](#)
- [5] Hailperin, T., “A set of axioms for logic,” *Journal of Symbolic Logic*, vol. 9 (1944), pp. 1–19. [Zbl 0060.02201](#). [MR 0009753](#). [573](#)
- [6] Hinnion, R., “Sur la théorie des ensembles de Quine,” Ph.D. dissertation, Université libre de Bruxelles, Brussels, 1975. [579](#)
- [7] Holmes, M. R., “The axiom of anti-foundation in Jensen’s ‘New Foundations with urelements,’” *Bulletin de la Societe Mathematique de la Belgique, Series B*, vol. 43 (1991), pp. 167–79. [Zbl 0755.03030](#). [MR 1314689](#). [576](#)
- [8] Holmes, M. R., *Elementary Set Theory with a Universal Set*, vol. 10 of *Cahiers du Centre de Logique*, Université Catholique de Louvain, Louvain-la-Neuve, Belgium, 1998. [Zbl 0959.03001](#). [MR 1759289](#). [573](#), [579](#)
- [9] Jensen, R. B., “On the consistency of a slight (?) modification of Quine’s ‘New Foundations,’” *Synthese*, vol. 19 (1969), pp. 250–63. [573](#)
- [10] Mac Lane, S., *Mathematics, Form and Function*, Springer, New York, 1986. [Zbl 0675.00001](#). [MR 0816347](#). [574](#)
- [11] Mathias, A. R. D., “The strength of Mac Lane set theory,” *Annals of Pure and Applied Logic*, vol. 110 (2001), pp. 107–234. [Zbl 1002.03045](#). [MR 1846761](#). [574](#), [577](#)
- [12] Potter, M., *Set Theory and Its Philosophy: A Critical Introduction*, Oxford University Press, New York, 2004. [Zbl 1066.03002](#). [MR 2050123](#). [577](#)
- [13] Quine, W. V. O., “New Foundations for Mathematical Logic,” *American Mathematical Monthly*, vol. 44 (1937), pp. 70–80. [Zbl 0016.19301](#). [MR 1523848](#). [572](#), [573](#)
- [14] Ramsey, F. P., “The foundations of mathematics,” *Proceedings of the London Mathematical Society (2)*, vol. 25 (1926), pp. 338–84. [Zbl 52.0046.01](#). [MR 1575288](#). [572](#)
- [15] Scott, D., “Definitions by abstraction in axiomatic set theory” (abstract 626t), *Bulletin of the American Mathematical Society*, vol. 61 (1955), p. 442. [577](#)

- [16] Scott, D., "Quine's individuals," pp. 111–15 in *Logic, Methodology and Philosophy of Science*, edited by E. Nagel, Stanford University Press, Stanford, 1962. [Zbl 0151.00904](#). [MR 0152444](#). [572](#)
- [17] Specker, E. P., "The axiom of choice in Quine's 'New Foundations for Mathematical Logic,'" *Proceedings of the National Academy of Sciences of the U. S. A.*, vol. 39 (1953), pp. 972–75. [Zbl 0051.03705](#). [MR 0059214](#). [574](#)
- [18] Wang, H., *Logic, Computers, and Sets*, Chelsea, New York, 1970. [Zbl 0212.31101](#). [MR 0269480](#). [572](#)
- [19] Whitehead, A. N., and B. Russell, *Principia Mathematica*, 3 vols., Cambridge University Press, Cambridge, 1927. [572](#), [575](#)
- [20] Wiener, N., "A simplification of the logic of relations," *Proceedings of the Cambridge Philosophical Society*, vol. 17 (1914), pp. 387–90. [Zbl 45.0122.16](#). [572](#)

Department of Mathematics  
Boise State University  
1910 University Drive  
Boise, Idaho 83725  
USA  
[holmes@math.boisestate.edu](mailto:holmes@math.boisestate.edu)