

A Note on the Axioms for Zilber's Pseudo-Exponential Fields

Jonathan Kirby

Abstract We show that Zilber's conjecture that complex exponentiation is isomorphic to his pseudo-exponentiation follows from the a priori simpler conjecture that they are elementarily equivalent. An analysis of the first-order types in pseudo-exponentiation leads to a description of the elementary embeddings, and the result that pseudo-exponential fields are precisely the models of their common first-order theory which are atomic over exponential transcendence bases. We also show that the class of all pseudo-exponential fields is an example of a nonfinitary abstract elementary class, answering a question of Kesälä and Baldwin.

1 Introduction

Zilber's *pseudo-exponential fields* were defined in Zilber [7] as models of certain axioms. Infinitary logic is used throughout the paper, which shows that the axioms are expressible by an $L_{\omega_1, \omega}(Q)$ -sentence. Anand Pillay emphasized to me the importance of understanding the first-order theory of these pseudo-exponential fields, which we denote by $T_{\mathbb{B}}$. In [4], Zilber and I gave an axiomatization of $T_{\mathbb{B}}$, assuming the Diophantine conjecture on intersections of subvarieties with tori (CIT). This note takes some steps towards understanding $T_{\mathbb{B}}$ unconditionally, in particular by describing the first-order types which are realized in pseudo-exponential fields. In Section 2, I show that the axioms for pseudo-exponential fields can be written in a first-order way, with two exceptions: an $L_{\omega_1, \omega}$ -sentence is needed to omit the type of a nonstandard integer, and the quantifier Q (there exist uncountably many) is essential for the countable closure property (CCP), although the axiom can be expressed in an $L(Q)$ -scheme, rather than just as an $L_{\omega_1, \omega}(Q)$ -sentence. These nonelementary axioms are

Received December 19, 2011; accepted May 28, 2012

2010 Mathematics Subject Classification: Primary 03C65; Secondary 03C48

Keywords: pseudo-exponentiation, exponential fields, Schanuel property, first-order theory, abstract elementary class

© 2013 by University of Notre Dame 10.1215/00294527-2143844

known to hold in \mathbb{C}_{exp} , so it follows immediately that if \mathbb{C}_{exp} satisfies the first-order theory of pseudo-exponential fields, then it satisfies the full $L_{\omega_1, \omega}(Q)$ -theory.

The main result of Zilber's paper is that there is a unique model of his axioms of cardinality 2^{\aleph_0} , which we call \mathbb{B} , and indeed of each uncountable cardinal κ . Thus we have the following.

Theorem 1 *If the complex exponential field \mathbb{C}_{exp} is elementarily equivalent to \mathbb{B} , then it is isomorphic to \mathbb{B} .*

The isomorphism is Zilber's main conjecture about \mathbb{C}_{exp} . However, Theorem 1 does not make proving the isomorphism any easier; rather it shows that proving elementary equivalence is just as far out of reach.

The class of pseudo-exponential fields naturally forms an abstract elementary class (AEC). Hyttinen and Kesälä [1] introduced the notion of a *finitary* AEC, and the class of pseudo-exponential fields has been used as an example of a nonfinitary AEC, but no proof of this fact has previously appeared in the literature. From the proof that the Q -quantifier cannot be eliminated, the next theorem follows quickly.

Theorem 2 *The category of pseudo-exponential fields (with CCP) together with closed embeddings is an abstract elementary class which is not finitary.*

Section 3 of this note uses the algebra of exponential fields developed in Kirby [3] and Kirby and Zilber [4] to address the issues of elementary embeddings and atomicity for pseudo-exponential fields. In fact here we can work with a broader class, not assuming the countable closure axiom, but unfortunately still narrower than the class of all models of $T_{\mathbb{B}}$. The main technical difficulty is that, without CIT, we do not understand what saturated models of $T_{\mathbb{B}}$ look like, and we do not have an unconditional quantifier-elimination result, so the proofs have to use less direct techniques, including a careful analysis of the types which are realized.

Theorem 3 *The strong embeddings between pseudo-exponential fields (not necessarily with CCP) are exactly the elementary embeddings. In particular, all pseudo-exponential fields, including those of finite exponential transcendence degree, are elementarily equivalent.*

This theorem complements [7, Theorem 5.13], which adds the additional hypothesis of infinite exponential transcendence degree but gets the stronger conclusion that strong embeddings are $L_{\omega_1, \omega}$ -embeddings. The stronger conclusion fails in general.

Finally, we show the following.

Theorem 4 *Each pseudo-exponential field (not necessarily with CCP) is an atomic model of $T_{\mathbb{B}}$ over an exponential transcendence base, in the language $\langle +, \cdot, \text{exp} \rangle$. In particular, $T_{\mathbb{B}}$ has a prime model. Conversely, every model of $T_{\mathbb{B}}$ which is atomic over an exponential transcendence base is a pseudo-exponential field.*

Zilber states in [7, Theorem 5.16] that the uncountable pseudo-exponential fields (with the countable closure property) are prime over exponential transcendence bases, but he uses an expanded language L^* which is an expansion of our language by $L_{\omega_1, \omega}$ -definitions, not by first-order definitions. So Theorem 4 does not follow directly from Zilber's methods.

2 The Axioms for Pseudo-Exponentiation

We give five axioms capturing Zilber's definition. Explanations of the terminology in axioms 4 and 5 are deferred to the more detailed discussions afterward. We use only the language $\langle +, \cdot, \exp \rangle$.

1. **ELA-field:** F is an algebraically closed field of characteristic zero, and its exponential map \exp is a homomorphism from its additive group to its multiplicative group, which is surjective. (The letters *ELA* stand for *exponential, logarithm, and algebraically closed*.)
2. **Standard kernel:** The kernel of the exponential map is an infinite cyclic group generated by a transcendental element τ .
3. **Schanuel property:** The *predimension function*

$$\delta(\bar{x}) := \text{td}(\bar{x}, \exp(\bar{x})) - \text{ldim}_{\mathbb{Q}}(\bar{x})$$

satisfies $\delta(\bar{x}) \geq 0$ for all tuples \bar{x} from F .

4. **Strong exponential-algebraic closedness:** If V is a rotund, additively and multiplicatively free subvariety of $\mathbb{G}_a^n \times \mathbb{G}_m^m$ defined over F and of dimension n , and \bar{a} is a finite tuple from F , then there is an \bar{x} in F such that $(\bar{x}, e^{\bar{x}}) \in V$ and is generic in V over \bar{a} .
5. **Countable closure property:** For each finite subset C of F , the exponential algebraic closure $\text{ecl}^F(C)$ of C in F is countable.

Clearly axiom 1 is first-order expressible.

2.1 Standard kernel Write $\ker(F) = \{x \in F \mid \exp(x) = 1\}$, the kernel of the exponential map. Define

$$Z(F) = \{r \in F \mid \forall x [x \in \ker \rightarrow rx \in \ker]\},$$

the *multiplicative stabilizer* of the kernel.

Axiom 2 can be split into three parts.

- 2a: The kernel is a cyclic Z -module.
- 2b: Every element of the kernel is transcendental over Z .
- 2c **Standard integers:** We have

$$(\forall r \in Z) \bigvee_{n \in \mathbb{N}} r = \underbrace{1 + \dots + 1}_n \vee r = \underbrace{1 + \dots + 1}_n = 0.$$

It is clear that axioms 2a and 2b are first-order expressible and that 2c is not, but it is given by a single $L_{\omega_1, \omega}$ -sentence. Equivalently, 2c can be viewed as omitting the incomplete type of a nonstandard integer.

2.2 The Schanuel property Axiom 3, the Schanuel property, is equivalent to the following axiom scheme, which is first-order expressible provided that axiom 2 holds, so we can quantify over the standard integers: for each $n \in \mathbb{N}$, for each subvariety $V \subseteq \mathbb{G}_a^n \times \mathbb{G}_m^n$ defined over \mathbb{Q} , of dimension $n - 1$,

$$(\forall x_1, \dots, x_n) (\exists \bar{m} \in \mathbb{Z}^n \setminus \{\bar{0}\}) \left[(\bar{x}, \exp(\bar{x})) \in V \rightarrow \sum_{i=1}^n m_i x_i = 0 \right].$$

2.3 Strong exponential-algebraic closedness Write G for the algebraic group $\mathbb{G}_a \times \mathbb{G}_m$. Each matrix $M \in \text{Mat}_{n \times n}(\mathbb{Z})$ defines a homomorphism $G^n \xrightarrow{M} G^n$ by acting as a linear map on \mathbb{G}_a^n and as a multiplicative map on \mathbb{G}_m^n . If $V \subseteq G^n$, we write $M \cdot V$ for its image. Note that if V is a subvariety of G^n , then so is $M \cdot V$.

An irreducible subvariety V of G^n is said to be *rotund* if and only if for every matrix $M \in \text{Mat}_{n \times n}(\mathbb{Z})$ we have $\dim M \cdot V \geq \text{rk } M$.

Suppose that (\bar{x}, \bar{y}) is a generic point of V over F , with the x_i being the coordinates from \mathbb{G}_a and the y_i the coordinates from \mathbb{G}_m . We say that V is *multiplicatively free* if and only if the y_i do not satisfy any equation of the form $\prod_{i=1}^n y_i^{m_i} = b$ with the $m_i \in \mathbb{Z}$, not all zero, and $b \in \mathbb{G}_m(F)$. Equivalently, the projection of V to \mathbb{G}_m^n does not lie in any coset of a proper algebraic subgroup of \mathbb{G}_m^n . Similarly, we say that V is *additively free* if and only if the x_i do not satisfy any equation of the form $\sum_{i=1}^n m_i x_i = a$ with the $m_i \in \mathbb{Z}$, not all zero, and $a \in F$.

Proposition *Axiom 4, strong exponential-algebraic closedness, is first-order expressible modulo axioms 1, 2, and 3.*

Proof We consider parametric families $(V_p)_{p \in P}$ of subvarieties of G^n , where P is some parameterizing variety. It is a well-known fact (part of the *fiber dimension theorem*) that the set of p such that V_p is irreducible and of dimension n is first-order definable in the field language. The property of being additively free is not definable in the field language since, for example, the subvariety of G^2 given by the equation

$$x_1 + px_2 = 0$$

is additively free if and only if $p \notin \mathbb{Q}$. However, it is definable as follows, allowing quantification over \mathbb{Z} :

$$(\forall \bar{m} \in \mathbb{Z}^n \setminus \{\bar{0}\}) \forall z \exists \bar{x} \left[\bar{x} \in V_p \wedge \sum_{i=1}^n m_i x_i \neq z \right].$$

It is easy to give similar definitions showing that rotundity and multiplicative freeness are definable allowing quantification over \mathbb{Z} . However, Theorem 3.2 of [7] shows that these two properties are even first-order definable in the field language. For a parametric family $(V_p)_{p \in P}$ of subvarieties of G^n , let P' be the set of $p \in P$ consisting of those p such that V_p is irreducible, of dimension n , rotund, and additively and multiplicatively free. Consider the following axiom scheme with one axiom for each family $(V_p)_{p \in P}$ and each natural number r :

$$(\forall p \in P') (\forall \bar{a} \in F^r) (\exists \bar{x} \in F^n) (\forall \bar{m} \in \mathbb{Q}^{n+r}) \left[(\bar{x}, e^{\bar{x}}) \in V_p \wedge \left(\sum_{i=1}^n m_i x_i + \sum_{i=1}^r m_{n+i} a_i = 0 \rightarrow \bigwedge_{i=1}^n m_i = 0 \right) \right].$$

This scheme is first-order expressible assuming that axiom 2 holds, so we can quantify over \mathbb{Z} and hence over \mathbb{Q} . It also follows from axiom 4, since that axiom gives an \bar{x} such that $(\bar{x}, e^{\bar{x}})$ is generic in V_p over \bar{a} . Since V_p is additively free, that \bar{x} does not satisfy any \mathbb{Q} -linear equation over \bar{a} , which is all that the scheme requires.

Now let V be any rotund, additively and multiplicatively free subvariety of G^n defined over F and of dimension n , and let \bar{a} be a finite tuple from F , as in the hypothesis of axiom 4. Then V is V_p for some parametric family $(V_p)_{p \in P}$ as above, with $p \in P'$. By extending \bar{a} , we may assume that V is defined over \bar{a} . By extending

again and using axiom 3, we may assume that $\delta(\bar{a}) \leq \delta(\bar{y}, \bar{a})$ for all tuples \bar{y} from F (since the values of $\delta(\bar{a}, \bar{y})$ lie in \mathbb{N} , so the minimum is attained). Thus for all \bar{y} , $\delta(\bar{y}/\bar{a}) := \delta(\bar{y}, \bar{a}) - \delta(\bar{a}) \geq 0$. Now invoke the axiom scheme to find an \bar{x} . We have $\text{ldim}_{\mathbb{Q}}(\bar{x}/\bar{a}) = n$, so we must have $\text{td}(\bar{x}, e^{\bar{x}}/\bar{a}, e^{\bar{a}}) \geq n$. But $\dim V = n$, so $(\bar{x}, e^{\bar{x}})$ must be generic in V over $(\bar{a}, e^{\bar{a}})$ and so a fortiori over \bar{a} . Thus axiom 4 is equivalent to this scheme, modulo axioms 1–3. \square

On [7, p. 87], Zilber remarks:

The definition [of strongly exponentially-algebraically closed] assumes a “slight saturatedness” of the exponentially-algebraically closed structure.

This remark had led me to assume that strong exponential-algebraic closedness was not first-order, even assuming the other axioms, so the above result was somewhat unexpected. Indeed the fact that strong exponential-algebraic closedness is first-order means that Zilber’s other notion of *exponential-algebraic closedness* is redundant for the construction of the exponential fields which do have standard kernel and the Schanuel property (although it is used in this note, in Section 3.1, where both these properties fail). In [4, Theorem 5.5] we prove that the notions of exponential-algebraic closedness and strong exponential-algebraic closedness coincide under the additional assumption that the Diophantine conjecture CIT is true (and assuming the other relevant axioms), and it would be interesting to know if the same result can be proved unconditionally.

2.4 The countable closure property In any exponential field F there is a pregeometry called exponential algebraic closure, which we write ecl^F . We give a quick account of its definition. Details can be found in Macintyre [6] or Kirby [2]. An exponential polynomial (without iterations of exponentiation) is a function of the form $f(\bar{X}) = p(\bar{X}, e^{\bar{X}})$, where $p \in F[X_1, \dots, X_n, Y_1, \dots, Y_n]$ is a polynomial. We can extend the formal differentiation of polynomials to exponential polynomials in a unique way such that $\frac{\partial e^X}{\partial X} = e^X$.

A *Khovanskii system of width n* consists of exponential polynomials f_1, \dots, f_n with equations

$$f_i(x_1, \dots, x_n) = 0 \quad \text{for } i = 1, \dots, n \tag{1}$$

and the inequation

$$\begin{vmatrix} \frac{\partial f_1}{\partial X_1} & \cdots & \frac{\partial f_1}{\partial X_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial X_1} & \cdots & \frac{\partial f_n}{\partial X_n} \end{vmatrix} (x_1, \dots, x_n) \neq 0, \tag{2}$$

where the differentiation here is the formal differentiation of exponential polynomials.

For any subset C of F , we define $a \in \text{ecl}^F(C)$ if and only if there are $n \in \mathbb{N}$, $a_1, \dots, a_n \in F$, and exponential polynomials f_1, \dots, f_n with coefficients from $\mathbb{Q}(C)$ such that $a = a_1$ and (a_1, \dots, a_n) is a solution to the Khovanskii system given by the f_i .

We say that $\text{ecl}^F(C)$ is the *exponential algebraic closure* of C in F . If $a \in \text{ecl}^F(C)$, we say that a is *exponentially algebraic* over C in F , and otherwise that it is *exponentially transcendental* over C in F .

Theorem 1.1 of [2] states that ecl^F is a pregeometry in any exponential field. We have stated axiom 5 in terms of this pregeometry. The definition of the pregeome-

try originally used by Zilber is different and only makes sense assuming axiom 2. However, when axiom 2 holds, the two definitions agree by [2, Theorem 1.3]. With Zilber’s original definition one can see that axiom 5 is expressible as an $L_{\omega_1, \omega}(Q)$ -sentence. Using this definition we can show the following.

Proposition *Axiom 5, the countable closure property, is expressible as an $L(Q)$ -scheme.*

Proof For the Khovanskii system on exponential polynomials $f = (f_1, \dots, f_n)$, write $\chi_f(\bar{x}, \bar{z})$ for the first-order formula expressing (1) and (2), where \bar{z} denotes the coefficients for the exponential polynomials. Then axiom 5 is expressed by the $L(Q)$ -sentences

$$(\forall \bar{z}) \neg (Qx_1)(\exists x_2, \dots, x_n) \chi_f(\bar{x}, \bar{z})$$

where f ranges over all finite lists of exponential polynomials with variables \bar{z} as coefficients. □

2.5 The complex exponential field and proof of Theorem 1 The complex exponential field \mathbb{C}_{exp} is the field of complex numbers equipped with the usual complex exponential function given by $\exp(z) = \sum_{n \in \mathbb{N}} \frac{z^n}{n!}$. Axioms 1 and 2 are chosen such that \mathbb{C}_{exp} satisfies them. Zilber noted in [7, Lemma 5.12] that \mathbb{C}_{exp} also satisfies axiom 5, the countable closure property. With the definition of ecl^F from Section 2.4, we can give a shorter proof. Given a finite subset C of \mathbb{C}_{exp} , there are only countably many Khovanskii systems with coefficients from $\mathbb{Q}(C)$. The inequation in a Khovanskii system says that the Jacobian of the functions f_1, \dots, f_n does not vanish, so, by the implicit function theorem, solutions to a Khovanskii system are isolated in the complex topology. Hence there are only countably many solutions to each system, so $\text{ecl}^{\mathbb{C}_{\text{exp}}}(C)$ is countable.

We have seen that axioms 3 and 4 are first-order expressible modulo axioms 1 and 2, and certainly \mathbb{C}_{exp} has cardinality 2^{\aleph_0} , so Theorem 1 follows.

2.6 Strong extensions We now summarize the definitions and results from [3] and [4] which we shall need.

In this note, a *partial exponential subfield* of an ELA-field F is a subfield $F_0 \subseteq F$ together with a \mathbb{Q} -linear subspace $D(F_0)$ of F_0 and the restriction of the exponential map \exp_F to $D(F_0)$, such that F_0 is generated as a subfield of F by $D(F_0) \cup \exp(D(F_0))$. Thus F_0 is determined by $D(F_0)$. For any finite tuple \bar{a} from F we define

$$\delta(\bar{a}/F_0) = \text{td}(\bar{a}, e^{\bar{a}}/F_0) - \text{ldim}_{\mathbb{Q}}(\bar{a}/D(F_0)),$$

and say that F_0 is *strong* in F , written $F_0 \triangleleft F$, if and only if for all $\bar{a} \in F$, $\delta(\bar{a}/F_0) \geq 0$.

If $B \subseteq F$ is a subset, then $\langle B \rangle_F$, the partial exponential subfield of F generated by B , is the partial exponential subfield F_0 with $D(F_0)$ equal to the \mathbb{Q} -linear span of B . We write $B \triangleleft F$ if and only if $\langle B \rangle_F \triangleleft F$.

If F satisfies the Schanuel property, or more generally, if there is $C \subseteq B$ with $C \triangleleft F$, then there is a smallest partial exponential subfield of F which contains B and is strong in F . We write it as $\lceil B \rceil_F$ and call it the *hull* of B in F . For any subset $B \subseteq F$, we write $\langle B \rangle_F^{\text{ELA}}$ for the smallest ELA-subfield of F containing $\ker(F) \cup B$, and we write $\lceil B \rceil_F^{\text{ELA}}$ for $\lceil \langle B \rangle_F \rceil_F^{\text{ELA}}$.

Proposition (A) *Suppose that F is an ELA-field with standard kernel and \bar{b} is a finite tuple from F such that $\bar{b} \triangleleft F$. Then there is $m \in \mathbb{N}^+$ such that the isomorphism type of $\langle \bar{b} \rangle_F^{\text{ELA}}$ is determined by the algebraic locus $V = \text{Loc}(\bar{b}/m, e^{\bar{b}/m} / \ker(F))$, a subvariety of G^n , and does not depend on F . Furthermore, $\langle \bar{b} \rangle_F^{\text{ELA}}$ is strong in F .*

Proof The existence of some $m \in \mathbb{N}^+$ such that the partial exponential subfield $\langle \bar{b} \rangle_F$ is determined up to isomorphism by V follows from the Thumback lemma (see [3, Fact 2.15], or see Zilber [8, Theorem 2] for a proof). Then [3, Theorem 2.18] applies to show that $\langle \bar{b} \rangle_F^{\text{ELA}}$ is uniquely determined and strong in F . \square

Proposition (B) *If $F \triangleleft M$ is a strong extension of ELA-fields such that $\ker(F) = \ker(M)$, M is generated as an ELA-field by F and a finite tuple $\bar{b} \in M^n$, and either F is countable or $\ker(F)$ is \aleph_0 -saturated, then there is $m \in \mathbb{N}^+$ such that the isomorphism type of M as an extension of F is determined by $V = \text{Loc}(\bar{b}/m, e^{\bar{b}/m}/F)$. Furthermore, \bar{b} can be chosen to be \mathbb{Q} -linearly independent over F , and then V is rotund, additively and multiplicatively free, and of dimension at least n . The extension is exponentially algebraic (i.e., $M = \text{ecl}^M(F)$) if and only if $\dim(V) = n$.*

In this case, we write M as $F|V$, “ F extended by V .”

Proof See [3, Sections 3, 5] for the case where F is countable. In the case where $\ker(F)$ is \aleph_0 -saturated, the analysis is exactly the same except that one needs to use [4, Theorem 3.3] in place of [3, Theorem 2.18]. \square

2.7 $L_{\omega_1, \omega}$ -theory Let Ψ be an $L_{\omega_1, \omega}$ -sentence expressing axioms 1–4. For any natural number n , it is easy to give an $L_{\omega_1, \omega}$ -sentence Φ_n specifying that F has exponential-transcendence degree equal to n .

From Section 2.6, we see that axiom 4, strong exponential-algebraic closedness, is equivalent (assuming axioms 1–3) to an existential closedness property for exponentially algebraic strong ELA-extensions, which do not extend the kernel. Lemma 5.9 of [3] shows that these extensions can be freely amalgamated, so, at least in the countable case, we can characterize the models as certain Fraïssé limits. Indeed any ELA-field F which is countable or has \aleph_0 -saturated kernel has a unique smallest strongly exponentially algebraically closed extension, which we write as F^\sim . For more details, see [3, Section 6].

In particular, each of the $L_{\omega_1, \omega}$ -sentences $\Psi_n := \Psi \wedge \Phi_n$ and $\Psi_\infty := \Psi \wedge \bigwedge_{n \in \mathbb{N}} \neg \Phi_n$ is countably categorical by the uniqueness of Fraïssé limits (or specifically by [3, Corollary 6.10]) and hence complete. For Ψ_∞ this was already proved in [7, Theorem 5.13]. Clearly these are the only completions of Ψ as $L_{\omega_1, \omega}$ -theories. In particular, Ψ_∞ gives the complete $L_{\infty, \omega}$ -theory of \mathbb{B} . We write B_n for the unique countable model of Ψ_n .

2.8 Necessity of Q and proof of Theorem 2 We now show that the Q -quantifier cannot be eliminated.

Proposition *The countable closure property is not expressible in $L_{\infty, \omega}$, even modulo axioms 1–4.*

Proof The idea is to find and use an exponentially algebraic regular (in fact, strongly minimal) type which is totally categorical and orthogonal to the generic (exponentially transcendental) type and to the kernel.

Let $F_0 = B_0$, and adjoin an extra element a such that $e^a = a$. This a generates a well-defined ELA-field extension $F_0|V$ of F_0 , where V is the subvariety of G given by the equation $x = y$. Let $F_1 = (F_0 | V)^\sim$, the strong exponential algebraic closure of $F_0|V$. By construction, F_1 satisfies axioms 1–5 and is an exponentially algebraic extension of F_0 , so it has the same exponential transcendence degree, zero. Thus $F_1 \cong F_0$.

Now suppose that \bar{b} is any finite tuple from F_0 , and let $A = [\bar{b}]_{F_0}^{\text{ELA}}$. Then $A \triangleleft F_0$, and since $F_0 \triangleleft F_1$ we also have $A \triangleleft F_1$. By the general Fraïssé construction, or more specifically by [3, Proposition 6.9], F_0 and F_1 are both isomorphic over A to its strong exponential algebraic closure A^\sim , and so in particular to each other. The same holds for any finite tuple \bar{b} , so it follows that the inclusion $F_0 \hookrightarrow F_1$ is an $L_{\infty,\omega}$ -embedding. Now we can iterate the construction to get a chain

$$F_0 \hookrightarrow F_1 \hookrightarrow F_2 \hookrightarrow \dots \hookrightarrow F_\alpha \hookrightarrow \dots$$

of length ω_1 . The union F_{ω_1} of this chain will be $L_{\infty,\omega}$ -equivalent to each element of the chain and hence to B_0 , but $\{a \in F_{\omega_1} \mid e^a = a\}$ has cardinality \aleph_1 , and each such element is exponentially algebraic, so the countable closure property fails. \square

We say that an embedding $F_1 \hookrightarrow F_2$ of pseudo-exponential fields is a *closed embedding* if and only if the image of F_1 is exponentially algebraically closed in F_2 , that is, $\text{ecl}^{F_2}(F_1) = F_1$. The category of all pseudo-exponential fields together with closed embeddings forms an AEC. The notion of an AEC being *finitary* was introduced by Hyttinen and Kesälä [1] and studied also by Kueker [5]. Using [5, Definition 3.1], an AEC \mathcal{K} is finitary if and only if whenever $M, N \in \mathcal{K}$ and $f : M \hookrightarrow N$ is an L -embedding such that, for every finite tuple \bar{a} from M , there is a \mathcal{K} -embedding $g : M \rightarrow N$ such that $g(\bar{a}) = f(\bar{a})$, then f is a \mathcal{K} -embedding.

In the above proof we showed that the inclusion map $F_0 \hookrightarrow F_1$ satisfied this last condition, but clearly it is not a closed embedding. Thus Theorem 2 is established. The proposition shows that our AEC is not closed under $L_{\omega_1,\omega}$ -equivalence, and one could use Kueker’s result that finitary AECs are closed under $L_{\omega_1,\omega}$ -equivalence to give another, less direct, proof of Theorem 2.

3 The First-Order Theory and Elementary Embeddings

3.1 The exponentially transcendental type Let f_1, \dots, f_n be exponential polynomials in variables x_1, \dots, x_n , and, as before, write $\chi_f(\bar{x})$ for the first-order formula corresponding to the Khovanskii system on the f_i , now suppressing the variables \bar{z} corresponding to coefficients of the f_i .

Given any exponential field F , and any set A of parameters from F , the exponentially transcendental type over A is the set of formulas

$$(\forall x_2, \dots, x_n) [\neg \chi_f(x, x_2, \dots, x_n)]$$

where n ranges over all positive natural numbers and f ranges over all n -tuples of exponential polynomials with coefficients from A . Write $p|_A(x)$ for this type over A . For the exponential fields under consideration it is a consistent partial type. In some cases, for example, \mathbb{R}_{exp} , the type is not complete.

Proposition *In $T_{\mathbb{B}}$, for any set of parameters A , the type $p|_A(x)$ is complete.*

Proof It is enough to prove the result for finite A . So let M be an \aleph_0 -saturated model of $T_{\mathbb{B}}$, and let A be a finite subset of M . Suppose that $a, b \in M$ are each exponentially transcendental over A . Let $M_0 = \text{ecl}^M(A)$. We will set up a back-and-forth system showing that $\text{tp}(a/M_0) = \text{tp}(b/M_0)$. In particular, $\text{tp}(a/A) = \text{tp}(b/A)$.

First, note that the partial E-field extensions of M_0 generated by a and b are isomorphic, since we have $\text{td}(a, e^a/M_0) = \text{td}(b, e^b/M_0) = 2$. Furthermore, since a and b are exponentially transcendental over M_0 , these partial E-fields are strong in M . By Proposition (B) in Section 2.6, there is an isomorphism

$$\theta_1 : \langle M_0, a \rangle_M^{\text{ELA}} \cong \langle M_0, b \rangle_M^{\text{ELA}}$$

between the ELA-subfields of M generated by $M_0 \cup \{a\}$ and $M_0 \cup \{b\}$, fixing M_0 and sending a to b . Furthermore, these ELA-subfields of M are strong in M .

Now suppose that we have n -tuples \bar{a} and \bar{b} in M , each \mathbb{Q} -linearly independent over M_0 , with $a_1 = a$ and $b_1 = b$ and an isomorphism

$$\theta_n : [M_0, \bar{a}]_M^{\text{ELA}} \cong [M_0, \bar{b}]_M^{\text{ELA}}$$

between the strong ELA-subfields of M generated by $M_0 \cup \bar{a}$, and $M_0 \cup \bar{b}$, fixing M_0 and sending a_i to b_i for $i = 1, \dots, n$.

Let $c \in M$. We want to find $d \in M$ and an isomorphism

$$\theta_{n+1} : [M_0, \bar{a}, c]_M^{\text{ELA}} \cong [M_0, \bar{b}, d]_M^{\text{ELA}}$$

extending θ_n . Write $F = [M_0, \bar{a}]_M^{\text{ELA}}$. By extending \bar{a} and \bar{b} , we may assume $M_0 \cup \bar{a} \triangleleft M$, so $F = \langle M_0, \bar{a} \rangle_M^{\text{ELA}}$, and similarly $F' := \langle M_0, \bar{b} \rangle_M^{\text{ELA}} \triangleleft M$.

If $c \in F$, then take $d = \theta_n(c)$. If c is exponentially transcendental over F , then, using \aleph_0 -saturation, we can find d exponentially transcendental over $A \cup \bar{b}$, which is therefore exponentially transcendental over $F' = [M_0, \bar{b}]_M^{\text{ELA}}$, and the same argument as above gives us θ_{n+1} .

Otherwise, since $F \triangleleft M$, there is a finite tuple \bar{c} of shortest length extending c such that $\delta(\bar{c}/F) = 0$. By Proposition (B) in Section 2.6, \bar{c} generates an ELA-extension of the form $F \triangleleft F|V$ where V is the locus of $(\bar{c}, e^{\bar{c}})$ over F , and V is rotund, additively and multiplicatively free, and of dimension equal to the length of the tuple \bar{c} . Let α be a finite tuple of parameters from F over which V is defined, and consider the variety V' defined by the same formula as V but with parameters $\beta = \theta_n(\alpha)$. Without loss of generality we may assume that \bar{b} is \mathbb{Q} -linearly independent over M_0 .

M was chosen to be an \aleph_0 -saturated model of $T_{\mathbb{B}}$. As such, we do not know that M is strongly exponentially algebraically closed, because we do not know that axiom 4 is unconditionally first-order. However, M does satisfy the axiom of exponential-algebraic closedness (see [4, Section 5.5]), which in combination with the \aleph_0 -saturation gives us $(\bar{d}, e^{\bar{d}}) \in V'(M)$, generic in V' over $\beta \cup \bar{b}$, and such that no \mathbb{Q} -linear combination of \bar{d} and \bar{b} is exponentially algebraic over A , because these conditions can be expressed by a partial type over the finite set $\beta \cup \bar{b} \cup A$.

Then \bar{d} is \mathbb{Q} -linearly independent over $M_0 \cup \bar{b}$, so $(\bar{d}, e^{\bar{d}})$ is generic in V' over $M_0 \cup \bar{b} \cup e^{\bar{b}}$, and, using Proposition (B) in Section 2.6 once more, $(\bar{d}, e^{\bar{d}})$ is generic in V' over F' . Hence the ELA-extension of F' generated by \bar{d} is of the form $F'|V'$, and so we have an isomorphism θ_{n+1} extending θ_n , as required. \square

3.2 Nonisolation of the exponentially transcendental type

Proposition *If $M \models T_{\mathbb{B}}$ and A is any set of parameters from M , then the type $p|_A(x)$ is not isolated.*

Proof By completeness of $p|_A(x)$ and the compactness theorem, if it were isolated by some formula, then it would be isolated by a finite subtype of the form

$$q(x, \bar{a}) = \bigwedge_{i=1}^r (\forall x_2, \dots, x_{n_i}) [\neg \chi_{f_i}(x, x_2, \dots, x_{n_i}, \bar{a})],$$

where \bar{a} is the finite tuple from A consisting of all coefficients from the exponential polynomials in the f_i . Let $N = \max\{n_i \mid i = 1, \dots, r\}$.

Define $e_0(x) = x$ and $e_{n+1}(x) = \exp(e_n(x))$ for $n \in \mathbb{N}$. Then in fact for any tuple \bar{b} in \mathbb{B} , there is $c \in \mathbb{B}$ such that $e_{N+1}(c) = c$ and $c, e_1(c), \dots, e_N(c)$ are algebraically independent over \bar{b} , by strong exponential-algebraic closedness. Then $\mathbb{B} \models q(c, \bar{b})$ because any tuple witnessing that c is exponentially algebraic must span the $(N + 1)$ -dimensional \mathbb{Q} -vector space spanned by $\{e_n(c) \mid n = 0, \dots, N\}$, but the Khovanskii systems in q look only at tuples of length up to N . Hence

$$T_{\mathbb{B}} \vdash \forall \bar{y} \exists x [q(x, \bar{y}) \wedge e_{N+1}(x) = x].$$

So there is $c \in M$ with $M \models q(c, \bar{a})$ but with c exponentially algebraic over \bar{a} . Hence $p|_A(x)$ is not isolated. \square

3.3 Isolated types Let $F \models \Psi$, and let X be an exponential transcendence base for F . Any element of \mathbb{Z} lies in the definable closure of \emptyset , and the kernel generator τ is model-theoretically algebraic, with $-\tau$ being the only conjugate. So the kernel of F is model-theoretically algebraic. Now let \bar{a} be an n -tuple from F , and let X_0 be the smallest subset of X such that $\bar{a} \in \text{ecl}(X_0)$. Let \bar{b} be an r -tuple which is a \mathbb{Q} -linear basis for $D([\!| X_0, \tau, \bar{a}]\!]_F)$ over $X_0 \cup \{\tau\}$, let $V = \text{Loc}(\bar{b}, e^{\bar{b}}/X_0, \tau)$, and let $M \in \text{Mat}_{n \times r}(\mathbb{Q})$ be such that $\bar{a} = M\bar{b}$. Consider the first-order formula $\varphi(\bar{x})$ given by

$$\exists \bar{y} [(\bar{y}, e^{\bar{y}}) \in V \wedge \bar{y} \text{ is } Q\text{-linearly independent over } X_0 \wedge \bar{x} = M\bar{y}],$$

where here Q is the field of fractions of the definable subring Z , not the quantifier. Since F has standard integers, $Q(F)$ will be \mathbb{Q} . Now if $F \models \varphi(\bar{a}')$ and \bar{b}' is a witness for \bar{y} , then $\text{ldim}_{\mathbb{Q}}(\bar{b}'/X_0, \tau) = r$, but $\dim V = r$ and $\delta(\bar{b}'/X_0) = 0$, so $(\bar{b}', e^{\bar{b}'})$ is generic in V over $X_0 \cup \{\tau\}$. Hence $\bar{b}' \in \text{ecl}(X_0)$, and, by Proposition (A) in Section 2.6, there is an isomorphism between $\langle X_0, \bar{b} \rangle_F^{\text{ELA}}$ and $\langle X_0, \bar{b}' \rangle_F^{\text{ELA}}$, preserving X_0 and τ and sending \bar{b} to \bar{b}' . Since $X \setminus X_0$ is exponentially algebraically independent over $\text{ecl}^F(X_0)$, this extends to an isomorphism between $\langle X_1, \bar{b} \rangle_F^{\text{ELA}}$ and $\langle X_1, \bar{b}' \rangle_F^{\text{ELA}}$ for any countable subset X_1 of X . Then by strong exponential-algebraic closedness of F , \bar{b} and \bar{b}' are back-and-forth equivalent over any such X_1 and hence have the same first-order type over X . Hence \bar{a} and \bar{a}' have the same type over X . Thus the formula φ isolates the first-order type of \bar{a} over X .

3.4 Elementary equivalence We can now prove the second part of Theorem 3, showing that all the B_n for $n \in \mathbb{N}$ are elementarily equivalent to \mathbb{B} . Let \bar{a} be an exponentially algebraically independent n -tuple in \mathbb{B} , and add parameters for \bar{a} . The type of

a nonstandard integer is a nonisolated (partial) type, and, by the above, the exponentially transcendental type over \bar{a} is also nonisolated. Hence, by the omitting types theorem, there is a countable model M of $T_{\mathbb{B}}$ containing \bar{a} which omits both types. Then $M \models \Psi_n$, and by countable categoricity of Ψ_n , M is isomorphic to B_n . Hence $B_n \models T_{\mathbb{B}}$.

3.5 Atomic models and proof of Theorem 4 From Section 3.3 above, the first-order type over an exponential transcendence base of any finite tuple from any model $F \models \Psi$ is isolated by a single first-order formula in the language $\langle +, \cdot, \exp \rangle$, so each such F is an atomic model of the expansion of $T_{\mathbb{B}}$ by parameters for the exponential transcendence base. In particular, B_0 is atomic over \emptyset , and a countable atomic model is prime, so B_0 is the prime model of $T_{\mathbb{B}}$. For the converse statement, any model of $T_{\mathbb{B}}$ which is atomic over an exponential transcendence base has standard integers and hence is a model of Ψ , that is, of axioms 1–4. That completes the proof of Theorem 4.

3.6 Elementary embeddings and the end of the proof of Theorem 3 If an embedding $F_1 \subseteq F_2$ of models of Ψ is not strong, then there are finite tuples $\bar{a} \triangleleft F_1$ and $\bar{b} \in F_2^n$ such that \bar{b} is \mathbb{Q} -linearly independent over \bar{a} and $\delta(\bar{b}/\bar{a}) < 0$, so $(\bar{b}, e^{\bar{b}})$ lies in some algebraic variety V defined over $(\bar{a}, e^{\bar{a}})$ of dimension less than n . The first-order formula expressing the existence of such a \bar{b} is true of \bar{a} in F_2 but false in F_1 , and hence the inclusion of F_1 in F_2 is not elementary.

Now suppose that $F_1 \triangleleft F_2$ is a strong extension of models of Ψ . We use the Tarski–Vaught test to show that $F_1 \preceq F_2$. So let \bar{a} be a tuple from F_1 , $b \in F_2$, and let $\varphi(\bar{x}, y)$ be a first-order formula such that $F_2 \models \varphi(\bar{a}, b)$. Extending \bar{a} if necessary, we may assume that $\bar{a} \triangleleft F_1$, and so $\bar{a} \triangleleft F_2$. Since the kernel does not extend, the ELA-subfield F_0 of F_2 generated by \bar{a} is contained in F_1 . If b is exponentially algebraic over \bar{a} , then it generates a strong ELA-extension of the form $F_0 \triangleleft F_0|V$ for some perfectly rotund V . There is an isomorphic strong ELA-extension inside F_1 by strong exponential-algebraic closedness of F_1 , and the element c corresponding to b under the isomorphism realizes the same principal formula as b in F_2 . In particular, $F_2 \models \varphi(\bar{a}, c)$. Otherwise b is exponentially transcendental over \bar{a} , and, since the exponentially transcendental type is approximated by isolated exponentially algebraic types as in Section 3.2 above, we can again find such a c in F_1 , depending on the formula φ . So $F_1 \preceq F_2$. That completes the proof of Theorem 3.

References

- [1] Hyttinen, T., and M. Kesälä, “Independence in finitary abstract elementary classes,” *Annals of Pure and Applied Logic*, vol. 143 (2006), pp. 103–38. [Zbl 1112.03026](#). [MR 2258625](#). [DOI 10.1016/j.apal.2006.01.009](#). [510, 516](#)
- [2] Kirby, J., “Exponential algebraicity in exponential fields,” *Bulletin of the London Mathematical Society*, vol. 42 (2010), pp. 879–90. [Zbl 1203.03050](#). [MR 2721747](#). [DOI 10.1112/blms/bdq044](#). [513, 514](#)
- [3] Kirby, J., “Finitely presented exponential fields,” to appear in *Algebra and Number Theory*, preprint, arXiv:0912.4019v4 [math.LO]. [510, 514, 515, 516](#)
- [4] Kirby, J., and B. Zilber, “Exponential fields and atypical intersections,” preprint, arXiv:1108.1075v1 [math.LO]. [509, 510, 513, 514, 515, 517](#)

- [5] Kueker, D. W., “Abstract elementary classes and infinitary logics,” *Annals of Pure and Applied Logic*, vol. 156 (2008), pp. 274–86. [Zbl 1155.03016](#). [MR 2484485](#). [DOI 10.1016/j.apal.2008.07.002](#). [516](#)
- [6] Macintyre, A. J., “Exponential algebra,” pp. 191–210 in *Logic and Algebra (Pontignano, 1994)*, vol. 180 of *Lecture Notes in Pure and Applied Mathematics*, Dekker, New York, 1996. [MR 1404940](#). [513](#)
- [7] Zilber, B., “Pseudo-exponentiation on algebraically closed fields of characteristic zero,” *Annals of Pure and Applied Logic*, vol. 132 (2005), pp. 67–95. [Zbl 1076.03024](#). [MR 2102856](#). [DOI 10.1016/j.apal.2004.07.001](#). [509](#), [510](#), [512](#), [513](#), [514](#), [515](#)
- [8] Zilber, B., “Covers of the multiplicative group of an algebraically closed field of characteristic zero,” *Journal of the London Mathematical Society (2)*, vol. 74 (2006), pp. 41–58. [Zbl 1104.03030](#). [MR 2254551](#). [DOI 10.1112/S0024610706023349](#). [515](#)

Acknowledgment

Some of the work for this paper was done while I was visiting the University of Helsinki, and I am grateful for their hospitality.

School of Mathematics
University of East Anglia
Norwich Research Park
Norwich NR4 7TJ
UK
jonathan.kirby@uea.ac.uk