

Banach J. Math. Anal. 10 (2016), no. 1, 169–208

<http://dx.doi.org/10.1215/17358787-3345137>

ISSN: 1735-8787 (electronic)

<http://projecteuclid.org/bjma>

## SHARP EXTENSIONS AND ALGEBRAIC PROPERTIES FOR SOLUTION FAMILIES OF VECTOR-VALUED DIFFERENTIAL EQUATIONS

LUCIANO ABADIAS,<sup>1</sup> CARLOS LIZAMA,<sup>2\*</sup> and PEDRO J. MIANA<sup>1</sup>

Communicated by L. P. Castro

ABSTRACT. In this paper we show the unexpected property that extension from local to global without loss of regularity holds for the solutions of a wide class of vector-valued differential equations, in particular for the class of fractional abstract Cauchy problems in the subdiffusive case. The main technique is the use of the algebraic structure of these solutions, which are defined by new versions of functional equations defining solution families of bounded operators. The convolution product and the double Laplace transform for functions of two variables are useful tools which we apply also to extend these solutions. Finally we illustrate our results with different concrete examples.

### 1. INTRODUCTION

Let  $A$  be a closed linear operator with domain  $D(A)$  defined in a complex Banach space  $X$ , and let  $0 < \tau \leq \infty$ . Suppose that  $A$  is the generator of a *local*  $C_0$ -semigroup  $\{T(t)\}_{t \in [0, \tau)}$  or, equivalently, that the first-order Cauchy problem

$$\begin{cases} u'(t) = Au(t) + x, & 0 \leq t < \tau, \\ u(0) = 0 \end{cases} \quad (1.1)$$

Copyright 2016 by the Tusi Mathematical Research Group.

Received Mar. 11, 2015; Accepted May 10, 2015.

\*Corresponding author.

2010 *Mathematics Subject Classification.* Primary 47D06; Secondary 44A10, 47D60, 44A35.

*Keywords.* abstract Cauchy problem, time translation, vector-valued solutions,  $(a, k)$ -regularized resolvent families, Laplace transform.

has a unique solution  $u \in C^1([0, \tau], X) \cap C([0, \tau], D(A))$  (i.e., it is *locally well-posed*). Then it is well known that  $A$  is the generator of a *global*  $C_0$ -semigroup  $\{T(t)\}_{t \geq 0}$  (i.e., the problem is *globally well posed*; see [2, Theorem 1.2], [1, Section 3.1], and also [29]). We observe that this dynamic behavior of the solution for the Cauchy problem (1.1) (i.e., the extension property from local to global without loss of regularity) heavily depends on the *translation in time property* of the Cauchy's functional equation, namely  $T(t+s) = T(t)T(s)$ ,  $t, s \geq 0$ . In contrast, this extension property is not more true for the class of local integrated semigroups (see [2, Example 4.6]). Furthermore, we have that if  $A$  generates a local  $(1*k)$ -convoluted semigroup on  $[0, \tau)$ , then  $A$  generates a local  $(1*k^{*n})$ -convoluted semigroup on an interval  $[0, n\tau)$  (see [8, Theorem 4.4], [23, Theorem 3.3]). In other words, in these cases there is evolution with jumps of regularity, and naturally the need of regularize the family of operators appears (in the sense of convolution) in order to have extension. In all of these cases, the property of translation in time of the associated functional equation is strongly connected with the problem of extension from a short to a long interval of definition for the corresponding family of operators.

Fractional diffusion equations are widely used to describe anomalous diffusion processes. From the point of view of operator-theoretical methods for partial differential equations, subdiffusion phenomena is modeled naturally by means of fractional Cauchy problems in the form

$$\begin{cases} D_t^\alpha u(t) = Au(t) + x, & 0 \leq t < \tau, \\ u(0) = 0, \end{cases} \quad (1.2)$$

where  $0 < \alpha < 1$  and the fractional derivative is taken in the Caputo sense (see [3], [24]). In [14] the existence of solutions of fractional Cauchy problems is studied in detail; it is also studied in [12] for the superdiffusive case  $1 < \alpha < 2$ . Suppose that  $A$  generates a local 1-parameter family of bounded operators that makes the equation (1.2) locally well posed. The natural question that arises is: Can (1.2) be globally well posed?

We point out that (1.2) is included in the more general Volterra-type equation

$$u(t) = k(t)x + A \int_0^t a(t-s)u(s) ds, \quad t \in (0, \tau), \quad (1.3)$$

for the special choice of kernel  $a(t) = k(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$ , for  $\alpha, t > 0$  and where  $\Gamma$  is the Euler gamma function. This Volterra-type equation in case  $k(t) \equiv 1$  has been deeply treated in the monograph [25] by Prüss. Further relevant studies have been done in the monographs [10] and [11] by Kostić. (See also [7]–[9], [15]–[18], and [23] for related work.) Therefore, we can set our problem in a more general context by classifying the classes of pairs  $(a, k)$  where extension of (1.3) from local to global and without loss of regularity holds.

We note that under certain conditions on the scalar-valued kernels  $a$  and  $k$ , *well-posedness* of the Volterra equation (1.3) is equivalent to the existence of a 1-parameter and strongly continuous family of bounded operators  $\{S(t)\}_{t > 0}$  that

satisfies a functional equation in the form

$$\begin{aligned} S(s) \int_0^t a(t-\tau)S(\tau) d\tau - S(t) \int_0^s a(s-\tau)S(\tau) d\tau \\ = k(s) \int_0^t a(t-\tau)S(\tau) d\tau - k(t) \int_0^s a(s-\tau)S(\tau) d\tau, \end{aligned} \quad (1.4)$$

for  $t, s > 0$  (see [17, Theorem 3.1]).

Explicitly, in the present article we will study the following questions.

(Q1) (Evolution with or without jumps of regularity; Sections 4 and 5). If  $A$  is the generator of a local regularized resolvent family on the interval  $(0, \tau)$ , is  $A$  also the generator of an local extended regularized resolvent family on the interval  $(0, (n+1)\tau)$  for  $n \in \mathbb{N}$ ? And in particular, for which class of pairs of kernels  $(a, k)$  do we have that if  $A$  is the generator of a local  $(a, k)$ -regularized resolvent family, then  $A$  is the generator of a global  $(a, k)$ -regularized resolvent family?

(Q2) (Time translation; Section 6). Determine the class of pairs  $(a, k)$  for which it is possible to find an equivalent and explicit expression for equation (1.4) in terms of the sum  $t + s$  instead of  $t$  and  $s$ .

We will first answer globally the problem of evolution (Q1), that is, the possibility to extend the family of operators  $S(t)$  from the interval  $(0, \tau)$  to the whole semiaxis  $(0, \infty)$ . More precisely, we prove the following: if  $A$  is the generator of a local  $(a, k)$ -regularized resolvent family on  $(0, \tau)$ , then  $A$  is the generator of a local  $(a, (a * k)^{*n} * k)$ -regularized resolvent family in  $(0, (n+1)\tau)$  (see Theorem 4.3). We note that this problem, which has been studied in a series of articles in the last few years, is settled here in a simple way, making transparent the process of regularization needed in each step of the extension. In particular, our result intersects the papers [2], [8], and [23] where the problem of extension for local integrated semigroups, local convoluted semigroups, and local convoluted cosine functions is studied, respectively. The question about extension for local convoluted semigroups and local convoluted cosine functions, which is resolved in [8] and [23], respectively, had been cited previously in the paragraph directly preceding [10, Theorem 1.2.7]. Results related to the extension of local  $C$ -regularized semigroups and local  $C$ -regularized cosine functions appeared for the first time in [29].

However, note that if we restrict this result to the  $C_0$ -semigroup family, we do not obtain evolution-conserving regularity. Under some conditions on  $a$  and  $k$ , we improve this result and obtain an extension without a jump of regularity in Theorem 5.1. We use this theorem to prove one of the main results in this paper concerning the fractional equation (1.2): if  $A$  is the generator of a local  $(g_\alpha, g_\alpha)$ -regularized resolvent family on  $[0, \tau)$ , with  $0 < \alpha < 1$ , then  $A$  is the generator of a global  $(g_\alpha, g_\alpha)$ -regularized resolvent family in  $[0, \infty)$  (see Corollary 5.2). These results recover and widely extend the property of evolution without jumps of regularity from the case of the solutions of first-order Cauchy problems to the case of fractional subdiffusive models.

In this paper, we are able to completely solve (Q2) by establishing an equivalent functional equation to (1.4), which defines global  $(a, k)$ -regularized resolvent families, in the following form:

$$\begin{aligned} & \int_t^{t+s} \int_0^r k(s+t-r)a(r-\tau)S(\tau) d\tau dr \\ & \quad - \int_0^s \int_0^r k(s+t-r)a(r-\tau)S(\tau) d\tau dr \\ & = \int_0^t \int_0^s a(t+s-r_1-r_2)S(r_1)S(r_2) dr_1 dr_2, \end{aligned} \tag{1.5}$$

for  $t, s \geq 0$  (see Theorem 6.1). The above formula widely solves the problem of time translation, not only extending all the several results existing in the literature, but also proposing and finding a better expression for older and new cases. For example, we will see that for  $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  and  $k(t) = 1$ , for  $t > 0$ , the border cases  $\alpha = 1$  and  $\alpha = 2$  are naturally included in our functional formula, unifying the cases  $0 < \alpha < 1$  and  $1 < \alpha < 2$  mentioned previously. We point out that functional equation (1.5) inspired the way to define extensions in local  $(a, k)$ -regularized resolvent family mentioned previously.

In the last few years, special interest has developed in the study of algebraic function equations (Q2) for  $(a, k)$ -regularized resolvent family only with  $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  for  $t > 0$  (due mainly for its connection with fractional differential equations). First results in this direction appeared in [13] where an equivalent functional equation to (1.4) in the case  $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  for  $0 < \alpha < 1$ , and  $k(t) = 1$ , ( $t > 0$ ), is given in [13, Formula (2.1)]. After that, a similar result for the case  $a(t) = k(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  is shown in [20, Proposition 2.2]. In the recent article [15], a further extension of a known result in the case  $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  and  $k(t) = \frac{t^\beta}{\Gamma(\beta+1)}$  was proved (see [15, Theorem 5]). We note that some restriction should be imposed for  $\alpha \geq 1$  (see Example 6.5). A further generalization, this time in case  $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  for  $0 < \alpha < 1$  and  $k(t) = \int_0^t K(s) ds$ , was successfully obtained in [18, Theorem 8]. Finally, we point out that in a very recent work [19], the authors discovered a functional equation which covers the case  $a(t) = \frac{t^{\alpha-1}}{\Gamma(\alpha)}$  and  $k(t) = 1$  ( $t > 0$ ) in the super diffusive case  $1 < \alpha < 2$  (see [19, Definition 3.1]).

To handle both questions (Q1), (Q2), we introduce an original technique in the context of scalar and vector-valued functions of two variables. In Section 2, we work with convolution product  $*_2$  (see formula (2.2)), and we prove some needed technical results that play a key role here. In Section 3, we see results about simple and double Laplace transform, properties that this transform verifies and in which appear the above convolutions products (see, for example, Theorem 3.4).

The double Laplace transform is an efficient and known tool to solve scalar differential equations in two variables (see, e.g., [5, Chapitre IV.15.3], [6], [26, pp. 226–228]). Other interesting applications of the double Laplace transform is to supply integral formulas (see [5, Chapitre IV.15.1]) and bilinear expansions (see

[5, Chapitre IV.15.2]). In [27], the structure of closed ideals of the convolution algebra  $L^1(\mathbb{R}^n)$  is studied and the Laplace transform for functions of several variables is also considered.

Finally, in Section 7, we illustrate our main results with some particular examples in which we consider some  $(a, k)$ -regularized resolvent families, some of which are related with the solution of fractional Cauchy problems, to conclude that the extension process is possible with or without jumps of regularity. We also give new functional equations obtained as a consequence of the formula (1.5) for known operators of the 1-parameter family:  $C_0$ -semigroups, cosine families, convoluted semigroups, and resolvent families.

## 2. CONVOLUTION PRODUCTS IN ONE AND TWO VARIABLES

In this section, we state some technical results for convolution products (in one and two variables) that we use to prove the relevant results in the following sections. The convolution product in several variables have been considered in some relevant fields in mathematical analysis (see, e.g., [27]). However, the convolution product in two variables is a new tool to apply to  $(a, k)$ -regularized resolvent families.

We denote  $\mathbb{R}_+ = [0, +\infty)$ ;  $\mathbb{R} = (-\infty, +\infty)$ ,  $\mathbb{R}_+^2 = \mathbb{R}_+ \times \mathbb{R}_+$  and  $\mathbb{R}^2 = \mathbb{R} \times \mathbb{R}$ . We consider the space of locally integrable functions in one and two variables,  $L_{\text{loc}}^1(\mathbb{R}_+)$ , and  $L_{\text{loc}}^1(\mathbb{R}_+^2)$ . The space  $\mathcal{C}^n(\mathbb{R}_+)$  is formed with continuous functions  $f : \mathbb{R}_+ \rightarrow \mathbb{C}$  such that  $f^{(j)}$  is continuous for  $0 \leq j \leq n$  for  $n \geq 0$ . Some of the above spaces will also be considered in  $(0, \tau)$  or  $[0, \tau)$  instead of  $\mathbb{R}_+$ , with  $\tau > 0$ .

Let  $f, g : \mathbb{R}_+ \rightarrow \mathbb{C}$ , we write  $f_t(s) := f(s+t)\chi_{[-s, +\infty)}(t)$  for  $t \in \mathbb{R}$ ;  $f^+ : \mathbb{R}_+^2 \rightarrow \mathbb{C}$  the function given by  $f^+(t, s) := f(t+s)$ ;  $f^- : \mathbb{R}^2 \rightarrow \mathbb{C}$  for the function given by  $f^-(t, s) := f(|t-s|)$ ;  $f \otimes g : \mathbb{R}_+^2 \rightarrow \mathbb{C}$  by  $f \otimes g(t, s) := f(t)g(s)$  for  $(t, s) \in \mathbb{R}_+^2$ ; and

$$f * g(t) = \int_0^t f(t-s)g(s) ds, \quad t > 0, \quad (2.1)$$

the usual convolution product, for the functions  $f, g$  where the product is convergent. We write  $f^{*2}$  instead of  $f * f$  and then  $f^{*n} = f * (f^{*(n-1)})$  for the case where  $n \geq 2$  is the  $n$ -fold convolution power of  $f$ .

For  $F, G : \mathbb{R}_+^2 \rightarrow \mathbb{C}$  given, we define the convolution product in two variables by

$$F *_2 G(t, s) := \int_0^t \int_0^s F(t-u, s-v)G(u, v) dv du, \quad t, s > 0, \quad (2.2)$$

whenever it is well defined. This product is commutative and associative (see [5, Formula (13.9)] and [26, Formula (3-18-19)]).

We define functions  $g_\alpha(t) := \frac{t^{\alpha-1}}{\Gamma(\alpha)}$ ,  $e_\lambda(t) := e^{-\lambda t}$ , and  $e_{\lambda, \mu}(t, s) := e^{-\lambda t - \mu s} = e_\lambda \otimes e_\mu(t, s)$  for  $\alpha \in \mathbb{R} \setminus \{0, -1, -2, -3, \dots\}$ ,  $\lambda, \mu \in \mathbb{C}$  and  $t, s > 0$ . Note that  $(e_\lambda)^+ = e_{\lambda, \lambda}$  for  $\lambda \in \mathbb{C}$ . It is direct to check the following well-known identities:

$$g_\alpha * g_\beta = g_{\alpha+\beta}, \quad \alpha, \beta > 0;$$

$$e_\lambda * e_\mu = \frac{1}{\lambda - \mu}(e_\mu - e_\lambda), \quad \lambda \neq \mu;$$

$$e_{\lambda, \lambda'} *_2 e_{\mu, \mu'} = \frac{1}{(\lambda - \mu)(\lambda' - \mu')}(e_{\lambda, \lambda'} - e_{\lambda, \mu'} - e_{\mu, \lambda'} + e_{\mu, \mu'}), \quad \lambda \neq \mu, \lambda' \neq \mu'.$$

The way that  $*$  and  $*_2$  interact with operators  $\otimes$ ,  $(\cdot)_t$ , and  $(\cdot)^+$  is shown in the next theorem.

**Theorem 2.1.** *Let  $f, g, h, j \in L^1_{\text{loc}}(\mathbb{R}_+)$ . Then we have the following:*

- (i)  $(f \otimes g) *_2 (h \otimes j) = (f * h) \otimes (g * j)$ ,
- (ii)  $(g^+ *_2 (f \otimes h))(t, s) = h * (f * g)_t(s) - f_t * (h * g)(s)$ , for  $t, s \geq 0$ ,
- (iii)

$$(f^+ *_2 g^+)(t, s) = \begin{cases} (f_t * \mathcal{M}(g))(s) + s(f_s * g_s)(t - s) + (\mathcal{M}(f) * g_t)(s), & 0 \leq s \leq t, \\ (f_s * \mathcal{M}(g))(t) + t(f_t * g_t)(s - t) + (\mathcal{M}(f) * g_s)(t), & 0 \leq t \leq s, \end{cases}$$

where  $\mathcal{M}(g)(s) := sg(s)$  for  $s \in \mathbb{R}_+$ .

*Proof.* The proof of part (i) is straightforward. To show (ii), note that if  $g \in L^1_{\text{loc}}(\mathbb{R}_+)$ , then  $g^+ \in L^1_{\text{loc}}(\mathbb{R}_+^2)$ . We change variables to obtain the following equalities:

$$\begin{aligned} & \int_0^t \int_0^s g(t + s - r_1 - r_2) f(r_1) h(r_2) dr_1 dr_2 \\ &= \int_0^t \int_0^s g(v + z) f(t - v) h(s - z) dv dz \\ &= \int_0^s h(s - z) \int_z^{t+z} f(t + z - u) g(u) du dz \\ &= \int_0^s h(s - z) \int_0^{t+z} f(t + z - u) g(u) du dz \\ &\quad - \int_0^s h(s - z) \int_0^z f(t + z - u) g(u) du dz. \end{aligned}$$

Now, we apply Fubini theorem and change of variable  $s - z = r - u$  to get

$$\begin{aligned} \int_0^s h(s - z) \int_0^z f(t + z - u) g(u) du dz &= \int_0^s g(u) \int_u^s f(t + z - u) h(s - z) dz du \\ &= \int_0^s g(u) \int_u^s f(t + s - r) h(r - u) dr du \\ &= \int_0^s f(t + s - r) \int_0^r h(r - u) g(u) du dr. \end{aligned}$$

We express the above integrals in terms of convolution products to conclude the claim. The proof of (iii) is similar to the proof of part (ii).  $\square$

**Corollary 2.2.** *Let  $f, g, h \in L^1_{\text{loc}}(\mathbb{R}_+)$ . Then*

- (i)  $(g^+ *_2 (f \otimes h))(t, s) = f * (h * g)_s(t) - h_s * (f * g)(t)$ , for  $t, s \geq 0$ ,

(ii)  $(g^+ *_2 (f \otimes f))(t, s) = f * (f * g)_t(s) - f_t * (f * g)(s)$ , for  $t, s \geq 0$ .

*Proof.* (i) We apply the identity  $(g^+ *_2 (f \otimes h))(t, s) = (g^+ *_2 (h \otimes f))(s, t)$  for  $(t, s) \in \mathbb{R}_+^2$  and Theorem 2.1(ii).  $\square$

The next lemma extends [8, Lemma 2.1] and will be applied several times in the present article.

**Lemma 2.3.** *Take  $0 \leq \tau \leq t$  and  $f, g, h \in L_{\text{loc}}^1(\mathbb{R}_+)$ . Then*

$$\begin{aligned} & \int_0^{t-\tau} h(t-s)(g * f)(s) ds + \int_0^\tau f(t-s)(g * h)(s) ds \\ &= (f * g * h)(t) - g^+ *_2 (f \otimes h)(t - \tau, \tau). \end{aligned}$$

*Proof.* We use Fubini's theorem and a change of variables to obtain

$$\begin{aligned} & (f * g * h)(t) - \int_0^\tau f(t-r) \int_0^r g(r-s)h(s) ds dr \\ &= \int_0^t f(r) \int_0^{t-r} g(t-r-s)h(s) ds dr - \int_{t-\tau}^t f(r) \int_0^{t-r} g(t-r-s)h(s) ds dr \\ &= \int_0^{t-\tau} f(r) \int_0^{t-r} g(t-r-s)h(s) ds dr \\ &= \int_0^\tau h(s) \int_0^{t-\tau} g(t-s-r)f(r) dr ds + \int_\tau^t h(s) \int_0^{t-s} g(t-s-r)f(r) dr ds \\ &= \int_0^\tau \int_0^{t-\tau} g(t-s-r)h(s)f(r) dr ds + \int_0^{t-\tau} h(t-s)(g * f)(s) ds, \end{aligned}$$

for  $t \in \mathbb{R}_+$ . This proves the claim.  $\square$

*Remark 2.4.* Let  $X$  be a Banach space, while

$$L_{\text{loc}}^1(\mathbb{R}_+, X) := \{f : \mathbb{R}_+ \rightarrow X : f \text{ is a Bochner integrable on } [0, \tau] \text{ for all } \tau > 0\}.$$

We also consider  $L_{\text{loc}}^1(\mathbb{R}_+^2, X)$  for functions defined in two variables. The definitions of  $*$  and  $*_2$ , (see (2.1) and (2.2)), Theorem 2.1, Corollary 2.2, and Lemma 2.3 hold in the case that one function is vector-valued into  $X$ . The proof of these analogous results involves the ideas already employed in the scalar case.

### 3. LAPLACE TRANSFORM IN ONE AND TWO VARIABLES

In this section, we study properties concerned with the Laplace transform for functions in the above spaces. We say that  $f \in L_{\text{loc}}^1(\mathbb{R}_+, X)$  is a Laplace-transformable function if there exists  $\omega_f \in \mathbb{R}$  such that the usual Laplace transform

$$\hat{f}(\lambda) := \int_0^\infty e^{-\lambda t} f(t) dt = \lim_{\tau \rightarrow \infty} \int_0^\tau e^{-\lambda t} f(t) dt, \quad \Re \lambda > \omega_f,$$

is well defined (see, e.g., [1, Section 1.4]). Let  $f : \mathbb{R}^+ \rightarrow X$  be absolutely continuous and differentiable almost everywhere. Note that in the scalar case  $X = \mathbb{C}$  any absolutely continuous function defined for  $t \geq 0$  ( $t > 0$ ) is differentiable almost

everywhere  $t \geq 0$  ( $t > 0$ ) because the space  $\mathbb{C}$  has the Radon Nykodim property. If  $\Re\lambda > 0$  and  $\hat{f}'(\lambda)$  exist, then  $\hat{f}(\lambda)$  exists and

$$\hat{f}'(\lambda) = \lambda\hat{f}(\lambda) - f(0) \quad (3.1)$$

(see [1, Corollary 1.6.6]).

Similarly, we say that  $F \in L_{\text{loc}}^1(\mathbb{R}_+^2, X)$  is a *double Laplace-transformable function* (or *2-Laplace-transformable*) if there exist  $\omega_{1,F}, \omega_{2,F} \in \mathbb{R}$  such that

$$\mathcal{L}_2(F)(\lambda, \mu) := \int_0^\infty \int_0^\infty e^{-\lambda t} e^{-\mu s} F(t, s) ds dt := \lim_{\tau \rightarrow \infty} \int_0^\tau \int_0^\tau e^{-\lambda t} e^{-\mu s} F(t, s) ds dt$$

converges for  $\Re\lambda > \omega_{1,F}$  and  $\Re\mu > \omega_{2,F}$  (see [5, Chapitre IV] and [26, Section 3.18]) in the scalar case; the Laplace transform  $\mathcal{L}_2$  is commonly called the *double Laplace transform*.

For further use we establish the following theorem where we include some known identities of Laplace transforms and double Laplace transforms.

**Theorem 3.1.** *Let  $f \in L_{\text{loc}}^1(\mathbb{R}_+, X)$  and  $g \in L_{\text{loc}}^1(\mathbb{R}_+)$  be Laplace-transformable functions. Then the following identities hold:*

- (i)  $\mathcal{L}_2(f^+)(\lambda, \mu) = \frac{1}{\mu - \lambda}(\hat{f}(\lambda) - \hat{f}(\mu))$  for  $\Re\lambda, \Re\mu > \omega_f$  with  $\lambda \neq \mu$ ,
- (ii)  $\mathcal{L}_2(f^-)(\lambda, \mu) = \frac{1}{\lambda + \mu}(\hat{f}(\lambda) + \hat{f}(\mu))$  for  $\Re\lambda, \Re\mu > \omega_f$  with  $\Re(\lambda + \mu) > 0$ ,
- (iii)  $\mathcal{L}_2(f \otimes g)(\lambda, \mu) = \hat{f}(\lambda)\hat{g}(\mu)$  for  $\Re\lambda > \omega_f$  and  $\Re\mu > \omega_g$ .

Let  $F \in L_{\text{loc}}^1(\mathbb{R}_+^2, X)$  and  $G \in L_{\text{loc}}^1(\mathbb{R}_+^2)$  be double Laplace-transformable functions. Then the following identity holds:

- (iv)  $\mathcal{L}_2(F *_2 G)(\lambda, \mu) = \mathcal{L}_2(G)(\lambda, \mu)\mathcal{L}_2(F)(\lambda, \mu)$ , for  $\Re\lambda > \max(\omega_{1,F}, \omega_{1,G})$  and  $\Re\mu > \max(\omega_{2,F}, \omega_{2,G})$ .

*Proof.* The proof of (i) appears in [26, pp. 221–222]; the proof of (ii) in [26, pp. 223–224] and the equality (iii) appears in [26, (3-18-4)]. Finally the equality (iv) is straightforward and it is commented in [26, (3-18-20)].  $\square$

In what follows, given an absolutely continuous and differentiable almost everywhere function  $c : (0, \infty) \rightarrow X$ , we denote by  $c'$  its derivative and  $c(0^+) := \lim_{t \rightarrow 0^+} c(t)$ , whenever both limits exist.

**Theorem 3.2.** *Let  $c \in L_{\text{loc}}^1(\mathbb{R}_+, X)$  be an absolutely continuous on  $(0, \infty)$ , differentiable almost everywhere, and Laplace-transformable function.*

- (i) *If  $(c')^+ : \mathbb{R}_+^2 \rightarrow X$  is 2-Laplace-transformable, then*

$$\mathcal{L}_2((c')^+)(\lambda, \mu) = \frac{1}{\mu - \lambda}(\lambda\hat{c}(\lambda) - \mu\hat{c}(\mu)), \quad \Re\lambda, \Re\mu > \omega_c, \lambda \neq \mu.$$

- (ii) *If  $(c')^- : \mathbb{R}_+^2 \rightarrow X$  is 2-Laplace-transformable and  $c(0^+)$  exists, then*

$$\mathcal{L}_2((c')^-)(\lambda, \mu) = \frac{1}{\mu + \lambda}(\lambda\hat{c}(\lambda) + \mu\hat{c}(\mu)) - \frac{2c(0^+)}{\lambda + \mu}, \quad \Re\lambda, \Re\mu > \omega_c, \Re(\lambda + \mu) > 0.$$



*Proof.* (i) We integrate by parts to obtain

$$\begin{aligned} & \int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} c'(t+s) ds dt \\ &= \int_0^\infty e^{-(\lambda-\mu)t} \int_t^\infty e^{-\mu v} c'(v) dv dt \\ &= \int_0^\infty e^{-(\lambda-\mu)t} \left( -c(t)e^{-\mu t} + \mu \int_t^\infty e^{-\mu v} c(v) dv \right) dt. \end{aligned}$$

We change the inner variable to get the equality

$$\int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} c'(t+s) ds dt = -\hat{c}(\lambda) + \mu \mathcal{L}_2(c^+)(\lambda, \mu) = \frac{\lambda \hat{c}(\lambda) - \mu \hat{c}(\mu)}{\mu - \lambda},$$

for  $\Re \lambda, \Re \mu > \omega_c$ , and  $\lambda \neq \mu$ . (ii) For  $\Re \lambda, \Re \mu > \omega_c$ , and  $\Re(\lambda + \mu) > 0$ , note that

$$\begin{aligned} \mathcal{L}_2((c')^-)(\lambda, \mu) &= \int_0^\infty e^{-\lambda t} \int_{-t}^\infty e^{-\mu(v+t)} (c')^-(t, v+t) dv dt \\ &= \int_0^\infty e^{-(\lambda+\mu)t} \int_0^\infty e^{-\mu v} c'(v) dv dt \\ &\quad + \int_0^\infty e^{-(\lambda+\mu)t} \int_0^t e^{\mu v} c'(v) dv dt, \end{aligned}$$

where we have changed the inner variable. We integrate by parts to get that

$$\begin{aligned} \int_0^\infty e^{-(\lambda+\mu)t} \int_0^\infty e^{-\mu v} c'(v) dv dt &= \frac{1}{\lambda + \mu} \int_0^\infty e^{-\mu v} c'(v) dv \\ &= \frac{1}{\lambda + \mu} [-c(0^+) + \mu \hat{c}(\mu)]. \end{aligned}$$

On the other hand, we use Fubini's theorem to obtain,

$$\begin{aligned} \int_0^\infty e^{-(\lambda+\mu)t} \int_0^t e^{\mu v} c'(v) dv dt &= \int_0^\infty e^{\mu v} c'(v) \int_v^\infty e^{-(\lambda+\mu)t} dt dv \\ &= \frac{1}{\lambda + \mu} \int_0^\infty e^{-\lambda v} c'(v) dv \\ &= \frac{1}{\lambda + \mu} [-c(0^+) + \lambda \hat{c}(\lambda)], \end{aligned}$$

and we conclude the proof of the theorem.  $\square$

*Remark 3.3.* In the case that the function  $c'$  is a Laplace-transformable function, the proof of Theorem 3.2 is a straightforward consequence of Theorem 3.1(i), Theorem 3.1(ii), and the equality (3.1). The interesting example  $c = g_{1-\alpha}$  with  $0 < \alpha < 1$  does not satisfy this condition; however, it is absolutely continuous on  $(0, \infty)$  and Laplace-transformable, and we can apply Theorem 3.2(i) directly.

The following inversion theorem allows us to express operators  $()^+$  and  $()^-$  in terms of double convolution products. These equalities play important roles in extension formulae which are obtained in the following sections.

**Theorem 3.4.** *Let  $a \in L^1_{\text{loc}}(\mathbb{R}_+)$  be a Laplace-transformable function, and suppose that there exists  $c \in L^1_{\text{loc}}(\mathbb{R}_+)$  absolutely continuous on  $(0, \infty)$  and Laplace-transformable such that*

$$(a * c)(t) = 1, \quad t > 0. \tag{3.2}$$

- (i) *If  $(c')^+$  is 2-Laplace-transformable, then  $a^+ = -((c')^+ *_2 (a \otimes a))$ .*
- (ii) *If  $(c')^-$  is 2-Laplace-transformable and  $c(0^+) = 0$ , then  $a^- = ((c')^- *_2 (a \otimes a))$ .*

*Proof.* (i) We use Theorem 3.1(i) and Theorem 3.2(i) to prove that

$$\begin{aligned} \mathcal{L}_2(a^+)(z, w) &= \frac{\hat{a}(w) - \hat{a}(z)}{z - w} = \left( \frac{z\hat{c}(z) - w\hat{c}(w)}{z - w} \right) \frac{1}{zw\hat{c}(w)\hat{c}(z)} \\ &= \left( \frac{z\hat{c}(z) - w\hat{c}(w)}{z - w} \right) \hat{a}(w)\hat{a}(z) = \mathcal{L}_2((c')^+ * _2 (a \otimes a))(z, w). \end{aligned}$$

Due to the uniqueness of the Laplace transform (see, e.g., [5, p. 346]), we conclude the equality. The proof of part (ii) is similar and involves Theorem 3.1(ii) and Theorem 3.2(ii). □

*Example 3.5.* In the case that  $c'$  is a Laplace-transformable function, we apply Corollary 2.2(ii) to obtain an alternative proof of Theorem 3.4(i). However, the interesting example  $a = g_\alpha$  for  $0 < \alpha < 1$  and  $c = g_{1-\alpha}$  does not satisfy this condition and the direct proof given in Theorem 3.4(i) is needed. Note that  $c' = g_{-\alpha}$  and that the equality  $g_\alpha^+ = -((g_{-\alpha})^+ * _2 (g_\alpha \otimes g_\alpha))$  is equivalent, after an algebraic manipulation, to the formula

$$\frac{\alpha \sin \alpha \pi}{\pi} \int_0^t \int_0^s \frac{u^{\alpha-1} v^{\alpha-1}}{(t + u + s - v)^{\alpha+1}} ds dv = (t + s)^{\alpha-1}, \quad t, s > 0.$$

Analogously, let  $S, T : \mathbb{R}_+ \rightarrow \mathcal{B}(X)$  be strongly continuous operator families such that  $S(\cdot)x, T(\cdot)x \in L^1_{\text{loc}}(\mathbb{R}_+, X)$  for any  $x \in X$ . The operators  $S, T$  are said to be Laplace-transformable functions if there exists  $\omega \in \mathbb{R}$  such that the Laplace transform of  $S$  (resp.,  $T$ )

$$\hat{S}(\lambda)x = \int_0^\infty e^{-\lambda t} S(t)x dt, \quad \Re \lambda > \omega,$$

converges for  $x \in X$  (see, e.g., [1, Definition 3.1.4]). For  $h \in \mathbb{R}$ . we will denote  $S_h$  the translation operator of  $S$  given by  $S_h(u) := S(u + h)\chi_{[-h, +\infty)}(u)$  for  $u \in \mathbb{R}$  and the convolution product between  $T$  and  $S$ ,  $T * S$  given by

$$(T * S)(t)x := \int_0^t T(t - s)S(s)x ds, \quad t > 0, x \in X.$$

If  $g \in L^1_{\text{loc}}(\mathbb{R}_+)$  is a Laplace-transformable scalar-valued function, then we define  $g * S$  by

$$(g * S)(t)x := \int_0^t g(t - s)S(s)x ds, \quad t > 0, x \in X,$$

and

$$\begin{aligned} & (g^+ *_2 (S \otimes S))(t, s)x \\ & := \int_0^t \int_0^s g(t + s - r_1 - r_2)S(r_1)S(r_2)x \, dr_1 \, dr_2, \quad t, s \geq 0, x \in X, \end{aligned}$$

where  $(S \otimes S)(t, s) := S(t)S(s)$  is the composition operator for  $t, s \geq 0$ . We recall the following identities given in [7, Lemma 4.1]: for  $\lambda > \mu > \omega$ ,

$$\hat{S}(\lambda)\hat{T}(\mu)x = \int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} S(t)T(s)x \, ds \, dt, \quad x \in X, \quad (3.3)$$

and

$$\frac{1}{\mu - \lambda}(\hat{S}(\lambda) - \hat{S}(\mu))x = \int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} S(t + s)x \, ds \, dt, \quad x \in X. \quad (3.4)$$

If  $g : \mathbb{R}^+ \rightarrow \mathbb{C}$  is Laplace-transformable, we also have

$$\frac{1}{\mu - \lambda}\hat{g}(\mu)[\hat{S}(\lambda) - \hat{S}(\mu)]x = \int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} (g * S_t)(s)x \, ds \, dt, \quad x \in X, \quad (3.5)$$

and

$$\frac{1}{\mu - \lambda}\hat{T}(\mu)[\hat{g}(\lambda) - \hat{g}(\mu)]x = \int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} (T * g_t)(s)x \, ds \, dt, \quad x \in X. \quad (3.6)$$

Defining  $S(t) = S(-t)$  for  $t < 0$ , we have

$$\begin{aligned} & \frac{1}{\mu + \lambda}(\hat{S}(\lambda) + \hat{S}(\mu))x \\ & = \int_0^\infty e^{-\lambda t} \int_0^\infty e^{-\mu s} S(t - s)x \, ds \, dt, \quad \lambda + \mu > 0, x \in X. \end{aligned} \quad (3.7)$$

In fact equations (3.3), (3.4), (3.5), and (3.6) are valid for  $\Re\lambda, \Re\mu > \omega$  with  $\lambda \neq \mu$ , and (3.7) for  $\Re\lambda, \Re\mu > \omega$  with  $\Re(\lambda + \mu) > 0$ .

The following theorem shows how some double Laplace transforms of the double convolution product is also related with the single Laplace transform.

**Proposition 3.6.** *Let  $g \in L^1_{\text{loc}}(\mathbb{R}_+)$  and  $S : \mathbb{R}_+ \rightarrow \mathcal{B}(X)$  be locally integrable and strongly continuous functions, both Laplace-transformable functions. Then the following identities hold:*

- (i)  $\mathcal{L}_2(g^+ *_2 (S \otimes S))(\lambda, \mu) = \frac{1}{\mu - \lambda}[\hat{g}(\lambda) - \hat{g}(\mu)]\hat{S}(\lambda)\hat{S}(\mu)$  for  $\Re\lambda, \Re\mu > \omega$  with  $\lambda \neq \mu$ ,
- (ii)  $\mathcal{L}_2(g^- *_2 (S \otimes S))(\lambda, \mu) = \frac{1}{\lambda + \mu}[\hat{g}(\lambda) + \hat{g}(\mu)]\hat{S}(\lambda)\hat{S}(\mu)$  for  $\Re\lambda, \Re\mu > \omega$  with  $\Re(\lambda + \mu) > 0$ .

*Proof.* It is sufficient to apply Theorem 3.1(iv), (i) (or (ii)), and (3.3).  $\square$

The proof of the next corollary is a straightforward consequence of Theorem 3.1(iv), Theorem 3.2, and (3.3).

**Corollary 3.7.** *Let  $c \in L^1_{\text{loc}}(\mathbb{R}_+)$  be an absolutely continuous on  $(0, \infty)$  and Laplace-transformable function, and let  $S : \mathbb{R}^+ \rightarrow \mathcal{B}(X)$  be a locally integrable and strongly continuous and Laplace-transformable operator-valued function.*

(i) If  $(c')^+ : \mathbb{R}_+^2 \rightarrow \mathbb{C}$  is 2-Laplace-transformable, then

$$\begin{aligned} & \mathcal{L}_2((c')^+ *_2 (S \otimes S))(\lambda, \mu) \\ &= \frac{1}{\mu - \lambda} [\lambda \hat{c}(\lambda) - \mu \hat{c}(\mu)] \hat{S}(\lambda) \hat{S}(\mu), \quad \Re \lambda, \Re \mu > \omega, \lambda \neq \mu. \end{aligned}$$

(ii) If  $(c')^- : \mathbb{R}_+^2 \rightarrow \mathbb{C}$  is 2-Laplace transformable and  $c(0^+)$  exists, then

$$\mathcal{L}_2((c')^- *_2 (S \otimes S))(\lambda, \mu) = \frac{1}{\lambda + \mu} [\lambda \hat{c}(\lambda) + \mu \hat{c}(\mu)] \hat{S}(\lambda) \hat{S}(\mu) - \frac{2c(0^+)}{\lambda + \mu} \hat{S}(\lambda) \hat{S}(\mu),$$

for  $\Re \lambda, \Re \mu > \omega$  with  $\Re(\lambda + \mu) > 0$ .

*Example 3.8.* Let  $c = g_{1-\alpha}$  for  $0 < \alpha < 1$  and  $\hat{c}(\lambda) = \frac{1}{\lambda^{1-\alpha}}$  for  $\Re \lambda > 0$ . Then  $c' = g_{-\alpha}$  and we obtain the identity

$$\mathcal{L}_2((c')^+ *_2 (S \otimes S))(\lambda, \mu) = \frac{1}{\mu - \lambda} [\lambda^\alpha - \mu^\alpha] \hat{S}(\lambda) \hat{S}(\mu) \quad (3.8)$$

by Corollary 3.7(i). In particular, we recover [13, Formula (2.8)].

#### 4. LOCAL $(a, k)$ -REGULARIZED RESOLVENT FAMILIES

In this section, we prove extension theorems for local  $(a, k)$ -regularized resolvent families. In the following we suppose that the function  $k$  satisfies  $k(t) \neq 0$  for all  $t \in (0, \sigma)$ , where  $\sigma$  is a sufficiently small positive number. We begin by recalling the following definition.

*Definition 4.1.* Let  $0 < \tau \leq \infty$ , let  $a, k \in L_{\text{loc}}^1([0, \tau))$  with  $k \in \mathcal{C}(0, \tau)$  so that  $k(t) \neq 0$  for all  $t \in (0, \sigma)$  ( $\sigma$  small), and let  $A$  be a closed operator. A strongly continuous operator family  $\{S(t)\}_{t \in (0, \tau)} \subset \mathcal{B}(X)$  is a local (resp., global in case  $\tau = \infty$ )  $(a, k)$ -regularized resolvent family generated by  $A$  if the following conditions are satisfied:

- (i)  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{k(t)} = x$  for all  $x \in X$ ,
- (ii)  $S(t)A \subset AS(t), t \in (0, \tau)$ ,
- (iii)  $(a * S)(t)x \in D(A)$  for  $t \in (0, \tau)$  and  $x \in X$ , and the following Volterra equation holds:

$$A(a * S)(t)x = S(t)x - k(t)x, \quad x \in X, t \in (0, \tau). \quad (4.1)$$

*Remark 4.2.* The reason why we do not consider directly the value of  $S(\cdot)$  at the origin is that  $k$  could have a singularity at the origin; for example,  $k(t) = g_{\beta+1}(t)$  has a singularity at 0 if  $-1 < \beta < 0$ .

In the rest of this article we will assume that the functions  $a, k$  are positive.

We note that loss of regularity arises because we treat with evolution equations corresponding to regularization of certain base equation. The typical example is the local  $\alpha$ -times integrated semigroups, for example, the evolution equation

$$u'(t) = Au(t) + g_\alpha(t)x,$$

where  $\alpha > 0$ . In this case, the base equation, where no loss of regularity happens, is the Cauchy problem

$$u'(t) = Au(t),$$

where  $A$  is the generator of a  $C_0$ -semigroup (i.e., it is known that a local  $C_0$ -semigroup can be extended without loss of regularity). In terms of  $(a, k)$ -regularized resolvent families, it means that if  $A$  is the generator of a local  $(1, 1)$ -regularized resolvent family on  $[0, \tau)$ , then  $A$  is also the generator of a  $(1, 1)$ -regularized resolvent family on  $[0, 2\tau)$  and so on. However, this property is no longer true in the general case of  $(a, k)$ -regularized resolvent families, where loss of regularity is present. This phenomena has been observed for the case of  $k$ -convoluted semigroups (see [8]; in particular, for  $\alpha$ -times integrated semigroups, see [2], [22]) and of  $k$ -convoluted cosine families (see [23]).

The next theorem shows the most general result about extension of  $(a, k)$ -regularized resolvent families. It shows that we can extend a local  $(a, k)$ -regularized resolvent family defined in  $(0, \tau)$  to get a  $(a, (k * a)^{*n} * a)$ -regularized resolvent family in  $(0, (n + 1)\tau)$ .

**Theorem 4.3.** *Let  $n \in \mathbb{N}$ ,  $0 < \tau \leq \infty$ ,  $a, k \in L^1_{\text{loc}}([0, (n + 1)\tau))$  with  $k \in \mathcal{C}(0, (n + 1)\tau)$ , and let  $\{S_1(t)\}_{t \in (0, \tau)}$  be a local  $(a, k)$ -regularized resolvent family generated by  $A$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined recursively by*

$$\begin{aligned} S_{n+1}(t)x &:= (k * a * S_n)(t)x, \quad x \in X, t \in (0, nT], \quad \text{and} \\ S_{n+1}(t)x &:= (a^+ *_2 (S_n \otimes S_1))(nT, t - nT)x + \int_0^{nT} k(t - r)(a * S_n)(r)x dr \\ &\quad + \int_0^{t-nT} ((k * a)^{*n-1} * k)(t - r)(a * S_1)(r)x dr, \end{aligned}$$

for  $x \in X$  and  $t \in (nT, (n + 1)T]$ , is a local  $(a, (k * a)^{*n} * k)$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  will generate a local  $(a, (k * a)^{*n} * k)$ -regularized resolvent family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)\tau)}$ .

*Proof.* Note that the family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  is strongly continuous:

$$\begin{aligned} \lim_{t \rightarrow (nT)^+} S_{n+1}(t)x &= \lim_{t \rightarrow (nT)^+} \left( (a^+ *_2 (S_n \otimes S_1))(nT, t - nT)x \right. \\ &\quad \left. + \int_0^{nT} k(t - r)(a * S_n)(r)x dr \right. \\ &\quad \left. + \int_0^{t-nT} ((k * a)^{*n-1} * k)(t - r)(a * S_1)(r)x dr \right). \end{aligned}$$

The first summand tends to 0 using Corollary 2.2(i) in the vectorial case, the second one tends to  $(k * a * S_n)(nT)x$ , and the last term goes to 0. Furthermore, for  $\varepsilon > 0$  there exists  $t > 0$  sufficiently small such that

$$\left\| \frac{S_n(s)x}{(k * a)^{*n-1} * k(s)} - x \right\| < \varepsilon, \quad \text{for } 0 < s < t.$$

Then

$$\begin{aligned}
& \left\| \frac{S_{n+1}(t)x}{(k * a)^{*n} * k(t)} - x \right\| \\
&= \left\| \frac{S_{n+1}(t)x - (k * a)^{*n} * k(t)x}{(k * a)^{*n} * k(t)} \right\| \\
&\leq \frac{1}{(k * a)^{*n} * k(t)} \int_0^t (k * a)(t-s) \|S_n(s)x - (k * a)^{*(n-1)} * k(s)x\| ds \\
&\leq \frac{1}{(k * a)^{*n} * k(t)} \\
&\quad \times \int_0^t (k * a)(t-s)(k * a)^{*(n-1)} * k(s) \left\| \frac{S_n(s)x}{(k * a)^{*(n-1)} * k(s)} - x \right\| ds \leq \varepsilon.
\end{aligned}$$

So,  $\lim_{t \rightarrow 0^+} \frac{S_{n+1}(t)x}{(k * a)^{*n} * k(t)} = x$  for all  $x \in X$ . Note that  $\{S_{n+1}(t)\}_{t \in (0, nT]}$  is a local  $(a, (k * a)^{*n} * k)$ -regularized resolvent family generated by  $A$  (see [16, Remark 2.4(4)]). Now let  $t \in (nT, (n+1)T]$  and  $x \in X$ . It is clear that  $S_{n+1}(t)A \subset AS_{n+1}(t)$ . We show that  $(a * S_{n+1})(t)x \in D(A)$ . Note

$$(a * S_{n+1})(t)x = \int_0^{nT} a(t-s)S_{n+1}(s)x ds + \int_{nT}^t a(t-s)S_{n+1}(s)x ds.$$

On the one hand,

$$\int_0^{nT} a(t-s)S_{n+1}(s)x ds = \int_0^{nT} a(t-s)(k * a * S_n)(s)x ds \in D(A),$$

since  $(a * S_n)(s)x \in D(A)$ . On the other hand,

$$\begin{aligned}
& \int_{nT}^t a(t-s)S_{n+1}(s)x ds \\
&= \int_{nT}^t a(t-s) \left( \int_0^{s-nT} \int_0^{nT} a(s-r_1-r_2)S_n(r_1)S_1(r_2)x dr_1 dr_2 \right) ds \\
&\quad + \int_{nT}^t a(t-s) \int_0^{nT} k(s-r)(a * S_n)(r)x dr ds \\
&\quad + \int_{nT}^t a(t-s) \int_0^{s-nT} ((k * a)^{*(n-1)} * k)(s-r)(a * S_1)(r)x dr ds.
\end{aligned} \tag{4.2}$$

Note that  $(a * S_n)(r)x, (a * S_1)(r)x \in D(A)$ . Finally,

$$\begin{aligned}
& \int_{nT}^t a(t-s) \int_0^{s-nT} \int_0^{nT} a(s-r_1-r_2)S_n(r_1)S_1(r_2)x dr_1 dr_2 ds \\
&= \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_{nT}^s a(u-r_1)S_1(s-u)x du ds dr_1 \\
&= \int_0^{nT} S_n(r_1) \int_{nT}^t a(u-r_1) \int_u^t a(t-s)S_1(s-u)x ds du dr_1
\end{aligned}$$

$$= \int_0^{nT} S_n(r_1) \int_{nT}^t a(u-r_1) \int_0^{t-u} a(t-u-v) S_1(v) x \, dv \, du \, dr_1 \in D(A)$$

since  $(a * S_1)(t-u) \in D(A)$ . To finish the proof, we prove that for  $t \in (nT, (n+1)T]$  and  $x \in X$ , the equality (4.1) is satisfied. First observe that

$$A(a * S_{n+1})(t)x = A \int_0^{nT} a(t-s)(k * a * S_n)(s)x \, ds + A \int_{nT}^t a(t-s) S_{n+1}(s)x \, ds.$$

We apply the operator  $A$  to the first summand of (4.2), and we obtain

$$\begin{aligned} & A \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_0^{s-nT} a(s-r_1-r_2) S_1(r_2)x \, dr_2 \, ds \, dr_1 \\ &= A \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_{nT}^s a(u-r_1) S_1(s-u)x \, du \, ds \, dr_1 \\ &= A \int_0^{nT} S_n(r_1) \int_{nT}^t a(u-r_1) \int_u^t a(t-s) S_1(s-u)x \, ds \, du \, dr_1 \\ &= A \int_0^{nT} S_n(r_1) \int_{nT}^t a(u-r_1) \int_0^{t-u} a(t-u-v) S_1(v)x \, dv \, du \, dr_1 \\ &= \int_0^{nT} S_n(r_1) \int_{nT}^t a(u-r_1) (S_1(t-u) - k(t-u))x \, du \, dr_1 \\ &= \int_0^{nT} S_n(r_1) \int_0^{t-nT} a(t-r_1-r_2) (S_1(r_2) - k(r_2))x \, dr_2 \, dr_1. \end{aligned}$$

In the second summand of (4.2), we write

$$\begin{aligned} & \int_0^{nT} (a * S_n)(r)x \int_{nT}^t a(t-s) k(s-r) \, ds \, dr \\ &= \int_0^{nT} (a * S_n)(r)x \left( \int_0^{t-r} a(t-r-u) k(u) \, du \right. \\ & \quad \left. - \int_0^{nT-r} a(t-r-u) k(u) \, du \right) dr. \end{aligned}$$

We apply the operator  $A$  to each of the above terms to get

$$\begin{aligned} & A \int_0^{nT} (a * S_n)(r)x \int_0^{t-r} a(t-r-u) k(u) \, du \, dr \\ &= A \int_0^{t-nT} k(u) \int_0^{nT} a(t-u-r) (a * S_n)(r)x \, dr \, du \\ & \quad + A \int_{t-nT}^t k(u) \int_0^{t-u} a(t-u-r) (a * S_n)(r)x \, dr \, du \\ &= \int_0^{t-nT} k(u) \int_0^{nT} a(t-u-r) (S_n(r)x - ((k * a)^{(n-1)} * k)(r)x) \, dr \, du \\ & \quad + \int_{t-nT}^t k(u) ((a * S_n)(t-u)x - (k * a)^{*n}(t-u)x) \, du \end{aligned}$$

$$\begin{aligned}
&= \int_0^{t-nT} k(u) \int_0^{nT} a(t-u-r)(S_n(r)x - ((k * a)^{*(n-1)} * k)(r)x) dr du \\
&\quad + \int_0^{nT} k(t-r)((a * S_n)(r)x - (k * a)^{*n}(r)x) dr,
\end{aligned}$$

and

$$\begin{aligned}
&A \int_0^{nT} (a * S_n)(r)x \int_0^{nT-r} a(t-r-u)k(u) du dr \\
&= A \int_0^{nT} k(u) \int_0^{nT-u} a(t-u-r)(a * S_n)(r)x dr du.
\end{aligned}$$

In the third summand of (4.2), we write

$$\begin{aligned}
&\int_0^{t-nT} (a * S_1)(r)x \int_{r+nT}^t a(t-s)((k * a)^{*(n-1)} * k)(s-r) ds dr \\
&= \int_0^{t-nT} (a * S_1)(r)x \left( \int_r^t a(t-s)((k * a)^{*(n-1)} * k)(s-r) ds \right. \\
&\quad \left. - \int_r^{r+nT} a(t-s)((k * a)^{*(n-1)} * k)(s-r) ds \right) dr.
\end{aligned}$$

We apply the operator  $A$  to each of the above terms to obtain

$$\begin{aligned}
&A \int_0^{t-nT} (a * S_1)(r)x \int_r^t a(t-s)((k * a)^{*(n-1)} * k)(s-r) ds dr \\
&= A \int_0^{t-nT} (a * S_1)(r)x \int_0^{t-r} a(t-r-u)((k * a)^{*(n-1)} * k)(u) du dr \\
&= A \int_0^{nT} ((k * a)^{*(n-1)} * k)(u) \int_0^{t-nT} a(t-u-r)(a * S_1)(r)x dr du \\
&\quad + A \int_{nT}^t ((k * a)^{*(n-1)} * k)(u) \int_0^{t-u} a(t-u-r)(a * S_1)(r)x dr du \\
&= A \int_0^{nT} ((k * a)^{*(n-1)} * k)(u) \int_0^{t-nT} a(t-u-r)(a * S_1)(r)x dr du \\
&\quad + \int_{nT}^t ((k * a)^{*(n-1)} * k)(u) ((a * S_1)(t-u)x - (a * k)(t-u)x) du \\
&= A \int_0^{nT} ((k * a)^{*(n-1)} * k)(u) \int_0^{t-nT} a(t-u-r)(a * S_1)(r)x dr du \\
&\quad + \int_0^{t-nT} ((k * a)^{*(n-1)} * k)(t-s)((a * S_1)(s)x - (a * k)(s)x) ds
\end{aligned}$$

and

$$A \int_0^{t-nT} (a * S_1)(r)x \int_r^{r+nT} a(t-s)((k * a)^{*(n-1)} * k)(s-r) ds dr$$



$$= A \int_0^{nT} ((k * a)^{*(n-1)} * k)(u) \int_0^{t-nT} a(t-u-r)(a * S_1)(r)x \, dr \, du.$$

Then we have

$$\begin{aligned} & A \int_0^{t-nT} (a * S_1)(r)x \int_{r+nT}^t a(t-s)((k * a)^{*(n-1)} * k)(s-r) \, ds \, dr \\ &= \int_0^{t-nT} ((k * a)^{*(n-1)} * k)(t-s)((a * S_1)(s)x - (a * k)(s)x) \, ds. \end{aligned}$$

Finally note that

$$\begin{aligned} & A \int_0^{nT} a(t-s)(k * a * S_n)(s)x \, ds \\ &= A \int_0^{nT} (a * S_n)(r)x \int_r^{nT} a(t-s)k(s-r) \, ds \, dr \\ &= A \int_0^{nT} (a * S_n)(r)x \int_0^{nT-r} a(t-r-u)k(u) \, du \, dr \\ &= A \int_0^{nT} k(u) \int_0^{nT-u} a(t-u-r)(a * S_n)(r)x \, dr \, du. \end{aligned}$$

We join together all summands to conclude that

$$\begin{aligned} A(a * S_{n+1})(t)x &= \int_0^{t-nT} \int_0^{nT} a(t-r_1-r_2)S_n(r_1)S_1(r_2)x \, dr_1 \, dr_2 \\ &+ \int_0^{nT} k(t-r)(a * S_n)(r)x \, dr \\ &+ \int_0^{t-nT} ((k * a)^{*(n-1)} * k)(t-r)(a * S_1)(r)x \, dr \\ &- \int_0^{t-nT} k(u) \int_0^{nT} a(t-u-r)((k * a)^{*(n-1)} * k)(r)x \, dr \, du \\ &- \int_0^{nT} k(t-r)(k * a)^{*n}(r)x \, dr \\ &- \int_0^{t-nT} ((k * a)^{*(n-1)} * k)(t-r)(a * k)(r)x \, dr \\ &= S_{n+1}(t)x - (k * a)^{*n}(t)x, \end{aligned}$$

where we have used Lemma 2.3.  $\square$

The expression of  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  is not unique; in the following theorem we show  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  in terms of  $\{S_j(t)\}_{t \in (0, jT]}$  for all  $1 \leq j \leq n$ . The proof is similar to the proof of Theorem 4.3 and therefore we omit it.

**Theorem 4.4.** *Let  $n \in \mathbb{N}$ , let  $0 < \tau \leq \infty$ , let  $a, k \in L^1_{\text{loc}}([0, (n+1)\tau))$  with  $k \in \mathcal{C}(0, (n+1)\tau)$ , and let  $\{S_1(t)\}_{t \in (0, \tau)}$  be a local  $(a, k)$ -regularized resolvent*

family generated by  $A$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined in Theorem 4.3 satisfies

$$\begin{aligned} S_{n+1}(t)x &:= ((k * a)^{* (n+1-j)} * S_j)(t)x, \quad x \in X, t \in (0, jT], \quad \text{and} \\ S_{n+1}(t)x &:= (a^+ *_2 (S_j \otimes S_{n+1-j}))(jT, t - jT)x \\ &\quad + \int_0^{jT} ((k * a)^{n-j} * k)(t - r)(a * S_j)(r)x \, dr \\ &\quad + \int_0^{t-jT} ((k * a)^{* (j-1)} * k)(t - r)(a * S_{n+1-j})(r)x \, dr, \end{aligned}$$

for  $x \in X$ ,  $1 \leq j \leq n$  and  $t \in (jT, (n+1)T]$  for any  $T < \tau$ .

The following result is related to [8, Theorem 4.4] and [23, Theorem 3.3]. However, note that both results are not included in this corollary.

**Corollary 4.5.** *Let  $n \in \mathbb{N}$ , let  $0 < \tau \leq \infty$ , and let  $\{S_1(t)\}_{t \in (0, \tau]}$  be a local  $(g_\alpha, g_{\beta+1})$ -regularized resolvent family generated by  $A$ , with  $\alpha > 0$  and  $\beta > -1$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined by*

$$S_{n+1}(t)x := (g_{\beta+\alpha+1} * S_n)(t)x, \quad x \in X,$$

for  $t \in (0, nT]$  and

$$\begin{aligned} S_{n+1}(t)x &:= (g_\alpha^+ *_2 (S_n \otimes S_1))(nT, t - nT)x + \int_0^{nT} g_{\beta+1}(t - r)(g_\alpha * S_n)(r)x \, dr \\ &\quad + \int_0^{t-nT} g_{n(\beta+1)+\alpha(n-1)}(t - r)(g_\alpha * S_1)(r)x \, dr, \end{aligned}$$

for  $x \in X$  and  $t \in (nT, (n+1)T]$  is a local  $(g_\alpha, g_{(n+1)(\beta+1)+n\alpha})$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  generates a local  $(g_\alpha, g_{(n+1)(\beta+1)+n\alpha})$ -regularized resolvent family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)\tau]}$ .

However, if we restrict for example to the  $\alpha$ -times integrated semigroup case, the above extension is not the sharpest extension. Then for certain cases of the functions  $a$  and  $k$  there exist sharper extensions from the point of view of the regularized Cauchy problems. The following theorem gives us this sharp extension for a class of  $(a, k)$ -regularized resolvent families. Although the idea of the proof is similar to the proof of Theorem 4.3, we have included it to make easier the reading because we use additional methods.

**Theorem 4.6.** *Let  $n \in \mathbb{N}$ ,  $0 < \tau \leq \infty$ ,  $a, k \in L^1_{\text{loc}}(\mathbb{R}_+)$  with  $k \in \mathcal{C}(0, \infty)$  be Laplace-transformable functions such that there exist  $b, c \in L^1_{\text{loc}}(\mathbb{R}_+)$  Laplace-transformable satisfying the condition that  $c$  be absolutely continuous on  $(0, \infty)$ ,  $(c')^+$  be 2-Laplace-transformable,*

$$(a * b)(t) = k(t), \quad (a * c)(t) = 1, \quad t > 0,$$

and  $\{S_1(t)\}_{t \in (0, \tau]}$  be a local  $(a, k)$ -regularized resolvent family generated by  $A$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined by

$$S_{n+1}(t)x := (b * S_n)(t)x, \quad x \in X,$$

for  $t \in (0, nT]$  and

$$\begin{aligned} S_{n+1}(t)x &:= \int_0^{nT} b(t-r)S_n(r)x \, dr + \int_0^{t-nT} b^{*n}(t-r)S_1(r)x \, dr \\ &\quad - ((c')^+ *_2 (S_n \otimes S_1))(nT, t-nT)x \end{aligned}$$

for  $x \in X$  and  $t \in (nT, (n+1)T]$  is a local  $(a, b^{*n} * k)$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  generates a local  $(a, b^{*n} * k)$ -regularized resolvent family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)\tau)}$ .

*Proof.* Similarly to the proof of Theorem 4.3,  $\lim_{t \rightarrow 0^+} \frac{S_{n+1}(t)x}{(b^{*n} * k)(t)} = x$  for  $x \in X$ . Note that  $\{S_{n+1}(t)\}_{t \in (0, nT]}$  is a local  $(a, b^{*n} * k)$ -regularized resolvent family generated by  $A$  (see again [16, Remark 2.4(4)]). Now let  $t \in (nT, (n+1)T]$  and let  $x \in X$ . It is clear that  $S_{n+1}(t)A \subset AS_{n+1}(t)$ , and, following the proof of Theorem 4.3, it is easy to see that  $(a * S_{n+1})(t)x \in D(A)$ .

Now we prove that for  $t \in (nT, (n+1)T]$  and  $x \in X$ , the equality (4.1) is satisfied. First observe that

$$A(a * S_{n+1})(t)x = A \int_0^{nT} a(t-s)(b * S_n)(s)x \, ds + A \int_{nT}^t a(t-s)S_{n+1}(s)x \, ds.$$

Note that

$$\begin{aligned} &\int_{nT}^t a(t-s)S_{n+1}(s)x \, ds \\ &= \int_{nT}^t a(t-s) \left( \int_0^{nT} b(s-r)S_n(r)x \, dr \right. \\ &\quad \left. + \int_0^{s-nT} b^{*n}(s-r)S_1(r)x \, dr \right) ds \\ &\quad - \int_{nT}^t a(t-s) \int_0^{s-nT} \int_0^{nT} c'(s-r_1-r_2)S_n(r_1)S_1(r_2)x \, dr_1 \, dr_2 \, ds. \end{aligned} \tag{4.3}$$

We apply the operator  $A$  to the third summand of (4.3), and we obtain

$$\begin{aligned} &-A \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_0^{s-nT} c'(s-r_1-r_2)S_1(r_2)x \, dr_2 \, ds \, dr_1 \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_{nT}^s c'(u-r_1)S_1(s-u)x \, du \, ds \, dr_1 \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1) \int_u^t a(t-s)S_1(s-u)x \, ds \, du \, dr_1 \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1) \int_0^{t-u} a(t-u-v)S_1(v)x \, dv \, du \, dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1)(S_1(t-u) - k(t-u))x \, du \, dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_0^{t-nT} c'(t-r_1-r_2)(S_1(r_2) - k(r_2))x \, dr_2 \, dr_1. \end{aligned}$$

In the first summand of (4.3), we write

$$\begin{aligned} & \int_0^{nT} S_n(r)x \int_{nT}^t a(t-s)b(s-r) ds dr \\ &= \int_0^{nT} S_n(r)x \int_0^{t-r} a(t-r-u)b(u) du dr \\ & \quad - \int_0^{nT} S_n(r)x \int_0^{nT-r} a(t-r-u)b(u) du dr. \end{aligned}$$

We apply the operator  $A$  to each of the above terms to get

$$\begin{aligned} & A \int_0^{nT} S_n(r)x \int_0^{t-r} a(t-r-u)b(u) du dr \\ &= A \int_0^{t-nT} b(u) \int_0^{nT} a(t-u-r)S_n(r)x dr du \\ & \quad + A \int_{t-nT}^t b(u) \int_0^{t-u} a(t-u-r)S_n(r)x dr du \\ &= A \int_0^{t-nT} b(u) \int_0^{nT} a(t-u-r)S_n(r)x dr du \\ & \quad + \int_{t-nT}^t b(u) (S_n(t-u)x - (b^{*(n-1)} * k)(t-u)x) du \\ &= A \int_0^{t-nT} b(u) \int_0^{nT} a(t-u-r)S_n(r)x dr du \\ & \quad + \int_0^{nT} b(t-r) (S_n(r)x - (b^{*(n-1)} * k)(r)x) dr, \end{aligned}$$

and

$$\begin{aligned} & \int_0^{nT} S_n(r)x \int_0^{nT-r} a(t-r-u)b(u) du dr \\ &= \int_0^{nT} b(u) \int_0^{nT-u} a(t-u-r)S_n(r)x dr du. \end{aligned}$$

In the second summand of (4.3), we write

$$\begin{aligned} & \int_0^{t-nT} S_1(r)x \int_{r+nT}^t a(t-s)b^{*n}(s-r) ds dr \\ &= \int_0^{t-nT} S_1(r)x \left( \int_r^t a(t-s)b^{*n}(s-r) ds - \int_r^{r+nT} a(t-s)b^{*n}(s-r) ds \right) dr. \end{aligned}$$

We apply the operator  $A$  to each of the above terms to obtain

$$\begin{aligned} & A \int_0^{t-nT} S_1(r)x \int_r^t a(t-s)b^{*n}(s-r) ds dr \\ &= A \int_0^{t-nT} S_1(r)x \int_0^{t-r} a(t-r-u)b^{*n}(u) du dr \end{aligned}$$

$$\begin{aligned}
&= A \int_0^{nT} b^{*n}(u) \int_0^{t-nT} a(t-u-r)S_1(r)x \, dr \, du \\
&\quad + A \int_{nT}^t b^{*n}(u) \int_0^{t-u} a(t-u-r)S_1(r)x \, dr \, du \\
&= A \int_0^{nT} b^{*n}(u) \int_0^{t-nT} a(t-u-r)S_1(r)x \, dr \, du \\
&\quad + \int_{nT}^t b^{*n}(u)(S_1(t-u)x - k(t-u)x) \, du \\
&= A \int_0^{nT} b^{*n}(u) \int_0^{t-nT} a(t-u-r)S_1(r)x \, dr \, du \\
&\quad + \int_0^{t-nT} b^{*n}(t-s)(S_1(s)x - k(s)x) \, ds
\end{aligned}$$

and

$$\begin{aligned}
&\int_0^{t-nT} S_1(r)x \int_r^{r+nT} a(t-s)b^{*n}(s-r) \, ds \, dr \\
&= \int_0^{nT} b^{*n}(u) \int_0^{t-nT} a(t-u-r)S_1(r)x \, dr \, du.
\end{aligned}$$

Then we have that

$$\begin{aligned}
&A \int_0^{t-nT} S_1(r)x \int_{r+nT}^t a(t-s)b^{*n}(s-r) \, ds \, dr \\
&= \int_0^{t-nT} b^{*n}(t-s)(S_1(s)x - k(s)x) \, ds.
\end{aligned}$$

Furthermore note that

$$\begin{aligned}
A \int_0^{nT} a(t-s)(b * S_n)(s)x \, ds &= A \int_0^{nT} S_n(r)x \int_r^{nT} a(t-s)b(s-r) \, ds \, dr \\
&= A \int_0^{nT} S_n(r)x \int_0^{nT-r} a(t-r-u)b(u) \, du \, dr \\
&= A \int_0^{nT} b(u) \int_0^{nT-u} a(t-u-r)S_n(r)x \, dr \, du.
\end{aligned}$$

We join together all summands to conclude that

$$\begin{aligned}
A(a * S_{n+1})(t)x &= S_{n+1}(t)x + A \int_0^{t-nT} b(u) \int_0^{nT} a(t-u-r)S_n(r)x \, dr \, du \\
&\quad - \int_0^{nT} b(t-r)(b^{*(n-1)} * k)(r)x \, dr - \int_0^{t-nT} b^{*n}(t-r)k(r)x \, dr \\
&\quad + \int_0^{nT} S_n(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)k(r_2) \, dr_2 \, dr_1.
\end{aligned}$$

Now we use induction. As  $\{S_n(t)\}_{t \in (0, nT]}$  is a local  $(a, b^{*(n-1)} * k)$ -regularized resolvent family generated by  $A$ , then

$$S_n(r_1)x = A(a * S_n)(r_1)x + (b^{*(n-1)} * k)(r_1)x = A(a * S_n)(r_1)x + (b^{*n} * a)(r_1)x,$$

and so

$$\begin{aligned} & \int_0^{nT} S_n(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)k(r_2) dr_2 dr_1 \\ &= A \int_0^{nT} (a * S_n)(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)(a * b)(r_2) dr_2 dr_1 \\ & \quad + \int_0^{nT} (b^{*n} * a)(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)(a * b)(r_2) dr_2 dr_1. \end{aligned}$$

On the one hand,

$$\begin{aligned} & A \int_0^{nT} (a * S_n)(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)(a * b)(r_2) dr_2 dr_1 \\ &= A \int_0^{nT} S_n(u)x \int_0^{t-nT} b(v) \\ & \quad \times \int_u^{nT} \int_v^{t-nT} a(r_1-u)a(r_2-v)c'(t-r_1-r_2) dr_2 dr_1 dv du \\ &= -A \int_0^{nT} S_n(u)x \int_0^{t-nT} a(t-u-v)b(v) dv du \end{aligned}$$

and, on the other hand,

$$\begin{aligned} & \int_0^{nT} (b^{*n} * a)(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)(a * b)(r_2) dr_2 dr_1 \\ &= \int_0^{nT} b^{*n}(u)x \int_0^{t-nT} b(v) \\ & \quad \times \int_u^{nT} \int_v^{t-nT} a(r_1-u)a(r_2-v)c'(t-r_1-r_2) dr_2 dr_1 dv du \\ &= - \int_0^{nT} b^{*n}(u)x \int_0^{t-nT} a(t-u-v)b(v) dv du, \end{aligned}$$

where we have applied Theorem 3.4(i). Applying Lemma 2.3, we get

$$\begin{aligned} A(a * S_{n+1})(t)x &= S_{n+1}(t)x - \int_0^{nT} b(t-r)(b^{*(n-1)} * k)(r)x dr \\ & \quad - \int_0^{t-nT} b^{*n}(t-r)k(r)x dr \\ & \quad - \int_0^{nT} b^{*n}(u)x \int_0^{t-nT} a(t-u-v)b(v) dv du \\ &= S_{n+1}(t)x - \int_0^{nT} b(t-r)(b^{*n} * a)(r)x dr \end{aligned}$$

$$\begin{aligned}
& - \int_0^{t-nT} b^{*n}(t-r)(a*b)(r)x \, dr \\
& - \int_0^{nT} b^{*n}(u)x \int_0^{t-nT} a(t-u-v)b(v) \, dv \, du \\
& = S_{n+1}(t)x - (b^{*(n+1)} * a)(t)x = S_{n+1}(t)x - (b^{*n} * k)(t)x.
\end{aligned}$$

Finally we check that the family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  is strongly continuous. It is direct to check that  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  is uniformly bounded on  $[nT - \varepsilon, nT + \varepsilon]$  for all  $0 < \varepsilon < T$ , and strongly continuous on  $(0, nT) \cup (nT, (n+1)T]$ . Note that for  $t \rightarrow (nT)^+$ , we have that

$$\begin{aligned}
S_{n+1}(t)x - (b^{*n} * k)(t)x & = A \left( \int_0^{nT} a(t-s)S_{n+1}(s)x \, ds \right) \\
& \quad + A \left( \int_{nT}^t a(t-s)S_{n+1}(s)x \, ds \right) \\
& \rightarrow A \left( \int_0^{nT} a(nT-s)S_{n+1}(s)x \, ds \right) \\
& = S_{n+1}(nT)x - (b^{*n} * k)(nT)x, \quad x \in X,
\end{aligned}$$

and we conclude that the family  $\{S_{n+1}(t)\}_{t \in [0, (n+1)T]}$  is strongly continuous.  $\square$

The following result extends [22, Theorem 2] because we obtain the sharp extension of  $(g_\alpha, g_{\beta+1})$ -regularized resolvent families when  $0 < \alpha < 1$  and  $\beta - \alpha > -1$ , and when  $\alpha \rightarrow 1^-$  we recover the  $\alpha$ -times integrated semigroup case, considered in [22]. More generally one could consider the case of  $K$ -convoluted resolvent families, that is  $(g_\alpha, (1 * K))$ -regularized resolvent families for  $0 < \alpha < 1$ , and compare it to the limit case when  $\alpha \rightarrow 1^-$  (see [8, Theorem 4.4]).

**Corollary 4.7.** *Let  $n \in \mathbb{N}$ , let  $0 < \tau \leq \infty$ , and let  $\{S_1(t)\}_{t \in (0, \tau)}$  be a local  $(g_\alpha, g_{\beta+1})$ -regularized resolvent family generated by  $A$  with  $0 < \alpha < 1$  and  $\beta - \alpha > -1$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined by*

$$S_{n+1}(t)x := (g_{\beta-\alpha+1} * S_n)(t)x, \quad x \in X,$$

for  $t \in (0, nT]$  and

$$\begin{aligned}
S_{n+1}(t)x & := \int_0^{nT} g_{\beta-\alpha+1}(t-r)S_n(r)x \, dr + \int_0^{t-nT} g_{n(\beta-\alpha+1)}(t-r)S_1(r)x \, dr \\
& \quad - ((g_{-\alpha})^+ * (S_n \otimes S_1))(nT, t - nT)x,
\end{aligned}$$

for  $x \in X$  and  $t \in (nT, (n+1)T]$  is a local  $(g_\alpha, g_{n(\beta-\alpha+1)+\beta+1})$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  generates a local  $(g_\alpha, g_{n(\beta-\alpha+1)+\beta+1})$ -regularized resolvent family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)\tau)}$ .

## 5. SOLUTIONS OF EVOLUTIONARY PROBLEMS WITHOUT JUMPS OF REGULARITY

In this section, we identify a wide class of evolution equations where no loss of regularity happens. It is interesting to note that it was not known until now if this

property goes beyond the cases of the heat and wave equations (i.e., the semigroup and cosine cases). We begin with the following result, which is subordinated to the semigroup case in the sense that we cannot go beyond  $\alpha > 1$  when we restrict to the particular case of  $(g_\alpha, g_\alpha)$ -regularized resolvent families (see the next corollary).

**Theorem 5.1.** *Let  $n \in \mathbb{N}$ ,  $0 < \tau \leq \infty$ ,  $a \in L^1_{\text{loc}}(\mathbb{R}_+)$  with  $a \in \mathcal{C}(0, \infty)$  be a Laplace-transformable function such that there exists  $c \in L^1_{\text{loc}}(\mathbb{R}_+)$  Laplace-transformable satisfying the condition that  $c$  is absolutely continuous on  $(0, \infty)$ ,  $(c')^+$  is 2-Laplace-transformable, and  $(a * c)(t) = 1$  for all  $t > 0$ , and let  $\{S_1(t)\}_{t \in (0, \tau)}$  be a local  $(a, a)$ -regularized resolvent family generated by  $A$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined by*

$$S_{n+1}(t)x := S_n(t)x, \quad x \in X,$$

for  $t \in (0, nT]$  and

$$S_{n+1}(t)x := -((c')^+ *_2 (S_n \otimes S_1))(nT, t - nT)x, \quad x \in X,$$

and  $t \in (nT, (n+1)T]$  is a local  $(a, a)$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  generates a global  $(a, a)$ -regularized resolvent family  $\{S(t)\}_{t \in (0, \infty)}$ .

*Proof.* Let us note here that  $\lim_{t \rightarrow 0^+} \frac{S_{n+1}(t)x}{a(t)} = x$  for  $x \in X$  and the family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  is strongly continuous. The proof of this fact is similar to Theorem 4.6. Obviously,  $\{S_{n+1}(t)\}_{t \in (0, nT]}$  is a local  $(a, a)$ -regularized resolvent family generated by  $A$ . Now let  $t \in (nT, (n+1)T]$  and  $x \in X$ . It is clear that  $S_{n+1}(t)A \subset AS_{n+1}(t)$ . We show that  $(a * S_{n+1})(t)x \in D(A)$ . Note that

$$(a * S_{n+1})(t)x = \int_0^{nT} a(t-s)S_n(s)x \, ds + \int_{nT}^t a(t-s)S_{n+1}(s)x \, ds, \quad x \in X.$$

On the one hand, note that  $\int_0^{nT} a(t-s)S_n(s)x \, ds \in D(A)$  (see (5.1) at the end of the proof). On the other hand,

$$\begin{aligned} & \int_{nT}^t a(t-s)S_{n+1}(s)x \, ds \\ &= - \int_{nT}^t a(t-s) \int_0^{s-nT} \int_0^{nT} c'(s-r_1-r_2)S_n(r_1)S_1(r_2)x \, dr_1 \, dr_2 \, ds \\ &= - \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_0^{s-nT} c'(s-r_1-r_2)S_1(r_2)x \, dr_2 \, ds \, dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_{nT}^s c'(u-r_1)S_1(s-u)x \, du \, ds \, dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1) \int_u^t a(t-s)S_1(s-u)x \, ds \, du \, dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1) \int_0^{t-u} a(t-u-v)S_1(v)x \, dv \, du \, dr_1 \in D(A) \end{aligned}$$



since  $(a * S_1)(t-u) \in D(A)$ . To finish the proof, we prove that for  $t \in (nT, (n+1)T]$  and  $x \in X$  the equality

$$A(a * S_{n+1}(t))x = S_{n+1}(t)x - a(t)x,$$

is verified. First observe that

$$A(a * S_{n+1})(t)x = A \int_0^{nT} a(t-s)S_n(s)x ds + A \int_{nT}^t a(t-s)S_{n+1}(s)x ds.$$

Now, we develop the second term applying change of variables and Fubini's theorem:

$$\begin{aligned} & A \int_{nT}^t a(t-s)S_{n+1}(s)x ds \\ &= -A \int_{nT}^t a(t-s) \int_0^{s-nT} \int_0^{nT} c'(s-r_1-r_2)S_n(r_1)S_1(r_2)x dr_1 dr_2 ds \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_0^{s-nT} c'(s-r_1-r_2)S_1(r_2)x dr_2 ds dr_1 \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t a(t-s) \int_{nT}^s c'(u-r_1)S_1(s-u)x du ds dr_1 \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1) \int_u^t a(t-s)S_1(s-u)x ds du dr_1 \\ &= -A \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1) \int_0^{t-u} a(t-u-v)S_1(v)x dv du dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_{nT}^t c'(u-r_1)(S_1(t-u) - a(t-u))x du dr_1 \\ &= - \int_0^{nT} S_n(r_1) \int_0^{t-nT} c'(t-r_1-r_2)(S_1(r_2) - a(r_2))x dr_2 dr_1, \end{aligned}$$

where we have used the fact that  $\{S_1(t)\}_{t \in (0, T]}$  is a local  $(a, a)$ -regularized resolvent family generated by  $A$ . Then

$$\begin{aligned} A(a * S_{n+1})(t) &= A \int_0^{nT} a(t-s)S_n(s)x ds + S_{n+1}(t)x \\ &\quad + \int_0^{nT} S_n(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1. \end{aligned}$$

As  $\{S_n(t)\}_{t \in (0, nT]}$  is a local  $(a, a)$ -regularized resolvent family generated by  $A$ , then

$$S_n(r_1)x = A(a * S_n)(r_1)x + a(r_1)x,$$

and

$$\begin{aligned} & \int_0^{nT} S_n(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1 \\ &= \int_0^{nT} (A(a * S_n)(r_1) + a(r_1))x \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1. \end{aligned}$$

On the one hand, we obtain the following identity by change of variables and Fubini's theorem,

$$\begin{aligned}
& A \int_0^{nT} (a * S_n)(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1 \\
&= A \int_0^{nT} \left( \int_0^{r_1} a(r_1-u)S_n(u)x du \right) \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1 \\
&= A \int_0^{nT} S_n(u)x \int_u^{nT} a(r_1-u) \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1 du \\
&= A \int_0^{nT} S_n(u)x \int_0^{nT-u} \int_0^{t-nT} c'(t-u-v-r_2)a(v)a(r_2) dr_2 dv du \\
&= A \int_0^{nT} S_n(u)x((c')^+ *_2(a \otimes a))(nT-u, t-nT) du \\
&= -A \int_0^{nT} a(t-u)S_n(u)x du,
\end{aligned} \tag{5.1}$$

where we have used Theorem 3.4. On the other hand, we use Theorem 3.4 again to get

$$\begin{aligned}
\int_0^{nT} a(r_1)x \int_0^{t-nT} c'(t-r_1-r_2)a(r_2) dr_2 dr_1 &= -((c')^+ *_2(a \otimes a))(t-nT, nT)x \\
&= -a(t)x.
\end{aligned}$$

We join all the terms and we obtain the result.  $\square$

The next result considers the special case of  $(g_\alpha, g_\alpha)$ -regularized resolvent families. Here we have to restrict to the range  $0 < \alpha < 1$  according to the given hypothesis in the above theorem. We observe that this condition is optimal in the following sense: when  $\alpha = 1$ , we are treating with the parabolic case, that is, the equation


$$\begin{cases} u'(t) = Au(t) + x, & t \in [0, \tau), x \in D(A), \\ u(0) = 0, \end{cases}$$

where  $A$  is the generator of a  $C_0$ -semigroup, or, equivalently, a  $(1, 1)$ -regularized resolvent family. We know that in this case no loss of regularity happens. Now, for  $0 < \alpha < 1$ , we have to consider the fractional order differential equation

$$\begin{cases} {}_R D_t^\alpha u(t) = Au(t) + g_\alpha(t)x, & t \in (0, \tau), x \in D(A), \\ (g_{1-\alpha} * u)(0) = 0, \end{cases} \tag{5.2}$$

where  ${}_R D_t^\alpha$  denotes the fractional derivative in the Riemann–Liouville sense, and  $A$  is the generator of a  $(g_\alpha, g_\alpha)$ -regularized resolvent family (see also [11, Example 2.1.38] and the paragraph preceding it). The following corollary shows that again no loss of regularity happens for equation (5.2). In passing, we conclude the remarkable fact that equation (5.2) is at the basis of the process of regularization for  $0 < \alpha < 1$ , where the solutions of the regularized problems correspond to the families of Corollary 4.7.

In the following picture we can see graphically the previous comments for  $(g_\alpha, g_{\beta+1})$ -regularized resolvent families with  $0 < \alpha < 1$ . Note that the straight line formed by the points  $(\alpha, \alpha - 1)$  corresponds to  $(g_\alpha, g_\alpha)$ -regularized families, which is the basis of the process of regularization for  $0 < \alpha < 1$ . For  $\alpha = 1$ , the point  $(1, 0)$  corresponds to a  $C_0$ -semigroup, and the points  $(1, \beta)$  correspond to  $\beta$ -times integrated semigroups for  $\beta > 0$ .



**Corollary 5.2.** *Let  $n \in \mathbb{N}$ , let  $0 < \tau \leq \infty$ , let  $0 < \alpha < 1$ , and let  $\{S_1(t)\}_{t \in (0, \tau)}$  be a local  $(g_\alpha, g_\alpha)$ -regularized resolvent family generated by  $A$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined by*

$$S_{n+1}(t)x = S_n(t)x, \quad x \in X,$$

for  $t \in (0, nT]$  and

$$S_{n+1}(t)x = \frac{\alpha}{\Gamma(1-\alpha)} \int_0^{t-nT} \int_0^{nT} \frac{S_n(r_1)S_1(r_2)x}{(t-r_1-r_2)^{1+\alpha}} dr_1 dr_2$$

for  $x \in X$  and  $t \in (nT, (n+1)T]$  is a local  $(g_\alpha, g_\alpha)$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  generates a global  $(g_\alpha, g_\alpha)$ -regularized resolvent family  $\{S(t)\}_{t \in (0, \infty)}$ .

Now, we consider a different class of  $(a, k)$ -regularized resolvent families such that we can solve the extension problem without loss of regularity.

**Theorem 5.3.** *Let  $n \in \mathbb{N}$ , let  $0 < \tau \leq \infty$ , and let  $a \in L^1_{\text{loc}}(\mathbb{R}_+)$  with  $a \in \mathcal{C}(0, \infty)$  be a Laplace-transformable function such that there exists  $c \in L^1_{\text{loc}}(\mathbb{R}_+)$  Laplace-transformable satisfying the condition that  $c$  is absolutely continuous, differentiable a.e.,  $c(0^+) = 0$ ,  $(c')^+$  and  $(c')^-$  are 2-Laplace-transformable, and  $(a * c)(t) = 1$  for all  $t > 0$ , and let  $\{S_1(t)\}_{t \in (0, \tau)}$  be a local  $(a * 1, a)$ -regularized resolvent family generated by  $A$ . Then the family of operators  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  defined by*

$$S_{n+1}(t)x := S_n(t)x, \quad x \in X,$$

for  $t \in (0, nT]$  and

$$S_{n+1}(t)x := -S_n(2nT - t)x + ((c')^- * {}_2(S_n \otimes S_1))(nT, t - nT)x$$

$$- ((c')^+ *_2 (S_n \otimes S_1))(nT, t - nT)x$$

for  $x \in X$  and  $t \in (nT, (n+1)T]$  is a local  $(a * 1, a)$ -regularized resolvent family generated by  $A$  for any  $T < \tau$ . Then  $A$  generates a global  $(a * 1, a)$ -regularized resolvent family  $\{S(t)\}_{t \in (0, \infty)}$ .

*Proof.* Note that  $\lim_{t \rightarrow 0^+} \frac{S_{n+1}(t)x}{a(t)} = x$  for  $x \in X$  and the family  $\{S_{n+1}(t)\}_{t \in (0, (n+1)T]}$  is strongly continuous (see the proof in Theorem 4.6); in particular,  $\{S_{n+1}(t)\}_{t \in (0, nT]}$  is a local  $(a * 1, a)$ -regularized resolvent family generated by  $A$ . Now let  $t \in (nT, (n+1)T]$  and  $x \in X$ . It is clear that  $S_{n+1}(t)A \subset AS_{n+1}(t)$ . Following the same ideas as in the proofs of the previous theorems, we conclude that  $(a * 1 * S_{n+1})(t)x \in D(A)$ .

To finish the proof, it remains for us to prove that for  $t \in (nT, (n+1)T]$  and  $x \in X$  the equality  $A(a * 1 * S_{n+1}(t))x = S_{n+1}(t)x - a(t)x$  is satisfied. First observe that

$$A(a * 1 * S_{n+1})(t)x = A \int_0^{nT} (a * 1)(t - s)S_n(s)x ds + A \int_{nT}^t (a * 1)(t - s)S_{n+1}(s)x ds.$$

Note that

$$\begin{aligned} & \int_{nT}^t (a * 1)(t - s)S_{n+1}(s)x ds \\ &= \int_{nT}^t (a * 1)(t - s) \left( -S_n(2nT - t)x \right. \\ & \quad + \int_0^{s-nT} \int_0^{nT} (c')^-(nT - r_1, s - nT - r_2)S_n(r_1)S_1(r_2)x dr_1 dr_2 \\ & \quad \left. - \int_0^{s-nT} \int_0^{nT} (c')^+(nT - r_1, s - nT - r_2)S_n(r_1)S_1(r_2)x dr_1 dr_2 \right) ds. \end{aligned}$$

We take the second term and apply the operator  $A$ , to obtain, using change of variables and Fubini's theorem, that

$$\begin{aligned} & A \int_0^{nT} S_n(r_1) \int_{nT}^t (a * 1)(t - s) \\ & \quad \times \int_0^{s-nT} (c')^-(nT - r_1, s - nT - r_2)S_1(r_2)x dr_2 ds dr_1 \\ &= A \int_0^{nT} S_n(r_1) \int_{nT}^t (a * 1)(t - s) \\ & \quad \times \int_{nT}^s (c')^-(nT - r_1, u - nT)S_1(s - u)x du ds dr_1 \\ &= A \int_0^{nT} S_n(r_1) \int_{nT}^t (c')^-(nT - r_1, u - nT) \\ & \quad \times \int_u^t (a * 1)(t - s)S_1(s - u)x ds du dr_1 \end{aligned}$$

$$\begin{aligned}
&= A \int_0^{nT} S_n(r_1) \int_{nT}^t (c')^-(nT - r_1, u - nT) \\
&\quad \times \int_0^{t-u} (a * 1)(t - u - v) S_1(v) x \, dv \, du \, dr_1 \\
&= \int_0^{nT} S_n(r_1) \int_{nT}^t (c')^-(nT - r_1, u - nT) (S_1(t - u) - a(t - u)) x \, du \, dr_1 \\
&= \int_0^{nT} S_n(r_1) \int_0^{t-nT} (c')^-(nT - r_1, t - nT - r_2) (S_1(r_2) - a(r_2)) x \, dr_2 \, dr_1 \\
&= ((c')^- * _2 (S_n \otimes S_1))(nT, t - nT) x \\
&\quad - \int_0^{nT} \int_0^{t-nT} (c')^-(nT - r_1, t - nT - r_2) a(r_2) S_n(r_1) x \, dr_2 \, dr_1,
\end{aligned}$$

where we have used the fact that  $\{S_1(t)\}_{t \in (0, T]}$  is a local  $(a * 1, a)$ -regularized resolvent family generated by  $A$ . Now, observe that, because  $\{S_n(t)\}_{t \in (0, nT]}$  is a local  $(a * 1, a)$ -regularized resolvent family generated by  $A$ , then  $S_n(r_1)x = A(a * 1 * S_n)(r_1)x + a(r_1)x$ , and

$$\begin{aligned}
&- \int_0^{nT} \int_0^{t-nT} (c')^-(nT - r_1, t - nT - r_2) a(r_2) S_n(r_1) x \, dr_2 \, dr_1 \\
&= - \int_0^{nT} \int_0^{t-nT} (c')^-(nT - r_1, t - nT - r_2) a(r_2) \\
&\quad \times (A(a * 1 * S_n)(r_1) + a(r_1)) x \, dr_2 \, dr_1.
\end{aligned}$$

On the one hand,

$$\begin{aligned}
&-A \int_0^{nT} \int_0^{t-nT} (c')^-(nT - r_1, t - nT - r_2) a(r_2) (a * 1 * S_n)(r_1) x \, dr_2 \, dr_1 \\
&= -A \int_0^{nT} (1 * S_n)(u) x \\
&\quad \times \int_0^{nT-u} \int_0^{t-nT} (c')^-(nT - u - v, t - nT - r_2) a(v) a(r_2) \, dr_2 \, dv \, du \\
&= -A \int_0^{nT} ((c')^- * _2 (a \otimes a))(nT - u, t - nT) (1 * S_n)(u) x \, du \\
&= -A \int_0^{nT} a^-(nT - u, t - nT) (1 * S_n)(u) x \, du \\
&= -A \left( \int_0^{2nT-t} a(2nT - t - u) (1 * S_n)(u) x \, du \right. \\
&\quad \left. + \int_{2nT-t}^{nT} a(t + u - 2nT) (1 * S_n)(u) x \, du \right),
\end{aligned}$$

and, on the other hand,

$$\begin{aligned} & - \int_0^{nT} \int_0^{t-nT} (c')^-(nT - r_1, t - nT - r_2) a(r_1) a(r_2) x \, dr_2 \, dr_1 \\ & = -((c')^- *_2 (a \otimes a))(nT, t - nT) x = -a^-(nT, t - nT) x \\ & = -a(2nT - t) x = A(a * 1 * S_n)(2nT - t) x - S_n(2T - t) x, \end{aligned}$$

where we have applied Theorem 3.4(ii) and the fact that  $\{S_n(t)\}_{t \in (0, nT]}$  is a local  $(a * 1, a)$ -regularized resolvent family generated by  $A$ . Then the second term is equal to

$$\begin{aligned} & A \int_0^{nT} S_n(r_1) \int_{nT}^t (a * 1)(t - s) \\ & \quad \times \int_0^{s-nT} (c')^-(nT - r_1, s - nT - r_2) S_1(r_2) x \, dr_2 \, ds \, dr_1 \\ & = ((c')^- *_2 (S_n \otimes S_1))(nT, t - nT) x - S_n(2T - t) x \\ & \quad - A \int_{2nT-t}^{nT} a(t + u - 2nT) (1 * S_n)(u) x \, du. \end{aligned}$$

Similarly, we repeat the process for the third term, and we get

$$\begin{aligned} & -A \int_0^{nT} S_n(r_1) \int_{nT}^t (a * 1)(t - s) \\ & \quad \times \int_0^{s-nT} (c')^+(nT - r_1, s - nT - r_2) S_1(r_2) x \, dr_2 \, ds \, dr_1 \\ & = -((c')^+ *_2 (S_n \otimes S_1))(nT, t - nT) x - a(t) x \\ & \quad - A \int_0^{nT} a(t - u) (1 * S_n)(u) x \, du, \end{aligned}$$

where we have applied Theorem 3.4(i). We join all the terms and we get

$$\begin{aligned} A(a * 1 * S_{n+1})(t) x & = S_{n+1}(t) x - a(t) x + A \left( \int_0^{nT} (a * 1)(t - s) S_n(s) x \, ds \right. \\ & \quad - \int_{nT}^t (a * 1)(t - s) S_n(2T - s) x \, ds \\ & \quad - \int_{2nT-t}^{nT} a(t + u - 2nT) (1 * S_n)(u) x \, du \\ & \quad \left. - \int_0^{nT} a(t - u) (1 * S_n)(u) x \, du \right). \end{aligned}$$

Note that

$$\begin{aligned} & \int_0^{nT} (a * 1)(t - s) S_n(s) x \, ds - \int_0^{nT} a(t - u) (1 * S_n)(u) x \, du \\ & = \int_{t-nT}^t (a * 1)(u) S_n(t - u) x \, du - \int_0^{nT} a(t - u) (1 * S_n)(u) x \, du \end{aligned}$$

$$\begin{aligned}
&= (1 * a)(t - nT)(1 * S_n)(nT)x \\
&= \int_{2nT-t}^{nT} a(t + u - 2nT)(1 * S_n)(u)x \, du - \int_0^{t-nT} (a * 1)(u)S_n(2T - t + u)x \, du \\
&= \int_{2nT-t}^{nT} a(t + u - 2nT)(1 * S_n)(u)x \, du - \int_{nT}^t (a * 1)(t - s)S_n(2T - s)x \, ds,
\end{aligned}$$

where we have used a change of variables and [23, Lemma 2.2] (note that this lemma is true when one of the two functions is a vector valued function). Then we conclude the result.  $\square$

## 6. ALGEBRAIC TIME TRANSLATION IDENTITIES FOR ( $a, k$ )-REGULARIZED RESOLVENT FAMILIES

In this section, applying Laplace transform methods, we solve the problem of time translation in case of global ( $a, k$ )-regularized resolvent families. Under certain conditions on the kernels ( $a, k$ ), we know that Definition 4.1 of ( $a, k$ )-regularized resolvent families is equivalent to the existence of a commutative and strongly continuous family of bounded and linear operators that satisfy  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{k(t)} = x$  for all  $x \in X$  and the functional equation

$$\begin{aligned}
S(s) \int_0^t a(t - \tau)S(\tau)x \, d\tau - S(t) \int_0^s a(s - \tau)S(\tau)x \, d\tau \\
= k(s) \int_0^t a(t - \tau)S(\tau)x \, d\tau - k(t) \int_0^s a(s - \tau)S(\tau)x \, d\tau,
\end{aligned} \tag{6.1}$$

for  $t, s \in (0, \tau)$  (see [17, Theorem 3.1]).

Let  $S(t)$  be an ( $a, k$ )-regularized resolvent family in a Banach space  $X$ . In what follows, we will suppose that the commutative and locally integrable family  $\{S(t)\}_{t>0}$  as well as the kernels  $a, k \in L^1_{\text{loc}}(\mathbb{R}^+)$  are Laplace-transformable, with  $k \in \mathcal{C}(0, \infty)$ . We note that an application of the double Laplace transform to (6.1) gives the following identity which appears in [17, Remark 3.2]:

$$\hat{S}(\lambda)\hat{S}(\mu)x = \frac{\hat{k}(\lambda)}{\hat{a}(\lambda)} \frac{1}{\frac{1}{\hat{a}(\lambda)} - \frac{1}{\hat{a}(\mu)}} \hat{S}(\mu)x - \frac{\hat{k}(\mu)}{\hat{a}(\mu)} \frac{1}{\frac{1}{\hat{a}(\lambda)} - \frac{1}{\hat{a}(\mu)}} \hat{S}(\lambda)x, \tag{6.2}$$

valid for all sufficiently large  $\Re\mu, \Re\lambda$ , and  $x \in X$ . Using the notation in the preceding section, and the above identity, we arrive at the following notable characterization.

**Theorem 6.1.** *A Laplace-transformable and strongly continuous family of bounded and linear operators  $\{S(t)\}_{t>0}$  is an ( $a, k$ )-regularized resolvent family if and only if  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{k(t)} = x$  for all  $x \in X$  and the following functional equation holds:*

$$(a^+ * {}_2(S \otimes S))(t, s)x = k * (a * S)_t(s)x - k_t * (a * S)(s)x, \quad t, s > 0, x \in X. \tag{6.3}$$

*Proof.* From (6.2) we get the equivalent identity

$$(\hat{a}(\mu) - \hat{a}(\lambda))\hat{S}(\lambda)\hat{S}(\mu)x = \hat{k}(\lambda)[\hat{a}(\mu)\hat{S}(\mu)x - \hat{a}(\lambda)\hat{S}(\lambda)x] + \hat{a}(\lambda)[\hat{k}(\lambda) - \hat{k}(\mu)]\hat{S}(\lambda)x$$

valid for all  $\Re\lambda, \Re\mu$  sufficiently large. In turn, the above identity is equivalent to

$$\begin{aligned} \frac{1}{\lambda - \mu} (\hat{a}(\mu) - \hat{a}(\lambda)) \hat{S}(\lambda) \hat{S}(\mu) x &= \frac{1}{\lambda - \mu} \hat{k}(\lambda) [(\widehat{a * S})(\mu) x - (\widehat{a * S})(\lambda) x] \\ &\quad + \frac{1}{\lambda - \mu} [\hat{k}(\lambda) - \hat{k}(\mu)] (\widehat{a * S})(\lambda) x. \end{aligned}$$

Using the identities (3.5) and (3.6), Proposition 3.6(i) and uniqueness of the Laplace transform, we have the result.  $\square$

An interesting particular case is the following corollary, which we quote here for further reference.

**Corollary 6.2.** *A Laplace-transformable and strongly continuous family of bounded and linear operators  $\{S(t)\}_{t>0}$  is an  $(a, a)$ -regularized resolvent family if and only if  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{a(t)} = x$  for all  $x \in X$  and the following functional equation holds:*

$$\begin{aligned} &\int_0^t \int_0^s a(t + s - r_1 - r_2) S(r_1) S(r_2) x \, dr_2 \, dr_1 \\ &= \int_0^t \int_0^s a(r_1) a(r_2) S(t + s - r_1 - r_2) x \, dr_1 \, dr_2 \end{aligned}$$

for all  $t, s > 0$  and  $x \in X$ .

*Proof.* We use Corollary 2.2 in Theorem 6.1 and the result is obtained directly.  $\square$

Our next results have the objective of extending and recovering some of the results mentioned in the Introduction. We will see that in order to do that, we need to impose regularity conditions on the kernels  $a$  and  $k$ , and therefore, the results are less general than our Theorem 6.1 above.

**Theorem 6.3.** *Let  $a, k \in L^1_{\text{loc}}(\mathbb{R}_+)$  be given, with  $k \in \mathcal{C}(0, \infty)$ . Suppose there exist functions  $b, c \in L^1_{\text{loc}}(\mathbb{R}_+)$  Laplace-transformable such that  $c$  is absolutely continuous on  $(0, \infty)$  and  $(c')^+$  is 2-Laplace-transformable, satisfying*

$$(a * b)(t) = k(t), \quad (a * c)(t) = 1, \quad t > 0.$$

*A Laplace-transformable and strongly continuous family of bounded and linear operators  $\{S(t)\}_{t>0}$  is an  $(a, k)$ -regularized resolvent family if and only if  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{k(t)} = x$  for all  $x \in X$  and the following functional equation holds:*

$$((c')^+ * {}_2(S \otimes S))(t, s)x = b_t * S(s)x - b * S_t(s)x, \quad t, s > 0, x \in X.$$

*Proof.* From (6.2) we obtain the equivalent identity

$$\begin{aligned} &\frac{1}{\mu - \lambda} (\lambda \hat{c}(\lambda) - \mu \hat{c}(\mu)) \hat{S}(\lambda) \hat{S}(\mu) x \\ &= \frac{1}{\mu - \lambda} \hat{b}(\lambda) \hat{S}(\mu) x - \frac{1}{\mu - \lambda} \hat{b}(\mu) \hat{S}(\lambda) x \\ &= \frac{1}{\mu - \lambda} \hat{b}(\lambda) [\hat{S}(\mu) x - \hat{S}(\lambda) x] + \frac{1}{\mu - \lambda} [\hat{b}(\lambda) - \hat{b}(\mu)] \hat{S}(\lambda) x. \end{aligned}$$



Hence, the result follows from Corollary 3.7(i) and formulas (3.5) and (3.6).  $\square$

*Example 6.4.* We set  $a := g_\alpha$  for  $0 < \alpha < 1$  and  $k(t) := \int_0^t K(s) ds$  for  $t > 0$ . In this case we can choose  $c = g_{1-\alpha}$  and  $b = g_{1-\alpha} * K$  satisfying the hypothesis. Therefore, we recover the functional equation

$$\begin{aligned} & \left( \int_t^{t+s} - \int_0^s \right) (g_{1-\alpha} * K)(t+s-\sigma) S(\sigma) x d\sigma \\ &= \alpha \int_0^t \int_0^s \frac{S(r_1) S(r_2) x}{(t+s-r_1-r_2)^{1+\alpha}} dr_1 dr_2, \end{aligned} \quad (6.4)$$

which appeared in [18, Theorem 8] and was mentioned in the Introduction. Here, we include as particular case the identity

$$\left( \int_t^{t+s} - \int_0^s \right) \frac{S(\tau) x}{(t+s-\tau)^\alpha} d\tau = \alpha \int_0^t \int_0^s \frac{S(r_1) S(r_2) x}{(t+s-r_1-r_2)^{1+\alpha}} dr_1 dr_2 \quad (6.5)$$

(see [13, Definition 3] and the more general identity developed in [15, Theorem 5]).

*Example 6.5.* Let  $a := g_\alpha$  and  $k := g_{\beta+1}$  where  $\alpha > 0, \beta > -1$ . This choosing of the pair  $(a, k)$  produces the theory of  $(\alpha, \beta)$ -ROF families introduced in [4] (see also [17, Example 3.10] for a more general approach in terms of  $(a, k)$ -regularized resolvent families). As mentioned in the introduction, a time translation formula for  $(\alpha, \beta)$ -ROF families was developed recently in [15]. Now observe that for  $a$  and  $k$  as before, we can choose  $c = g_{1-\alpha}$  whenever  $0 < \alpha < 1$ , and  $b = g_{\beta-\alpha+1}$  whenever  $\beta - \alpha > -1$  to obtain

$$a * c = g_\alpha * g_{1-\alpha} = 1 \quad \text{and} \quad a * b = g_\alpha * g_{\beta-\alpha+1} = g_{\beta+1}.$$

Therefore, the hypothesis of Theorem 6.3 are satisfied and, in consequence, we recover the formula

$$\begin{aligned} & \left( \int_t^{t+s} \int_0^s \right) (s+t-r)^{\beta-\alpha} S(r) x dr \\ &= \alpha \frac{\Gamma(\beta-\alpha+1)}{\Gamma(1-\alpha)} \int_0^t \int_0^s \frac{S(r_1) S(r_2) x}{(t+s-r_1-r_2)^{1+\alpha}} dr_1 dr_2, \end{aligned} \quad (6.6)$$

whenever  $\alpha \in \mathbb{R}_+ \setminus \mathbb{N}_0, \beta \in \mathbb{R}_+$  and  $\beta - \alpha > -1$  discovered in [15, Theorem 5], but now under the restrictions  $0 < \alpha < 1$  and  $\beta - \alpha > -1$ . We observe that our result correct the above formula where a more relaxed condition on  $\alpha$  is assumed, namely  $\alpha \in \mathbb{R}_+ \setminus \mathbb{N}_0$ . However, we note that for  $\alpha \geq 1$  the double integral on the right hand side of (6.6) diverges, as can be easily seen.

Our next result widely extends the well known semigroup functional equation to a more general class of strongly continuous families of operators. They are connected with integral equations of Volterra-type, as we will see in the next section.

**Theorem 6.6.** *Let  $a \in L_{\text{loc}}^1(\mathbb{R}_+)$  be given, with  $a \in \mathcal{C}(0, \infty)$ . Suppose there exists  $c \in L_{\text{loc}}^1(\mathbb{R}_+)$  Laplace-transformable function such that  $c$  is absolutely continuous on  $(0, \infty)$  and such that  $(c')^+$  is 2-Laplace-transformable, satisfying  $(a * c)(t) = 1$  for all  $t > 0$ . A Laplace-transformable and strongly continuous family of bounded*

and linear operators  $\{S(t)\}_{t>0}$  is an  $(a, a)$ -regularized resolvent family if and only if  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{a(t)} = x$  for all  $x \in X$  and the following functional equation holds:

$$S(t+s)x = -((c')^+ *_2 (S \otimes S))(t, s)x, \quad t, s > 0, x \in X.$$

*Proof.* From (6.2) we get the equivalent identity

$$\frac{1}{\lambda - \mu} (\lambda \hat{c}(\lambda) - \mu \hat{c}(\mu)) \hat{S}(\lambda) \hat{S}(\mu)x = \frac{1}{\lambda - \mu} (\hat{S}(\mu)x - \hat{S}(\lambda)x).$$

Hence, the result follows from Corollary 3.7(i) and formula (3.4).  $\square$

*Example 6.7.* If  $a = g_\alpha$ , then  $\hat{a}(\lambda) = \frac{1}{\lambda^\alpha}$  and hence we can choose  $c = g_{1-\alpha}$  with  $0 < \alpha < 1$  satisfying the hypothesis. Note that this example recovers the functional equation

$$S(t+s)x = \frac{\alpha}{\Gamma(1-\alpha)} \int_0^t \int_0^s \frac{S(r_1)S(r_2)x}{(t+s-r_1-r_2)^{1+\alpha}} dr_1 dr_2, \quad (6.7)$$

for  $0 < \alpha < 1$  as stated in [20, Definition 2.1(ii)].

Now we consider more relaxed assumptions on the functions  $a$  and  $k$  than the previous theorems. In contrast, the obtained functional equations are more involved and difficult to handle. The advantage is that it permits us to extend the range from  $0 < \alpha < 1$  to  $1 < \alpha < 2$ , extending recent results and producing new formulas in cases where they were not known.

**Theorem 6.8.** *Let  $a, k \in L^1_{\text{loc}}(\mathbb{R}_+)$  be given, with  $k \in \mathcal{C}(0, \infty)$ . Suppose there exist  $b, c \in L^1_{\text{loc}}(\mathbb{R}_+)$  Laplace-transformable functions satisfying  $(a * c)(t) = t$  and  $(a * b)(t) = (1 * k)(t)$  for all  $t > 0$ . A Laplace-transformable and strongly continuous family of bounded and linear operators  $\{S(t)\}_{t>0}$  is an  $(a, k)$ -regularized resolvent family if and only if  $\lim_{t \rightarrow 0^+} \frac{S(t)x}{k(t)} = x$  for all  $x \in X$  and the following functional equation holds:*

$$\begin{aligned} & b * (1 * S)_t(s)x - b_t * (1 * S)(s)x \\ &= (c * S)(t)(1 * S)(s)x \\ &+ (1 * S)(t)(c * S)(s)x - (c^+ *_2 (S \otimes S))(t, s)x, \end{aligned}$$

for  $t, s > 0$ , and  $x \in X$ .

*Proof.* From (6.2) we obtain the equivalent identity

$$\frac{\hat{k}(\lambda)}{\hat{a}(\lambda)} \hat{S}(\mu)x - \frac{\hat{k}(\mu)}{\hat{a}(\mu)} \hat{S}(\lambda)x = \left( \frac{1}{\hat{a}(\lambda)} - \frac{1}{\hat{a}(\mu)} \right) \hat{S}(\lambda) \hat{S}(\mu)x.$$

We multiply the identity by  $\frac{1}{(\lambda-\mu)\lambda\mu}$ , and we obtain that

$$\begin{aligned} \frac{1}{\lambda - \mu} \left( \frac{\hat{b}(\lambda)}{\mu} \hat{S}(\mu)x - \frac{\hat{b}(\mu)}{\lambda} \hat{S}(\lambda)x \right) &= \left( \frac{\hat{c}(\lambda)}{\mu} + \frac{\hat{c}(\mu)}{\lambda} \right) \hat{S}(\lambda) \hat{S}(\mu)x \\ &- \frac{1}{\lambda - \mu} (\hat{c}(\mu) - \hat{c}(\lambda)) \hat{S}(\lambda) \hat{S}(\mu)x. \end{aligned}$$

Note that

$$\begin{aligned} & \frac{1}{\lambda - \mu} \left( \frac{\hat{b}(\lambda)}{\mu} \hat{S}(\mu)x - \frac{\hat{b}(\mu)}{\lambda} \hat{S}(\lambda)x \right) \\ &= \frac{1}{\lambda - \mu} \left( \hat{b}(\lambda) (\widehat{(1 * S)}(\mu)x) - \widehat{(1 * S)}(\lambda)x - (\hat{b}(\mu) - \hat{b}(\lambda)) \widehat{(1 * S)}(\lambda)x \right). \end{aligned}$$

The result follows from Proposition 3.6(i) and formulas (3.5) and (3.6).  $\square$

*Example 6.9.* Set  $a = g_\alpha$ , with  $1 < \alpha < 2$  and  $k = g_1$ . In this case, we can choose  $b = c = g_{2-\alpha}$  to satisfy  $a * c = g_2$  and  $a * b = 1 * k$ . Therefore we recover the functional equation

$$\begin{aligned} & \left( \int_t^{t+s} - \int_0^s \right) \int_0^\sigma \frac{S(\tau)x}{(t+s-\sigma)^{\alpha-1}} d\tau d\sigma \\ &= \int_0^t \int_0^s \frac{S(\sigma)S(\tau)x}{(t-\sigma)^{\alpha-1}} d\tau d\sigma \\ &+ \int_0^t \int_0^s \frac{S(\sigma)S(\tau)x}{(s-\tau)^{\alpha-1}} d\tau d\sigma - \int_0^t \int_0^s \frac{S(\sigma)S(\tau)x}{(t+s-\sigma-\tau)^{\alpha-1}} d\tau d\sigma \end{aligned} \quad (6.8)$$

developed in [19, Definition 3.1] and cited in the introduction.

*Example 6.10.* Let  $a = g_\alpha$  and  $k = g_{\beta+1}$  where  $\alpha > 0, \beta > -1$ . In this case we obtain a new functional equation for  $(\alpha, \beta)$ -ROF families (introduced in [4]) in contrast with those developed in [15]. See also Example 6.5 for a correction on the assumptions on  $\alpha$  and  $\beta$ . Indeed, we can choose  $c = g_{2-\alpha}$  and  $b = g_{\beta-\alpha+2}$  under the assumptions  $0 < \alpha < 2$  and  $\beta - \alpha > -2$ .

## 7. EXAMPLES, APPLICATIONS, AND FINAL COMMENTS

**7.1. Multiplication local regularized families in  $L^p(\mathbb{R})$ .** We consider the Lebesgue space  $L^p(\mathbb{R})$ ,  $1 \leq p \leq \infty$ , and  $a = g_\alpha$  with  $\alpha \in (0, 2)$ . Define the multiplication operator

$$Af(x) := (1 + x + ix^2)^\alpha, \quad x \in \mathbb{R}, f \in L^p(\mathbb{R}),$$

with maximal domain in  $L^p(\mathbb{R})$ . Assume  $s \in (1, 2)$ ,  $\delta = \frac{1}{s}$  and  $K_\delta(t) = \mathcal{L}^{-1}(e^{-\lambda^\delta})(t)$ ,  $t > 0$ , where  $\mathcal{L}^{-1}$  is the inverse Laplace transform. Then  $A$  generates a global  $(a, K_\delta)$ -regularized resolvent family in  $L^p(\mathbb{R})$ . Furthermore, when  $s = 2$  there exists  $\tau > 0$  such that  $A$  generates a local  $(a, K_{\frac{1}{2}})$ -regularized resolvent family on  $[0, \tau)$  (see [9, Example 2.31]). Then we can apply Theorem 4.3, and conclude that  $A$  generates a local  $(a, (K_{\frac{1}{2}} * a)^{*n} * K_{\frac{1}{2}})$ -regularized resolvent family on  $(0, (n+1)\tau)$ , for all  $n \in \mathbb{N}$ .

**7.2. Local regularized families in sequence spaces.** Let  $l^2(\mathbb{N}) = \{x = (x_m)_{m=1}^\infty \subset \mathbb{C} : \sum_{m=1}^\infty |x_m|^2 < \infty\}$  be the Hilbert space of all square-summable sequences with the norm  $\|x\| = (\sum_{m=1}^\infty |x_m|^2)^{\frac{1}{2}}$ . We take  $\tau > 0$ , and

$$a_m = \frac{m}{\tau} + i \left( \left( \frac{e^m}{m} \right)^2 - \left( \frac{m}{\tau} \right)^2 \right)^{\frac{1}{2}}, \quad m \in \mathbb{N},$$

where  $i$  is the imaginary identity. We note that for all  $n \in \mathbb{N}$ , the sequence  $(a_m)_{m=1}^\infty$  generates a local  $n$ -times integrated semigroup on  $l^2(\mathbb{N})$  for  $t \in [0, n\tau)$ , see [28, pp. 75–76].

The Mittag–Leffler functions are defined by

$$E_{\alpha,\beta}(z) := \sum_{n=0}^\infty \frac{z^n}{\Gamma(\alpha n + \beta)}, \quad \alpha, \beta > 0, z \in \mathbb{C}.$$

For short,  $E_\alpha := E_{\alpha,1}$ . We take  $0 < \alpha < 2$ . Observe that the function  $E_\alpha$  is a  $(g_\alpha, 1)$ -regularized resolvent family.

For any  $\beta \in \mathbb{R}^+$ ,  $(T_{\alpha,\beta}(t))_{t \in (0, \beta\tau)}$ , defined by

$$T_{\alpha,\beta}(t)x = \left( \frac{1}{\Gamma(\beta)} \int_0^t (t-s)^{\beta-1} E_\alpha((a_m s)^\alpha) x_m ds \right)_{m=1}^\infty, \quad \text{for } x \in l^2(\mathbb{N}),$$

is a local  $(g_\alpha, g_{\beta+1})$ -regularized resolvent family on  $l^2(\mathbb{N})$ :

Note that  $a_m \in \mathbb{C}^+$ , the set of imaginary numbers with positive real part. Then for all  $s \geq 0$  and  $0 < \alpha < 2$ ,  $|\arg(a_m s)^\alpha| \leq \frac{\alpha\pi}{2}$ . So, the asymptotic expansion in [3, (1.27)] and the continuity of the Mittag–Leffler function imply that there are constants  $c, C$  such that  $ce^{a_m s} \leq E_\alpha((a_m s)^\alpha) \leq Ce^{a_m s}$ . Observe that

$$\|T_{\alpha,\beta}(t)\| = \sup_{m \in \mathbb{N}} \left| \frac{1}{\Gamma(\beta)} \int_0^t (t-s)^{\beta-1} E_\alpha((a_m s)^\alpha) ds \right| < \infty$$

if and only if

$$\sup_{m \in \mathbb{N}} \left| \frac{1}{\Gamma(\beta)} \int_0^t (t-s)^{\beta-1} e^{a_m s} ds \right| < \infty,$$

which happens if only if  $0 \leq t < \beta\tau$  (see [22, Example 1]). It is clear that  $\{T_{\alpha,\beta}(t)\}_{t \in (0, \beta\tau)}$  is strongly continuous and verifies any functional equation associated to  $(g_\alpha, g_{\beta+1})$ -regularized families. The case  $\alpha = 1$  is made in [22].

**7.3. A new class of regularized families without jumps of regularity.** Let  $0 < \tau \leq \infty$  and  $b \in L^1_{\text{loc}}(\mathbb{R}^+)$ . We define  $a = b * b$ . Suppose that  $\{R_1(t)\}_{t \in (0, \tau)}$  is a local  $(a, a)$ -regularized resolvent family generated by  $A$ , such that  $A$  verifies condition (H5) of [9] (e.g.,  $A$  densely defined). Then, by [9, Theorem 2.34], we have that

$$\mathcal{A} \equiv \begin{pmatrix} 0 & I \\ A & 0 \end{pmatrix}$$

is the generator of a local  $(b, b^{*3})$ -regularized resolvent family  $\{S_1(t)\}_{t \in (0, \tau)}$  given explicitly by

$$S_1(t) = \begin{pmatrix} (b * R_1)(t) & (a * R_1)(t) \\ R_1(t) - a(t)I & (b * R_1)(t) \end{pmatrix}, \quad 0 < t < \tau.$$

Now, we suppose that there is a  $c \in L^1_{\text{loc}}(\mathbb{R}^+)$  Laplace-transformable such that  $c$  is absolutely continuous, differentiable a.e., and such that  $(c')^+$  is 2-Laplace-transformable, and satisfying  $(a * c)(t) = 1$  for all  $t > 0$ . Then we conclude that  $A$  generates a global  $(a, a)$ -regularized resolvent family  $\{R(t)\}_{t \in (0, \infty)}$  which extends  $\{R_1(t)\}_{t \in (0, \tau)}$ , see Theorem 5.1. Then, we can extend  $\{S_1(t)\}_{t \in (0, \tau)}$  without loss

of regularity; that is,  $\mathcal{A}$  generates a global  $(b, b^{*3})$ -regularized resolvent family  $\{S(t)\}_{t \in (0, \infty)}$  given by

$$S(t) = \begin{pmatrix} (b * R)(t) & (a * R)(t) \\ R(t) - a(t)I & (b * R)(t) \end{pmatrix}, \quad 0 < t.$$

In the particular, case  $b = g_{\frac{\alpha}{2}}$  with  $0 < \alpha < 1$ ,  $a = g_{\alpha}$  and  $\{R_1(t)\}_{t \in (0, \tau)}$  be a local  $(g_{\alpha}, g_{\alpha})$ -regularized resolvent family generated by  $A$ , we can extend  $\{S_1(t)\}_{t \in (0, \tau)}$  without loss of regularity; that is,  $\mathcal{A}$  generates a global  $(g_{\frac{\alpha}{2}}, g_{\frac{3\alpha}{2}})$ -regularized resolvent family  $\{S(t)\}_{t > 0}$  such that  $S(t) = S_1(t)$  for  $0 < t < \tau$ .

**7.4. Applications to obtain new functional equations.** We give several examples of the abstract results in Section 6. They show that we can recover, extend and produce new functional equations that in some cases are interesting for their own nature.

*Example 7.1* ( $C_0$ -semigroups). Choose  $a(t) = 1$  and  $k(t) = 1$  for  $t > 0$ . Then, for  $x \in X$ , we obtain

$$\begin{aligned} & \int_0^t \int_0^s S(\tau_1)S(\tau_2)x \, d\tau_2 \, d\tau_1 \\ &= \int_t^{t+s} \int_0^r S(\tau)x \, d\tau \, dr - \int_0^s \int_0^r S(\tau)x \, d\tau \, dr, \quad t, s > 0. \end{aligned}$$

Taking the derivative one time with respect to the variable  $t$  we obtain

$$S(t) \int_0^s S(\tau_2)x \, d\tau_2 = \int_0^{t+s} S(\tau)x \, d\tau - \int_0^t S(\tau)x \, d\tau, \quad t, s > 0,$$

which was introduced in [17, Example 3.4]. Hence taking the derivative with respect to the variable  $s$  we get the Cauchy formula  $S(s)S(t) = S(t+s)$ . Observe that in this way we deduce easily Cauchy's functional equation from the formula in Theorem 6.1.

*Example 7.2* (Cosine families). Choose  $a(t) = t$  and  $k(t) = 1$  for  $t > 0$ . Then, for  $x \in X$  and  $t, s > 0$  we have

$$\int_0^t \int_0^s (t+s-r_1-r_2)S(r_1)S(r_2)x \, dr_2 \, dr_1 = \left( \int_t^{t+s} - \int_0^s \right) \int_0^r (r-\tau)S(\tau)x \, d\tau \, dr.$$

To see directly why the above formula is equivalent to the D'Alembert functional equation  $S(t+s) + S(|t-s|) = 2S(t)S(s)$  we proceed as in the above example, first taking derivative with respect to the variable  $t$ , to obtain

$$\begin{aligned} & S(t) \int_0^s (s-r_2)S(r_2)x \, dr_2 + \int_0^t S(r_1) \int_0^s S(r_2)x \, dr_2 \, dr_1 \\ &= \int_0^{t+s} (t+s-\tau)S(\tau)x \, d\tau - \int_0^t (t-\tau)S(\tau)x \, d\tau, \end{aligned}$$

and then taking derivative with respect to the variable  $s$ , to have

$$S(t) \int_0^s S(r_2)x \, dr_2 + \int_0^t S(r_1)S(s)x \, dr_1 = \int_0^{t+s} S(\tau)x \, d\tau, \quad t, s > 0.$$

By [21, Theorem 2], the above functional equation is equivalent to the cosine functional equation.

*Example 7.3* (Convoluted semigroups). Choosing  $a(t) = 1$  for  $t > 0$  and  $k \in C^2(\mathbb{R}_+)$  and proceeding as in the above examples, we obtain a new functional equation for convoluted semigroups:

$$\begin{aligned} S(t)S(s)x &= k(0)S(t+s)x + k'(0) \int_0^{t+s} S(\tau)x \, d\tau - k'(s) \int_0^t S(\tau)x \, d\tau \\ &\quad - k'(t) \int_0^s S(\tau)x \, d\tau + \int_t^{t+s} k''(t+s-r) \int_0^r S(\tau)x \, d\tau \, dr \\ &\quad - \int_0^s k''(t+s-r) \int_0^r S(\tau)x \, d\tau \, dr, \end{aligned}$$

for  $t, s > 0$  and  $x \in X$ . Setting  $k(t) := (1 - \epsilon) + \epsilon t$  for  $0 \leq \epsilon \leq 1$ , and  $t \geq 0$ , this formula shows an interesting fact: How continuously moves the functional equation from the case of semigroups to the case of 1-times integrated semigroup as  $\epsilon$  varies from 0 to 1:

$$S(t)S(s)x = (1 - \epsilon)S(t+s)x + \epsilon \left( \int_t^{t+s} - \int_0^s \right) S(\tau)x \, d\tau, \quad t, s > 0, x \in X.$$

*Example 7.4* (Resolvent families). Now we take  $k(t) = 1$  for  $t > 0$ ;  $a \in C^2(\mathbb{R}_+)$  and we proceed as above. We get this (new) functional equation:

$$\begin{aligned} a(0)S(t)S(s)x &+ S(t) \int_0^s a'(s-\tau)S(\tau)x \, d\tau + S(s) \int_0^t a'(t-\tau)S(\tau)x \, d\tau \\ &+ 2 \int_0^t \int_0^s a''(t+s-r_1-r_2)S(r_1)S(r_2)x \, dr_2 \, dr_1 \\ &= a(0)S(t+s)x + \int_0^s a'(t+s-\tau)S(\tau)x \, d\tau, \end{aligned}$$

for  $x \in X$ . For  $0 \leq \epsilon \leq 1$ , we set  $a(t) := (1 - \epsilon) + \epsilon t$ , and we see how the formula continuously moves from the semigroup to the cosine family cases when  $\epsilon$  goes from 0 to 1:

$$(1 - \epsilon)[S(t)S(s)x - S(t+s)x] = \epsilon \left[ \int_0^{t+s} S(\tau)x \, d\tau - \int_0^s S(\tau)x \, d\tau - \int_0^t S(\tau)x \, d\tau \right],$$

for  $x \in X$ . Note the intriguing case  $\epsilon = 1/2$  where the difference between both functional equations is the same, which indicates that in some suitable norm the study of the topology of the set of all functional equations satisfying (1.4) should be relevant.

**Acknowledgments.** We thank the anonymous referees for several comments, ideas, and references which have helped improve the final version of this article.

Abadias's work and Miana's work have been partially supported by a grant from Project MTM2013-42105-P (DGI-FEDER) of the MCYTTS, by a grant from Project E-64, D. G. Aragón, and by a grant UZCUD2014-CIE-09 from the Universidad de Zaragoza, Spain. Lizama's work has been partially supported by a grant

from DICYT of the Universidad de Santiago de Chile Project CONICYT-PIA ACT1112 Stochastic Analysis Research Network, by a grant from FONDECYT 1140258, and by a grant from the Ministerio de Educación CEI Iberus (Spain).

## REFERENCES

1. W. Arendt, C. J. K. Batty, M. Hieber, and F. Neubrander, *Vector-valued Laplace Transforms and Cauchy Problems*, 2nd ed., Monogr. Math. **96**, Birkhäuser, Basel, 2011. [MR2798103](#). [DOI 10.1007/978-3-0348-0087-7](#). [170](#), [175](#), [176](#), [178](#)
2. W. Arendt, O. El-Mennaoui, and V. Kéyantuo, *Local integrated semigroups: Evolution with jumps of regularity*, J. Math. Anal. Appl. **186** (1994), no. 2, 572–595. [Zbl 0833.47034](#). [MR1293012](#). [DOI 10.1006/jmaa.1994.1318](#). [170](#), [171](#), [181](#)
3. E. Bajlekova, *Fractional evolution equations in Banach spaces*, Ph.D. dissertation, Technische Universiteit Eindhoven, Eindhoven, The Netherlands, 2001. [Zbl 0989.34002](#). [MR1868564](#). [170](#), [204](#)
4. C. Chen and M. Li, *On fractional resolvent operator functions*, Semigroup Forum **80** (2010), no. 1, 121–142. [Zbl 1185.47040](#). [MR2580614](#). [DOI 10.1007/s00233-009-9184-7](#). [201](#), [203](#)
5. V. Ditkine and A. Proudnikov, *Calcul opérationnel*, 2nd ed., Mir, Moscow, 1983. [MR0734900](#). [172](#), [173](#), [176](#), [178](#)
6. J. C. Jaeger, *The solution of boundary value problems by a double Laplace transformation*, Bull. Amer. Math. Soc. **46** (1940), 687–693. [MR0002453](#). [172](#)
7. V. Keyantuo, C. Lizama, and P. J. Miana, *Algebra homomorphisms defined via convoluted semigroups and cosine functions*, J. Funct. Anal. **257** (2009), no. 11, 3454–3487. [Zbl 1190.47045](#). [MR2571434](#). [DOI 10.1016/j.jfa.2009.07.017](#). [170](#), [179](#)
8. V. Keyantuo, P. J. Miana, and L. Sánchez-Lajusticia, *Sharp extensions for convoluted solutions of abstract Cauchy problems*, Integral Equations Operator Theory **77** (2013), no. 2, 211–241. [Zbl 1301.47064](#). [MR3101782](#). [DOI 10.1007/s00020-013-2076-y](#). [170](#), [171](#), [175](#), [181](#), [186](#), [191](#)
9. M. Kostić, *(a, k)-regularized C-resolvent families: Regularity and local properties*, Abstr. Appl. Anal. (2009), article ID 858242. [MR2533573](#). [DOI 10.1155/2009/858242](#). [170](#), [203](#), [204](#)
10. M. Kostić, *Generalized Semigroups and Cosine Functions*, Matematički Institut SANU, Belgrade, 2011. [MR2790849](#). [170](#), [171](#)
11. M. Kostić, *Abstract Volterra Integro-differential Equations*, CRC, Boca Raton, New York, 2015. [170](#), [194](#)
12. K. Li and J. Peng, *Fractional abstract Cauchy problems*, Integral Equations Operator Theory **70** (2011), no. 3, 333–361. [Zbl 1242.34008](#). [MR2806472](#). [DOI 10.1007/s00020-011-1864-5](#). [170](#)
13. K. Li and J. G. Peng, *A novel characteristic of solution operator for the fractional abstract Cauchy problem*, J. Math. Anal. Appl. **385** (2012), no. 2, 786–796. [Zbl 1231.47039](#). [MR2834853](#). [DOI 10.1016/j.jmaa.2011.07.009](#). [172](#), [180](#), [201](#)
14. M. Li, C. Chen, and F. B. Li, *On fractional powers of generators of fractional resolvent families*, J. Funct. Anal. **259** (2010), no. 10, 2702–2726. [Zbl 1203.47021](#). [MR2679023](#). [DOI 10.1016/j.jfa.2010.07.007](#). [170](#)
15. Y.-N. Li and H.-R. Sun, *Integrated fractional resolvent operator function and fractional abstract Cauchy problem*, Abstr. Appl. Anal. **2014**, article ID 430418. [MR3166612](#). [DOI 10.1155/2014/430418](#). [170](#), [172](#), [201](#), [203](#)
16. C. Lizama, *Regularized solutions for abstract Volterra equations*, J. Math. Anal. Appl. **243** (2000), no. 2, 278–292. [Zbl 0952.45005](#). [MR1741524](#). [DOI 10.1006/jmaa.1999.6668](#). [170](#), [182](#), [187](#)
17. C. Lizama and F. Poblete, *On a functional equation associated with (a, k)-regularized resolvent families*, Abstr. Appl. Anal. **2012**, article ID 495487. [MR2947741](#). [170](#), [171](#), [199](#), [201](#), [205](#)



18. Z.-D. Mei, J.-G. Peng, and J.-H. Gao, *Convolved fractional  $C$ -semigroups and fractional abstract Cauchy problems*, Abstr. Appl. Anal. **2014**, article ID 357821. [MR3166602](#). DOI [10.1155/2014/357821](#). 170, 172, 201
19. Z.-D. Mei, J.-G. Peng, and J.-X. Jia, *A new characteristic property of Mittag-Leffler functions and fractional cosine functions*, Studia Math. **220** (2014), no. 2, 119–140. [MR3168756](#). DOI [10.4064/sm220-2-2](#). 172, 203
20. Z.-D. Mei, J.-G. Peng, and Y. Zhang, *A characteristic of fractional resolvents*, Fract. Calc. Appl. Anal. **16** (2013), no. 4, 777–790. [MR3124335](#). DOI [10.2478/s13540-013-0048-1](#). 172, 202
21. Z.-D. Mei, J.-G. Peng, and Y. Zhang, *On a characteristic of cosine functions*, Semigroup Forum **88** (2014), no. 1, 221–226. [MR3164161](#). DOI [10.1007/s00233-013-9524-5](#). 206
22. P. J. Miana, *Local and global solutions of well-posed integrated Cauchy problems*, Studia Math. **187** (2008), no. 3, 219–232. [Zbl 1156.47044](#). [MR2417455](#). DOI [10.4064/sm187-3-2](#). 181, 191, 204
23. P. J. Miana and V. Poblete, *Sharp extensions for convolved solutions of wave equations*, Differential Integral Equations **28** (2015), nos. 3–4, 309–332. [MR3306565](#). 170, 171, 181, 186, 199
24. I. Podlubny, *Fractional Differential Equations*, Math. Sci. Eng. **198**, Academic, San Diego, 1999. [MR1658022](#). 170
25. J. Prüss, *Evolutionary Integral Equations and Applications*, Monogr. Math. **87**, Birkhäuser, Basel, 1993. [MR1238939](#). DOI [10.1007/978-3-0348-8570-6](#). 170
26. I. H. Sneddon, *The Use of Integral Transforms*, McGraw-Hill, New York, 1972. [Zbl 0237.44001](#). 172, 173, 176
27. E. Strouse, *Closed ideals in convolution algebras and the Laplace transform*, Michigan Math. J. **35** (1988), no. 2, 185–196. [Zbl 0684.46045](#). [MR0959266](#). DOI [10.1307/mmj/1029003746](#). 173
28. N. Tanaka and N. Okazawa, *Local  $C$ -semigroups and local integrated semigroups*, Proc. London Math. Soc. (3) **61** (1990), no. 1, 63–90. [MR1051099](#). DOI [10.1112/plms/s3-61.1.63](#). 204
29. S. W. Wang and M. C. Gao, *Automatic extensions of local regularized semigroups and local regularized cosine functions*, Proc. Am. Math. Soc. **127** (1999), no. 6, 1651–1663. [Zbl 0916.47031](#). [MR1600157](#). DOI [10.1090/S0002-9939-99-04859-5](#). 170, 171

<sup>1</sup>DEPARTAMENTO DE MATEMÁTICAS, INSTITUTO UNIVERSITARIO DE MATEMÁTICAS Y APLICACIONES, UNIVERSIDAD DE ZARAGOZA, ZARAGOZA, SPAIN.

*E-mail address:* [labadias@unizar.es](mailto:labadias@unizar.es)

*E-mail address:* [pjmiana@unizar.es](mailto:pjmiana@unizar.es)

<sup>2</sup>DEPARTAMENTO DE MATEMÁTICA Y CIENCIA DE LA COMPUTACIÓN, UNIVERSIDAD DE SANTIAGO DE CHILE, CASILLA 307-CORREO 2, SANTIAGO-CHILE, CHILE.

*E-mail address:* [carlos.lizama@usach.cl](mailto:carlos.lizama@usach.cl)