

## THE EVENTUAL STABILITY OF DEPTH, ASSOCIATED PRIMES AND COHOMOLOGY OF A GRADED MODULE

MARC CHARDIN, JEAN-PIERRE JOUANOLOU AND AHAD RAHIMI

**1. Introduction.** The asymptotic stability of several homological invariants of graded pieces of a graded module has attracted quite a lot of attention over the last decades. An early important result was the proof by Brodmann of the eventual stabilization of associated primes of the powers of an ideal in a Noetherian ring ([1]).

We provide in this text several stability results together with estimates of the degree from which it stabilizes. One of our initial goals was to obtain a simple proof of the tameness result of Brodmann in [2] for graded components of cohomology over rings of dimension at most two. This is achieved in the last section and gives a slight generalization of what is known, as our result (Theorem 7.4) applies to Noetherian rings of dimension at most two that are either local or the epimorphic image of a Gorenstein ring. Recall that Cutkosky and Herzog provided examples in [3] showing that tameness does not hold over rings of dimension three (even over such nice local rings).

Besides this result, we establish, for a graded module  $M$  over a polynomial ring  $S$  (in finitely many variables, with its standard grading) over a commutative ring  $R$ , stability results for the depth and cohomological dimension of graded pieces with respect to a finitely generated  $R$ -ideal  $I$ . It follows from our results that the cohomological dimension of  $M_\mu$  with respect to  $I$  is constant for  $\mu > \text{reg}(M)$ , and the depth with respect to  $I$  is at least equal to its eventual value for  $\mu > \text{reg}(M)$  and stabilizes when it reaches this value for some  $M_\mu$  with  $\mu > \text{reg}(M)$ . See Propositions 3.1 and 4.9 for more precise results.

Recall that  $\text{reg}(M) \in \mathbf{Z}$  when  $M \neq 0$  is finitely generated and  $R$  is Noetherian.

When  $R$  is Noetherian,  $\mathfrak{p} \in \text{Spec}(R)$  is associated to  $M_\mu$  for some  $\mu$  if and only if  $\mathfrak{p} = \mathfrak{P} \cap R$  for  $\mathfrak{P}$  associated to  $M$  in  $S$ , and the sets of

---

Received by the editors on March 13, 2012, and in revised form on August 1, 2012.

DOI:10.1216/JCA-2013-5-1-63 Copyright ©2013 Rocky Mountain Mathematics Consortium

associated primes of  $M_\mu$  are non decreasing for  $\mu > \text{reg}(M)$ . It implies that this set eventually stabilizes when  $M$  is finitely generated.

Before we establish these regularity results in Sections 4 and 5, we prove several facts about depth and cohomological dimension with respect to a finitely generated ideal and about Castelnuovo-Mumford regularity of a graded module. Our definition of depth agrees with the one introduced by Northcott. These results are stated in a quite general setting, and self-contained proofs are given. Our arguments are often at least as simple as the ones proposed under stronger hypotheses in classical references. We are in particular careful about separating statements where a finiteness hypothesis is needed (notably in terms of finite generation, finite presentation, or Noetherianity) from others that do not require it. We show that several basic results on regularity hold without any finiteness hypothesis, and that many results on the asymptotic behaviour hold for modules of finite regularity.

In Section 6, we give fairly general duality statements that encapsulate the Herzog-Rahimi spectral sequence we use in the last section to derive tameness from our previous stability results.

**1. Local cohomology and depth.** Let  $A$  be a commutative ring (with unit) and  $M$  an  $A$ -module. If  $a = (a_1, \dots, a_r)$  is an  $r$ -tuple of elements of  $A$ ,  $K^\bullet(a; M)$  is the Koszul complex and  $H^i(a; M)$  its  $i$ th cohomology module. Also,  $\mathcal{C}_a^\bullet(M)$  is the Čech complex. This complex is isomorphic to  $\varinjlim_n K^\bullet(a_1^n, \dots, a_r^n; M)$ . If  $a$  and  $b$  generates two ideals with same radical, then  $H^i(\mathcal{C}_a^\bullet(M)) \simeq H^i(\mathcal{C}_b^\bullet(M))$  for all  $i$ . Moreover this isomorphism is graded (of degree 0) if  $A$ ,  $M$  and the ideals generated by  $a$  and  $b$  are graded. This, for instance, follows from [7, 1.2.3 and 1.4.1]. It can also be proved in an elementary way as follows: first notice that it is sufficient to prove that, if  $y \in \sqrt{(x_1, \dots, x_t)}$ , then  $H^i(\mathcal{C}_{(x_1, \dots, x_t)}^\bullet(M)) \simeq H^i(\mathcal{C}_{(x_1, \dots, x_t, y)}^\bullet(M))$ , second show that  $\mathcal{C}_{(x_1, \dots, x_t)}^\bullet(M_y)$  is acyclic if  $y \in \sqrt{(x_1, \dots, x_t)}$ , and conclude using that  $\mathcal{C}_{(x_1, \dots, x_t, y)}^\bullet(M)$  is the mapping cone of the natural map  $\mathcal{C}_{(x_1, \dots, x_t)}^\bullet(M) \rightarrow \mathcal{C}_{(x_1, \dots, x_t)}^\bullet(M_y)$ .

We will denote by  $H_I^i(M)$  the  $i$ th homology module of  $H^i(\mathcal{C}_a^\bullet(M))$ , if  $a$  generates the ideal  $I$ .

The  $i$ th right derived functor of the left exact functor  $H_I^0$  coincides with the functor  $T^i(-) := \varinjlim_n \text{Ext}_A^i(A/I^n, -)$ . It coincides with  $H_I^i$  if

and only if  $H_I^i(N) = 0$  whenever  $i > 0$  and  $N$  is injective, and this holds if  $A$  is Noetherian or  $I$  is generated by a regular sequence.

If  $X := \text{Spec}(A)$  and  $Y := V(I) \subset X$ , one has an isomorphism

$$H_I^i(M) \simeq H_Y^i(X, \widetilde{M}).$$

Indeed, the Serre affineness theorem and the Cartan-Leray theorem (see, e.g., [9, 14], and [6, 5.9.1]) provide isomorphisms

$$(1) \quad H_Y^i(X, \widetilde{M}) \simeq H^i(\mathcal{C}_{(a_1, \dots, a_r)}^\bullet(M)) \simeq H^i(M \otimes_A^{\mathbf{L}} \mathcal{C}_{(a_1, \dots, a_r)}^\bullet(A)),$$

as  $\mathcal{C}_{(a_1, \dots, a_r)}^\bullet(A)$  is a complex of flat modules. These isomorphisms show that the functor  $M \mapsto H_I^i(M)$  commutes with direct sums and filtered inductive limits and provide a spectral sequence

$$(2) \quad E_2^{p,q} = \text{Tor}_{-p}^A(M, H_I^q(A)) \implies H_I^{p+q}(M).$$

Also notice that the isomorphism  $H_I^i(M) \simeq \lim_{\substack{\longrightarrow \\ n}} H^i(a_1^n, \dots, a_r^n; M)$  shows that any element of  $H_I^i(M)$  is annihilated by a power of the ideal  $I$ .

**Definition 1.1.** If  $I$  is a finitely generated  $A$ -ideal and  $M$  an  $A$ -module, we set

$$\text{depth}_I(M) := \max\{p \in \mathbf{N} \cup \{+\infty\} \mid H_I^i(M) = 0, \text{ for all } i < p\},$$

and

$$\text{cd}_I(M) := \max\{p \in \mathbf{N} \cup \{-\infty\} \mid H_I^p(M) \neq 0\}.$$

In case there might be an ambiguity on the ring over which  $I$  and  $M$  are considered, we will use the notations  $\text{depth}_I^A(M)$  and  $\text{cd}_I^A(M)$ .

Notice that, for any  $A$ -module  $M$ ,  $\text{cd}_I(M)$  is bounded above by the minimal number of generators of any ideal  $J$  such that  $\sqrt{J} = \sqrt{I}$  (this number is called the arithmetic rank of  $I$  in  $A$ ,  $\text{ara}_A(I)$ ).

**Lemma 1.2.** *If  $I$  is generated by  $a = (a_1, \dots, a_r)$ ,*

$$\text{depth}_I(M) = \max\{p \in \mathbf{N} \cup \infty \mid H^i(a; M) = 0, \text{ for all } i < p\}.$$

*Proof.* Let  $d := \max\{p \mid H^i(a; M) = 0, \text{ for all } i < p\}$ . Recall that, for positive integers  $l_i$ ,  $H^i(a_1^{l_1}, \dots, a_r^{l_r}; M) = 0$  if and only if  $H^i(a; M) = 0$ . It follows that  $H^i_I(M) = \varinjlim_n H^i(a_1^n, \dots, a_r^n; M) = 0$  if  $H^i(a; M) = 0$ . Notice that  $d = \infty$  if and only if  $d > r$ , in which case  $H^i(a; M) = H^i_I(M) = 0$  for all  $i$ . Hence  $\text{depth}_I(M) \geq d$ . We now assume  $d < \infty$ . As  $I$  annihilates  $H^i(a; M)$  for any  $i$ , the totalization of the complex  $\mathcal{C}_I^\bullet K^\bullet(a; M)$  has cohomology isomorphic to the one of  $K^\bullet(a; M)$ . It provides a spectral sequence

$$'E_1^{p,q} = H_I^q K^p(a; M) \implies H^{p+q}(a; M).$$

As  $H_I^q K^p(a; M) = 0$  for  $q < d$ , this in turn provides a natural into map  $H^d(a; M) \rightarrow H_I^d(M)$  which shows that  $\text{depth}_I(M) \leq d$ .  $\square$

**Corollary 1.3.** *If  $I$  is a finitely generated  $A$ -ideal, then for any  $A$ -module  $M$ ,*

$$\text{depth}_I(M) = \min_{\mathfrak{p} \in V(I)} \{\text{depth}_{I_{\mathfrak{p}}}(M_{\mathfrak{p}})\}.$$

To show that this notion agrees with the one introduced by Northcott, we first prove a lemma.

**Lemma 1.4.** *Let  $N$  be an  $A$ -module and  $a \in I$  a non zero divisor on  $N$ . Then*

$$\text{depth}_I(N/aN) = \text{depth}_I(N) - 1.$$

*Proof.* Consider the exact sequence

$$0 \longrightarrow N \xrightarrow{\times a} N \longrightarrow N/aN \longrightarrow 0$$

and the induced long exact sequence on cohomology with support in  $I$ ,

$$\dots \rightarrow H_I^{i-1}(N) \rightarrow H_I^{i-1}(N/aN) \rightarrow H_I^i(N) \xrightarrow{\times a} H_I^i(N) \rightarrow \dots,$$

and let  $r := \text{depth}_I(N)$ . The above sequence shows that  $\text{depth}_I(N/aN) \geq r - 1$ . Furthermore, if  $r < \infty$ ,  $H_I^{r-1}(N/aN) = 0$  if and only if the multiplication by  $a$  is injective on  $H_I^r(N)$ . But this does not hold since any element of  $H_I^r(N)$  is annihilated by a power of  $a$  and  $H_I^r(N) \neq 0$  by definition.  $\square$

We will also use a version of the Dedekind-Mertens lemma, that we now recall in its general form, together with immediate corollaries that are useful in this text.

**Theorem 1.5** (Generalized Dedekind-Mertens lemma) [8, 3.2.1]. *Let  $A$  be a ring,  $M$  be a  $A$ -module and  $\underline{T}$  a set of variables. For  $P \in A[\underline{T}]$  and  $Q \in M[\underline{T}]$ , let  $c(P)$  be the  $A$ -ideal generated by the coefficients of  $P$ ,  $c(Q)$  the submodule of  $M$  generated by the coefficients of  $Q$  and  $\ell(Q)$  the number of non zero coefficients of  $Q$ .*

*Then one has the equality*

$$c(P)^{\ell(Q)-1}c(PQ) = c(P)^{\ell(Q)}c(Q).$$

*In particular, the kernel of the multiplication by  $P$  in  $M[\underline{T}]$  is supported in  $V(c(P))$ .*

**Corollary 1.6.** *Let  $A$  be a ring,  $I = (a_0, \dots, a_p)$  an  $A$ -ideal and  $M$  an  $A$ -module. Set  $\xi := a_0 + a_1T + \dots + a_pT^p \in A[T]$ . Then*

$$\ker(M[T] \xrightarrow{\times \xi} M[T]) \subset H_I^0(M[T]) = H_I^0(M)[T].$$

Let  $S = R[X_1, \dots, X_n]$  be a polynomial ring over a commutative ring  $R$ , and set  $S_+ := (X_1, \dots, X_n)$ .

**Corollary 1.7.** *Let  $M$  be a graded  $S$ -module. Set  $\ell := T_1X_1 + \dots + T_nX_n$  with  $\deg T_i = 0$ . Then the kernel of the map,*

$$M[T_1, \dots, T_n] \xrightarrow{\times \ell} M[T_1, \dots, T_n](1)$$

*is a graded  $S[T_1, \dots, T_n]$ -submodule of  $H_{S_+}^0(M)[T_1, \dots, T_n]$ .*

**Corollary 1.8.** *Consider indeterminates  $(U_{i,j})_{1 \leq i, j \leq n}$ ,  $\xi_i := \sum_{1 \leq j \leq n} U_{i,j} X_j$ ,  $\Delta := \det (U_{i,j})_{1 \leq i, j \leq n}$  and  $R' := R[(U_{i,j})_{1 \leq i, j \leq n}]_{\Delta}$ . Let  $S' := R'[X_1, \dots, X_n]$ , and set  $M' := M \otimes_R R'$  for any  $S$ -module  $M$ . Then  $(\xi_1, \dots, \xi_n)$  is  $M'$ -regular iff  $V(S_+') = V(\xi_1, \dots, \xi_n)$ .*

The following proposition shows that the above definition of depth agrees with the one introduced by Northcott in [10].

**Proposition 1.9.** *Let  $r \geq 1$  be an integer and  $I$  a finitely generated  $A$ -ideal. The following are equivalent:*

- (1)  $\text{depth}_I(M) \geq r$ ,
- (2) *There exists a faithfully flat extension  $B$  of  $A$  and a regular sequence  $f_1, \dots, f_r$  on  $B \otimes_A M$  contained in  $IB$ ,*
- (3) *There exists a polynomial extension  $B$  of  $A$  and a regular sequence  $f_1, \dots, f_r$  on  $B \otimes_A M$  contained in  $IB$ ,*
- (4) *There exists a regular sequence  $f_1, \dots, f_r$  on  $M[T_1, \dots, T_r]$ , where the  $T_i$ 's are variables, contained in  $IA[T_1, \dots, T_r]$ .*

*Proof.* The implications (4)  $\Rightarrow$  (3)  $\Rightarrow$  (2) are trivial. Furthermore, (2) implies that  $H_{IB}^i(B \otimes_A M) = 0$  for  $i < r$  using Lemma 1.4, which in turn implies (1) since  $H_{IB}^i(B \otimes_A M) \simeq B \otimes_A H_I^i(M)$  because  $B$  is flat over  $A$ .

Finally, (1) implies (4) by induction on  $r$ , using Lemma 1.4 and Corollary 1.6.  $\square$

*Remark 1.10.* Let  $r \geq 1$  be an integer and  $I$  a finitely generated  $A$ -ideal. If  $\text{depth}_I(M) \geq r$  and  $f_1, \dots, f_s \in IB$  is a regular sequence on  $B \otimes_A M$  for some flat extension  $B$  of  $A$ , then  $s \leq r$  and there exists a faithfully flat extension  $C$  of  $B$  and  $f_{s+1}, \dots, f_r \in IC$  such that  $f_1, \dots, f_r$  is regular on  $C \otimes_A M$ .

**2. Castelnuovo-Mumford regularity.** Let  $S$  be a finitely generated standard graded algebra over a commutative ring  $R$ . Recall that, for a graded  $S$ -module  $M$ ,

$$\text{reg}(M) := \sup\{\mu \mid \exists i, H_{S_+}^i(M)_{\mu-i} \neq 0\},$$

with the convention that  $\sup \emptyset = -\infty$ .

**Lemma 2.1.** *Let  $M$  be a graded  $S$ -module. Consider the following properties:*

- (i)  $M_\mu = 0$  for  $\mu \gg 0$ ,
- (ii)  $M = H_{S_+}^0(M)$ ,
- (iii)  $H_{S_+}^i(M) = 0$  for  $i > 0$ .

*Then (i)  $\Rightarrow$  (ii)  $\Rightarrow$  (iii), (ii)  $\Rightarrow$  (i) if  $M$  is finitely generated or  $\text{reg}(M) < \infty$ , and (iii)  $\Rightarrow$  (ii) if  $M_\mu = 0$  for  $\mu \ll 0$ .*

*Proof.* (i)  $\Rightarrow$  (ii) is clear since any homogeneous element in  $M$  is killed by a power of any element of  $S_+$  if (i) holds. If (ii) holds, then  $M_x = 0$  for any  $x \in S_+$ ; hence, the Čech complex on generators of  $S_1$  (as an  $R$ -module) is concentrated in homological degree 0, which shows (iii).

If (ii) holds, any element in  $M$  is killed by a power of  $S_+$ ; hence, if  $M$  is finitely generated by  $(m_t)_{t \in T}$ , any generator  $m_t$  is killed by  $S_+^{N_t}$ , for some  $N_t \in \mathbf{N}$ . It then follows that any element in  $M$  of degree bigger than  $\max_{t \in T} \{\deg(m_t) + N_t\}$  is 0. If  $\text{reg}(M) < \infty$ , (ii)  $\Rightarrow$  (i) follows trivially from the definition of  $\text{reg}(M)$ .

If (iii) holds, set  $N := M/H_{S_+}^0(M)$ . The exact sequence  $0 \rightarrow H_{S_+}^0(M) \rightarrow M \rightarrow N \rightarrow 0$  gives rise to a long exact sequence in local cohomology showing that  $H_{S_+}^i(N) = 0$  for all  $i$ . As  $\text{depth}_{S_+}(N) = +\infty$ , Lemma 1.2 shows that  $N = \hat{S}_+ N$ . This implies that  $N = 0$  as  $N_\mu = 0$  for  $\mu \ll 0$ .  $\square$

The following two propositions extend classical results on regularity.

**Proposition 2.2.** *Let  $\ell \geq 1$  and  $m$  be integers, and let  $M$  be a graded  $S$ -module.*

*If  $H_{S_+}^i(M)_{m-i} = 0$  for  $i \geq \ell$ , then  $H_{S_+}^i(M)_{\mu-i} = 0$  for  $\mu \geq m$  and  $i \geq \ell$ .*

*Assume that  $H_{S_+}^i(M)_{m-i} = 0$  for all  $i$ , and let  $\mu \geq m$ . Then  $H_{S_+}^i(M)_{\mu-i} = 0$  for  $i > 0$  and  $H_{S_+}^0(M)_\mu = M_\mu/S_1 M_{\mu-1}$ .*

*Proof.* We may assume that  $S = R[X_1, \dots, X_n]$  is a polynomial ring over  $R$ . We then prove the assertion by induction on  $n$ .

When  $n = 0$ ,  $M = H_{S_+}^0(M)$  and  $H_{S_+}^i(M) = 0$  for  $i \neq 0$ , and the claim follows in both cases.

Next assume that  $n \geq 1$ , and the assertion is true for  $n - 1$  over any commutative ring. Let

$$\xi := X_1 T^{n-1} + \cdots + X_{n-1} T + X_n \in R[T, X_1, \dots, X_n].$$

The Dedekind-Mertens Lemma implies that

$$\ker(\xi : M[T](-1) \longrightarrow M[T]) \subset H_{S_+}^0(M[T]),$$

by Corollary 1.6. But  $H_{S_+}^i(N[T]) = H_{S_+}^i(N)[T]$  for any  $S$ -module  $N$  and any  $i$ . Hence, replacing  $R$  by  $R[T]$ ,  $M$  by  $M[T]$  and  $X_n$  by  $\xi$ , we may assume that

$$K := \ker(X_n : M(-1) \longrightarrow M) \subset H_{S_+}^0(M).$$

We then have  $H_{S_+}^i(K) = 0$  for  $i \neq 0$  and the exact sequence

$$0 \longrightarrow K \longrightarrow M(-1) \xrightarrow{\times X_n} M \longrightarrow Q \longrightarrow 0$$

induces for all  $i$  an exact sequence

$$H_{S_+}^i(M)(-1) \xrightarrow{\times X_n} H_{S_+}^i(M) \longrightarrow H_{S_+}^i(Q) \longrightarrow H_{S_+}^{i+1}(M)(-1).$$

For  $i + j = m$ , the equalities

$$H_{S_+}^i(M)_j = 0 \quad \text{and} \quad H_{S_+}^{i+1}(M)(-1)_j = H_{S_+}^{i+1}(M)_{j-1} = 0$$

imply  $H_{S_+}^i(Q)_j = 0$ . Hence,  $H_{S_+}^i(Q)_j = 0$  for  $i \geq \ell$  and  $i + j = m$ . As  $Q$  is annihilated by  $X_n$ , setting  $\mathfrak{n} := (X_1, \dots, X_{n-1})$ , one has  $H_{S_+}^i(Q) = H_{\mathfrak{n}}^i(Q)$  for all  $i$ . Applying the recursion hypothesis to the  $R[X_1, \dots, X_{n-1}]$ -module  $Q$ , it follows that

$$H_{S_+}^i(Q)_j = H_{\mathfrak{n}}^i(Q)_j = 0, \quad \text{for all } i \geq \ell, \text{ for all } j \geq m - i.$$

Hence,  $X_n : H_{S_+}^i(M)_{j-1} \rightarrow H_{S_+}^i(M)_j$  is onto for  $i \geq \ell$  and  $i + j \geq m$ . As  $H_{S_+}^i(M)_{m-i} = 0$  for  $i \geq \ell$ , this proves our claim.  $\square$



*Remark 2.3.* Notice that the exact sequence  $0 \rightarrow S_+M \rightarrow M \rightarrow M/S_+M \rightarrow 0$  induces an exact sequence

$$\begin{aligned} 0 \longrightarrow H_{S_+}^0(S_+M) \longrightarrow H_{S_+}^0(M) \longrightarrow M/S_+M \\ \longrightarrow H_{S_+}^1(S_+M) \longrightarrow H_{S_+}^1(M) \longrightarrow 0. \end{aligned}$$

**Proposition 2.4.** *If  $S = R[X_1, \dots, X_n]$  is a polynomial ring, then for any graded  $S$ -module  $M$ :*

- (i)  $\text{reg}(M) = \sup\{\mu \mid \exists i, \text{Tor}_i^S(M, R)_{\mu+i} \neq 0\}$ ,
- (ii)  $\text{reg}(M) = \sup\{\mu \mid \exists i, H_i(X_1, \dots, X_n; M)_{\mu+i} \neq 0\}$ ,
- (iii)  $\text{reg}(M) = \sup\{\mu \mid \exists j, H^j(X_1, \dots, X_n; M)_{\mu-j} \neq 0\}$ ,
- (iv)  $\text{reg}(M) = \sup\{\mu \mid \exists j, \text{Ext}_S^j(R, M)_{\mu-j} \neq 0\}$ .

In particular,  $M$  is generated in degrees at most  $\text{reg}(M)$  (when  $\text{reg}(M) = -\infty$ , it means that  $M$  is generated in degrees at most  $\mu$ , for any  $\mu \in \mathbf{Z}$ ).

*Proof.* We first show this equality if  $\text{reg}(M) < \infty$ . Let  $K_\bullet(M) := K_\bullet(X_1, \dots, X_n; M)$  and  $K^\bullet(M) := K^\bullet(X_1, \dots, X_n; M)$ . As  $K_\bullet(M) = K_\bullet(S) \otimes_S M$ ,  $K_\bullet(M) = \text{Hom}_S(K_\bullet(S), M)$  and  $K_\bullet(S)$  is a free  $S$ -resolution of  $R$ , it follows that  $H_i(K_\bullet(M)) \simeq \text{Tor}_i^S(M, R)$  and  $H^j(K^\bullet(M)) \simeq \text{Ext}_S^j(R, M)$ . Furthermore, the complexes  $K^\bullet(M)$  and  $K_\bullet(M)$  are isomorphic, up to a shift in homological degree and internal degree:  $K^\bullet(M) \simeq K_{n-\bullet}(M)(n)$ , proving that

$$\begin{aligned} \text{Ext}_S^j(R, M)_{\mu-j} &\simeq H^j(K^\bullet(M))_{\mu-j} \\ &\simeq H_{n-j}(K_\bullet(M))_{\mu-j+n} \simeq \text{Tor}_{n-j}^S(M, R)_{\mu-j+n} \end{aligned}$$

and the equivalence of the four items.

To prove (ii), the double complex  $C_{(X_1, \dots, X_n)}^\bullet K_\bullet(M)$  gives rise to two spectral sequences whose first terms are, respectively,

$$\begin{aligned} {}'E_1^{p,q} &= K_{-p}(X_1, \dots, X_n; H_{S_+}^q(M)), \\ {}'E_2^{p,q} &= \text{Tor}_{-p}^S(H_{S_+}^q(M), R) \end{aligned}$$

and

$$\begin{aligned} {}''E_1^{p,q} &= \mathcal{C}_{(X_1, \dots, X_n)}^p \operatorname{Tor}_{-q}^S(M, R), \\ {}''E_2^{p,q} &= H_{S_+}^p(\operatorname{Tor}_{-q}^S(M, R)). \end{aligned}$$

Recall that  $K_\bullet(M)_{X_i}$  is acyclic for any  $i$ , hence  ${}''E_1^{p,q} = 0$  for  $p \neq 0$ , which implies that  ${}''E_\infty^{p,q} = 0$  for  $p \neq 0$  and  ${}''E_\infty^{0,q} \simeq \operatorname{Tor}_{-q}^S(M, R)$ .

On the other hand,  $({}'E_1^{p,q})_\mu = 0$  for  $\mu > \operatorname{reg}(M) - q - p$  as  $H_{S_+}^q(M)$  lives in degrees at most  $\operatorname{reg}(M) - q$ . It follows first that  $\operatorname{Tor}_{-j}^S(M, R)$  lives in degrees at most  $\operatorname{reg}(M) - j$  showing that  $\operatorname{Tor}_i^S(M, R)_{\mu+i} = 0$  for  $\mu > \operatorname{reg}(M)$ .

To conclude, choose  $j$  such that  $H_{S_+}^j(M)_{\operatorname{reg}(M)-j} \neq 0$ . Set  $\mu := \operatorname{reg}(M) - j + n$  and notice that  $({}'E_1^{p,q})_\mu = 0$  when  $p + q = -n + j + 1$ . As  ${}'E_1^{p,q} = 0$  for  $p < -n$ , it follows that  $0 \neq H_{S_+}^j(M)_{\operatorname{reg}(M)-j} = ({}'E_1^{-n,j})_\mu \simeq ({}'E_\infty^{-n,j})_\mu$ , which shows that  ${}''E_\infty^{0,j-n} \simeq \operatorname{Tor}_{n-j}^S(M, R)_{\operatorname{reg}(M)+n-j} \neq 0$ .

To finish the proof, we must show that  $\operatorname{reg}(M) < \infty$  if there exists  $\mu_0$  such that  $\operatorname{Tor}_i^S(M, R)_\mu = 0$  for all  $i$  and  $\mu \geq \mu_0$ .

We first show that, in this case, there exists a graded free  $S$ -resolution  $F_\bullet$  of  $M$  with  $F_i = \bigoplus_{j \in \mathcal{I}_i} S(-d_{ij})$ , and  $d_{ij} < \mu_0$  for all  $i$  and  $j$ . Notice that, if  $M$  is graded and  $(M/S_+M)_{>\nu} = 0$ , then  $M$  is generated in degree at most  $\nu$ , showing the existence of a graded epimorphism  $\phi : F_0 \rightarrow M$  with  $F_0$  as claimed. The exact sequence,

$$0 \longrightarrow \ker(\phi) \longrightarrow F_0 \longrightarrow M \longrightarrow 0,$$

gives rise to another

$$\operatorname{Tor}_1^S(M, R) \longrightarrow \ker(\phi)/S_+\ker(\phi) \longrightarrow F_0/S_+F_0,$$

and proves the existence of  $F_1$ , as claimed, such that  $F_1 \rightarrow \ker(\phi)$  is a graded epimorphism. As  $\operatorname{Tor}_j^S(M, R) \simeq \operatorname{Tor}_{j-1}^S(\ker(\phi), R)$ , for  $j \geq 2$ , the conclusion follows by induction on  $i$ .

Finally, it suffices to remark that, if  $F_\bullet$  is a graded resolution as above, then  $H_{S_+}^i(M)_\mu \simeq H_{n-i}(H_{S_+}^n(F_\bullet)_\mu)$  vanishes in degree bigger than  $-n + \max_{i,j} \{d_{ij}\} \leq -n + \mu_0$ , as  $H_{S_+}^n(F_\bullet)_\mu = 0$  for all  $i$  for such a  $\mu$ . Hence,  $\operatorname{reg}(M) \leq \mu_0$ .  $\square$

**Lemma 2.5.** *For any graded  $S$ -module  $N$ ,*

$$\operatorname{reg}(N) = \sup_{\mathfrak{p} \in \operatorname{Spec}(R)} \{\operatorname{reg}(N \otimes_R R_{\mathfrak{p}})\}.$$

*Furthermore,  $\operatorname{reg}(N) = \operatorname{reg}(N \otimes_R R_{\mathfrak{p}})$  for some  $\mathfrak{p} \in \operatorname{Spec}(R)$  if  $\operatorname{reg}(N) < \infty$ .*

*Proof.* For  $\mathfrak{p} \in \operatorname{Spec}(R)$ ,

$$H_{S_+}^i(N \otimes_R R_{\mathfrak{p}})_{\mu} \simeq (H_{S_+}^i(N)_{\mu}) \otimes_R R_{\mathfrak{p}},$$

from which the claim directly follows.  $\square$

### 3. Depth of the graded components of a graded module.

As in the previous section,  $S$  is a finitely generated standard graded algebra over a commutative ring  $R$ .

**Proposition 3.1.** *Let  $I$  be a finitely generated  $R$ -ideal,  $M$  a graded  $S$ -module and  $d$  an integer. Assume that  $\operatorname{depth}_I(M_{\mu}) \geq d$  for  $\mu \gg 0$ . Then:*

- (i)  $\operatorname{depth}_I(M_{\mu}) \geq d$  for any  $\mu$  such that  $H_{S_+}^q(M)_{\mu} = 0$  for  $q < d$ ,
- (ii) if  $\operatorname{depth}_I(M_{\mu}) = d$  for some  $\mu$  such that  $H_{S_+}^q(M)_{\mu} = 0$  for  $q \leq d$ , then  $\operatorname{depth}_I(M_{\nu}) \leq d$  for any  $\nu \geq \mu$ .

*Proof.* Let  $a = (a_1, \dots, a_r)$  be generators of  $I$ . Recall that  $H^p(a; M)_{\mu} = H^p(a; M_{\mu})$ , as  $I$  is an  $R$ -ideal. Hence, if  $\operatorname{depth}_I(M_{\mu}) \geq d$  for  $\mu \gg 0$ , the  $S$ -modules  $H^p(a; M)$  are supported in  $V(S_+)$  for  $p < d$ , by Lemma 1.2 and (i)  $\Rightarrow$  (ii) in Lemma 2.1. Comparing the two spectral sequences obtained from the double complex  $\mathcal{C}_{S_+}^{\bullet} K^{\bullet}(a; M)$ , computing the hypercohomology  $\mathbf{H}^{\bullet}$  of  $K^{\bullet}$ , we obtain on the one hand that  $\mathbf{H}^i \simeq H^i(a; M)$  for  $i < d$  and  $\mathbf{H}^d \simeq H_{S_+}^0(H^d(a; M))$ . On the other hand, one has a spectral sequence

$${}'E_1^{p,q} = H_{S_+}^q K^p \implies \mathbf{H}^{p+q},$$

which shows that  $(\mathbf{H}^i)_{\mu} = 0$  for  $i < d$  if  $H_{S_+}^q(M)_{\mu} = 0$  for  $q < d$ , proving (i).

For (ii), the condition  $H_{S_+}^q(M)_\mu = 0$  for  $q \leq d$  implies that  $H_{S_+}^0(H^d(a; M))_\mu = H_{S_+}^0(H^d(a; M_\mu)) = 0$ . Hence, if  $H^d(a; M_\mu) \neq 0$ , then  $0 \neq (S_+)^{\nu-\mu}H^d(a; M)_\mu \subseteq H^d(a; M)_\nu$ , which shows (ii) by Lemma 1.2.  $\square$

**Corollary 3.2.** *Let  $I$  be an  $R$ -ideal and  $M$  a graded  $S$ -module. Then  $\text{depth}_I(M_{\geq \mu})$  is independent of  $\mu$  for  $\mu > \text{reg}(M) - \text{depth}_{S_+}(M)$ .*

**Theorem 3.3.** *Let  $I$  be an  $R$ -ideal and  $M$  a graded  $S$ -module with  $\text{reg}(M) < \infty$ . Set  $r := \text{reg}(M) - \text{depth}_{S_+}(M)$ ,  $d := \min_{\nu > r} \{\text{depth}_I(M_\nu)\} = \text{depth}_I(M_{> r})$  and  $\mu_0 := \inf\{\nu > r \mid \text{depth}_I(M_\nu) = d\}$ .*

*Then  $\text{depth}_I(M_\mu) = d$  for all  $\mu \geq \mu_0$ .*

*Proof.* We may assume that  $d < \infty$ . By definition of  $d$ ,  $\text{depth}_I(M_\mu) \geq d$  for  $\mu \geq \mu_0$ , as  $\mu_0 > r$ . On the other hand,  $\text{depth}_I(M_\mu) \leq d$  for  $\mu \geq \nu$  by Proposition 3.1 (ii). The conclusion follows.  $\square$

**4. Cohomological dimension.** Let  $A$  be a commutative ring (with unit),  $I$  a finitely generated ideal and  $M$  an  $A$ -module. First remark that it follows from (1) that, if  $M$  is annihilated by an ideal  $J$ , for instance, if  $J = \text{ann}_A(M)$ , considering  $M$  as an  $A/J$ -module one has

$$(3) \quad \text{cd}_I^A(M) = \text{cd}_{(I+J)/J}^{A/J}(M).$$

Furthermore,

**Proposition 4.1.** *Let  $M$  be an  $A$ -module.*

(a)

$$\text{cd}_I(M) \leq \max_{\substack{ECM \\ E f.g.}} \{\text{cd}_I(E)\}.$$

(b)

$$\text{cd}_I(M) \leq \text{cd}_I(A/\text{ann}_A M) \leq \text{cd}_I(A).$$

(c) *If  $M$  is finitely generated, then*

$$\text{cd}_I(M) = \text{cd}_I(A/\text{ann}_A M).$$

*Proof.* (a)  $M$  is the filtered inductive limit of its submodules of finite type (for the inclusion), and local cohomology commutes with filtered inductive limits.

(b) The spectral sequence (2) shows that  $\text{cd}_I(N) \leq \text{cd}_I(A)$  for any module  $N$ , in particular  $\text{cd}_I(A/\text{ann}_A(M)) \leq \text{cd}_I(A)$ . Together with (3) applied with  $J := \text{ann}_A(M)$ , we get  $\text{cd}_I^A(M) = \text{cd}_{(I+J)/J}^{A/J}(M) \leq \text{cd}_{(I+J)/J}^{A/J}(A/J) = \text{cd}_I(A/J)$ .

(c) According to (b), it suffices to show that  $\text{cd}_I(A/\text{ann}_A M) \leq \text{cd}_I(M)$ . Replacing  $A$  by  $A/\text{ann}_A(M)$ , we may assume that  $M$  is faithful.

We will show that  $\text{cd}_I(M) \leq r$  implies  $\text{cd}_I(A) \leq r$ . This is clear for  $r \geq \text{ara}_A(I)$ , and we now perform a descending recursion on  $r$ . Assume that this is true for  $r + 1$ . If  $\text{cd}_I(M) \leq r$ , by recursion hypothesis we know that  $\text{cd}_I(A) \leq r + 1$ ; hence, spectral sequence (2) implies that

$$0 = H_I^{r+1}(M) = M \otimes_A H_I^{r+1}(A).$$

As  $M$  is faithful and of finite type, [13, 4.3] shows that  $H_I^{r+1}(A) = 0$ , which implies that  $\text{cd}_I(A) \leq r$ .  $\square$

The following corollary has been proved by Dibaei and Vahidi in the Noetherian case in [4, 2.2].

**Corollary 4.2.** *Let  $M$  be an  $A$ -module and  $I, J$  two finitely generated ideals. Then*

$$\text{cd}_{I+J}(M) \leq \text{cd}_I(A/\text{ann}_A M) + \text{cd}_J(M),$$

and  $\text{cd}_{I+J}(M) \leq \text{cd}_I(M) + \text{cd}_J(M)$  if  $M$  is finitely generated.

*Proof.* We may assume that  $M$  is faithful. If  $a$  is a finite set of generators of  $I$  and  $b$  a finite set of generators of  $J$ , the double complex with components  $\mathcal{C}_a^i(A) \otimes_A \mathcal{C}_b^j(M)$  gives rise to a spectral sequence with second term  $H_I^i(H_J^j(M))$  that abuts to  $H_{I+J}^{i+j}(M)$ . Hence,

$$\text{cd}_{I+J}(M) \leq \max\{i + j \mid H_I^i(H_J^j(M)) \neq 0\} \leq \text{cd}_I(A) + \text{cd}_J(M),$$

by Proposition 4.1 (b).

Furthermore, by Proposition 4.1 (c),  $\text{cd}_I(A) = \text{cd}_I(M)$  whenever  $M$  is faithful and finitely generated.  $\square$

**Corollary 4.3.** *Let  $M$  be a finitely generated  $A$ -module and  $\mathfrak{a}$  an ideal of  $A$  such that  $\mathfrak{a}^t M = 0$  for some  $t$ . Then*

$$\text{cd}_I(M) = \text{cd}_I(M/\mathfrak{a}M).$$

*Proof.* Let  $d := \text{cd}_I(M) = \text{cd}_I(A/\text{ann}_A(M))$  (by Proposition 4.1 (c)). As  $M/\mathfrak{a}M$  is annihilated by  $\text{ann}_A(M)$ , it follows from Proposition 4.1 (b) that  $\text{cd}_I(M/\mathfrak{a}M) \leq d$ . Furthermore, Proposition 4.1 (b) shows that the functor  $N \mapsto H_I^d(N)$  restricted to the category of  $A$ -modules annihilated by  $\text{ann}_A(M)$  is right exact. It implies that  $H_I^d(M/\mathfrak{a}M) = H_I^d(M) \otimes_A A/\mathfrak{a} = H_I^d(M)/\mathfrak{a}H_I^d(M)$ .

If  $\text{cd}_I(M/\mathfrak{a}M) < d$  it implies

$$H_I^d(M) = \mathfrak{a}H_I^d(M) = \mathfrak{a}^2H_I^d(M) = \cdots = \mathfrak{a}^tH_I^d(M) = 0,$$

which contradicts the definition of  $d$ .  $\square$

**Proposition 4.4.** *Let  $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$  be an exact sequence of  $A$ -modules. Then*

$$\text{cd}_I(M) \leq \max\{\text{cd}_I(M'), \text{cd}_I(M'')\} \leq \text{cd}_I(A/\text{ann}_A(M)).$$

*Furthermore, all inequalities are equalities if  $M$  is finitely generated.*

*Proof.* First, the exact sequences

$$H_I^i(M') \longrightarrow H_I^i(M) \longrightarrow H_I^i(M''), \quad i \in \mathbf{Z},$$

show the inequality on the left. As both  $A$ -modules  $M'$  and  $M''$  are annihilated by  $\text{ann}_A(M)$ , the inequality on the right follows from (3) and Proposition 4.1 (b).

Finally, the extreme terms are equal according to Proposition 4.1 (c) if  $M$  is finitely generated.  $\square$

*Remark 4.5.* If  $A$  is a domain, distinct from its field of fractions  $K$  and  $I$  a proper finitely generated ideal, then  $A$  is a submodule of  $K$  such that

$$-\infty = \text{cd}_I(K) < 0 \leq \text{cd}_I(A).$$

**Corollary 4.6.** *If  $M$  is a Noetherian  $A$ -module, then*

$$\begin{aligned} \text{cd}_I(M) &= \max_{\mathfrak{p} \in \text{Supp}_A(M)} \text{cd}_I(A/\mathfrak{p}) \\ &= \max_{\mathfrak{p} \in \text{Ass}_A(M)} \text{cd}_I(A/\mathfrak{p}) \\ &= \max_{\mathfrak{p} \in \text{Min}_A(M)} \text{cd}_I(A/\mathfrak{p}). \end{aligned}$$

*Proof.* Let  $\text{Min}_A(M)$  be the minimal primes in the support of  $M$ . Every  $\mathfrak{p} \in \text{Supp}_A(M)$  contains some  $\mathfrak{q} \in \text{Min}_A(M)$ , and the canonical epimorphism  $A/\mathfrak{q} \rightarrow A/\mathfrak{p}$  gives an inequality  $\text{cd}_I(A/\mathfrak{p}) \leq \text{cd}_I(A/\mathfrak{q})$  by Proposition 4.4. It follows that

$$\max_{\mathfrak{p} \in \text{Supp}_A(M)} \text{cd}_I(A/\mathfrak{p}) = \max_{\mathfrak{p} \in \text{Min}_A(M)} \text{cd}_I(A/\mathfrak{p}).$$

On the other hand,  $\text{Min}_A(M) \subset \text{Ass}_A(M)$ , and for  $\mathfrak{p} \in \text{Ass}_A(M)$ , the existence of a monomorphism  $A/\mathfrak{p} \rightarrow M$  implies by Proposition 4.4 that  $\text{cd}_I(A/\mathfrak{p}) \leq \text{cd}_I(M)$ . Hence,

$$\max_{\mathfrak{p} \in \text{Min}_A(M)} \text{cd}_I(A/\mathfrak{p}) \leq \max_{\mathfrak{p} \in \text{Ass}_A(M)} \text{cd}_I(A/\mathfrak{p}) \leq \text{cd}_I(M),$$

and it remains to show that  $\text{cd}_I(M) \leq \max_{\mathfrak{p} \in \text{Supp}_A(M)} \text{cd}_I(A/\mathfrak{p})$ .

Observe that, if  $M \neq 0$ , it admits a finite filtration by cyclic modules  $A/\mathfrak{p}_i$  ( $1 \leq i \leq t$ ) with  $\mathfrak{p}_i \in \text{Supp}_A(M)$  for all  $i$ . Again, by Proposition 4.4, we obtain

$$\text{cd}_I(M) = \max_{1 \leq i \leq t} \{\text{cd}_I(A/\mathfrak{p}_i)\} \leq \max_{\mathfrak{p} \in \text{Supp}_A(M)} \{\text{cd}_I(A/\mathfrak{p})\}.$$

This concludes the proof.  $\square$

The following result generalizes the main theorem of [5], which applies in the case of two finitely generated modules over a Noetherian ring.

**Proposition 4.7.** *Let  $M$  and  $N$  be  $A$ -modules. Assume  $M$  is finitely presented, and  $\text{Supp}_A(N) \subset \text{Supp}_A(M)$ . Then*

$$\text{cd}_I(N) \leq \text{cd}_I(M).$$

*Proof.* Let  $E$  be a finitely generated submodule of  $N$ . The inclusion  $\text{Supp}_A(N) \subset \text{Supp}_A(M)$  implies that

$$(4) \quad \sqrt{\text{ann}_A(M)} \subset \sqrt{\text{ann}_A(E)}.$$

As  $M$  is finitely generated, the Fitting ideal  $\text{Fitt}_A^0(M)$  has the same radical as  $\text{ann}_A(M)$  and contains a power of  $\text{ann}_A(M)$ . Furthermore, as  $M$  is finitely presented, this ideal is finitely generated. It then follows from (4) that there exists  $t$  such that  $\text{ann}_A(M)^t \subset \text{ann}_A(E)$ . By Corollary 4.3 and Proposition 4.1 (b) and (c), one has

$$\text{cd}_I(E) \leq \text{cd}_I(E/\text{ann}_A(M)E) \leq \text{cd}_I(A/\text{ann}_A(M)) = \text{cd}_I(M).$$

The conclusion follows by Proposition 4.1 (a) applied to the  $A$ -module  $N$ .  $\square$

Now, let  $S$  be a finitely generated standard graded algebra over a commutative ring  $R$ .

**Definition 4.8.** For a graded  $S$ -module  $M$ ,

$$a_{S_+}^i(M) := \sup\{\mu \mid H_{S_+}^i(M)_\mu \neq 0\},$$

so that  $\text{reg}(M) = \max_i\{a_{S_+}^i(M) + i\}$ .

**Proposition 4.9.** *Let  $M$  be a finitely generated graded  $S$ -module and  $I$  a finitely generated  $R$ -ideal. Then*

- (a)  $\text{cd}_I(M_\mu)$  is a non decreasing function of  $\mu$  for  $\mu > a_{S_+}^0(M)$ ,
- (b)  $\text{cd}_I(M_\mu)$  is constant for  $\mu \geq \text{reg}(M) + n - \text{depth}_{S_+}(M)$  if  $n > 0$ .

*Proof.* We may, and will, assume that  $\text{cd}_{S_+}(M) > 0$ , as the proposition is immediate when  $\text{cd}_{S_+}(M) = 0$  by Lemma 2.1. We also remark



that it suffices to prove the claim after the faithfully flat base change  $R \rightarrow R'$ ; hence, we may further assume (making an invertible linear change of coordinates) by Remark 1.8 that the sequence  $(X_1, \dots, X_n)$  is  $M$ -regular off  $V(S_+)$ .

In particular, the kernel of the map  $M \rightarrow M(1)$  induced by multiplication by  $X_n$  is contained in  $H_{S_+}^0(M)$ . It follows an injection  $M_\nu \rightarrow M_{\nu+1}$  for  $\nu > a_{S_+}^0(M)$  which proves (a) by Proposition 4.4 as  $M_{\nu+1}$  is finitely generated over  $R$ .

To prove (b), we consider the two converging spectral sequences arising from the double complex  $\mathcal{C}_I^\bullet K_\bullet(X_1, \dots, X_n; M)$ . They have as respective second terms

$$\mathrm{Tor}_{-p}^S(H_I^q(M), R) \quad \text{and} \quad H_I^p(\mathrm{Tor}_{-q}^S(M, R)).$$

Let  $d := \max_{\nu > a_{S_+}^0(M)} \{\mathrm{cd}_I(M_\nu)\}$ . We may assume  $d \geq 0$ . It follows from the comparison of the spectral sequences that

$$\mathrm{Tor}_0^S(H_I^d(M), R)_{\nu+1} = H_I^d(M_{\nu+1})/S_1 H_I^d(M_\nu) = 0$$

if  $H_I^{d+i}(M_{\nu-i}) = 0$  for  $1 \leq i \leq n-1$  and  $H_I^{d+i}(\mathrm{Tor}_i^S(M, R)_{\nu+1}) = 0$  for all  $i$ .

But  $\mathrm{Tor}_i^S(M, R)_{\nu+1} = 0$  for  $\nu \geq \mathrm{reg}(M) + i$  and  $\mathrm{Tor}_i^S(M, R) = 0$  for  $i > n - \mathrm{depth}_{S_+}(M)$  by Lemma 1.2.

It follows that  $H_I^d(M_\nu) = 0$  implies  $H_I^d(M_{\nu+1}) = 0$  if

$$\begin{aligned} \nu &\geq \max\{a_{S_+}^0(M) + n, \mathrm{reg}(M) + n - \mathrm{depth}_{S_+}(M)\} \\ &= \mathrm{reg}(M) + n - \mathrm{depth}_{S_+}(M). \end{aligned}$$

This implies (b), in view of (a).  $\square$

**5. Associated primes of the graded components of a graded module.** Let  $S$  be a standard graded Noetherian algebra over a commutative ring  $R$ .

**Theorem 5.1.** *Let  $M$  be a graded  $S$ -module. Then*

$$\bigcup_{\mu \in \mathbf{Z}} \mathrm{Ass}_R(M_\mu) = \{\mathfrak{P} \cap R, \mathfrak{P} \in \mathrm{Ass}_S(M)\}.$$

*Proof.* For  $\mu \in \mathbf{Z}$ , let  $\mathfrak{p} \in \text{Ass}_R(M_\mu)$ . There exists an  $x \in M_\mu$  with  $\mathfrak{p} = \text{ann}_R(x)$ . Hence,  $\mathfrak{p}R_\mathfrak{p} = \text{ann}_{R_\mathfrak{p}}(x)$ . Let  $\mathfrak{Q}$  be an  $S \otimes_R R_\mathfrak{p}$ -ideal, maximal among those of the form  $\text{ann}_{S \otimes_R R_\mathfrak{p}}(y)$ ,  $y \in M \otimes_R R_\mathfrak{p}$ , that contains  $\text{ann}_{R_\mathfrak{p}}(x)$ . The ideal  $\mathfrak{Q}$  is associated to  $M \otimes_R R_\mathfrak{p}$ ; hence,  $\mathfrak{P} := \mathfrak{Q} \cap S$  is associated to  $M$  and  $\mathfrak{P} \cap R = \mathfrak{p}$ . One inclusion follows.

Conversely, let  $\mathfrak{P}$  be an ideal associated to  $M$ . We need to show that  $\mathfrak{p} := \mathfrak{P} \cap R$  is associated to  $M_\mu$  for some  $\mu$ . This will be the case if  $\mathfrak{p}R_\mathfrak{p}$  is associated  $(M_\mu)_\mathfrak{p}$ , so that we may assume that  $R$  is local with maximal ideal  $\mathfrak{p}$ . Let  $m \neq 0$  in  $M$  such that  $\mathfrak{P}m = 0$ . If  $m_\nu$  is the degree  $\nu$  component of  $m$ , one has  $\mathfrak{p}m_\nu = 0$ . Hence, choosing  $\mu$  such that  $m_\mu \neq 0$ , one has  $\mathfrak{p} \subseteq \text{ann}_R(m_\mu)$ ; hence,  $\mathfrak{p} = \text{ann}_R(m_\mu)$ , as  $\mathfrak{p}$  is maximal.  $\square$

**Theorem 5.2.** *Let  $M$  be a graded  $S$ -module. If  $H_{S_+}^0(M)_\nu = 0$ , then  $\text{Ass}_R(M_\nu) \subseteq \text{Ass}_R(M_\mu)$  for all  $\mu \geq \nu$ .*

*Proof.* Let  $x_1, \dots, x_n$  be generators of  $S_1$  as an  $R$ -module, and let  $T_1, \dots, T_n$  be variables. Set  $P := \sum_{|\alpha|=\mu-\nu} x^\alpha T^\alpha$ . The polynomial  $P$  is of bidegree  $(\mu - \nu, \mu - \nu)$  for the bigraduation defined by setting  $\deg(x_i) := (1, 0)$  and  $\deg(T_i) := (0, 1)$ , and  $c(P) = (S_+)^{\mu-\nu}$ . Theorem 1.5 shows that kernel  $K$  of the map

$$M[\underline{T}] \xrightarrow{\times P} M[\underline{T}]$$

is a submodule of  $H_{S_+}^0(M)[\underline{T}]$ . Hence,  $K$  vanishes in bidegree  $(\nu, \theta)$  for any  $\theta$ . In particular, it provides an injective map

$$M_\nu = M[\underline{T}]_{\nu,0} \xrightarrow{\times P} M[\underline{T}]_{\mu,\mu-\nu}.$$

As  $M[\underline{T}]_{\mu,\mu-\nu}$  is a finite direct sum of copies of  $M_\mu$ , it follows that any associated prime of  $M_\nu$  is associated to  $M_\mu$ .  $\square$

**Theorem 5.3.** *Let  $M$  be a finitely generated graded  $S$ -module and  $\mathcal{A}$  the finite set  $\cup_{\mu \in \mathbf{Z}} \text{Ass}_R(M_\mu)$  (Theorem 5.1). Set*

$$j(M) := \max_{\mathfrak{p} \in \mathcal{A}} \{a_{S_+}^0(H_\mathfrak{p}^0(M \otimes_R R_\mathfrak{p}))\} \leq a_{S_+}^0(M).$$

*Then, for any ideal  $\mathfrak{p} \in \text{Spec}(R)$ ,  $\ell_{R_\mathfrak{p}}(H_\mathfrak{p}^0(M_\mu \otimes_R R_\mathfrak{p}))$  is a nondecreasing function of  $\mu$ , for  $\mu > j(M)$ .*

*Proof.* First notice that  $H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}}) = 0$  if  $\mathfrak{p} \notin \mathcal{A}$ . Let  $\mathfrak{p} \in \mathcal{A}$ . We will prove that  $\ell_{R_{\mathfrak{p}}}(H_{\mathfrak{p}}^0(M_{\mu} \otimes_R R_{\mathfrak{p}}))$  is a nondecreasing function of  $\mu$ , for  $\mu > a_{S_+}^0(H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}}))$ . The proof of Theorem 5.2, applied to the  $S \otimes_R R_{\mathfrak{p}}$ -module  $H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})$ , with  $P := \sum x_i T_i$  provides an injective morphism of  $R_{\mathfrak{p}}[\underline{T}]$ -modules

$$H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})_{\mu}[\underline{T}] \longrightarrow H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})_{\mu+1}[\underline{T}].$$

Let  $R_{\mathfrak{p}}(\underline{T}) := S^{-1}R_{\mathfrak{p}}[\underline{T}]$  with  $S$  be the multiplicative system of polynomials whose coefficient ideal is the unit ideal. The above injection induces an injective morphism of  $R_{\mathfrak{p}}(\underline{T})$ -modules of finite length

$$H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})_{\mu} \otimes_{R_{\mathfrak{p}}} R_{\mathfrak{p}}(\underline{T}) \longrightarrow H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})_{\mu+1} \otimes_{R_{\mathfrak{p}}} R_{\mathfrak{p}}(\underline{T}).$$

For any  $R_{\mathfrak{p}}$ -module  $N$  of finite length, the  $R_{\mathfrak{p}}(\underline{T})$ -module  $N \otimes_{R_{\mathfrak{p}}} R_{\mathfrak{p}}(\underline{T})$  is a module of the same length as the  $R_{\mathfrak{p}}$ -module  $N$ . The conclusion follows.  $\square$

**Corollary 5.4.** *Let  $M$  be a finitely generated graded  $S$ -module. Then, for any  $\mathfrak{p} \in \text{Spec}(R)$  of height 0,  $\ell_{R_{\mathfrak{p}}}(M_{\mu} \otimes_R R_{\mathfrak{p}})$  is a nondecreasing function of  $\mu$ , for  $\mu > a_{S_+}^0(M)$ . In particular,  $M_{\mu} \otimes_R R_{\mathfrak{p}} = 0$  for all  $\mu > \text{reg}(M)$  or  $M_{\mu} \otimes_R R_{\mathfrak{p}} \neq 0$  for all  $\mu > \text{reg}(M)$ .*

*Proof.* Notice that  $M \otimes_R R_{\mathfrak{p}} = H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})$  as  $\mathfrak{p}$  is of height 0. Also recall that  $M$  (hence,  $M \otimes_R R_{\mathfrak{p}}$ ) is generated in degrees at most  $\text{reg}(M)$ .  $\square$

**Lemma 5.5.** *Let  $M$  be a graded  $S$ -module generated in degree at most  $B$ . Then*

- (i)  $M_{\mu} \neq 0 \Rightarrow M_{\mu+1} \neq 0$ , if  $\mu > a_{S_+}^0(M)$ ,
- (ii)  $M_{\mu} \neq 0 \Leftrightarrow M_{\mu+1} \neq 0$  if  $\mu > \max\{B, a_{S_+}^0(M)\}$ ,
- (iii)  $M_{\mu} \neq 0 \Leftrightarrow M_{\mu+1} \neq 0$  if  $\mu > \text{reg}(M)$ .

*Proof.* (i) follows from Theorem 5.2 as  $R$  is Noetherian. Now (ii) and (iii) follow from (i), as  $M_{\mu} = 0 \Rightarrow M_{\mu+1} = 0$  for  $\mu \geq B$ , and  $M$  is generated in degrees at most  $\text{reg}(M)$ .  $\square$

*Remark 5.6.* The proof of Theorem 5.2 shows that the above lemma holds without assuming that  $R$  is Noetherian.

## 6. Duality results.

**6.1. Preliminaries on  $R\text{Hom}$ .** Let  $(X, \mathcal{O}_X)$  be a ringed space and  $Y$  closed in  $X$ . For any complex  $K^\bullet \in D(X)$ , let  $C(K^\bullet)$  be the Godement resolution of  $K^\bullet$ , and set:

$$\begin{aligned} R\Gamma(X, K^\bullet) &:= C(K^\bullet) \text{ and } R\Gamma_Y(X, K^\bullet) := \Gamma_Y(X, C(K^\bullet)) \text{ in } D(X), \\ R\Gamma(X, K^\bullet) &:= \Gamma(X, C(K^\bullet)) \text{ and } R\Gamma_Y(X, K^\bullet) := \Gamma_Y(X, C(K^\bullet)) \text{ in } D(\Gamma(X, \mathcal{O}_X)). \end{aligned}$$

Notice that a flasque resolution of  $K^\bullet$  in  $D^+(X)$  (e.g., an injective resolution) can be used in place of  $C(K^\bullet)$  if  $K^\bullet \in D^+(X)$ .

We set  $H^i(X, K^\bullet) := H^i(R\Gamma(X, K^\bullet))$ ,  $H_Y^i(X, K^\bullet) := H^i(R\Gamma_Y(X, K^\bullet))$  and  $\mathbf{H}^i(X, K^\bullet) := H^i(R\Gamma(X, K^\bullet))$ , which coincides with the usual notations for  $\mathcal{O}_X$ -modules when considered as complexes concentrated in degree 0.

If there exists a  $d$  such that, for any  $\mathcal{O}_X$ -module  $\mathcal{E}$ ,  $H^i(X, \mathcal{E}) = 0$  (respectively  $H_Y^i(X, \mathcal{E}) = 0$ ) for  $i > d$ , then any flasque resolution of  $K^\bullet$  can be used in place of  $C(K^\bullet)$  to compute  $H^i(X, K^\bullet)$  and  $\mathbf{H}^i(X, K^\bullet)$  (respectively,  $H_Y^i(X, K^\bullet)$ ).

Given  $K^\bullet$  in  $D(X)$  and  $L^\bullet$  in  $D^+(X)$ , one checks that the class in  $D(X)$  (respectively in  $D(\Gamma(X, \mathcal{O}_X))$ ) of the complex  $\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet)$  (respectively,  $\text{Hom}_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet)$ ) is independent of the choice of an injective resolution  $I^\bullet$  of  $L^\bullet$  in  $D^+(X)$ , and one set

$$R\text{Hom}_{\mathcal{O}_X}^\bullet(K^\bullet, L^\bullet) := \text{Hom}_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet)$$

and

$$R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, L^\bullet) := \mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet).$$

When the components of the complex  $K^\bullet$  are locally free  $\mathcal{O}_X$ -modules, there is a quasi-isomorphism

$$R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, L^\bullet) \simeq \mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, L^\bullet).$$

For a pair  $(\mathcal{E}, \mathcal{F})$  of  $\mathcal{O}_X$ -modules, one defines

$$\mathrm{Hom}_Y(\mathcal{E}, \mathcal{F}) := \Gamma_Y(X, \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{E}, \mathcal{F})) = \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{E}, \Gamma_Y \mathcal{F})$$

and

$$\mathcal{H}om_Y(\mathcal{E}, \mathcal{F}) := \Gamma_Y \mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \mathcal{F}) = \mathcal{H}om_{\mathcal{O}_X}(\mathcal{E}, \Gamma_Y \mathcal{F}),$$

and then extends these definitions to pairs of complexes of  $\mathcal{O}_X$ -modules, as usual.

Assume  $L^\bullet$  is bounded below. Given a bounded below injective resolution  $I^\bullet$  of  $L^\bullet$ , the components of the complex  $\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet)$  are flasques; hence,

$$\begin{aligned} R\Gamma_Y(R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, L^\bullet)) &= R\Gamma_Y(\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet)) \\ &= \Gamma_Y(\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, I^\bullet)) \\ &= \mathcal{H}om_Y^\bullet(K^\bullet, I^\bullet), \end{aligned}$$

in the following cases:

- (a)  $K^\bullet$  is bounded below,
- (b) there exists a  $d$  such that, for any  $\mathcal{O}_X$ -module  $\mathcal{E}$ ,  $H_Y^i(X, \mathcal{E}) = 0$  for  $i > d$ .

In cases (a) and (b),  $\mathcal{H}om_Y^\bullet(K^\bullet, I^\bullet)$  is independent of the choice of  $I^\bullet$  (up to an isomorphism in  $D(X)$ ), and one sets

$$R\Gamma_Y^\bullet(K^\bullet, L^\bullet) := \mathcal{H}om_Y^\bullet(K^\bullet, I^\bullet).$$

As the components of  $\Gamma_Y I^\bullet$  are injective  $\mathcal{O}_X$ -modules,  $\mathcal{H}om_Y^\bullet(K^\bullet, I^\bullet) \simeq \mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \Gamma_Y I^\bullet) \simeq R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, R\Gamma_Y L^\bullet)$ . In other words,

$$R\Gamma_Y^\bullet(K^\bullet, L^\bullet) \simeq R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, R\Gamma_Y L^\bullet)$$

if (a) or (b) holds.

**6.2. Some spectral sequences.** Let  $A$  be a commutative ring (with unit) and  $I$  a finitely generated  $A$ -ideal. Set  $(X, \mathcal{O}_X) := (\mathrm{Spec}(A), \tilde{A})$  and  $Y := V(I) \subset X$ .

Assume that there exists an  $n$  such that

$$H_I^i(A) = H_Y^i(X, \mathcal{O}_X) = 0, \quad \text{for } i \neq n.$$

Then  $R\Gamma_Y(\mathcal{O}_X) \simeq \mathbf{H}_Y^n(\mathcal{O}_X)[-n]$  in  $D^+(X, \mathcal{O}_X)$ . Given a complex  $K^\bullet \in D^-(X, \mathcal{O}_X)$ , it follows that

$$\begin{aligned} R\Gamma_Y(R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \mathcal{O}_X)) &\simeq R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, R\Gamma_Y(\mathcal{O}_X)) \\ &\simeq R\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \mathbf{H}_Y^n(\mathcal{O}_X))[-n] \end{aligned}$$

in  $D(X, \mathcal{O}_X)$ . Such an isomorphism holds for  $K^\bullet \in D(X, \mathcal{O}_X)$  when  $X$  has finite homological dimension, hence, for instance, if  $A$  is Noetherian of finite dimension.

Assuming further that the components of  $K^\bullet$  are locally free  $\mathcal{O}_X$ -modules of finite type, and under one of the two hypotheses above, one has

$$\begin{aligned} R\Gamma_Y(\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \mathcal{O}_X)) &\simeq \mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \mathbf{H}_Y^n(\mathcal{O}_X))[-n] \\ &\text{in } D(X, \mathcal{O}_X). \end{aligned}$$

This provides a spectral sequence

$$\begin{aligned} E_2^{p,q} &= \mathbf{H}_Y^p(H^q \mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \mathcal{O}_X)) \\ &\implies H^{p+q-n}(\mathcal{H}om_{\mathcal{O}_X}^\bullet(K^\bullet, \mathbf{H}_Y^n(\mathcal{O}_X))), \end{aligned}$$

in the two cases above.

As the  $\mathcal{O}_X$ -modules taking place in this spectral sequence are quasi-coherent, and the components of  $K^\bullet$  are finitely presented (recall that  $\widetilde{\text{Hom}}_A(\widetilde{M}, \widetilde{N}) = \mathcal{H}om_{\mathcal{O}_X}(\widetilde{M}, \widetilde{N})$  if  $M$  is finitely presented) the above results show the following proposition.

**Proposition 6.1.** *Let  $A$  be a commutative ring and  $I$  a finitely generated  $A$ -ideal. Assume that there exists an  $n$  such that*

$$H_I^i(A) = H_Y^i(X, \mathcal{O}_X) = 0 \quad \text{for } i \neq n.$$

*Then, for any bounded below complex of projective  $A$ -modules of finite type (respectively, for any complex of projective  $A$ -modules of finite*

type if  $A$  is Noetherian of finite dimension)  $K^\bullet$ , there exists a spectral sequence

$$E_2^{p,q} = H_I^p(H^q \text{Hom}_A^\bullet(K^\bullet, A)) \implies H^{p+q-n}(\text{Hom}_A^\bullet(K^\bullet, H_I^n(A))).$$

**Corollary 6.2.** *Assume  $(A, \mathfrak{m})$  is local Gorenstein of dimension  $n$ . Then, for any complex  $K^\bullet$  of free  $A$ -modules of finite type, there is a spectral sequence*

$$E_2^{p,q} = H_{\mathfrak{m}}^p(H^q \text{Hom}_A^\bullet(K^\bullet, A)) \implies H^{p+q-n}(\text{Hom}_A^\bullet(K^\bullet, H_{\mathfrak{m}}^n(A))).$$

*Proof.* Under the hypotheses of Corollary 6.2,  $H_{\mathfrak{m}}^i(A) = 0$  for  $i \neq n$  and  $H_{\mathfrak{m}}^n(A)$  is injective.  $\square$

**Example 6.3.** In the context of Corollary 6.2, taking for  $K^\bullet$  the dual of a resolution of finitely generated module  $M$  by free modules of finite type, gives the local duality

$$H_{\mathfrak{m}}^p(M) \simeq \text{Hom}_A(\text{Ext}_A^{n-p}(M, A), H_{\mathfrak{m}}^n(A)).$$

**Corollary 6.4.** *Assume  $I$  is generated by a weakly regular sequence of length  $n$ . Then, for any bounded below complex  $K^\bullet$  of projective  $A$ -modules of finite type, there is a spectral sequence*

$$E_2^{p,q} = H_I^p(H^q \text{Hom}_A^\bullet(K^\bullet, A)) \implies H^{p+q-n}(\text{Hom}_A^\bullet(K^\bullet, H_I^n(A))).$$

**Example 6.5.** Let  $R$  be a commutative ring,  $X_i$  for  $1 \leq i \leq n$  indeterminates, set  $A := R[X_1, \dots, X_n]$ , with its standard grading and  $\mathfrak{p} := (X_1, \dots, X_n)$ . As  $H_{\mathfrak{p}}^n(A) \simeq (X_1 \cdots X_n)^{-1} R[X_1^{-1}, \dots, X_n^{-1}]$ , it follows that, for any graded free  $A$ -module of finite type  $F$  and every integer  $\nu$ , the pairing

$$\begin{aligned} \text{Hom}_A(F, H_{\mathfrak{p}}^n(A))_{-\nu-n} \otimes_R F_\nu &\longrightarrow H_{\mathfrak{p}}^n(A)_{-n} \simeq R \\ (u : F \rightarrow H_{\mathfrak{p}}^n(A)(-\nu - n)) \otimes_R x &\longmapsto u(x) \end{aligned}$$

defines a perfect duality between  $R$ -modules of finite type, and this duality is functorial in the free graded  $A$ -module  $F$ . It gives, for each integer  $\nu$ , an isomorphism of complexes of  $R$ -modules

$$\mathrm{Hom}_A^\bullet(K^\bullet, H_{\mathfrak{p}}^n(A))_{-\nu-n} \simeq \mathrm{Hom}_R^\bullet(K_{-\nu}^\bullet, R).$$

Together with Corollary 6.4, it gives for any  $\nu$  a spectral sequence of  $R$ -modules

$$E_2^{p,q} = H_{\mathfrak{p}}^p(H^q \mathrm{Hom}_A^\bullet(K^\bullet, A))_\nu \implies H^{p+q-n}(\mathrm{Hom}_R^\bullet(K_{-\nu-n}^\bullet, R)).$$

Replacing  $K^\bullet$  by its dual  $F^\bullet$ , one deduces of a spectral sequence

$$E_2^{p,q} = H_{\mathfrak{p}}^p(H^q(F^\bullet)_\nu) \implies H^{p+q-n}(\mathrm{Hom}_R^\bullet(\mathrm{Hom}_A^\bullet(F^\bullet, A)_{-\nu-n}, R)).$$

For instance, if  $M$  is a graded  $A$ -module admitting a resolution by free modules of finite rank, taking for  $K^\bullet$  such a resolution, that one may assume to be graded, a spectral sequence follows:

$$E_2^{p,q} = H_{\mathfrak{p}}^p(\mathrm{Ext}_A^q(M, A))_\nu \implies \mathrm{Ext}_R^{p+q-n}(M_{-\nu-n}, R).$$

**6.3. The Herzog-Rahimi spectral sequences.** We keep notations as in the preceding subsection. For any graded complex  $K^\bullet$  whose components are of finite type, we have established an isomorphism

$$R\Gamma_{\mathfrak{p}}(\mathrm{Hom}_A^\bullet(K^\bullet, A)) \simeq \mathrm{Hom}_A^\bullet(K^\bullet, H_{\mathfrak{p}}^n(A))[-n] \quad \text{in } D(A),$$

whenever  $K^\bullet$  is bounded above or  $A$  is Noetherian of finite dimension.

In each of these cases, for any integer  $\nu$ , the isomorphisms

$$R\Gamma_{\mathfrak{p}}(\mathrm{Hom}_A^\bullet(K^\bullet, A))_\nu \simeq \mathrm{Hom}_R^\bullet(K_{-\nu-n}^\bullet, R)[-n] \quad \text{in } D(R)$$

follow. Hence, if  $F^\bullet$  is a graded complex of finite free  $A$ -modules, and either  $A$  is Noetherian of finite dimension or  $F^\bullet$  is bounded below, one has the isomorphisms

$$\begin{aligned} R\Gamma_{\mathfrak{p}}(F^\bullet)_\nu &\simeq \mathrm{Hom}_R^\bullet(\mathrm{Hom}_A^\bullet(F^\bullet, A)_{-\nu-n}, R)[-n] \\ &\simeq R\mathrm{Hom}_R^\bullet(\mathrm{Hom}_A^\bullet(F^\bullet, A)_{-\nu-n}, R)[-n] \quad \text{in } D(R). \end{aligned}$$



Now, assume further that  $(R, \mathfrak{m})$  is local Gorenstein of dimension  $d$ . The above isomorphisms then give

$$\begin{aligned} R\mathrm{Hom}_R^\bullet(R\Gamma_{\mathfrak{p}}(F^\bullet)_\nu, H_{\mathfrak{m}}^d(R)) \\ &\simeq R\mathrm{Hom}_R^\bullet(R\Gamma_{\mathfrak{p}}(F^\bullet)_\nu, R\Gamma_{\mathfrak{m}}(R))[d] \\ &\simeq R\Gamma_{\mathfrak{m}}(R\mathrm{Hom}_R^\bullet(R\Gamma_{\mathfrak{p}}(F^\bullet)_\nu, R))[d] \\ &\simeq R\Gamma_{\mathfrak{m}}(R\mathrm{Hom}_R^\bullet(R\mathrm{Hom}_R^\bullet(\mathrm{Hom}_A^\bullet(F^\bullet, A)_{-\nu-n}, R), R))[n+d] \\ &\simeq R\Gamma_{\mathfrak{m}}(\mathrm{Hom}_A^\bullet(F^\bullet, A)_{-\nu-n})[n+d] \end{aligned}$$

As  $R$  is Gorenstein,  $H_{\mathfrak{m}}^d(R)$  is the injective envelope of the residue field of  $R$ , and we obtain a spectral sequence

$$\begin{aligned} E_2^{p,q} &= H_{\mathfrak{m}}^p(H^q(\mathrm{Hom}_A^\bullet(F^\bullet, A)_{-\nu-n})) \\ &\implies \mathrm{Hom}_R(H^{n+d-p-q}(R\Gamma_{\mathfrak{p}}(F^\bullet)_\nu), H_{\mathfrak{m}}^d(R)). \end{aligned}$$

If  $M$  is a graded  $A$ -module with free resolution  $F^\bullet$ , this spectral sequence takes the form

$$E_2^{p,q} = H_{\mathfrak{m}}^p(\mathrm{Ext}_A^q(M, A)_{-\nu-n}) \implies \mathrm{Hom}_R(H_{\mathfrak{p}}^{n+d-p-q}(M)_\nu, H_{\mathfrak{m}}^d(R)),$$

which is the Herzog-Rahimi spectral sequence, as  $\omega_A \simeq A(-n)$  in this situation.

**7. Tameness of local cohomology over Noetherian rings.** In this section  $S$  is a finitely generated standard graded algebra over an epimorphic image  $R$  of a Gorenstein ring.

**Theorem 7.1.** *Let  $(R, \mathfrak{m})$  be a local Noetherian Gorenstein ring of dimension  $d$ ,  $S$  a finitely generated standard graded Cohen-Macaulay algebra over  $R$  and  $M$  a finitely generated graded  $S$ -module. Set  $-\vee := \mathrm{Hom}_R(-, H_{\mathfrak{m}}^d(R))$ . Then, there is a spectral sequence*

$$E_2^{i,j} = H_{\mathfrak{m}}^i(\mathrm{Ext}_S^j(M, \omega_S)_\gamma) \implies (H_{S_+}^{\dim S - (i+j)}(M)_{-\gamma})^\vee.$$

*Proof.* See [12, Section 4] or the previous section.  $\square$

For an  $R$ - or an  $S$ -module  $M$ , we set

$$H_{[i]}^0(M) := \bigcup_{\substack{I \subseteq R \\ \dim(\bar{R}/I) \leq i}} H_I^0(M) \subseteq M.$$

We will need the following facts about the functor  $H_{[i]}^0(-)$ ,

**Lemma 7.2.** *Let  $M$  be a graded  $S$ -module. Then*

- (i)  $H_{[i]}^0(M)_\gamma = H_{[i]}^0(M_\gamma)$ .
- (ii) *If  $\mathfrak{p}$  a prime ideal of  $R$  with  $\dim(R/\mathfrak{p}) = i$ , then*

$$H_{[i]}^0(M) \otimes_R R_{\mathfrak{p}} = H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}}).$$

*Proof.* Claim (i) follows from the fact that, for any  $R$ -ideal  $I$ ,  $H_I^0(M)_\gamma = H_I^0(M_\gamma)$ . For (ii), recall that inductive limits commute with tensor products, and notice that, if  $\dim(R/I) \leq i$ , then  $H_{I_{\mathfrak{p}}}^0(M \otimes_R R_{\mathfrak{p}}) = 0$  if  $I \not\subseteq \mathfrak{p}$ , and  $H_{I_{\mathfrak{p}}}^0(M \otimes_R R_{\mathfrak{p}}) = H_{\mathfrak{p}}^0(M \otimes_R R_{\mathfrak{p}})$  otherwise.  $\square$

**Theorem 7.3.** *Let  $S$  be a polynomial ring in  $n$  variables over an equidimensional Gorenstein ring  $R$  of dimension  $d$ . Let  $M$  be a finitely generated graded  $S$ -module.*

*Then there exist  $A, B, C, D, E$  defined below such that:*

- (1) (a) *For  $\gamma > A$ ,  $\dim H_{S_+}^i(M)_{-\gamma} = d \Rightarrow \dim H_{S_+}^i(M)_{-\gamma-1} = d$ .*
- (b) *For  $\gamma > B$ ,  $\dim H_{S_+}^i(M)_{-\gamma} = d \Leftrightarrow \dim H_{S_+}^i(M)_{-\gamma-1} = d$ .*
- (2) (a) *For  $\gamma > C$ ,  $\dim H_{S_+}^i(M)_{-\gamma} \geq d - 1 \Rightarrow \dim H_{S_+}^i(M)_{-\gamma-1} \geq d - 1$ .*
- (b) *For  $\gamma > D$ ,  $\dim H_{S_+}^i(M)_{-\gamma} \geq d - 1 \Leftrightarrow \dim H_{S_+}^i(M)_{-\gamma-1} \geq d - 1$ .*
- (3) *For  $\gamma > E$ ,  $\dim H_{S_+}^i(M)_{-\gamma} \geq d - 2 \Rightarrow \dim H_{S_+}^i(M)_{-\gamma-1} \geq d - 2$ .*

*In particular, there exists a  $\gamma_0$  such that either  $\dim H_{S_+}^i(M)_\gamma$  is constant for  $\gamma < \gamma_0$  of value at least  $d - 2$ , or  $\dim H_{S_+}^i(M)_\gamma < d - 2$  for any  $\gamma < \gamma_0$ .*

Set  $a_j^i := \text{end}(H_{S_+}^0(H_{[d-j]}^0(\text{Ext}_S^i(M, \omega_S)))) \leq \text{end}(H_{S_+}^0(\text{Ext}_S^i(M, \omega_S)))$   
and  $r_j^i := \text{reg}(H_{[d-j]}^0(\text{Ext}_S^i(M, \omega_S)))$ . Then one has:

$$\begin{aligned} A &:= a_0^{n-i}, & B &:= r_0^{n-i}, \\ C &:= \max\{a_1^{n-i+1}, a_0^{n-i}\}, & D &:= \max\{r_1^{n-i+1}, r_0^{n-i}\} \end{aligned}$$

and

$$E := \max\{a_0^{n-i}, r_0^{n-i+1} - 1, r_2^{n-i+1} - 2, a_2^{n-i+2}\}.$$

*Proof.* Recall that, if  $N$  is a finitely generated  $R$ -module,  $\dim N < r$  if and only if  $N_{\mathfrak{p}} = 0$  for all  $\mathfrak{p} \in \text{Spec}(R)$  with  $\dim(R/\mathfrak{p}) = r$ . Furthermore, it follows from Lemma 7.2 and from Lemma 2.5 and its proof that, for any  $\mathfrak{p} \in \text{Spec}(R)$  with  $\dim(R/\mathfrak{p}) = r$  and for any  $\ell$ , one has

$$a_{S_+}^\ell(H_{\mathfrak{p}}^0(\text{Ext}_{S \otimes_R R_{\mathfrak{p}}}^i(M \otimes_R R_{\mathfrak{p}}, \omega_{S \otimes_R R_{\mathfrak{p}}})) \leq a_{S_+}^\ell(H_{[r]}^0(\text{Ext}_S^i(M, \omega_S))).$$

Therefore,  $\text{reg}(H_{\mathfrak{p}}^0(\text{Ext}_{S \otimes_R R_{\mathfrak{p}}}^i(M \otimes_R R_{\mathfrak{p}}, \omega_{S \otimes_R R_{\mathfrak{p}}})) \leq r_{n-r}^i$ , and, as a consequence, it suffices to prove (1) (a) and (1) (b) when  $\dim R = 0$ , (2) (a) and (2) (b) when  $\dim R = 1$  and (3) when  $\dim R = 2$ .

Notice that  $H_{[d]}^0(N) = N$  for any  $S$ -module  $N$ .

If  $\dim R = 0$ , by Theorem 7.1,  $(H_{S_+}^i(M)_{-\gamma})^\vee \simeq \text{Ext}_S^{n-i}(M, \omega_S)_\gamma$ , and the result follows from Lemma 5.5 (i) and (iii).

If  $\dim R = 1$ , Theorem 7.1 provides exact sequences

$$\begin{aligned} 0 \rightarrow H_{\mathfrak{m}}^1(\text{Ext}_S^{n-i}(M, \omega_S)_\gamma) \rightarrow (H_{S_+}^i(M)_{-\gamma})^\vee \\ \rightarrow H_{\mathfrak{m}}^0(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma) \rightarrow 0. \end{aligned}$$

The result follows applying Lemma 5.5 (i) and (iii) to  $H_{\mathfrak{m}}^0(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma)$  and Corollary 5.4 to  $\text{Ext}_S^{n-i}(M, \omega_S)_\gamma$ . Indeed,  $H_{\mathfrak{m}}^1(\text{Ext}_S^{n-i}(M, \omega_S)_\gamma)$  is zero if and only if  $\dim \text{Ext}_S^{n-i}(M, \omega_S)_\gamma < 1$ .

We now assume that  $\dim R = 2$ . In this case, Theorem 7.1 provides a spectral sequence which converges to  $(H_{S_+}^\bullet(M)_{-\gamma})^\vee$ ,

$$\begin{array}{ccccc}
& & H_m^0(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma) & & H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S)_\gamma) \\
& & \swarrow & & \swarrow \\
\cdots & & & & \\
& \psi_\gamma^{n-i+1} & H_m^1(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma) & \psi_\gamma^{n-i+2} & H_m^1(\text{Ext}_S^{n-i+2}(M, \omega_S)_\gamma) \\
& & \swarrow & & \swarrow \\
H_m^2(\text{Ext}_S^{n-i}(M, \omega_S)_\gamma) & & H_m^2(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma) & & \cdots
\end{array}$$

It provides a filtration  $F_*^0 \subseteq F_*^1 \subseteq F_*^2 = (H_{S_+}^i(M)_{-*})^\vee$ , by graded  $S$ -modules, such that  $F_\gamma^2/F_\gamma^1 \simeq \ker(\psi_\gamma^{n-i+2})$ ,  $F_\gamma^1/F_\gamma^0 \simeq H_m^1(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma)$  and  $F_\gamma^0 \simeq \text{coker}(\psi_\gamma^{n-i+1})$ .

We will show that the three modules satisfy :

- (i)  $F_\gamma^0 \neq 0 \Rightarrow F_{\gamma+1}^0 \neq 0$ , if  $\gamma > a_0^{n-i}$ ,
- (ii)  $F_\gamma^1/F_\gamma^0 \neq 0 \Rightarrow F_{\gamma+1}^1/F_{\gamma+1}^0 \neq 0$  if  $\gamma > \max\{r_0^{n-i+1}-1, r_2^{n-i+1}-2\}$ ,
- (iii)  $F_\gamma^2/F_\gamma^1 \neq 0 \Rightarrow F_{\gamma+1}^2/F_{\gamma+1}^1 \neq 0$  if  $\gamma > a_2^{n-i+2}$ .

For (i), notice that  $\text{coker}(\psi_\gamma^{n-i+1}) = 0$  if and only if  $H_m^2(\text{Ext}_S^{n-i}(M, \omega_S)_\gamma) = 0$ , hence if and only if  $\dim(\text{Ext}_S^{n-i}(M, \omega_S)_\gamma) < 2$ . Hence, (i) follows from Corollary 5.4.

For (ii), let  $N := \text{Ext}_S^{n-i+1}(M, \omega_S)/H_m^0(\text{Ext}_S^{n-i+1}(M, \omega_S))$ . The exact sequence

$$0 \longrightarrow H_m^0(\text{Ext}_S^{n-i+1}(M, \omega_S)) \longrightarrow \text{Ext}_S^{n-i+1}(M, \omega_S) \longrightarrow N \longrightarrow 0$$

shows that  $H_m^0(N) = 0$  (hence,  $\text{depth}(N) \geq 1$ ),

$$F^1/F^0 = H_m^1(\text{Ext}_S^{n-i+1}(M, \omega_S)) \simeq H_m^1(N)$$

and

$$\begin{aligned}
a_{S_+}^1(N) &\leq \max\{a_{S_+}^1(\text{Ext}_S^{n-i+1}(M, \omega_S)), a_{S_+}^2(H_m^0(\text{Ext}_S^{n-i+1}(M, \omega_S)))\} \\
&\leq \max\{r_0^{n-i+1} - 1, r_2^{n-i+1} - 2\}.
\end{aligned}$$

Hence, Proposition 3.1 (ii) implies (ii).

For (iii), let  $\alpha \in H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S)_\gamma) = H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S))_\gamma$ .

Let  $x_1, \dots, x_t$  be generators of  $S_1$  as an  $R$ -module, and set  $\ell := \sum_i x_i T_i \in S[\underline{T}]$ . For  $\gamma > a_{S_+}^0(H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S)))$ ,  $\ell\alpha$  is not zero in  $H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S)_{\gamma+1})[\underline{T}]$  by Theorem 1.5 (see also the proof of Theorem 5.3). The commutative diagram

$$\begin{array}{ccc} H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S)_\gamma) & \xrightarrow{\times \ell} & H_m^0(\text{Ext}_S^{n-i+2}(M, \omega_S)_{\gamma+1})[\underline{T}] \\ \psi_\gamma^{n-i+2} \downarrow & & \downarrow \psi_{\gamma+1}^{n-i+2} \otimes_S 1_{S[\underline{T}]} \\ H_m^2(\text{Ext}_S^{n-i+1}(M, \omega_S)_\gamma) & \xrightarrow{\times \ell} & H_m^2(\text{Ext}_S^{n-i+1}(M, \omega_S)_{\gamma+1})[\underline{T}] \end{array}$$

then shows that  $\psi_\gamma^{n-i+2}(\alpha) \neq 0$  if  $\psi_{\gamma+1}^{n-i+2}$  is injective. Hence,  $\psi_\gamma^{n-i+2}$  is injective if  $\psi_{\gamma+1}^{n-i+2}$  is. Claim (iii) follows.  $\square$

**Theorem 7.4.** *Let  $S$  be a Noetherian standard graded algebra over a commutative ring  $R$ . Assume  $R$  has dimension at most two and either  $R$  is an epimorphic image of a Gorenstein ring or  $R$  is local. Let  $M$  be a finitely generated graded  $S$ -module.*

*Then there exists a  $\gamma_0$  such that, for any  $i$ ,*

$$\{H_{S_+}^i(M)_\gamma = 0 \text{ for } \gamma < \gamma_0\} \quad \text{or} \quad \{H_{S_+}^i(M)_\gamma \neq 0 \text{ for } \gamma < \gamma_0\}.$$

*Proof.* First, if  $R$  is local, then we can complete  $R$  to reduce to the case where  $R$  is a quotient of a regular ring (by Cohen structure theorem); hence, an epimorphic image of a Gorenstein ring.

As a Gorenstein ring is a finite product of equidimensional Gorenstein rings, and each such ring is itself a quotient of a Gorenstein ring of any bigger dimension,  $R$  is also a quotient of an equidimensional Gorenstein ring  $R'$ . We further remark that  $R$  is the epimorphic image of  $R'/K$ , where  $K$  is generated by a regular sequence of length  $\dim R' - 2$  in  $R'$ .

Thus, we may, and will, assume that  $R$  is an equidimensional Gorenstein ring of dimension at most two. Now  $S$  is an epimorphic image of a polynomial ring in a finite number of variables over  $R$ , so that we may, and will, also assume that  $S$  is a polynomial ring over  $R$ .

The result then follows from Theorem 7.3.  $\square$

## REFERENCES

1. Markus Brodmann, *Asymptotic stability of Ass* ( $M/I^n M$ ), Proc. Amer. Math. Soc. **74** (1979), 16–18.
2. ———, *A cohomological stability result for projective schemes over surfaces*, J. reine angew. Math. **606** (2007), 179–192.
3. S. Dale Cutkosky and Jürgen Herzog, *Failure of tameness for local cohomology*, J. Pure Appl. Algebra **211** (2007), 428–432.
4. Mohammad T. Dibaei and Alireza Vahidi, *Artinian and non-Artinian local cohomology modules*, Canad. Math. Bull. **54** (2011), 619–629.
5. K. Divaani-Aazar, R. Naghipour and M. Tousi, *Cohomological dimension of certain algebraic varieties*, Proc. Amer. Math. Soc. **130** (2002), 3537–3544.
6. Roger Godement, *Topologie algébrique et théorie des faisceaux*, Actual.Sci.Industr. **1252**, Hermann, Paris, 1973.
7. Alexandre Grothendieck, *Éléments de géométrie algébrique*, III. *Étude cohomologique des faisceaux cohérents*, I, Inst. Hautes Études Sci. Publ. Math. **11**, 1961.
8. Jean-Pierre Jouanolou, *Monoides*, Publ. l’Inst. Rech. Math. Avanc., Université Louis Pasteur, 1986.
9. George R. Kempf, *Some elementary proofs of basic theorems in the cohomology of quasicoherent sheaves*, Rocky Mountain J. Math. **10** (1980), 637–645.
10. D.G. Northcott, *Finite free resolutions*, Cambridge University Press, Cambridge, 1976.
11. A. Ooishi, *Castelnuovo’s regularity of graded rings and modules*, Hiroshima Math. J. **12** (1982), 627–644.
12. Ahad Rahimi, *Local cohomology of bigraded modules*, Ph. D. thesis, Univ. Essen, 2007.
13. M. Stokes, *Some dual homological results for modules over commutative rings*, J. Pure Appl. Algebra **65** (1990), 153–162.
14. A.A. Suslin, *Triviality of certain cohomology groups*, Z. Nauch. Sem. Lenin. Otdel. Mat. Inst. Steklov. (LOMI) **94** (1979), 114–115; J. Math. Sci. **19** (1982), 1048–1049 (in English).

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UPMC, BOITE 247, 4, PLACE JUSSIEU,  
F-75252 PARIS CEDEX, FRANCE

**Email address:** [chardin@math.jussieu.fr](mailto:chardin@math.jussieu.fr)

IRMA, UNIVERSITÉ DE STRASBOURG, 7, RUE RENÉ DESCARTES F-67084 STRASBOURG, FRANCE

**Email address:** [jean-pierre.jouanolou@math.unistra.fr](mailto:jean-pierre.jouanolou@math.unistra.fr)

DEPARTMENT OF MATHEMATICS, RAZI UNIVERSITY, KERMANSHAH, IRAN AND  
SCHOOL OF MATHEMATICS, IPM, P. O. BOX 19395-5746, TEHRAN, IRAN

**Email address:** [ahad.rahimi@razi.ac.ir](mailto:ahad.rahimi@razi.ac.ir)