

## ANOTHER DEFINITION OF AN EULER CLASS GROUP OF A NOETHERIAN RING

MANOJ K. KESHARI AND SATYA MANDAL

**1. Introduction.** All the rings are assumed to be commutative Noetherian and all the modules are finitely generated.

Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $L$  be a projective  $A$ -module of rank 1. In [3], Bhatwadekar and Sridharan defined an abelian group, called the Euler class group of  $A$  with respect to  $L$  which is denoted by  $E(A, L)$ . To the pair  $(P, \chi)$ , where  $P$  is a projective  $A$ -module of rank  $n$  with determinant  $L$  and  $\chi : L \xrightarrow{\sim} \wedge^n P$  an isomorphism, called an  $L$ -orientation of  $P$ , they attached an element of  $E(A, L)$  which is denoted by  $e(P, \chi)$ . One of the main result in [3] is that  $P$  has a unimodular element if and only if  $e(P, \chi)$  is zero in  $E(A, L)$ .

We will define the Euler class group of  $A$  with respect to a projective  $A$ -module  $F = Q \oplus A$  of rank  $n$ , denoted by  $E(A, F)$ . To the pair  $(P, \chi)$ , where  $P$  is a projective  $A$ -module of rank  $n$  and  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  is an isomorphism, called an  $F$ -orientation of  $P$ , we associate an element of the Euler class group, denoted by  $e(P, \chi)$  and prove the following result:  $P$  has a unimodular element if and only if  $e(P, \chi)$  is zero in  $E(A, F)$ . Note that, when  $F = L \oplus A^{n-1}$ ,  $E(A, F)$  is the same as the Euler class group  $E(A, L)$  defined in [3].

**2. Preliminaries.** Let  $A$  be a ring, and let  $M$  be an  $A$ -module. For  $m \in M$ , we define  $O_M(m) = \{\varphi(m) \mid \varphi \in \text{Hom}_A(M, A)\}$ . We say that  $m$  is *unimodular* if  $O_M(m) = A$ . The set of all unimodular elements of  $M$  will be denoted by  $\text{Um}(M)$ . Note that, if a projective  $A$ -module  $P$  has a unimodular element, then  $P \xrightarrow{\sim} P_1 \oplus A$ .

Let  $P$  be a projective  $A$ -module. Given an element  $\varphi \in P^*$  and an element  $p \in P$ , we define an endomorphism  $\varphi_p$  as the composite

---

The first author supported by BOYSCAST Fellowship 2008–09 of Department of Science and Technology, India.

Received by the editors on December 19, 2009, and in revised form on June 14, 2010.

$P \xrightarrow{\varphi} A \xrightarrow{p} P$ . If  $\varphi(p) = 0$ , then  $\varphi_p^2 = 0$  and hence  $1 + \varphi_p$  is a unipotent automorphism of  $P$ .

By a *transvection*, we mean an automorphism of  $P$  of the form  $1 + \varphi_p$ , where  $\varphi(p) = 0$  and either  $\varphi$  is unimodular in  $P^*$  or  $p$  is unimodular in  $P$ . We denote by  $\text{EL}(P)$  the subgroup of  $\text{Aut}(P)$  generated by all the transvections of  $P$ . Note that  $\text{EL}(P)$  is a normal subgroup of  $\text{Aut}(P)$ .

Recall that, if  $A$  is a ring of dimension  $n$  and if  $P$  is a projective  $A$ -module of rank  $n$ , then any surjection  $\alpha : P \twoheadrightarrow J$  is called a *generic* surjection of  $P$  if  $J$  is an ideal of  $A$  of height  $n$ .

The following result is due to Bhatwadekar and Roy ([2, Proposition 4.1]):

**Proposition 2.1.** *Let  $B$  be a ring, and let  $I$  be an ideal of  $B$ . Let  $P$  be a projective  $B$ -module. Then any element of  $\text{EL}(P/IP)$  can be lifted to an automorphism of  $P$ .*

We state some results from [3] for later use.

**Lemma 2.2** [3, Lemma 3.0]. *Let  $A$  be a ring of dimension  $n$ , and let  $P$  be a projective  $A$ -module of rank  $n$ . Let  $\lambda : P \twoheadrightarrow J_0$  and  $\mu : P \twoheadrightarrow J_1$  be two surjections, where  $J_0$  and  $J_1$  are ideals of  $A$  of height  $n$ . Then there exists an ideal  $I$  of  $A[T]$  of height  $n$  and a surjection  $\alpha(T) : P[T] \twoheadrightarrow I$  such that  $I(0) = J_0$ ,  $I(1) = J_1$ ,  $\alpha(0) = \lambda$  and  $\alpha(1) = \mu$ .*

For a rank 1 projective  $A$ -module  $L$  and  $P' = L \oplus A^{n-1}$ , the following result is proved in [3, Proposition 3.1]. Since the same proof works in our case, we omit the proof.

**Proposition 2.3.** *Let  $A$  be a ring of dimension  $n \geq 2$  such that  $(n - 1)!$  is a unit in  $A$ . Let  $P$  and  $P' = Q \oplus A$  be projective  $A$ -modules of rank  $n$ , and let  $\chi : \wedge^n P \xrightarrow{\sim} \wedge^n P'$  be an isomorphism. Suppose that  $\alpha(T) : P[T] \twoheadrightarrow I$  is a surjection, where  $I$  is an ideal of  $A[T]$  of height  $n$ . Then there exists a homomorphism  $\phi : P' \rightarrow P$ , an ideal  $K$  of  $A$  of height  $\geq n$  which is comaximal with  $(I \cap A)$  and a surjection  $\rho(T) : P'[T] \twoheadrightarrow I \cap KA[T]$  such that the following holds:*

- (i)  $\wedge^n(\phi) = u\chi$ , where  $u = 1$  modulo  $I \cap A$ .

- (ii)  $(\alpha(0) \circ \phi)(P') = I(0) \cap K.$
- (iii)  $(\alpha(T) \circ \phi(T)) \otimes A[T]/I = \rho(T) \otimes A[T]/I.$
- (iv)  $\rho(0) \otimes A/K = \rho(1) \otimes A/K.$

**Theorem 2.4** (Addition principle [3, Theorem 3.2]). *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $J_1, J_2$  be two comaximal ideals of  $A$  of height  $n$ . Let  $P = P_1 \oplus A$  be a projective  $A$ -module of rank  $n$ , and let  $\Phi : P \twoheadrightarrow J_1$  and  $\Psi : P \twoheadrightarrow J_2$  be two surjections. Then, there exists a surjection  $\Theta : P \twoheadrightarrow J_1 \cap J_2$  such that  $\Phi \otimes A/J_1 = \Theta \otimes A/J_1$  and  $\Psi \otimes A/J_2 = \Theta \otimes A/J_2$ .*

**Theorem 2.5** (Subtraction principle [3, Theorem 3.3]). *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $J$  and  $J'$  be two comaximal ideals of  $A$  of height  $\geq n$  and  $n$ , respectively. Let  $P$  and  $P' = Q \oplus A$  be projective  $A$ -modules of rank  $n$ , and let  $\chi : \wedge^n P \xrightarrow{\sim} \wedge^n P'$  be an isomorphism. Let  $\alpha : P \twoheadrightarrow J \cap J'$  and  $\beta : P' \twoheadrightarrow J'$  be surjections. Let “bar” denote reduction modulo  $J'$ , and let  $\bar{\alpha} : \bar{P} \twoheadrightarrow J'/J'^2$  and  $\bar{\beta} : \bar{P}' \twoheadrightarrow J'/J'^2$  be surjections induced from  $\alpha$  and  $\beta$ , respectively. Suppose there exists an isomorphism  $\delta : \bar{P} \xrightarrow{\sim} \bar{P}'$  such that  $\bar{\beta}\delta = \bar{\alpha}$  and  $\wedge^n(\delta) = \bar{\chi}$ . Then there exists a surjection  $\theta : P \twoheadrightarrow J$  such that  $\theta \otimes A/J = \alpha \otimes A/J$ .*

**Lemma 2.6** [3, Proposition 6.7]. *Let  $A$  be a ring of dimension  $n$ , and let  $P, P'$  be stably isomorphic projective  $A$ -modules of rank  $n$ . Then there exists an ideal  $J$  of  $A$  of height  $\geq n$  such that  $J$  is a surjective image of both  $P$  and  $P'$ . Further, given any ideal  $K$  of height  $\geq 1$ ,  $J$  can be chosen to be comaximal with  $K$ .*

We state the following result from [1, Proposition 2.11] for later use.

**Proposition 2.7.** *Let  $A$  be a ring, and let  $I$  be an ideal of  $A$  of height  $n$ . Let  $f \in A$  be a non-zerodivisor modulo  $I$ , and let  $P = P_1 \oplus A$  be a projective  $A$ -module of rank  $n$ . Let  $\alpha : P \rightarrow I$  be a linear map such that the induced map  $\alpha_f : P_f \twoheadrightarrow I_f$  is a surjection. Then, there exists  $\Psi \in \text{EL}(P_f^*)$  such that:*

- (i)  $\beta = \Psi(\alpha) \in P^*$  and
- (ii)  $\beta(P)$  is an ideal of  $A$  of height  $n$  contained in  $I$ .

**3. Euler class group  $E(A, F)$ .** Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . We define the Euler class group of  $A$  with respect to  $F$  as follows:

Let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements. Let  $\alpha$  and  $\beta$  be two surjections from  $F/JF$  to  $J/J^2$ . We say that  $\alpha$  and  $\beta$  are *related* if there exists an automorphism  $\sigma$  of  $F/JF$  of determinant 1 such that  $\alpha\sigma = \beta$ . Clearly, this is an equivalence relation on the set of all surjections from  $F/JF$  to  $J/J^2$ . Let  $[\alpha]$  denote the equivalence class of  $\alpha$ . We call  $[\alpha]$  a *local  $F$ -orientation* of  $J$ .

Since  $\dim A/J = 0$ ,  $\mathrm{SL}_{A/J}(F/JF) = \mathrm{EL}(F/JF)$  and, therefore, by (2.1), the canonical map from  $\mathrm{SL}_A(F)$  to  $\mathrm{SL}_{A/J}(F/JF)$  is surjective. Hence, if a surjection  $\alpha : F/JF \rightarrow J/J^2$  can be lifted to a surjection  $\Delta : F \rightarrow J$ , then so can any other surjection  $\beta$  equivalent to  $\alpha$ .

A local  $F$ -orientation  $[\alpha]$  is called a *global  $F$ -orientation* of  $J$  if the surjection  $\alpha$  can be lifted to a surjection from  $F$  to  $J$ . From now on, we shall identify a surjection  $\alpha$  with the equivalence class  $[\alpha]$  to which  $\alpha$  belongs.

Let  $\mathcal{M}$  be a maximal ideal of  $A$  of height  $n$ , and let  $\mathcal{N}$  be an  $\mathcal{M}$ -primary ideal such that  $\mathcal{N}/\mathcal{N}^2$  is generated by  $n$  elements. Let  $w_{\mathcal{N}}$  be a local  $F$ -orientation of  $\mathcal{N}$ . Let  $G$  be the free abelian group on the set of pairs  $(\mathcal{N}, w_{\mathcal{N}})$ , where  $\mathcal{N}$  is a  $\mathcal{M}$ -primary ideal and  $w_{\mathcal{N}}$  is a local  $F$ -orientation of  $\mathcal{N}$ .

Let  $J = \cap \mathcal{N}_i$  be the intersection of finitely many  $\mathcal{M}_i$ -primary ideals, where  $\mathcal{M}_i$  are distinct maximal ideals of  $A$  of height  $n$ . Assume that  $J/J^2$  is generated by  $n$  elements, and let  $w_J$  be a local  $F$ -orientation of  $J$ . Then  $w_J$  gives rise, in a natural way, to local  $F$ -orientations  $w_{\mathcal{N}_i}$  of  $\mathcal{N}_i$ . We associate to the pair  $(J, w_J)$ , the element  $\sum(\mathcal{N}_i, w_{\mathcal{N}_i})$  of  $G$ .

Let  $H$  be the subgroup of  $G$  generated by the set of pairs  $(J, w_J)$ , where  $J$  is an ideal of  $A$  of height  $n$  and  $w_J$  is a global  $F$ -orientation of  $J$ .

We define the *Euler class group* of  $A$  with respect to  $F$ , denoted by  $E(A, F)$ , as the quotient group  $G/H$ .

Let  $P$  be a projective  $A$ -module of rank  $n$ , and let  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  be an isomorphism. We call  $\chi$  an  *$F$ -orientation* of  $P$ . To the pair  $(P, \chi)$ , we associate an element  $e(P, \chi)$  of  $E(A, F)$  as follows:

Let  $\lambda : P \twoheadrightarrow J_0$  be a generic surjection of  $P$  and let “bar” denote reduction modulo the ideal  $J_0$ . Then, we obtain an induced surjection  $\bar{\lambda} : \bar{P} \twoheadrightarrow J_0/J_0^2$ . Since  $\dim A/J_0 = 0$ , every projective  $A/J_0$ -module of constant rank is free. Hence, we choose an isomorphism  $\bar{\gamma} : F/J_0F \xrightarrow{\sim} P/J_0P$  such that  $\wedge^n(\bar{\gamma}) = \bar{\chi}$ . Let  $w_{J_0}$  be the local  $F$ -orientation of  $J_0$  given by  $\bar{\lambda} \circ \bar{\gamma} : F/J_0F \twoheadrightarrow J_0/J_0^2$ . Let  $e(P, \chi)$  be the image in  $E(A, F)$  of the element  $(J_0, w_{J_0})$  of  $G$ . We say that  $(J_0, w_{J_0})$  is obtained from the pair  $(\lambda, \chi)$ . We will show that the assignment sending the pair  $(P, \chi)$  to the element  $e(P, \chi)$  of  $E(A, F)$  is well defined.

Let  $\mu : P \twoheadrightarrow J_1$  be another generic surjection of  $P$ . By (2.2), there exists a surjection  $\alpha(T) : P[T] \twoheadrightarrow I$ , where  $I$  is an ideal of  $A[T]$  of height  $n$  with  $\alpha(0) = \lambda$ ,  $I(0) = J_0$ ,  $\alpha(1) = \mu$  and  $I(1) = J_1$ . Using (2.3), we get an ideal  $K$  of  $A$  of height  $n$  and a local  $F$ -orientation  $w_K$  of  $K$  such that  $(I(0), w_{I(0)}) + (K, w_K) = 0 = (I(1), w_{I(1)}) + (K, w_K)$  in  $E(A, F)$ . Therefore,  $(J_0, w_{J_0}) = (J_1, w_{J_1})$  in  $E(A, F)$ . Therefore,  $e(P, \chi)$  is well defined in  $E(A, F)$ .

We define the *Euler class* of  $(P, \chi)$  to be  $e(P, \chi)$ .

For a projective  $A$ -module  $L$  of rank 1 and  $F = L \oplus A^{n-1}$ , the following result is proved in [3, Proposition 4.1]. Since the same proof works in our case, we omit the proof.

**Proposition 3.1.** *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $J, J_1, J_2$  be ideals of  $A$  of height  $n$  such that  $J$  is comaximal with  $J_1$  and  $J_2$ . Let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . Assume that  $\alpha : F \twoheadrightarrow J \cap J_1$  and  $\beta : F \twoheadrightarrow J \cap J_2$  are surjections with  $\alpha \otimes A/J = \beta \otimes A/J$ . Suppose there exists an ideal  $J_3$  of height  $n$  such that:*

- (i)  $J_3$  is comaximal with  $J, J_1$  and  $J_2$  and
- (ii) there exists a surjection  $\gamma : F \twoheadrightarrow J_3 \cap J_1$  with  $\alpha \otimes A/J_1 = \gamma \otimes A/J_1$ .

Then there exists a surjection  $\lambda : F \twoheadrightarrow J_3 \cap J_2$  with  $\lambda \otimes A/J_3 = \gamma \otimes A/J_3$  and  $\lambda \otimes A/J_2 = \beta \otimes A/J_2$ .

Using (2.4), (2.5) and (3.1), and following the proof of [3, Theorem 4.2], the next result follows.

**Theorem 3.2.** *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . Let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements. Let  $w_J : F/JF \twoheadrightarrow J/J^2$  be*

a local  $F$ -orientation of  $J$ . Suppose that the image of  $(J, w_J)$  is zero in  $E(A, F)$ . Then  $w_J$  is a global  $F$ -orientation of  $J$ .

Using (3.2) and (2.5), and following the proof of [3, Corollary 4.3], the next result follows.

**Corollary 3.3.** *Let  $A$  be a ring of dimension  $n \geq 2$ . Let  $P$  and  $F = Q \oplus A$  be projective  $A$ -modules of rank  $n$ , and let  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  be an  $F$ -orientation of  $P$ . Let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements, and let  $w_J$  be a local  $F$ -orientation of  $J$ . Suppose  $e(P, \chi) = (J, w_J)$  in  $E(A, F)$ . Then there exists a surjection  $\alpha : P \twoheadrightarrow J$  such that  $(J, w_J)$  is obtained from  $(\alpha, \chi)$ .*

Using (3.2) and (3.3), and following the proof of [3, Theorem 4.4], the next result follows.

**Corollary 3.4.** *Let  $A$  be a ring of dimension  $n \geq 2$ . Let  $P$  and  $F = Q \oplus A$  be projective  $A$ -modules of rank  $n$ , and let  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  be an  $F$ -orientation of  $P$ . Then  $e(P, \chi) = 0$  in  $E(A, F)$  if and only if  $P$  has a unimodular element.*

Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . Let “bar” denote reduction modulo the nil radical  $N$  of  $A$ , and let  $\overline{A} = A/N$  and  $\overline{F} = F/NF$ . Let  $J$  be an ideal of  $A$  of height  $n$  with primary decomposition  $J = \cap \mathcal{N}_i$ . Then  $\overline{J} = (J + N)/N$  is an ideal of  $\overline{A}$  of height  $n$  with primary decomposition  $\overline{J} = \cap \overline{\mathcal{N}}_i$ . Moreover, any surjection  $w_J : F/JF \twoheadrightarrow J/J^2$  induces a surjection  $\overline{w}_{\overline{J}} : \overline{F}/\overline{J}\overline{F} \twoheadrightarrow \overline{J}/\overline{J}^2 = (J + N)/(J^2 + N)$ . Hence, the assignment sending  $(J, w_J)$  to  $(\overline{J}, \overline{w}_{\overline{J}})$  gives rise to a group homomorphism  $\Phi : E(A, F) \twoheadrightarrow E(\overline{A}, \overline{F})$ .

As a consequence of (3.2), we get the following result, the proof of which is same as of [3, Corollary 4.6].

**Corollary 3.5.** *The homomorphism  $\Phi : E(A, F) \rightarrow E(\overline{A}, \overline{F})$  is an isomorphism.*

**4. Some results on  $E(A, F)$ .** Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . Let  $J$  be an ideal of  $A$  of height  $n$ , and let  $w_J : F/JF \rightarrow J/J^2$  be a surjection. Let  $\bar{b} \in A/J$  be a unit. Then, composing  $w_J$  with an automorphism of  $F/JF$  of determinant  $\bar{b}$ , we get another local  $F$ -orientation of  $J$ , which we denote by  $\bar{b}w_J$ . Further, if  $w_J$  and  $\tilde{w}_J$  are two local  $F$ -orientations of  $J$ , then it is easy to see that  $\tilde{w}_J = \bar{b}w_J$  for some unit  $\bar{b} \in A/J$ .

We recall the following two results from [3, Lemmas 2.7 and 2.8], respectively.

**Lemma 4.1.** *Let  $A$  be a ring, and let  $P$  be a projective  $A$ -module of rank  $n$ . Assume  $0 \rightarrow P_1 \rightarrow A \oplus P \xrightarrow{(b, -\alpha)} A \rightarrow 0$  is an exact sequence. Let  $(a_0, p_0) \in A \oplus P$  be such that  $a_0b - \alpha(p_0) = 1$ . Let  $q_i = (a_i, p_i) \in P_1$  for  $i = 1, \dots, n$ . Then:*

(i) *the map  $\delta : \wedge^n P_1 \rightarrow \wedge^n P$  given by  $\delta(q_1 \wedge \dots \wedge q_n) = a_0(p_1 \wedge \dots \wedge p_n) + \sum_1^n (-1)^i a_i(p_0 \wedge \dots \wedge p_{i-1} \wedge p_{i+1} \wedge \dots \wedge p_n)$  is an isomorphism.*

(ii)  $\delta(bq_1 \wedge \dots \wedge q_n) = p_1 \wedge \dots \wedge p_n$ .

**Lemma 4.2.** *Let  $A$  be a ring, and let  $P$  be a projective  $A$ -module of rank  $n$ . Assume  $0 \rightarrow P_1 \rightarrow A \oplus P \xrightarrow{(b, -\alpha)} A \rightarrow 0$  is an exact sequence. Then:*

(i) *The map  $\beta : P_1 \rightarrow A$  given by  $\beta(q) = c$ , where  $q = (c, p)$ , has the property that  $\beta(P_1) = \alpha(P)$ .*

(ii) *The map  $\Phi : P \rightarrow P_1$  given by  $\Phi(p) = (\alpha(p), bp)$  has the property that  $\beta \circ \Phi = \alpha$  and  $\delta \circ \wedge^n \Phi$  is a scalar multiplication by  $b^{n-1}$ , where  $\delta$  is as in (4.1).*

The following result can be deduced from (4.1) and (4.2). Briefly it says that, if there exists a projective  $A$ -module  $P$  of rank  $n$  with an  $F$ -orientation  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  such that  $e(P, \chi) = (J, w_J)$ , and if  $\bar{a} \in A/J$  is a unit, then there exists another projective  $A$ -module  $P_1$  with  $[P_1] = [P]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_1 : \wedge^n F \xrightarrow{\sim} \wedge^n P_1$  of  $P_1$  such that  $e(P_1, \chi_1) = (J, \bar{a}^{n-1}w_J)$ .

**Lemma 4.3.** *Let  $A$  be a ring of dimension  $n \geq 2$ . Let  $P$  and  $F = Q \oplus A$  be projective  $A$ -modules of rank  $n$ , and let  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  be an  $F$ -orientation of  $P$ . Let  $\alpha : P \rightarrow J$  be a generic surjection of  $P$ ,*

and let  $(J, w_J)$  be obtained from  $(\alpha, \chi)$ . Let  $a, b \in A$  with  $ab = 1$  modulo  $J$ , and let  $P_1$  be the kernel of the surjection  $(b, -\alpha) : A \oplus P \rightarrow A$ . Let  $\beta : P_1 \rightarrow J$  be as in (4.2), and let  $\chi_1$  be the  $\overline{F}$ -orientation of  $P_1$  given by  $\delta^{-1}\chi$ , where  $\delta$  is as in (4.1). Then  $(J, \overline{a^{n-1}w_J})$  is obtained from  $(\beta, \chi_1)$ .

Using the above results and following the proof of [3, Lemmas 5.3, 5.4 and 5.5], respectively, the next three results follow. Note that in these results we need  $F = Q \oplus A^2$ .

**Lemma 4.4.** *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A^2$  be a projective  $A$ -module of rank  $n$ . Let  $J$  be an ideal of  $A$  of height  $n$ , and let  $w_J : F/JF \rightarrow J/J^2$  be a surjection. Suppose  $w_J$  can be lifted to a surjection  $\alpha : F \rightarrow J$ . Let  $\bar{a} \in A/J$  be a unit, and let  $\theta$  be an automorphism of  $F/JF$  with determinant  $\bar{a}^2$ . Then the surjection  $w_J \circ \theta : F/JF \rightarrow J/J^2$  can be lifted to a surjection  $\gamma : F \rightarrow J$ .*

**Lemma 4.5.** *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A^2$  be a projective  $A$ -module of rank  $n$ . Let  $J$  be an ideal of  $A$  of height  $n$ , and let  $w_J$  be a local  $F$ -orientation of  $J$ . Let  $\bar{a} \in A/J$  be a unit. Then  $(J, w_J) = (J, \overline{a^2}w_J)$  in  $E(A, F)$ .*

**Lemma 4.6.** *Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . Let  $J$  be an ideal of  $A$  of height  $n$ , and let  $w_J$  be a local  $F$ -orientation of  $J$ . Suppose  $(J, w_J) \neq 0$  in  $E(A, F)$ . Then there exists an ideal  $J_1$  of height  $n$  which is comaximal with  $J$  and a local  $F$ -orientation  $w_{J_1}$  of  $J_1$  such that  $(J, w_J) + (J_1, w_{J_1}) = 0$  in  $E(A, F)$ . Further, given any ideal  $K$  of  $A$  of height  $\geq 1$ ,  $J_1$  can be chosen to be comaximal with  $K$ .*

The following result is similar to [3, Lemma 5.6].

**Lemma 4.7.** *Let  $A$  be an affine domain of dimension  $n \geq 2$  over a field  $k$ , and let  $f$  be a non-zero element of  $A$ . Let  $F = Q \oplus A^2$  be a projective  $A$ -module of rank  $n$ , and let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements. Suppose that  $(J, w_J) \neq 0$  in  $E(A, F)$ , but the image of  $(J, w_J)$  is zero in  $E(A_f, F_f)$ .*



Then there exists an ideal  $J_2$  of  $A$  of height  $n$  such that  $(J_2)_f = A_f$  and  $(J, w_J) = (J_2, w_{J_2})$  in  $E(A, F)$ .

*Proof.* Since  $(J, w_J) \neq 0$  in  $E(A, F)$ , but its image is zero in  $E(A_f, F_f)$ , we see that  $f$  is not a unit in  $A$ . By (4.6), we can choose an ideal  $J_1$  of height  $n$  which is comaximal with  $Jf$  such that  $(J, w_J) + (J_1, w_{J_1}) = 0$  in  $E(A, F)$ . Since the image of  $(J, w_J)$  is zero in  $E(A_f, F_f)$ , it follows that the image of  $(J_1, w_{J_1})$  is zero in  $E(A_f, F_f)$ .

By (3.2), there exists a surjection  $\alpha : F_f \twoheadrightarrow (J_1)_f$  such that  $\alpha \otimes A_f / (J_1)_f = (w_{J_1})_f$ . Choose a positive integer  $k$  such that  $f^{2k}\alpha : F \rightarrow J_1$ . Since  $f$  is a unit modulo  $J_1$ , by (4.5),  $(J_1, w_{J_1}) = (J_1, f^{2kn}w_{J_1})$  in  $E(A, F)$ . By (2.7), there exists a  $\Psi \in \text{EL}(F_f^*)$  such that  $\beta = \Psi(\alpha) \in F^*$  and  $\beta(F) \subset J_1$  is an ideal of height  $n$ . Thus,  $\beta(F) = J_1 \cap J_2$ , where  $J_2$  is an ideal of  $A$  of height  $n$  such that  $(J_2)_f = A_f$ . Hence,  $J_1 + J_2 = A$ . From the surjection  $\beta$ , we get  $(J_1, w_{J_1}) + (J_2, w_{J_2}) = 0$  in  $E(A, F)$ . Since  $(J, w_J) + (J_1, w_{J_1}) = 0$  in  $E(A, F)$ , it follows that  $(J, w_J) = (J_2, w_{J_2})$  in  $E(A, F)$ . This proves the result.  $\square$

Using (3.3), (4.5) and (4.7), and following the proof of [3, Lemma 5.8], the following result can be proved.

**Lemma 4.8.** *Let  $A$  be an affine domain of dimension  $n \geq 2$  over a field  $k$ . Let  $P$  and  $F = Q \oplus A^2$  be projective  $A$ -modules of rank  $n$  with  $\wedge^n P \xrightarrow{\sim} \wedge^n F$ . Let  $f$  be a non-zero element of  $A$ . Assume that every generic surjection ideal of  $P$  is a surjective image of  $F$ . Then every generic surjection ideal of  $P_f$  is a surjective image of  $F_f$ .*

Using the above results and following the proof of [3, Theorem 5.9], the next result follows.

**Theorem 4.9.** *Let  $A$  be an affine domain of dimension  $n \geq 2$  over a real closed field  $k$ . Let  $P$  and  $F = Q \oplus A^2$  be projective  $A$ -modules of rank  $n$  with  $\wedge^n P \xrightarrow{\sim} \wedge^n F$ . Assume that every generic surjection ideal of  $P$  is a surjective image of  $F$ . Then  $P$  has a unimodular element.*

*In particular, if  $L = \wedge^n P$  and every generic surjection ideal of  $P$  is a surjective image of  $L \oplus A^{n-1}$ , then  $P$  has a unimodular element.*

**5. Weak Euler class group.** Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $F = Q \oplus A$  be a projective  $A$ -module of rank  $n$ . We define the weak Euler class group  $E_0(A, F)$  of  $A$  with respect to  $F$  as follows:

Let  $\mathcal{S}$  be the set of ideals  $\mathcal{N}$  of  $A$  such that  $\mathcal{N}/\mathcal{N}^2$  is generated by  $n$  elements, where  $\mathcal{N}$  is an  $\mathcal{M}$ -primary ideal for some maximal ideal  $\mathcal{M}$  of  $A$  of height  $n$ . Let  $G$  be the free abelian group on the set  $\mathcal{S}$ .

Let  $J = \cap \mathcal{N}_i$  be the intersection of finitely many ideals  $\mathcal{N}_i$ , where  $\mathcal{N}_i$  is an  $\mathcal{M}_i$ -primary and the  $\mathcal{M}_i$ 's are distinct maximal ideals of  $A$  of height  $n$ . Assume that  $J/J^2$  is generated by  $n$  elements. We associate to  $J$  the element  $\sum \mathcal{N}_i$  of  $G$ . We denote this element by  $(J)$ .

Let  $H$  be the subgroup of  $G$  generated by elements of the type  $(J)$ , where  $J$  is an ideal of  $A$  of height  $n$  which is a surjective image of  $F$ .

We set  $E_0(A, F) = G/H$ .

Let  $P$  be a projective  $A$ -module of rank  $n$  such that  $\wedge^n P \xrightarrow{\sim} \wedge^n F$ . Let  $\lambda : P \rightarrow J_0$  be a generic surjection of  $P$ . We define  $e(P) = (J_0)$  in  $E_0(A, F)$ . We will show that this assignment is well defined.

Let  $\mu : P \rightarrow J_1$  be another generic surjection of  $P$ . By (2.2), there exists a surjection  $\alpha(T) : P[T] \rightarrow I$ , where  $I$  is an ideal of  $A[T]$  of height  $n$  with  $\alpha(0) = \lambda$ ,  $I(0) = J_0$ ,  $\alpha(1) = \mu$  and  $I(1) = J_1$ . Now, as before, using (2.3), we see that  $(J_0) = (J_1)$  in  $E_0(A, F)$ . This shows that  $e(P)$  is well defined.

Note that there is a canonical surjection from  $E(A, F)$  to  $E_0(A, F)$  obtained by forgetting the orientations.

We state the following result which follows from (4.3) and (4.5).

**Lemma 5.1.** *Let  $A$  be a ring of even dimension  $n$ . Let  $P$  and  $F = Q \oplus A^2$  be projective  $A$ -modules of rank  $n$ , and let  $\chi : \wedge^n F \xrightarrow{\sim} \wedge^n P$  be an  $F$ -orientation of  $P$ . Let  $e(P, \chi) = (J, w_J)$  in  $E(A, F)$ , and let  $\tilde{w}_J$  be another local  $F$ -orientation of  $J$ . Then there exists a projective  $A$ -module  $P_1$  with  $[P_1] = [P]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_1$  of  $P_1$  such that  $e(P_1, \chi_1) = (J, \tilde{w}_J)$  in  $E(A, F)$ .*

**Proposition 5.2.** *Let  $A$  be a ring of even dimension  $n$ , and let  $F = Q \oplus A^2$  be a projective  $A$ -module of rank  $n$ . Let  $J_1$  and  $J_2$  be two comaximal ideals of  $A$  of height  $n$ , and let  $J_3 = J_1 \cap J_2$ . If any two of*

$J_1, J_2$  and  $J_3$  are surjective images of projective  $A$ -modules of rank  $n$  which are stably isomorphic to  $F$ , then so is the third one.

*Proof.* (i) Let  $P_1$  and  $P_2$  be two projective  $A$ -modules of rank  $n$  with  $[P_1] = [P_2] = [F]$  in  $K_0(A)$ , and let  $\psi_1 : P_1 \twoheadrightarrow J_1$  and  $\psi_2 : P_2 \twoheadrightarrow J_2$  be two surjections. Choose  $F$ -orientations  $\chi_1$  and  $\chi_2$  of  $P_1$  and  $P_2$ , respectively. Then  $e(P_1, \chi_1) = (J_1, w_{J_1})$  and  $e(P_2, \chi_2) = (J_2, w_{J_2})$  in  $E(A, F)$ .

By (2.6), there exists an ideal  $J'_1$  of height  $n$  which is a surjective image of both  $P_1$  and  $F$ . Hence,  $e(P_1, \chi_1) = (J_1, w_{J_1}) = (J'_1, w_{J'_1})$  in  $E(A, F)$  for some local  $F$ -orientation  $w_{J'_1}$  of  $J'_1$ . Similarly, there exists an ideal  $J'_2$  of height  $n$  which is a surjective image of both  $P_2$  and  $F$ . Hence,  $e(P_2, \chi_2) = (J_2, w_{J_2}) = (J'_2, w_{J'_2})$  in  $E(A, F)$  for some local  $F$ -orientation  $w_{J'_2}$  of  $J'_2$ . Further, we may assume that  $J'_1 + J'_2 = A$ . Let  $(J_1, w_{J_1}) + (J_2, w_{J_2}) = (J_3, w_{J_3})$  in  $E(A, F)$ .

Let  $J'_3 = J'_1 \cap J'_2$ . By the addition principle (2.4),  $J'_3$  is a surjective image of  $F$  and  $(J'_1, w_{J'_1}) + (J'_2, w_{J'_2}) = (J'_3, w_{J'_3})$  in  $E(A, F)$ . Hence,  $(J'_3, w_{J'_3}) = (J_3, w_{J_3})$ . Since  $J'_3$  is a surjective image of  $F$ , by (5.1), there exists a projective  $A$ -module  $P_3$  with  $[P_3] = [F]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_3$  of  $P_3$  such that  $e(P_3, \chi_3) = (J'_3, w_{J'_3}) = (J_3, w_{J_3})$  in  $E(A, F)$ . By (3.3), there exists a surjection  $\psi_3 : P_3 \twoheadrightarrow J_3$  such that  $(\psi_3, \chi_3)$  induces  $(J_3, w_{J_3})$ . This proves the first part.

(ii) Now assume that  $J_1$  and  $J_3$  are surjective images of  $P'_1$  and  $P_3$ , respectively, where  $P'_1$  and  $P_3$  are projective  $A$ -modules of rank  $n$  with  $[P'_1] = [P_3] = [F]$  in  $K_0(A)$ .

Let  $e(P_3, \chi_3) = (J_3, w_3)$  for some  $F$ -orientation  $\chi_3$  of  $P_3$ , and let  $(J_3, w_3) = (J_1, w_1) + (J_2, w_2)$  in  $E(A, F)$ . Let  $e(P'_1, \chi'_1) = (J_1, w'_1)$  for some  $F$ -orientation  $\chi'_1$  of  $P'_1$ . By (5.1), there exists a projective  $A$ -module  $P_1$  of rank  $n$  with  $[P_1] = [P'_1]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_1$  of  $P_1$  such that  $e(P_1, \chi_1) = (J_1, w_1)$  in  $E(A, F)$ .

By (2.6), there exists an ideal  $J_4$  of height  $n$  which is a surjective image of  $F$  and  $P_1$ , both, and is comaximal with  $J_2$  such that  $e(P_1, \chi_1) = (J_1, w_1) = (J_4, w_4)$ . Write  $J_5 = J_4 \cap J_2$ . Assume that  $(J_4, w_4) + (J_2, w_2) = (J_5, w_5)$  in  $E(A, F)$ . Then we have  $e(P_3, \chi_3) = (J_3, w_3) = (J_5, w_5)$  in  $E(A, F)$ .

Since  $J_4$  is a surjective image of  $F$ , we get  $e(F, \chi) = (J_4, \tilde{w}_4) = 0$  for some  $\chi$ . If  $(J_4, \tilde{w}_4) + (J_2, w_2) = (J_5, \tilde{w}_5)$ , then  $(J_2, w_2) = (J_5, \tilde{w}_5)$ .

Since  $e(P_3, \chi_3) = (J_5, w_5)$ , by (5.1), there exists a projective  $A$ -module  $\tilde{P}_3$  of rank  $n$  with  $[\tilde{P}_3] = [P_3]$  in  $K_0(A)$  such that  $e(\tilde{P}_3, \tilde{w}_3) = (J_5, \tilde{w}_5) = (J_2, w_2)$ . Hence, by (3.3),  $J_2$  is a surjective image of  $\tilde{P}_3$  which is stably isomorphic to  $F$ . This completes the proof.  $\square$

**Proposition 5.3.** *Let  $A$  be a ring of even dimension  $n$ , and let  $F = Q \oplus A^2$  be a projective  $A$ -module of rank  $n$ . Let  $J$  be an ideal of  $A$  of height  $n$ . Then  $(J) = 0$  in  $E_0(A, F)$  if and only if  $J$  is a surjective image of a projective  $A$ -module of rank  $n$  which is stably isomorphic to  $F$ .*

*Proof.* Let  $J_1$  be an ideal of  $A$  of height  $n$ . Assume that  $J_1$  is surjective image of a projective  $A$ -module of rank  $n$  which is stably isomorphic to  $F$ . Assume  $(J_1, w_{J_1})$  is a non-zero element of  $E(A, F)$ . We will show that there exist height  $n$  ideals  $J_2$  and  $J_3$  with local  $F$ -orientations  $w_{J_2}$  and  $w_{J_3}$  respectively such that:

- (i)  $J_2, J_3$  are comaximal with any given ideal of height  $\geq 1$ ,
- (ii)  $(J_1, w_{J_1}) = -(J_2, w_{J_2}) = (J_3, w_{J_3})$  in  $E(A, F)$  and
- (iii)  $J_2, J_3$  are surjective images of projective  $A$ -modules of rank  $n$  which are stably isomorphic to  $F$ .

By (4.6), there exists an ideal  $J_2$  of height  $n$  which is comaximal with  $J_1$  and any given ideal of height  $\geq 1$  such that  $(J_1, w_{J_1}) + (J_2, w_{J_2}) = 0$  in  $E(A, F)$ . By (3.2),  $J_1 \cap J_2$  is a surjective image of  $F$ . By (5.2),  $J_2$  is a surjective image of a projective  $A$ -module of rank  $n$  which is stably isomorphic to  $F$ .

Repeating the above with  $(J_2, w_{J_2})$ , we get an ideal  $J_3$  of height  $n$  which is comaximal with any given ideal of height  $\geq 1$  such that  $(J_2, w_{J_2}) + (J_3, w_{J_3}) = 0$  in  $E(A, F)$ . Further,  $J_3$  is a surjective image of a projective  $A$ -module of rank  $n$  which is stably isomorphic to  $F$ . Thus, we have  $(J_1, w_{J_1}) = -(J_2, w_{J_2}) = (J_3, w_{J_3})$  in  $E(A, F)$ . This proves the above claim.

From the above discussion, we see that, given any element  $h$  in kernel of the canonical map  $\Phi : E(A, F) \rightarrow E_0(A, F)$ , there exists an ideal  $\tilde{J}$  of height  $n$  such that  $\tilde{J}$  is a surjective image of a projective  $A$ -module of rank  $n$  which is stably isomorphic to  $F$  and  $h = (\tilde{J}, w_{\tilde{J}})$  in  $E(A, F)$ . Moreover,  $\tilde{J}$  can be chosen to be comaximal with any ideal of height  $\geq 1$ .

Now assume  $(J) = 0$  in  $E_0(A, F)$ . Choose some local  $F$ -orientation  $w_J$  of  $J$ . Then  $(J, w_J) \in \ker(\Phi)$ . From the previous paragraph, we get that there exists an ideal  $K$  of height  $n$  comaximal with  $J$  such that  $-(J, w_J) = (K, w_K)$  in  $E(A, F)$ . Further,  $K$  is a surjective image of a projective  $A$ -module which is stably isomorphic to  $F$ .

Since  $(J, w_J) + (K, w_K) = 0$  in  $E(A, F)$ , by (3.2),  $J \cap K$  is surjective image of  $F$ . By (5.2),  $J$  is a surjective image of a projective  $A$ -module of rank  $n$  which is stably isomorphic to  $F$ .

Conversely, assume that  $J$  is a surjective image of a projective  $A$ -module  $P$  of rank  $n$  which is stably isomorphic to  $F$ . Let  $\chi$  be a  $F$ -orientation of  $P$ . Then  $e(P, \chi) = (J, w_J)$  in  $E(A, F)$ . By (2.6), there exists an ideal  $I$  of height  $n$  which is a surjective image of both  $P$  and  $F$ . Then  $e(P, \chi) = (J, w_J) = (I, w_I)$  in  $E(A, F)$ . Therefore,  $(J) = (I)$  in  $E_0(A, F)$ , and hence  $(J) = 0$  in  $E_0(A, F)$ . This completes the proof.  $\square$

**Proposition 5.4.** *Let  $A$  be a ring of even dimension  $n$ , and let  $F = Q \oplus A^2$  and  $P$  be projective  $A$ -modules of rank  $n$  with  $\wedge^n P \xrightarrow{\sim} \wedge^n F$ . Then  $e(P) = 0$  in  $E_0(A, F)$  if and only if  $[P] = [P_1 \oplus A]$  in  $K_0(A)$  for some projective  $A$ -module  $P_1$  of rank  $n - 1$ .*

*Proof.* Assume that  $[P] = [P_1 \oplus A]$  in  $K_0(A)$ . By (2.6), there exists an ideal  $J$  of  $A$  of height  $n$  which is a surjective image of both  $P$  and  $P_1 \oplus A$ . Hence,  $e(P_1 \oplus A, \chi) = (J, w_J) = 0$  in  $E(A, F)$ , by (3.4). Hence,  $J$  is a surjective image of  $F$ . By (5.3),  $e(P) = (J) = 0$  in  $E_0(A, F)$ .

Conversely, assume that  $e(P) = 0$  in  $E_0(A, F)$ . Let  $\psi : P \twoheadrightarrow J$  be a generic surjection of  $P$ , and let  $e(P, \chi) = (J, w_J)$  in  $E(A, F)$  for some  $F$ -orientation  $\chi$  of  $P$ . Since  $e(P) = (J) = 0$  in  $E_0(A, F)$ , by (5.3),  $J$  is a surjective image of a projective  $A$ -module  $P_1$  with  $[P_1] = [F]$  in  $K_0(A)$ . By (2.6), there exists a height  $n$  ideal  $J_1$  which is a surjective image of both  $P_1$  and  $F$ . Let  $e(P_1, \chi_1) = (J, \tilde{w}_J) = (J_1, w_{J_1})$  for some  $F$ -orientation  $\chi_1$  of  $P_1$ .

By (5.1), there exists a rank  $n$  projective  $A$ -module  $P_2$  with  $[P_2] = [P]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_2$  of  $P_2$  such that  $e(P_2, \chi_2) = (J, \tilde{w}_J) = (J_1, w_{J_1})$  in  $E(A, F)$ . Since  $J_1$  is a surjective image of  $F$ ,  $(J_1, \tilde{w}_{J_1}) = 0$  in  $E(A, F)$  for some local  $F$ -orientation  $\tilde{w}_{J_1}$  of  $J_1$ . By (5.1), there exists a projective  $A$ -module  $P_3$  with  $[P_3] = [P_2]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_3$  of  $P_3$  such that  $e(P_3, \chi_3) = (J_1, \tilde{w}_{J_1}) = 0$  in  $E(A, F)$ . Hence,  $P_3 = P_4 \oplus A$ , by (3.4). Therefore,  $[P] = [P_2] = [P_4 \oplus A]$  in  $K_0(A)$ . This completes the proof.  $\square$

**Proposition 5.5.** *Let  $A$  be a ring of even dimension  $n$ . Let  $P$  and  $F = Q \oplus A^2$  be projective  $A$ -modules of rank  $n$  with  $\wedge^n P \xrightarrow{\sim} \wedge^n F$ . Suppose that  $e(P) = (J)$  in  $E_0(A, F)$ , where  $J$  is an ideal of  $A$  of height  $n$ . Then there exists a projective  $A$ -module  $P_1$  of rank  $n$  such that  $[P] = [P_1]$  in  $K_0(A)$  and  $J$  is a surjective image of  $P_1$ .*

*Proof.* Since  $P/J P$  is free and  $J/J^2$  is generated by  $n$  elements, we get a surjection  $\bar{\psi} : P/J P \twoheadrightarrow J/J^2$ . By [3, Corollary 2.14], we can lift  $\bar{\psi}$  to a surjection  $\psi : P \twoheadrightarrow J \cap J_1$ , where  $J_1$  is a height  $n$  ideal comaximal with  $J$ . Let  $e(P, \chi) = (J, w_J) + (J_1, w_{J_1})$  in  $E(A, F)$  for some  $F$ -orientation  $\chi$  of  $P$ .

Since  $e(P) = (J) = (J \cap J_1)$  in  $E_0(A, F)$ ,  $(J_1) = 0$  in  $E_0(A, F)$ . By (5.3),  $J_1$  is a surjective image of a projective  $A$ -module  $P_2$  of rank  $n$  which is stably isomorphic to  $F$ . By (5.1), there exists a rank  $n$  projective  $A$ -module  $P_3$  with  $[P_2] = [P_3]$  in  $K_0(A)$  and an  $F$ -orientation  $\chi_3$  of  $P_3$  such that  $e(P_3, \chi_3) = (J_1, w_{J_1})$  in  $E(A, F)$ .

By (2.6), there exists an ideal  $J_2$  of height  $n$  which is comaximal with  $J$  and is a surjective image of both  $F$  and  $P_3$ . Assume that  $e(P_3, \chi_3) = (J_1, w_{J_1}) = (J_2, w_{J_2})$  in  $E(A, F)$ . Hence,  $e(P, \chi) = (J, w_J) + (J_2, w_{J_2}) = (J \cap J_2, w_{J \cap J_2})$ . By (3.3), there exists a surjection  $\phi : P \twoheadrightarrow J \cap J_2$ . Since  $J_2$  is a surjective image of  $F$ , we get  $(J_2, \tilde{w}_{J_2}) = 0$  for some local  $F$ -orientation  $\tilde{w}_{J_2}$  of  $J_2$ . Let  $(J, w_J) + (J_2, \tilde{w}_{J_2}) = (J \cap J_2, \tilde{w}_{J \cap J_2})$ . By (4.3), there exists rank  $n$  projective  $A$ -module  $P_1$  with  $[P] = [P_1]$  in  $K_0(A)$  and  $e(P_1, \chi_1) = (J \cap J_2, \tilde{w}_{J \cap J_2}) = (J, w_J)$  in  $E(A, F)$  for some  $F$ -orientation  $\chi_1$  of  $P_1$ . By (3.3), there exists a surjection  $\alpha : P_1 \twoheadrightarrow J$ . This proves the result.  $\square$

The proof of the following result is similar to [3, Proposition 6.5]; hence, we omit it.

**Proposition 5.6.** *Let  $A$  be a ring of even dimension  $n$ , and let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements. Let  $F = Q \oplus A^2$  be a projective  $A$ -module of rank  $n$ , and let  $\tilde{w}_J : F/J F \twoheadrightarrow J/J^2$  be a surjection. Suppose that the element  $(J, \tilde{w}_J)$  of  $E(A, F)$  belongs to the kernel of the canonical homomorphism  $E(A, F) \twoheadrightarrow E_0(A, F)$ . Then there exists a projective  $A$ -module  $P_1$  of rank  $n$  such that  $[P_1] = [F]$  in  $K_0(A)$  and  $e(P_1, \chi_1) = (J, \tilde{w}_J)$  in  $E(A, F)$  for some  $F$ -orientation  $\chi_1$  of  $P_1$ .*

**6. Application.** Let  $A$  be a ring of dimension  $n \geq 2$ , and let  $L$  be a projective  $A$ -module of rank 1. Let  $F = Q \oplus A$  be a projective

$A$ -module of rank  $n$  with determinant  $L$ . The group  $E(A, L)$  defined by Bhatwadekar and Sridharan [3] is the same as  $E(A, L \oplus A^{n-1})$ . We will define a map  $\Delta : E(A, L) \rightarrow E(A, F)$ .

Let  $w_J : L/JL \oplus (A/J)^{n-1} \twoheadrightarrow J/J^2$  be a surjection. Since  $\dim A/J = 0$ ,  $Q/JQ$  is isomorphic to  $L/JL \oplus (A/J)^{n-2}$ . Choose an isomorphism  $\theta : Q/JQ \xrightarrow{\sim} L/JL \oplus (A/J)^{n-2}$  of determinant one. Let  $\tilde{w}_J = w_J \circ (\theta, id) : Q/JQ \oplus A/J \twoheadrightarrow J/J^2$  be a surjection.

Assume that  $w_J$  can be lifted to a surjection  $\Phi : L \oplus A^{n-1} \twoheadrightarrow J$ . Write  $\Phi = (\Phi_1, a)$ . We may assume that  $\Phi_1(L \oplus A^{n-2}) = K$  is an ideal of height  $n - 1$ . Further, we may assume that the isomorphism  $\theta : Q/JQ \xrightarrow{\sim} L/JL \oplus (A/J)^{n-2}$  is induced from an isomorphism  $\theta' : Q/KQ \xrightarrow{\sim} L/KL \oplus (A/K)^{n-2}$  (i.e.,  $\theta' \otimes A/J = \theta$ ).

Let  $(\Phi_2, a) : Q \oplus A \rightarrow J = (K, a)$  be a lift of  $\tilde{w}_J$ . Then  $\Phi_2 \otimes A/K : Q/KQ \twoheadrightarrow K/K^2$  is a surjection. Let  $\phi_2 : Q \rightarrow K$  be a lift of  $\Phi_2 \otimes A/K$ . Then  $\phi_2(Q) + K^2 = K$ . Hence, there exists an  $e \in K^2$  with  $e(1 - e) \in \phi_2(Q)$  such that  $\phi_2(Q) + Ae = K$ . Now it is easy to check that  $\phi_2(Q) + Aa = \phi_2(Q) + (e + (1 - e)a)A = K + Aa = J$  and  $(\phi_2, e + (1 - e)a) : Q \oplus A \twoheadrightarrow J$  is a lift of  $\tilde{w}_J$ .

Hence, we have shown that, if  $w_J$  can be lifted to a surjection from  $L \oplus A^{n-1} \twoheadrightarrow J$ , then  $\tilde{w}_J$  can be lifted to a surjection from  $Q \oplus A$  to  $J$ . Further, if we choose a different isomorphism  $\theta_1 : Q/JQ \oplus A/J \xrightarrow{\sim} L/JL \oplus (A/J)^{n-1}$  of determinant one and  $w_1 = w_J \circ \theta_1 : Q/JQ \oplus A/J \twoheadrightarrow J/J^2$ , then  $\tilde{w}_J$  and  $w_1$  are connected by an element of  $EL(Q/JQ \oplus A/J)$ . Hence, if we define  $\Delta : E(A, L) \rightarrow E(A, F)$  by  $\Delta(w_J) = \tilde{w}_J$ , then this map is well defined. It is easy to see that  $\Delta$  is a group homomorphism.

Similarly, we can define a map  $\Delta_1 : E(A, F) \rightarrow E(A, L)$ , and it is easy to show that  $\Delta \circ \Delta_1 = id$  and  $\Delta_1 \circ \Delta = id$ . Hence, we get the following interesting result:

**Theorem 6.1.** *Let  $A$  be a ring of dimension  $n \geq 2$ . Let  $L$  and  $F = Q \oplus A$  be projective  $A$ -modules of ranks 1 and  $n$ , respectively, with  $\wedge^n F \xrightarrow{\sim} L$ . Then  $E(A, L)$  is isomorphic to  $E(A, F)$ .*

Let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements. Further, assume that there exists a surjection  $\alpha : L \oplus A^{n-1} \twoheadrightarrow J$ . We will show that  $J$  is also a surjective image of  $F = Q \oplus A$ . Let  $w_J$  be the local  $L$ -orientation of  $J$  induced from

$\alpha$ . Then  $(J, w_J) = 0$  in  $E(A, L)$ . Hence,  $\Delta(J, w_J) = (J, \tilde{w}_J) = 0$  in  $E(A, F)$ . Hence, by (3.2),  $J$  is a surjective image of  $F$ .  $\square$

We define a map  $\tilde{\Delta} : E_0(A, L) \rightarrow E_0(A, F)$  by  $(J) \mapsto (J)$ . The above discussion shows that  $\tilde{\Delta}$  is well defined. Similarly, we can define a map  $\tilde{\Delta}_1 : E_0(A, F) \rightarrow E_0(A, L)$  such that  $\tilde{\Delta} \circ \tilde{\Delta}_1 = id$  and  $\tilde{\Delta}_1 \circ \tilde{\Delta} = id$ . Thus we get the following interesting result:

**Theorem 6.2.** *Let  $A$  be a ring of dimension  $n \geq 2$ . Let  $L$  and  $F = Q \oplus A$  be projective  $A$ -modules of ranks 1 and  $n$ , respectively, with  $\wedge^n F \xrightarrow{\sim} L$ . Then  $E_0(A, L)$  is isomorphic to  $E_0(A, F)$ .*

Since, by [3, 6.8],  $E_0(A, L)$  is canonically isomorphic to  $E_0(A, A)$ , we get the surprising result that  $E_0(A, F)$  is canonically isomorphic to  $E_0(A, A^n)$  for any projective  $A$ -module  $F = Q \oplus A$  of rank  $n$ .

We end with the following result which follows from (5.3).

**Proposition 6.3.** *Let  $A$  be a ring of even dimension  $n$ , and let  $J$  be an ideal of  $A$  of height  $n$  such that  $J/J^2$  is generated by  $n$  elements. Let  $L$  and  $P$  be projective  $A$ -modules of ranks 1 and  $n$ , respectively, such that  $P$  is stably isomorphic to  $L \oplus A^{n-1}$ . Then  $J$  is surjective image of  $P$  if and only if, given any projective  $A$ -module  $Q$  of rank  $n-2$  with determinant  $L$ , there exists a projective  $A$ -module  $P_1$  which is stably isomorphic to  $Q \oplus A^2$  such that  $J$  is surjective image of  $P_1$ .*

## REFERENCES

1. S.M. Bhatwadekar and M.K. Keshari, *A question of Nori: Projective generation of ideals*, K-Theory **28** (2003), 329–351.
2. S.M. Bhatwadekar and A. Roy, *Some theorems about projective modules over polynomial rings*, J. Algebra **86** (1984), 150–158.
3. S.M. Bhatwadekar and Raja Sridharan, *The Euler class group of a Noetherian ring*, Compos. Math. **122** (2000), 183–222.

DEPARTMENT OF MATHEMATICS, IIT MUMBAI, MUMBAI - 400076, INDIA  
**Email address:** keshari@math.iitb.ac.in

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF KANSAS, 1460 JAYHAWK BLVD,  
 LAWRENCE, KS 66045  
**Email address:** mandal@math.ku.edu