

Rank Gradients of Infinite Cyclic Covers of 3-Manifolds

JASON DEBLOIS, STEFAN FRIEDL, & STEFANO VIDUSSI

ABSTRACT. Given a 3-manifold M with no spherical boundary components, and a primitive class $\phi \in H^1(M; \mathbb{Z})$, we show that the following are equivalent:

- (1) ϕ is a fibered class,
- (2) the rank gradient of (M, ϕ) is zero,
- (3) the Heegaard gradient of (M, ϕ) is zero.

1. Introduction

A *directed 3-manifold* is a pair (M, ϕ) where M is a compact, orientable, connected 3-manifold with toroidal or empty boundary, and $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$ is a primitive class, that is, ϕ viewed as a homomorphism $\pi_1(M) \rightarrow \mathbb{Z}$ is an epimorphism. We say that a directed 3-manifold (M, ϕ) *fibers over* S^1 if there exists a fibration $p : M \rightarrow S^1$ such that the induced map $p_* : \pi_1(M) \rightarrow \pi_1(S^1) = \mathbb{Z}$ coincides with ϕ . We refer to such ϕ as a *fibered class*.

It is well known that the pair $(\pi_1(M), \phi : \pi_1(M) \rightarrow \mathbb{Z})$ determines whether ϕ is fibered or not. Indeed, it follows from Stallings’ theorem [St62] (together with the resolution of the Poincaré conjecture) that ϕ is a fibered class if and only if $\text{Ker}(\phi : \pi_1(M) \rightarrow \mathbb{Z})$ is finitely generated.

Stallings’ theorem can be generalized in various directions (see e.g. [FV12, Theorem 5.2], [SW09a; SW09b], and [FSW13]). Our main result gives a new fibering criterion, which is also a strengthening of Stallings’ theorem. In order to state our result, we need the notion of rank gradient, which was first introduced by Lackenby [La05]. Given a finitely generated group π , we denote by $\text{rk}(\pi)$ the *rank* of π , that is, the minimal number of generators of π . If (M, ϕ) is a directed 3-manifold, then we write

$$\pi_n = \text{Ker}(\pi_1(M) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}_n),$$

and we refer to

$$\text{rg}(M, \phi) := \liminf_{n \rightarrow \infty} \frac{1}{n} \text{rk}(\pi_n)$$

as the *rank gradient* of (M, ϕ) . (In the notation of [La05] this is the rank gradient of $(\pi_1 M, \{\pi_n\})$.)

Received January 8, 2013. Revision received January 10, 2014.

J. DeBlois was partially supported by NSF grant DMS-1240329. S. Vidussi was partially supported by NSF grant DMS-0906281.

If ϕ is a fibered class, then ϕ is dual to a fiber S of a fibration $M \rightarrow S^1$ (a connected surface), and it is straightforward to show that $\text{rk}(\pi_n) \leq 1 + \text{genus}(S)$ for any n (see e.g. Lemma 2.3). In particular, $\text{rg}(M, \phi) = 0$.

Our main result now says that the converse to this statement holds. More precisely, we will prove the following theorem.

THEOREM 1.1. *Let (M, ϕ) be a directed 3-manifold. Then the following three statements are equivalent:*

- (1) *the class ϕ is fibered,*
- (2) *the sequence $\text{rk}(\pi_n)$, $n \in \mathbb{N}$, is bounded,*
- (3) *$\text{rg}(M, \phi) = 0$.*

It follows from the discussion preceding the theorem that it suffices to prove that (3) implies (1). In fact, we will present two quite different approaches to the proof of this statement.

The first, discussed in Section 3, uses tools from geometric group theory: acylindrical accessibility and the finite height property. It applies only to closed hyperbolic manifolds but has the advantage of generalizing more readily to the broader setting of hyperbolic groups, where the separability results used for the general case are not currently available. Moreover, with more work, this approach yields explicit lower bounds on the rank gradient. In the sequel the first author [De13] refines Theorem 1.1 in this way for M closed and hyperbolic, bounding $\text{rg}(M, \phi)$ below in terms of the Thurston norm of a nonfibered class ϕ .

The second proof, discussed in Section 4, uses the recent proof (see [FV12]) that given any nonfibered directed 3-manifold (M, ϕ) , there exists a twisted Alexander polynomial that vanishes. This proof in turn relies on the recent results of Wise [Wi09; Wi12a; Wi12b].

To describe our second result, we need to introduce the notion of Heegard gradient. A *Heegaard surface* for a compact 3-manifold M is an embedded separating surface $S \subset M$ such that the two components of M cut along S are compression bodies. The minimal genus of a Heegaard surface is called the *Heegaard genus* $h(M)$ of M . Given a class $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$, we can then define the *Heegaard gradient* $\text{hg}(M, \phi)$ in a similar fashion to the rank gradient. We refer to Section 2.2 for more details. In that section we will also see that the subsequent theorem is a straightforward consequence of Theorem 1.1.

THEOREM 1.2. *Let (M, ϕ) be a directed 3-manifold. Then the following three statements are equivalent:*

- (1) *ϕ is fibered,*
- (2) *the sequence $h(M_n)$, $n \in \mathbb{N}$, is bounded,*
- (3) *$\text{hg}(M, \phi) = 0$.*

This theorem was proved by Lackenby [La06, Theorem 1.11] for closed hyperbolic 3-manifolds. To the best of our knowledge, the general case has not been proved before. The equivalence presented in the abstract immediately follows

from Theorems 1.1 and 1.2 if M has empty or toroidal boundary. In the general case see Lemma 4.5.

The equivalence between vanishing of rank and Heegaard gradients holds with no restriction on boundaries. This is proved at the end of Section 4.

THEOREM 1.3. *For a compact, orientable, connected 3-manifold M and a primitive class $\phi \in H^1(M; \mathbb{Z})$, $\text{rg}(M, \phi) = 0$ if and only if $\text{hg}(M, \phi) = 0$.*

We will now formulate the last theorem of the paper. Recall that a group π is normally generated by a subset $S \subset \pi$ if π is the smallest normal subgroup of π that contains S . We define the *normal rank* $n(\pi)$ of π to be the smallest cardinality of a normal generating set of π . The first part of this theorem can also be viewed as a strengthening of Stallings' fibering theorem.

THEOREM 1.4. (1) *If (M, ϕ) is a nonfibered directed 3-manifold, then $\text{Ker}(\phi)$ admits a finite index subgroup with infinite normal rank.*

(2) *There exists a nonfibered directed 3-manifold (M, ϕ) such that $\text{Ker}(\phi)$ has finite normal rank.*

CONVENTION. Unless it says specifically otherwise, all groups are assumed to be finitely generated, all manifolds are assumed to be orientable, connected, and compact, and all 3-manifolds are assumed to have empty or toroidal boundary.

2. The Rank Gradient and the Heegaard Gradient

2.1. The Rank Gradient

We start out with the following well-known lemma.

LEMMA 2.1. *Let G be a finitely generated group, and let $H \subset G$ be a finite index subgroup. Then*

$$\text{rk}(H) \leq [G : H] \cdot (\text{rk}(G) - 1) + 1 \leq [G : H] \cdot \text{rk}(G). \quad (1)$$

Proof. Let $\alpha : F \rightarrow G$ be an epimorphism where F is a free group of rank $\text{rk}(G)$. Note that $\alpha^{-1}(H)$ is a subgroup of F of index $d := [G : H]$. It follows from elementary properties of the free group that $\alpha^{-1}(H)$ is a free group of rank

$$d \cdot (\text{rk}(F) - 1) = [G : H] \cdot (\text{rk}(G) - 1).$$

Since α restricts to an epimorphism from the free group $\alpha^{-1}(H)$ onto H , it now follows that

$$\text{rk}(H) \leq [G : H] \cdot (\text{rk}(G) - 1) + 1 \leq [G : H] \cdot \text{rk}(G). \quad \square$$

We now let π be a finitely generated group and let $\phi : \pi \rightarrow \mathbb{Z}$ be a homomorphism. Then we write

$$\pi_n := \text{Ker}(\pi_1(M) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}_n),$$

and we refer to

$$\text{rg}(\pi, \phi) := \liminf_{n \rightarrow \infty} \frac{1}{n} \text{rk}(\pi_n)$$

as the *rank gradient* of (π, ϕ) . It is a consequence of (1) that this limit does indeed exist. (Note that Lackenby defines the rank gradient using $\frac{1}{n}(\text{rk}(\pi_n) - 1)$ instead of $\frac{1}{n} \text{rk}(\pi_n)$, but it is clear that this gives rise to the same limit.)

The following lemma is now an immediate consequence of (1) and the definitions.

LEMMA 2.2. *Let π be a finitely generated group, and let $\phi : \pi \rightarrow \mathbb{Z}$ be a homomorphism.*

(1) *If $\alpha : \Gamma \rightarrow \pi$ is an epimorphism, then*

$$\text{rg}(\Gamma, \phi \circ \alpha) \geq \text{rg}(\pi, \phi).$$

(2) *If $\Gamma \subset \pi$ is a finite index subgroup, then*

$$\text{rg}(\Gamma, \phi) \leq [\pi : \Gamma] \cdot \text{rg}(\pi, \phi).$$

The following two lemmas show that Theorem 1.1 is indeed a strengthening of Stallings' fibering theorem.

LEMMA 2.3. *Let π be a finitely generated group, and let $\phi : \pi \rightarrow \mathbb{Z}$ be an epimorphism. If $\text{Ker}(\phi)$ is generated by k elements, then for any $n \in \mathbb{N}$, we have $\text{rk}(\pi_n) \leq k + 1$; in particular, $\text{rg}(\pi, \phi) = 0$.*

Proof. We write $K = \text{Ker}(\phi)$. Note that the epimorphism $\phi : \pi \rightarrow \mathbb{Z} = \langle t \rangle$ splits since $\langle t \rangle$ is in particular a free group. We can thus view π as a semidirect product $\pi = \langle t \rangle \rtimes K$. Under this identification, we furthermore have that $\pi_n = \langle t^n \rangle \rtimes K$. In particular, if $\{g_1, \dots, g_k\}$ is a generating set for K , then $\{t^n, g_1, \dots, g_k\}$ is a generating set for π_n . \square

LEMMA 2.4. *There exist a finitely presented group π and an epimorphism $\phi : \pi \rightarrow \mathbb{Z}$ such that $\text{Ker}(\phi)$ is infinitely generated but such that $\text{rk}(\pi_n) \leq 2$ for all n .*

Proof. We consider the semidirect product

$$\pi := \langle t \rangle \rtimes \mathbb{Z} \left[\begin{array}{c} 1 \\ 2 \end{array} \right]$$

where t^n acts on $\mathbb{Z}[1/2]$ by multiplication by 2^n together with the epimorphism $\phi : \pi \rightarrow \mathbb{Z}$ defined by $\phi(t^n) = n$ and $\phi(a) = 0$ for $a \in \mathbb{Z}[1/2]$. It is clear that $\text{Ker}(\phi) = \mathbb{Z}[1/2]$ is not finitely generated. On the other hand, it is straightforward to see that

$$\pi_n = \langle t^n \rangle \rtimes \mathbb{Z} \left[\begin{array}{c} 1 \\ 2 \end{array} \right]$$

is generated by t^n and $1 \in \mathbb{Z}[1/2]$. We thus showed that $\text{rk}(\pi_n) \leq 2$ for all n . \square

This raises the following question.

QUESTION 2.5. Do there exist a finitely presented group π and a homomorphism $\phi : \pi \rightarrow \mathbb{Z}$ such that $\text{rg}(\pi, \phi) = 0$ but such that the sequence $\text{rk}(\pi_n)$ is unbounded?

We conclude this section with the following elementary lemma.

LEMMA 2.6. *Let F be a free group on k generators, and $\phi : F \rightarrow \mathbb{Z}$ an epimorphism. Then*

$$\text{rg}(F, \phi) = k - 1.$$

The statement of the lemma already appears in [La05], but for the reader's convenience, we provide a proof.

Proof of Lemma 2.6. It is well known that any subgroup of F of index n is a free group on $n(k - 1) + 1$ generators. (This follows, for example, for an elementary argument using Euler characteristics of finite covers of graphs.) The lemma is now an immediate consequence of this observation. \square

2.2. The Heegaard Gradient

We now recall several basic definitions and facts on Heegaard splittings of 3-manifolds. We refer to [Jo] and [Sc02] for more details. We start out with several definitions:

- (1) A *compression body* H is the result of gluing disjoint 2-handles to $\Sigma \times [0, 1]$, where Σ is a closed surface, along $\Sigma \times 1$ and then capping off some spherical boundary components with 3-balls. We then write $\partial_+ H = \Sigma \times 0$ and $\partial_- H = \partial H \setminus \partial_+ H$. Note that a compression body with $\partial_- H = \emptyset$ is a handlebody.
- (2) A *Heegaard surface* for a 3-manifold M is an embedded separating surface $S \subset M$ such the two components of M cut along S are compression bodies H_1 and H_2 with $\partial_+ H_1 = \Sigma = \partial_+ H_2$.

Note that every compact 3-manifold admits a Heegaard surface (see e.g. [Sc02, Section 2]). In the following we refer to the minimal genus of a Heegaard surface as the *Heegaard genus* $h(M)$ of M .

Furthermore, given a directed 3-manifold (M, ϕ) with corresponding cyclic covers M_n , $n \in \mathbb{N}$, we define, following [La06], the *Heegaard gradient* of (M, ϕ) to be

$$\text{hg}(M, \phi) := \liminf_{n \rightarrow \infty} \frac{1}{n} h(M_n).$$

Note that if $p : \tilde{M} \rightarrow M$ is a k -fold cover, then the preimage of a Heegaard surface is again a Heegaard surface; it now follows easily that $h(\tilde{M}) \leq k \cdot h(M)$. We therefore see in particular that the Heegaard gradient is well defined.

We summarize a few key properties of the Heegaard genus in a lemma.

LEMMA 2.7. *Let M be a 3-manifold. Then the following hold:*

(1)

$$\text{rk}(\pi_1(M)) \leq \begin{cases} h(M) & \text{if } M \text{ is closed,} \\ 2h(M) & \text{otherwise.} \end{cases}$$

(2) If $\phi \in H^1(M; \mathbb{Z})$ is a primitive class, then

$$\text{rg}(\pi_1(M), \phi) \leq \begin{cases} \text{hg}(M) & \text{if } M \text{ is closed,} \\ 2 \text{hg}(M) & \text{otherwise.} \end{cases}$$

(3) If $\phi \in H^1(M; \mathbb{Z})$ is a primitive fibered class, then

$$h(M) \leq 2 \cdot \text{genus of the fiber} + 1.$$

REMARK. (1) Note that there exist closed 3-manifolds with $\text{rk}(\pi_1(M)) < h(M)$. In fact, there exist examples of such 3-manifolds that are Seifert fibered [BZ84], graph manifolds [We03; SWe07], and hyperbolic [Li11]. On the other hand, Souto [So08] and Namazi and Souto [NS09] showed that $\text{rk}(\pi_1(N)) = h(N)$ for hyperbolic 3-manifolds that are “sufficiently complicated” in a certain sense.

(2) To the best of our knowledge, it is not known whether there exist closed 3-manifold pairs (M, ϕ) with $\text{rg}(M, \phi) < \text{hg}(M, \phi)$.

(3) Note that Theorem 1.2 is an immediate consequence of Theorem 1.1 and Lemma 2.7.

Proof of Lemma 2.7. First note that if M is a closed 3-manifold and Σ is a Heegaard surface of genus g , then the compression bodies obtained by cutting M along Σ are in fact handlebodies. We can thus view M as the result of gluing together two handlebodies H_1, H_2 with g 1-handles each. In particular, we can build M out of H_1 by adding g 2-handles and one 3-handle. Since $\pi_1(H_1)$ is generated by g elements, it follows that $\text{rk}(\pi_1(M)) \leq g$. This evidently implies (1) and (2) for closed 3-manifolds.

If M is any 3-manifold and Σ is a Heegaard surface of genus g , then we can view M as the result of gluing 2-handles and 3-handles to $\Sigma \times [-1, 1]$. It follows that $\pi_1(M)$ is generated by a generating set for $\pi_1(\Sigma)$, that is, $\text{rk}(\pi_1(M)) \leq 2g$. This evidently implies (1) and (2) for 3-manifolds that are not closed.

We now turn to the proof of (3). Suppose that Σ is the fiber of a fibration $M \rightarrow S^1$. We can then identify M with $(\Sigma \times [0, 1]) / (x, 0) \sim (f(x), 1)$ for some self-diffeomorphism f of Σ . If M is a closed 3-manifold, then we pick two disjoint disks D_1 and D_2 on Σ . Then

$$(\Sigma \setminus (D_1 \cup D_2)) \times 0 \cup (\Sigma \setminus (D_1 \cup D_2)) \times \frac{1}{2} \cup \partial D_1 \times \left[0, \frac{1}{2}\right] \cup \partial D_2 \times \left[\frac{1}{2}, 1\right]$$

is a surface of genus $2g + 1$, and it is in fact a Heegaard surface for M : it cuts M into

$$\begin{aligned} & (\Sigma - \text{int } D_1) \times \left[0, \frac{1}{2}\right] \cup (\text{int } D_2) \times \left[\frac{1}{2}, 1\right] \quad \text{and} \\ & (\Sigma - \text{int } D_2) \times \left[\frac{1}{2}, 1\right] \cup (\text{int } D_1) \times \left[0, \frac{1}{2}\right], \end{aligned}$$

each the union of a 1-handle with a handlebody of the form (bounded surface) \times (interval).

If M is not closed, then Σ has nontrivial boundary, and M has toroidal boundary $\partial\Sigma \times [0, 1]/(x, 0) \sim (f(x), 1)$. Let η be a closed, f -invariant tubular neighborhood of $\partial\Sigma$ in Σ , so $N = (\eta \times [0, 1])/ (x, 0) \sim (f(x), 1)$ is a tubular neighborhood of ∂M in M , and let D_1 be a disk in Σ disjoint from η . Taking $H_1 = (\Sigma - \text{int } D_1) \times [0, 1/2] \cup N$, we claim that the frontier S of H_1 in M is a Heegaard surface.

A maximal collection of disjoint, nonparallel, nonboundary-parallel arcs embedded in $\Sigma - (\text{int } D_1 \sqcup \eta)$ that each begin and end on ∂D_1 gives rise, by crossing with $[0, \frac{1}{2}]$, to a collection \mathcal{D} of disjoint compressing disks for S in H_1 with the property that $H_1 - (S \cup \bigcup \{D \in \mathcal{D}\})$ retracts onto ∂M . Thus, H_1 is a compression body. It is easy to see that the other side H_2 of S in M is a handlebody of the form described in the closed case, and the claim follows. \square

3. Proof of the Main Theorem for Closed Hyperbolic 3-Manifolds

Given a finitely generated group Γ acting on a tree T , an ‘‘accessibility’’ principle relates the combinatorics of $\Gamma \backslash T$ to the structure of Γ . *Acylindrical* accessibility, introduced by Sela [Se99], does not require prior knowledge of the structure of vertex or edge stabilizers, but only that their action on T is ‘‘nice enough’’:

DEFINITION. The action $\Gamma \times T \rightarrow T$ is *k-acylindrical* if no $g \in \Gamma - \{1\}$ fixes a segment of length greater than k and *k-cylindrical* otherwise.

We will later on make use of the following theorem of R. Weidmann.

THEOREM 3.1 (Weidmann [We02]). *Let Γ be a noncyclic, freely indecomposable, finitely generated group, and $\Gamma \times T \rightarrow T$ a k-acylindrical, minimal (i.e. leaving no proper subtree invariant) action. Then $\Gamma \backslash T$ has at most $1 + 2k(\text{rk}(\Gamma) - 1)$ vertices.*

We will use the height of edge stabilizers, a notion from [GMR98], to bound cylindricity of the action under consideration.

DEFINITION. The *height* of an infinite subgroup Λ in Γ is k if there is a collection of k essentially distinct conjugates of Λ such that the intersection of all the elements of the collection is infinite and k is maximal possible. (The conjugate of Λ by γ is *essentially distinct* from the conjugate by γ' if $\Lambda\gamma \neq \Lambda\gamma'$.)

LEMMA 3.2. *Suppose that a torsion-free group Γ acts on a tree T on the left, transitively on edges. If the stabilizer Λ of an edge e_0 has height k in Γ , then the action of Γ on T is k-acylindrical.*

Proof. Because the action is transitive on edges, each edge stabilizer is conjugate to Λ , and the conjugates corresponding to distinct edges are essentially distinct: for an edge $e \neq e_0$, the stabilizer of e in Γ is $\gamma^{-1}\Lambda\gamma$, where $\gamma \in \Gamma$ satisfies $\gamma \cdot e = e_0$. Every element $\lambda\gamma$ of $\Lambda\gamma$ thus also satisfies $(\lambda\gamma) \cdot e = e_0$.

Now suppose that $\gamma \in \Gamma - \{1\}$ fixes an edge arc of length n . Then γ , hence also the subgroup $\langle \gamma \rangle$ that it generates, is in the intersection of the conjugates corresponding to the edges of this arc. Since Γ is torsion-free, $\langle \gamma \rangle$ is infinite, and Λ has height at least n . But Λ has height k , so it follows that $\Gamma \times T \rightarrow T$ is k -acylindrical. \square

Let (M, ϕ) be a directed 3-manifold. We pick a properly embedded oriented surface S in M dual to ϕ of minimal complexity. (Here, recall that the complexity of a surface S with connected components $S_1 \cup \dots \cup S_k$ is defined as $\chi_-(S) = \sum_{i=1}^k \max\{-\chi(S_i), 0\}$.) We also pick a tubular neighborhood $S \times [-1, 1]$ of S in M .

We view S^1 as the topological space underlying a graph G , with a single vertex v and a single edge e . Note that there exists a canonical continuous map $p : M \rightarrow G$ given by sending $S \times (-1, 1) \rightarrow (-1, 1) \rightarrow e$ and by sending every point in $M \setminus S \times (-1, 1)$ to v . The induced map $p_* : \pi_1(M) \rightarrow \pi_1(G) = \mathbb{Z}$ is precisely the map given by $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$.

We denote by G_0 the graph that has one vertex for each component of $M \setminus S \times (-1, 1)$ and one edge for each component of $S \times [-1, 1]$ with the obvious attaching maps. Note that there exist canonical maps $q : M \rightarrow G_0$ and $r : G_0 \rightarrow G$ that make the following diagram commute:

$$\begin{array}{ccc} & M & \\ q \swarrow & & \searrow p \\ G_0 & \xrightarrow{r} & G. \end{array}$$

It is clear from the definitions that all the maps induce epimorphisms on fundamental groups. In particular, G_0 is not a tree, and hence its Euler characteristic $\chi(G_0)$ is nonpositive. If $\chi(G_0)$ is negative, then the conclusion of Theorem 1.1 requires no machinery.

LEMMA 3.3. *Let (M, ϕ) be a directed 3-manifold. If the graph G_0 has $\chi(G_0) < 0$, then $\text{rg}(M, \phi) \geq -\chi(G_0)$.*

Proof. Recall that $q_* : \pi_1(M) \rightarrow \pi_1(G_0)$ is an epimorphism; it thus follows from Lemma 2.2 that

$$\text{rg}(M, \phi) \geq \text{rg}(\pi_1(G_0), r_*).$$

The lemma is now an immediate consequence of Lemma 2.6. \square

Recall that G_0 is the underlying graph of a graph-of-spaces decomposition of M , with vertex spaces the components of $M \setminus S \times (-1, 1)$ and edge spaces those of S . (We use the perspective on graphs of groups and spaces from [SWa79]; for definitions, see p. 155 there. See also [Ser80] and [Tr80].) This has an associated left action of π on a tree T , without involutions, such that each vertex stabilizer is conjugate to $\pi_1(M)$ for some component X of $\overline{M - S \times (-1, 1)}$ and each edge stabilizer to $\pi_1(S_0)$ for some component S_0 of S (see [SWa79, pp. 166–167]).

Using this, we can now prove the nontrivial implication of Theorem 1.1 for closed hyperbolic 3-manifolds.

THEOREM 3.4. *Let (M, ϕ) be a directed 3-manifold where M is a closed hyperbolic 3-manifold. If ϕ is nonfibered, then $\text{rg}(M, \phi) > 0$.*

Proof. Let (M, ϕ) be a directed 3-manifold where M is a closed hyperbolic 3-manifold and where ϕ is nonfibered. We write $\pi = \pi_1(M)$, and we pick a surface S of minimal complexity dual to ϕ . Since M is hyperbolic, we can and will assume that no component of S is a sphere or a torus. We denote by G_0 the graph defined previously.

On account of Lemma 3.3 we may also assume that the graph G_0 has Euler characteristic 0. We will show below that S is connected and nonseparating, that is, $G_0 = G$. Assuming this for the moment, let us prove the result.

Since S is not a fiber surface, $\pi_1(S)$ is a quasi-Fuchsian subgroup of π (see e.g. [Bon86]). Therefore, by the main theorem of [GMRS98], $\pi_1(S)$ has finite height in π (cf. Corollary on [GMRS98, p. 322]), so by Lemma 3.2 the $\pi_1(M)$ -action on the tree determined by S is k -acylindrical for some $k \in \mathbb{N}$. This action has quotient G , with one edge and vertex, so in particular it is minimal. Since M is hyperbolic and closed, π is also noncyclic, freely indecomposable, and finitely generated.

For each $n \in \mathbb{N}$, π_n also acts on T , with quotient a graph G_n with n edges and vertices. The action of π_n inherits k -acylindricity from that of π , and since π_n has finite index in π , its action is also minimal. It now follows from Theorem 3.1 that

$$\text{rk}(\pi_n) \geq \frac{n-1}{2k} + 1. \quad (2)$$

We thus see that $\text{rg}(M, \phi) > 0$.

We return to showing that $G_0 = G$, assuming that $\chi(G_0) = 0$. Since G_0 has Euler characteristic zero, it is homotopy equivalent to its minimal-length closed edge path, call it γ . Each edge of G_0 that is not in γ is contained in a subtree T_0 of G_0 that intersects γ at a single vertex v_0 with the property that $T_0 - \{v_0\}$ is a component of $G_0 - \{v_0\}$. Since T_0 is a subtree, the component of S corresponding to any edge in T_0 is nullhomologous. Removing such a component thus reduces the complexity of S , so the fact that S has minimal complexity implies that there are none, that is, $G_0 = \gamma$.

We claim also that all edges of G_0 point in the same direction. Note that identifying $\pi_1(G)$ with \mathbb{Z} requires choosing an orientation for e . This in turn gives an orientation to the interval fibers of each component of $S \times [-1, 1]$ or, equivalently, an orientation to each edge of G_0 . If these do not all point in the same direction, at least one vertex v_0 of γ is the initial vertex of each edge containing it. The sum of the components of S corresponding to these edges is trivial in homology, again contradicting the fact that S has minimal complexity. The claim follows and implies that G_0 covers G . But $p_* = \phi$ maps onto $\pi_1(G)$, so we must have $G_0 = G$. \square

4. Proof of the Main Theorem for 3-Manifolds with Empty or Toroidal Boundary

4.1. Twisted Alexander Polynomials

In this section we quickly recall the definition of twisted Alexander polynomials. This invariant was initially introduced by Lin [Lin01], Wada [Wa94], and Kirk and Livingston [KL99]. We refer to the survey paper [FV10] for a detailed presentation.

Let M be a 3-manifold, let $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$, and let $\alpha : \pi_1(M) \rightarrow G$ be an epimorphism onto a finite group G . We write $\pi = \pi_1(M)$. We can now define a left $\mathbb{Q}[\pi]$ -module structure on $\mathbb{Q}[G] \otimes_{\mathbb{Q}} \mathbb{Q}[t^{\pm 1}] =: \mathbb{Q}[G][t^{\pm 1}]$ as follows:

$$g \cdot (v \otimes p) := (\alpha(g) \cdot v) \otimes (t^{\phi(g)} p),$$

where $g \in \pi$ and $v \otimes p \in \mathbb{Q}[G] \otimes_{\mathbb{Q}} \mathbb{Q}[t^{\pm 1}] = \mathbb{Q}[G][t^{\pm 1}]$.

Denote by \tilde{M} the universal cover of M . We then use the representation $\alpha \otimes \phi$ to regard $\mathbb{Q}[G][t^{\pm 1}]$ as a left $\mathbb{Q}[\pi]$ -module. The chain complex $C_*(\tilde{M})$ is also a left $\mathbb{Q}[\pi]$ -module via deck transformations. Using the natural involution $g \mapsto g^{-1}$ on the group ring $\mathbb{Q}[\pi]$, we can view $C_*(\tilde{M})$ as a right $\mathbb{Q}[\pi]$ -module. We can therefore consider the tensor products

$$C_*^{\phi \otimes \alpha}(M; \mathbb{Q}[G][t^{\pm 1}]) := C_*(\tilde{M}) \otimes_{\mathbb{Q}[\pi]} \mathbb{Q}[G][t^{\pm 1}],$$

which form a complex of $\mathbb{Q}[t^{\pm 1}]$ -modules. We then consider the $\mathbb{Q}[t^{\pm 1}]$ -modules

$$H_*^{\phi \otimes \alpha}(M; \mathbb{Q}[G][t^{\pm 1}]) := H_*(C_*^{\phi \otimes \alpha}(M; \mathbb{Q}[G][t^{\pm 1}])).$$

When ϕ is understood, we will drop it from the notation; similarly, if α is the trivial representation to $\text{GL}(1, \mathbb{Q})$, then we will also drop it from the notation. We will later on also consider the modules $H_*(M; \mathbb{Q}(t))$ and $H_*(M; \mathbb{Q}[t^{\pm 1}]/(t^k - 1))$, which are defined analogously.

Since M is compact and since $\mathbb{Q}[t^{\pm 1}]$ is a PID, we have an isomorphism

$$H_1^{\phi \otimes \alpha}(M; \mathbb{Q}[G][t^{\pm 1}]) \cong \bigoplus_{i=1}^r \frac{\mathbb{Q}[t^{\pm 1}]}{p_i(t)} \mathbb{Q}[t^{\pm 1}]$$

for some $p_i \in \mathbb{Q}[t^{\pm 1}]$. We define the *twisted Alexander polynomial* as follows:

$$\Delta_{M, \phi}^{\alpha} := \prod_{i=1}^r p_i(t) \in \mathbb{Q}[t^{\pm 1}].$$

Note that $\Delta_{M, \phi}^{\alpha} \in \mathbb{Q}[t^{\pm 1}]$ is well defined up to multiplication by a unit in $\mathbb{Q}[t^{\pm 1}]$. We also adopt the convention that we drop α from the notation if α is the trivial representation to $\text{GL}(1, \mathbb{Q})$.

We will later on make use of the following two facts about (twisted) Alexander polynomials.

LEMMA 4.1. *Let (M, ϕ) be a directed 3-manifold, and let $\alpha : \pi_1(M) \rightarrow G$ be an epimorphism onto a finite group. We denote by $p : \tilde{M} \rightarrow M$ the corresponding finite cover. We write $\tilde{\phi} := p^*\phi$. Then*

$$\Delta_{M, \phi}^\alpha = \Delta_{\tilde{M}, \tilde{\phi}}.$$

The lemma thus says that we can view a twisted Alexander polynomial of a directed 3-manifold (M, ϕ) as an untwisted Alexander polynomial of a corresponding cover of M . The lemma is a straightforward consequence of the Shapiro lemma. We refer to [FV08a, Lemma 3.3] or [FV10, Section 3] for details.

LEMMA 4.2. *Let (M, ϕ) be a directed 3-manifold with $\Delta_{M, \phi} = 0$. Let $n \in \mathbb{Z}$, and let $\pi_n := \text{Ker}(\pi \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n)$. Then*

$$b_1(\pi_n) \geq n.$$

Proof. First note that the assumption that $\Delta_{M, \phi} = 0$ implies that $H_1(M; \mathbb{Q}[t^{\pm 1}]) \cong \mathbb{Q}[t^{\pm 1}] \oplus H$ for some $\mathbb{Q}[t^{\pm 1}]$ -module H . It now follows from the universal coefficient theorem that for any n , we have a short exact sequence

$$\begin{aligned} 0 &\rightarrow H_1(M; \mathbb{Q}[t^{\pm 1}]) \otimes_{\mathbb{Q}[t^{\pm 1}]} \frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1} \\ &\rightarrow H_1\left(M; \frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1}\right) \\ &\rightarrow \text{Tor}_{\mathbb{Q}[t^{\pm 1}]}\left(H_0(M; \mathbb{Q}[t^{\pm 1}]), \frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1}\right) \rightarrow 0. \end{aligned}$$

Since $H_1(M; \mathbb{Q}[t^{\pm 1}]) \cong \mathbb{Q}[t^{\pm 1}] \oplus H$, it follows that

$$\dim_{\mathbb{Q}}\left(H_1\left(M; \frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1}\right)\right) \geq \dim_{\mathbb{Q}}\left(\frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1}\right) = n.$$

Recall that we assumed that ϕ is primitive, which implies that the map $\pi \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/n$ is surjective. We can thus apply Shapiro's lemma, which in this case states that

$$H_1(\pi_n; \mathbb{Q}) \cong H_1\left(\pi; \frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1}\right) \cong H_1\left(M; \frac{\mathbb{Q}[t^{\pm 1}]}{t^n - 1}\right). \quad \square$$

4.2. Twisted Alexander Polynomials and Fibered Classes

Let (M, ϕ) be a directed 3-manifold, and let $\alpha : \pi_1(M) \rightarrow G$ be an epimorphism onto a finite group. If ϕ is fibered, then it was shown by many authors at varying levels of generality that $\Delta_{M, \phi}^\alpha$ is monic, in particular, nonzero. We refer to [Ch03; KM05; GKM05; Ki07; FK06; Fr13] for details.

In [FV12], extending earlier results in [FV08b; FV11a; FV11b], the following converse was proved.

THEOREM 4.3. *Let (M, ϕ) be a directed 3-manifold. If $\phi \in H^1(M)$ is nonfibered, then there exists an epimorphism $\alpha : \pi_1(M) \rightarrow G$ onto a finite group G such that*

$$\Delta_{M,\phi}^\alpha = 0.$$

The proof of this theorem relies heavily on the result of Wise [Wi12a; Wi12b] that subgroups of hyperbolic 3-manifolds that are carried by an embedded surface are separable. (See also [AFW12] for precise references.)

4.3. Proof of Theorem 1.1

As discussed in the introduction, the proof of Theorem 1.1 reduces to the proof of the following theorem.

THEOREM 4.4. *If (M, ϕ) is a directed 3-manifold which is not fibered, then $\text{rg}(M, \phi) > 0$.*

Proof. Let (M, ϕ) be a directed 3-manifold such that ϕ is not fibered. We have to show that $\text{rg}(M, \phi) > 0$. By Theorem 4.3, there exists an epimorphism $\alpha : \pi_1(M) \rightarrow G$ onto a finite group G such that

$$\Delta_{M,\phi}^\alpha = 0.$$

We write $\pi := \pi_1(M)$ and $\tilde{\pi} := \text{Ker}(\alpha)$, and we denote by \tilde{M} the cover corresponding to $\tilde{\pi}$. Note that $\phi(\tilde{\pi}) = d\mathbb{Z}$ for some $d \neq 0 \in \mathbb{Z}$. We write $\tilde{\phi} := \frac{1}{d}p^*(\phi) \in \text{Hom}(\tilde{\pi}, \mathbb{Z}) = H^1(\tilde{M}; \mathbb{Z})$. Note that $\tilde{\phi}$ is a primitive class.

For any $n \in \mathbb{N}$, we furthermore write

$$\pi_n := \text{Ker}\{\pi \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}_n\} \quad \text{and} \quad \tilde{\pi}_n := \text{Ker}\{\tilde{\pi} \xrightarrow{\tilde{\phi}} \mathbb{Z} \rightarrow \mathbb{Z}_n\}.$$

We now have the following claim.

CLAIM. For any $n \in \mathbb{N}$, the group $\tilde{\pi}_n$ is a subgroup of π_{dn} of index at most $[\pi : \tilde{\pi}]$.

Note that for any $n \in \mathbb{N}$, we have

$$\tilde{\pi}_n := \text{Ker}\{\tilde{\pi} \xrightarrow{\tilde{\phi}} \mathbb{Z} \rightarrow \mathbb{Z}_n\} = \text{Ker}\{\tilde{\pi} \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}_{dn}\}.$$

We thus see that the group $\tilde{\pi}_n$ is indeed a subgroup of π_{dn} . We thus have the equalities

$$[\pi : \tilde{\pi}] \cdot [\tilde{\pi} : \tilde{\pi}_n] = [\pi : \tilde{\pi}_n] = [\pi : \pi_{dn}] \cdot [\pi_{dn} : \tilde{\pi}_n].$$

It now follows from $[\tilde{\pi} : \tilde{\pi}_n] = n$ and $[\pi : \pi_{dn}] = dn$ that

$$[\pi_{dn} : \tilde{\pi}_n] = \frac{1}{d}[\pi : \tilde{\pi}] \leq [\pi : \tilde{\pi}].$$

This concludes the proof of the claim.

It follows from Lemmas 4.1 that $\Delta_{\tilde{M}, p^*\phi} = 0$, which in turn implies that $\Delta_{\tilde{M}, \tilde{\phi}} = 0$. It now follows from Lemma 4.2 that

$$b_1(\tilde{\pi}_n) \geq n \quad \text{for any } n. \tag{3}$$

For any n , we thus have by (1) and (3) that

$$\frac{1}{n} \text{rk}(\pi_n) \geq \frac{1}{dn} \text{rk}(\pi_{dn}) \geq \frac{1}{dn} \frac{1}{[\pi : \tilde{\pi}]} \text{rk}(\tilde{\pi}_n) \geq \frac{1}{dn} \frac{1}{[\pi : \tilde{\pi}]} b_1(\tilde{\pi}_n) \geq \frac{1}{d} \frac{1}{[\pi : \tilde{\pi}]}.$$

It thus follows that $\text{rg}(M, \phi) > 0$ as desired. \square

Recall that we assumed throughout the paper that M is a compact 3-manifold with empty or toroidal boundary. The statement of Theorem 1.1 does not hold if M has a spherical boundary component. Indeed, if (M, ϕ) is a fibered directed 3-manifold, then deleting a 3-ball gives rise to a 3-manifold with the same fundamental group but which is no longer fibered. It is therefore reasonable to restrict ourselves to 3-manifolds with no spherical boundary components. Extending verbatim the definition of rank gradient to this context, it is straightforward to see that the statement of Theorem 1.1 applies also to this slightly more general case:

LEMMA 4.5. *Let M be a compact 3-manifold with no spherical boundary components that has at least one nontoroidal boundary component. Then M is not fibered, and for any primitive $\phi \in H^1(M; \mathbb{Z})$, we have $\text{rg}(M, \phi) > 0$.*

Proof. If M fibers over S^1 , then the boundary components also have to fiber over S^1 , which means that all boundary components have to be tori.

Now let M be a compact 3-manifold that has at least one nontoroidal boundary component F , and let $\phi \in H^1(M; \mathbb{Z}) = \text{Hom}(\pi_1(M), \mathbb{Z})$ be a primitive element. We have to show that $\text{rg}(M, \phi) > 0$.

We denote by $d \in \mathbb{Z}_{\geq 0}$ the unique element such that $\phi(\pi_1(F)) = d\mathbb{Z}$. We first suppose that $d > 0$. Given $n \in \mathbb{N}$, we consider the finite cover M_{nd} of M corresponding to $\pi_1(M) \xrightarrow{\phi} \mathbb{Z} \rightarrow \mathbb{Z}/nd$, and we furthermore consider the cover F_n of F corresponding to $\pi_1(F) \rightarrow \pi_1(M) \xrightarrow{\phi} d\mathbb{Z} \rightarrow d\mathbb{Z}/nd \cong \mathbb{Z}/n$. Note that by the assumption that d is positive the cover F_n is a connected cover of F . By the multiplicativity of the Euler characteristic under finite covers we see that

$$b_1(F_n) - 2 = n(b_1(F) - 2).$$

Since F is nonspherical and nontoroidal, we see in particular that $b_1(F) \geq 2n$.

Note that M_{nd} contains d copies of F_n as boundary components. By the standard half-live-half-die argument coming from Poincaré duality, we deduce that

$$b_1(M_{nd}) \geq \frac{1}{2}(d \cdot b_1(F_n)) = dn.$$

It is now obvious that $\text{rg}(M, \phi) \geq 1$.

The case that $d = 0$ is proved almost in the same way. We leave the details to the reader. \square

Proof of Theorem 1.3. For a compact 3-manifold M , let \widehat{M} be obtained from M by filling all spherical boundary components with balls. The inclusion map $M \rightarrow \widehat{M}$ takes Heegaard surfaces to Heegaard surfaces and induces an isomorphism of fundamental groups. Moreover, every Heegaard surface for \widehat{M} may be

isotoped into M by an innermost disk argument, giving a Heegaard surface there. In particular, the Heegaard genus of M equals that of \widehat{M} as does the rank of π_1 .

Since both spheres and balls lift to covers (having trivial π_1), if $M' \rightarrow M$ is a finite-degree cover, then \widehat{M}' is the cover of \widehat{M} corresponding to $\pi_1 M' < \pi_1 M = \pi_1 \widehat{M}$. It follows that the rank and Heegaard gradients of any family of covers $\{M_n \rightarrow M\}$ may be computed in the corresponding family $\{\widehat{M}_n \rightarrow \widehat{M}\}$, reducing Theorem 1.3 to our prior results. \square

5. Normal Generating Sets

In this section we will prove Theorem 1.4, whose statement we recall for the reader's convenience:

THEOREM 5.1. (1) *If (M, ϕ) is a nonfibered directed 3-manifold, then $\text{Ker}(\phi)$ admits a finite index subgroup with infinite normal rank.*

(2) *There exists a nonfibered directed 3-manifold (M, ϕ) such that $\text{Ker}(\phi)$ has finite normal rank.*

Proof. We first note that if π is any group, then any set of elements that normally generates π is also a generating set of $H_1(\pi; \mathbb{Z})$. It thus follows that

$$n(\pi) \geq b_1(\pi).$$

If (M, ϕ) is a nonfibered directed 3-manifold, then by Theorem 1.1 there exists an epimorphism $\alpha : \pi_1(M) \rightarrow G$ onto a finite group G such that

$$\Delta_{M, \phi}^\alpha = 0.$$

We write $\tilde{\pi} := \text{Ker}(\alpha)$, and we denote by \tilde{M} the cover of M corresponding to $\tilde{\pi}$. As in the proof of Theorem 4.4, we note that $\phi(\tilde{\pi}) = d\mathbb{Z}$ for some $d \neq 0 \in \mathbb{Z}$, and we write $\tilde{\phi} := \frac{1}{d}p^*(\phi) \in \text{Hom}(\tilde{\pi}, \mathbb{Z}) = H^1(\tilde{M}; \mathbb{Z})$. Note that by Lemma 4.1 $\Delta_{M, \phi}^\alpha = 0$ implies that $\Delta_{\tilde{M}, \tilde{\phi}} = 0$. This in turn is equivalent to saying that $H_1(\tilde{M}; \mathbb{Q}[t^{\pm 1}])$ is not a $\mathbb{Q}[t^{\pm 1}]$ -torsion, that is,

$$\dim(H_1(\tilde{\phi}\text{-cover of } \tilde{M}; \mathbb{Q})) = \infty.$$

We thus see that $b_1(\text{Ker}(\alpha \times \phi)) = \infty$, that is, $n(\text{Ker}(\alpha \times \phi)) = \infty$. Since $\text{Ker}(\alpha \times \phi)$ is a finite index subgroup of $\text{Ker}(\phi)$, this concludes the proof of (1).

We now turn to the proof of (2). Let (N, ψ) be a fibered directed 3-manifold with $N \neq S^1 \times D^2$. We denote the fiber surface by S and the monodromy by φ . We can then identify N with $(S \times [0, 1]) / (x, 0) \sim (\varphi(x), 1)$. We pick an essential simple closed curve C on $S \times \frac{1}{2}$, and we pick an open tubular neighborhood νC of C in $S \times (0, 1)$. We furthermore pick a nontrivial knot $K \subset S^3$. We then consider the 3-manifold

$$M := (N \setminus \nu C) \cup (S^3 \setminus \nu K),$$

where we glue the meridian of K to a push-off of C in $S \times \frac{1}{2}$, and where we glue the longitude of K to a meridian of C . We denote by $\phi \in H^1(M; \mathbb{Z})$ the class dual to $S \times 0 \subset M$.

We claim that (M, ϕ) has all the desired properties. We denote by \tilde{M} the infinite cyclic cover of M corresponding to ϕ . Given $i \in \mathbb{Z}$, we write

$$W_i := ((S \times [0, 1] \setminus \nu C) \cup (S^3 \setminus \nu K)) \times i.$$

Note that we can canonically identify \tilde{M} with

$$\left(\bigcup_{i \in \mathbb{Z}} W_i \right) / (x, i) \sim (\varphi(x), i + 1).$$

Also note that \tilde{M} contains the incompressible tori $\partial \overline{\nu C} \times i$; in particular, $\pi_1(\tilde{M})$ is not a surface group. It thus follows that ϕ is not a fibered class.

We now denote by Γ the smallest normal subgroup of $\pi_1(\tilde{M}) = \text{Ker}(\phi)$ that contains $\pi_1(S \times 0)$. We are done with the proof of Theorem 1.4 once we showed that $\Gamma = \text{Ker}(\phi)$. First note that C , and hence the meridian of $K \times 0$, lies in Γ . Since the meridian of K normally generates $\pi_1(S^3 \setminus \nu K)$, it follows that the longitude of $K \times 0$ also lies in Γ . It is now straightforward to see that $\pi_1(W_0) \subset \Gamma$. This in particular implies that $\pi_1(S \times 1)$ lies in Γ . But then the same argument as above shows that $\pi_1(W_1) \subset \Gamma$. Iterating this argument, we see that $\pi_1(W_i)$ lies in Γ for all $i \in \mathbb{N}$. Almost the same argument also shows that $\pi_1(W_i)$ lies in Γ for all $i \in \mathbb{Z}_{\leq 0}$. It now follows that $\pi_1(\tilde{M})$ is contained in Γ . \square

ACKNOWLEDGMENTS. We wish to thank Dan Silver for a helpful conversation. We are also grateful to the referee for many helpful comments.

References

- [AFW12] M. Aschenbrenner, S. Friedl, and H. Wilton, *3-manifold groups*, preprint, 2012, [arXiv:1205.0202](https://arxiv.org/abs/1205.0202).
- [BZ84] M. Boileau and H. Zieschang, *Heegaard genus of closed orientable Seifert 3-manifolds*, *Invent. Math.* 76 (1984), 455–468.
- [Bon86] F. Bonahon, *Bouts des variétés hyperboliques de dimension 3*, *Ann. of Math.* 124 (1986), 71–158.
- [Ch03] J. Cha, *Fibred knots and twisted Alexander invariants*, *Trans. Amer. Math. Soc.* 355 (2003), 4187–4200.
- [De13] J. DeBlois, *Explicit rank bounds for cyclic covers*, preprint, 2013, [arXiv:1310.7823](https://arxiv.org/abs/1310.7823).
- [Fr13] S. Friedl, *Twisted Reidemeister torsion, the Thurston norm and fibered manifolds*, preprint, 2013, to be published by *Geometriae Dedicata*.
- [FK06] S. Friedl and T. Kim, *The Thurston norm, fibered manifolds and twisted Alexander polynomials*, *Topology* 45 (2006), 929–953.
- [FSW13] S. Friedl, D. Silver, and S. Williams, *Splittings of knot groups*, preprint, 2013.
- [FV08a] S. Friedl and S. Vidussi, *Twisted Alexander polynomials and symplectic structures*, *Amer. J. Math.* 130 (2008), no. 2, 455–484.
- [FV08b] ———, *Symplectic $S^1 \times N^3$, surface subgroup separability, and vanishing Thurston norm*, *J. Amer. Math. Soc.* 21 (2008), 597–610.
- [FV10] ———, *A survey of twisted Alexander polynomials*, *The mathematics of knots: theory and application* (M. Banagl, D. Vogel, eds.), *Contributions in Mathematical and Computational Sciences*, pp. 45–94, 2010.

- [FV11a] ———, *Twisted Alexander polynomials detect fibered 3-manifolds*, *Ann. of Math.* 173 (2011), 1587–1643.
- [FV11b] ———, *Twisted Alexander polynomials and fibered 3-manifolds*, *Low-Dimensional and Symplectic Topology*, *Proc. Sympos. Pure Math.* 82 (2011), 111–130.
- [FV12] ———, *A vanishing theorem for twisted Alexander polynomials with applications to symplectic 4-manifolds*, *J. Eur. Math. Soc.* 16 (2013), no. 6, 2027–2041.
- [GMRS98] R. Gitik, M. Mitra, E. Rips, and M. Sageev, *Widths of subgroups*, *Trans. Amer. Math. Soc.* 350 (1998), no. 1, 321–329.
- [GKM05] H. Goda, T. Kitano, and T. Morifuji, *Reidemeister torsion, twisted Alexander polynomial and fibred knots*, *Comment. Math. Helv.* 80 (2005), no. 1, 51–61.
- [Jo] J. Johnson, *Notes on Heegaard splittings*, (<http://www.math.okstate.edu/~jjohnson/notes.pdf>).
- [KL99] P. Kirk and C. Livingston, *Twisted Alexander invariants, Reidemeister torsion and Casson–Gordon invariants*, *Topology* 38 (1999), no. 3, 635–661.
- [KM05] T. Kitano and T. Morifuji, *Divisibility of twisted Alexander polynomials and fibred knots*, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 4 (2005), no. 1, 179–186.
- [Ki07] T. Kitayama, *Normalization of twisted Alexander invariants*, preprint, 2007, [arXiv:0705.2371](https://arxiv.org/abs/0705.2371).
- [La05] M. Lackenby, *Expanders, rank and graphs of groups*, *Israel J. Math.* 146 (2005), 357–370.
- [La06] ———, *Heegaard splittings, the virtually Haken conjecture and property (τ)* , *Invent. Math.* 164 (2006), no. 2, 317–359.
- [Li11] T. Li, *Rank and genus of 3-manifolds*, *J. Amer. Math. Soc.* 26 (2013), no. 3, 777–829.
- [Lin01] X. S. Lin, *Representations of knot groups and twisted Alexander polynomials*, *Acta Math. Sin. (Engl. Ser.)* 17 (2001), no. 3, 361–380.
- [NS09] H. Namazi and J. Souto, *Heegaard splittings and pseudo-Anosov maps*, *Geom. Funct. Anal.* 19 (2009), 1195–1228.
- [Sc02] M. Scharlemann, *Heegaard splittings of compact 3-manifolds*, *Handbook of geometric topology* (R. Daverman, R. Sherr, eds.), pp. 921–953, North-Holland, Amsterdam, 2002.
- [SWe07] J. Schultens and R. Weidmann, *On the geometric and the algebraic rank of graph manifolds*, *Pacific J. Math.* 231 (2007), 481–510.
- [SWa79] P. Scott and T. Wall, *Topological methods in group theory*, *Homological Group Theory*, *Proc. Sympos.*, Durham, 1977, *London Math. Soc. Lecture Note Ser.* 65 (1979), 137–203.
- [Se99] Z. Sela, *Endomorphisms of hyperbolic groups. I. The Hopf property*, *Topology* 38 (1999), 301–321.
- [Ser80] J.-P. Serre, *Trees*, Springer-Verlag, Berlin–New York, 1980.
- [SW09a] D. Silver and S. Williams, *Nonfibered knots and representation shifts*, *Proceedings of the Postnikov Memorial Conference*, *Banach Center Publ.* 85 (2009), 101–107.
- [SW09b] ———, *Twisted Alexander polynomials and representation shifts*, *Bull. Lond. Math. Soc.* 41 (2009), 535–540.
- [So08] J. Souto, *The rank of the fundamental group of certain hyperbolic 3-manifolds fibering over the circle*, *Geom. Topol. Monogr.* 14 (2008), 505–518.
- [St62] J. Stallings, *On fibering certain 3-manifolds*, *Topology of 3-manifolds and related topics* (*Proc. The Univ. of Georgia Institute*, 1961), pp. 95–100, Prentice-Hall, Englewood Cliffs, N.J., 1962.

- [Tr80] M. Tretkoff, *A topological approach to the theory of groups acting on trees*, J. Pure Appl. Algebra 16 (1980), 323–333.
- [Wa94] M. Wada, *Twisted Alexander polynomial for finitely presentable groups*, Topology 33 (1994), no. 2, 241–256.
- [We02] R. Weidmann, *The Nielsen method for groups acting on trees*, Proc. Lond. Math. Soc. (3) 85 (2002), 93–118.
- [We03] T. Weidmann, *Some 3-manifolds with 2-generated fundamental group*, Arch. Math. (Basel) 81 (2003), no. 5, 589–595.
- [Wi09] D. Wise, *The structure of groups with a quasiconvex hierarchy*, Electron. Res. Announc. Math. Sci. 16 (2009), 44–55.
- [Wi12a] ———, *The structure of groups with a quasi-convex hierarchy*, 189 pages, preprint, 2012, downloaded on October 29, 2012 from (<http://www.math.mcgill.ca/wise/papers.html>).
- [Wi12b] ———, *From riches to RAAGs: 3-manifolds, right-angled Artin groups, and cubical geometry*, CBMS Regional Conference Series in Mathematics, 2012.

J. DeBlois
Department of Mathematics
University of Pittsburgh
jdeblois@pitt.edu

S. Friedl
Fakultät für Mathematik
Universität Regensburg
Germany
sfriedl@gmail.com

S. Vidussi
Department of Mathematics
University of California
Riverside, CA 92521
USA
svidussi@math.ucr.edu