

# A Generalization of Sankaran and LVMB Manifolds

LAURENT BATTISTI & KARL OELJEKLAUS

ABSTRACT. We describe the construction of a new class of non-Kähler compact complex manifolds. They can be seen as a generalization of Sankaran, OT, and LVMB manifolds. Moreover, we give properties of these new spaces. Their Kodaira dimension is  $-\infty$ , and under a mild condition, their algebraic dimension is equal to zero.

## 0. Introduction

In this article, we construct a new family of non-Kähler complex compact manifolds by a combination of methods of Bosio [5] and Sankaran [12]. The class of manifolds constructed in this paper appears as a generalization of already known examples of non-Kähler manifolds, namely LVMB and Sankaran manifolds (ibid) along with OT manifolds [9].

Although the field of non-Kähler geometry remains relatively unexplored, there is, nevertheless, a constant progress. New classes of non-Kähler compact complex manifolds have been constructed and studied recently.

The first example is given by the class of LVMB manifolds. Their construction is due to Bosio [5] and can be summarized as follows. Given a family of subsets of  $\{0, \dots, n\}$  all having  $2m + 1$  elements (where  $n$  and  $m$  are integers such that  $2m \leq n$ ) and a family of  $n + 1$  linear forms on  $\mathbb{C}^m$  satisfying technical conditions, we can find an open subset  $U$  of  $\mathbb{P}_n(\mathbb{C})$  and an action of a complex Lie group  $G \cong \mathbb{C}^m$  on  $\mathbb{P}_n(\mathbb{C})$  such that the quotient  $U/G$  is a compact complex manifold. These manifolds generalize Hopf and Calabi–Eckmann manifolds, and they also generalize a class of manifolds due to Meersseman [8], called LVM manifolds.

OT manifolds were constructed in [9] by the second author and M. Toma. We start by choosing an algebraic number field  $K$  having  $s > 0$  (resp.  $2t > 0$ ) real (resp. complex) embeddings. Then, for a nice choice of a subgroup  $A$  of the groups of units  $\mathcal{O}_K^*$  of  $K$ , the quotient  $X(K, A)$  of  $\mathbb{H}^s \times \mathbb{C}^t$  under the action of  $A \times \mathcal{O}_K$  is a complex compact manifold. The required condition on  $A$  is that the projection on the first  $s$  coordinates of its image through the logarithmic representation of units

$$\begin{aligned} \ell : \mathcal{O}_K^* &\longrightarrow \mathbb{R}^{s+t}, \\ a &\longmapsto (\ln |\sigma_1(a)|, \dots, \ln |\sigma_s(a)|, 2 \ln |\sigma_{s+1}(a)|, \dots, 2 \ln |\sigma_{s+t}(a)|) \end{aligned} \tag{1}$$

---

Received May 5, 2014. Revision received September 18, 2014.

The second author is partially supported by the ANR project MNGNK, decision #ANR-10-BLAN-0118.

is a full lattice in  $\mathbb{R}^s$ . Such an  $A$  is called **admissible**. When there exists no proper intermediate field extension  $\mathbb{Q} \subset K' \subset K$  with  $A \subset \mathcal{O}_{K'}^*$ , we say that the manifold  $X(K, A)$  is of **simple type**.

In [12], G. Sankaran studies the action of a discrete group  $W$  isomorphic to  $\mathbb{Z}^b$  on an open subset  $U$  of a toric manifold, chosen to obtain a compact quotient  $U/W$ . The group  $W$  is a subgroup of the group of units of a number field  $K$ , and the construction of the infinite fan defining the toric manifold narrowly depends on  $W$ . This construction generalizes the previous ones due to Inoue, Kato, Sankaran himself, and Tsuchihashi. In the following, we will refer to the manifolds constructed in [12] as *Sankaran manifolds*.

Our construction combines the methods of Bosio and Sankaran. First, we take the quotient of a well-chosen  $n$ -dimensional toric manifold  $X_\Delta$  under the action of a complex Lie subgroup  $G \cong \mathbb{C}^t$  of  $(\mathbb{C}^*)^n$  such that the quotient  $X$  of  $X_\Delta$  by  $G$  is a (not necessarily compact) manifold and then we choose a suitable open subset of  $X$  on which a discrete group  $W$  acts and gives a compact quotient. The group  $W$  is a subgroup of the group of units of a number field  $K$  having  $s > 0$  real and  $2t > 0$  complex embeddings, with  $s + 2t = n$ . Of course, the choices of  $W$  and  $\Delta$  are strongly related as in Sankaran's case. We obtain an  $(s + t)$ -dimensional non-Kähler complex compact manifold.

The paper is organized as follows: In Section 1, we introduce the notation and results that will be needed in the sequel.

The next section is devoted to a description of the construction itself, which is subdivided into two steps. At the end of Section 2, some remarks are made on the structure of the manifolds, and we show how they generalize other known classes of manifolds.

In Section 3, we provide results on invariants and geometric properties of the new manifolds. We will prove that their Kodaira dimension is  $-\infty$ , that they are not Kähler, and finally show that their algebraic dimension is zero under a mild technical assumption. Moreover, we compute the second Betti number of OT manifolds under the same assumption, a fact that was already known for OT manifolds of simple type.

We conclude the paper by giving a concrete example of a three-dimensional manifold.

## 1. Preliminaries

Before starting the construction, we need to introduce or recall notation and results. Most of them can be found in Sankaran's paper [12], and others come from [9] and [2]. Detailed proofs are only provided when modifications are needed to fit in with the current context.

The section is divided into four parts as follows: First, we settle the number-theoretic ground of our construction. Then, we study complex Lie groups that will arise later, in particular, a certain Cousin group, that is, a complex Lie group without nonconstant holomorphic functions. Furthermore, we state some facts

about manifolds with corners associated to a fan and finally construct a special fan.

1.1. Number-Theoretic Notation and Results

Let  $K$  be a number field having  $n = s + 2t$  distinct embeddings in  $\mathbb{C}$ , and let  $\sigma_1, \dots, \sigma_s$  be the  $s$  real, and  $\sigma_{s+1}, \dots, \sigma_n$  be the  $2t$  complex (nonreal) ones. Up to reordering these embeddings, we can assume that  $\sigma_{s+i} = \bar{\sigma}_{s+t+i}$  for  $1 \leq i \leq t$ . We also require that both  $s$  and  $t$  are strictly positive.

Define the map

$$\begin{aligned} \sigma_K : K &\longrightarrow \mathbb{R}^s \times \mathbb{C}^{2t}, \\ k &\longmapsto (\sigma_1(k), \dots, \sigma_s(k), \sigma_{s+1}(k), \dots, \sigma_n(k)). \end{aligned}$$

Denote by  $\mathcal{O}_K$  the set of algebraic integers of  $K$ . The image  $\sigma_K(\mathcal{O}_K)$  is a lattice of rank  $n$  in  $\mathbb{R}^s \times \mathbb{C}^{2t} \subset \mathbb{C}^n$ . Let  $\mathcal{O}_K^*$  be the group of units of  $\mathcal{O}_K$ . Since  $s \geq 1$ , Dirichlet’s unit theorem tells us that  $\mathcal{O}_K^*$  is isomorphic to  $\{\pm 1\} \times \mathbb{Z}^{s+t-1}$ . The group  $\mathcal{O}_K^*$  acts on  $\mathbb{C}^n$  by componentwise multiplication:

$$\eta \cdot (z_1, \dots, z_n) := (\sigma_1(\eta)z_1, \dots, \sigma_n(\eta)z_n).$$

For all  $\eta \in \mathcal{O}_K^*$ , set  $\eta_i := \sigma_i(\eta)$  if  $i = 1, \dots, s$  and  $\eta_i := |\sigma_i(\eta)|$  if  $i = s + 1, \dots, s + t$ . Denote by  $\mathcal{O}_K^{*,+}$  the set of units  $\eta \in \mathcal{O}_K^*$  such that  $\eta_i > 0$  for all  $i = 1, \dots, s$ . The following theorem is due to Sankaran. For the convenience of the reader, we give its proof.

**THEOREM 1.1** ([12], Theorem 2.1). *For all  $b \in \{1, \dots, s\}$ , there exists a subgroup  $W < \mathcal{O}_K^{*,+}$  of rank  $b$  satisfying the following condition, called “Assumption C”<sup>1</sup>: for all  $\eta \neq 1 \in W$ , either there is  $i \leq b$  such that  $\eta_i > \eta_j$  for every  $j > b$ , or there is  $i \leq b$  such that  $\eta_i < \eta_j$  for every  $j > b$ .*

*Proof.* Set  $g := s + t - b$  and to every  $\eta \in \mathcal{O}_K^{*,+}$  associate an element of  $\mathbb{R}^{bg}$  via the map

$$\begin{aligned} \varphi_b : \mathcal{O}_K^{*,+} &\longrightarrow \mathbb{R}^{bg}, \\ \eta &\longmapsto \begin{pmatrix} \ln(\eta_1/\eta_{b+1}) & \cdots & \ln(\eta_1/\eta_{s+t}) \\ \vdots & \ln(\eta_i/\eta_{b+j}) & \vdots \\ \ln(\eta_b/\eta_{b+1}) & \cdots & \ln(\eta_b/\eta_{s+t}) \end{pmatrix}. \end{aligned}$$

For  $i = 1, \dots, b$ , let  $\pi_{i,b}$  be the projection

$$\begin{aligned} \pi_{i,b} : \mathbb{R}^{bg} &\longrightarrow \mathbb{R}^g, \\ (t_{i,j}) &\longmapsto (t_{i,j})_{j=1, \dots, g} \end{aligned}$$

that is,  $\pi_{i,b}$  maps an element of  $\mathbb{R}^{bg}$  written in matrix form to its  $i$ th row.

<sup>1</sup>In [12], Sankaran formulates two others assumptions (A and B), which we do not use here.

The fact that  $W$  satisfies Assumption C means that for all  $\eta \neq 1 \in W$ , there exists  $i \in \{1, \dots, b\}$  such that

$$\pi_{i,b}(\eta) \in Q_b := \{(x_1, \dots, x_g) \in \mathbb{R}^g \mid \text{all } x_j \text{ are nonzero and of same sign}\}.$$

Denote by  $U_b$  the vector space  $\varphi_b(\mathcal{O}_K^{*,+}) \otimes \mathbb{R} \cong \mathbb{R}^{s+t-1} \subset \mathbb{R}^{bg}$ . In order to prove the existence of a group  $W$  of rank  $b$  verifying Assumption C, it is enough to find a  $b$ -dimensional linear subspace  $A \subset U_b$  that is generated by elements of  $\varphi_b(\mathcal{O}_K^{*,+})$  and such that for every  $x \neq 0 \in A$ , there exists  $i \in \{1, \dots, b\}$  with  $\pi_{i,b}(x) \in Q_b$ . We proceed by induction.

For  $b = 1$ , the result is clear since it is enough to take  $A$  to be any line passing through the origin and any point of  $Q_1$ .

Assume that the result is proved at rank  $b - 1$ : there exists a  $(b - 1)$ -dimensional linear subspace  $A'$  of  $U_{b-1}$  having the following property:

$$\text{for all } x \in A', \text{ there exists } i \in \{1, \dots, b - 1\} \text{ such that } \pi_{i,b-1}(x) \in Q_{b-1}.$$

In this case, for all  $x \in A'$ , the integer  $i$  satisfies  $\pi_{i,b}(x) \in Q_b$ .

Now, fix a projection along  $A'$ , say  $\pi_{A'} : U_b \rightarrow \mathbb{R}^g$ . There is a map  $L \in GL_g(\mathbb{R})$  such that the following diagram commutes:

$$\begin{array}{ccc} & U_b & \\ \pi_{b,b} \swarrow & & \searrow \pi_{A'} \\ \mathbb{R}^g \cong \pi_{b,b}(U_b) & \overset{L}{\dashrightarrow} & U_b/A' \cong \mathbb{R}^g. \end{array}$$

Now we consider  $Q_{A'} := L(Q_b)$ . Take a line  $\ell \subset Q_{A'} \cup \{0\}$ ; then the subspace  $A := \pi_{A'}^{-1}(\ell)$  fits our purpose. To see this, pick an  $x \in A$ . Either we have  $x \in A'$ , and in this case, there is  $i \in \{1, \dots, b\}$  such that  $\pi_{i,b}(x) \in Q_b$ , or  $\pi_{A'}(x) \in \ell \setminus \{0\} \subset Q_{A'}$  and  $\pi_{A'}(x) = L(\pi_{b,b}(x))$ , which means that  $\pi_{b,b}(x) \in Q_b$ .  $\square$

### 1.2. A Cousin Group

Consider now the linear subspace

$$H := \{(0, \dots, 0, z_{s+t+1}, \dots, z_{s+t+t}) \mid z_{s+t+1}, \dots, z_{s+t+t} \in \mathbb{C}\} \cong \mathbb{C}^t$$

of  $\mathbb{C}^n$  and call  $\pi_H$  the projection from  $\mathbb{C}^n$  to  $\mathbb{C}^{s+t}$  with respect to  $H$  given by the  $s + t$  first coordinates.

LEMMA 1.2. *The restriction of  $\pi_H$  to  $\sigma_K(\mathcal{O}_K)$  is injective.*

*Proof.* It is sufficient to check that  $H \cap \sigma_K(\mathcal{O}_K) = \{0\}$ , which is straightforward because all  $\sigma_i$  are embeddings.  $\square$

Now let  $W$  be as in the previous theorem for some  $b \in \{1, \dots, s\}$ . Notice that the action of  $W$  on  $\mathbb{C}^n$  (resp.  $\mathbb{C}^n/H \cong \mathbb{C}^{s+t}$ ) induces an action on  $\sigma_K(\mathcal{O}_K)$  (resp.  $\pi_H(\sigma_K(\mathcal{O}_K))$ ).



The group  $H$  is a closed Lie subgroup of  $\mathbb{C}^n/\sigma_K(\mathcal{O}_K) \cong (\mathbb{C}^*)^n$  via the following map  $\iota$ :

$$H \longrightarrow (\mathbb{C}^*)^n, \tag{2}$$

$$\begin{pmatrix} 0 \\ \vdots \\ 0 \\ z_{s+t+1} \\ \vdots \\ z_{s+t+t} \end{pmatrix} \longmapsto \begin{pmatrix} \exp(2i\pi \sum_{i=1}^t h_{i,1} z_{s+t+i}) \\ \vdots \\ \exp(2i\pi \sum_{i=1}^t h_{i,n} z_{s+t+i}) \end{pmatrix},$$

where  $h_{i,j}$  is the  $j$ th coordinate of the vector  $h_i$ . We still denote by  $H$  the image of  $H$  under this map. We shall see that the group  $(\mathbb{C}^n/\sigma_K(\mathcal{O}_K))/H$  only has trivial holomorphic functions.

A connected complex Lie group admitting no nonconstant holomorphic functions is called a *Cousin group* or a *toroidal group*. We have the following:

**COROLLARY 1.5.** *The quotient of the complex Lie group  $\mathbb{C}^n/\sigma_K(\mathcal{O}_K) \cong (\mathbb{C}^*)^n$  by  $H$  is a Cousin group, which we call  $C_0$ . We denote by  $p$  the quotient map  $p : \mathbb{C}^n/\sigma_K(\mathcal{O}_K) \rightarrow C_0$ .*

*Proof.* By Lemma 1.2 the quotient of  $\mathbb{C}^n/\sigma_K(\mathcal{O}_K)$  by  $H$  is isomorphic to the quotient of  $\mathbb{C}^{s+t} = \mathbb{C}^n/H$  by  $\pi_H(\sigma_K(\mathcal{O}_K))$ , which is a Cousin group by Lemma 2.4 of [9]. □

**LEMMA 1.6.** *The subgroups  $H$  and  $(\mathbb{S}^1)^n$  of  $\mathbb{C}^n/\sigma_K(\mathcal{O}_K)$  intersect trivially.*

*Proof.* An element of this intersection corresponds to an element of the linear subspace  $H$  satisfying the following equation:

$$P_{\mathcal{B}, \mathcal{B}_K} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ z_{s+t+1} \\ \vdots \\ z_{s+t+t} \end{pmatrix},$$

where  $P_{\mathcal{B}, \mathcal{B}_K}$  is the change-of-basis matrix from the canonical basis to  $\mathcal{B}_K$  and all  $x_i$  are real. The  $i$ th and  $(i + t)$ th components of every vector of the basis  $\mathcal{B}_K$  are conjugated (for  $i = s + 1, \dots, s + t$ ). Since all  $x_i$  are real numbers, the  $i$ th and  $(i + t)$ th components of the vector  $(0, \dots, 0, z_{s+t+1}, \dots, z_{s+t+t})$  are also conjugated, and hence  $z_{s+t+1} = \dots = z_{s+t+t} = 0$ . □

**REMARK 1.7.** The quotient of  $(\mathbb{C}^*)^n$  by  $(\mathbb{S}^1)^n$  is given by the map

$$\text{ord} : (\mathbb{C}^*)^n \longrightarrow \mathbb{R}^n,$$

$$(z_1, \dots, z_n) \longmapsto (-\ln |z_1|, \dots, -\ln |z_n|).$$

With equation (2), we see that the space  $\text{ord}(H) \cong \mathbb{R}^{2t}$  is generated by the  $2t$  vectors  $\Re(h_j)$  and  $\Im(h_j)$  (for  $j \in \{1, \dots, t\}$ ); in other words,  $\text{ord}(H) = \hat{H}$ .

### 1.3. Manifolds with Corners

In [2], we can find the definition of the *manifold with corners* of a (not necessarily rational) fan  $\Delta$  of  $\mathbb{R}^n$ , denoted by  $\mathcal{M}c(\Delta)$ . Heuristically, this is a partial compactification of  $\mathbb{R}^n$  obtained in the following way: for every cone  $\sigma$  in  $\Delta$ , a complementary subspace of  $\text{span}(\sigma)$  “at infinity” is sent in the direction of  $\sigma$ . We also define a topology on this space, and when  $\Delta$  is rational, this space is (homeomorphic to) the manifold with corners of  $\Delta$  in the usual toric-geometrical sense, that is,  $X_\Delta/(\mathbb{S}^1)^n$ . This construction was inspired by the first chapter of [1].

We need the following two lemmas.

LEMMA 1.8 ([2], Lemma 1.17). *Let  $\sigma$  be a cone of a fan  $\Delta$ . Then the closure  $S$  of  $\sigma \subset \mathcal{M}c(\Delta)$  in  $\mathcal{M}c(\Delta)$  is compact.*

LEMMA 1.9 ([2], Lemma 2.3). *Let  $\Delta$  be a fan in  $\mathbb{R}^n$ , and  $E \cong \mathbb{R}^k$  be a linear subspace of  $\mathbb{R}^n$ . The action of  $E$  on  $\mathcal{M}c(\Delta)$  is proper if and only if the restriction of the quotient map  $\pi : \mathbb{R}^n \rightarrow \mathbb{R}^n/E$  to the support  $|\Delta|$  of  $\Delta$  is injective.*

We state the following definition and proposition, the proof of which is straightforward.

DEFINITION 1.10. Let  $\Delta$  be a (not necessarily rational) fan of  $\mathbb{R}^n$ , and  $G$  a discrete group of  $GL_n(\mathbb{R})$ . We say that  $G$  **acts** on  $\Delta$  if for every cone  $\sigma \in \Delta$  and every  $g \in G$ , we have  $g(\sigma) \in \Delta$ . This action is called **free** if  $g(\sigma) \neq \sigma$  when  $g$  is not the identity of  $G$  and  $\sigma \neq \{0\}$  is a cone of  $\Delta$ . This action is called **properly discontinuous** if for every  $\sigma \in \Delta$ , the set  $\{g \in G \mid (g(\sigma) \cap \sigma) \setminus \{0\} \neq \emptyset\}$  is finite.

For a fan  $\Delta$ , we set  $|\Delta|^* := |\Delta| \setminus \{0\}$ , and  $|\Delta|^{*c}$  is the complement of this set in  $\mathbb{R}^n$ .

PROPOSITION 1.11. *Let  $\Delta$  be a (not necessarily rational) fan, and  $G$  a discrete group acting freely and properly discontinuously on  $\Delta$ . Then this action induces an action of  $G$  on  $\mathcal{M}c(\Delta) \setminus |\Delta|^{*c}$ , which is also free and properly discontinuous.*

REMARK 1.12. Notice that  $\mathcal{M}c(\Delta) \setminus |\Delta|^{*c} = (\mathcal{M}c(\Delta) \setminus \mathbb{R}^n) \cup |\Delta|^*$ , that is, this set is the support of  $\Delta$  (except 0) to which we add the components “at infinity” of  $\mathcal{M}c(\Delta)$ .

### 1.4. A Suitable Fan

In the vector space  $E$ , we construct a rational fan  $\Delta$  with respect to the lattice  $\mathcal{O}_K \subset E$ . First, this fan has to be  $W$ -invariant. Furthermore, the image of its support under the projecting map  $\pi_{\hat{H}}$  must be included in an open degenerate proper cone  $\Omega \subset \pi_{\hat{H}}(E) \cong \mathbb{R}^s$  invariant under the action of  $W$ . *Degenerate* means that the closure of the cone  $\Omega$  contains a nontrivial linear subspace of  $\pi_{\hat{H}}(E)$ . If these

conditions are satisfied, then there is an action of  $W$  on the toric manifold associated to this fan.

The projection  $\pi_{\tilde{H}}$  is injective on  $\sigma_K(\mathcal{O}_K)$  and  $W$ -equivariant. Assume that there is a fan  $\Delta'$  in  $\pi_{\tilde{H}}(E)$  that is generated by elements of  $\pi_{\tilde{H}}(\sigma_K(\mathcal{O}_K))$  and whose support is such a cone  $\Omega$ . Take its pullback via the map  $\pi_{\tilde{H}}$ . This is the desired fan  $\Delta$ .

To construct the fan  $\Delta'$  in  $\pi_{\tilde{H}}(E)$ , we follow the steps of Sankaran in [12] (Thm. 2.5). Again, we only give proofs of results when they need modifications, and all the others adapt readily.

Let  $\Omega$  be an open degenerate cone in  $\pi_{\tilde{H}}(E)$  invariant under  $W$ . We start by describing the maximal vector subspace contained in its closure.

LEMMA 1.13 ([12], Lemma 1.2). *The vector space  $N$  of maximal dimension contained in the closure of  $\Omega$  is of the form*

$$N = \{(x_1, \dots, x_s) \in \pi_{\tilde{H}}(E) \mid x_{i_1} = \dots = x_{i_{s-h}} = 0\}$$

with  $h > 0$ .

REMARK 1.14. Up to renumbering the coordinates of  $\pi_{\tilde{H}}(E)$  and all the  $\sigma_i$  simultaneously, we may assume that  $N$  is the set

$$\{(x_1, \dots, x_s) \in \pi_{\tilde{H}}(E) \mid x_1 = \dots = x_{s-h} = 0\}.$$

This allows us to describe  $\Omega$  explicitly.

LEMMA 1.15 ([12], Lemma 1.3). *We have  $\Omega = N \times L_+$ , where*

$$L = \{x \in \pi_{\tilde{H}}(E) \mid x_{s-h+1} = \dots = x_s = 0\}$$

and

$$L_+ = \{x \in L \mid \pm x_i > 0, i \in \{1, \dots, s-h\}\}.$$

Up to composition with a suitable element of  $\mathcal{O}_K^*$ , we may assume that  $L_+$  is the set  $\{x \in L \mid x_i > 0, i \in \{1, \dots, s-h\}\}$ .

Now, we take an integer  $h \in \{1, \dots, s-1\}$  and consider the expression of  $\Omega$  given by Lemma 1.15. To ensure that  $L_+/W$  is a compact manifold, we choose the rank  $b$  of  $W$  to be equal to the dimension  $s-h$  of  $L$ . In particular, we have  $1 \leq b < s$ . Recall that given an integer  $b \in \{1, \dots, s-1\}$ , Theorem 1.1 guarantees the existence of a subgroup  $W$  of rank  $b$  of  $\mathcal{O}_K^{*,+}$  satisfying Assumption C.

PROPOSITION 1.16 ([12], Prop. 1.4). *The action of  $W$  on  $\Omega$  is free and properly discontinuous.*

PROPOSITION 1.17 ([12], Prop. 2.2). *If  $C \subset \Omega \cup \{0\}$  is a nondegenerate closed cone, then, for all  $\eta \in W$ , we have  $\bigcap_{a \in \mathbb{Z}} \eta^a C \subset L_+$ .*

REMARK 1.18. Let us recall the idea of Sankaran’s proof: there is a real number  $\delta > 0$  such that

$$C \subset C_\delta := \left\{ v \in \Omega \mid \sum_{i=1}^b v_i^2 \geq \delta \sum_{j>b} v_j^2 \right\}. \tag{3}$$

It is hence enough to establish the result for  $C_\delta$ . This can be done by observing that there exist  $N > 1$  and  $a \in \mathbb{N}$  such that  $\eta^k C_\delta \subset C_{N^k \delta}$  for  $k > a$ . A crucial remark for the next part of the construction is that the constants  $a$  and  $N$  can be chosen independently of  $\eta \in W$ . For further details, see the proof of Theorem 2.4 in [12]. Assumption C and the fact that  $b < s$  are used in this proof.

THEOREM 1.19 ([12], Thm. 2.5). *There exists an infinite fan  $\Delta'$  of  $\mathbb{R}^s$  stable under the action of  $W$  such that its support  $|\Delta'|$  is  $(\Omega \setminus L_+) \cup \{0\}$ , all its cones are generated by elements of  $\pi_{\tilde{H}}(\mathcal{O}_K)$ , and  $\Delta'/W$  is a finite set of cones.*

REMARK 1.20. In particular,  $\Delta'$  is constructed in such a way that the action of  $W$  on it is free and properly discontinuous in the sense of Definition 1.10.

## 2. Construction of the New Manifolds

Now we are able to start the construction. It is divided into two steps. First, we choose a group  $W$  as in Theorem 1.1 and get an infinite fan on which  $W$  acts thanks to Theorem 1.19. We take the quotient  $X$  of its associated toric manifold by the complex Lie group  $H$  and check that we have an action of  $W$  on the complex manifold  $X$ .

The second step is to find a suitable open subset  $U$  of  $X$  such that its quotient under the action of  $W$  becomes a compact manifold.

### 2.1. Step 1

Let  $K$  be a number field with  $s > 0$  real and  $2t > 0$  complex embeddings. Choose an integer  $b \in \{1, \dots, s - 1\}$ . Theorem 1.1 shows the existence of a subgroup  $W$  of  $\mathcal{O}_K^{*,+}$  of rank  $b$  satisfying Assumption C.

Take a fan  $\Delta'$  in  $\mathbb{R}^s$  as in Theorem 1.19. Denote by  $\Delta$  its preimage  $\pi_{\tilde{H}}^{-1}(\Delta')$ , which is a rational fan of  $E$  with respect to  $\mathcal{O}_K$ , and consider its associated toric manifold denoted by  $X_\Delta$ . The group  $H$  acts as a closed subgroup of  $(\mathbb{C}^*)^n$  (see equation (2)) on  $X_\Delta$ , and we have the following:

LEMMA 2.1. *The action of  $H$  on  $X_\Delta$  is free and proper.*

*Proof.* To check that this action is proper, we first observe that the group  $\tilde{H} = \text{ord}(H) \cong H$  acts properly on  $\text{Mc}(\Delta)$ . This is a consequence of Lemma 1.9 and the fact that the map  $\pi_{\tilde{H}} : E \rightarrow E/\tilde{H}$  is injective on  $|\Delta|$  by construction. Finally, it is clear that the action of  $H$  on  $X_\Delta$  is proper if and only if the action of  $\tilde{H}$  on  $\text{Mc}(\Delta)$  is proper because  $(\mathbb{S}^1)^n$  is compact.

The isotropy group of a point  $x \in X_\Delta$  is a compact subgroup of  $H \cong \mathbb{C}^t$  and hence trivial. □

The preceding lemma tells us that the quotient  $X := X_\Delta/H$  is a complex manifold, and since  $(\mathbb{C}^*)^n$  is abelian, the following diagram is commutative:

$$\begin{array}{ccc}
 X_\Delta & \xrightarrow{(\mathbb{S}^1)^n} & \mathcal{M}c(\Delta) \\
 \downarrow H \cong \mathbb{C}^t & & \downarrow \text{ord}(H) = \tilde{H} \cong \mathbb{R}^{2t} \\
 X & \xrightarrow[\rho((\mathbb{S}^1)^n) \cong (\mathbb{S}^1)^n]{q} & \mathcal{M}c(\pi_{\tilde{H}}(\Delta))
 \end{array} \tag{4}$$

The action of  $(\mathbb{S}^1)^n$  on  $X_\Delta$  descends to  $X$  via the group  $\rho((\mathbb{S}^1)^n) \cong (\mathbb{S}^1)^n$ . We denote by  $q : X \rightarrow \mathcal{M}c(\pi_{\tilde{H}}(\Delta))$  the quotient map for this action. A detailed proof of the fact that  $\mathcal{M}c(\Delta)/\tilde{H}$  and  $\mathcal{M}c(\pi_{\tilde{H}}(\Delta))$  are homeomorphic can be found in Section 2.2.2, pp. 558–559 of [2].

LEMMA 2.2. *The action of  $W$  on  $X_\Delta$  descends to an action on  $X$  which we denote  $(\eta, x) \mapsto \eta \cdot x$ .*

*Proof.* We shall prove that the action of  $W$  normalizes the subgroup  $H$  of  $(\mathbb{C}^*)^n$ .

Let  $\eta \in W$ ,  $x \in X_\Delta$ , and  $z = (z_1, \dots, z_t) \in H$ . First,  $\eta$  is equivariant with respect to

$$\begin{aligned}
 \tilde{\eta} : (\mathbb{C}^*)^n &\longrightarrow (\mathbb{C}^*)^n, \\
 (t_1, \dots, t_n) &\longmapsto (t_1^{a_{1,1}} \dots t_n^{a_{1,n}}, \dots, t_1^{a_{i,1}} \dots t_n^{a_{i,n}}, \dots, t_1^{a_{n,1}} \dots t_n^{a_{n,n}}),
 \end{aligned}$$

where  $(a_{i,j})$  is the matrix of the linear map  $\eta$  in the basis  $\mathcal{B}_K$ ; see, for instance, [6], Theorem 6.4 on p. 244.

This means that  $\eta(\iota(z)x) = \tilde{\eta}(\iota(z))\eta(x)$ . The map  $\iota$  is defined in equation (2). Moreover,  $\tilde{\eta}(\iota(z)) = \iota(\sigma_{s+t+1}(\eta)z_1, \dots, \sigma_{s+t+1}(\eta)z_t)$ , and hence we have

$$\eta(\iota(z)x) = \iota(\sigma_{s+t+1}(\eta)z_1, \dots, \sigma_{s+t+1}(\eta)z_t)\eta(x). \quad \square$$

LEMMA 2.3. *The group  $W$  acts on  $\mathcal{M}c(\Delta)$ , and this action descends to  $\mathcal{M}c(\pi_{\tilde{H}}(\Delta))$ . Moreover, the map  $q$  is equivariant with respect to the action of  $W$ .*

*Proof.* For an element  $x \in X_\Delta$  (resp.  $y \in \mathcal{M}c(\Delta)$ ), we denote by  $H \cdot x$  and  $\tilde{H} \cdot y$  the  $H$ -orbit of  $x$  and the  $\tilde{H}$ -orbit of  $y$ , respectively. The proof of the previous lemma shows that the action of  $W$  descends to  $\mathcal{M}c(\Delta)$ . The projection map of  $E \subset \mathcal{M}c(\Delta)$  with respect to  $\tilde{H}$  commutes with the action of  $W$  (see Remark 1.4), and hence there is an action of  $W$  on  $\mathcal{M}c(\pi_{\tilde{H}}(\Delta))$ . Denote this action again by  $(\eta, \tilde{H} \cdot y) \mapsto \eta \cdot \tilde{H} \cdot y$ . Lets us prove that  $q$  is  $W$ -equivariant.

Let  $H \cdot x \in X$  and  $\eta \in W$ . By Lemma 2.2 and the commutativity of diagram (4) we have  $q(\eta \cdot H \cdot x) = q(H \cdot \eta(x)) = \tilde{H} \cdot \text{ord}(\eta(x))$ . We first prove that  $\text{ord}(\eta(x)) = \eta(\text{ord}(x))$  on the dense open set  $(\mathbb{C}^*)^n$  of  $X_\Delta$ .

We have

$$\text{ord}(x) = -\frac{1}{2\pi}(\ln |x_1|, \dots, \ln |x_n|)$$

and

$$\begin{aligned} \text{ord}(\eta(x)) &= -\frac{1}{2\pi} (\ln(|x_1^{a_{1,1}} \cdots x_n^{a_{1,n}}|), \dots, \ln(|x_1^{a_{n,1}} \cdots x_n^{a_{n,n}}|)) \\ &= -\frac{1}{2\pi} \left( \sum_{i=1}^n a_{1,i} \ln |x_i|, \dots, \sum_{i=1}^n a_{n,i} \ln |x_i| \right) \\ &= \eta(\text{ord}(x)). \end{aligned}$$

By the density of  $(\mathbb{C}^*)^n$  in  $X_\Delta$ , this equality holds on  $X_\Delta$ . Hence,

$$q(\eta \cdot H.x) = \tilde{H} \cdot \text{ord}(\eta(x)) = \tilde{H} \cdot \eta(\text{ord}(x)) = \eta \cdot (\tilde{H} \cdot \text{ord}(x)). \quad \square$$

### 2.2. Step 2

The last step consists of finding an open subset  $U$  of  $X$  on which the action of  $W$  is free and properly discontinuous. This shows that the quotient  $U/W$  is a complex manifold. It turns out that the action of  $W$  on the manifold with corners  $\mathcal{Mc}(\pi_{\tilde{H}}(\Delta))$  carries enough information for our purpose. In this space, we find an open subset on which the action of  $W$  is free and properly discontinuous, with a compact fundamental domain. This proves the compactness of the quotient. Then, we pull everything back to  $X$  via the map  $q$ .

Let  $\eta_1, \dots, \eta_b$  be generators of  $W$ . Taking their inverses if necessary, we may assume that they all satisfy the following:

**ASSUMPTION  $\mathbb{C}^+$ .** For every  $i \in \{1, \dots, b\}$  and every nondegenerate closed cone  $C \subset \Omega \cup \{0\}$ , we have  $\bigcap_{a \in \mathbb{N}} \eta_i^a C \subset L_+$ . (Remark 1.18.)

Define the affine subspace

$$H_0 := \{(\underbrace{1, \dots, 1}_{b \text{ times}}, x_1, \dots, x_{s-b}) \mid x_1, \dots, x_{s-b} \in \mathbb{R}\} \subset E/\tilde{H}$$

and set  $H_i := \eta_i(H_0)$  for  $i = 1, \dots, b$ . Note  $B$  the (unbounded) convex envelope of  $H_0, H_1, \dots, H_b$ .

**LEMMA 2.4.** Let  $B_b$  be the image of  $B$  by the projection on the  $b$  first coordinates of  $E/\tilde{H}$ . Then  $\bigcup_{\eta \in W} \eta(B_b) = (\mathbb{R}_*^+)^b$ , and the union  $\bigcup_{\eta \in W \setminus W_{>1}} \eta(B_b)$  is bounded, where  $W_{>1}$  is the set of elements of  $W$  having at least one of their  $b$  first coordinates greater than or equal to 1.


*Proof.* For the first statement, notice that  $W$  is a lattice of  $\mathbb{R}^b$  via the map  $w \mapsto (\ln \eta_1, \dots, \ln \eta_b)$ . The second one comes from the fact that every element  $\eta$  of  $W \setminus W_{>1}$  satisfies  $\eta_i < 1$  for all  $i \in \{1, \dots, b\}$ . □

**LEMMA 2.5.** The action of  $W$  on  $\mathcal{U} := \mathcal{Mc}(\pi_{\tilde{H}}(\Delta)) \setminus |\Omega|^{*c}$  is free, properly discontinuous, and admits a compact fundamental domain. The same holds for the action of  $W$  on the preimage  $U := q^{-1}(\mathcal{U})$  in  $X$ .

*Proof.* The fact that the action is free and properly discontinuous is a consequence of Proposition 1.11. By Theorem 1.19 there is a finite set of cones  $\Sigma := \{\sigma_1, \dots, \sigma_d\}$  such that  $W \cdot \Sigma = \Delta'$ . We still assume that  $\eta_1, \dots, \eta_b$  is a family of generators of  $W$  satisfying assumption  $C^+$  and  $B$  is the set defined directly before Lemma 2.4. Let us consider the closure  $\bar{D}$  in  $\mathcal{M}c(\pi_{\tilde{H}}(\Delta))$  of the set

$$D := \underbrace{\left( \bigcup_{\eta \in W^+} \eta(|\Sigma|) \cap B \right)}_{D_1} \cup \underbrace{\left( |\Sigma| \cap \bigcup_{\eta \in W_{>1}} \eta(B) \right)}_{D_2}.$$

Here  $|\Sigma|$  is the union of the cones of  $\Sigma$ , and  $W^+$  is the set of elements of  $W$  satisfying assumption  $C^+$  along with the identity. Notice in particular that  $W^+ \subset W_{>1}$ . Figure 1 gives a picture of  $\bar{D}$  for  $s = 2$  and  $b = 1$ .



**Figure 1** The fundamental domain  $\bar{D}$

To prove the lemma, it is enough to show that  $\bar{D}$  is a compact fundamental domain for the action of  $W$ .

For compactness, first notice that  $D_1$  is bounded. This is a consequence of Remark 1.18. Indeed, there is an upper bound for the  $b$  first coordinates of points in  $D_1$  because of the definition of  $B$ , whereas for the  $s - b$  last coordinates, we observe that the inclusion  $\eta^a C_\delta \subset C_{N^a \delta}$  holds for  $a \in \mathbb{N}$  large enough,  $\delta > 0$ , and  $N > 1$ . Hence, the elements of  $D_1$  lie in a cone  $C_\delta$  for well-chosen  $\delta > 0$ . Equation (3) shows that a subset of  $C_\delta$  that is bounded in the  $b$  first coordinates is also bounded in the  $s - b$  last ones.

Since  $\Sigma$  is a finite set, the closure of  $D_2$  is the union of the closures of  $\sigma \cap \bigcup_{\eta \in W_{>1}} \eta(B)$  for  $\sigma \in \Sigma$ . Such a set is contained in  $\sigma \setminus B(0, C)$  for some constant  $C > 0$ . Indeed, let  $x = (x_1, \dots, x_s) \in B$  and  $\eta \in W_{>1}$ . For all  $\eta \in W_{>1}$ , there exists an integer  $i \leq b$  such that  $\eta_i \geq 1$ , so without loss of generality we may assume that  $\eta_1 \geq 1$ . The following inequalities hold:

$$\|\eta(x)\|^2 = \sum_{i=1}^s \eta_i^2 x_i^2 \geq \eta_1^2 x_1^2 \geq x_1^2 \geq \min_{y \in B} y_1^2 > 0.$$

The last inequality is a consequence of the fact that the projection of  $B$  on the first  $b$  coordinates is a compact set in  $\mathbb{R}^b$  having no point with a zero coordinate. All the constants  $C_i := \min_{y \in B} y_i^2$  for  $i = 1, \dots, b$  are strictly positive. Let  $C$  be the square root of the smallest of these constants. We have  $\|\eta(x)\| \geq C$ . Finally, the closure of  $D_2$  is compact by Lemma 1.8.

In order to finish the proof, we check that  $\bar{D}$  is a fundamental domain for the action of  $W$ .

First, observe that for all  $x \in \Omega \setminus L_+$ , there are two elements  $\eta$  and  $\eta' \in W$  such that  $x \in \eta(|\Sigma|) \cap \eta'(B)$ . We have either  $\eta^{-1}\eta' \in W^+ \subset W_{>1}$  or  $\eta\eta'^{-1} \in W^+$ . This respectively implies that either  $\eta^{-1}(x) \in |\Sigma| \cap \eta^{-1}\eta'(B)$  or  $\eta'^{-1}(x) \in \eta\eta'^{-1}(|\Sigma|) \cap B$ . If  $x \in L_+$ , Lemma 2.4 gives the conclusion since  $B \cap L_+ \subset \bar{D}$ .

For the remaining case, take a point  $x = y + \infty \cdot \tau$  in  $\bar{D}$  on a component at infinity of  $\text{Mc}(\pi_{\tilde{H}}(\Delta))$ . The second conclusion of Lemma 2.4 implies then that there is a constant  $C' > C$  such that for every cone  $\sigma$  of  $\pi_{\tilde{H}}(\Delta)$ ,

$$\sigma \setminus B(0, C') = \left( \sigma \cap \bigcup_{\eta \in W_{>1}} \eta(B) \right) \setminus B(0, C').$$

Now let  $\pi$  be the projection from  $\mathbb{R}^s$  to  $(\mathbb{R}^s)^\tau$  with respect to  $L(\tau)$ , where  $(\mathbb{R}^s)^\tau \oplus L(\tau) = \mathbb{R}^s$ . Let also  $\sigma$  be the  $s$ -dimensional cone having  $\tau$  as a face satisfying  $\pi(y) \in \pi(\sigma)$ , and let  $g$  be an element of the group  $W$  such that  $g(\sigma) \in \Sigma$ . Such a  $g$  always exists because of the definition of  $\Sigma$ . We have

$$g(x) = g(y) + \infty \cdot g(\tau).$$

Up to replacing  $y$  by  $y + w$  with  $w \in L(\tau)$ , we may assume that  $y \in \sigma$ ; hence,  $g(y) \in g(\sigma)$ , and the conclusion follows. We only need to prove the existence of  $\sigma$ . For this, take a generating set  $\{v_1, \dots, v_k\}$  of  $\tau$ , so that  $\tau$  is written as

$$\tau = \mathbb{R}_{\geq 0}v_1 + \dots + \mathbb{R}_{\geq 0}v_k.$$

Consider the open ball of radius  $\varepsilon > 0$  centered at  $v_0 := v_1 + \dots + v_k \in \tau$ . Since the support of  $\pi_{\tilde{H}}(\Delta)$  is  $(\Omega \setminus L_+) \cup \{0\}$ , there exists  $\varepsilon > 0$  small enough such that  $B(v_0, \varepsilon) \subset \Omega \setminus L_+$ . Then, by shrinking  $\varepsilon$  if necessary, we may assume that  $B(v_0, \varepsilon)$  only intersects the cones containing  $\tau$  as a face in view of the freeness of the  $W$ -action. Write

$$B(v_0, \varepsilon) = B(v_0, \varepsilon) \cap \bigcup_{\substack{\sigma > \tau \\ \dim \sigma = s}} \sigma.$$

The image of  $B(v_0, \varepsilon)$  under the projection  $\pi$  is an open neighborhood of  $0 \in \pi(\mathbb{R}^s)$ , and the set  $\pi(\bigcup_{\substack{\sigma > \tau \\ \dim \sigma = s}} \sigma)$  is a union of cones, so it is invariant under homothetic transforms of positive ratio. Hence,  $\pi(\bigcup_{\substack{\sigma > \tau \\ \dim \sigma = s}} \sigma) = \pi(\mathbb{R}^s)$ , there is a cone  $\sigma \in \pi_{\tilde{H}}(\Delta)$  admitting  $\tau$  as a face, and  $\pi(y) \in \pi(\sigma)$ .  $\square$

REMARK 2.6. In a similar fashion, we can describe a compact fundamental domain showing that Sankaran manifolds are compact. Actually, in [12, p. 47], the author mentions this possibility to obtain compactness, although it did not seem feasible in general.

Finally, the previous lemma implies the following:

PROPOSITION 2.7. *The quotient  $Y := U/W$  is a complex compact manifold of dimension  $s + t$ .*

### 2.3. Remarks

(a) The open subset  $U \subset X$  intersects the Cousin group  $C_0$  defined in Corollary 1.5. Note that  $U_0 := U \cap C_0$  is a dense open subset of  $U$ . The quotient of  $U_0$  by  $W$  is not compact, and this is the reason for adding divisors at infinity to  $U_0$ . In fact, the irreducible components of these divisors are generalized LVMB manifolds in the sense of Theorem 5.2 of [2].

To see this, first observe that the preimage of  $U \setminus U_0$  in  $X_\Delta$  is the set  $\bigcup_{\sigma \in \Delta \setminus \{0\}} \text{orb}(\sigma)$ . Let  $\sigma = \mathbb{R}_{\geq 0}v$  be a one-dimensional cone of  $\Delta$  with  $v \in \sigma_K(\mathcal{O}_K)$ . The fan of the toric manifold  $\overline{\text{orb}(\sigma)}$ , denoted by  $\Delta/\sigma$ , is obtained by taking the image of every cone of  $\Delta$  having  $\sigma$  as a face under the projection map with respect to  $\text{span}(\sigma)$ ; see Section VI.4 in [6]. Denote by  $\pi_\sigma : E \cong \mathbb{R}^n \rightarrow E/\text{span}(\sigma) \cong \mathbb{R}^{n-1}$  this projection. We get a finite fan in  $E/\text{span}(\sigma)$ , and since  $H$  trivially intersects  $\sigma_K(\mathcal{O}_K)$ , the images of  $H$  and  $\sigma_K(\mathcal{O}_K)$  under  $\pi_\sigma$  have a trivial intersection in  $E/\text{span}(\sigma)$ . Denote by  $\bar{\pi}_{\tilde{H}}$  the projection from  $E/\text{span}(\sigma)$  with respect to  $\pi_\sigma(H)$ ; this is an injective map on  $|\Delta/\sigma|$ . The fan  $\bar{\pi}_{\tilde{H}}(\Delta/\sigma)$  is complete because the closure of  $\sigma$  in  $\mathcal{M}c(\Delta)$  is compact by Lemma 1.8. All the hypotheses of Theorem 5.2 of [2] are satisfied.

(b) Our construction generalizes other known classes of manifolds.

(i) As for the case  $b = 0$ , which corresponds to the case that  $W$  is trivial, we recover the LVMB manifolds.

(ii) It is also possible to extend the construction to the case  $b = s$ . In this case, the fan  $\Delta$  then has to be the trivial fan, and this leads to the description of OT manifolds; see [9] and  $Y$  is the compact quotient of  $C_0$  by  $W$ .

(iii) When replacing the linear subspace  $H$  by the trivial space  $\{0\}$ , the rank  $b$  of  $W$  varies between 1 and  $s + t - 1$ , and we get the construction of Sankaran manifolds as in [12].

### 3. Invariants and Geometric Properties

LEMMA 3.1. *The open set  $U$  admits no nonconstant holomorphic functions.*

*Proof.* The manifold  $X$  contains the complex Lie group  $C_0$  as an open dense subset; see Corollary 1.5. Since  $U_0 = U \cap C_0$  is an open set of  $C_0$  stable under the action of the maximal compact subgroup  $(\mathbb{S}^1)^n$  of  $C_0$ , it has no nonconstant holomorphic functions. □

The following lemma will help us to compute the Kodaira dimension of our manifolds.

LEMMA 3.2. *Let  $G \cong \mathbb{C}^t$  be a closed subgroup of  $(\mathbb{C}^*)^n$ , and  $\Omega$  be an open subset of an  $n$ -dimensional toric manifold  $Z$ . Assume that the action of  $G$  on  $\Omega$  is proper and that the quotient  $\Omega/G$  admits no nonconstant holomorphic functions. Then the Kodaira dimension of  $\Omega/G$  is  $-\infty$ .*

*Proof.* Let  $\pi : \Omega \rightarrow \Omega/G$  be the quotient map. Choose a basis  $v_1, \dots, v_n$  of the Lie algebra of  $(\mathbb{C}^*)^n$  such that  $v_1, \dots, v_t$  is a basis of the Lie algebra of  $G$ . Define the vector fields  $v_1^*, \dots, v_n^*$  on  $\Omega/G$  by  $v_i^* := d\pi_y(v_i(y))$  for  $y \in \pi^{-1}(x)$  and consider the wedge product  $\tau := v_{t+1}^* \wedge \dots \wedge v_n^* \in \Gamma_{\mathcal{O}}(\Omega/G, K_{\Omega/G}^{-1})$ . Notice that  $\tau$  is nonzero on  $\pi((\mathbb{C}^*)^n \cap \Omega)$  and vanishes on  $\pi((Z \setminus (\mathbb{C}^*)^n) \cap \Omega)$ . If the Kodaira dimension of  $\Omega/G$  is not  $-\infty$ , then there is a positive integer  $k$  such that there exists a nontrivial section  $s \in \Gamma_{\mathcal{O}}(\Omega/G, K_{\Omega/G}^{\otimes k})$ . Then  $s(\tau^k)$  is a holomorphic function of  $\Omega/G$ , hence a constant equal to zero, because  $\tau^k$  only vanishes in the complement of  $\pi((\mathbb{C}^*)^n \cap \Omega)$ . □

PROPOSITION 3.3. *The Kodaira dimension of  $Y$  is  $-\infty$ .*

*Proof.* Use the previous lemma with  $G = H$ ,  $Z = X_{\Delta}$ , and  $\Omega = \pi^{-1}(U)$ , where  $\pi : X_{\Delta} \rightarrow X$  is the quotient map under the action of  $H$ . □

As a consequence of the previous lemma, we also have the following:

COROLLARY 3.4. *The Kodaira dimension of an LVMB manifold is  $-\infty$ .*

PROPOSITION 3.5. *One has the minoration  $\dim H^1(Y, \mathcal{O}) \geq b$ .*

*Proof.* Let  $\rho : W \rightarrow \mathbb{C}$  be a group homomorphism. We will associate to this homomorphism a principal  $\mathbb{C}$ -bundle above  $Y$  and show that if this bundle is trivial,

then  $\rho$  is also trivial, which will lead to the desired inequality. Consider the action of  $W$  on the product  $U \times \mathbb{C}$  given by

$$\eta.(u, z) := (\eta(u), z + \rho(\eta))$$

for all  $\eta \in W$ ,  $u \in U$ , and  $z \in \mathbb{C}$ . The quotient  $F := (U \times \mathbb{C})/W$  is a principal  $\mathbb{C}$ -bundle above  $Y$ . Indeed, the action of  $\xi \in \mathbb{C}$  on an element  $[u, z] := W.(u, z) \in F$  is given by  $\xi.[u, z] := [u, z + \xi]$ . If  $F$  is trivial, then it must have a global section, that is, there is a holomorphic function  $f : U \rightarrow \mathbb{C}$  satisfying the equality  $f(\eta(u)) = f(u) - \rho(\eta)$  for all  $\eta \in W$  and  $u \in U$ . By Lemma 3.1 the function  $f$  is constant, and this implies  $\rho \equiv 0$ .  $\square$

**COROLLARY 3.6.** *The fundamental group of the complex manifold  $Y$  is isomorphic to  $W$ , and hence  $b_1(Y) = b$ . Moreover,  $Y$  is non-Kähler.*

*Proof.* As for the fundamental group, the result follows from the fact that  $Y$  is a  $\mathbb{Z}^b$ -quotient of the simply connected complex manifold  $U$ . By the previous proposition, if  $Y$  is Kähler, then  $b_1(Y) \geq 2b$ , a contradiction.  $\square$

Recall that an element  $\alpha$  of  $\mathcal{O}_K^*$  is called a **reciprocal unit** if  $\alpha^{-1}$  is a conjugate of  $\alpha$  over  $\mathbb{Q}$ . In the following, we shall prove that the algebraic dimension of  $Y$  is zero if there is such a unit in  $W$ .

**PROPOSITION 3.7.** *If the group  $W$  contains at least one nonreciprocal unit, then  $\dim H^2(U_0/W, \mathbb{Z}) = \binom{b}{2}$ .*

*Proof.* The open set  $U_0/W$  of  $Y$  is the quotient of its contractible universal covering  $\mathbb{H}^b \times \mathbb{C}^{s+t-b}$  by the group  $W \times \mathcal{O}_K$ ; hence, we have  $H^2(U/W, \mathbb{Q}) \cong H^2(W \times \mathcal{O}_K, \mathbb{Q})$ . As in [9], we use the Lyndon–Hochschild–Serre spectral sequence associated to the short exact sequence

$$0 \rightarrow \mathcal{O}_K \rightarrow W \times \mathcal{O}_K \rightarrow W \rightarrow 0.$$

We have  $E_2^{p,q} = H^p(W, H^q(\mathcal{O}_K, \mathbb{Q})) \Rightarrow H^{p+q}(W \times \mathcal{O}_K, \mathbb{Q})$  and the following exact sequence:

$$\begin{aligned} 0 \rightarrow H^1(W, \mathbb{Q}^{\mathcal{O}_K}) \rightarrow H^1(W \times \mathcal{O}_K, \mathbb{Q}) \rightarrow \overbrace{H^1(\mathcal{O}_K, \mathbb{Q})^W}^{E_2^{0,1}} \\ \rightarrow H^2(W, \mathbb{Q}^{\mathcal{O}_K}) \rightarrow H^2(W \times \mathcal{O}_K, \mathbb{Q})_1 \rightarrow \underbrace{H^1(W, H^1(\mathcal{O}_K, \mathbb{Q}))}_{E_2^{1,1}}, \end{aligned}$$

where  $H^2(W \times \mathcal{O}_K, \mathbb{Q})_1$  is defined by the exact sequence

$$0 \rightarrow H^2(W \times \mathcal{O}_K, \mathbb{Q})_1 \rightarrow H^2(W \times \mathcal{O}_K, \mathbb{Q}) \rightarrow \underbrace{H^2(\mathcal{O}_K, \mathbb{Q})^W}_{E_2^{0,2}}.$$

See, for instance, [11].

If we prove that  $E_2^{0,1} = E_2^{0,2} = E_2^{1,1} = 0$ , then the result follows. For the fact that  $E_2^{0,1} = E_2^{1,1} = 0$ , the proof adapts readily from [9], Proposition 2.3, so we will not repeat it here.

As for the group  $E_2^{0,2} = H^2(\mathcal{O}_K, \mathbb{Q})^W \cong \text{Alt}^2(\mathcal{O}_K, \mathbb{Q})^W$ , recall that an element of  $\text{Alt}^2(\mathcal{O}_K, \mathbb{Q})^W$  is of the form  $\gamma = \sum_{i < j} a_{i,j} \sigma_i \wedge \sigma_j$  with  $a_{i,j} \in \mathbb{C}$ . Moreover, the  $W$ -invariance of  $\gamma$  means that for every pair  $(i, j)$  such that  $a_{i,j} \neq 0$ , we have  $\sigma_i(\eta)\sigma_j(\eta) = 1$  for all  $\eta \in W$ . Now  $W$  contains a nonreciprocal unit  $\eta_0$ , and therefore the relation  $\sigma_i(\eta_0)\sigma_j(\eta_0) = 1$  can never hold for any choice of  $i < j$ . Hence,  $\gamma$  is trivial, and so is the group  $E_2^{0,2}$ .  $\square$

**PROPOSITION 3.8.** *If the group  $W$  contains at least one nonreciprocal unit, then  $U_0/W$  contains no complex hypersurface; in particular, the algebraic dimension of  $Y$  is zero.*

*Proof.* We have the following commutative diagram:

$$\begin{array}{ccc}
 U_0 & \xrightarrow{W \cong \mathbb{Z}^b} & U_0/W \\
 \mathbb{R}^{s-b} \times (\mathbb{S}^1)^n \downarrow q & & \downarrow q' \mathbb{R}^{s-b} \times (\mathbb{S}^1)^n \\
 (\mathbb{R}_{>0})^b & \xrightarrow{W} & (\mathbb{S}^1)^b
 \end{array} \tag{5}$$

The open set  $U_0$  is diffeomorphic to  $(\mathbb{R}_{>0})^b \times \mathbb{R}^{s-b} \times (\mathbb{S}^1)^n$ ,  $W$  acts properly discontinuously on  $(\mathbb{R}_{>0})^b$ , and  $U_0/W$  is an  $\mathbb{R}^{s-b} \times (\mathbb{S}^1)^n$ -bundle over  $(\mathbb{S}^1)^b$ . Because  $W$  contains at least one nonreciprocal unit, the previous proposition gives that the map  $(q')^* : H^2((\mathbb{S}^1)^b, \mathbb{Z}) \rightarrow H^2(U_0/W, \mathbb{Z})$  is injective and the proof of Proposition 3.4 of [3] adapts readily. Finally,  $Y$  only has a finite number of divisors, namely those added at infinity to compactify  $U_0/W$ , and hence its algebraic dimension is equal to zero.  $\square$

**REMARK 3.9.** The second Betti number of an OT manifold of simple type was already computed in Proposition 2.3 of [9]. In fact, the proof of Proposition 3.7 shows that we can replace the simplicity condition with the assumption that  $s$  is odd.

**COROLLARY 3.10.** *Let  $X = X(K, A)$  be an OT manifold. Let  $s > 0$ , and  $2t > 0$  be the number of real and complex embeddings of  $K$ , respectively. Then, if  $s$  is odd, then  $\dim_{\mathbb{R}} H^2(X, \mathbb{R}) = \binom{s}{2}$ .*

*Proof.* Notice that this result holds with the same proof as before, under the assumption that the group  $A \cong \mathbb{Z}^s$  contains at least one nonreciprocal unit. If  $s$  is odd, then  $K$  cannot contain any reciprocal unit. Indeed if  $K$  contained such a unit, say  $\alpha$ , then the degree of  $\mathbb{Q}(\alpha)$  over  $\mathbb{Q}$  would be even and divide the degree of  $K$ .  $\square$

#### 4. An Example

To conclude this paper, we describe a concrete example of a manifold obtained by the given construction. In what follows, we continue to use the same notation.

**DEFINITION 4.1.** A **Salem number** is a real algebraic integer  $\gamma > 1$  such that all its conjugates are of modulus smaller than or equal to 1, with equality for a least one of them.

**REMARK 4.2.** By using the fact that it admits a root of modulus 1, we easily prove that the minimal polynomial  $P$  of a Salem number  $\gamma$  is palindromic, that is, it satisfies  $X^{\deg P} P(1/X) = P(X)$ . This implies that the roots of  $P$  are  $\gamma$ ,  $1/\gamma$ , and complex numbers of absolute value 1. In particular, the minimal polynomial of a Salem number has degree at least 4, and a Salem number is necessarily a unit of even degree.

**EXAMPLE 4.3.** (See [4], p. 85) It is possible to describe all polynomials of degree 4 that are minimal polynomials of a Salem number. These are the polynomials with integer coefficients of the form  $X^4 + q_1 X^3 + q_2 X^2 + q_1 X + 1$  with  $2(q_1 - 1) < q_2 < -2(q_1 + 1)$ . The minimal polynomial of the smallest Salem number of degree 4 is  $X^4 - X^3 - X^2 - X + 1$ .

Now we consider the polynomial  $P(X) := X^4 - X^3 - X^2 - X + 1$ . Its roots are  $\alpha \approx 1,722$  (truncated value),  $\alpha^{-1}$ ,  $\beta$ , and  $\bar{\beta}$ , where  $\beta$  is a complex number of modulus 1, and  $\Im(\beta) > 0$ . Note  $\sigma_1, \sigma_2, \tau_1$ , and  $\bar{\tau}_1$  the associated embeddings of the field  $K := \mathbb{Q}[X]/\langle P \rangle$  in  $\mathbb{R}$  (for the first two ones) and  $\mathbb{C}$  (for the last two ones).

We can check by computation that the family  $(1, \alpha, \alpha^2, \alpha^3)$  forms an integral basis of  $K$ ; see Section 2.6 of [13] for a method. The image under the map  $\sigma_K$  of this family is the following basis of  $\mathbb{C}^4$ :

$$\mathcal{B}_K = \left( \begin{pmatrix} 1 \\ 1 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \alpha \\ \alpha^{-1} \\ \beta \\ \bar{\beta} \end{pmatrix}, \begin{pmatrix} \alpha^2 \\ \alpha^{-2} \\ \beta^2 \\ \bar{\beta}^2 \end{pmatrix}, \begin{pmatrix} \alpha^3 \\ \alpha^{-3} \\ \beta^3 \\ \bar{\beta}^3 \end{pmatrix} \right).$$

Moreover,  $\alpha$  and  $1 - \alpha$  are two fundamental units of  $\mathcal{O}_K^*$ . Since  $s = 2$ , we have  $\Omega = N \times L_+$  with  $h = \dim N = 1 = \dim L = b$  (see Lemma 1.13 for notations).

It is clear that the two subgroups  $W := \langle \alpha \rangle_{\mathbb{Z}}$  and  $\bar{W} := \langle 1 - \alpha \rangle_{\mathbb{Z}}$  of  $\mathcal{O}_K^{*,+}$  satisfy Assumption C and they both are of rank  $b = 1$ .

The action of  $W$  on  $\mathbb{C}^4$  is given by the following diagonal matrix:

$$M := \begin{pmatrix} \alpha & 0 & 0 & 0 \\ 0 & \alpha^{-1} & 0 & 0 \\ 0 & 0 & \beta & 0 \\ 0 & 0 & 0 & \bar{\beta} \end{pmatrix}.$$

In the basis  $\mathcal{B}_K$ , the matrix of the linear map associated to  $M$  is the companion matrix of  $P$ :

$$C := \begin{pmatrix} 0 & 0 & 0 & -1 \\ 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Here,  $H$  is the subgroup  $\{(0, 0, 0, z) \in \mathbb{C}^4 \mid z \in \mathbb{C}\}$  of  $\mathbb{C}^4$ . In the basis  $\mathcal{B}_K$ ,  $H$  is the set  $\{(-\beta z, \beta(1 - \beta)z, (\bar{\beta} - 1)z, z) \mid z \in \mathbb{C}\}$ , and the embedding  $\iota : H \rightarrow (\mathbb{C}^*)^4$  is given by

$$\iota(z) = (e^{-2i\pi\beta z}, e^{2i\pi\beta(1-\beta)z}, e^{2i\pi(\bar{\beta}-1)z}, e^{2i\pi z}).$$

The cone  $\Omega \subset \mathbb{R}^2$  is the open half-plane  $\mathbb{R}_{>0} \times \mathbb{R}$ . We define a fan of  $\mathbb{R}^2$  whose cones are generated by elements of  $\pi_{\tilde{H}}(\mathcal{O}_K)$  and which is invariant under the action of  $W$  as follows.

Let  $\sigma_1$  be the cone positively generated by the vectors  $(1, 1)$  and  $(\alpha, \alpha^{-1})$ ,  $\sigma_2$  be the cone generated by  $(1, -1)$  and  $(\alpha, -\alpha^{-1})$ , and let  $\Delta'$  be the fan generated by  $W \cdot \{\sigma_1, \sigma_2\}$ . It is clear that  $(\Omega \setminus (\mathbb{R}_{>0} \times \{0\})) \cup \{0\}$  is the support of  $\Delta'$ . For a picture of the situation in  $\mathbb{R}^2$ , we refer to Figure 1.

In this example, the divisors that we add to  $U_0$  so that its quotient by  $W$  is compact are chains of Hopf surfaces.

Indeed, let  $\sigma \in \Delta \subset E \cong \mathbb{R}^4$  be a one-dimensional cone, say  $\sigma$  is generated by  $(\alpha, \alpha^{-1}, \beta, \bar{\beta})$  for simplicity. By the first remark of Section 2.3 the quotient of  $\text{orb}(\sigma)$  by  $H$  is a generalized LVMB manifold. The fan  $\Delta/\sigma$  of  $\mathbb{R}^3$  is generated by the images of the two two-dimensional cones containing  $\sigma$  under  $\pi_\sigma : \mathbb{R}^4 \rightarrow \mathbb{R}^4/\text{span}(\sigma)$ . Up to a linear isomorphism, we can assume that this is a subfan of the fan of  $\mathbb{P}^3(\mathbb{C})$ . In other words,  $\text{orb}(\sigma)/H$  is an LVMB manifold. If we adopt the notation of [7], this situation corresponds to the case  $n = 4, k = 3, n_1 = n_2 = 1$ , and  $n_3 = 2$ ; hence,  $\text{orb}(\sigma)/H$  is a Hopf surface (Sect. 4(b), p. 260, *ibid.*).

Also, notice that  $\text{orb}(\sigma)/H$  is necessarily a Hopf surface by Potters' theorem on the classification of almost homogeneous complex compact surfaces [10].

Hence, we have built a complex manifold of dimension 3 that consists of a dense open set that is the quotient of a Cousin group by a discrete group compactified by two cycles of Hopf surfaces. A computation shows that each of these Hopf surfaces contains the two elliptic curves  $\mathbb{C}/\langle 1, \beta \rangle$  and  $\mathbb{C}/\langle 1, 1 - \bar{\beta} \rangle$ . Now since  $1 - \bar{\beta} = (\beta - 1)/\beta$ , these two elliptic curves are isomorphic.

However, because  $W$  only contains reciprocal units, we cannot apply Proposition 3.8, and hence we do not know yet the algebraic dimension of this manifold. On the other hand, if we carry the construction with the group  $\bar{W}$  instead of  $W$ , then we know that the resulting manifold has its algebraic dimension equal to zero.

ACKNOWLEDGMENTS. The first author would like to thank Prof. A. T. Huckleberry and Prof. P. Heinzner for invitation as a postdoc to Bochum's Ruhr-Universität.

## References

- [1] A. Ash, D. Mumford, M. Rapoport, and Y.-S. Tai, *Smooth compactifications of locally symmetric varieties*, 2nd edition, Cambridge Math. Lib., Cambridge University Press, Cambridge, 2010.
- [2] L. Battisti, *LVMB manifolds and quotients of toric varieties*, Math. Z. 275 (2013), no. 1, 549–568.
- [3] L. Battisti and K. Oeljeklaus, *Holomorphic line bundles over domains in Cousin groups and the algebraic dimension of OT-manifolds*, Proc. Edinb. Math. Soc. (2013, to appear), [arXiv:1306.3047v1](https://arxiv.org/abs/1306.3047v1).
- [4] M.-J. Bertin, A. Decomps-Guilloux, M. Grandet-Hugot, M. Pathiaux-Delefosse, and J.-P. Schreiber, *Pisot and Salem numbers*, Birkhäuser Verlag, Basel, 1992.
- [5] F. Bosio, *Variétés complexes compactes: une généralisation de la construction de Meersseman et López de Medrano-Verjovsky*, Ann. Inst. Fourier (Grenoble) 51 (2001), no. 5, 1259–1297.
- [6] G. Ewald, *Combinatorial convexity and algebraic geometry*, Grad. Texts in Math., 168, Springer-Verlag, New York, 1996.
- [7] S. López de Medrano and A. Verjovsky, *A new family of complex, compact, non-symplectic manifolds*, Bull. Braz. Math. Soc. (N.S.) 28 (1997), no. 2, 253–269.
- [8] L. Meersseman, *A new geometric construction of compact complex manifolds in any dimension*, Math. Ann. 317 (2000), no. 1, 79–115.
- [9] K. Oeljeklaus and M. Toma, *Non-Kähler compact complex manifolds associated to number fields*, Ann. Inst. Fourier (Grenoble) 55 (2005), no. 1, 161–171.
- [10] J. Potters, *On almost homogeneous compact complex analytic surfaces*, Invent. Math. 8 (1969), 244–266.
- [11] C. H. Sah, *Cohomology of split group extensions*, J. Algebra 29 (1974), 255–302.
- [12] G. K. Sankaran, *A class of non-Kähler complex manifolds*, Tohoku Math. J. (2) 41 (1989), no. 1, 43–64.
- [13] I. Stewart and D. Tall, *Algebraic number theory and Fermat’s last theorem*, 3rd edition, A K Peters, Ltd., Natick, MA, 2002.

L. Battisti  
 Fakultät für Mathematik  
 Ruhr-Universität Bochum  
 D-44780 Bochum  
 Germany

[laurent.battisti@yahoo.com](mailto:laurent.battisti@yahoo.com)  
[laurent.battisti@rub.de](mailto:laurent.battisti@rub.de)

K. Oeljeklaus  
 CNRS, Centrale Marseille, I2M,  
 UMR 7373  
 Aix-Marseille Université  
 CMI, 39, Rue F. Joliot-Curie  
 F-13453 Marseille  
 France

[karl.oeljeklaus@univ-amu.fr](mailto:karl.oeljeklaus@univ-amu.fr)