

SHAPE VIA MULTI-NETS

By

Zvonko ČERIN

Abstract. We give in this paper a description of a new category related to shape category. We consider families of multi-valued functions between topological spaces which we call multi-nets. In a well-controlled way functions of a multi-net more and more resemble single-valued functions. We introduce a notion of homotopy for multi-nets and a composition of homotopy classes. The resultant homotopy category of multi-nets \mathcal{HM} is naturally equivalent to the shape category provided we restrict to spaces which have ANR-resolutions with onto projections. However, the homotopy category of multi-nets is interesting because it provides an intrinsic method of studying global properties of spaces. Our idea is to extend Borsuk's approach based on fundamental sequences to arbitrary topological spaces in analogy with Sanjurjo's description of shape category of compact metric spaces in terms of upper semi-continuous multi-valued functions.

Introduction

The subject of this paper belongs to the part of geometric topology which is known under the name shape theory. The method of our investigations is through the use of multi-valued functions. Our motivation is a desire to get a new description of the shape category which will be an extension to arbitrary topological spaces of Sanjurjo's approach to shape theory of compact metric spaces via upper semi-continuous multi-valued functions [10].

The classical homotopy theory studies the equivalence relation of homotopy for maps. Recall that maps (i. e., continuous single-valued functions) f and g between topological spaces X and Y are called homotopic provided there is a map h from the product $X \times I$ of X with the unit closed segment $I = [0, 1]$ into

1980 *Mathematics, Subject Classification* (1985 *Revision*). Primary 54B25, 54F45, 54C56.
Key words and phrases. homotopy, shape theory, multi-valued function, approximate resolution, expansion, σ -close, σ -small.

Received July 29, 1993.

Y such that $h(x, 0)=f(x)$ and $h(x, 1)=g(x)$ for every $x \in X$. The homotopy category \mathcal{H} has as objects topological spaces and as morphisms homotopy classes of maps. The homotopy classes are composed by composing representatives and the identity morphisms are homotopy classes of the identity maps.

The equivalence relation of homotopy for maps leads to a useful and rich theory only when we restrict to spaces with nice local properties like polyhedra and absolute neighbourhood retracts. The problems arise in the definition above when the space Y is such that there are not many maps from $X \times I$ into Y so that the properties of Y are preventing identification of maps which ought to be identified. In other words, the definition of homotopy is too rigid because the function h must be continuous and single-valued and because it must take values in the space Y .

This has led K. Borsuk to modify homotopy theory so that the new theory which he calls shape theory agrees with the old on absolute neighbourhood retracts and that it gives much better results for spaces with complicated local structure where the old theory is inadequate. The modification of Borsuk relies on the idea to relinquish the insistence in the definition of homotopy that the map h goes precisely into the space Y . The obvious alternative method which was undertaken by Sanjurjo in [9] and [10] and further followed in this paper is to give up with the requirement that the function h is continuous and/or single-valued while retaining the desirable condition that it takes values in the space Y . In order to properly honour these two diverse methods we shall call them the Borsuk approach and the Sanjurjo approach to shape theory. We use names outer shape theory and inner shape theory.

In the original Borsuk's description [2] of shape category Sh_B of compact metric spaces, the spaces X and Y are considered as closed subsets of the Hilbert cube Q and maps from X into Y are replaced with fundamental sequences. Recall that a fundamental sequence φ from X into Y is a sequence $\{\varphi_i\}_{i=1}^{\infty}$ of maps $\varphi_i: Q \rightarrow Q$ such that for every neighbourhood U of Y in Q there is a neighbourhood V of X in Q and an index i with the property that the restrictions $\varphi_i|_V$ and $\varphi_j|_V$ are homotopic in U for every $j > i$. The role of the homotopy relation plays the following notion. Fundamental sequences φ and ψ from X into Y are called homotopic provided for every neighbourhood U of Y in Q there is a neighbourhood V of X in Q and an index i with the property that the restrictions $\varphi_j|_V$ and $\psi_j|_V$ are homotopic in U for every $j \geq i$. This is an equivalence relation $[\varphi]$ denotes the homotopy class of a fundamental sequence φ , and homotopy classes are composed by the rule $[\psi] \circ [\varphi] = [\psi \circ \varphi]$, where $\psi \circ \varphi$ is a fundamental sequence formed by compositions $\psi_i \circ \varphi_i$. The category Sh_B has

compact metric spaces as objects and homotopy classes of fundamental sequences as morphisms.

In spite of its simplicity and clear geometric flavour, Borsuk's theory relies too much on the Hilbert cube and the use of open neighbourhoods of subsets so that the extension of it to wider classes of spaces proved to be a formidable problem. This was accomplished by many authors so that now we have different descriptions of shape category Sh . Its objects are topological spaces while its morphisms are rather awkward constructions involving things such as Morita's ANR-expansions, Grothendick's pro-categories, and intricate concepts of category theory (see [7]). All these efforts belong to the outer shape theory because they use some outside objects in order to study global properties of spaces. In particular, these approaches all require the use of absolute neighbourhood retracts. In this paper we propose to follow for arbitrary topological spaces Borsuk's geometric method based on fundamental sequences as closely as possible without any reference to absolute neighbourhood retracts.

Instead of fundamental sequences we consider multi-nets. The other steps are identical. We define a notion of homotopy for multi-nets and the morphisms are simply homotopy classes of multi-nets. This idea has previously been used by Sanjurjo in [9] and [10] to get an analogous description of Sh_B . The crux of this approach is to use functions which are not continuous and/or single-valued. Our investigation started with attempts to extend Sanjurjo's method to arbitrary topological spaces.

The difference in approach is that we do not require multi-valued functions to be upper semi-continuous though it is possible with only minor modifications to build up the appropriate category where this requirement is fulfilled.

The key tools are given as Lemmas 2 and 3 which provide replacement of a small multi-valued functions (as defined in Definition 2) by a close (see Definition 3) continuous single-valued function and necessary transitivity of the notion of small homotopy (from Definition 4).

The multi-nets and their homotopy is given in Definitions 5 and 6. The most difficult part is to find the correct notion of composition for homotopy classes of multi-nets. This is accomplished in the first three claims and summarized in Theorem 2.

With the description of the new category \mathcal{HM} thus completed, the rest of the paper deals with setting up a functor θ from our homotopy category of multi-nets into the shape category (see Theorem 3 and Claims 4-6).

Finally, in Theorem 4 and Claims 7-10, we show that the homotopy category of multi-nets is naturally equivalent to the shape category on spaces hav-

ing ANR-resolutions with onto projections. This is done by describing the inverse ζ of the functor θ .

The present paper is only the first in a series where we shall attempt to do large portions of inner shape theory using small multi-valued functions. This approach is particularly suitable for some problems. It's obvious merit is that it does not need any outside objects (like a nice ambient space or an inverse limit expansion into nice spaces). In conclusion, this paper lies foundations for the study of the homotopy category \mathcal{HM} of multi-nets and establishes some connections between \mathcal{HM} and the shape category. In the paper "Shape theory intrinsically" we shall prove by far more complicated arguments that the categories \mathcal{HM} and \mathcal{Sh} are equivalent.

Small Multi-valued functions

In this section we shall introduce notions that are required for our theory and prove two useful technical results.

Let \hat{Y} denote the collection of all normal covers of a topological space Y [1]. With respect to the refinement relation $>$ the set \hat{Y} is a directed set. Two normal covers σ and τ of Y are equivalent provided $\sigma > \tau$ and $\tau < \sigma$. In order to simplify our notation we denote a normal cover and it's equivalence class by the same symbol. Consequently, \hat{Y} also stands for the associated quotient set.

Let \check{Y} denote the collection of all finite subsets c of \hat{Y} which have a unique (with respect to the refinement relation) maximal element $\check{c} \in \hat{Y}$. We consider \check{Y} ordered by the inclusion relation and regard \hat{Y} as a subset of single-element subsets of \check{Y} . Notice that \check{Y} is a cofinite directed set [7, p. 11].

We shall repeatedly use the following lemma (see [7, p. 9]).

LEMMA 1. *Let $\{f_1, \dots, f_n\}$ be a finite collection of functions from a cofinite directed set $(M, <)$ into a directed set $(L, <)$. Then there is an increasing function $g: M \rightarrow L$ such that $g(x) > f_1(x), \dots, f_n(x)$ for every $x \in M$.*

The next two definitions introduce precisely a type of multi-valued functions that we shall use.

DEFINITION 1. Let X and Y be topological spaces. By a *multi-valued function* or an *M-function* $F: X \rightarrow Y$ we mean a rule which associates a non-empty subset $F(x)$ of Y to every point x of X . Let $M(X, Y)$ denote all *M-functions* from X into Y .

DEFINITION 2. Let $F: X \rightarrow Y$ be a multi-valued function and let $\alpha \in \hat{X}$ and $\gamma \in \hat{Y}$. We shall say that F is an (α, γ) -map provided for every $A \in \alpha$ there is a $C_A \in \gamma$ with $F(A) \subset C_A$. On the other hand, F is γ -small provided there is an $\alpha \in \hat{X}$ such that F is an (α, γ) -map.

The following is a notion of closeness for multi-valued functions that is needed in this approach to shape theory.

DEFINITION 3. Let $F, G: X \rightarrow Y$ be multi-valued functions and let $\gamma \in \hat{Y}$. We shall say that F and G are γ -close and we write $F \stackrel{\gamma}{=} G$ provided for every $x \in X$ there is a $C_x \in \gamma$ with $F(x) \cup G(x) \subset C_x$.

The following lemma is not needed in the description of the category \mathcal{AM} but only in setting up a functor θ from the category \mathcal{AM} into the shape category Sh . This is a very useful approximation result which shows that a sufficiently small multi-valued functions into an approximate polyhedron can be replaced by a continuous single-valued function.

Recall [7] that an *approximate polyhedron* is a topological space Y with the property that for every $\sigma \in \hat{Y}$ there is a polyhedron P and maps $u: Y \rightarrow P$ and $d: P \rightarrow Y$ with $id_Y \stackrel{\sigma}{=} d \circ u$.

LEMMA 2. For every normal cover σ of an approximate polyhedron Y there is a normal cover τ of Y such that every τ -small multi-valued function $F: X \rightarrow Y$ from a topological space X into Y there is a normal cover ρ of X with the property that for every canonical map $p: X \rightarrow N(\rho)$ from X into the nerve $N(\rho)$ of ρ there is a single-valued continuous function $f: N(\rho) \rightarrow Y$ with $F \stackrel{\sigma}{=} f \circ p$.

PROOF OF LEMMA 2. Let $\lambda \in \sigma^*$ and $\nu \in \lambda^*$, where σ^* denotes the set of all normal covers τ of Y such that the star $st(\tau)$ of τ refines σ . Choose a simplicial polytope P with the metric topology and maps $u: Y \rightarrow P$ and $d: P \rightarrow Y$ with

$$(1) \quad id_Y \stackrel{\nu}{=} d \circ u.$$

Let $\varepsilon = d^{-1}(\nu) \in \hat{P}$. Let $\eta \in \varepsilon^*$. Since P is an ANR [6, p. 106], there is a refinement π of η with the property that every partial realization in P of a simplicial polytope K with the Whitehead topology relative to π defined on a subpolytope L of K which contains all vertices of K extends to a full realization of K in P relative to ν [6, p. 122]. Let $\xi \in \pi^*$ and let $\tau \in \hat{Y}$ be a common refinement of ν and $u^{-1}(\xi)$.

Consider a τ -small multi-valued function $F: X \rightarrow Y$. Choose a $\beta \in \hat{X}$ such that F is a (β, τ) -map. Let $\{\lambda_B | B \in \beta\}$ be a partition of unity subordinated to β , and let $\{\mu_B | B \in \beta\}$ be its locally finite improvement [4, p. 354]. Let $\rho = \{\mu_B^{-1}((0, 1]) | B \in \beta\}$. Hence, for every $R \in \rho$ there is a $T_R \in \tau$, an $N_R \in \nu$, a $K_R \in \xi$, $y_R \in Y$ and a $z_R \in P$ with $F(R) \subset T_R$, $T_R \subset N_R$, $u(T_R) \subset K_R$, $y_R \in T_R$, $z_R \in K_R$ and $z_R = u(y_R)$. Let $p: X \rightarrow N(\rho)$ be a canonical map of X into the nerve $N(\rho)$ of ρ (see [4]).

Define a function $\varphi: N(\rho)^0 \rightarrow P$ by the rule $\varphi(R) = z_R$ for every $R \in \rho$. This function is continuous and it provides a partial realization of $N(\rho)$ in P relative to the cover π .

Indeed, let $\delta = \langle A, B, \dots, Z \rangle$ be a simplex of $N(\rho)$. We shall find a member of π which contains the set $\varphi(N(\rho)^0 \cap \delta)$, i. e., the set $\{z_A, \dots, z_Z\}$. Suppose $x \in A \cap \dots \cap Z$. Since $F(x)$ is non-empty, the sets T_A, \dots, T_Z and therefore also the sets K_A, \dots, K_Z have non-empty intersection. Since ξ is a star-refinement of π , it is clear that some member P_δ of π contains their union.

Let $\psi: N(\rho) \rightarrow Y$ be a full realization in P of $N(\rho)$ relative to η . Let f denote the composition $d \circ \psi$. Then f is the required continuous single-valued function.

Let $x \in X$ and suppose that A, \dots, Z are all members of ρ which contain the point x . Then $p(x)$ lies in the simplex δ of $N(\rho)$ determined by these sets. It follows that a member E_x of η contains both $\psi \circ p(x)$ and points z_A, \dots, z_Z . Since ξ refines η and η is a star-refinement of ε , there is a member N_x of ν with $d(E_x \cup K_A \cup \dots \cup K_Z) \subset N_x$. On the other hand, from (1) we get the existence of members N_A, \dots, N_Z of ν such that N_C contains both y_C and $d(z_C)$ for every $C = A, \dots, Z$. It follows that

$$f \circ p(x) \in N_x \quad d(z_A) \in N_x \cap N_A, \quad y_A \in N_A \cap T_A, \quad F(x) \subset T_A.$$

Hence, some member of σ contains both $f \circ p(x)$ and $F(x)$. \square

The following definition is the most important for this paper and our approach to inner shape theory.

DEFINITION 4. Let $F, G: X \rightarrow Y$ be multi-valued functions between topological spaces and let γ be a normal cover of the space Y . We shall say that F and G are γ -homotopic and write $F \stackrel{\gamma}{\cong} G$ provided there is a γ -small multi-valued function H from the product $X \times I$ of X and the unit segment $I = [0, 1]$ into Y such that $F(x) \subset H(x, 0)$ and $G(x) \subset H(x, 1)$ for every $x \in X$. We shall say that H is a γ -homotopy that joins F and G or that it realizes the relation

(or homotopy) $F \stackrel{\gamma}{\cong} G$.

The following lemma gives an adequate substitute for transitivity of the homotopy relation for maps. It will be used later many times.

LEMMA 3. *Let $F, G, H: X \rightarrow Y$ be multi-valued functions. Let $\sigma \in \hat{Y}$ and $\tau \in \sigma^*$. If $F \stackrel{\tau}{\cong} G$ and $G \stackrel{\tau}{\cong} H$, then $F \stackrel{\sigma}{\cong} H$.*

PROOF OF LEMMA 3. Let $K, L: X \times I \rightarrow Y$ be τ -small multi-valued functions such that

$$(2) \quad F(x) \subset K(x, 0), \quad G(x) \subset K(x, 1) \cap L(x, 0), \quad H(x) \subset L(x, 1)$$

for every $x \in X$. Define $M: X \times I \rightarrow Y$ by

$$M(x, t) = \begin{cases} K(x, 2t), & x \in X, 0 \leq t < 1/2 \\ K(x, 1) \cup L(x, 0), & x \in X, t = 1/2 \\ L(x, 2t-1), & x \in X, 1/2 < t \leq 1. \end{cases}$$

Clearly, by (2), $F(x) \subset M(x, 0)$ and $H(x) \subset M(x, 1)$ for every $x \in X$. Hence, it remains to see that M is σ -small.

Since both K and L are τ -small, there are normal covers α and β of $X \times I$ so that for every $A \in \alpha$ there is a $T_A^K \in \tau$ with $K(A) \subset T_A^K$ and for every $B \in \beta$ there is a $T_B^L \in \tau$ with $L(B) \subset T_B^L$. Let a normal cover γ be a common refinement of α and β . Then for every $C \in \gamma$ we can find $T(C), W(C) \in \tau$ with

$$(3) \quad K(C) \subset T(C) \quad \text{and} \quad L(C) \subset W(C).$$

We now use [4, p. 358], to get a normal cover $\varepsilon \in \hat{X}$ together with the function $r: \varepsilon \rightarrow \{2, 3, 4, \dots\}$ such that every set $E \times [t_{2i}, t_{2i+4}]$ is contained in a member $C_{E,i}$ of γ , where $E \in \varepsilon$, $i=0, 1, 2, \dots, rE-2$, and $t_j = j/4rE$ for every $j=0, 1, \dots, 4rE$.

We define for each $E \in \varepsilon$ an open over $|E|$ of I as follows:

$$|E| = \{V_1, V_2, \dots, V_{4rE-1}\},$$

where $V_1 = [0, t_2)$, $V_2 = (t_1, t_3)$, $V_3 = (t_2, t_4)$, \dots , $V_{4rE-1} = (t_{4rE-2}, 1]$.

Since $\{\{E \times V \mid V \in |E|\} \mid E \in \varepsilon\}$ is a normal cover of $X \times I$, our proof will be completed provided we show that for every $E \in \varepsilon$ and every $V \in |E|$ there is a member of σ containing $M(E \times V)$.

If $V = V_i$, for $i \neq 2rE$, this follows from (3).

Let $V = V_{2rE}$. Then $M(E \times V) = K(E \times (t_{4rE-2}, 1]) \cup L(E \times [0, t_2))$. But, $K(E \times (t_{4rE-2}, 1]) \subset T(C_{E, 2rE-1})$ and $L(E \times [0, t_2)) \subset W(C_{E, 0})$. As E is a non-empty set,

there is an $x \in E$. The relation (2) shows that the non-empty set $G(x)$ lies in the intersection of sets $T(C_{E, 2rE-1})$ and $W(C_{E, 0})$. Hence, a member of σ contains $M(E \times V)$. \square

Multi-nets

The following two definitions correspond to Borsuk's definitions of fundamental sequences and homotopy for fundamental sequences.

DEFINITION 5. Let X and Y be topological spaces. By a *multi-net* or an *M-net* from X into Y we shall mean a collection $\varphi = \{F_c \mid c \in \tilde{Y}\}$ of multi-valued functions $F_c: X \rightarrow Y$ such that for every $\gamma \in \hat{Y}$ there is a $c \in \tilde{Y}$ with $F_d \stackrel{\gamma}{\cong} F_c$ for every $d > c$. We use functional notation $\varphi: X \rightarrow Y$ to indicate that φ is a multi-net from X into Y . Let $MN(X, Y)$ denote all multi-nets $\varphi: X \rightarrow Y$.

DEFINITION 6. Two multi-nets $\varphi = \{F_c\}$ and $\psi = \{G_c\}$ between topological spaces X and Y are *homotopic* provided for every $\gamma \in \hat{Y}$ there is a $c \in \tilde{Y}$ such that $F_d \stackrel{\gamma}{\cong} G_d$ for every $d > c$.

It follows from Lemma 3 that the relation of homotopy is an equivalence relation on the set $MN(X, Y)$. The homotopy class of a multi-net φ is denoted by $[\varphi]$ and the set of all homotopy classes by $\mathcal{HM}(X, Y)$.

Our first goal is to define a composition for homotopy classes of multi-nets and to establish its associativity.

Let $\varphi = \{F_c\}: X \rightarrow Y$ be a multi-net. For every $c \in \tilde{Y}$ there is an $\bar{f}(c) \in \tilde{Y}$ such that for all $d, e > \bar{f}(c)$ there is a normal cover $\bar{f}(c, d, e)$ of $X \times I$ and an $(\bar{f}(c, d, e), \tilde{c})$ -map joining F_d and F_e .

Let $C = \{(c, d, e) \mid c \in \tilde{Y}, d, e > \bar{f}(c)\}$. Then C is a subset of $\tilde{Y} \times \tilde{Y} \times \tilde{Y}$ that becomes a cofinite directed set when we define that $(c, d, e) > (c', d', e')$ iff $c > c', d > d'$ and $e > e'$.

Now, let $f: \tilde{Y} \rightarrow \tilde{Y}$ be an increasing function such that $f(c) > \bar{f}(c)$, c for every $c \in \tilde{Y}$. We shall use the same notation f for an increasing function $f: C \rightarrow \widehat{X \times I}$ such that $f(c, d, e) > \bar{f}(c, d, e)$ for every $(c, d, e) \in C$. Let $(c, d, e) \in C$. For the normal cover $f(c, d, e)$ of $X \times I$, by [4, p. 358], there is a normal cover $\varepsilon = \hat{f}(c, d, e)$ of X and a function $r = \check{f}(c, d, e): \varepsilon \rightarrow \{2, 3, 4, \dots\}$ such that every set $E \times [(i-1)/rE, (i+1)/rE]$, where $E \in \varepsilon$ and $i = 1, 2, \dots, rE-1$, is contained in a member of $f(c, d, e)$.

Let $\tilde{f}: C \rightarrow \hat{X}$ be an increasing function with $\tilde{f}(c, d, e) > \hat{f}(c, d, e)$ for every

$(c, d, e) \in \mathcal{C}$. We shall use the shorter notation $\tilde{f}(c)$ and $f(c)$ for the covers $\tilde{f}(c, f(c), f(c))$ and $f(c, f(c), f(c))$.

CLAIM 1. *There is an increasing function $f^*: \tilde{Y} \rightarrow \hat{X}$ such that*

- (1) $f^*(c) > \tilde{f}(c)$ for every $c \in \tilde{Y}$, and
- (2) f^* is cofinal in \tilde{f} , i.e., for every $(c, d, e) \in \mathcal{C}$ there is an $m \in \tilde{Y}$ with $f^*(m) > \tilde{f}(c, d, e)$.

PROOF OF CLAIM 1. Let $\mathcal{D} = \{\tilde{f}(c, d, e) \mid (c, d, e) \in \mathcal{C}\}$.

If \tilde{Y} is a finite set, then \mathcal{D} is a finite collection of elements of \hat{X} . Let $\alpha \in \hat{X}$ be a common refinement of all members of \mathcal{D} . Let $f^*: \tilde{Y} \rightarrow \hat{X}$ be a constant function into α .

If \tilde{Y} is an infinite set, then the cardinality of \mathcal{D} does not exceed the cardinality of \tilde{Y} . Hence, there is a surjection $g: \tilde{Y} \rightarrow \mathcal{D}$. Let $f^*: \tilde{Y} \rightarrow \hat{X}$ be an increasing function such that $f^*(c) > g(c), \tilde{f}(c)$ for every $c \in \tilde{Y}$. \square

The above discussion shows that every multi-net $\varphi: X \rightarrow Y$ determines eight functions denoted by $\bar{f}, f, \hat{f}, \tilde{f}$ and f^* . With the help of these functions we shall define the composition of homotopy classes of multi-nets as follows.

Let $\varphi = \{F_c\}: X \rightarrow Y$ and $\psi = \{G_s\}: Y \rightarrow Z$ be multi-nets. Let $\lambda = \{H_s\}$, where $H_s = G_{g(s)} \circ F_{f(g^*(s))}$ for every $s \in \tilde{Z}$.

CLAIM 2. *The collection λ is a multi-net from X into Z .*

PROOF OF CLAIM 2. Let $\sigma \in \hat{Z}$. We must find an $s \in \tilde{Z}$ such that

$$(13) \quad H_t \overset{\sigma}{\cong} H_s \quad \text{for every } t > s.$$

Let $\tau \in \sigma^{*2}$, where σ^{*n} denotes the set of all normal covers τ of Z such that the n -th star $st^n(\tau)$ of τ refines σ . Let $s = \{\tau\} \in \tilde{Z}$.

Consider an index $t > s$. We shall find an index $c \in \tilde{Y}$ so that

$$(14) \quad H_t \overset{\tau}{\cong} G_x \circ F_c,$$

$$(15) \quad G_x \circ F_c \overset{\tau}{\cong} G_y \circ F_c,$$

and

$$(16) \quad G_y \circ F_c \overset{\tau}{\cong} H_s,$$

where $x = g(t)$ and $y = g(s)$. Repeated use of Lemma 3 will give (13) from the relations (14)–(16).

Invoking the property (2) of Claim 1, choose a $u > t$ so that $r > \tilde{g}(s, p, q)$, where $r = g^*(u)$, $p = g(s)$, and $q = g(t)$. Let $c = f(r)$. Since $q > p > \bar{g}(s)$ and $\tilde{s} = \tau$, there is a $(g(s, p, q), \tau)$ -map $L : Y \times I \rightarrow Z$ joining G_p and G_q . But, F_c is joined to itself by an r -small homotopy. It follows that $L \circ (F_c \times id_I)$ is a τ -small homotopy realizing the relation (15).

On the other hand, G_x is a $(\tilde{g}(t), \tau)$ -map while $F_{f(g^*(t))}$ and F_c are joined by a $g^*(t)$ -small homotopy K . The property (1) of Claim 1 implies that $G_{g(t)} \circ K$ is a τ -small homotopy which realizes the relation (14). In an analogous fashion one can show that (16) is also true. \square

We now define the composition of homotopy classes of multi-nets by the rule $[\{G_s\}] \circ [\{F_c\}] = [\{G_{g(s)} \circ F_{f(g^*(s))}\}]$.

CLAIM 3. *The composition of homotopy classes of multi-nets is well-defined.*

PROOF OF CLAIM 3. Let $\kappa = \{K_c\}$ and $\lambda = \{L_s\}$ be multi-nets homotopic to φ and ψ , respectively, and let $\mu = \{M_s\}$, where $M_s = L_{l(s)} \circ K_{k(l^*(s))}$ for every $s \in \tilde{Z}$. We must show that multi-nets χ and μ are homotopic. In other words, that for every $\sigma \in \hat{Z}$ there is an $s \in \tilde{Z}$ such that

$$(17) \quad H_t \stackrel{\tau}{\cong} M_t \quad \text{for every } t > s.$$

Let $\sigma \in \hat{Z}$. Let $\tau \in \sigma^{*4}$. Let $s = \{\tau\} \in \tilde{Z}$. In order to prove (17), we shall argue that for every $t > s$ we can find indices $c \in \tilde{Y}$ and $u \in \tilde{Z}$ such that

$$(18) \quad H_t \stackrel{\tau}{\cong} G_x \circ F_c,$$

$$(19) \quad G_x \circ F_c \stackrel{\tau}{\cong} G_u \circ F_c,$$

$$(20) \quad G_u \circ F_c \stackrel{\tau}{\cong} L_u \circ F_c,$$

$$(21) \quad L_u \circ F_c \stackrel{\tau}{\cong} L_u \circ K_c,$$

$$(22) \quad L_u \circ K_c \stackrel{\tau}{\cong} L_y \circ K_c,$$

$$(23) \quad L_y \circ K_c \stackrel{\tau}{\cong} M_t,$$

where we put $x = g(t)$ and $y = l(t)$. From the relations (18)–(23) with the help of Lemma 3 we shall get (17).

We shall now describe how big c and u must be chosen for relations (18), (19), (20) and (21) to hold separately. The relations (22) and (23) are analogous to relations (19) and (18), respectively. We leave to the reader the task of making a cumulative choice for c and u which accomplishes our goal. It is

important to notice that u is selected first while c is selected only once u is already known.

Add (18). Since G_x is a $(\tilde{g}(t), \tau)$ -map and $g^*(t)$ refines the cover $\tilde{g}(t)$, by the property (1) of Claim 1, it suffices to take $c > f(g^*(t))$.

Add (19). If $u > x$, then G_x and G_u are joined by a $(g(u), \tau)$ -map $P: Y \times I \rightarrow Z$. Let $c > f(g^*(t))$. Then F_c is $g^*(u)$ -small. Since $g^*(u) > \tilde{g}(u)$, it follows that $P \circ (F_c \times id_I)$ is a τ -small homotopy joining the left and the right side of the relation (19).

Add (20). Since $\phi \cong \lambda$, there is a $u \in \tilde{Z}$, a normal cover η of $Y \times I$, and an (η, τ) -map $S: Y \times I \rightarrow Z$ joining G_u and L_u . Let ξ be a normal cover of Y obtained by the application of [4, p. 358] to the cover η . If $c > f(\xi)$, then F_a is ξ -small so that $S \circ (F_c \times id_I)$ is a τ -small homotopy joining compositions which appear in (20).

Add (21). Let $u > y$. Then L_u is an $(\tilde{l}(u), \tau)$ -map. Since φ and κ are homotopic, there is an index $c \in C$ so that F_c and K_c are joined by an $\tilde{l}(u)$ -small homotopy $T: X \times I \rightarrow Y$. The composition $L_u \circ T$ realizes the relation (21). \square

THEOREM 1. *The composition of homotopy classes of multi-nets is associative.*

PROOF OF THEOREM 1. Let $\varphi = \{F_c\}$, $\psi = \{G_s\}$ and $\chi = \{H_p\}$ be multi-nets from X into Y , from Y into Z , and from Z into W , respectively. Let $\mu = \{M_s\}$, $\nu = \{N_p\}$, $\kappa = \{K_p\}$ and $\lambda = \{L_p\}$, where $M_s = G_{g(s)} \circ F_{f(g^*(s))}$ for every $s \in \tilde{Z}$ and $N_p = H_{h(p)} \circ G_{g(h^*(p))}$, $K_p = H_{h(p)} \circ M_{m(h^*(p))}$, and $L_p = N_{n(p)} \circ F_{f(n^*(p))}$, for every $p \in \tilde{W}$. We must show that κ and λ are homotopic, i. e., that for every $\pi \in \tilde{W}$ there is a $p \in \tilde{W}$ such that

$$(24) \quad K_q \overset{\pi}{\cong} L_q \quad \text{for every } q > p.$$

Let $\pi \in \tilde{W}$. Let $\rho \in \pi^{*4}$. Let $p = \{\rho\} \in W$. In order to prove (24), we shall show that for every $q > p$ we can find indices $c \in \tilde{Y}$ and $s \in \tilde{Z}$ such that

$$(25) \quad K_q \overset{\rho}{\cong} H_x \circ G_y \circ F_c,$$

$$(26) \quad H_x \circ G_y \circ F_c \overset{\rho}{\cong} H_x \circ G_s \circ F_c,$$

$$(27) \quad H_x \circ G_s \circ F_c \overset{\rho}{\cong} H_z \circ G_s \circ F_c,$$

$$(28) \quad H_z \circ G_s \circ F_c \overset{\rho}{\cong} N_w \circ F_c,$$

and

$$(29) \quad N_w \circ F_c \stackrel{\rho}{\cong} L_q,$$

where $x=h(q)$, $y=g(m(h^*(q)))$, $z=h(n(q))$ and $w=n(q)$. Repeated use of Lemma 3 will give (24) from the relations (25)–(29).

The method of proof is similar to the proof of Claim 3. We shall only describe for each of the relations (25)–(29) how large the indices u and c must be in order that this homotopy holds. An easy exercise of putting together all these selections is once again left to the reader. Since relations (28) and (29) are analogous with relations (26) and (25), respectively, it suffices to consider only relations (25)–(27).

Add (25). Observe that H_x is an (α, ρ) -map while G_y is a (β, α) -map, where $\alpha=\tilde{h}(q)$, $\beta=h^*(q)$, and $\gamma=\tilde{g}(m(\beta))$. Let $\delta=g^*(m(\beta))$. If $c>f(\delta)$, then $F_{f(\delta)}$ and F_c are joined by a δ -small homotopy $P: X \times I \rightarrow Y$. But, δ refines γ by the property (1) of Claim 1. Hence, $H_x \circ G_y \circ P$ is a ρ -small homotopy between K_q and $H_x \circ G_y \circ F_c$.

Add (26). As above, H_x is an (α, ρ) -map. Since $m(s)>s$ for every $s \in \tilde{Z}$, we get $y>g(\beta)$. Therefore, if we take $s>y$, then G_y and G_s are joined by an (ε, β) -map $Q: Y \times I \rightarrow Z$, where $\varepsilon=g(\beta)$. However, β refines α so that $H_x \circ Q$ is an (ε, ρ) -map. Let η be a normal cover of Y associated to ε by [4, p. 358]. Finally, for $c>f(\eta)$ we see that $H_x \circ Q \circ (F_c \times id_I)$ realizes the relation (26).

Add (27). Since $n(r)>r$ for every $r \in \tilde{W}$, we get $z>x$ so that H_x and H_z are joined by an (η, ρ) -map $T: Z \times I \rightarrow W$, where η denotes the normal cover $h(z)$ of $Z \times I$. Let $\xi=h^*(z)$ and let $s>g(\xi)$. Then G_s is a $(\tilde{g}(s), \xi)$ -map. Let $\zeta=g^*(s)$ and take $c>f(\zeta)$. The composition $T \circ ((G_s \circ F_c) \times id_I)$ realizes the relation (27). \square

The category \mathcal{AM}

For a topological space X , let $\iota^X = \{I_a\}: X \rightarrow X$ be the identity multi-net defined by $I_a = id_X$ for every $a \in \tilde{X}$. It is easy to show that for every multi-net $\varphi: X \rightarrow Y$ the following relations hold.

$$[\varphi] \circ [\iota^X] = [\varphi] = [\iota^Y] \circ [\varphi]$$

We can summarize the above with the following theorem.

THEOREM 2. *The topological spaces as objects together with the homotopy classes of multi-nets as morphisms and the composition of homotopy classes form the category \mathcal{AM} .*

There is an obvious functor J from the category \mathcal{Top} of topological spaces and continuous maps into the category \mathcal{AM} . On objects the functor J is the identity while on morphisms it associates to a map $f: X \rightarrow Y$ the homotopy class of a multi-net $\underline{f} = \{F_c\}: X \rightarrow Y$, where $F_c = f$ for every $c \in \tilde{Y}$.

Our first main result can be stated as follows. Let Sh be the shape category of arbitrary topological spaces and let $S: \mathcal{Top} \rightarrow Sh$ be the shape functor [7].

THEOREM 3. *There is a functor θ from the category \mathcal{AM} into the shape category Sh such that $S = \theta \circ J$.*

Description of the functor θ

The functor θ will leave the objects unchanged. In order to explain how θ effects the morphisms we must work much harder. First we encounter the dilemma of selecting the right description of shape morphisms among the many that exist in the literature.

In the rest of this paper, let X, Y and Z be topological spaces and let

$$\mathbf{p} = \{p^a\}: X \longrightarrow \mathcal{X} = \{X_a, \varepsilon_a, p_b^a, A\},$$

$$\mathbf{q} = \{q^c\}: Y \longrightarrow \mathcal{Y} = \{Y_c, \xi_c, q_a^c, C\},$$

and

$$\mathbf{r} = \{r^m\}: Z \longrightarrow \mathcal{Z} = \{Z_m, \nu_m, r_n^m, M\}$$

be uniform commutative approximate resolutions of X, Y and Z where each X_a, Y_c and Z_m is a polyhedron, $st^3(\varepsilon_a)$ -close maps into X_a , $st^3(\xi_c)$ -close maps into Y_c and $st^3(\nu_m)$ -close maps into Z_m are homotopic, and $A = (A, >)$, $C = (C, >)$, and $M = (M, >)$ are infinite cofinite directed sets with cardinalities greater or equal to cardinalities of \tilde{X}, \tilde{Y} and \tilde{Z} , respectively. The existence of such approximate resolutions follows from [8] and [11].

We can associate with the approximate resolutions \mathbf{p}, \mathbf{q} and \mathbf{r} the underlying expansions in the sense of Morita [7]

$$|\mathbf{p}| = \{p^a\}: X \longrightarrow |\mathcal{X}| = \{X_a, p_b^a, A\},$$

$$|\mathbf{q}| = \{q^c\}: Y \longrightarrow |\mathcal{Y}| = \{Y_c, q_a^c, C\},$$

and

$$|\mathbf{r}| = \{r^m\}: Z \longrightarrow |\mathcal{Z}| = \{Z_m, r_n^m, M\}.$$

It is well-known that shape morphisms from X into Y could be considered as equivalence classes of morphisms of inverse systems $|\mathcal{X}|$ and $|\mathcal{Y}|$ (see [7] and [11]). More precisely, the set $Sh(X, Y)$ of all shape morphisms between

spaces X and Y can be identified with the set $pro\text{-}\mathcal{A}Pol(|\mathcal{X}|, |\mathcal{Y}|)$ of all morphisms in the Grothendick's pro-category $pro\text{-}\mathcal{A}Pol$ of the homotopy of polyhedra $\mathcal{A}Pol$ between the objects $|\mathcal{X}|$ and $|\mathcal{Y}|$. In our description of what θ does on morphisms of the category $\mathcal{A}M$ we shall view shape morphisms in this way.

Let $\varphi = \{F_s\}_{s \in \tilde{Y}} : X \rightarrow Y$ be a multi-net. By Lemma 2, we can find a refinement η_c of ξ_c so that for every $st^2(\eta_c)$ -small multi-valued function $K : W \rightarrow Y_c$ there is a normal cover ρ of W with the property that for every canonical map $r : W \rightarrow N(\rho)$ there is a map $k : N(\rho) \rightarrow Y_c$ with $K \stackrel{\xi_c}{=} k \circ r$. Let $\pi_c = (q^c)^{-1}(\eta_c)$.

Choose an index $l_c \in \tilde{Y}$ so that

$$(31) \quad F_s \stackrel{\pi_c}{\cong} F_t \quad \text{for all } s, t > l_c$$

Let $\lambda : C \rightarrow \tilde{Y}$ be an increasing function such that $\lambda(c) > l_c$, $\{\pi_c\}$, $v(c)$ for every $c \in C$, where $v : C \rightarrow \tilde{Y}$ is a surjection. We shall need later the fact that the function λ is cofinal, i.e., that for every $s \in \tilde{Y}$ there is a $d \in C$ with $\lambda(d) > s$.

Observe that $F_{\lambda(c)}$ is π_c -small. Hence, $q^c \circ F_{\lambda(c)}$ is η_c -small. Let ρ be a normal cover of X such that $q^c \circ F_{\lambda(c)}$ is a (ρ, η_c) -map. Let $r : X \rightarrow N(\rho)$ be a canonical map. The way in which we selected the cover η_c gives a map k which satisfies

$$(32) \quad k \circ r \stackrel{\xi_c}{=} q^c \circ F_{\lambda(c)}.$$

Let $\zeta = k^{-1}(\xi_c)$. By the property (R1) for the approximate resolution p , there is an index $f(c) \in A$ and a map $g : X_{f(c)} \rightarrow N(\rho)$ with $r \stackrel{\zeta}{=} g \circ p^{f(c)}$. Hence,

$$(33) \quad k \circ r \stackrel{\xi_c}{=} k \circ g \circ p^{f(c)}.$$

Let $f^c = k \circ g : X_{f(c)} \rightarrow Y_c$. The relations (32) and (33) together imply

$$(34) \quad f^c \circ p^{f(c)} \stackrel{st(\xi_c)}{=} q^c \circ F_{\lambda(c)}.$$

CLAIM 4. *The pair $\mathbf{f} = (f, \{f^c | c \in C\})$ is a morphism between inverse systems $|\mathcal{X}|$ and $|\mathcal{Y}|$.*

PROOF OF CLAIM 4. We must show that for every pair c, d of elements of C with $d > c$ it is possible to find an $a > f(c), f(d)$ so that

$$(35) \quad f^c \circ p_a^{f(c)} \cong q_a^c \circ f^d \circ p_a^{f(d)}.$$

Since $\lambda(d) > \lambda(c) > l_c$, by (31), the functions $F_{\lambda(c)}$ and $F_{\lambda(d)}$ can be joined by π_c -small homotopy $H : X \times I \rightarrow Y$. Hence, $q^c \circ F_{\lambda(c)}$ and $q^c \circ F_{\lambda(d)}$ are joined by the η_c -

small homotopy $q^c \circ H$. It follows that there is a single-valued continuous function $K: X \times I \rightarrow Y$ with

$$(36) \quad K \stackrel{\xi_c}{=} q^c \circ H.$$

The way in which we constructed f^d , the relation $q^c = q_a^c \circ q^d$, and the uniformity property of q give

$$(37) \quad q^c \circ F_{\lambda(a)} \stackrel{st(\xi_c)}{=} q_a^c \circ f^d \circ p^{f(d)}.$$

We know that

$$(38) \quad q^c \circ F_{\lambda(c)}(x) \subset q^c \circ H(x, 0) \quad \text{and} \quad q^c \circ F_{\lambda(a)}(x) \subset q^c \circ H(x, 1)$$

for every $x \in X$. Combining relations (36), (34) and (38) we obtain

$$(39) \quad K_0 \stackrel{st^2(\xi_c)}{=} f^c \circ p^{f(c)},$$

while (36), (37) and (38) imply

$$(40) \quad K_1 \stackrel{st^2(\xi_c)}{=} q_a^c \circ f^d \circ p^{f(d)},$$

where $K_0, K_1: X \rightarrow Y$ are defined by $K_0(x) = K(x, 0)$ and $K_1(x) = K(x, 1)$ for every $x \in X$. But, the assumption about ξ_c gives that the maps appearing in (39) and (40) are homotopic. Hence,

$$(41) \quad f^c \circ p_b^{f(c)} \circ p^b \cong q_a^c \circ f^d \circ p_b^{f(d)} \circ p^b,$$

where $b > f(c), f(d)$. However, the system $|\mathcal{X}|$ satisfies the condition (E2) from the reference [7, p. 48], so that an $a > b$ for which (35) holds surely exists. \square

Now we can define that θ acts on morphisms of the category \mathcal{M} (i. e., on homotopy classes of multi-nets) by the rule $\theta([\varphi]) = [f]$, where $[f]$ denotes the equivalence class of f with respect to the equivalence relation \sim (see [7, p. 6]).

CLAIM 5. *The function θ is well-defined i.e., it does not depend on the choices of $\varphi, \lambda,$ and f^c in our description of f .*

PROOF OF CLAIM 5. Suppose that $\psi = \{G_c\}: X \rightarrow Y$ is multi-net homotopic to φ and let the morphism $g = (g, \{g^c | c \in C\})$ of inverse systems $|\mathcal{X}|$ and $|\mathcal{Y}|$ be constructed from ψ by the above procedure using in it μ instead of λ . We must show that f and g are equivalent, i. e., that for every $c \in C$ there is an $a > f(c), g(c)$ with

$$(42) \quad f^c \circ p_a^{f(c)} \cong g^c \circ p_a^{g(c)}.$$

Let a $c \in C$ be given. Since φ and ψ are homotopic multi-nets, there is an

index $s_c \in \tilde{Y}$ such that

$$(43) \quad F_t \stackrel{\pi_c}{\cong} G_u \quad \text{for all } t, u > s_c.$$

Since the functions λ and μ are increasing and cofinal, there is a $d > c$ such that $\lambda(d), \mu(d) > s_c$. From (43), we get

$$(44) \quad F_{\lambda(d)} \stackrel{\pi_c}{\cong} G_{\mu(d)}$$

Let $H: X \times I \rightarrow Y$ be a π_c -small multi-valued function with $F_{\lambda(d)}(x) \subset H(x, 0)$ and $G_{\mu(d)}(x) \subset H(x, 1)$ for every $x \in X$. Hence, $q^c \circ H$ is an η_c -small multi-valued function and

$$(45) \quad q^c \circ F_{\lambda(d)}(x) \subset q^c \circ H(x, 0) \quad \text{and} \quad q^c \circ G_{\mu(d)}(x) \subset q^c \circ H(x, 1)$$

for every $x \in X$. Just as in the proof of Claim 4 there is a single-valued continuous function $N: X \times I \rightarrow Y_c$ with

$$(46) \quad N \stackrel{\xi_c}{\cong} q^c \circ H.$$

On the other hand, since $\lambda(d) > \lambda(c) > l_c$ the functions $F_{\lambda(c)}$ and $F_{\lambda(d)}$ are joined by a π_c -small homotopy $L: X \times I \rightarrow Y$. It follows that $q^c \circ L$ is an η_c -small homotopy which satisfies

$$(47) \quad q^c \circ F_{\lambda(c)}(x) \subset q^c \circ L(x, 0) \quad \text{and} \quad q^c \circ F_{\lambda(d)}(x) \subset q^c \circ L(x, 1)$$

for every $x \in X$. Pick a single-valued continuous function $M: X \times I \rightarrow Y_c$ with

$$(48) \quad M \stackrel{\xi_c}{=} q^c \circ L.$$

Similarly, there is a single-valued continuous function $P: X \times I \rightarrow Y_c$ together with a π_c -small homotopy $R: X \times I \rightarrow Y$ such that

$$(49) \quad q^c \circ G_{\mu(c)}(x) \subset q^c \circ R(x, 0) \quad \text{and} \quad q^c \circ F_{\mu(c)}(x) \subset q^c \circ R(x, 1)$$

for every $x \in X$, and

$$(50) \quad P \stackrel{\xi_c}{=} q^c \circ R.$$

In analogy with (35), we also have

$$(51) \quad g^c \circ p^{g(c)} \stackrel{st(\xi_c)}{=} q^c \circ G_{\mu(c)}.$$

Let M_0, M_1, N_0, N_1, P_0 and P_1 be maps defined from maps M, N and P as we defined K_0 and K_1 from K in the proof of Claim 4.

The relations (34) and (48) imply that $f^c \circ p^{f(c)}$ and M_0 are $st^2(\xi_c)$ -close maps

into Y_c . It follows that they are homotopic. Similarly, the maps $g^c \circ p^{g(c)}$ and P_1 are homotopic.

The maps M_1 and N_0 are also homotopic because from relations (45)–(48) we see that both are ξ_c -close to the function $q^c \circ F_{\lambda(c)}$. The maps N_1 and P_0 are homotopic because of a similar reason.

We conclude from the last two paragraphs that maps $f^c \circ g^{f(c)}$ and $g^c \circ p^{g(c)}$ are homotopic. Just as in the proof of Claim 4, with the help of the condition (E2), we can conclude that there exists an $a > f(c)$, $g(c)$ so that (42) holds. \square

CLAIM 6.

- (1) Let $\iota = \{(id_X)_{x \in \tilde{X}}\}$ be the identity multi-net on a space X . Then the morphism $i: |\mathcal{X}| \rightarrow |\mathcal{X}|$ associated to ι by our description of θ is the identity morphism $(id_A, \{(id_X)_a \mid a \in A\})$.
- (2) Let $\varphi = \{F_i\}: X \rightarrow Y$ and $\psi = \{G_u\}: Y \rightarrow Z$ be multi-nets. Then

$$\theta([\psi] \circ [\varphi]) = \theta([\psi]) \circ \theta([\varphi]).$$

- (3) θ is a functor and the relation $S = \theta \circ J$ holds.

PROOF OF CLAIM 6 (2). Let $\eta = \{H_u\}: X \rightarrow Z$, where $H_u = G_{g(u)} \circ F_{f(g^*(u))}$ for every $u \in \tilde{Z}$. Let $\mathbf{f} = (f, \{f^c\}_{c \in C})$, $\mathbf{g} = (g, \{g^m\}_{m \in M})$, and $\mathbf{h} = (h, \{h^m\}_{m \in M})$ be obtained from φ, ψ and η by the above procedure. We must show that \mathbf{h} and $\mathbf{g} \circ \mathbf{f}$ are homotopic. Since $\mathbf{g} \circ \mathbf{f} = (f \circ g, \{g^m \circ f^{g(m)}\})$, this amounts to show that for every $m \in M$ there is an $a > t$, x such that

$$(52) \quad h^m \circ p_a^x \cong g^m \circ f^v \circ p_a^t,$$

where $t = f \circ g(m)$, $x = h(m)$ and $v = g(m)$.

Once again, our method is to show that

$$(53) \quad h^m \circ p^x \cong g^m \circ f^v \circ p^t,$$

and then use the condition (E2) to get the required index.

In order to establish (53), we shall argue that there are large enough indices $b \in C$ and $n \in M$ such that

$$(54) \quad h^m \circ p^x \stackrel{st(\nu_m)}{=} r^m \circ H_y,$$

$$(55) \quad r^m \circ H_y \stackrel{\mu_m}{\cong} r^m \circ G_z \circ F_c,$$

$$(56) \quad r^m \circ G_z \circ F_c \stackrel{\mu_m}{\cong} r^m \circ G_n \circ F_c,$$

$$(57) \quad r^m \circ G_n \circ F_c \stackrel{\mu_m}{\cong} r^m \circ G_w \circ F_c,$$

$$(58) \quad r^m \circ G_w \circ F_c \stackrel{st^2(\nu_m)}{=} g^m \circ q^v \circ F_c,$$

$$(59) \quad g^m \circ q^v \circ F_c \stackrel{\mu_m}{\cong} g^m \circ q^v \circ F_u,$$

and

$$(60) \quad g^m \circ q^v \circ F_u \stackrel{st(\mu_m)}{=} g^m \circ f^v \circ p^t,$$

where $y = \gamma(m)$, $z = g(\gamma(m))$, $w = \kappa(m)$, $u = \lambda(g(m))$, μ_m is analogous to η_c and γ , κ and λ are functions used in constructing h , g and f , respectively.

Suppose for a moment that the relations (54)–(57) hold. From (55)–(57) it follows that there is a $st^2(\mu_m)$ -small multi-valued function $K: X \times I \rightarrow Z$ such that

$$(61) \quad r^m \circ H_y(x) \subset K(x, 0) \quad \text{and} \quad r^m \circ G_w \circ F_c(x) \subset K(x, 1)$$

for every $x \in X$. Similarly, from (59), it follows that there is a μ_m -small multi-valued function $L: X \times I \rightarrow Z$ with

$$(62) \quad g^m \circ q^v \circ F_c(x) \subset L(x, 0) \quad \text{and} \quad g^m \circ q^v \circ F_u(x) \subset L(x, 1)$$

for every $x \in X$. Let B and D be single-valued continuous functions such that

$$(63) \quad B \stackrel{\nu_m}{=} K \quad \text{and} \quad D \stackrel{\nu_m}{=} L$$

From (54), (61) and (63), we get that maps $h^m \circ p^x$ and B_0 are $st^3(\nu_m)$ -close. Hence,

$$(64) \quad h^m \circ p^x \cong B_0.$$

Similarly, from (58) and (61)–(63), it follows that the two maps B_1 and D_0 are $st^3(\nu_m)$ -close. Hence,

$$(65) \quad B_1 \cong D_0.$$

Finally, from (60), (62) and (63), we obtain that maps D_1 and $g^m \circ f^v \circ p^t$ are $st^3(\nu_m)$ -close. Hence,

$$(66) \quad D_1 \cong g^m \circ f^v \circ p^t.$$

The relations (64)–(66) together imply the relation (53). Thus it remains to explain why (54)–(60) hold. We shall describe what choice of c and n make each of these relations true and leave to the reader to put together all choices to pick them so that all are true simultaneously.

Add (54). This follows from the way in which h was constructed (it corresponds to the relation (34)).

Add (55). Observe that $H_y = G_z \circ F_s$, where $s = f(g^*(y))$. Since G_z is a

$(\tilde{g}(y, z, z), \tilde{y})$ -map (and therefore also a $(g^*(y), \tilde{y})$ -map because $g^*(y)$ refines $\tilde{g}(y, z, z)$) and \tilde{y} refines $\rho_m = (r^m)^{-1}(\mu^m)$, it suffices to take $c > s$ because then F_s and F_c are joined by a $g^*(y)$ -small homotopy $Q: X \times I \rightarrow Y$ so that $r^m \circ G_z \circ Q$ is a μ_m -small homotopy joining $r^m \circ H_y$ and $r^m \circ G_z \circ F_c$.

Add (56). Let $n > z$. Then G_z and G_n are joined by a $(g(y, z, n), \tilde{y})$ -map $R: Y \times I \rightarrow Z$. Hence, if $c > g^*(n)$, then $r^m \circ R \circ (F_c \times id_I)$ is a μ_m -small homotopy joining $r^m \circ G_z \circ F_c$ and $r^m \circ G_n \circ F_c$.

Add (57). Let $n > w$. Then G_n and G_w are joined by a ρ_m -small homotopy $T: Y \times I \rightarrow Z$. Let ω be a normal cover of $Y \times I$ such that T is an (ω, ρ_m) -map and let ζ be a normal cover of Y obtained from ω by application of [4, p. 358]. Let $c > f(\{\zeta\})$. Then F_c is a ζ -small multi-valued function so that $r^m \circ T \circ (F_c \times id_I)$ is a μ_m -small homotopy joining $r^m \circ G_n \circ F_c$ and $r^m \circ G_w \circ F_c$.

Add (58). First we observe that

$$(67) \quad r^m \circ G_w = g^m \circ q^v.$$

The relation (67) is just the version of the relation (34) for ϕ . Choose a normal cover π of Y such that G_w is a (π, ρ_m) -map. Let $c > f(\{\pi\})$. Then F_c is π -small and the composition $r^m \circ G_w \circ F_c$ is μ_m -small. Let $\beta = (g^m \circ q^v)^{-1}(\mu_m)$. Let $c > f(\{\beta\})$. Then F_c is β -small and the composition $g^m \circ q^v \circ F_c$ is also μ_m -small. With this information on the size of both sides appearing in (58), from (67), we can get (58).

Add (59). We can assume that $\xi_v > (g^m)^{-1}(\mu_m)$ for every $m \in M$. It might be necessary to pass from a given set of ξ_c 's to the new ones by an inductive argument on number of predecessors in order to accomplish this. Let $c > u$. Then F_c and F_u are joined by a π_v -small homotopy $U: X \times I \rightarrow Y$ and $g^m \circ q^v \circ U$ is a μ_m -small homotopy between $g^m \circ q^v \circ F_c$ and $g^m \circ q^v \circ F_u$.

Add (60). The relation (34) for $c = v$ reads

$$q^v \circ F_u = f^v \circ p^t.$$

Since $\xi_v > (g^m)^{-1}(\mu_m)$, we get from this the relation (60). □

PROOF OF CLAIM 6 (3). That θ is a functor follows from the previous discussion. It remains to see that $S = \theta \circ J$. Let $f: X \rightarrow Y$ be a map, i.e., a morphism of the category $\mathcal{T}op$. For each $c \in C$, there is a $\mu_c \in \xi_c^{*2}$ such that q is also a commutative uniform approximate resolution of Y into the approximate inverse system $q' = \{Y_c, \mu_c, q_c^a, C\}$. By Theorem (6.3) in [8], there is an approximate map $f: \mathcal{X} \rightarrow q'$ such that (p, q, f) is an approximate resolution of f . By Lemma (5.6) in [8], we get

$$(68) \quad f^c \circ p^{f(c)} \stackrel{\xi_c}{=} q^c \circ f.$$

Let $\varphi_f = \{F_i\} : X \rightarrow Y$ be a multi-net, where $F_i = f$ for every $i \in \tilde{Y}$. Then $[\varphi_f] = J(f)$. In applying the procedure from the description of θ to the multi-net φ_f we can take for λ a constant function and the above morphism f . The relation (68) implies that $S = \theta \circ J$. Indeed, the induced morphisms satisfy

$$(69) \quad |f| \circ |p| = |q| \circ f.$$

Since there is a unique morphism which satisfies (69), namely the morphism $S(f)$, we get $S(f) = \theta(J(f))$. \square

Inverse of θ

We shall now prove that on spaces which admit ANR-resolutions with the onto projections (that we call O-spaces) the functor θ is a category isomorphism.

DEFINITION 7. A space X is called an *O-space* provided there is an ANR-resolution $p = \{p^a\} : X \rightarrow \{X_a, p_b^a, A\}$ in the sense of Mardesić [7], where each projection p^a is an onto map.

At present we do not know what is the real extend of O-spaces. From results in [11], it follows that inverse limits of inverse systems of compact Hausdorff spaces with onto bonding maps are O-spaces. In particular, all compact metric spaces are O-spaces. One can easily check that the examples of non-degenerate regular spaces with the property that every real valued map on them is constant [5, p. 160] provide examples of spaces that are not O-spaces.

THEOREM 4. *Let X be a topological space and let Y be an O-space. Then the function $\theta : \mathcal{AM}(X, Y) \rightarrow Sh(X, Y)$ is a bijection.*

In order to prove Theorem 4, we shall construct the function $\zeta : Sh(X, Y) \rightarrow \mathcal{AM}(X, Y)$ which will be the inverse for the function θ . The description of ζ and the verification of its properties is given below in Claims 7-10.

Construction of the function ζ

Let $f = (f, \{f^c\}_{c \in C})$ be a morphism between inverse systems $|\mathcal{X}|$ and $|\mathcal{Y}|$. Let $s \in \tilde{Y}$. Recall that s is a finite set of normal covers of Y with the unique maximal element $\tilde{s} \in \hat{Y}$. By the condition (B1) for the approximate resolution

q [8], there is an index $c(s) \in C$ such that

$$(70) \quad (q^c)^{-1}(\xi_c) \quad \text{refines } \tilde{s} \text{ for every } c > c(s).$$

Let $\gamma: \tilde{Y} \rightarrow C$ be an increasing function with $\gamma(s) > c(s)$ for every $s \in \tilde{Y}$. Let $\varphi = \{F_s\}_{s \in \tilde{Y}}$, where $F_s = (q^{\gamma(s)})^{-1} \circ f^{\gamma(s)} \circ p^{\gamma(s)}$.

CLAIM 7. *The family φ is a multi-net from X into Y .*

PROOF OF CLAIM 7. Let a $\sigma \in \hat{Y}$ be given. We must show that there is a $c \in \tilde{Y}$ such that

$$(71) \quad F_t \overset{\sigma}{\cong} F_s \quad \text{for every } t > s.$$

Let $s = \{\sigma\} \in \hat{Y}$. Let $t > s$. Put $m = \gamma(t)$, $n = \gamma(s)$, $v = f(m)$ and $w = f(n)$. Since $m > n$ and f is a morphism of inverse systems, there is an $a > v, w$ and a map $K: X_a \times I \rightarrow Y_n$ with

$$(72) \quad K(x, 0) = q_m^n \circ f^m \circ p_a^v(x) \quad \text{and} \quad K(x, 1) = f^m \circ p_a^w(x)$$

for every $x \in X_a$. Let $L = (q^n)^{-1} \circ K \circ (p^a \times id_I)$. Then $L: X \times I \rightarrow Y$ is a σ -small homotopy. Moreover, for every $x \in X$, from (72), we get

$$(73) \quad L(x, 0) = (q^n)^{-1} \circ q_m^n \circ f^m \circ p_a^v \circ p^a(x) \supset (q^m)^{-1} \circ f^m \circ p^v(x) = F_t(x),$$

and

$$(74) \quad L(x, 1) = (q^n)^{-1} \circ f^n \circ p_a^w \circ p^a(x) = (q^n)^{-1} \circ f^n \circ p^w(x) = F_s(x).$$

Hence, L is a σ -small homotopy between F_t and F_s . \square

Now we can define the function ζ by the rule $\zeta([\mathbf{f}]) = [\varphi]$.

CLAIM 8. *The function ζ is well-defined, i.e., the value $\zeta([\mathbf{f}])$ does not depend on the choice of the representative f of the equivalence class $[\mathbf{f}]$ and on the choice of the function γ in our description of φ .*

PROOF OF CLAIM 8. Let $\mathbf{g} = (g, \{g^c\}_{c \in C}) \in [\mathbf{f}]$. Let $\psi = \{G_s\}_{s \in \tilde{Y}}$ be constructed from \mathbf{g} by the above procedure using the increasing function $\mu: \tilde{Y} \rightarrow C$. We must show that φ and ψ are homotopic, i.e., that for every $\sigma \in \hat{Y}$ there is an $s \in \tilde{Y}$ such that

$$(75) \quad F_t \overset{\sigma}{\cong} G_t \quad \text{for every } t > s.$$

Let a $\sigma \in \hat{Y}$ be given. Let $\tau \in \sigma^{*2}$. Put $s = \{\tau\} \in \tilde{Y}$. Pick an increasing function $\delta: \tilde{Y} \rightarrow C$ such that $\delta(t) > \gamma(t)$, $\mu(t)$ for every $t \in \tilde{Y}$.

Let $t > s$. Let $m = \gamma(t)$, $n = \mu(t)$, $k = \delta(t)$, $u = f(m)$, $v = f(k)$, $y = g(n)$ and $z = g(k)$. Since $k > m$, there is an $a > u, v$ with $f^m \circ p_a^u \cong q_k^m \circ f^k \circ p_a^v$. As in the proof of Claim 7, we can conclude from here that

$$(76) \quad F_t \stackrel{\tau}{\cong} K,$$

where $K = (q^k)^{-1} \circ f^k \circ p^v$. Similarly, we obtain

$$(77) \quad L \stackrel{\tau}{\cong} G_t,$$

where $L = (q^k)^{-1} \circ g^k \circ p^z$. Since f and g are equivalent, there is a $b > v, z$ and a homotopy $H: X_b \times I \rightarrow Y_k$ with $H(x, 0) = f^k \circ p_b^v(x)$ and $H(x, 1) = g^k \circ p_b^z(x)$ for every $x \in X_a$. It follows that the composition $(q^k)^{-1} \circ H \circ (p^b \times id_I)$ is a τ -small homotopy joining K and L . This together with (77) and (76) implies (75). \square

CLAIM 9. For every morphism $f = (f, \{f^c\}_{c \in C}): |\mathcal{X}| \rightarrow |\mathcal{Y}|$ we have $|f| = \theta \circ \zeta([f])$.

PROOF OF CLAIM 9. For every $s \in \tilde{Y}$ choose an index $c(s) \in C$ such that (70) holds. Let $\gamma: \tilde{Y} \rightarrow C$ be an increasing function with $\gamma(s) > c(s)$ for every $s \in \tilde{Y}$. Let $\delta: C \rightarrow \tilde{Y}$ be a function such that $\delta(c) \in \gamma^{-1}(c)$ whenever $\gamma^{-1}(c) \neq \emptyset$. Let $\varphi = \{F_s\}_{s \in \tilde{Y}}$, where $F_s = (q^{\gamma(s)})^{-1} \circ f^{\gamma(s)} \circ p^{\delta(s)}$. With respect to φ we now choose η_c, π_c and l_c as we did in the description of the function θ . Hence, we can assume that $(q^{\gamma(m)})^{-1} \circ f^{\gamma(m)} \circ p^{\delta(m)} \stackrel{\pi_c}{\cong} (q^{\gamma(n)})^{-1} \circ f^{\gamma(n)} \circ p^{\delta(n)}$ whenever $m, n > l_c$.

Next, we shall select a cofinal increasing function $\lambda: C \rightarrow \tilde{Y}$ such that $\lambda(c) > l_c, \{\pi_c\}, \delta(c)$ for every $c \in C$. Let $u = \lambda(c)$, $v = \gamma(u)$ and $w = f(v)$. Then $v > c$ and

$$q^c \circ F_u = q^c \circ (q^v)^{-1} \circ f^v \circ p^w = q_v^c \circ f^v \circ p^w.$$

Hence, in the next step, (i. e., the selection of the index “ $f(c)$ ” and the single-valued continuous function “ f^c ”) we can take some $z = g(c)$ with $z > w$ and the map $g^c = q_v^c \circ f^v \circ p_z^w$. It remains to check that the morphisms f and $g = (g, \{g^c\}_{c \in C})$ are equivalent. In other words, that for every $c \in C$ we can find an $a > f(c), g(c)$ with

$$f^c \circ p_a^{f(c)} \cong g^c \circ p_a^{g(c)} = q_v^c \circ f^v \circ p_z^w \circ p_a^z.$$

But, this follows from the fact that $v > c$ and f is a morphism of inverse systems. \square

DEFINITION 8. Let σ be a normal cover of a space Y . Two multi-valued functions $F, G: X \rightarrow Y$ are σ -hooked provided for every $x \in X$ there is an $S_x \in \sigma$ such that S_x has non-empty intersection with both $F(x)$ and $G(x)$.

Observe that σ -close multi-valued functions are σ -hooked.

LEMMA 4. *Let $F, G: X \rightarrow Y$ be multi-valued functions and let σ be a normal cover of Y . If F and G are σ -small and σ -hooked, then $F \stackrel{st(\sigma)}{\cong} G$.*

PROOF OF LEMMA 4. Since F and G are σ -small, there is a normal cover η of X such that for every $E \in \eta$ there are $S_E, T_E \in \sigma$ with $F(E) \subset S_E$ and $G(E) \subset T_E$. Define a function $H: X \times I \rightarrow Y$ by the rule $H(x, t) = F(x) \cup G(x)$ for every $x \in X$ and every $t \in I$. Let $\xi = \{E \times I \mid E \in \eta\}$. Clearly, ξ is a normal cover of $X \times I$. We shall check that H is a $(\xi, st(\delta))$ -map. This would imply that H is a $st(\sigma)$ -small homotopy joining F and G .

Then $H(K) = F(E) \cup G(E) = S_E \cup T_E$, for a member $K = E \times I$ of ξ and $E \in \eta$. But, since F and G are σ -hooked, for every $x \in E$ there is an $R_x \in \sigma$ with $R_x \cap F(x) \neq \emptyset$ and $R_x \cap G(x) \neq \emptyset$. Hence, $H(K) \subset st(R_x, \sigma)$. \square

CLAIM 10. *For every multi-net $\varphi = \{F_s\}_{s \in \tilde{Y}}: X \rightarrow Y$ we have $\zeta \circ \theta([\varphi]) = [\varphi]$.*

PROOF OF CLAIM 10. We first perform steps from the description of the functor θ to get $C_c, \pi_c, l_c, \lambda, f$, and the maps f^c . Then we perform steps from the description of ζ to get indices $c(s)$, the function γ , and a multi-net $\psi = \{G_s\}_{s \in \tilde{Y}}$, where G_s is the composition $(q^{\gamma(s)})^{-1} \circ f^{\gamma(s)} \circ p^{f(\gamma(s))}$. We must show that multi-nets φ and ψ are homotopic, i.e., that for every $\sigma \in \hat{Y}$ there is an $s \in \tilde{Y}$ with

$$(78) \quad F_t \stackrel{\sigma}{\cong} G_t \quad \text{for every } t > s.$$

Let a $\sigma \in \hat{Y}$ be given. Let $\tau \in \sigma^{*3}$. Since φ is a multi-net there is an $s > \{\tau\}$ such that

$$(79) \quad F_r \stackrel{\sigma}{\cong} F_t \quad \text{for all } r, t > s.$$

Let $t > s$. We shall prove that there is a large enough index $c \in C$ with the property that

$$(80) \quad F_t \stackrel{\tau}{\cong} F_u,$$

$$(81) \quad F_u \stackrel{st^2(\tau)}{\cong} (q^c)^{-1} \circ f^c \circ p^v,$$

and

$$(82) \quad (q^c)^{-1} \circ f^c \circ p^v \stackrel{\tau}{\cong} G_t,$$

where $u = \lambda(c)$ and $v = f(c)$. The relations (80)–(82) and Lemma 3 imply (78).

Add (80). Since λ is a cofinal function, there is a $c \in C$ so that $\lambda(c) > s$.

Then (80) is a consequence of (79).

Add (81). Let $c > \gamma(\{\tau\})$. It follows that $(q^c)^{-1}(\xi_c)$ refines τ so that $(q^c)^{-1}(st(\xi_c))$ refines $st(\tau)$. Hence, from the relation (34), we get

$$(83) \quad (q^c)^{-1} \circ q^c \circ F_u \stackrel{st(\tau)}{=} (q^c)^{-1} \circ f^c \circ p^v.$$

But, the composition on the left side of (83) is a π_c -small multi-valued function. Since $\pi_c = (q^c)^{-1}(\eta_c)$ refines τ , by Lemma 4, from (83) we get (81).

Add (82). Let $w = \gamma(t)$ and $z = f(w)$. Let $c > w, \gamma(\{\tau\})$. Since $(f, \{f^a\}_{a \in c})$ is a morphism of inverse systems, there is an $a > v, z$ and a homotopy $H: X_a \times I \rightarrow Y_c$ with $H(x, 0) = q_c^v \circ f^c \circ p_a^z(x)$ and $H(x, 1) = f^w \circ p_a^z(x)$ for every $x \in X_a$. It follows that $(q^c)^{-1} \circ H \circ (p^a \times id_I)$ is a τ -small homotopy joining the left and the right side of (82). \square

REMARK. It is possible to use only multi-valued functions that are upper semi-continuous or to require that in addition images of points are compact. With these functions we shall get a similar result but the space Y is further restricted to spaces that admit ANR-resolutions with closed and perfect projections, respectively.

References

- [1] R. A. Aló and M. L. Shapiro, Normal Topological Spaces, Cambridge Univ. Press, Cambridge, 1972.
- [2] K. Borsuk, Concerning homotopy properties of compacta, Fund. Math. **62** (1968), 223–245.
- [3] K. Borsuk, Theory of Shape, Monografie Matem. 59, Polish Scientific Publishers, Warszawa, 1977.
- [4] A. Dold, Lectures on Algebraic Topology, Springer-Verlag, Berlin, 1972.
- [5] R. Engelking, General Topology, Monografie Matem. 60, Polish Scientific Publishers, Warszawa, 1977.
- [6] S. T. Hu, Theory of Retracts, Wayne State University Press, Detroit, 1965.
- [7] S. Mardešić and J. Segal, Shape Theory, North Holland, Amsterdam, 1982.
- [8] S. Mardešić and T. Watanabe, Approximate resolutions of spaces and mappings, Glasnik Mat. **24** (1989), 583–633.
- [9] J. Sanjurjo, Shape morphisms and small multivalued maps, Math. Japonica **35** (1990), 713–717.
- [10] J. Sanjurjo, An intrinsic description of shape, Trans. Amer. Math. Soc. **329** (1992), 625–636.
- [11] T. Watanabe, Approximate Shape, I, Tsukuba J. Math. **11** (1987), 17–59.

Kopernikova 7, 10020 Zagreb,
Croatia, Europe
E-mail: zcerin@x400.srce.hr