

AFFINE INNER AUTOMORPHISMS OF $SU(2)$

By

U-Hang KI and Joon-Sik PARK¹

Abstract. We show which inner automorphisms of $(SU(2), g)$ with an arbitrary left invariant metric g into itself are affine transformations, and obtain affine transformations of $(SU(2), g)$ which are not harmonic, and study geodesics of $(SU(2), g)$ with some conditions.

0. Introduction

It is interesting to show which diffeomorphisms between two Riemannian manifolds are affine transformations. In this paper, we treat the case $(SU(2), g)$ with a left invariant Riemannian metric g . It is well known that every inner automorphism of G a compact connected semisimple Lie group into itself is both affine and harmonic with respect to a bi-invariant Riemannian metric g_0 on G . However, we here deal with an arbitrary left invariant metric g on $SU(2)$, and show which inner automorphisms of $SU(2)$ are affine transformations of $(SU(2), g)$ into itself.

On the other hand we study geodesics in $(SU(2), g)$. In case of naturally reductive homogeneous space, it is well known that geodesics are orbits of 1-parameter subgroups. R. Dohira (cf. [1]) studied geodesics in reductive homogeneous spaces satisfying certain conditions. Using Dohira's Theorem, we give a complete description of geodesics in $(SU(2), g)$ satisfying some conditions.

In §1, we obtain necessary and sufficient conditions for inner automorphisms A_x , ($x \in SU(2)$), of $(SU(2), g)$, to be affine transformations (cf. Proposition 1.3–1.5). Moreover, in Theorem 1.7 and 1.8, we show that for any left invariant but not bi-invariant Riemannian metric g on $SU(2)$, there always exist on $(SU(2), g)$

Keywords: affine transformations, harmonic maps, left invariant metrics on Lie groups.

MS classification: 58E20, 53C42, 58G25.

¹The second author wishes to acknowledge the financial support of the Korean Research Foundation made in the program year of 1997

Received June 30, 1998

both a non-affine inner automorphism, and a non-harmonic but affine inner automorphism.

In §2, using R. Dohira's Theorem we give a complete description of geodesics in $(SU(2), g)$ satisfying certain conditions (cf. Theorem 2.1). Finally we get necessary and sufficient conditions for arbitrary given geodesics in $(SU(2), g)$ with certain left invariant metric g to be closed (cf. Theorem 2.3).

§1. Affine Inner Automorphisms of $(SU(2), g)$

Let B be the Killing form of the Lie algebra $\mathfrak{su}(2)$ of $SU(2)$. Then the Killing form satisfies

$$(1.1) \quad B(X, Y) = 4 \operatorname{Trace}(XY), \quad (X, Y \in \mathfrak{su}(2)).$$

We define an inner product $\langle \cdot, \cdot \rangle_0$ on $\mathfrak{su}(2)$ by

$$(1.2) \quad \langle \cdot, \cdot \rangle_0 := -B(X, Y), \quad (X, Y \in \mathfrak{su}(2)).$$

The following lemma is known (cf. [5, Lemma 1.1, p. 154]):

LEMMA 1.1. *Let g be a left invariant Riemannian metric. Let $\langle \cdot, \cdot \rangle$ be an inner product on $\mathfrak{su}(2)$ defined by $\langle X, Y \rangle := g_e(X_e, Y_e)$, where $X, Y \in \mathfrak{su}(2)$ and e is the identity matrix of $SU(2)$. Then there exist an orthonormal basis (X_1, X_2, X_3) of $\mathfrak{su}(2)$ with respect to $\langle \cdot, \cdot \rangle_0$ such that*

$$(1.3) \quad \begin{cases} [X_1, X_2] = (1/\sqrt{2})X_3, & [X_2, X_3] = (1/\sqrt{2})X_1, \\ [X_3, X_1] = (1/\sqrt{2})X_2, & \langle X_i, X_j \rangle = \delta_{ij}a_i^2, \end{cases}$$

where a_i , $(1 \leq i \leq 3)$, are positive real numbers determined by the given left invariant Riemannian metric g of $SU(2)$.

Let ∇ be the Riemannian connection on $(SU(2), g)$. Here g is an arbitrary given left invariant Riemannian metric in $SU(2)$. Let (X_1, X_2, X_3) be left invariant vector fields related to B and g which appear in Lemma 1.1.

An inner automorphism $A_x : (SU(2), g) \rightarrow (SU(2), g)$, $(x \in SU(2))$, is an affine transformation if and only if

$$(1.4) \quad Ad(x)\nabla_{X_i}X_j = \nabla_{Ad(x)X_i}Ad(x)X_j, \quad (i, j = 1, 2, 3).$$

With respect to the Riemannian connection, we have

$$(1.5) \quad \begin{aligned} 2 \cdot g(\nabla_X Y, Z) &= X \cdot g(Y, Z) + Y \cdot g(X, Z) - Z \cdot g(X, Y) \\ &\quad + g([X, Y], Z) + g([Z, X], Y) + g(X, [Z, Y]) \end{aligned}$$

for all vector fields X, Y, Z . In this section, for simplicity we put

$$(1.6) \quad \begin{cases} F_1 := (2\sqrt{2})^{-1}(a_1^2 - a_2^2)a_3^{-2}, & F_2 := (2\sqrt{2})^{-1}(a_2^2 - a_3^2)a_1^{-2}, \\ F_3 := (2\sqrt{2})^{-1}(a_3^2 - a_1^2)a_2^{-2}, \end{cases}$$

From (1.3) and (1.5), we get

$$(1.7) \quad \begin{cases} \nabla_{X_i} X_i = 0 \quad (i = 1, 2, 3), & \nabla_{X_1} X_2 = \{(2\sqrt{2})^{-1} - F_1\} X_3, \\ \nabla_{X_1} X_3 = -\{(2\sqrt{2})^{-1} + F_3\} X_2, & \nabla_{X_2} X_3 = \{(2\sqrt{2})^{-1} - F_2\} X_1. \end{cases}$$

We put

$$(1.8) \quad Y_i := 2\sqrt{2}X_i, \quad (i = 1, 2, 3).$$

Then, from (1.3) and (1.8) we have

$$(1.9) \quad [Y_1, Y_2] = 2Y_3, \quad [Y_2, Y_3] = 2Y_1, \quad [Y_3, Y_1] = 2Y_1.$$

In order to prove the following Propositions, we get:

LEMMA 1.2. *For an inner automorphism A_x , ($x \in SU(2)$),*

$$\nabla_{Ad(x)X_i} Ad(x)X_j = Ad(x)(\nabla_{X_i} X_j)$$

if and only if

$$\nabla_{Ad(x)X_j} Ad(x)X_i = Ad(x)(\nabla_{X_j} X_i), \quad (i, j = 1, 2, 3).$$

PROPOSITION 1.3. *An inner automorphism A_x , ($x = \exp(rY_1)$, $r \in \mathbf{R}$), of $(SU(2), g)$ is an affine transformation if and only if $a_2 = a_3$ or $\sin(2r) = 0$, that is,*

$$(1.10) \quad a_2 = a_3 \quad \text{or} \quad r \in \{(n\pi)/2 \mid n \text{ is an integer}\}.$$

PROOF. Using (1.3), (1.8) and (1.9), we have

$$(1.11) \quad \begin{cases} Ad(x)X_1 = X_1, \\ Ad(x)X_2 = \cos(2r)X_2 + \sin(2r)X_3, \\ Ad(x)X_3 = \cos(2r)X_3 - \sin(2r)X_2. \end{cases}$$

Putting $\phi := Ad(x)$, from (1.6), (1.7) and (1.11) we get

$$(1.12) \quad \left\{ \begin{aligned} \phi^{-1}(\nabla_{\phi X_1} \phi X_1) &= 0, \quad \phi^{-1}(\nabla_{\phi X_2} \phi X_2) = -\phi^{-1}(\nabla_{\phi X_3} \phi X_3) = -F_2 \sin(4r) X_1, \\ \phi^{-1}(\nabla_{\phi X_1} \phi X_2) &= -2^{-1}(F_1 + F_2) \sin(4r) X_2 \\ &\quad + \{(2\sqrt{2})^{-1} + F_3 \sin^2(2r) - F_1 \cos^2(2r)\} X_3, \\ \phi^{-1}(\nabla_{\phi X_1} \phi X_3) &= \{F_1 \sin^2(2r) - F_3 \cos^2(2r) - (2\sqrt{2})^{-1}\} X_2 \\ &\quad + 2^{-1}(F_1 + F_3) \sin(4r) X_3, \\ \phi^{-1}(\nabla_{\phi X_2} \phi X_3) &= \{(2\sqrt{2})^{-1} - F_2 \cos(4r)\} X_1. \end{aligned} \right.$$

Hence, we find from (1.6), (1.7), (1.12) and Lemma 1.2 that A_x is an affine transformation if and only if

$$(1.13) \quad a_2 = a_3 \quad \text{or} \quad \sin(2r) = 0. \quad \text{q.e.d.}$$

PROPOSITION 1.4. *An inner automorphism A_x , ($x = \exp(rY_2)$, $r \in \mathbf{R}$), of $(SU(2), g)$ is an affine transformation if and only if*

$$(1.14) \quad a_3 = a_1 \quad \text{or} \quad r \in \{(n\pi)/2 \mid n \text{ is an integer}\}.$$

PROOF. Using (1.3), (1.8) and (1.9), we have

$$(1.15) \quad \begin{cases} Ad(x)X_1 = \cos(2r)X_1 - \sin(2r)X_3, \\ Ad(x)X_2 = X_2, \quad Ad(x)X_3 = \sin(2r)X_1 + \cos(2r)X_3. \end{cases}$$

From (1.6), (1.7) and (1.15), we have

$$(1.16) \quad \left\{ \begin{aligned} \phi^{-1}(\nabla_{\phi X_1} \phi X_1) &= -\phi^{-1}(\nabla_{\phi X_3} \phi X_3) = F_3 \sin(4r) X_2, \quad \phi^{-1}(\nabla_{\phi X_2} \phi X_2) = 0, \\ \phi^{-1}(\nabla_{\phi X_1} \phi X_2) &= 2^{-1}(F_1 + F_2) \sin(4r) X_1 \\ &\quad + \{(2\sqrt{2})^{-1} + F_2 \sin^2(2r) - F_1 \cos^2(2r)\} X_3, \\ \phi^{-1}(\nabla_{\phi X_1} \phi X_3) &= -\{(2\sqrt{2})^{-1} + F_3 \cos(4r)\} X_2, \\ \phi^{-1}(\nabla_{\phi X_2} \phi X_3) &= \{(2\sqrt{2})^{-1} + F_1 \sin^2(2r) - F_2 \cos^2(2r)\} X_1 \\ &\quad - 2^{-1}(F_1 + F_2) \sin(4r) X_3, \end{aligned} \right.$$

where $\phi := Ad(x)$. We know from (1.6), (1.7), (1.16) and Lemma 1.2 that A_x is an affine transformation if and only if

$$(1.17) \quad a_1 = a_3 \quad \text{or} \quad \sin(2r) = 0. \quad \text{q.e.d.}$$

PROPOSITION 1.5. *An inner automorphism A_x , ($x = \exp(rY_3)$, $r \in \mathbf{R}$), is an affine transformation if and only if*

$$(1.18) \quad a_1 = a_2 \quad \text{or} \quad r \in \{(n\pi)/2 \mid n \text{ is an integer}\}.$$

PROOF. We get from (1.3), (1.8) and (1.9)

$$(1.19) \quad \begin{cases} Ad(x)(X_1) = \cos(2r)X_1 + \sin(2r)X_2, \\ Ad(x)(X_2) = \cos(2r)X_2 - \sin(2r)X_1, \quad Ad(x)(X_3) = X_3. \end{cases}$$

From (1.6), (1.7) and (1.19), we obtain

$$(1.20) \quad \begin{cases} \phi^{-1}(\nabla_{\phi X_1} \phi X_1) = -\phi^{-1}(\nabla_{\phi X_2} \phi X_2) = -F_1 \sin(4r)X_3, \\ \phi^{-1}(\nabla_{\phi X_3} \phi X_3) = 0, \quad \phi^{-1}(\nabla_{\phi X_1} \phi X_2) = \{(2\sqrt{2})^{-1} - F_1 \cos(4r)\}X_3, \\ \phi^{-1}(\nabla_{\phi X_1} \phi X_3) = -2^{-1}(F_2 + F_3) \sin(4r)X_1 \\ \quad + \{F_2 \sin^2(2r) - F_3 \cos^2(2r) - (2\sqrt{2})^{-1}\}X_2, \\ \phi^{-1}(\nabla_{\phi X_2} \phi X_3) = \{(2\sqrt{2})^{-1} + F_3 \sin^2(2r) - F_2 \cos^2(2r)\}X_1 \\ \quad + 2^{-1}(F_2 + F_3) \sin(4r)X_2, \end{cases}$$

where $\phi := Ad(x)$. Using (1.6), (1.7) and Lemma 1.2, we obtain this proposition. q.e.d.

Since the metric g of $(SU(2), g)$ is bi-invariant iff $a_1 = a_2 = a_3$, we obtain from Proposition 1.3, 1.4 and 1.5:

THEOREM 1.6. *An inner automorphism A_x of $(SU(2), g)$ for any $x \in SU(2)$ is an affine transformation if and only if the metric g of $(SU(2), g)$ is bi-invariant.*

Harmonic maps of a compact Riemannian manifold into another Riemannian manifold are the extrema (cf. [2, 6]). In the case of $(SU(2), g)$, the following Lemma (cf. [4]) is known:

LEMMA 1.7. *A necessary and sufficient condition for an inner automorphism A_x of $(SU(2), g)$ to be harmonic is*

$$\begin{cases} a_2 = a_3 \text{ or } \sin(4r) = 0, & \text{in case of } x = \exp(rY_1) \text{ and } r \in \mathbf{R}, \\ a_1 = a_3 \text{ or } \sin(4r) = 0, & \text{in case of } x = \exp(rY_2) \text{ and } r \in \mathbf{R}, \\ a_1 = a_2 \text{ or } \sin(4r) = 0, & \text{in case of } x = \exp(rY_3) \text{ and } r \in \mathbf{R}. \end{cases}$$

From Propositions 1.3–1.5 and Lemma 1.7, we get:

THEOREM 1.8. *Assume that a left invariant metric g of $(SU(2), g)$ is not bi-invariant. Then, there always exists harmonic inner automorphisms A_x of $(SU(2), g)$ which are not affine transformations.*

REMARK. An affine transformation between two Riemannian manifolds is harmonic.

Moreover, from (1.11), (1.15), (1.19) and Propositions 1.3–1.5, we have

COROLLARY 1.9. *If an inner automorphism A_x for $x \in SU(2)$ such that*

$$\begin{cases} \exp(rY_1) & \text{if } a_2 \neq a_3, \\ \exp(rY_2) & \text{if } a_3 \neq a_1, \\ \exp(rY_3) & \text{if } a_1 \neq a_2, \end{cases}$$

is an affine transformation, then A_x is an isometry.

§2. Geodesics in $(SU(2), g)$

We retain the notations as in §1. R. Dohira’s Theorem and Corollary which appear in [1] can be stated in our case $(SU(2), g)$ as follows:

THEOREM 2.1. *Assume $a_2 = a_3$. Let $\sigma(t)$ be a geodesic in $(SU(2), g)$ such that*

$$\sigma(0) = e, \quad \dot{\sigma}(0) = \sum_{i=1}^3 k_i Y_i \quad (\text{each } k_i \in \mathbf{R}).$$

Then

$$(2.1) \quad \sigma(t) = \exp(t(k_2 Y_2 + k_3 Y_3 + a_1^2 a_2^{-2} k_1 Y_1)) \exp(t(1 - a_1^2 a_2^{-2}) k_1 Y_1).$$

PROOF. We put $\{Y_2, Y_3\}_R =: \mathfrak{m}_1$ and $\{Y_1\}_R =: \mathfrak{m}_2$. Then

$$(2.2) \quad [\mathfrak{m}_1, \mathfrak{m}_1] \subset \mathfrak{m}_2, \quad [\mathfrak{m}_1, \mathfrak{m}_2] \subset \mathfrak{m}_1,$$

$$(2.3) \quad g([X, Y], Z) + a_1^2 a_2^{-1} g(X, [Z, Y]) = 0$$

for each $X, Y \in \mathfrak{m}_1, Z \in \mathfrak{m}_2$. In view of Dohira's Theorem (cf. [1]), we can get this Theorem.

COROLLARY 2.2. *Assume $a_2 = a_3$. A geodesic in $(SU(2), g)$ which intersects itself is a closed geodesic.*

Using Theorem 2.1 and Corollary 2.2, we obtain

THEOREM 2.3. *Let $r \in \mathbf{R} \setminus \{(n\pi)/2 \mid n \text{ is an integer}\}$ and $x = \exp(rY_1)$. Assume A_x is an affine transformation. Then, a geodesic $\sigma(t)$ in $(SU(2), g)$ with condition $\sigma(0) = e$ and $\dot{\sigma}(0) = \sum_{i=1}^3 k_i Y_i$ is closed if and only if there exist a real number $L (\in \mathbf{R} \setminus \{0\})$ satisfying*

$$(2.4) \quad \begin{cases} \cos(EL) = \cos((a_1^2 a_2^{-2} - 1)k_1 L), \\ a_1^2 a_2^{-2} k_1 \sin(EL) = E \sin((a_1^2 a_2^{-2} - 1)k_1 L), \\ k_2 \sin(EL) = 0, \text{ and } k_3 \sin(EL) = 0, \end{cases}$$

where $E := \sqrt{(k_2^2 + k_3^2 + a_2^{-4} k_1^2)}$.

PROOF. In this proof, we put $c := a_1^2 a_2^{-2}$. Then, from Proposition 1.3 and Theorem 2.1 we have

$$(2.5) \quad \sigma(t) = \exp(t(k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)) \exp((1 - c)k_1 t Y_1).$$

And then, if $\sigma(t)$ is closed, by Corollary 2.2 we know that there exist real numbers $L (\in \mathbf{R} \setminus \{0\})$ satisfying

$$(2.6) \quad \exp(L(k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)) = \exp((c - 1)k_1 L Y_1).$$

We may assume (cf. [5, Proof of Lemma 1.1, p. 154]) that (Y_1, Y_2, Y_3) , which appears in (1.8), satisfies

$$(2.7) \quad Y_i^{4n} = e, \quad Y_i^{4n+1} = Y_i, \quad Y_i^{4n+2} = -e, \quad Y_i^{4n+3} = -Y_i, \quad (i = 1, 2, 3),$$

for every non-negative integer n , and

$$(2.8) \quad Y_1 Y_2 = -Y_2 Y_1 = Y_3, \quad Y_2 Y_3 = -Y_3 Y_2 = Y_1, \quad Y_3 Y_1 = -Y_1 Y_3 = Y_2.$$

Using (2.7) and (2.8), we get

(2.9)

$$\left\{ \begin{array}{l} (k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)^{4n} = (k_2^2 + k_3^2 + c^2 k_1^2)^{2n} e, \\ (k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)^{4n+1} = (k_2^2 + k_3^2 + c^2 k_1^2)^{2n} (k_2 Y_2 + k_3 Y_3 + ck_1 Y_1), \\ (k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)^{4n+2} = -(k_2^2 + k_3^2 + c^2 k_1^2)^{2n+1} e, \\ (k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)^{4n+3} = -(k_2^2 + k_3^2 + c^2 k_1^2)^{2n+1} (k_2 Y_2 + k_3 Y_3 + ck_1 Y_1), \end{array} \right.$$

for every non-zero integer n . From (2.7), we have

$$(2.10) \quad \exp((c-1)k_1 L Y_1) = \cos((c-1)k_1 L) I_2 + \sin((c-1)k_1 L) Y_1.$$

By the help of (2.9), we obtain

$$(2.11) \quad \begin{aligned} \exp(L(k_2 Y_2 + k_3 Y_3 + ck_1 Y_1)) \\ = \cos(EL) I_2 + ck_1 E^{-1} \sin(EL) Y_1 + k_2 E^{-1} \sin(EL) Y_2 \\ + k_3 E^{-1} \sin(EL) Y_3. \end{aligned}$$

Comparing (2.10) with (2.11), we can get this Theorem.

q.e.d.

From this Theorem, we obtain the following:

COROLLARY 2.4. *Assume that $k_1 k_2 k_3 \neq 0$ and the metric g in $(SU(2), g)$ with $a_2 = a_3$ is not bi-invariant. Then, if $k_1^{-1}(a_1^2 a_2^{-2} - 1) \sqrt{(k_2^2 + k_3^2 + a_1^4 a_2^{-4} k_1^2)}$ is not a rational number, the geodesic $\sigma(t)$ with $\sigma(0) = e$ and $\dot{\sigma}(0) = \sum_{i=1}^3 k_i Y_i$ is not closed.*

References

- [1] R. Dohira, Geodesics in reductive homogeneous spaces, *Tsukuba J. Math.*, **19** (1995), 233–243.
- [2] J. Eells and L. Lemaire, *Selected Topics in Harmonic Maps*, CBMS Regional Conf., 1981.
- [3] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, New York, 1978.
- [4] Joon-Sik Park, Harmonic inner automorphisms of compact connected semisimple Lie groups, *Tohoku Math. J.*, **42** (1990), 83–91.
- [5] K. Sugahara, The sectional curvature and the diameter estimate for the left invariant metrics on $SU(2, \mathbb{C})$ and $SO(3, \mathbb{R})$, *Math. Japonica*, **26** (1981), 153–159.
- [6] H. Urakawa, Stability of harmonic maps and eigenvalues of Laplacian, *Trans. Amer. Math. Soc.*, **301** (1987), 557–589.

Topology and Geometry Research Center
Kyungpook National University
Taegu 702-701, Korea

Department of Mathematics
Pusan University of Foreign Studies
55-1, Uam Dong Nam-Gu, Pusan,
608-738, Korea