

Spin, statistics, orientations, unitarity

THEO JOHNSON-FREYD

A topological quantum field theory is *hermitian* if it is both oriented and complex-valued, and orientation-reversal agrees with complex conjugation. A field theory *satisfies spin-statistics* if it is both spin and super, and 360° -rotation of the spin structure agrees with the operation of flipping the signs of all fermions. We set up a framework in which these two notions are precisely analogous. In this framework, field theories are defined over $\mathbf{VECT}_{\mathbb{R}}$, but rather than being defined in terms of a single tangential structure, they are defined in terms of a bundle of tangential structures over $\mathrm{Spec}(\mathbb{R})$. Bundles of tangential structures may be étale-locally equivalent without being equivalent, and hermitian field theories are nothing but the field theories controlled by the unique nontrivial bundle of tangential structures that is étale-locally equivalent to Orientations. This bundle owes its existence to the fact that $\pi_1^{\text{ét}}(\mathrm{Spec}(\mathbb{R})) = \pi_1 \mathbf{BO}(\infty)$. We interpret Deligne’s “existence of super fiber functors” theorem as implying that $\pi_2^{\text{ét}}(\mathrm{Spec}(\mathbb{R})) = \pi_2 \mathbf{BO}(\infty)$ in a categorification of algebraic geometry in which symmetric monoidal categories replace commutative rings. One finds that there are eight bundles of tangential structures étale-locally equivalent to Spins, one of which is distinguished; upon unpacking the meaning of a field theory with that distinguished tangential structure, one arrives at a field theory that is both hermitian and satisfies spin-statistics. Finally, we formulate in our framework a notion of reflection-positivity and prove that if an étale-locally-oriented field theory is reflection-positive then it is necessarily hermitian, and if an étale-locally-spin field theory is reflection-positive then it necessarily both satisfies spin-statistics and is hermitian. The latter result is a topological version of the famous spin-statistics theorem.

14A22, 57R56, 81T50

0 Introduction

The main result of this article is a topological version of the spin-statistics theorem. The usual spin-statistics theorem (see Streater and Wightman [25]) asserts that in a unitary quantum field theory on Minkowskian spacetime, the fields of the theory live in a supervector space, the even (or bosonic) fields are integer spin representations of the Lorentz group, and the odd (or fermionic) fields are half-integer spin representations. In

other words, the spin of a particle agrees with its parity. Here unitarity is actually two conditions: a hermiticity condition (asserting that the determinant- (-1) component of the Lorentz group acts complex-antilinearly) and a reflection-positivity condition related to the requirement that the Hamiltonian of the quantum field theory have positive spectrum.

To formulate a version in the functorial setting of topological quantum field theory, we need

- to have orientations and spin structures on our source bordism category,
- to have complex supervector spaces in our target category, but to be able to talk about complex-antilinear maps as well as antisuper maps (ie maps that treat even and odd parts differently),
- to be able to link these structures on source and target categories.

We will solve all three problems by introducing generalizations of oriented and spin \mathbb{R} -linear field theories (we generally drop the words “topological” and “quantum”) that we call étale-locally-oriented and étale-locally-spin. Étale-locally-oriented and étale-locally-spin field theories admit a natural notion of “reflection-positivity” (defined in terms of a certain “integration” map taking in an étale-locally-oriented or -spin field theory and producing an unoriented \mathbb{R} -linear field theory). With this technology in place, our main result is the following version of the spin-statistics theorem:

Theorem 0.1 *Every once-extended étale-locally-spin reflection-positive topological quantum field theory is hermitian (hence unitary) and satisfies spin-statistics.*

By definition, a field theory is *unextended* if it is defined in codimensions 0 and 1, and *once-extended* if it is defined in codimensions 0, 1, and 2. [Corollary 4.8](#), which we prove only in outline, extends [Theorem 0.1](#) to more-than-once-extended field theories. Freed and Hopkins prove a similar spin-statistics theorem in [\[12, Theorem 11.3\]](#), but there are notable differences between the approach used there and the one used in this paper.

As a warm-up to [Theorem 0.1](#), in [Section 1](#) we develop in detail the notions of étale-local orientation and reflection-positivity in the context of unextended field theories. The following analog of [Theorem 0.1](#) follows almost immediately from the definitions:

Theorem 0.2 *Every unextended étale-locally-oriented reflection-positive field theory is hermitian.*

The parallel between [Theorem 0.1](#) and [Theorem 0.2](#) is an indication of the second main theme of this paper, which is to argue that hermiticity and spin-statistics phenomena arise from the same source. Note also that we reverse part of the logic from the standard spin-statistics theorem: as usually presented, hermiticity is a required assumption in order to imply spin-statistics; in our version, hermiticity and spin-statistics are both forced by reflection-positivity.

In order to define étale-locally-oriented manifolds, we consider local structures on manifolds that range over not (as in the case of orientations) sets, but schemes over \mathbb{R} . There are precisely two local structures that are étale-locally-over- $\text{Spec}(\mathbb{R})$ isomorphic to orientations. The two versions of étale-local-orientations are usual-orientations and hermitian structures; the latter are characterized by the property that the scheme of hermitian structures on a point is $\text{Spec}(\mathbb{C})$ and that the restriction map

$$\{\text{hermitian structures on } [0, 1]\} \rightarrow \{\text{hermitian structures on } \{0, 1\}\}$$

is the “antidiagonal” map $\text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(\mathbb{C}) \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\mathbb{C})$ sending λ to $(\lambda, \bar{\lambda})$. Hermitian structures owe their existence to the fact that the absolute Galois group of \mathbb{R} happens to be the same as the group $\pi_0 \text{O}(\infty)$ of connected components of the orthogonal group.

Each étale-local-orientation leads to a version of étale-locally-oriented field theory: in addition to the usual (unextended) oriented bordism category $\text{BORD}_{d-1,d}^{\text{Or}}$, there is a hermitian bordism category $\text{BORD}_{d-1,d}^{\text{Her}}$ which is not a category but rather a stack of categories over $\text{Spec}(\mathbb{R})$; the two types of field theories are symmetric monoidal functors of stacks of categories $\text{BORD}_{d-1,d}^{\text{Or}} \rightarrow \text{VECT}_{\mathbb{R}}$ and $\text{BORD}_{d-1,d}^{\text{Her}} \rightarrow \text{VECT}_{\mathbb{R}}$, where $\text{VECT}_{\mathbb{R}}$ is enhanced to the stack of categories QCOH . As such, our notion of étale-locally-oriented field theory involves infusing both the source and target categories with \mathbb{R} -algebraic geometry. The two versions unpack to \mathbb{R} -linear oriented field theories and to hermitian field theories in the usual sense.

Our definition of étale-locally-spin structures requires a categorification of (some basic notions from) real algebraic geometry. We begin this program in [Section 2](#). Our main contribution here is to categorify the notion of field and to interpret Deligne’s existence of fiber functors [\[9\]](#) as asserting that the categorified algebraic closure of \mathbb{R} is not \mathbb{C} but rather the category $\text{SUPERVECT}_{\mathbb{C}}$ of complex supervector spaces. (As we will use a slight modification of the main result of [\[9\]](#), we include a complete proof.)

Remark 0.3 As is already apparent, we will be working both with fields in the sense of commutative algebra and field theories in the sense of physics, and English includes an unfortunate terminological conflict. We don’t have a good solution to this problem, but will stick to the following convention: “field” used as a noun means “field in the

sense of algebra”; “field theory” means “(classical or quantum) functorial topological field theory in the sense of physics”.

We also prove that the extension $\text{VECT}_{\mathbb{R}} \hookrightarrow \text{SUPERVECT}_{\mathbb{C}}$ is Galois, and use this fact to categorify the notion of étale-local. There are precisely eight types of étale-locally-spin structures, of which one is distinguished by the following coincidence: the categorified absolute Galois group of \mathbb{R} is canonically equivalent to the Picard groupoid $\pi_{\leq 1}\text{O}(\infty)$. This distinguished version incorporates both hermiticity and spin-statistics phenomena. In summary, we find that the second row of the following table is a categorification of the first:

algebraic closure	tangential structure	Galois group	physical phenomenon
$\mathbb{R} \hookrightarrow \mathbb{C}$	$\text{SO}(d) \hookrightarrow \text{O}(d)$	$\text{Gal}(\mathbb{C}/\mathbb{R}) = \pi_0\text{O}(\infty)$	hermiticity
$\text{VECT}_{\mathbb{R}} \hookrightarrow \text{SUPERVECT}_{\mathbb{C}}$	$\text{Spin}(d) \rightarrow \text{O}(d)$	$\text{Gal}(\text{SUPERVECT}_{\mathbb{C}}/\mathbb{R}) = \pi_{\leq 1}\text{O}(\infty)$	spin-statistics

Our categorification result suggests the following conjecture:

Conjecture 0.4 *There is an infinitely categorified version of commutative algebra, and in it the infinitely categorified absolute Galois group of \mathbb{R} is $\text{O}(\infty)$.*

Remark 0.5 The papers Ganter and Kapranov [13] and Kapranov [15] suggest that rather than $\text{O}(\infty)$, it is the sphere spectrum that controls supermathematics. Very low homotopy groups cannot distinguish between various important spectra. The connection with topological quantum field theory focused on in this paper provides a reason to prefer $\text{O}(\infty)$.

We prove [Theorem 0.1](#) in [Section 3](#), which also contains examples of various types of étale-locally-spin field theories. We end the paper in [Section 4](#) by outlining how to extend our étale-locally-structured cobordism categories to the fully-extended ∞ -categorical world of Lurie [20].

1 Oriented, hermitian, and unitary field theories

This section serves as an extended warm-up to the remainder of the paper. We will develop in a 1-categorical setting the notions of “étale-locally-oriented” and “reflection-positive” and prove [Theorem 0.2](#), which asserts that étale-locally-oriented reflection-positive topological quantum field theories are necessarily hermitian.

The functorial framework for quantum field theory, as formulated by Atiyah and Segal in [1; 23], is well-known. Fix a dimension d and construct a symmetric monoidal

category $\text{BORD}_{d-1,d}$ whose objects are $(d-1)$ -dimensional closed smooth manifolds, morphisms are d -dimensional smooth cobordisms up to isomorphism, and the symmetric monoidal structure is disjoint union. An (unextended) *unoriented* or *unstructured* \mathbb{R} -linear d -dimensional functorial topological quantum field theory is a symmetric monoidal functor $\text{BORD}_{d-1,d} \rightarrow \text{VECT}_{\mathbb{R}}$. We will henceforth drop the words “functorial topological quantum”.

In general, one does not care simply about unstructured field theories. Let MAN_d denote the site of d -dimensional (possibly open) manifolds and local diffeomorphisms, with covers the surjections. If \mathcal{X} is a category with limits, an \mathcal{X} -valued local structure is a sheaf $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$. A local structure is *topological* if it takes isotopic (among local diffeomorphisms) maps of manifolds to equal morphisms in \mathcal{X} .

The reason for considering local structures valued in general categories is because, in examples, the collection of \mathcal{G} -structures on a manifold M is not just a set but carries more algebraic or analytic structure. For example, Stolz and Teichner [24] require local structures valued in supermanifolds. We will focus on the case when \mathcal{G} is valued in the category $\text{SCH}_{\mathbb{R}}$ of schemes over \mathbb{R} . (In fact, all of our examples will take values in the subcategory $\text{AFSCH}_{\mathbb{R}}$ of affine schemes.)

The following is an easy exercise:

Lemma 1.1 *Suppose $d \geq 1$. There are precisely two isotopy classes of local diffeomorphisms $\mathbb{R}^d \rightarrow \mathbb{R}^d$ (the identity and orientation-reversal), and so if \mathcal{G} is an \mathcal{X} -valued topological local structure, then $\mathcal{G}(\mathbb{R}^d)$ has an action by $\mathbb{Z}/2$. The assignment $\mathcal{G} \mapsto \mathcal{G}(\mathbb{R}^d)$ gives an equivalence of categories between the category of \mathcal{X} -valued topological local structures and the category $\mathcal{X}^{\mathbb{Z}/2}$ of objects in \mathcal{X} equipped with a $\mathbb{Z}/2$ -action. \square*


Example 1.2 The topological local structure \mathcal{G}_X corresponding to a $\mathbb{Z}/2$ -set $X \in \text{SETS}^{\mathbb{Z}/2}$ can be constructed as follows. For any manifold M , let $\text{Or}_M \rightarrow M$ denote the orientation double cover; then $\mathcal{G}_X(M) = \text{maps}_{\mathbb{Z}/2}(\text{Or}_M, X)$, where $\text{maps}_{\mathbb{Z}/2}$ denotes continuous $\mathbb{Z}/2$ -equivariant functions. If \mathcal{X} has limits, then for $X \in \mathcal{X}^{\mathbb{Z}/2}$ the formula $\text{maps}_{\mathbb{Z}/2}(\text{Or}_M, X)$ continues to make sense, and again defines the topological local structure corresponding to X .

The most important example is when $X = \mathbb{Z}/2$ is the *trivial* $\mathbb{Z}/2$ -torsor given by the translation action of $\mathbb{Z}/2$ on itself. Then $\mathcal{G}_{\mathbb{Z}/2} = \text{Or}$ is the sheaf $\text{Or}(M) = \{\text{orientations of } M\}$.

Given a SETS -valued topological local structure \mathcal{G} , there is a \mathcal{G} -structured bordism category $\text{BORD}_{d-1,d}^{\mathcal{G}}$, an object of which consists of a closed $(d-1)$ -manifold N

together with an element of $\mathcal{G}(N \times \mathbb{R})$, and whose morphisms are d -dimensional cobordisms similarly equipped with \mathcal{G} -structure. If \mathcal{G} is a SETS-valued topological local structure, a \mathcal{G} -structured \mathbb{R} -linear d -dimensional field theory is a symmetric monoidal functor $\text{BORD}_{d-1,d}^{\mathcal{G}} \rightarrow \text{VECT}_{\mathbb{R}}$. It will be useful to unpack the construction of $\text{BORD}_{d-1,d}^{\mathcal{G}}$ in order to have a more explicit description of \mathcal{G} -structured field theories. The following logic is used in [20, Section 3.2] to reduce the “ \mathcal{G} -structured cobordism hypothesis” to the unstructured case; see also [22, Section 3.5].


Let $\text{SPANS}(\text{SETS})$ denote the symmetric monoidal category whose objects are sets and whose morphisms are isomorphism classes of *correspondences*, ie diagrams of shape $X \leftarrow A \rightarrow Y$; composition is by fibered product and the symmetric monoidal structure is by cartesian product. A \mathcal{G} -structured classical field theory is a symmetric monoidal functor $\text{BORD}_{d-1,d}^{\mathcal{G}} \rightarrow \text{SPANS}(\text{SETS})$. Every SETS-valued topological local structure \mathcal{G} defines an unstructured classical field theory $\tilde{\mathcal{G}}: \text{BORD}_{d-1,d} \rightarrow \text{SPANS}(\text{SETS})$:



Functoriality for $\tilde{\mathcal{G}}: \text{BORD}_{d-1,d} \rightarrow \text{SPANS}(\text{SETS})$ follows from the sheaf axiom for \mathcal{G} .

Unpacking the definitions results in the following:

Lemma 1.3 *Let $\text{SPANS}(\text{SETS}; \text{VECT}_{\mathbb{R}})$ denote the symmetric monoidal category whose objects are pairs (X, V) where $X \in \text{SETS}$ and V is a vector bundle over X , and for which a morphism from (X, V) to (Y, W) is an isomorphism class of diagrams $X \xleftarrow{f} A \xrightarrow{g} Y$ together with a vector bundle map $f^*V \rightarrow g^*W$. Then a \mathcal{G} -structured field theory is the same data as a choice of lift:*



Suppose that \mathcal{G} is a topological local structure valued not in SETS but in $\text{SCH}_{\mathbb{R}}$. Our strategy will be to take Lemma 1.3 as the model for the definition of \mathcal{G} -structured field theory. To do this, note that $\text{VECT}_{\mathbb{R}}$ is naturally an object of \mathbb{R} -algebraic geometry. Indeed, there is a stack of categories on $\text{SCH}_{\mathbb{R}}$, namely $\text{QCOH}: \text{Spec}(A) \mapsto \text{MOD}_A$,

whose category of global sections is nothing but $\text{QCOH}(\text{Spec}(\mathbb{R})) = \text{VECT}_{\mathbb{R}}$. We can therefore define:

Definition 1.4 Let \mathcal{G} be a topological local structure valued in schemes over \mathbb{R} , thought of as a classical field theory

$$\tilde{\mathcal{G}}: \text{BORD}_{d-1,d} \rightarrow \text{SPANS}(\text{SCH}_{\mathbb{R}}).$$

Let $\text{SPANS}(\text{SCH}_{\mathbb{R}}; \text{QCOH})$ denote the symmetric monoidal category whose objects are pairs (X, V) where X is a scheme over \mathbb{R} and $V \in \text{QCOH}(X)$, in which a morphism from (X, V) to (Y, W) is (an isomorphism class of) a correspondence of schemes $X \xleftarrow{f} A \xrightarrow{g} Y$ together with a map of quasicoherent sheaves $f^*V \rightarrow g^*W$, in which composition is by fibered product, and in which the symmetric monoidal structure is $\times_{\text{Spec}(\mathbb{R})}$. A \mathcal{G} -structured field theory is a choice of lift:

$$\begin{array}{ccc} & & \text{SPANS}(\text{SCH}_{\mathbb{R}}; \text{QCOH}) \\ & \nearrow \text{dashed arrow} & \downarrow \text{forget the QCOH-data} \\ \text{BORD}_{d-1,d} & \xrightarrow{\tilde{\mathcal{G}}} & \text{SPANS}(\text{SCH}_{\mathbb{R}}) \end{array}$$

Any topological local structure \mathcal{G} valued in SETS defines a topological local structure, which we will also call \mathcal{G} , valued in $\text{SCH}_{\mathbb{R}}$, via the symmetric monoidal inclusion $\text{SETS} \hookrightarrow \text{SCH}_{\mathbb{R}}$, $S \mapsto S \times \text{Spec}(\mathbb{R})$. In this case, the notion of \mathcal{G} -structured field theory from Definition 1.4 agrees with the usual notion in terms of symmetric monoidal functors $\text{BORD}_{d-1,d}^{\mathcal{G}} \rightarrow \text{VECT}_{\mathbb{R}}$, since $\text{QCOH}(S \times \text{Spec}(\mathbb{R})) = \{\text{real vector bundles on } S\}$.

We will focus on four examples of topological local structures \mathcal{G} valued in $\text{SCH}_{\mathbb{R}}$, two of which come from topological local structures valued in SETS . We will unpack a bit about the values of \mathcal{G} -structured field theories in all four cases to make everything explicit.

Example 1.5 An unstructured or unoriented field theory is a “ $\text{Spec}(\mathbb{R})$ -structured” one, where $\text{Spec}(\mathbb{R})(M) = \text{Spec}(\mathbb{R})$ for all manifolds M . Let Z be an unstructured field theory. If M a closed d -dimensional manifold, then $Z(M) \in \mathcal{O}(\text{Spec}(\mathbb{R})) = \mathbb{R}$. If N is a closed $(d-1)$ -dimensional manifold, then $Z(N) \in \text{QCOH}(\text{Spec}(\mathbb{R})) = \text{VECT}_{\mathbb{R}}$. Consider the macaroni cobordisms $N \times \smile: N \sqcup N \rightarrow \emptyset$ and $N \times \frown: \emptyset \rightarrow N \sqcup N$. The first defines a symmetric pairing $Z(N \times \smile): Z(N) \otimes Z(N) \rightarrow \mathbb{R}$ and the second a symmetric copairing $\mathbb{R} \rightarrow Z(N) \otimes Z(N)$. The zig-zag equations $\mathcal{S} = \mathcal{I}$ and $\mathcal{Z} = \mathcal{I}$ require this pairing and copairing to be inverse to each other, and are equivalent to making $V = Z(N)$ into a symmetrically self-dual vector space over \mathbb{R} , ie we have $\varphi: V \xrightarrow{\sim} V^*$ with $\varphi^* \circ \varphi = \text{id}_V$.

Example 1.6 An oriented field theory is one with topological local structure $\text{Or} = \mathcal{G}_{\mathbb{Z}/2}$ from Example 1.2, thought of as being valued in $\text{SCH}_{\mathbb{R}}$ via $S \mapsto S \times \text{Spec}(\mathbb{R})$. Orientations are distinguished among all topological local structures by Lemma 1.1: they correspond to the trivial $\mathbb{Z}/2$ -torsor. We will review the basic structure enjoyed by an oriented field theory Z .

Let M be a connected closed d -dimensional manifold. Then $Z(M)$ is a function on $\text{Or}(M) \times \text{Spec}(\mathbb{R})$. If M is nonorientable, then $\text{Or}(M) = \emptyset$ and $Z(M)$ is no data. If M is orientable, then $\text{Or}(M) \times \text{Spec}(\mathbb{R}) \cong \text{Spec}(\mathbb{R}) \sqcup \text{Spec}(\mathbb{R})$, the two points corresponding to the two orientations of M , and $Z(M)$ is an element of $\mathcal{O}(\text{Spec}(\mathbb{R}) \sqcup \text{Spec}(\mathbb{R})) = \mathbb{R} \times \mathbb{R}$, ie a pair of numbers (indexed by the two orientations of M).

Suppose now that N is a closed connected $(d-1)$ -dimensional manifold. Again if N is nonorientable, $\text{Or}(N)$ is empty and Z assigns no data. If N is orientable, $Z(N)$ is a sheaf on $\text{Or}(N) \times \text{Spec}(\mathbb{R}) \cong \text{Spec}(\mathbb{R}) \sqcup \text{Spec}(\mathbb{R})$, ie a pair (V, V') of real vector spaces, one for each orientation of N . These vector spaces are not independent. Rather, the macaroni cobordisms $N \times \mathcal{J}: N \sqcup N \rightarrow \emptyset$ and $N \times \mathcal{C}: \emptyset \rightarrow N \sqcup N$ each admit two orientations, which induce orientations of their boundaries such that the two copies of N have opposite orientations. Some definition-unpacking shows that the data of $Z(N \times \mathcal{J})$ is nothing but a linear map $V \otimes_{\mathbb{R}} V' \rightarrow \mathbb{R}$, and the data of $Z(N \times \mathcal{C})$ is a linear map $\mathbb{R} \rightarrow V \otimes_{\mathbb{R}} V'$. The zig-zag equations assert that $Z(N \times \mathcal{J})$ and $Z(N \times \mathcal{C})$ make V and V' into dual vector spaces.

Example 1.7 Lemma 1.1 distinguishes a second topological local structure valued in $\text{SCH}_{\mathbb{R}}$. Specifically, there is a canonical nontrivial $\mathbb{Z}/2$ -torsor over $\text{Spec}(\mathbb{R})$, namely $\text{Spec}(\mathbb{C})$ with the complex conjugation action. We will suggestively write $\text{Her}: \text{MAN}_d \rightarrow \text{SCH}_{\mathbb{R}}$ for this topological local structure, and call $\text{Her}(M)$ the scheme of hermitian structures on M . One easily sees that for any manifold M ,

$$\text{Her}(M) = \text{Or}(M) \times_{\mathbb{Z}/2} \text{Spec}(\mathbb{C}),$$

where $\mathbb{Z}/2$ acts on $\text{Or}(M)$ by orientation-reversal and on $\text{Spec}(\mathbb{C})$ by complex conjugation, and $\times_{\mathbb{Z}/2}$ denotes the coequalizer of these actions. A hermitian field theory is étale-locally-oriented in the sense that Her and Or are both valued in schemes étale over $\text{Spec}(\mathbb{R})$ and are étale-locally isomorphic as topological local structures over $\text{Spec}(\mathbb{R})$, since they pull back to isomorphic topological local structures along $\text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(\mathbb{R})$. Since there are precisely two $\mathbb{Z}/2$ -torsors over $\text{Spec}(\mathbb{R})$, there are precisely two topological local structures étale-locally isomorphic to Or , ie precisely two kinds of étale-locally-oriented field theory.

We now justify the name “hermitian”. Suppose that Z is a Her-structured field theory and M is a closed d -dimensional manifold. If M is not orientable, then $\text{Her}(M) = \emptyset$ is the empty scheme and $Z(M)$ is no data. If M is orientable and nonempty, then $\text{Her}(M)$ is noncanonically isomorphic to the disjoint union of $2^{|\pi_0 M| - 1}$ copies of $\text{Spec}(\mathbb{C})$. In particular, if M is connected and orientable, then either orientation of M determines an isomorphism $\text{Her}(M) \cong \text{Spec}(\mathbb{C})$. Thus, either choice of orientation identifies $Z(M) \in \mathcal{O}(\text{Her}(M))$ with a complex number. The two choices of orientation determine isomorphisms that differ by complex conjugation. So one can think of Z as assigning to every oriented manifold a complex number, subject to the condition that orientation-reversal agrees with complex conjugation. Finally, if $M = \emptyset$, then $\text{Her}(M) = \text{Spec}(\mathbb{R})$ and $Z(M) = 1$.

Suppose now that N is a closed connected $(d-1)$ -dimensional manifold. Again, if N is nonorientable, then $\text{Her}(N) = \emptyset$ and $Z(N)$ is no data. If N is orientable, $Z(N)$ is a vector bundle on $\text{Her}(N) \cong \text{Spec}(\mathbb{C})$, ie a complex vector space. The values of the macaroni $Z(N \times \curvearrowright)$ and $Z(N \times \curveleft)$ now are bundles of linear maps over $\text{Her}(N \times \curvearrowright) \cong \text{Her}(N \times \curveleft) \cong \text{Spec}(\mathbb{C})$. The domain and codomain of $Z(N \times \curvearrowright)$ are given by pulling back $Z(N \sqcup N)$ and $Z(\emptyset)$ along the restrictions

$\text{Her}(N \times \curvearrowright) \rightarrow \text{Her}(N \sqcup N) = \text{Her}(N) \times_{\text{Spec } \mathbb{R}} \text{Her}(N)$ and $\text{Her}(N \times \curveleft) \rightarrow \text{Her}(\emptyset)$,
and similarly for $Z(N \times \curveleft)$. Unpacking gives

$$\begin{aligned} Z(N \times \curvearrowright) &\in \text{hom}_{\mathbb{R}}(Z(N \times \{\text{pt}\}) \otimes_{\mathbb{R}} Z(N \times \{\text{pt}\}), \mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}, \\ Z(N \times \curveleft) &\in \text{hom}_{\mathbb{R}}(\mathbb{R}, Z(N \times \{\text{pt}\}) \otimes_{\mathbb{R}} Z(N \times \{\text{pt}\})) \otimes_{\mathbb{R}} \mathbb{C}. \end{aligned}$$

The restriction map

$$\text{Spec}(\mathbb{C}) = \text{Her}(N \times \curvearrowright) \rightarrow \text{Her}(N) \times_{\text{Spec } \mathbb{R}} \text{Her}(N) = \text{Spec}(\mathbb{C}) \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\mathbb{C})$$

is the antidiagonal map $\lambda \mapsto (\lambda, \bar{\lambda})$, and so $Z(N \times \curvearrowright)$ is a sesquilinear pairing on $Z(N)$. It follows from the zig-zag equations that $Z(N \times \curvearrowright)$ and $Z(N \times \curveleft)$ identify the \mathbb{C} -linear dual vector space $Z(N)^*$ to $Z(N) \in \text{Vect}_{\mathbb{C}}$ with the complex conjugate space $\overline{Z(N)}$. Finally, the symmetry of $N \times \curvearrowright$ translates into the requirement that the sesquilinear pairing on $Z(N)$ is symmetric, or equivalently the isomorphism $\varphi: Z(N)^* \xrightarrow{\sim} \overline{Z(N)}$ satisfies $\bar{\varphi}^* \circ \varphi = \text{id}$. It is in this sense that hermitian field theories are “hermitian”.

Example 1.8 In addition to $\text{Her}: \text{MAN}_d \rightarrow \text{SCH}_{\mathbb{R}}$, there is another topological local structure whose value on \mathbb{R}^d is $\text{Spec}(\mathbb{C})$, namely the one corresponding via [Lemma 1.1](#) to $\text{Spec}(\mathbb{C})$ with the trivial $\mathbb{Z}/2$ -action. We will simply call this topological local structure $\text{Spec}(\mathbb{C})$. It satisfies $\text{Spec}(\mathbb{C})(M) = \text{Spec}(\mathbb{C})^{\pi_0 M}$ for every manifold M . When one unpacks the notion of $\text{Spec}(\mathbb{C})$ -structured field theory, one finds that they

are nothing but *complex-linear* unstructured field theories. For example, the values of $\text{Spec}(\mathbb{C})$ -structured field theories on closed connected $(d-1)$ - and d -dimensional manifolds are objects of $\text{QCOH}(\text{Spec}(\mathbb{C})) = \text{VECT}_{\mathbb{C}}$ and elements of $\mathcal{O}(\text{Spec}(\mathbb{C})) = \mathbb{C}$, respectively.

Example 1.7 provided one of two reasons why hermitian field theories are distinguished: they correspond to the unique nontrivial torsor over $\text{Spec}(\mathbb{R})$ for the group $\mathbb{Z}/2 = \pi_0 \text{hom}_{\text{MAN}_d}(\mathbb{R}^d, \mathbb{R}^d)$. **Theorem 0.2** provides the second reason, by asserting that of the two types of étale-locally-oriented field theories, only hermiticity is compatible with reflection-positivity. We now define reflection-positivity and prove **Theorem 0.2**.

Definition 1.9 A d -dimensional unstructured (ie $\text{Spec}(\mathbb{R})$ -structured) field theory $Z: \text{BORD}_{d-1,d} \rightarrow \text{VECT}_{\mathbb{R}}$ is *reflection-positive* if the nondegenerate symmetric pairing $Z(N \times \cdot): Z(N) \otimes Z(N) \rightarrow \mathbb{R}$ is positive-definite for every closed $(d-1)$ -dimensional manifold N .

Most of the physics literature, including Atiyah’s original definition of functorial topological field theory from [1], includes hermiticity directly in the definition of quantum field theory. As such, reflection-positivity is usually posed as the requirement that the hermitian form on the complex vector space $Z(N)$ should be positive-definite. For nontopological quantum field theories defined on Minkowski $\mathbb{R}^{d-1,1}$, reflection-positivity is a stronger condition assuring the existence of an analytic continuation to imaginary time $\mathbb{R}^{d-1} \times i\mathbb{R}_{\geq 0}$, and reflection refers to reflection in the time axis. Positive-definiteness of the Hilbert space is what remains when interpreting this stronger condition for topological field theories.

From the point of view of this paper, the nonhermitian version of reflection-positivity in **Definition 1.9** is the most primitive. The hermitian version arises as follows. Suppose first that Z is not hermitian but oriented. One can produce an unstructured field theory $\int_{\text{Or}} Z$ from Z by integrating out the choice of orientation:

$$\int_{\text{Or}} Z: M \mapsto \int_{\sigma \in \text{Or}(M)} Z(M, \sigma).$$

Here the integral is a finite sum of numbers when M is d -dimensional and a finite direct sum when $\dim M < d$. In particular, for N a connected $(d-1)$ -dimensional manifold, $(\int_{\text{Or}} Z)(N) = Z(N) \oplus Z(N)^*$ with the obvious symmetric pairing.

Let Z^* denote the orientation-reversal of the field theory Z . There is a canonical equivalence $\int_{\text{Or}} Z \cong \int_{\text{Or}} Z^*$. It follows that \int_{Or} makes sense not just for oriented field theories but for any étale-locally-oriented field theory. Indeed, suppose Z is not

oriented but hermitian. Using the isomorphism $\text{Her} \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\mathbb{C}) \cong \text{Or} \times \text{Spec}(\mathbb{C})$, one sees that the base-changed field theory $Z_{\mathbb{C}} = Z \otimes_{\mathbb{R}} \mathbb{C}$ is naturally oriented and \mathbb{C} -linear, and so $\int_{\text{Or}} Z_{\mathbb{C}}$ makes sense as a \mathbb{C} -linear unstructured field theory. But the hermiticity of Z defines a Galois action on $\int_{\text{Or}} Z_{\mathbb{C}}$, describing how it descends to an \mathbb{R} -linear unstructured field theory $\int_{\text{Or}} Z$. One finds that, for Z a hermitian field theory and N a connected $(d-1)$ -dimensional manifold, $(\int_{\text{Or}} Z)(N)$ is nothing but the underlying real vector space of the hermitian vector space $Z(N)$; the symmetric pairing is twice the real part of the hermitian pairing on $Z(N)$.

The usual notion of reflection-positivity is then captured by the following:

Definition 1.10 An étale-locally-oriented field theory Z is *reflection-positive* if the unoriented field theory $\int_{\text{Or}} Z$ is reflection-positive. A field theory is *unitary* if it is reflection-positive and hermitian.

With this notion, the proof of [Theorem 0.2](#) is immediate:

Proof of Theorem 0.2 If V is a nonzero real vector space, $V \oplus V^*$ is never positive-definite. \square

Remark 1.11 One can also integrate a $\text{Spec}(\mathbb{C})$ -structured field theory to a $\text{Spec}(\mathbb{R})$ -structured one. One finds that $\int_{\text{Spec}(\mathbb{C})} c = 2 \text{Re}(c)$ for $c \in \mathcal{O}(\text{Spec}(\mathbb{C}))$, and that the integral of a complex vector space $V \in \text{VECT}_{\mathbb{C}}$ is the underlying real vector space of V . If Z is a $\text{Spec}(\mathbb{C})$ -structured field theory, then $Z(N \times \mathbb{D})$ is a \mathbb{C} -linear symmetric pairing on the complex vector space $Z(N)$, and $\int_{\text{Spec}(\mathbb{C})} Z(N \times \mathbb{D})$ is twice its real part, thought of as a symmetric pairing on the real vector space $\int_{\text{Spec}(\mathbb{C})} Z(N)$. The real part of a complex-linear symmetric pairing is never positive-definite.

2 A categorified Galois extension

[Section 1](#) illustrated the important role that algebraic geometry and Galois theory play in explaining the origin of hermitian phenomena in quantum field theory. The goal of this section and the next is to tell a similar story concerning super phenomena of fermions and spinors. Explicitly, \mathbb{C} appeared because it is the algebraic closure of \mathbb{R} . This section will explain that $\text{SUPERVECT}_{\mathbb{C}}$ is the categorified algebraic closure of $\text{VECT}_{\mathbb{R}}$. This is essentially Deligne’s “existence of super fiber functors” theorem from [\[9\]](#). We state this result as [Theorem 2.7](#) and provide details of its proof, as our phrasing is somewhat different from that of [\[9\]](#).

A convenient setting for categorified \mathbb{R} -linear algebra is provided by the bicategory $\text{PRES}_{\mathbb{R}}$ of \mathbb{R} -linear locally presentable categories, \mathbb{R} -linear cocontinuous functors,

and natural transformations: direct sums play the role of addition and quotients play the role of subtraction. Two of the many ways that $\text{PRES}_{\mathbb{R}}$ is convenient are that it admits all limits and colimits [5] and that it has a natural symmetric monoidal structure $\boxtimes = \boxtimes_{\mathbb{R}}$ satisfying a hom-tensor adjunction [16]. The unit object for \boxtimes is $\text{VECT}_{\mathbb{R}}$. Basic examples of \mathbb{R} -linear locally presentable categories include the categories MOD_A of A -modules for any \mathbb{R} -algebra A ; the tensor product enjoys $\text{MOD}_A \boxtimes \text{MOD}_B \simeq \text{MOD}_{A \otimes B}$.

Definition 2.1 A categorified commutative \mathbb{R} -algebra is a symmetric monoidal object in $\text{PRES}_{\mathbb{R}}$.

We embed noncategorified commutative \mathbb{R} -algebras among categorified commutative \mathbb{R} -algebras with the following lemma, whose proof is a straightforward exercise (see [8, Proposition 2.3.9]):

Lemma 2.2 The assignment taking a commutative \mathbb{R} -algebra R to the categorified commutative \mathbb{R} -algebra $(\text{MOD}_R, \otimes_R)$ and an \mathbb{R} -algebra homomorphism $f: R \rightarrow S$ to extension of scalars $\otimes_R S: \text{MOD}_R \rightarrow \text{MOD}_S$ defines a fully faithful embedding of the category of commutative \mathbb{R} -algebras into the bicategory of categorified commutative \mathbb{R} -algebras. □

We turn now to categorifying the notion of algebraic closure. Algebraic closures of fields are determined by a weak universal property ranging over only finite-dimensional algebras. Summarizing the story over \mathbb{R} , we have:

- Lemma 2.3** (0) \mathbb{C} is a nonzero finite-dimensional commutative \mathbb{R} -algebra.
- (1) Every map $\mathbb{C} \rightarrow A$ of nonzero finite-dimensional commutative \mathbb{R} -algebras is an injection.
 - (2) If A is a nonzero finite-dimensional commutative \mathbb{R} -algebra, then there exists a map $A \rightarrow \mathbb{C}$ of commutative \mathbb{R} -algebras.
 - (3) Items (0)–(2) determine \mathbb{C} uniquely up to nonunique isomorphism. □

Of course, (0)–(1) are equivalent to the statement that \mathbb{C} is a *field*, and (2) is equivalent to the statement that \mathbb{C} is *algebraically closed*. We categorify these notions in turn.

Definition 2.4 A strongly generating set in an \mathbb{R} -linear locally presentable category \mathcal{C} is a set of objects in \mathcal{C} that generate \mathcal{C} under colimits. The category \mathcal{C} is *finite-dimensional* if it admits a finite strongly generating set $\{C_1, \dots, C_n\}$ such that all hom-spaces between generators $\text{hom}(C_i, C_j)$ are finite-dimensional and moreover every

generator C_i is *compact projective* in \mathcal{C} , in the sense that $\text{hom}(C_i, -): \mathcal{C} \rightarrow \text{VECT}_{\mathbb{R}}$ is cocontinuous.

A categorified commutative \mathbb{R} -algebra $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ is *finite-dimensional* as a categorified commutative \mathbb{R} -algebra if the underlying \mathbb{R} -linear category of \mathcal{C} is finite-dimensional and moreover every projective object $P \in \mathcal{C}$ is dualizable.

Compact projectivity, sometimes called tininess, is a strong but reasonable finiteness condition to impose on an object. There are many definitions of projectivity that agree for abelian categories but diverge for locally presentable but not necessarily abelian categories; ours is one of the stronger possible choices. If \mathcal{C} is a finite-dimensional \mathbb{R} -linear locally presentable category, then \mathcal{C} is automatically equivalent to the category MOD_A of modules for a finite-dimensional associative algebra A (eg one can take $A = \text{End}(\bigoplus_i C_i)$).

Finite-dimensionality as a categorified algebra is stronger than just finite-dimensionality of the underlying category. The condition that compact projectivity implies dualizability expresses a compatibility between internal and external notions of finite-dimensionality in a symmetric monoidal category, which otherwise might badly diverge [18]. Indeed, $P \in \text{MOD}_A$ is compact projective exactly when the functor $\otimes_{\mathbb{R}} P: \text{VECT}_{\mathbb{R}} \rightarrow \text{MOD}_A$ has a right adjoint of the form $\otimes_A P^{\vee}$ for some left A -module P^{\vee} , whereas, for $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ a symmetric monoidal category, $P \in \mathcal{C}$ is dualizable when the functor $\otimes P: \mathcal{C} \rightarrow \mathcal{C}$ has a right adjoint of the form $\otimes P^*$ for some $P^* \in \mathcal{C}$.

To check that $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ is finite-dimensional as a categorified commutative \mathbb{R} -algebra, it suffices to check that the underlying \mathbb{R} -linear category \mathcal{C} is finite-dimensional and that each generator C_i is dualizable.

Definition 2.4 explains how to categorify item (0) from **Lemma 2.3**. With it in hand, we may categorify the notion of algebraically closed field by following items (1)–(2):

Definition 2.5 A *categorified field* is a nonzero finite-dimensional categorified commutative \mathbb{R} -algebra $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ such that every 1-morphism $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \dots)$ of nonzero categorified commutative \mathbb{R} -algebras is faithful and injective on isomorphism classes of objects.

A finite-dimensional categorified field $(\mathcal{C}, \otimes, \dots)$ is *algebraically closed* if for every nonzero finite-dimensional categorified commutative \mathbb{R} -algebra $(\mathcal{B}, \otimes, \dots)$, there exists a 1-morphism $F: (\mathcal{B}, \otimes_{\mathcal{B}}, \dots) \rightarrow (\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ of categorified commutative \mathbb{R} -algebras.

Lemma 2.6 A *finite-dimensional commutative \mathbb{R} -algebra R is a field if and only if $(\text{MOD}_R, \otimes_R, \dots)$ is a categorified field.*

Proof It is clear that if $(\text{MOD}_R, \otimes_R, \dots)$ is a categorified field, then R is a field, simply by using the faithfulness assumption and 1–morphisms to categorified algebras of the form $(\text{MOD}_S, \otimes_S, \dots)$.

Conversely, suppose R is a field and $F: \text{MOD}_R \rightarrow \mathcal{C}$ is any \mathbb{R} –linear functor. Suppose F is not faithful. Then there is a nonzero morphism $f: X \rightarrow Y$ in MOD_R with $F(f) = 0$. Using the fact that in MOD_R all exact sequences split, one can show that $F(\text{im}(f)) = 0$, from which it follows that $F(R) = 0$. If F is symmetric monoidal, $F(R) \cong \mathbb{1}_{\mathcal{C}}$ is the monoidal unit in \mathcal{C} , and so \mathcal{C} is the zero category. This verifies the faithfulness condition in [Definition 2.5](#).

Suppose that $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ is a finite-dimensional categorified commutative algebra over \mathbb{R} , and let $\mathbb{1}_{\mathcal{C}}$ denote its monoidal unit. Any $\lambda \in \text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}})$ defines a natural endomorphism of the identity functor on \mathcal{C} via

$$\lambda|_X = \lambda \otimes \text{id}_X: X = \mathbb{1}_{\mathcal{C}} \otimes_{\mathcal{C}} X \rightarrow \mathbb{1}_{\mathcal{C}} \otimes_{\mathcal{C}} X = X,$$

and clearly $\lambda|_{\mathbb{1}} = \lambda$. Since \mathcal{C} is finite-dimensional, it is equivalent to MOD_A for a finite-dimensional associative algebra A ; then the algebra of natural endomorphisms of the identity functor is nothing but the center $Z(A) \subseteq A$. It follows that $\text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}) \subseteq Z(A)$ is finite-dimensional. Suppose that $\mathbb{1}_{\mathcal{C}} \in \mathcal{C}$ corresponded to an infinite-dimensional A –module M_A . Then $\text{End}_A(M_A) = \text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}})$ would be infinite-dimensional, as it is the subalgebra of $\text{End}_{\mathbb{R}}(M)$ cut out by finitely many equations (imposing compatibility with multiplication by a basis in the finite-dimensional algebra A). It follows that $\mathbb{1}_{\mathcal{C}}$ corresponds to a finite-dimensional A –module, and so $\mathbb{1}_{\mathcal{C}}$ is a *compact* object in \mathcal{C} in the sense that $\text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, -): \mathcal{C} \rightarrow \text{VECT}_{\mathbb{R}}$ preserves infinite direct sums.

If R is a field, every object in MOD_R is isomorphic to $R^{\oplus \alpha}$ for some cardinal α . Let $F: (\text{MOD}_R, \otimes_R, \dots) \rightarrow \mathcal{C}$ be a cocontinuous symmetric monoidal functor. On objects it takes $R \in \text{MOD}_R$ to $\mathbb{1}_{\mathcal{C}}$, and so takes $R^{\oplus \alpha}$ to $\mathbb{1}_{\mathcal{C}}^{\oplus \alpha}$. Since $\mathbb{1}_{\mathcal{C}}$ is compact, $\text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}}, \mathbb{1}_{\mathcal{C}}^{\oplus \alpha}) = \text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}})^{\oplus \alpha}$ is $(\dim(\text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}})) \times \alpha)$ –dimensional over \mathbb{R} . Since $\dim(\text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}})) < \infty$, the cardinal α is determined by the cardinal $\dim(\text{End}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}})) \times \alpha$. This verifies the injectivity-on-objects condition in [Definition 2.5](#). □

We now describe the categorified algebraic closure of \mathbb{R} . Recall that the symmetric monoidal category $\text{SUPERVECT}_{\mathbb{C}}$ of *supervector spaces* over \mathbb{C} is by definition equivalent as a monoidal category, but not as a symmetric monoidal category, to the category $\text{REP}_{\mathbb{C}}(\mathbb{Z}/2)$ of complex representations of the group $\mathbb{Z}/2$. Let \mathbb{J} denote the sign representation, also called the *odd line*. In $\text{REP}_{\mathbb{C}}(\mathbb{Z}/2)$, the symmetry $\mathbb{J} \otimes \mathbb{J} \rightarrow \mathbb{J} \otimes \mathbb{J}$ is multiplication by $+1$; in $\text{SUPERVECT}_{\mathbb{C}}$ the symmetry is -1 . The rest of the symmetry is determined from this law by the axioms of a symmetric monoidal category.

The following is, with just a few changes of context, the main result of [9]; because of these few changes, we review the proof.

Theorem 2.7 $\text{SUPERVECT}_{\mathbb{C}}$ is the unique (up to nonunique equivalence) finite-dimensional algebraically closed categorified field over \mathbb{R} .

Proof To show that $\text{SUPERVECT}_{\mathbb{C}}$ is a categorified field, one proceeds as in the proof of Lemma 2.6. We need the following additional observation. Let $F: \text{SUPERVECT}_{\mathbb{C}} \rightarrow \mathcal{C}$ be a morphism of finite-dimensional categorified commutative \mathbb{R} -algebras, and let $\mathbb{J}_{\mathcal{C}} = F(\mathbb{J})$ denote the image of the odd line. Then $\mathbb{J}_{\mathcal{C}}$ has self-braiding -1 whereas $\mathbb{1}_{\mathcal{C}}$ has self-braiding $+1$, from which it follows that $\mathbb{1}_{\mathcal{C}}$ and $\mathbb{J}_{\mathcal{C}}$ are not isomorphic. On the other hand, tensoring with $\mathbb{J}_{\mathcal{C}}$ induces an autoequivalence of \mathcal{C} , and so $\mathbb{J}_{\mathcal{C}}$, like $\mathbb{1}_{\mathcal{C}}$, is compact and nonzero. From these facts, it follows that F is faithful and that one can recover the isomorphism type of an object $V = \mathbb{1}^{\oplus\alpha} \oplus \mathbb{J}^{\oplus\beta} \in \text{SUPERVECT}_{\mathbb{C}}$ from the vector space $\text{hom}_{\mathcal{C}}(\mathbb{1}_{\mathcal{C}} \oplus \mathbb{J}_{\mathcal{C}}, F(V))$.

We next verify that, assuming $\text{SUPERVECT}_{\mathbb{C}}$ is algebraically closed, it is the unique such category. Suppose that \mathcal{C} is another algebraically closed finite-dimensional categorified field over \mathbb{R} . Then there are symmetric monoidal functors $\mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ and $\text{SUPERVECT}_{\mathbb{C}} \rightarrow \mathcal{C}$, both faithful and injective on objects. Their composition $\text{SUPERVECT}_{\mathbb{C}} \rightarrow \mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ is full and essentially surjective as it necessarily takes $\mathbb{1}$ to $\mathbb{1}$ and \mathbb{J} to \mathbb{J} . Thus the functor $\mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ is essentially surjective and full (fullness uses that $\mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ is injective on objects).

Finally, we prove that $\text{SUPERVECT}_{\mathbb{C}}$ is algebraically closed. Let \mathcal{C} be a nonzero finite-dimensional categorified commutative \mathbb{R} -algebra. We must construct a 1-morphism $\mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$. By including $\mathcal{C} \rightarrow \mathcal{C} \boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}}$ if necessary, we may assume without loss of generality that \mathcal{C} receives a 1-morphism $\text{SUPERVECT}_{\mathbb{C}} \rightarrow \mathcal{C}$. As above, we will denote the images under this 1-morphism of $\mathbb{1}, \mathbb{J} \in \text{SUPERVECT}_{\mathbb{C}}$ by $\mathbb{1}_{\mathcal{C}}, \mathbb{J}_{\mathcal{C}}$, respectively.

We will need the following notion. Let λ be a partition of $n \in \mathbb{N}$ and V_{λ} the corresponding irrep of the symmetric group \mathbb{S}_n . Recall that, for any \mathbb{C} -linear symmetric monoidal category $(\mathcal{C}, \otimes, \dots)$ containing direct sums and splittings of idempotents, the Schur functor $S_{\lambda}: \mathcal{C} \rightarrow \mathcal{C}$ is the (nonlinear) functor $X \mapsto (X^{\otimes n} \otimes V_{\lambda})_{\mathbb{S}_n}$, where \mathbb{S}_n acts on $X^{\otimes n}$ via the symmetry on \mathcal{C} , and $(-)\mathbb{S}_n$ denotes the functor of coinvariants. S_{λ} is natural for symmetric monoidal \mathbb{C} -linear functors.

Choose a strong projective generator $P \in \mathcal{C}$. (In the notation of Definition 2.4, one can for example take $P = \bigoplus_i C_i$.) Then the underlying category of \mathcal{C} is equivalent to the category of $\text{End}_{\mathcal{C}}(P)$ -modules, and the subcategory of compact objects of \mathcal{C}

is the abelian category of finite-dimensional $\text{End}_{\mathcal{C}}(P)$ -modules. In particular, every compact object has finite length. As shown in the proof of Lemma 2.6, $\mathbb{1}_{\mathcal{C}}$ is compact, from which it follows that all dualizable objects are compact. Since P is dualizable by assumption, $P^{\otimes n}$ is also dualizable and hence compact.

We claim that there exists some λ such that $S_{\lambda}(P) = 0$. Indeed, suppose that there were not. Then, as in [9, Paragraph 1.20], the isomorphism $P^{\otimes n} \cong \bigoplus_{|\lambda|=n} V_{\lambda} \otimes S_{\lambda}(P)$ would imply that

$$\text{length}(P^{\otimes n}) \geq \sum_{|\lambda|=n} \dim V_{\lambda} \geq \left(\sum (\dim V_{\lambda})^2 \right)^{1/2} = (n!)^{1/2},$$

which grows more quickly than any geometric series. Suppose that $X, Y, M \in \mathcal{C}$ are compact objects and E is an extension of X by Y . Then, as in [9, Lemma 4.8], right exactness of the tensor functor implies

$$\text{length}(E \otimes M) \leq \text{length}(E \otimes X) + \text{length}(E \otimes Y).$$

From this, the lengths of the tensor products of simple objects, and the fact that finite-dimensional algebras admit only finitely many simple modules, one can bound the growth of $\text{length}(P^{\otimes n})$ by some geometric series.

Given a commutative algebra object $A \in \mathcal{C}$, let $\mathbb{1}_A$ and \mathbb{J}_A denote the images of $\mathbb{1}_{\mathcal{C}}$ and $\mathbb{J}_{\mathcal{C}}$ under the extension-of-scalars functor $\otimes_A: \mathcal{C} \rightarrow \{A\text{-modules in } \mathcal{C}\}$. Note that \otimes_A makes $\{A\text{-modules in } \mathcal{C}\}$ into a categorified commutative \mathbb{R} -algebra. Following [9, Proposition 2.9], we will find a nonzero commutative algebra $A \in \mathcal{C}$ such that $P \otimes A \cong \mathbb{1}_A^{\oplus r} \oplus \mathbb{J}_A^{\oplus s}$ for some $r, s \in \mathbb{N}$. Supposing we have done so, let R be the commutative superalgebra whose even part is $\text{End}(\mathbb{1}_A) \cong \text{End}(\mathbb{J}_A)$ and whose odd part is $\text{hom}(\mathbb{J}_A, \mathbb{1}_A) \cong \text{hom}(\mathbb{1}_A, \mathbb{J}_A)$, ie the “endomorphism superalgebra” of $\mathbb{1}_A$. Since P is a compact projective generator of \mathcal{C} and $P \otimes A \cong \mathbb{1}_A^{\oplus r} \oplus \mathbb{J}_A^{\oplus s}$, the symmetric monoidal category $\{A\text{-modules in } \mathcal{C}\}$ is strongly generated as a category by $\mathbb{1}_A$ and \mathbb{J}_A , and so is equivalent to the category SUPERMOD_R of R -modules in $\text{SUPERVECT}_{\mathcal{C}}$; this equivalence is then manifestly symmetric monoidal.

Suppose by induction that we have found a nonzero commutative algebra object $A \in \mathcal{C}$ such that $P \otimes A \cong \mathbb{1}_A^{\oplus r'} \oplus \mathbb{J}_A^{\oplus s'} \oplus P'$ for some $P' \in \{A\text{-modules in } \mathcal{C}\}$. Then P' is a summand of a dualizable object and hence dualizable. If $\text{Sym}^n P' = \bigwedge^n P' = 0$ for all sufficiently large n , then $P' = 0$ by [9, Corollary 1.7 and Lemma 1.17]. If on the other hand $\text{Sym}^n P' \neq 0$ for all n (resp. $\bigwedge^n P' \neq 0$ for all n), then [9, Lemma 2.8], which does not assume the category to be rigid, constructs a nonzero A -algebra A' such that $P' \otimes A' \cong \mathbb{1}_{A'} \oplus P''$ (resp. $P' \otimes A' \cong \mathbb{J}_{A'} \oplus P''$). We iterate, continually splitting off $\mathbb{1}_{A'}$ s and $\mathbb{J}_{A'}$ s. The iteration must terminate as otherwise $S_{\lambda}(P) \neq 0$ for all λ [9, Corollary 1.9].

Thus we have found a nonzero commutative superalgebra R and a morphism $\mathcal{C} \rightarrow \text{SUPERMOD}_R$ of categorified commutative \mathbb{R} -algebras. We can choose a field \mathbb{L} that receives a map from R and extend scalars further so as to build a linear cocontinuous symmetric monoidal functor $\mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{L}}$. Moreover, since $\text{End}_{\mathcal{C}}(P)$ is finite-dimensional over \mathbb{C} , the functor $\mathcal{C} \rightarrow \text{SUPERVECT}_{\mathbb{L}}$ factors through $\text{SUPERVECT}_{\mathbb{K}}$ for some intermediate field $\mathbb{C} \subseteq \mathbb{K} \subseteq \mathbb{L}$ which is finite-dimensional over \mathbb{C} . But since \mathbb{C} is algebraically closed, the only such field is $\mathbb{K} = \mathbb{C}$. \square

Remark 2.8 The fact that $\text{SUPERVECT}_{\mathbb{C}}$ is algebraically closed explains its central role in categorified representation theory [17; 13].

Remark 2.9 The categorified algebraic closure of $\overline{\mathbb{F}}_p$ is not yet known. When $p > 2$, Ostrik [21] conjectures that the answer is a characteristic- p version of quantum $\text{SU}(2)$ at level $p - 2$ called VER_p . Etingof has conjectured that the categorified algebraic closure of $\overline{\mathbb{F}}_2$ is a nonsemisimple characteristic-2 version of SUPERVECT , described by the triangular Hopf algebra $\overline{\mathbb{F}}_2[x]/(x^2)$ with $\Delta(x) = 1 \otimes x + x \otimes 1$ and R -matrix $R = 1 \otimes 1 + x \otimes x$.

We now use the algebraic closure $\text{VECT}_{\mathbb{R}} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ to categorify the notion of torsor over $\text{Spec}(\mathbb{R})$. We first show that $\text{VECT}_{\mathbb{R}} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ is ‘‘Galois’’. Let $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots)$ be a categorified commutative \mathbb{R} -algebra. A \mathcal{C} -module is an \mathbb{R} -linear locally presentable category $\mathcal{V} \in \text{PRES}_{\mathbb{R}}$ together with an action of \mathcal{C} on \mathcal{V} which is cocontinuous in each variable. A morphism of finite-dimensional \mathcal{C} -modules is a cocontinuous strong module functor. Since \mathcal{C} is commutative, the bicategory $\text{MOD}_{\mathcal{C}}$ of finite-dimensional \mathcal{C} -modules carries a symmetric monoidal structure $\boxtimes_{\mathcal{C}}$. See for example Definitions 2.1, 2.6 and 3.2 of [10].

Let $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \dots)$ be a 1-morphism of categorified commutative \mathbb{R} -algebras. Such a map makes $(\mathcal{D}, \otimes_{\mathcal{D}}, \dots)$ into a commutative algebra object in $\text{MOD}_{\mathcal{C}}$. Let $\text{Aut} = \text{Aut}_{\mathcal{C}}(\mathcal{D})$ denote the group of \mathcal{C} -linear symmetric monoidal automorphisms of \mathcal{D} . We will denote by $\text{MOD}_{\mathcal{D} \rtimes \text{Aut}}$ the bicategory of \mathcal{D} -modules equipped with a \mathcal{C} -linear Aut -action such that the \mathcal{D} -action is Aut -equivariant. It is a symmetric monoidal bicategory with symmetric monoidal structure given by the tensor product of underlying \mathcal{D} -modules. The scalar extension functor $\boxtimes_{\mathcal{C}} \mathcal{D}: \text{MOD}_{\mathcal{C}} \rightarrow \text{MOD}_{\mathcal{D}}$ factors canonically through $\text{MOD}_{\mathcal{D} \rtimes \text{Aut}}$:

$$\text{MOD}_{\mathcal{C}} \xrightarrow{\boxtimes_{\mathcal{C}} \mathcal{D}} \text{MOD}_{\mathcal{D} \rtimes \text{Aut}} \xrightarrow{\text{forget}} \text{MOD}_{\mathcal{D}}.$$

The functor $\boxtimes_{\mathcal{C}} \mathcal{D}: \text{MOD}_{\mathcal{C}} \rightarrow \text{MOD}_{\mathcal{D} \rtimes \text{Aut}}$ has a right adjoint $(-)^{\text{Aut}}: \text{MOD}_{\mathcal{D} \rtimes \text{Aut}} \rightarrow \text{MOD}_{\mathcal{C}}$ given by taking the Aut -fixed points of a module $\mathcal{V} \in \text{MOD}_{\mathcal{D} \rtimes \text{Aut}}$.

Definition 2.10 An extension of categorified fields $(\mathcal{C}, \otimes_{\mathcal{C}}, \dots) \rightarrow (\mathcal{D}, \otimes_{\mathcal{D}}, \dots)$ is *Galois* if

$$\boxtimes_{\mathcal{C}} \mathcal{D}: \text{MOD}_{\mathcal{C}} \rightleftarrows \text{MOD}_{\mathcal{D} \rtimes \text{Aut}} : (-)^{\text{Aut}}$$

is an equivalence of bicategories.

We will prove:

Theorem 2.11 *The extension $\text{VECT}_{\mathbb{R}} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ is Galois.*

Remark 2.12 For comparison, the extensions

$$\text{VECT}_{\mathbb{F}_p} \rightarrow \text{VER}_p \quad \text{and} \quad \text{VECT}_{\mathbb{F}_2} \rightarrow \text{REP}(\mathbb{F}[x]/(x^2))$$

from Remark 2.9 are not Galois (except for when $p = 3$). Indeed, the latter is “purely inseparable”, and the maximal “separable” subextension of $\text{VECT}_{\mathbb{F}_p} \rightarrow \text{VER}_p$ is $\text{SUPERVECT}_{\mathbb{F}_p}$.

We henceforth write $\text{GAL}(\mathbb{R}) = \text{Aut}_{\mathbb{R}}(\text{SUPERVECT}_{\mathbb{C}})$, and call it the *categorified absolute Galois group of \mathbb{R}* . We first calculate it:

Lemma 2.13 *The categorified absolute Galois group of \mathbb{R} is $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$.*

Proof Since categorified commutative \mathbb{R} –algebras form a bicategory, $\text{GAL}(\mathbb{R})$ is a group object in homotopy 1–types. A symmetric monoidal autoequivalence of $\text{SUPERVECT}_{\mathbb{C}}$ consists of a functor $F: \text{SUPERVECT}_{\mathbb{C}} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ and some compatible isomorphisms. We can canonically trivialize the isomorphisms $F(\mathbb{1}) \cong \mathbb{1}$ and $F(\mathbb{1} \otimes X) \cong \mathbb{1} \otimes F(X)$, and so the only remaining datum is an isomorphism $\phi: F(\mathbb{J} \otimes \mathbb{J}) \xrightarrow{\cong} F(\mathbb{J}) \otimes F(\mathbb{J})$, of which there are \mathbb{C}^{\times} –many. The functor F admits symmetric monoidal natural automorphisms that are trivial on $\mathbb{1}$ but act on \mathbb{J} by $\alpha \in \mathbb{C}^{\times}$. Under such an automorphism, the map ϕ transforms to $\phi\alpha^2$. Thus we find that

$$\text{Aut}(F) \cong \ker(\mathbb{C}^{\times} \xrightarrow{\alpha \mapsto \alpha^2} \mathbb{C}^{\times}) \cong \mathbb{Z}/2.$$

F induces an automorphism of $\mathbb{C} = \text{End}(\mathbb{1})$. If this is the identity, then F is monoidally equivalent to the identity; otherwise, F is monoidally equivalent to extension of scalars along the complex conjugation map $\mathbb{C} \rightarrow \mathbb{C}$. Thus $\pi_0(\text{GAL}(\mathbb{R})) \cong \mathbb{Z}/2$, and the computation above shows that each connected component is a $\text{B}(\mathbb{Z}/2)$. These fit together via the Galois action of $\mathbb{Z}/2$ on

$$\ker(\mathbb{C}^{\times} \xrightarrow{\alpha \mapsto \alpha^2} \mathbb{C}^{\times}),$$

and so $\text{GAL}(\mathbb{R})$ is a split extension $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$. Direct calculation verifies that it is the trivial extension; one can also show via standard techniques that there are no nontrivial split extensions of $\mathbb{Z}/2$ by $\text{B}(\mathbb{Z}/2)$. \square

The nontrivial element in $\pi_1 \text{B}(\mathbb{Z}/2)$ acts on $\text{SUPERVECT}_{\mathbb{C}}$ as the natural transformation of the identity commonly called $(-1)^f$, where f stands for “fermion number”.

Proof of Theorem 2.11 The bicategory $\text{MOD}_{\text{VECT}_{\mathbb{R}}}$ is nothing but $\text{PRES}_{\mathbb{R}}$ itself. Given $\mathcal{V} \in \text{PRES}_{\mathbb{R}}$, its image under $\boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}}$ in $\text{MOD}_{\text{SUPERVECT}_{\mathbb{C}}} \rtimes \text{Aut}$ can be described as follows. The objects of $\mathcal{V} \boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}}$ are formal direct sums $V_0 \oplus \mathbb{J} V_1$ where V_0 and V_1 are objects of \mathcal{V} . The morphisms are

$$\text{hom}(V_0 \oplus \mathbb{J} V_1, W_0 \oplus \mathbb{J} W_1) = \text{hom}_{\mathcal{V}}(V_0, W_0) \otimes \mathbb{C} \oplus \text{hom}_{\mathcal{V}}(V_1, W_1) \otimes \mathbb{C}.$$

$\text{SUPERVECT}_{\mathbb{C}}$ acts on $\mathcal{V} \boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}}$ in the obvious way. The action of $\text{Aut}_{\mathbb{R}}(\text{SUPERVECT}_{\mathbb{C}}) = \mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ is via complex conjugation and $(-1)^f$, just as it is on $\text{SUPERVECT}_{\mathbb{C}}$. The fixed-points of this action are therefore the purely even objects — those of the form $V_0 \oplus \mathbb{J} 0$ — equipped with a \mathbb{C} -antilinear involutive automorphism of V_0 . The fact that $\mathbb{R} \rightarrow \mathbb{C}$ is Galois then implies that the composition $(-)^{\text{Aut}} \circ (\boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}})$ is equivalent to the identity.

It remains to verify that $(\boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}}) \circ (-)^{\text{Aut}}$ is equivalent to the identity. Let \mathcal{V} be a $\text{SUPERVECT}_{\mathbb{C}}$ -module. Then \mathcal{V} comes equipped with an endofunctor $\mathbb{J} \otimes: \mathcal{V} \rightarrow \mathcal{V}$, given by the action of the odd line $\mathbb{J} \in \text{SUPERVECT}_{\mathbb{C}}$, satisfying $(\mathbb{J} \otimes)^2 \cong \text{id}$, and for each $X \in \text{VECT}_{\mathbb{C}}$ an endofunctor $X \otimes: \mathcal{V} \rightarrow \mathcal{V}$. The data of an Aut -action on \mathcal{V} compatible with these actions consists of: an endofunctor $V \mapsto \bar{V}$, squaring to the identity, such that $\overline{X \otimes V} \cong \bar{X} \otimes \bar{V}$ for $X \in \text{VECT}_{\mathbb{C}}$; and a natural automorphism θ of the identity functor, squaring to the identity, such that $\theta_{\mathbb{J} \otimes V} = -\text{id}_{\mathbb{J}} \otimes \theta_V$. Let’s say that $V \in \mathcal{V}$ is *purely even* if $\theta_V = +1$ and *purely odd* if $\theta_V = -1$. Then every $V \in \mathcal{V}$ canonically decomposes into a direct sum $V = V_0 \oplus V_1$ of purely even and purely odd submodules. The Aut -fixed points are the purely even submodules $V = V_0$ equipped with isomorphisms $V \cong \bar{V}$. Note that $\mathbb{J} \otimes$ interchanges purely even and purely odd objects, and so

$$\begin{aligned} \mathcal{V} &\simeq \{\text{purely even objects in } \mathcal{V}\} \boxplus \{\text{purely odd objects in } \mathcal{V}\} \\ &\simeq \{\text{purely even objects in } \mathcal{V}\} \boxtimes_{\mathbb{C}} \text{SUPERVECT}_{\mathbb{C}}. \end{aligned}$$

Finally, since $\mathbb{R} \rightarrow \mathbb{C}$ is Galois, restricting from $\{\text{purely even objects in } \mathcal{V}\}$ to those with $V \cong \bar{V}$ gives an \mathbb{R} -linear category whose tensor product with \mathbb{C} is exactly $\{\text{purely even objects in } \mathcal{V}\}$. \square

Remark 2.14 Theorem 2.11 implies that the full list of categorified field extensions of \mathbb{R} consists of the familiar categories $\text{VECT}_{\mathbb{R}}$, $\text{VECT}_{\mathbb{C}}$, $\text{SUPERVECT}_{\mathbb{R}}$ and $\text{SUPERVECT}_{\mathbb{C}}$, and a less-familiar category that deserves to be called $\text{SUPERVECT}_{\mathbb{H}}$. The first four are the fixed-points for the obvious subgroups $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$, $\text{B}(\mathbb{Z}/2)$, $\mathbb{Z}/2$ and $\{1\}$ of the categorified Galois group $\text{GAL}(\mathbb{R})$. The last is the fixed-points for the nonobvious inclusion $\mathbb{Z}/2 \hookrightarrow \mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ which is the identity on the first component and the nontrivial map $\mathbb{Z}/2 \rightarrow \text{B}(\mathbb{Z}/2)$ on the second component (corresponding to the nontrivial class in $\text{H}^2(\text{B}(\mathbb{Z}/2); \mathbb{Z}/2)$). As a category, $\text{SUPERVECT}_{\mathbb{H}} \simeq \text{VECT}_{\mathbb{R}} \boxplus \text{MOD}_{\mathbb{H}}$, hence the name. The monoidal structure involves the Morita equivalence $\mathbb{H} \otimes_{\mathbb{R}} \mathbb{H} \simeq \mathbb{R}$.

We can now categorify the usual classification of torsors in terms of Galois actions.

Definition 2.15 Let G be a finite Picard groupoid. A categorified G -torsor over \mathbb{R} is a nonzero G -equivariant categorified commutative \mathbb{R} -algebra \mathcal{T} such that the functor

$$\mathcal{T} \boxtimes_{\mathbb{R}} \mathcal{T} \rightarrow \text{maps}(G, \mathcal{T}), \quad t_1 \boxtimes t_2 \mapsto (g \triangleright t_1) \otimes t_2$$

is an equivalence, where $\text{maps}(G, \mathcal{T})$ denotes the categorified commutative algebra of \mathcal{T} -valued functors on the underlying groupoid of G , \triangleright denotes the action of G on \mathcal{T} , and \otimes denotes the multiplication in \mathcal{T} .

Proposition 2.16 Let $\text{GAL}(\mathbb{R}) = \text{Aut}_{\mathbb{R}}(\text{SUPERVECT}_{\mathbb{C}})$ denote the categorified absolute Galois group of \mathbb{R} . For each finite categorified group G , there is a natural-in- G equivalence

$$\{\text{categorified } G\text{-torsors over } \mathbb{R}\} \simeq \text{maps}(\text{B GAL}(\mathbb{R}), \text{BG}).$$

The proof is just as in the uncategorified situation:

Proof Let \mathcal{T} be a categorified G -torsor over \mathbb{R} . Then $\mathcal{T}' = \mathcal{T} \boxtimes_{\mathbb{R}} \text{SUPERVECT}_{\mathbb{C}}$ is a G -torsor over $\text{SUPERVECT}_{\mathbb{C}}$. Since $\text{SUPERVECT}_{\mathbb{C}}$ is algebraically closed, we can choose a symmetric monoidal functor $F: \mathcal{T}' \rightarrow \text{SUPERVECT}_{\mathbb{C}}$. Let \boxtimes' denote $\boxtimes_{\text{SUPERVECT}_{\mathbb{C}}}$. The equivalence $\mathcal{T}' \boxtimes' \mathcal{T}' \mapsto \text{maps}(G, \mathcal{T}')$ making \mathcal{T}' into a torsor over $\text{SUPERVECT}_{\mathbb{C}}$ fits into a commutative square

$$\begin{array}{ccc} \mathcal{T}' \boxtimes' \mathcal{T}' & \xrightarrow{\sim} & \text{maps}(G, \mathcal{T}') \\ \downarrow \text{id} \boxtimes' F & & \downarrow F \circ \\ \mathcal{T}' & \xrightarrow{t \mapsto (g \mapsto F(g \triangleright t))} & \text{maps}(G, \text{SUPERVECT}_{\mathbb{C}}) \end{array}$$

in which the downward arrows are both equivalent to $\boxtimes_{\mathcal{T}'}$ SUPERVECT $_{\mathbb{C}}$. It follows that \mathcal{T}' is a trivial G -torsor over SUPERVECT $_{\mathbb{C}}$.

Theorem 2.11 then provides an equivalence of homotopy 2-types

$$\begin{aligned} & \{\text{categorified } G\text{-torsors over } \mathbb{R}\} \\ & \simeq \{\text{GAL}(\mathbb{R})\text{-actions on the trivial } G\text{-torsor over SUPERVECT}_{\mathbb{C}} \\ & \quad \text{compatible with the action on SUPERVECT}_{\mathbb{C}}\}. \end{aligned}$$

But $\text{Aut}_{\mathbb{R}}(\text{maps}(G, \text{SUPERVECT}_{\mathbb{C}})) \simeq G \times \text{GAL}(\mathbb{R})$, and the equivariance requirement is equivalent to the requirement that the morphism $\text{GAL}(\mathbb{R}) \rightarrow G \times \text{GAL}(\mathbb{R})$ is the identity on the second component. Therefore we are left with maps $\text{GAL}(\mathbb{R}) \rightarrow G$ up to equivalences given by inner automorphism. \square

3 Spin and spin-statistics field theories

With the categorified Galois extension $\text{VECT}_{\mathbb{R}} \rightarrow \text{SUPERVECT}_{\mathbb{C}}$ from [Section 2](#) in hand, we are equipped to categorify the story from [Section 1](#). The uncategorified story related orientations with hermiticity; the categorified story will relate spin and statistics.

Recall that a *spin structure* on a d -dimensional manifold M is a $\text{Spin}(d)$ -principal bundle $P \rightarrow M$ together with an isomorphism $P \times_{\text{Spin}(d)} \mathbb{R}^d \cong TM$. The collection $\text{Spins}(M)$ of spin structures on M is not naturally a set, but rather a groupoid. We therefore extend without further comment the notion of *topological local structure* valued in a bicategory \mathcal{X} to be a sheaf $\text{MAN}_d \rightarrow \mathcal{X}$ that takes homotopies between maps in MAN_d to isomorphisms between maps in \mathcal{X} and homotopies between homotopies to equalities between isomorphisms. Generalizing [Lemma 1.1](#), we have:

Lemma 3.1 *Let \mathcal{X} be a bicategory with limits. Topological local structures on MAN_d valued in \mathcal{X} are equivalent to objects of \mathcal{X} equipped with an action by the Picard groupoid $\pi_{\leq 1} \text{hom}_{\text{MAN}_d}(\mathbb{R}^d, \mathbb{R}^d) = \pi_{\leq 1} \text{O}(d)$. When $d \geq 3$, this Picard groupoid is canonically equivalent to $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$.* \square

Remark 3.2 The existence of an identification $\pi_{\leq 1} \text{O}(d) \cong \mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$, $d \geq 3$, is the same as the standard assertion that the k -invariant connecting $\pi_1 \text{BO}(\infty)$ and $\pi_2 \text{BO}(\infty)$ vanishes. However, the group $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ admits a nontrivial group automorphism, given by the identity on each factor and the nontrivial group map $\mathbb{Z}/2 \rightarrow \text{B}(\mathbb{Z}/2)$, corresponding to the nontrivial element of $\text{H}^2(\text{B}(\mathbb{Z}/2); \mathbb{Z}/2) = \mathbb{Z}/2$, mixing the factors. Thus there are two inequivalent identifications $\pi_{\leq 1} \text{O}(\infty) \cong \mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$. To pick one is the same as to pick a splitting of the projection

$\pi_{\leq 1}O(d) \rightarrow \pi_0O(d) = \mathbb{Z}/2$. There is a canonical choice: the “stable” splitting $\mathbb{Z}/2 \rightarrow O(d)$ sending the nontrivial element of $\mathbb{Z}/2$ to the matrix

$$\begin{pmatrix} -1 & & & \\ & +1 & & \\ & & +1 & \\ & & & \ddots \end{pmatrix},$$

called “T” in the physics literature. Corresponding to the two splittings $\mathbb{Z}/2 \rightarrow \pi_{\leq 1}O(d)$ are two projections $\pi_{\leq 1}O(d) \rightarrow B(\mathbb{Z}/2)$, the kernels of which are the two *pin* groups $\text{Pin}^{\pm}(d)$.

Example 3.3 We recall two standard facts about spin structures. First, given any spin structure on a d -dimensional manifold M , let $P \rightarrow M$ denote the corresponding $\text{Spin}(d)$ -bundle. Then $P \times_{\text{Spin}(d)} \text{Pin}^+(d)$ is a $\text{Pin}^+(d)$ -bundle over M with a distinguished sheet. The other sheet of $P \times_{\text{Spin}(d)} \text{Pin}^+(d)$ also defines a $\text{Spin}(d)$ -bundle over M , corresponding to the *orientation-reversal* of the original spin structure. Second, any spin structure on M admits a square-1 automorphism which acts on the bundle $P \rightarrow M$ by multiplication by the nontrivial central element of $\text{Spin}(d)$ coming from 360° -rotation in $\text{SO}(d)$. The mapping cylinder of this automorphism is the product spin manifold $M \times \mathfrak{A}$, where \mathfrak{A} denotes the nontrivial-rel-boundary spin structure on the interval $[0, 1]$. We will also use the name “ \mathfrak{A} ” to denote the automorphism of the spin structure. Together, orientation-reversal and \mathfrak{A} define an action of $\mathbb{Z}/2 \times B(\mathbb{Z}/2)$ on $\text{Spins}(M)$.

When $M = \mathbb{R}^d$, the orientation reversal and 360° -rotation action of $\mathbb{Z}/2 \times B(\mathbb{Z}/2)$ witness $\text{Spins}(\mathbb{R}^d)$ as the trivial $(\mathbb{Z}/2 \times B(\mathbb{Z}/2))$ -torsor. When $d \geq 3$, orientation-reversal and 360° -rotation make up the full group $\pi_{\leq 1}O(d) = \mathbb{Z}/2 \times B(\mathbb{Z}/2)$, and so Spins is the topological local structure corresponding to the trivial $\pi_{\leq 1}O(d)$ -torsor via [Lemma 3.1](#). When $d < 3$, the canonical inclusion $X \mapsto \begin{pmatrix} X & \\ & 1 \end{pmatrix}$ of $O(d)$ into $O(3)$ provides an action of $\pi_{\leq 1}O(d)$ on $\pi_{\leq 1}O(3) \cong \mathbb{Z}/2 \times B(\mathbb{Z}/2)$, which in turn corresponds to the topological local structure Spins .

We now move to an algebrogeometric setting in which there are interesting topological local structures that are étale-locally equivalent to Spins in the way that Her was étale-locally equivalent to Or. Ordinary algebraic geometry does not suffice, since $\text{Spec}(\mathbb{C})$ is étale-contractible in the ordinary sense. Instead, since groupoids are a categorification of sets, we work with a categorification of schemes:

Definition 3.4 The bicategory $\text{CATAFFSCH}_{\mathbb{R}}$ of *categorified affine schemes over \mathbb{R}* is opposite to the bicategory of categorified commutative \mathbb{R} -algebras in the sense of

Definition 2.1. We will write $\text{Spec}(\mathcal{C})$ for the categorified affine scheme corresponding to a categorified commutative algebra \mathcal{C} .

Lemma 2.2 provides a fully faithful inclusion of the category $\text{AFFSCH}_{\mathbb{R}}$ of uncategorified affine schemes into $\text{CATAFFSCH}_{\mathbb{R}}$; in particular, we identify $\text{Spec}(\mathbb{R})$ with $\text{Spec}(\text{VECT}_{\mathbb{R}})$. The details of notions like “nonaffine categorified scheme” and “categorified étale topology” have yet to be worked out, and are the subject of joint work in progress by A Chirvasitu, E Elmanto and the author. **Theorem 2.11** suggests that $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}}) \rightarrow \text{Spec}(\mathbb{R})$ is a “categorified étale cover” and **Theorem 2.7** suggests that $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ is “categorified étale contractible”. In particular, we will say that categorified affine schemes X and Y are *étale-locally equivalent* if their pullbacks $X \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ and $Y \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ are equivalent as categorified affine schemes over $\text{SUPERVECT}_{\mathbb{C}}$. This in particular implies that for any Picard groupoid G , the geometric notion of categorified G -torsors over $\text{Spec}(\mathbb{R})$, defined as G -objects over $\text{Spec}(\mathbb{R})$ étale-locally equivalent to G acting on itself, agrees with the algebraic notion from **Definition 2.15**, which by **Lemma 2.13** and **Proposition 2.16** are classified by maps $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2) \rightarrow G$.

Now note the following coincidence: there is a canonical equivalence $\pi_{\leq 1}\text{O}(d) \cong \mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2) \cong \text{GAL}(\mathbb{R})$, and hence a canonical categorified $\pi_{\leq 1}\text{O}(d)$ -torsor, when $d \geq 3$. This torsor is nothing but the categorified affine scheme $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ equipped with its $\text{GAL}(\mathbb{R})$ -action.

Definition 3.5 The sheaf of *hermitian spin-statistics structures* is the sheaf

$$\text{HerSpinStats}: \text{MAN}_d \rightarrow \text{CATAFFSCH}_{\mathbb{R}}$$

such that $\text{HerSpinStats}(\mathbb{R}^d) = \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$, on which $\pi_{\leq 1}\text{O}(d) \cong \text{GAL}(\mathbb{R})$ acts via the Galois action.

Lemma 3.6 For any manifold M ,

$$\text{HerSpinStats}(M) = \frac{\text{Spins}(M) \times \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})}{\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)},$$

where $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ acts on $\text{Spins}(M)$ by orientation-reversal and by \mathfrak{A} from **Example 3.3**, and it acts on $\text{SUPERVECT}_{\mathbb{C}}$ by complex conjugation and by $(-1)^f$ from **Lemma 2.13**. \square

Lemma 3.6 begins to justify the phrase “hermitian spin-statistics structure” from **Definition 3.5**: that orientation-reversal acts by complex conjugation is the essence of

hermiticity, and that \mathfrak{A} acts by $(-1)^f$ is a version of spin-statistics as it is used in physics.

To further justify the name, we should study hermitian spin-statistics field theories directly. The definition of hermitian spin-statistics field theory will be a direct analog of hermitian field theory from [Section 1](#).


Let $\text{BORD}_{d-2,d-1,d}$ denote the once-extended d -dimensional bordism bicategory constructed by Schommer-Pries in [\[22\]](#). Given any topological local structure valued in groupoids $\mathcal{G}: \text{MAN}_d \rightarrow \text{GPOIDS}$, [\[22\]](#) also explains how to build a symmetric monoidal bicategory $\text{BORD}_{d-2,d-1,d}^{\mathcal{G}}$ of bordisms with \mathcal{G} -structure. A *once-extended \mathcal{G} -structured field theory* is then a symmetric monoidal functor $Z: \text{BORD}_{d-2,d-1,d}^{\mathcal{G}} \rightarrow \mathcal{V}$ for some symmetric monoidal bicategory \mathcal{V} of “categorified vector spaces”.

We will take $\mathcal{V} = \text{ALG}_{\mathbb{R}}$ to be the symmetric monoidal “Morita” bicategory of associative algebras, bimodules and intertwiners. Just as $\text{VECT}_{\mathbb{R}}$ had a natural extension to the stack QCOH of categories over $\text{SCH}_{\mathbb{R}}$, so too $\text{ALG}_{\mathbb{R}}$ has a natural extension allowing for “bundles” or “sheaves” of algebras over any categorified affine scheme: given a categorified commutative \mathbb{R} -algebra \mathcal{C} , set $\text{ALG}(\text{Spec}(\mathcal{C})) = \text{ALG}(\mathcal{C})$ to be the symmetric monoidal bicategory of algebra objects in \mathcal{C} , bimodule objects in \mathcal{C} , and intertwiners in \mathcal{C} . Although we have not defined, and will not use, any topology on $\text{CATAFFSCH}_{\mathbb{R}}$, and so cannot say precisely what it means to be a stack of bicategories, it is not hard to find a bicategory object internal to $\text{CATAFFSCH}_{\mathbb{R}}$ that represents $\text{ALG}(-)$, and so $\text{ALG}(-)$ is certainly a stack of bicategories in any subcanonical topology.

Remark 3.7 The Eilenberg–Watts theorem [\[11; 26\]](#) identifies $\text{ALG}_{\mathbb{R}}$ with the full subcategory of $\text{PRES}_{\mathbb{R}}$ whose objects admit a compact projective generator. The correct target for once-extended *nontopological* quantum field theory is more likely the larger $\text{PRES}_{\mathbb{R}}$. But it is reasonable to expect that every *topological* field theory factors through $\text{ALG}_{\mathbb{R}}$, since it is expected that only categories equivalent to MOD_A , $A \in \text{ALG}_{\mathbb{R}}$, are sufficiently dualizable (see [\[6\]](#)). Indeed, one should expect more: topological field theories should factor through the subcategory of $\text{ALG}_{\mathbb{R}}$ whose objects are finite-dimensional algebras and whose morphisms are finite-dimensional bimodules. This subcategory is equivalent to the bicategory $\text{MOD}_{\text{PRES}_{\mathbb{R}}}$ of finite-dimensional $\text{PRES}_{\mathbb{R}}$ -modules from [Section 2](#). More generally, for \mathcal{C} a finite-dimensional categorified commutative ring, the bicategory $\text{MOD}_{\mathcal{C}}$ of finite-dimensional \mathcal{C} -modules is a subcategory of $\text{ALG}(\mathcal{C})$, which is a subcategory of the bicategory of all \mathcal{C} -modules.

Definition 3.8 Let $\text{SPANS}_2(\text{CATAFFSCH}_{\mathbb{R}})$ denote the symmetric monoidal bicategory whose objects are categorified affine schemes, 1-morphisms are spans $X \leftarrow A \rightarrow Y$,

and 2–morphisms are spans-between-spans:



Composition is by fibered product, and the symmetric monoidal structure is the cartesian product in $\text{CATAFFSCH}_{\mathbb{R}}$. Let \mathcal{G} be a topological local structure valued in $\text{CATAFFSCH}_{\mathbb{R}}$; it defines a symmetric monoidal functor

$$\tilde{\mathcal{G}}: \text{BORD}_{d-2,d-1,d} \rightarrow \text{SPANS}_2(\text{CATAFFSCH}_{\mathbb{R}}).$$

Let $\text{SPANS}_2(\text{CATAFFSCH}_{\mathbb{R}}; \text{ALG})$ be the symmetric monoidal bicategory whose objects are a categorified affine scheme X together with an algebra $V \in \text{ALG}(X)$, whose 1–morphisms are spans $X \xleftarrow{f} A \xrightarrow{g} Y$ together with a bimodule between f^*V and g^*W in $\text{ALG}(A)$, and whose 2–morphisms are spans of spans together with an intertwiner between pulled-back bimodules. A \mathcal{G} –structured field theory is a lift:

$$\begin{array}{ccc} & \text{SPANS}_2(\text{CATAFFSCH}_{\mathbb{R}}; \text{ALG}) & \\ & \nearrow \tilde{\mathcal{G}} & \downarrow \text{forget the ALG-data} \\ \text{BORD}_{d-2,d-1,d} & \longrightarrow & \text{SPANS}_2(\text{CATAFFSCH}_{\mathbb{R}}) \end{array}$$

Example 3.9 We now continue to justify the name “hermitian spin-statistics” from Lemma 3.6. Let Z be a d –dimensional HerSpinStats–structured field theory. We will unpack its values on various manifolds.

Suppose first that M is a closed d –dimensional manifold. Considered as an element of $\text{BORD}_{d-2,d-1,d}$, M is an endo-2–morphism of the identity 1–morphism of the unit object. Then $Z(M)$ is an endo-2–morphism of the identity 1–morphism of the unit object in $\text{ALG}(\text{HerSpinStats}(M))$, ie a function $Z(M) \in \mathcal{O}(\text{HerSpinStats}(M))$. Any choice of spin structure for M determines a map $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}}) \rightarrow \text{HerSpinStats}(M)$, and these maps together cover $\text{HerSpinStats}(M)$ as the spin structure varies over M . Thus the data of $Z(M)$ is the data of an element of $\mathcal{O}(\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})) = \mathbb{C}$ for each spin structure on M . By the construction of HerSpinStats from Lemma 3.6, two spin structures on M with reversed orientation lead to complex-conjugate values of $Z(M)$. This is a manifestation of the hermiticity of Z .

To see spin-statistics phenomena, consider next the case of a closed $(d-1)$ –dimensional manifold N . Then $Z(N)$ is an endo-1–morphism of the unit object in the category

$\text{ALG}(\text{HerSpinStats}(N))$, ie a vector bundle on $\text{HerSpinStats}(N)$. Again, any spin structure on N allows this vector bundle to be pulled back to a vector bundle on $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$, and so $Z(N)$ assigns a complex supervector space to each spin structure on N . In addition to the hermiticity requirement that orientation-reversed spin structures map to complex-conjugate supervector spaces, there is another relation between these supervector spaces and the spin structures. Indeed, fix a spin structure σ on N , and let $Z(N, \sigma)$ denote the corresponding complex supervector space. Consider the spin cobordism $(N, \sigma) \times \mathfrak{A}$. This spin structure picks out a particular map $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}}) \rightarrow \text{HerSpinStats}(N \times [0, 1])$, along which $Z(N \times [0, 1])$ pulls back to a map $Z((N, \sigma) \times \mathfrak{A}): Z(N, \sigma) \rightarrow Z(N, \sigma)$. But $(N, \sigma) \times \mathfrak{A}$ is simply the mapping cylinder of the 360° -rotation of σ , and [Lemma 3.6](#) identifies 360° -rotation with $(-1)^f$. All together, we find that $Z((N, \sigma) \times \mathfrak{A})$ is required to evaluate to $(-1)^f: Z(N, \sigma) \rightarrow Z(N, \sigma)$.

Similar discussion applies also in codimension-2, and hermitian spin-statistics field theories unpack to spin field theories $\text{BORD}_{d-2, d-1, d}^{\text{Spins}} \rightarrow \text{ALG}(\text{SUPERVECT}_{\mathbb{C}})$ such that the actions of $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ on the source and target categories are intertwined. The phrase “spin-statistics” refers to the identification $\mathfrak{A} = (-1)^f$. In a spin field theory the (-1) -eigenstates of \mathfrak{A} are called *spinors* and in a super field theory the (-1) -eigenstates of $(-1)^f$ are called *fermions*, so “spin-statistics” can be equivalently described as the assertion that the classes of spinors and fermions agree.

By construction, HerSpinStats is an *étale-locally-spin* topological local structure in the sense that

$\text{HerSpinStats} \times_{\text{Spec}(\mathbb{R})} \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ and $\text{Spins} \times \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ are equivalent. Since $\text{GAL}(\mathbb{R}) = \mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ and Spins corresponds to the trivial $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ -torsor, [Proposition 2.16](#) asserts that the set of inequivalent topological local structures étale-locally-equivalent to Spins is equivalent to

$$\pi_0 \text{ maps } (\text{B}(\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)), \text{B}(\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2))),$$

which can be easily computed as

$$\begin{aligned} \text{H}^1(\text{B}(\mathbb{Z}/2); \mathbb{Z}/2) \times \text{H}^2(\text{B}(\mathbb{Z}/2); \mathbb{Z}/2) \times \text{H}^1(\text{B}^2(\mathbb{Z}/2); \mathbb{Z}/2) \times \text{H}^2(\text{B}^2(\mathbb{Z}/2); \mathbb{Z}/2) \\ \cong (\mathbb{Z}/2)^3, \end{aligned}$$

and so there are exactly eight different choices. Whether the corresponding field theories are oriented or hermitian is controlled by the component $\mathbb{Z}/2 \rightarrow \mathbb{Z}/2$ relating complex conjugation with orientation-reversal. Whether the field theories are spin or spin-statistics is controlled by the component $\text{B}(\mathbb{Z}/2) \rightarrow \text{B}(\mathbb{Z}/2)$ relating $(-1)^f$ with \mathfrak{A} .

But once these choices are made, there is still the choice of map $\mathbb{Z}/2 \rightarrow \mathbf{B}(\mathbb{Z}/2)$ — the possible choices are parametrized by $H^2(\mathbb{Z}/2; \mathbb{Z}/2) \cong \mathbb{Z}/2$ — which adjusts how orientation-reversal behaves on fermions.

There are also topological local structures \mathcal{G} satisfying $\mathcal{G}(\mathbb{R}^d) = \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$ but in which part or all of $\pi_{\leq 1} \mathbf{O}(d)$ acts trivially, analogous to the \mathbb{C} -linear unstructured field theories from [Example 1.8](#). We now illustrate a few of the possible choices to emphasize that spin and statistics are not intrinsically linked, even in the presence of hermiticity. We will then prove [Theorem 0.1](#) showing that spin and statistics are linked when an extra reflection-positivity hypothesis is imposed. In order to construct examples of field theories with various topological local structures, we focus on the case when $d = 2$, since then we can use Schommer-Pries's classification of 2-dimensional field theories from [\[22\]](#).

Example 3.10 A *hermitian spin field theory* is an \mathbb{R} -linear field theory with local structure $\text{Spins} \times_{\mathbb{Z}/2} \text{Spec}(\mathbb{C})$. Unpacking the definition, a hermitian spin field theory is a nonsuper \mathbb{C} -linear spin field theory such that orientation-reversal agrees with complex conjugation. In terms of simultaneously-spin-and-super field theories, \mathfrak{R} acts nontrivially but $(-1)^f$ acts trivially.

Two-dimensional \mathbb{C} -linear spin field theories in ALG are classified by finite-dimensional complex semisimple algebras A equipped with a trivialization $\varphi: A^* \otimes_A A^* \xrightarrow{\sim} A$ of A - A bimodules, where A^* denotes the linear dual bimodule to A , such that the two maps $\varphi \otimes \text{id}: A^* \otimes_A A^* \otimes_A A^* \xrightarrow{\sim} A \otimes_A A^* = A^*$ and $\text{id} \otimes \varphi: A^* \otimes_A A^* \otimes_A A^* \xrightarrow{\sim} A^* \otimes_A A = A^*$ agree. The hermiticity requirement unpacks to having a (\mathbb{C} -antilinear) stellar structure, ie a Morita equivalence $A^{\text{op}} \cong \bar{A}$, where \bar{A} is the complex-conjugate algebra, satisfying certain requirements [\[22, Section 3.8.6\]](#). Stellar structures are the Morita-equivariant version of $*$ -structures, and any $*$ -structure defines a stellar structure. Hermiticity requires that φ be real.

For example, we can take $A = \mathbb{C}$ with its standard $*$ -algebra structure, and choose the trivialization $\varphi: \mathbb{C} = \mathbb{C}^* \otimes_{\mathbb{C}} \mathbb{C}^* \xrightarrow{\sim} \mathbb{C}$ to be multiplication by -1 . Either trivialization $\pm\sqrt{-1}: \mathbb{C}^* \rightarrow \mathbb{C}$ presents the \mathbb{C} -linear field theory defined by A as the underlying spin field theory of an oriented field theory over \mathbb{C} . But as a hermitian spin theory, the field theory defined by A is fundamentally spin, since neither $\pm\sqrt{-1}$ is real.

Example 3.11 A *hermitian super field theory* is an \mathbb{R} -linear field theory with local structure $\text{Or} \times_{\mathbb{Z}/2} \text{Spec}(\text{SUPERVECT}_{\mathbb{C}}) \cong \text{Her} \times \text{Spec}(\text{SUPERVECT}_{\mathbb{R}})$, ie an oriented field theory valued in $\text{SUPERVECT}_{\mathbb{C}}$ such that orientation-reversal agrees with complex conjugation. In terms of simultaneously-spin-and-super field theories, $(-1)^f$ acts nontrivially but \mathfrak{R} acts trivially.

Two-dimensional hermitian super field theories are classified by symmetric Frobenius stellar superalgebras. In particular, every symmetric Frobenius \ast -superalgebra determines a hermitian super field theory. Consider the complex superalgebra $\mathbb{C}\text{Cliff}(2) = \mathbb{C}\langle x, y \rangle / (x^2 = y^2 = 1, [x, y] = 0)$, where x and y are odd. It admits a \ast -structure in which $x^\ast = x\sqrt{-1}$ and $y^\ast = y\sqrt{-1}$. Then xy is imaginary and even, and $\mathbb{C}\text{Cliff}(2)$ admits a symmetric Frobenius \ast -superalgebra structure in which $\text{tr}(xy) = \sqrt{-1}$ and $\text{tr}(1) = \text{tr}(x) = \text{tr}(y) = 0$.

As a complex Frobenius superalgebra, $\mathbb{C}\text{Cliff}(2)$ is Morita-equivalent to \mathbb{C} , and so the \mathbb{C} -linear oriented super field theory defined by $\mathbb{C}\text{Cliff}(2)$ is the superification of a purely bosonic theory. But the Morita equivalence $\mathbb{C}\text{Cliff}(2) \simeq \mathbb{C}$ is not compatible with the stellar structure, and so the corresponding hermitian super field theory defined by $\mathbb{C}\text{Cliff}(2)$ is fundamentally super.

Example 3.12 Two-dimensional spin-statistics field theories are classified by finite-dimensional semisimple “twisted-symmetric” Frobenius superalgebras. Specifically, let A be a finite-dimensional semisimple superalgebra arising as $Z(\{\text{pt}\})$ for some 2-dimensional field theory. Then 360° rotation acts by the dual bimodule $Z(\mathfrak{R}) = {}_A A_A^\ast$. Let ${}_A(-1)_A^f$ denote the bimodule A with actions $a \triangleright m \triangleleft b = am(-1)^{|b|}b$; it is the bimodule corresponding to the algebra automorphism $(-1)^f : A \rightarrow A$. The spin-statistics data “ $\mathfrak{R} = (-1)^f$ ” then corresponds to a bimodule isomorphism $\phi : {}_A A_A^\ast \xrightarrow{\sim} {}_A(-1)_A^f$.

Consider the trace $\text{tr}(a) = \langle \phi^{-1}(1_A), a \rangle$, where $\langle \cdot, \cdot \rangle : A^\ast \otimes A \rightarrow \mathbb{C}$ denotes the canonical pairing. This trace is not symmetric. In a symmetric Frobenius superalgebra, the trace should satisfy $\text{tr}(ab) = (-1)^{|a||b|} \text{tr}(ba)$. Instead, the trace pairing above satisfies $\text{tr}(ab) = (-1)^{|a| \cdot (|b|+1)} \text{tr}(ba) = \text{tr}(ba)$, where the second equality follows from the fact that tr , being an even map, vanishes on odd elements. Thus not the superalgebra A but rather the underlying nonsuper algebra $\text{Forget}(A)$ is symmetric Frobenius.

Real spin-statistics field theories are classified by twisted-symmetric Frobenius superalgebras in $\text{SUPERVECT}_{\mathbb{R}}$. Hermitian spin-statistics field theories are classified by twisted-symmetric Frobenius stellar superalgebras in $\text{SUPERVECT}_{\mathbb{C}}$, where the isomorphism ϕ is real.

The Clifford algebras $\mathbb{C}\text{Cliff}(n) = \mathbb{C}\langle x_1, \dots, x_n \rangle / ([x_j, x_k] = 2\delta_{jk})$ admit twisted-symmetric Frobenius \ast -superalgebra structures. As in [Example 3.11](#), we can give $\mathbb{C}\text{Cliff}(n)$ a \ast -structure by declaring $x_j^\ast = x_j\sqrt{-1}$. When n is odd, there is an isomorphism of superalgebras $\mathbb{C}\text{Cliff}(n) \cong \mathbb{C}\text{Cliff}(1) \otimes \text{Mat}_{\mathbb{C}}(2^{(n-1)/2})$, where $\text{Mat}_{\mathbb{C}}(m)$ is the purely-even algebra of $m \times m$ complex matrices, and so we can define the trace $\text{tr} : \mathbb{C}\text{Cliff}(n) \rightarrow \mathbb{C}$ to be the matrix trace on the even part (and to vanish on the odd part). This tr is twisted-symmetric and real and so defines a 2-dimensional spin-statistics hermitian field theory.

When n is even, the isomorphism $\text{Forget}(\text{Cliff}(n)) \cong \text{Mat}_{\mathbb{C}}(2^{n/2})$ defines a twisted-symmetric Frobenius structure on $\text{Cliff}(n)$. When n is even, $\text{Cliff}(n)$ also admits a nontwisted symmetric Frobenius structure; [Example 3.11](#) describes the case $n = 2$.

Example 3.13 A *twisted-hermitian spin field theory* is like a hermitian spin field theory except that rather than the canonical action of $\mathbb{Z}/2$ on Spins , we twist the action by the nontrivial map $\mathbb{Z}/2 \rightarrow \text{B}(\mathbb{Z}/2)$. This unpacks to the requirement that the trivialization φ in [Example 3.10](#) be purely imaginary.

A *twisted-hermitian super field theory* is like a hermitian super field theory except that rather than the canonical action of $\mathbb{Z}/2$ on $\text{SUPERVECT}_{\mathbb{C}}$, we twist the action by the nontrivial map $\mathbb{Z}/2 \rightarrow \text{B}(\mathbb{Z}/2)$. These are classified not by symmetric Frobenius stellar superalgebras, but by symmetric Frobenius *twisted-stellar* superalgebras. These are defined analogously to stellar superalgebras but with one modification. For any superalgebra A , consider the superalgebra A' defined by $x \cdot' y = (-1)^{|x| \cdot |y|} xy$. A stellar structure on A includes a Morita equivalence between the opposite superalgebra A^{op} and the complex conjugate superalgebra \bar{A} . A twisted-stellar structure instead makes A^{op} equivalent to \bar{A}' . A special case is that of twisted- $*$ -superalgebras. In a $*$ -superalgebra, $x \mapsto x^*$ must be an algebra antiautomorphism, which in $\text{SUPERVECT}_{\mathbb{C}}$ means that $(xy)^* = (-1)^{|x| \cdot |y|} y^* x^*$. In a twisted- $*$ -superalgebra, we have instead $(xy)^* = y^* x^*$ for elements of arbitrary parity. Examples of twisted- $*$ -superalgebras include $\text{Cliff}(n)$ for arbitrary n with $x_i^* = x_i$.

The nontrivial automorphism of $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ mentioned in [Remark 3.2](#) defines a second $\mathbb{Z}/2 \times \text{B}(\mathbb{Z}/2)$ -torsor over $\text{Spec}(\mathbb{R})$ with total space $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$. The corresponding topological local structure controls *twisted-hermitian spin-statistics field theories*. The twisted- $*$ -superalgebras $\text{Cliff}(n)$ with their twisted-symmetric Frobenius structures from [Example 3.12](#) provide examples of twisted hermitian spin-statistics field theories.

Twisted-real spin-statistics field theories are classified by twisted-symmetric Frobenius algebra objects in the category $\text{SUPERVECT}_{\mathbb{H}}$ from [Remark 2.14](#).

We now extend the notion of reflection-positivity from [Definitions 1.9](#) and [1.10](#) to the étale-locally-spin case. Following the physics literature, and in disagreement with Freed and Hopkins [12], we declare that reflection-positivity of an extended field theory can be detected in codimension one:

Definition 3.14 An extended unstructured field theory $Z: \text{BORD}_{d-2, d-1, d} \rightarrow \text{ALG}_{\mathbb{R}}$ is *reflection-positive* if its restriction $Z|_{\text{BORD}_{d-1, d}}: \text{BORD}_{d-1, d} \rightarrow \text{VECT}_{\mathbb{R}}$ to an unextended field theory is reflection-positive in the sense of [Definition 1.9](#), ie if the

symmetric pairing $Z(N \times \mathbb{S}) : Z(N)^{\otimes 2} \rightarrow \mathbb{R}$ is positive-definite for every closed $(d-1)$ -dimensional manifold N .

In [Definition 1.10](#) we defined reflection-positivity for étale-locally-oriented field theories in terms of integration over the space of étale-local orientations. We now extend that logic to étale-locally-spin field theories. Consider first the case when Z is spin. For any manifold M , $\text{Spins}(M)$ is a finite groupoid, and so Baez and Dolan [4] define an integration map

$$\int_{\text{Spins}(M)} : \mathcal{O}(\text{Spins}(M)) \rightarrow \mathbb{R}, \quad f \mapsto \sum_{x \in \pi_0 \text{Spins}(M)} f(x) / |\pi_1(\text{Spins}(M), x)|.$$

When V is a bundle over $\text{Spins}(M)$, $\int_{\text{Spins}(M)} V$ is the space of coinvariants of V .

Example 3.15 Given a two-dimensional nonhermitian spin field theory Z corresponding to the algebra $Z(\{\text{pt}\}) = A$ and trivialization $\varphi : A^* \otimes_A A^* \xrightarrow{\sim} A$, one can compute the unstructured field theory $\int_{\text{Spins}} Z$ in two steps. First, one can integrate over the fibers of the projection $\text{Spins} \rightarrow \text{Or}$. The corresponding oriented field theory $\int_{\text{Spins}/\text{Or}} Z$ is controlled by the symmetric Frobenius algebra $B = A \oplus A^*$ with multiplication $(a \oplus \alpha) \cdot (b \oplus \beta) = (ab + \varphi(\alpha \otimes \beta)) \oplus (a\beta + \alpha b)$ and Frobenius structure $\text{tr}(a \oplus \alpha) = \alpha(1)$. Second, one can integrate over the choice of orientation, producing the unstructured field theory controlled by $B \oplus B^{\text{op}}$ with the obvious algebraic $*$ -structure.

The construction \int_{Spins} makes sense for any étale-locally-spin field theory: if Z has local structure \mathcal{G} where $\mathcal{G}(\{\text{pt}\})$ is a categorified $\mathbb{Z}/2 \times \mathbb{B}(\mathbb{Z}/2)$ -torsor, then the base change $Z_{\mathcal{G}}$ of Z along $\mathcal{G}(\{\text{pt}\}) \rightarrow \text{Spec}(\mathbb{R})$ is a Galois-equivariant spin field theory over $\mathcal{G}(\{\text{pt}\})$; thus $\int_{\text{Spins}} Z_{\mathcal{G}}$ is a Galois-equivariant unstructured field theory and so descends to $\text{Spec}(\mathbb{R})$.

Example 3.16 Suppose that Z is a two-dimensional spin-statistics field theory, either hermitian or oriented. In order to treat both oriented and hermitian field theories, we first study the \mathbb{C} -linear spin-statistics field theory $Z_{\mathbb{C}} = Z \otimes_{\mathbb{R}} \mathbb{C}$.

As in [Example 3.12](#), $Z_{\mathbb{C}}$ is determined by a finite-dimensional semisimple \mathbb{C} -linear superalgebra A together with a bimodule isomorphism $\phi : {}_A A_A^* \xrightarrow{\sim} {}_A (-1)_A^f$. Let \tilde{Z} denote the $\text{SUPERVECT}_{\mathbb{C}}$ -valued spin field theory determined by A together with the isomorphism

$$\varphi = \phi \otimes \phi : A^* \otimes_A A^* \xrightarrow{\sim} (-1)^f \otimes_A (-1)^f \cong A.$$

Integrate \tilde{Z} to a $\text{SUPERVECT}_{\mathbb{C}}$ -valued oriented field theory $\int_{\text{Spins}/\text{Or}} \tilde{Z}$ controlled by the superalgebra algebra $B = A \oplus A^*$. Then \tilde{Z} canonically descends to a nonsuper \mathbb{C} -linear oriented field theory $\int_{\text{Spins}/\text{Or}} Z_{\mathbb{C}}$. Indeed, the isomorphism $\phi : A^* \xrightarrow{\sim} (-1)_A^f$

identifies B with the semidirect product for the parity-reversal action $A \rtimes \mathbb{Z}/2 = A \oplus A\epsilon$ where $\epsilon = \epsilon^{-1}$ is even and $\epsilon a = (-1)^{|a|}a\epsilon$. In particular, the bimodule ${}_B(-1)_B^f$ is canonically trivialized by $b \mapsto b\epsilon$. Let $\text{Forget}(B)$ denote the underlying nonsuper algebra of B . The trivialization ${}_B(-1)_B^f \cong {}_B B_B$ determines a Morita equivalence, namely $B/(1 - \epsilon) \oplus \Pi B/(1 + \epsilon)$, between B and $\text{Forget}(B)$. The nonsuper functor $\int_{\text{Spins}/\text{Or}} Z_{\mathbb{C}}$ assigns $\text{Forget}(B)$ to the point.

Finally, because the entire construction is equivariant under complex conjugation, if Z was real, then $\int_{\text{Spins}/\text{Or}} Z_{\mathbb{C}}$ naturally descends to a real oriented field theory, and if Z was hermitian, then $\int_{\text{Spins}/\text{Or}} Z_{\mathbb{C}}$ is naturally hermitian. Let us describe the hermitian case, as it is the more interesting one. In terms of algebras, if Z was hermitian, then A is stellar. By declaring that ϵ is real, B also becomes stellar, and hence so too is the Morita-equivalent purely even algebra $\text{Forget}(B)$. This stellar structure defines $\int_{\text{Spins}/\text{Or}} Z$ as a hermitian field theory. In most examples, the stellar structure on A comes from a $*$ -structure. In this case, B is also $*$. After tracing through the equivalences, one finds that the induced $*$ -structure on $\text{Forget}(B)$ is $b \mapsto b^* \epsilon^{|b|}$.

Remark 3.17 One can also understand [Example 3.16](#) in terms of categories of modules. The Morita class of the superalgebra $A = \tilde{Z}(\{\text{pt}\})$ is determined by the supercategory $\mathcal{A} = \text{SUPERMOD}_A$. $\tilde{Z}(\mathfrak{A})$ defines an action of $\mathbb{Z}/2$ on \mathcal{A} , and $\mathcal{B} = \text{SUPERMOD}_B$ is the supercategory of fixed points for this action. Being supercategories, \mathcal{A} and \mathcal{B} carry endo-superfunctors $(-1)_{\mathcal{A}}^f$ and $(-1)_{\mathcal{B}}^f$ which are the identity on objects and even morphisms but act by $(-1)^f$ on odd morphisms. The spin-statistics data $\tilde{Z}(\mathfrak{A}) \cong (-1)_{\mathcal{A}}^f$ provides a trivialization of $(-1)_{\mathcal{B}}^f$. This is precisely the data needed to factor $\mathcal{B} \simeq \mathcal{B}_{\text{ev}} \boxtimes \text{SUPERVECT}_{\mathbb{C}}$, where \mathcal{B}_{ev} is the plain category consisting of the even objects of \mathcal{B} , ie objects for which the trivialization $(-1)_{\mathcal{B}}^f \cong \text{id}$ acts as the identity.

The Morita equivalence between B and $\text{Forget}(B)$ in [Example 3.16](#) identifies \mathcal{B}_{ev} with $\text{MOD}_{\text{Forget}(B)}$. A straightforward calculation shows that \mathcal{B}_{ev} is also the underlying nonsuper category \mathcal{A}_0 of \mathcal{A} , ie the one with the same objects and even morphisms but with odd morphisms forgotten. Since the restriction to \mathcal{A}_0 of $(-1)_{\mathcal{A}}^f$ is trivial, and since we started with an isomorphism of superfunctors $(-1)_{\mathcal{A}}^f \cong \mathfrak{A}$, on the category \mathcal{A}_0 we have $\mathfrak{A} \cong \text{id}$. This is another way to see that $\mathcal{A}_0 = \mathcal{B}_{\text{ev}}$ defines an oriented field theory.

Definition 3.18 An étale-locally-spin field theory Z is *reflection-positive* if the unstructured field theory $\int_{\text{Spins}} Z = \int_{\text{Or}} \int_{\text{Spins}/\text{Or}} Z$ is reflection-positive.

We can now prove [Theorem 0.1](#), which asserts that all extended étale-locally-spin reflection-positive field theories are hermitian and satisfy spin-statistics.

Proof of Theorem 0.1 An étale-locally-spin field theory either satisfies spin-statistics or is spin-but-not-super. It suffices to show that if Z is a nonzero spin-but-not-super field theory then it is not reflection-positive; hermiticity will follow from Theorem 0.2. (A field theory is *zero* if it sends to the zero object all nonempty cobordisms. The zero field theory is vacuously reflection-positive and makes sense for all topological local structures.)

Suppose that Z is a nonzero spin-but-not-super field theory and consider the \mathbb{C} -linear spin-but-not-super field theory $Z_{\mathbb{C}} = Z \otimes_{\mathbb{R}} \mathbb{C}$. Let P be a connected oriented $(d-2)$ -dimensional manifold and let $\text{Spins} / \text{Or}(P)$ denote the groupoid of spin structures on P compatible with the chosen orientation. Since Z is nonzero, we can find P such that $Z(P, \sigma) \neq 0$ for at least one $\sigma \in \text{Spins} / \text{Or}(P)$. Then in particular $\text{Spins} / \text{Or}(P) \neq \emptyset$ and so $\text{Spins} / \text{Or}(P)$ is a torsor for $B(\mathbb{Z}/2) \times H^1(P; \mathbb{Z}/2)$.

Each choice of $\sigma \in \text{Spins} / \text{Or}(P)$ determines a dimensional reduction of $Z_{\mathbb{C}}$ to the two-dimensional \mathbb{C} -linear spin-but-not-super field theory $Z_{\mathbb{C}}(- \times (P, \sigma))$. By the classification of two-dimensional field theories [22], $A = Z_{\mathbb{C}}(\{\text{pt}\} \times (P, \Sigma))$ is a finite-dimensional semisimple algebra over \mathbb{C} , and so up to Morita equivalence we can assume $A = \mathbb{C}^{\oplus n}$ for some n . A bimodule isomorphism $A^* \otimes_A A^* \xrightarrow{\sim} A$ cannot permute the direct summands, and so the field theory $Z_{\mathbb{C}}(- \times (P, \sigma))$ is equivalent to a direct sum $\bigoplus_{i=1}^n Y_{\sigma}^{(i)}$ of complex-linear spin field theories each of which satisfies $A^{(i)} = Y_{\sigma}^{(i)}(\{\text{pt}\}) = \mathbb{C}$.

The two-dimensional spin-but-not-super field theory $Y_{\sigma}^{(i)}$ then satisfies

$$\int_{\text{Spins} / \text{Or}} Y_{\sigma}^{(i)}(\{\text{pt}\}) = A^{(i)} \oplus (A^{(i)})^* = \mathbb{C}[x]/(x^2 = 1)$$

with $\text{tr}(a + bx) = b$, and the complex Hilbert space is $\int_{\text{Spins} / \text{Or}} Y_{\sigma}^{(i)}(S^1) = \mathbb{C}^2$ with purely off-diagonal inner product. Thus

$$\int_{\text{Spins} / \text{Or}} Z(P \times S^1) = \int_{\sigma \in \text{Spins} / \text{Or}(P)} \bigoplus_i \mathbb{C}^2$$

is a nonzero direct sum of Hilbert spaces with purely off-diagonal inner product. Such an inner product cannot be positive-definite. □

Example 3.19 The hermitian spin-statistics field theory Z_n defined by $\text{Cliff}(n)$ from Example 3.12 is reflection-positive. Indeed, when n is odd, Example 3.16 implies that the hermitian field theory $\int_{\text{Spins} / \text{Or}} Z_n$ is controlled by

$$\text{Forget}(\text{Cliff}(n) \times \mathbb{Z}/2) \cong \text{Mat}_{\mathbb{C}}(2^{(n+1)/2}).$$

As discussed before [Definition 1.10](#), the Hilbert space $(\int_{\text{Spins}} Z_n)(S^1)$ is then the underlying real vector space of $(\int_{\text{Spins}/\text{Or}} Z_n)(S^1) = \mathbb{C}$ equipped with the real part of its hermitian pairing, which comes in turn from the $*$ -structure on $\text{Mat}_{\mathbb{C}}(2^{(n+1)/2})$. But $\langle v, v \rangle = |v|^2 \langle 1, 1 \rangle = |v|^2 \text{tr}(1) = |v|^2 2^{(n+1)/2} > 0$, so Z_n is reflection-positive. When n is even, $(\int_{\text{Spins}} Z_n)(S^1) \cong \mathbb{C}^{\oplus 2}$ with its positive-definite hermitian form, where the first copy of \mathbb{C} comes from “a boson on S^1 with its trivial spin structure” and the second from “a fermion on S^1 with its nontrivial spin structure”.

When n is even, $\text{Cliff}(n)$ also admits a symmetric Frobenius structure, and so defines a hermitian nonspin super field theory Z'_n . We can mimic [Remark 1.11](#) and integrate over $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$. The corresponding Hilbert space $(\int_{\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})} Z'_n)(S^1)$ is again a copy of $\mathbb{C}^{\oplus 2}$, but this time with the indefinite hermitian inner product.

4 Extension to higher categories

This last section explains how to extend the ideas in this paper to the higher-categorical setting championed by Lurie [\[20\]](#). We will assume familiarity with (∞, n) -categories and give only an outline of the necessary constructions. Following the by-now standard notation in the ∞ -categorical literature, we let SPACES denote the ∞ -category of topological spaces. For the remainder of this paper, let MAN_d denote the $(\infty, 1)$ -category coming from the topological category of d -dimensional smooth manifolds and local diffeomorphisms. Given an $(\infty, 1)$ -category \mathcal{X} with limits, a *topological local structure on d -dimensional manifolds valued in \mathcal{X}* is a sheaf $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$. We will not specify precisely the meaning of “sheaf”; one version is spelled out by Ayala [\[2\]](#). (Although that paper begins with “geometric” local structures, its main theorem asserts that the cobordism category it constructs from a geometric local structure \mathcal{F} depends only on the corresponding topological local structure $\tau\mathcal{F}$.) We will care most about the case when \mathcal{X} is an ∞ -topos, for example the ∞ -topos of sheaves of spaces on a site like $\text{SCH}_{\mathbb{R}}$ or $\text{CATAFFSCH}_{\mathbb{R}}$.

Generalizing [Lemma 1.1](#) and [Lemma 3.1](#), the following standard fact follows from the existence of good open covers together with the homotopy equivalence $\text{O}(d) \simeq \text{hom}_{\text{MAN}_d}(\mathbb{R}^d, \mathbb{R}^d)$; see Ayala and Francis [\[3\]](#):

Lemma 4.1 *The $(\infty, 1)$ -category of \mathcal{X} -valued topological local structures on d -dimensional manifolds is equivalent to the $(\infty, 1)$ -category $\mathcal{X}^{\text{O}(d)}$ of \mathcal{X} -objects equipped with an action by the topological group $\text{O}(d)$, the equivalence being given by sending a sheaf $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$ to $\mathcal{G}(\mathbb{R}^d) \in \mathcal{X}$. \square*

The sheaf corresponding to an object $X \in \mathcal{X}^{O(d)}$ can be constructed as follows. Given any d -manifold $M \in \text{MAN}_d$, define $X(M) = \text{maps}_{O(d)}(\text{Fr}_M, X) \in \mathcal{X}$, where $\text{Fr}_M \rightarrow M$ denotes the frame bundle, $\text{maps}_{O(d)}$ denotes $O(d)$ -equivariant maps, and \mathcal{X} is tensored over SPACES since it is an ∞ -topos. Then $X(\mathbb{R}^d) \simeq X$ in $\mathcal{X}^{O(d)}$. A special case is when $X \in \mathcal{X}$ is equipped with the trivial $O(d)$ action. Then $X(M) \simeq \text{maps}(M, X)$ is the *classical topological sigma-model with target X* .

Example 4.2 Let $G \rightarrow O(d)$ be a map of topological groups. A G -tangentia structure on a d -dimensional manifold M is a G -principal bundle $P \rightarrow M$ with an equivalence $P \times_G O(d) \simeq \text{Fr}_M$ of $O(d)$ -bundles. The sheaf $\text{MAN}_d \rightarrow \text{SPACES}$ of G -tangentia structures is classified by the quotient $O(d)/G$ with its natural $O(d)$ -action.

We now explain how to build an (∞, d) -category $\text{BORD}_d^{\mathcal{G}} = \text{BORD}_{0, \dots, d}^{\mathcal{G}}$ of “ \mathcal{G} -structured bordisms” for each topological local structure \mathcal{G} . For a suitable target \mathcal{V} , a *fully-extended \mathcal{G} -structured quantum field theory* will be a symmetric monoidal functor $\text{BORD}_d^{\mathcal{G}} \rightarrow \mathcal{V}$; the precise statement is in [Definition 4.4](#). We let BORD_d denote the “unstructured” bordism category whose construction is thoroughly outlined by Lurie [20], and for which all details have been provided by Calaque and Scheimbauer [7]. We will not review the construction of BORD_d itself.

Let \mathcal{Y} be an $(\infty, 1)$ -category with finite limits; for example, $\mathcal{Y} = \mathcal{X}$ an ∞ -topos. For each d , Haugseng [14] builds from \mathcal{Y} a symmetric monoidal (∞, d) -category $\text{SPANS}_d(\mathcal{Y})$. (The case when $\mathcal{Y} = \text{SPACES}$ is outlined, under the name FAM_d , in [20].) The objects of $\text{SPANS}_d(\mathcal{Y})$ are those of \mathcal{Y} , but a 1-morphism from X to Y in $\text{SPANS}_d(\mathcal{Y})$ is a *span* $X \leftarrow A \rightarrow Y$ in \mathcal{Y} , and higher morphisms are spans-between-spans. Following [14], we call symmetric monoidal functors $\text{BORD}_d \rightarrow \text{SPANS}_d(\mathcal{Y})$ *classical* (unstructured, fully extended) field theories valued in \mathcal{Y} .

Every \mathcal{Y} -valued topological local structure \mathcal{G} determines a classical field theory $\tilde{\mathcal{G}}$ (and the celebrated cobordism hypothesis of [20] implies that all classical field theories arise from topological local structures). Indeed, given a k -dimensional manifold M for $k \leq d$, set $\tilde{\mathcal{G}}(M) = \mathcal{G}(M \times \mathbb{R}^{d-k})$; if M has boundary, first glue on a “collar” $M \rightsquigarrow M \cup_{\partial M} (\partial M \times \mathbb{R}_{\geq 0})$. Then if M is a cobordism from N_1 to N_2 , the restriction maps $\mathcal{G}(M) \rightarrow \mathcal{G}(N_1)$ and $\mathcal{G}(M) \rightarrow \mathcal{G}(N_2)$ make $\mathcal{G}(M)$ into a span, and functoriality for the assignment $\tilde{\mathcal{G}}: \text{BORD}_d \rightarrow \text{SPANS}_d(\mathcal{Y})$ follows from the sheaf axiom for \mathcal{G} .

Remark 4.3 In the model of BORD_d from [7], k -morphisms are not k -dimensional manifolds, but rather d -dimensional manifolds properly submersed over \mathbb{R}^{d-k} . When using that model, one can directly define $\tilde{\mathcal{G}}: \text{BORD}_d \rightarrow \text{SPANS}_d(\mathcal{Y})$ simply as $\tilde{\mathcal{G}}(M) = \mathcal{G}(M)$.

Let $\{pt\} \in \mathcal{Y}$ denote the terminal object and $\mathcal{Y}_{\{pt\}}$ the “undercategory” of *pointed objects* $\{pt\} \rightarrow X$ in \mathcal{Y} . The logic of [20] is to construct $\text{BORD}_d^{\mathcal{G}}$ for \mathcal{G} a SPACES–valued topological local structure and then observe that there is a pullback square of symmetric monoidal (∞, d) –categories, where the vertical arrows are the obvious forgetful functors:

$$\begin{array}{ccc}
 \text{BORD}_d^{\mathcal{G}} & \longrightarrow & \text{SPANS}_d(\text{SPACES}_{\{pt\}}) \\
 \text{Forget } \mathcal{G} \downarrow & \lrcorner & \downarrow \text{forget the pointing} \\
 \text{BORD}_d & \xrightarrow{\mathcal{G}} & \text{SPANS}_d(\text{SPACES})
 \end{array}$$

Indeed, a \mathcal{G} –structured manifold M is nothing but a manifold M together with a pointing of the space $\mathcal{G}(M)$. We will reverse the logic and interpret the above pullback square as the definition of $\text{BORD}_d^{\mathcal{G}}$. Some care must be taken when replacing SPACES by an ∞ –topos \mathcal{X} , as in general very few objects $X \in \mathcal{X}$ admit “global” points $\{pt\} \rightarrow X$. The correct approach is to work with symmetric monoidal (∞, d) –categories “internal to \mathcal{X} ”; for the definition, see Haugseng [14] and Li-Bland [19].

Definition 4.4 Let \mathcal{X} be an ∞ –topos. By [19, Theorem 4.3], the symmetric monoidal (∞, d) –category $\text{SPANS}_d(\mathcal{X})$ constructed in [14] underlies an internal symmetric monoidal (∞, d) –category in \mathcal{X} , which in an abuse of notation we will also call $\text{SPANS}_d(\mathcal{X})$; the same argument implies also that $\text{SPANS}_d(\mathcal{X}_{\{pt\}})$ is naturally an internal symmetric monoidal (∞, d) –category in \mathcal{X} . Via the unique topos map $\text{SPACES} \rightarrow \mathcal{X}$, also view BORD_d as an internal symmetric monoidal (∞, d) –category in \mathcal{X} .

Let $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$ be an \mathcal{X} –valued topological local structure and $\tilde{\mathcal{G}}: \text{BORD}_d \rightarrow \text{SPANS}_d(\mathcal{X})$ the corresponding classical field theory. It extends canonically to a functor of internal symmetric monoidal (∞, d) –categories. The (∞, d) –category $\text{BORD}_d^{\mathcal{G}}$ of \mathcal{G} –structured bordisms is by definition the following pullback of internal symmetric monoidal (∞, d) –categories:

$$\begin{array}{ccc}
 \text{BORD}_d^{\mathcal{G}} & \longrightarrow & \text{SPANS}_d(\mathcal{X}_{\{pt\}}) \\
 \downarrow & \lrcorner & \downarrow \text{forget the pointing} \\
 \text{BORD}_d & \xrightarrow{\mathcal{G}} & \text{SPANS}_d(\mathcal{X})
 \end{array}$$

Let \mathcal{V} be a symmetric monoidal (∞, d) –category internal to \mathcal{X} . A \mathcal{G} –structured field theory valued in \mathcal{V} is a functor $\text{BORD}_d^{\mathcal{G}} \rightarrow \mathcal{V}$ of internal symmetric monoidal (∞, d) –categories.

The need to work with internal categories in Definition 4.4 is in some sense unavoidable: “functors internal to \mathcal{X} ” is the appropriate language with which to impose that a functor be “smooth” for families parametrized by objects of \mathcal{X} . But one can also describe \mathcal{G} –structured field theories “externally” in terms of the lifting problems in Definition 1.4 and Definition 3.8. Given an ∞ –topos \mathcal{X} and a symmetric monoidal (∞, d) –category \mathcal{V} internal to \mathcal{X} , the papers [14; 19] construct a symmetric monoidal (∞, d) –category $\text{SPANS}_d(\mathcal{X}; \mathcal{V})$ whose k –morphisms are “bundles of k –morphisms in \mathcal{V} over k –fold spans in \mathcal{X} ”. Such a notion makes sense exactly because \mathcal{V} is internal to \mathcal{X} : by definition, a *bundle of k –morphisms in \mathcal{V} over $X \in \mathcal{X}$* is a map from X to the \mathcal{X} –object of k –morphisms in \mathcal{V} . After unpacking adjunctions, one finds:

Proposition 4.5 *Let \mathcal{X} be an ∞ –topos, $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$ a topological local structure, and \mathcal{V} a symmetric monoidal (∞, d) –category internal to \mathcal{X} . Then the data of a \mathcal{G} –structured field theory $\text{BORD}_d^{\mathcal{G}} \rightarrow \mathcal{V}$ is the same as the data of a lift:*

$$\begin{array}{ccc}
 & \text{SPANS}_d(\mathcal{X}; \mathcal{V}) & \\
 & \nearrow & \downarrow \text{forget the } \mathcal{V}\text{-data} \\
 \text{BORD}_d & \xrightarrow{\tilde{\mathcal{G}}} & \text{SPANS}_d(\mathcal{X})
 \end{array}
 \quad \square$$

Corollary 4.6 *Let \mathcal{X} be an ∞ –topos and \mathcal{V} an internal-to- \mathcal{X} symmetric monoidal (∞, d) –category with duals in the sense of [14]. Let $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$ be a topological local structure, and $\mathcal{G}(\{\text{pt}\}) = \mathcal{G}(\mathbb{R}^d)$ the corresponding object in $\mathcal{X}^{O(d)}$. Assuming the cobordism hypothesis, \mathcal{G} –structured field theories valued in \mathcal{V} are classified by $O(d)$ –equivariant bundles of \mathcal{V} –objects over $\mathcal{G}(\{\text{pt}\})$. \square*

We conclude by extending the examples from this paper. Note that under Lemma 4.1, the sheaves Or and Spins of orientations and spin structures correspond, respectively, to the actions of $O(d)$ on the 0– and 1–truncations $\pi_{\leq 0}O(\infty)$ and $\pi_{\leq 1}O(\infty)$, or equivalently to the trivial torsors for these groups. Any ∞ –topos \mathcal{X} admits a notion of torsor for topological groups: $X \in \mathcal{X}^G$ is a G –torsor if the map $G \times X \rightarrow X \times X$, $(g, x) \mapsto (gx, x)$ is an equivalence. An \mathcal{X} –valued topological local structure $\mathcal{G}: \text{MAN}_d \rightarrow \mathcal{X}$ is *locally Or* (resp. *locally Spins*) if $\mathcal{G}(\mathbb{R}^d)$ is a torsor for $\pi_{\leq 0}O(\infty)$ (resp. $\pi_{\leq 1}O(\infty)$). Suppose \mathcal{X} is the ∞ –topos of sheaves (valued in SPACES) on some site (with some subcanonical topology) containing the category $\text{AFFSCH}_{\mathbb{R}}$ of affine schemes over \mathbb{R} . Then there is a canonical \mathcal{X} –valued topological local structure $\text{Her}: \text{MAN}_d \rightarrow \mathcal{X}$ whose value on \mathbb{R}^d is $\text{Spec}(\mathbb{C})$. If \mathcal{X} is the ∞ –topos of sheaves on some site containing $\text{CATAFFSCH}_{\mathbb{R}}$, then similarly there is a canonical topological local structure $\text{HerSpinStats}: \mathbb{R}^d \mapsto \text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$.

For \mathcal{V} a suitable target symmetric monoidal (∞, d) -category internal to \mathcal{X} , we can then define *hermitian* and *hermitian spin-statistics* field theories as being Her- and HerSpinStats-structured field theories in the sense of [Definition 4.4](#). Note that the details of the ∞ -topos \mathcal{X} are largely irrelevant: given [Proposition 4.5](#), what matters for hermitian and spin-statistics field theories are the symmetric monoidal (∞, d) -categories of X -points of \mathcal{V} for X ranging over the possible values $\text{Spec}(\mathbb{R})$, $\text{Spec}(\mathbb{C})$, $\text{Spec}(\text{SUPERVECT}_{\mathbb{C}})$, ... of Her and HerSpinStats.

One standard criterion for deciding whether a proposed target \mathcal{V} is suitable is that “near the top” \mathcal{V} should look like VECT. More precisely, any symmetric monoidal (∞, d) -category \mathcal{V} determines a symmetric monoidal $(\infty, 1)$ -category $\Omega^{d-1}\mathcal{V}$ of endomorphisms of the identity $(d-2)$ -morphism on the identity $(d-3)$ -morphism on ... on the unit object in \mathcal{V} . The passage $\mathcal{V} \mapsto \Omega^{d-1}\mathcal{V}$ makes sense also for internal categories. The “looks like VECT near the top” criterion then says that for R a commutative \mathbb{R} -algebra, the $\text{Spec}(R)$ -points of $\Omega^{d-1}\mathcal{V}$ should be MOD_R , and that for \mathcal{C} a categorified commutative \mathbb{R} -algebra, the $\text{Spec}(\mathcal{C})$ -points of $\Omega^{d-1}\mathcal{V}$ should be \mathcal{C} itself. This assures, for example, that if $Z: \text{BORD}_d^{\text{Her}} \rightarrow \mathcal{V}$ is a fully-extended hermitian field theory, then its restriction

$$Z|_{\text{BORD}_{d-1,d}^{\text{Her}}} : \text{BORD}_{d-1,d}^{\text{Her}} = \Omega^{d-1}\text{BORD}_d \rightarrow \Omega^{d-1}\mathcal{V} = \text{VECT}$$

unpacks to a hermitian unextended field theory in the sense of [Definition 1.4](#).


An extended field theory $Z: \text{BORD}_d \rightarrow \mathcal{V}$ is *reflection-positive* if the unextended field theory $Z|_{\text{BORD}_{d-1,d}}$ is reflection-positive in the sense of [Definition 1.9](#). (This is different from the notion in [\[12\]](#) of reflection-positivity for extended field theories, which requires extra positivity data to be specified in high codimension.) [Definition 1.10](#) and [Definition 3.18](#) then apply to extended hermitian and spin-statistics field theories.

One could worry that restricting a field theory just to its top part is too much loss of information. The following observation is due to Chris Schommer-Pries:

Lemma 4.7 *Let \mathcal{V} be some symmetric monoidal (∞, d) -category with a zero object, and $Z: \text{BORD}_d^{\mathcal{G}} \rightarrow \mathcal{V}$ be a \mathcal{G} -structured extended field theory for some topological local structure \mathcal{G} . Suppose that the unextended field theory $Z|_{\text{BORD}_{d-1,d}^{\mathcal{G}}}$ is zero in the sense that it vanishes on all nonempty inputs. (Symmetric monoidality forces $Z(\emptyset)$ to be the unit object of $\Omega^{d-1}\mathcal{V}$.) Then Z is zero.*

Proof A k -morphism F is zero if and only if its identity $(k+1)$ -morphism id_F is zero. It therefore suffices to show that for N an arbitrary \mathcal{G} -structured $(d-1)$ -dimensional cobordism, $Z(N \times [0, 1]): Z(N) \rightarrow Z(N)$ is the zero d -morphism. But

the \mathcal{G} -structured cobordism $N \times [0, 1]$ can be factored through $N \sqcup S^{d-1}$ where the sphere S^{d-1} is given the \mathcal{G} -structure that extends to the disk D^d :



By assumption, $Z(S^{d-1}) = 0$, since S^{d-1} is closed, and so

$$Z(N \sqcup S^{d-1}) \cong Z(N) \otimes Z(S^{d-1}) = 0.$$

Only a zero morphism can factor through a zero object, and so $Z(N \times [0, 1]) = 0$. \square

Only the zero field theory is compatible with multiple topological local structures. **Lemma 4.7** assures that if a \mathcal{G} -structured fully extended field theory Z is not zero, then neither is its restriction $Z|_{\text{BORD}_{d-2,d-1,d}^{\mathcal{G}}}$ to a once-extended theory, and so $Z|_{\text{BORD}_{d-2,d-1,d}^{\mathcal{G}}}$ detects the local structure \mathcal{G} . Along with **Theorem 0.1**, we conclude:

Corollary 4.8 *Reflection-positive étale-locally-spin fully-extended field theories are necessarily unitary and satisfy spin-statistics.* \square

Acknowledgements I would like to thank K Costello and D Gaiotto for asking the questions that led to this note, D Freed for emphasizing the importance of positivity in the spin-statistics theorem, A Chirvasitu and E Elmanto for many discussions about Galois theory and locally presentable categories, and C Schommer-Pries for providing the proof of **Lemma 4.7**. I would like particularly to thank the referee for their considerable engagement with the organization and presentation of the ideas in this paper; their comments and suggestions led to many improvements. Some of this work occurred while I was a visitor to the Perimeter Institute for Theoretical Physics in Waterloo, Ontario, where I was treated to generous hospitality and a great work environment. Research at Perimeter Institute is supported by the Government of Canada through Industry Canada and by the Province of Ontario through the Ministry of Economic Development & Innovation. This research is also supported by the NSF grant DMS-1304054.

References

- [1] **M Atiyah**, *Topological quantum field theories*, Inst. Hautes Études Sci. Publ. Math. 68 (1988) 175–186 [MR](#)
- [2] **D Ayala**, *Geometric cobordism categories*, PhD thesis, Stanford University (2009) [MR](#)
Available at <http://search.proquest.com/docview/304999097>
- [3] **D Ayala, J Francis**, *Factorization homology of topological manifolds*, J. Topol. 8 (2015) 1045–1084 [MR](#)
- [4] **J C Baez, J Dolan**, *From finite sets to Feynman diagrams*, from “Mathematics unlimited—2001 and beyond” (B Engquist, W Schmid, editors), Springer, Berlin (2001) 29–50 [MR](#)
- [5] **G J Bird**, *Limits in 2-categories of locally presentable categories*, PhD thesis, University of Sydney (1984) Available at <http://maths.mq.edu.au/~street/BirdPhD.pdf>
- [6] **M Brandenburg, A Chirvasitu, T Johnson-Freyd**, *Reflexivity and dualizability in categorified linear algebra*, Theory Appl. Categ. 30 (2015) 808–835 [MR](#)
- [7] **D Calaque, C Scheimbauer**, *A note on the (∞, n) -category of cobordisms*, preprint (2015) [arXiv](#)
- [8] **A Chirvasitu, T Johnson-Freyd**, *The fundamental pro-groupoid of an affine 2-scheme*, Appl. Categ. Structures 21 (2013) 469–522 [MR](#)
- [9] **P Deligne**, *Catégories tensorielles*, Mosc. Math. J. 2 (2002) 227–248 [MR](#)
- [10] **C L Douglas, C Schommer-Pries, N Snyder**, *The balanced tensor product of module categories*, preprint (2014) [arXiv](#)
- [11] **S Eilenberg**, *Abstract description of some basic functors*, J. Indian Math. Soc. 24 (1960) 231–234 [MR](#)
- [12] **D S Freed, M J Hopkins**, *Reflection positivity and invertible topological phases*, preprint (2016) [arXiv](#)
- [13] **N Ganter, M Kapranov**, *Symmetric and exterior powers of categories*, Transform. Groups 19 (2014) 57–103 [MR](#)
- [14] **R Haugseng**, *Iterated spans and “classical” topological field theories*, preprint (2014) [arXiv](#)
- [15] **M Kapranov**, *Supergeometry in mathematics and physics*, preprint (2015) [arXiv](#)
- [16] **G M Kelly**, *Basic concepts of enriched category theory*, London Math. Soc. Lecture Note Ser. 64, Cambridge University Press (1982) [MR](#)
- [17] **A Kleshchev**, *Linear and projective representations of symmetric groups*, Cambridge Tracts in Mathematics 163, Cambridge University Press (2005) [MR](#)

- [18] **L G Lewis, Jr.** *When projective does not imply flat, and other homological anomalies*, Theory Appl. Categ. 5 (1999) 202–250 [MR](#)
- [19] **D Li-Bland**, *The stack of higher internal categories and stacks of iterated spans*, preprint (2015) [arXiv](#)
- [20] **J Lurie**, *On the classification of topological field theories*, from “Current developments in mathematics” (D Jerison, B Mazur, T Mrowka, W Schmid, R Stanley, S-T Yau, editors), Int. Press, Somerville, MA (2009) 129–280 [MR](#)
- [21] **V Ostrik**, *On symmetric fusion categories in positive characteristic*, preprint (2015) [arXiv](#)
- [22] **C J Schommer-Pries**, *The classification of two-dimensional extended topological field theories*, PhD thesis, University of California, Berkeley (2009) [MR](#) Available at <https://arxiv.org/abs/1112.1000>
- [23] **G Segal**, *The definition of conformal field theory*, from “Topology, geometry and quantum field theory” (U Tillmann, editor), London Math. Soc. Lecture Note Ser. 308, Cambridge University Press (2004) 421–577 [MR](#)
- [24] **S Stolz, P Teichner**, *Supersymmetric field theories and generalized cohomology*, from “Mathematical foundations of quantum field theory and perturbative string theory” (H Sati, U Schreiber, editors), Proc. Sympos. Pure Math. 83, Amer. Math. Soc., Providence, RI (2011) 279–340 [MR](#)
- [25] **R F Streater, A S Wightman**, *PCT, spin and statistics, and all that*, W A Benjamin, New York (1964) [MR](#)
- [26] **C E Watts**, *Intrinsic characterizations of some additive functors*, Proc. Amer. Math. Soc. 11 (1960) 5–8 [MR](#)

Department of Mathematics, Northwestern University
2033 Sheridan Road, Evanston, IL 60208, United States

theojf@math.northwestern.edu

<http://math.northwestern.edu/~theojf>

Received: 28 October 2015 Revised: 6 June 2016