

# ANNALS OF K-THEORY

vol. 4 no. 4 2019

**On the intersection motive of certain Shimura varieties:  
the case of Siegel threefolds**

Jörg Wildeshaus



A JOURNAL OF THE K-THEORY FOUNDATION

# On the intersection motive of certain Shimura varieties: the case of Siegel threefolds

Jörg Wildeshaus

We construct a Hecke-equivariant Chow motive whose realizations equal intersection cohomology of Siegel threefolds with regular algebraic coefficients. As a consequence, we are able to define Grothendieck motives for Siegel modular forms.

## 0. Introduction

The purpose of this paper is the construction and analysis of the *intersection motive* of Kuga–Sato families over a Siegel threefold relative to its Satake–(Baily–Borel) compactification. As in earlier work on Hilbert–Blumenthal varieties [Wildeshaus 2012b], Picard surfaces [Wildeshaus 2015], and more generally, Picard varieties of arbitrary dimension [Cloître 2017], the use of the formalism of *weight structures* [Bondarko 2010] proves to be successful for dealing with a problem, for which explicit geometrical methods seem inefficient.

However, Siegel threefolds present a characteristic feature different from the cases treated so far: the dimension of the boundary of their Satake–(Baily–Borel) compactification is equal to one. In particular, it is strictly positive.

As a consequence, the context of *geometrical motives*, i.e., motives over a point, is no longer adapted to the problem. Let us explain why.

The present construction, as the preceding ones, depends on *absence of weights*  $-1$  and  $0$  in the *boundary motive*. To prove absence of weights, the idea remains, as previously, to employ *realizations*. But then, realizations need to detect weights (and therefore, their absence). One may expect this to be true in general; let us agree to refer to that principle as *weight conservativity*. To date, weight conservativity is *proved* for the restriction of the (generic)  $\ell$ -adic realization to the category of *motives of abelian type* of characteristic zero [Wildeshaus 2018b].

---

Partially supported by the *Agence Nationale de la Recherche*, project “Régulateurs et formules explicites”.

*MSC2010*: primary 14G35; secondary 11F32, 11F46, 14C25, 14F20, 14F25.

*Keywords*: Siegel threefolds, weight structures, intersection motive, motives for Siegel modular forms.

However, unless the boundary of the Baily–Borel compactification of a given Shimura variety  $M$  is of dimension zero, its boundary motive, as well as the boundary motive of any Kuga–Sato family  $B$  over  $M$ , is in general not of abelian type; this is in any case true if  $M$  is a Siegel threefold. Concretely, this means that even if the realization of the boundary motive were proved to avoid weights  $-1$  and  $0$ , we could not formally conclude that the same is true for the boundary motive itself.

This is where *relative motives*, together with the *formalism of six operations*, enter. Denoting by  $j$  the open immersion of  $M$  into its Baily–Borel compactification  $M^*$ , by  $i$  its closed complement, and by  $\mathbb{1}_M$  the structural motive over  $M$ , there is an exact triangle

$$i_*i^*j_*\mathbb{1}_M[-1] \rightarrow j!\mathbb{1}_M \rightarrow j_*\mathbb{1}_M \rightarrow i_*i^*j_*\mathbb{1}_M$$

of motives over  $M^*$ . The boundary motive of  $M$  is isomorphic to the dual of the direct image of  $i_*i^*j_*\mathbb{1}_M$  under the structure morphism of  $M^*$ . More generally, the boundary motive of  $B$  is isomorphic to the dual of the direct image of  $i_*i^*j_*\pi_*(\mathbb{1}_B)$ , where  $\pi : B \rightarrow M$  denotes the projection of the Kuga–Sato family  $B$  to its base.

It is then true that the relative motive  $i_*i^*j_*\pi_*(\mathbb{1}_B)$  over  $M^*$  is of abelian type.

This suggests our strategy of proof. First, identify the  $\ell$ -adic realization of  $i_*i^*j_*\pi_*(\mathbb{1}_B)$ , or more generally, of  $i_*i^*j_*\mathcal{V}$ , for direct factors  $\mathcal{V}$  of  $\pi_*(\mathbb{1}_B)$ ; in the cases where weights  $0$  and  $1$  are avoided, weight conservativity tells us that  $i_*i^*j_*\mathcal{V}$  itself avoids weights  $0$  and  $1$ . Second, apply the direct image  $a_*$  associated to the structure morphism  $a$  of  $M^*$ . It is proper, therefore, the functor  $a_*$  is *weight exact*. In particular, if  $i_*i^*j_*\mathcal{V}$  avoids weights  $0$  and  $1$ , then so does  $a_*i_*i^*j_*\mathcal{V}$ . The corresponding direct factor of the boundary motive of  $B$  thus avoids weights  $-1$  and  $0$ .

It may be useful to remark that if  $M$  is a Hilbert–Blumenthal or Picard variety, then there is essentially no difference between  $i_*i^*j_*\mathcal{V}$  and its direct image under  $a$ , since the latter is of relative dimension zero on the boundary of  $M^*$ .

The passage from geometrical motives to relative motives necessitates a certain number of technical adjustments. For better legibility, we decided to separate these from the present text. The result is [Wildeshaus 2018a]; it contains in particular the identification of the boundary motive and the dual of  $a_*i_*i^*j_*\pi_*(\mathbb{1}_B)$  mentioned above.

Compared to the cases treated earlier, another feature of the boundary of Siegel threefolds is new: its canonical stratification is not reduced to a single type of strata. Indeed, in the boundary, one finds a closed stratum of dimension zero, the so-called *Siegel stratum*, and its complement, the so-called *Klingen stratum*, which is a disjoint union of (open) modular curves. Control of the weights avoided by the restrictions of the  $\ell$ -adic realization  $R_\ell(i_*i^*j_*\pi_*(\mathbb{1}_B))$  of  $i_*i^*j_*\pi_*(\mathbb{1}_B)$  to the two strata is related to but does not a priori determine the weights avoided by  $R_\ell(i_*i^*j_*\pi_*(\mathbb{1}_B))$ .

In fact, the precise relation is given by a long exact localization sequence. Its control is not obvious. In an earlier attempt, we succeeded to identify sufficiently many terms in this sequence, and (above all) certain morphisms, to prove absence of weights 0 and 1. This approach is technically difficult; moreover, it does not use the auto-duality property of the coefficients. Indeed, the device dual to the localization sequence is the colocalization sequence; even when the coefficients are auto-dual, the two sequences cannot be related. It turns out that both problems admit the same solution. Namely, the theory of *intermediate extensions* allows one to represent  $R_\ell(i^*j_*\pi_*(\mathbb{1}_B))$  as an extension of two “halves”, one dual to the other, and both related to the intermediate extension  $j_{!*}\pi_*(\mathbb{1}_B)$ . This observation is equally integrated in [Wildeshaus 2018a]; for our purposes, its concrete interest is to divide by two the number of cohomological degrees for which absence of weights has to be tested, and to reduce the number of morphisms in the localization sequence, which need to be identified, to zero.

The rôle of the intermediate extension is not only technical. It turns out that the dual of its direct image under  $a$  is canonically isomorphic to the *interior motive*, which according to [Wildeshaus 2009] can be defined as soon as the boundary motive avoids weights  $-1$  and  $0$ . This motivates the slight change of terminology in the title, as compared to the earlier work mentioned above [Wildeshaus 2012b; 2015; Cloître 2017].

Let us now give a more detailed account of the content of the present article. Section 1 contains the statement of our main result, Theorem 1.6. Denote by  $\mathrm{GSp}_{4,\mathbb{Q}}$  the group of symplectic similitudes of a fixed four-dimensional  $\mathbb{Q}$ -vector space  $V$ . As will be recalled, irreducible representations of  $\mathrm{GSp}_{4,\mathbb{Q}}$  are indexed by weights  $\underline{\alpha}$  depending on three integral parameters:  $\underline{\alpha} = \alpha(k_1, k_2, r)$ . The weight  $\underline{\alpha}$  is dominant if and only if  $k_1 \geq k_2 \geq 0$ ; it is regular if and only if  $k_1 > k_2 > 0$ . Denote by  $V_{\underline{\alpha}}$  the irreducible representation of highest weight  $\underline{\alpha}$ . According to the main result from [Ancona 2015] (which will be recalled in Theorem 1.4), there is a Chow motive  ${}^\alpha\mathcal{V}$  over the Siegel threefold  $M$  whose cohomological (Hodge theoretic or  $\ell$ -adic) realizations equal the classical *canonical construction*  $\mu(V_{\underline{\alpha}})$ . Part (a) of Theorem 1.6 then states that  $i^*j_*{}^\alpha\mathcal{V}$  is of abelian type. Part (b) asserts that if  $\underline{\alpha}$  is regular, then  $i^*j_*{}^\alpha\mathcal{V}$  avoids weights 0 and 1. It has recently become increasingly important to determine the precise interval containing  $[0, 1]$  of weights avoided by  $i^*j_*{}^\alpha\mathcal{V}$ . Theorem 1.6(b) gives a complete answer: putting  $k := \min(k_1 - k_2, k_2)$ , the motive  $i^*j_*{}^\alpha\mathcal{V}$  avoids all the weights between  $-k + 1$  and  $k$ , while both weights  $-k$  and  $k + 1$  do occur. Interestingly, this result does not depend on the level of the Siegel threefold. We then list the main consequences of this result (Corollaries 1.7, 1.8, 1.9, 1.11, 1.13), applying the theory developed in [Wildeshaus 2018a].

Section 2 is devoted to the proof of Theorem 1.6. As in previous cases, our control of smooth *toroidal compactifications* of  $M$  is sufficiently explicit to verify

that, as stated in [Theorem 1.6\(a\)](#), the motive  $i^* j_*^\alpha \mathcal{V}$  is indeed of abelian type. Given this result, and weight conservativity of the restriction of the  $\ell$ -adic realization  $R_\ell$ , part (b) of [Theorem 1.6](#) may be checked on the image of  $i^* j_*^\alpha \mathcal{V}$  under  $R_\ell$ . Given that  ${}^\alpha \mathcal{V}$  realizes to give  $\mu(V_\alpha)$ , the restriction of  $R_\ell(i^* j_*^\alpha \mathcal{V})$  to the (Siegel and Klingen) strata can be computed following a standard pattern, employing Pink's and Kostant's theorems. This computation ([Theorem 2.3](#)) is considerably simplified by results of [\[Lemma 2015\]](#). It remains to glue the information coming from the strata, in order to get control of the weights on the whole boundary. The part of [Theorem 1.6\(b\)](#) asserting that weights  $-k$  and  $k + 1$  occur in  $R_\ell(i^* j_*^\alpha \mathcal{V})$  ([Proposition 2.9](#)) is the single ingredient requiring a proof longer than any other.

In the final [Section 3](#), we give the necessary ingredients to perform the construction of the Grothendieck motive associated to a (Siegel) automorphic form with coefficients in an irreducible representation with regular highest weight ([Definition 3.5](#)). This is the analogue for Siegel threefolds of the main result from [\[Scholl 1990\]](#). On the level of Galois representations, our definition coincides with Weissauer's [\[2005, Theorem I\]](#). We also recover Urban's result [\[2005, Théorème 1\]](#) on characteristic polynomials associated to Frobenii ([Corollary 3.7](#)).

**Conventions.** We make use of the triangulated  $\mathbb{Q}$ -linear categories  $DM_{\mathbb{B},c}(X)$  of *constructible Beilinson motives* over  $X$  [\[Cisinski and Déglise 2009, Definition 15.1.1\]](#), indexed by schemes  $X$  over  $\text{Spec } \mathbb{Q}$ , which are separated and of finite type. As in [\[Cisinski and Déglise 2009\]](#), the symbol  $\mathbb{1}_X$  is used to denote the unit for the tensor product in  $DM_{\mathbb{B},c}(X)$ . We employ the full formalism of six operations developed in [\[loc. cit.\]](#). The reader may choose to consult [\[Hébert 2011, Section 2\]](#) or [\[Wildeshaus 2012a, Section 1\]](#) for concise presentations of this formalism.

Beilinson motives can be endowed with a canonical weight structure, thanks to the main results from [\[Hébert 2011\]](#); see [\[Bondarko 2010, Proposition 6.5.3\]](#) for the case  $X = \text{Spec } k$ , for a field  $k$  of characteristic zero. We refer to it as the *motivic weight structure*. Following [\[Wildeshaus 2012a, Definition 1.5\]](#), the category  $CHM(X)_{\mathbb{Q}}$  of *Chow motives* over  $X$  is defined as the heart  $DM_{\mathbb{B},c}(X)_{w=0}$  of the motivic weight structure on  $DM_{\mathbb{B},c}(X)$ .

A scheme is said to be *nilregular* if the underlying reduced scheme is regular in the usual sense.

## 1. Statement of the main result

In order to state our main result ([Theorem 1.6](#)), let us introduce the situation we are going to consider. The  $\mathbb{Q}$ -scheme  $M^K$  is a *Siegel threefold*, and the Chow motive  ${}^\alpha \mathcal{V}$  over  $M^K$  is associated to a *dominant weight*  $\underline{\alpha} = (k_1, k_2, r) \in \mathbb{Z}^3$ ,  $k_1 \geq k_2 \geq 0$  (see below for the precise normalizations). Denote by  $j$  the open immersion of  $M^K$  into its *Satake–(Baily–Borel) compactification*  $(M^K)^*$ , and by  $i : \partial(M^K)^* \hookrightarrow (M^K)^*$

the immersion of the complement of  $M^K$  in  $(M^K)^*$  (with the reduced scheme structure, say). Recall the following.

**Definition 1.1** (cf. [Wildeshaus 2018a, Definition 2.1(a)]). Let  $CHM(M^K)_{\mathbb{Q}, \partial w \neq 0, 1}$  denote the full subcategory of  $CHM(M^K)_{\mathbb{Q}}$  of objects  $V$  such that  $i^* j_* V$  is without weights 0 and 1.

**Theorem 1.6** implies that in our setting, the motive  ${}^\alpha \mathcal{V} \in CHM(M^K)_{\mathbb{Q}}$  belongs to  $CHM(M^K)_{\mathbb{Q}, \partial w \neq 0, 1}$  if and only if  $\alpha$  is *regular*:  $k_1 > k_2 > 0$ . More precisely, putting  $k := \min(k_1 - k_2, k_2)$ , the motive  $i^* j_* {}^\alpha \mathcal{V}$  is without weights  $-k + 1, -k + 2, \dots, k$ . The proof of **Theorem 1.6** is given in **Section 2**. It is an application of [Wildeshaus 2018a, Theorem 4.4]; in order to verify the hypotheses of the latter, we heavily rely on results from [Lemma 2015].

Fix a four-dimensional  $\mathbb{Q}$ -vector space  $V$ , together with a  $\mathbb{Q}$ -valued nondegenerate symplectic bilinear form  $J$ .

**Definition 1.2.** The group scheme  $G$  over  $\mathbb{Q}$  is defined as the group of symplectic similitudes

$$G := \mathrm{GSp}(V, J) \subset \mathrm{GL}(V).$$

Thus,  $G$  is reductive, and for any  $\mathbb{Q}$ -algebra  $R$ , the group  $G(R)$  equals

$$\{g \in \mathrm{GL}(V \otimes_{\mathbb{Q}} R) : \exists \lambda(g) \in R^*, J(g \bullet, g \bullet) = \lambda(g) \cdot J(\bullet, \bullet)\}.$$

In particular, the similitude norm  $\lambda(g)$  defines a canonical morphism

$$\lambda : G \rightarrow \mathbb{G}_{m, \mathbb{Q}}.$$

The group  $G$  is split over  $\mathbb{Q}$ , and its center  $Z(G)$  equals  $\mathbb{G}_{m, \mathbb{Q}} \subset \mathrm{GL}(V)$  (inclusion of scalar automorphisms). Maximal  $\mathbb{Q}$ -split tori, together with an inclusion into a Borel subgroup of  $G$ , are in bijection with symplectic  $\mathbb{Q}$ -bases of  $V$ , in which  $J$  acquires the  $4 \times 4$ -matrix

$$\begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix},$$

also denoted by  $J$ . Here as in the sequel, we denote by  $I_2$  the  $2 \times 2$ -matrix representing the identity. Fix one such basis  $(e_1, e_2, e_3, e_4)$ , use it to identify  $G$  with the subgroup  $\mathrm{GSp}_{4, \mathbb{Q}}$  of  $\mathrm{GL}_{4, \mathbb{Q}}$  of matrices  $g$  satisfying the relation

$${}^t g J g = \lambda(g) \cdot J,$$

the maximal split torus with the subgroup  $T$  of diagonal matrices

$$\{\mathrm{diag}(a, b, a^{-1}q, b^{-1}q) \in \mathrm{GL}_{4, \mathbb{Q}}\},$$

and the Borel subgroup with the subgroup of matrices stabilizing the flag of totally isotropic subspaces  $(e_1)_{\mathbb{Q}} \subset (e_1, e_2)_{\mathbb{Q}}$  of  $V$ . We consider triplets  $(k_1, k_2, r) \in \mathbb{Z}^3$

satisfying the congruence relation

$$r \equiv k_1 + k_2 \pmod{2}.$$

To such a triplet, let us associate the (representation-theoretic) weight

$$\alpha(k_1, k_2, r) : T \rightarrow \mathbb{G}_{m, \mathbb{Q}}, \quad \text{diag}(a, b, a^{-1}q, b^{-1}q) \mapsto a^{k_1} b^{k_2} q^{-(r+k_1+k_2)/2}.$$

Note that restriction of  $\alpha(k_1, k_2, r)$  to  $T \cap \text{Sp}(V, J)$  corresponds to the projection onto  $(k_1, k_2)$ . In particular, the weight  $\alpha(k_1, k_2, r)$  is dominant if and only if  $k_1 \geq k_2 \geq 0$ ; it is regular if and only if  $k_1 > k_2 > 0$ . Note also that the composition of  $\alpha(k_1, k_2, r)$  with the cocharacter

$$\mathbb{G}_{m, \mathbb{Q}} \rightarrow T, \quad x \mapsto \text{diag}(x, x, x, x)$$

equals

$$\mathbb{G}_{m, \mathbb{Q}} \rightarrow \mathbb{G}_{m, \mathbb{Q}}, \quad x \mapsto x^{-r}.$$

The character  $\lambda$  on  $T$  equals  $\alpha(0, 0, -2)$ , and  $\det = \lambda^2$ .

**Definition 1.3.** The analytic space  $\mathcal{H}$  is defined as the subspace of  $M_2(\mathbb{C})$  of those complex  $2 \times 2$ -matrices, which are symmetrical, and whose imaginary part is (positive or negative) definite:

$$\mathcal{H} := \{\tau \in M_2(\mathbb{C}) : {}^t\tau = \tau \text{ and } \text{Im}(\tau) \text{ definite}\}.$$

The group of real points  $G(\mathbb{R})$  acts on  $\mathcal{H}$  by analytical automorphisms [Pink 1989, Example 2.7]. In fact,  $(G, \mathcal{H})$  are *pure Shimura data* [Pink 1989, Definition 2.1]. Their *reflex field* [Pink 1989, Section 11.1] equals  $\mathbb{Q}$ . Given that  $Z(G) = \mathbb{G}_{m, \mathbb{Q}}$ , the Shimura data  $(G, \mathcal{H})$  satisfy condition (+) from [Wildeshaus 2007, Section 5].

Let us now fix additional data:

- (A) an open compact subgroup  $K$  of  $G(\mathbb{A}_f)$  which is neat [Pink 1989, Section 0.6],
- (B) a triplet  $(k_1, k_2, r) \in \mathbb{Z}^3$  satisfying the congruence

$$r \equiv k_1 + k_2 \pmod{2},$$

and in addition,

$$k_1 \geq k_2 \geq 0.$$

In other words, the character  $\underline{\alpha} := \alpha(k_1, k_2, r)$  is dominant.

These data are used as follows. The Shimura variety  $M^K := M^K(G, \mathcal{H})$  is smooth over  $\mathbb{Q}$ . This is the Siegel threefold of level  $K$ . According to [Pink 1989, Theorem 11.16], it admits an interpretation as modular space of abelian surfaces with additional structures. In particular, there is a universal family  $\mathcal{B}$  of abelian surfaces over  $M^K$ .

The following result holds in the general context of (smooth) Shimura varieties of *PEL*-type.

**Theorem 1.4** [Ancona 2015, Théorème 8.6]. *There is a  $\mathbb{Q}$ -linear tensor functor*

$$\tilde{\mu} : \text{Rep}(G) \rightarrow \text{CHM}^s(M^K)_{\mathbb{Q}}$$

*from the Tannakian category  $\text{Rep}(G)$  of algebraic representations of  $G$  in finite dimensional  $\mathbb{Q}$ -vector spaces to the  $\mathbb{Q}$ -linear category  $\text{CHM}^s(M^K)_{\mathbb{Q}}$  of smooth Chow motives over  $M^K$  (see [Levine 2009, Definition 5.16]). It has the following properties.*

- (a) *The composition of  $\tilde{\mu}$  with the cohomological Hodge theoretic realization is isomorphic to the canonical construction functor  $\mu_{\mathbb{H}}$  (e.g., [Wildeshaus 1997, Theorem 2.2]) to the category of admissible graded-polarizable variations of Hodge structure on  $M_{\mathbb{C}}^K$ .*
- (b) *The composition of  $\tilde{\mu}$  with the cohomological  $\ell$ -adic realization is isomorphic to the canonical construction functor  $\mu_{\ell}$  (e.g., [Wildeshaus 1997, Chapter 4]) to the category of lisse  $\ell$ -adic sheaves on  $M^K$ .*
- (c) *The functor  $\tilde{\mu}$  commutes with Tate twists.*
- (d) *The functor  $\tilde{\mu}$  maps the representation  $V$  to the dual of the Chow motive  $\pi_*^1 \mathbb{1}_B$  over  $M^K$ .*

Here, we denote by  $\pi_*^m \mathbb{1}_B$  the  $m$ -th *Chow-Künneth component* of the Chow motive  $\pi_* \mathbb{1}_B$  over  $M^K$  [Deninger and Murre 1991, Theorem 3.1].

*Proof.* Parts (a), (c) and (d) are identical to [Ancona 2015, Théorème 8.6].

As for part (b), repeat the proof of [loc. cit.], observing that the  $\ell$ -adic analogue of [Ancona 2015, Proposition 8.5] holds (the base change to  $\mathbb{Q}_{\ell}$  of the subgroup  $G_1$  of  $G$  coincides with the Lefschetz group).  $\square$

Given that the representation on  $V$  is faithful, it follows that any object in the image of  $\tilde{\mu}$  is isomorphic to a direct sum of direct factors of Tate twists of the Chow motive  $\pi_{n_i, * \mathbb{1}_B^{n_i}}$  associated to  $B^{n_i}$ , for suitable  $n_i \in \mathbb{N}$ , where  $\pi_{n_i} : B^{n_i} \rightarrow M^K$  denotes the  $n_i$ -fold fibre product of  $B$  over  $M^K$ .

**Definition 1.5.** (a) Denote by  $V_{\alpha} \in \text{Rep}(G)$  the irreducible representation of highest weight  $\alpha$ .

(b) Define  ${}^{\alpha}\mathcal{V} \in \text{CHM}^s(M^K)_{\mathbb{Q}} \subset \text{CHM}(M^K)_{\mathbb{Q}}$  as

$${}^{\alpha}\mathcal{V} := \tilde{\mu}(V_{\alpha}).$$

Given that  $V_{\alpha}$  is of weight  $r$ , the cohomological realizations of  ${}^{\alpha}\mathcal{V}$  equal zero in (classical, i.e., nonperverse) degrees  $\neq r$ , and  $\mu_{\mathbb{H}}(V_{\alpha})$  (in the Hodge theoretic setting) or  $\mu_{\ell}(V_{\alpha})$  (in the  $\ell$ -adic setting) in degree  $r$ .



Denote by  $j : M^K \hookrightarrow (M^K)^*$  the open immersion of  $M^K$  into its Satake–(Baily–Borel) compactification, by  $i : \partial(M^K)^* \hookrightarrow (M^K)^*$  its complement, and by  $\Phi$  the natural stratification of  $\partial(M^K)^*$  (the latter will be made explicit in the beginning of Section 2). Here is our main result.

**Theorem 1.6.** (a) *The motive  $i^* j_*^\alpha \mathcal{V} \in DM_{\mathbb{B},c}(\partial(M^K)^*)$  is a  $\Phi$ -constructible motive of abelian type over  $\partial(M^K)^*$  (see Definition 2.1).*

(b) *The motive  $i^* j_*^\alpha \mathcal{V}$  is without weights*

$$-k + 1, -k + 2, \dots, k,$$

where  $k := \min(k_1 - k_2, k_2)$ . Both weights  $-k$  and  $k + 1$  do occur in  $i^* j_*^\alpha \mathcal{V}$ . In particular,  ${}^\alpha \mathcal{V}$  belongs to the subcategory  $CHM(M^K)_{\mathbb{Q}, \partial w \neq 0,1}$  of  $CHM(M^K)_{\mathbb{Q}}$  if and only if  $\underline{\alpha}$  is regular.

Theorem 1.6 should be compared to [Wildeshaus 2012b, Theorem 3.5], [Wildeshaus 2015, Theorem 3.8], and [Cloître 2017, Theorem 3.6, Proposition 3.8, Proposition 3.9] (see also [Wildeshaus 2018a, Remark 5.8(b)]), which treat the cases of Hilbert–Blumenthal varieties, of Picard surfaces, and of Picard varieties of arbitrary dimension, respectively.

Theorem 1.6 is proved in Section 2. For the rest of the present section, assume that  $k = \min(k_1 - k_2, k_2) \geq 1$ , i.e.,  $k_1 > k_2 > 0$ . Given that according to Theorem 1.6(b), the motive  ${}^\alpha \mathcal{V}$  belongs to  $CHM(M^K)_{\mathbb{Q}, \partial w \neq 0,1}$ , the intersection motive of  $M^K$  relative to  $(M^K)^*$  with coefficients in  ${}^\alpha \mathcal{V}$  is at our disposal: by [Wildeshaus 2018a, Definition 3.7], it equals

$$a_* j_{!*} {}^\alpha \mathcal{V} \in CHM(\mathbb{Q})_{\mathbb{Q}},$$

where  $a : (M^K)^* \rightarrow \text{Spec } \mathbb{Q}$  is the structure morphism of  $(M^K)^*$ . By abuse of language, let us abbreviate, and refer to  $a_* j_{!*} {}^\alpha \mathcal{V}$  as the intersection motive with coefficients in  ${}^\alpha \mathcal{V}$ . Let us list the main corollaries of Theorem 1.6.

**Corollary 1.7.** *Denote by  $a$  and  $\tilde{a}$  the structure morphisms of  $(M^K)^*$  and  $M^K$ , respectively, and by  $m$  the natural transformation  $j_! \rightarrow j_*$ . Assume  $k_1 > k_2 > 0$ , i.e.,  $k \geq 1$ .*

(a) *The motive  $\tilde{a}_! {}^\alpha \mathcal{V} \in DM_{\mathbb{B},c}(\mathbb{Q})$  is without weights  $-k, -k + 1, \dots, -1$ , and the motive  $\tilde{a}_* {}^\alpha \mathcal{V} \in DM_{\mathbb{B},c}(\mathbb{Q})$  is without weights  $1, 2, \dots, k$ . More precisely, the exact triangles*

$$a_* i_{*!} i^* j_{!*} {}^\alpha \mathcal{V}[-1] \rightarrow \tilde{a}_! {}^\alpha \mathcal{V} \rightarrow a_* j_{!*} {}^\alpha \mathcal{V} \rightarrow a_* i_{*!} i^* j_{!*} {}^\alpha \mathcal{V}$$

and

$$a_* j_{!*} {}^\alpha \mathcal{V} \rightarrow \tilde{a}_* {}^\alpha \mathcal{V} \rightarrow a_* i_{*!} i^! j_{!*} {}^\alpha \mathcal{V}[1] \rightarrow a_* j_{!*} {}^\alpha \mathcal{V}[1]$$

are weight filtrations (of  $\tilde{a}_! {}^\alpha \mathcal{V}$ ) avoiding weights  $-k, -k + 1, \dots, -1$ , and (of  $\tilde{a}_* {}^\alpha \mathcal{V}$ ) avoiding weights  $1, 2, \dots, k$ , respectively.

(b) *The intersection motive  $a_* j_{!*}^{\alpha} \mathcal{V} \in \text{CHM}(\mathbb{Q})_{\mathbb{Q}}$  behaves functorially with respect to both  $\tilde{a}_!^{\alpha} \mathcal{V}$  and  $\tilde{a}_*^{\alpha} \mathcal{V}$ . In particular, any endomorphism of  $\tilde{a}_!^{\alpha} \mathcal{V}$  or of  $\tilde{a}_*^{\alpha} \mathcal{V}$  induces an endomorphism of  $a_* j_{!*}^{\alpha} \mathcal{V}$ .*

(c) *Let  $\tilde{a}_!^{\alpha} \mathcal{V} \rightarrow N \rightarrow \tilde{a}_*^{\alpha} \mathcal{V}$  be a factorization of the morphism  $a_* m : \tilde{a}_!^{\alpha} \mathcal{V} \rightarrow \tilde{a}_*^{\alpha} \mathcal{V}$  through a Chow motive  $N \in \text{CHM}(\mathbb{Q})_{\mathbb{Q}}$ . Then the intersection motive  $a_* j_{!*}^{\alpha} \mathcal{V}$  is canonically identified with a direct factor of  $N$ , with a canonical direct complement.*

*Proof.* Given [Theorem 1.6](#), parts (a), (b) and (c) follow from [\[Wildeshaus 2018a, Theorem 3.4\]](#), [\[Wildeshaus 2018a, Theorem 3.5\]](#) and [\[Wildeshaus 2009, Corollary 2.5\]](#), respectively.  $\square$

The equivariance statement from [Corollary 1.7\(b\)](#) applies in particular to endomorphisms coming from the Hecke algebra  $\mathfrak{H}(K, G(\mathbb{A}_f))$  associated to the neat open compact subgroup  $K$  of  $G(\mathbb{A}_f)$ . Recall that by what was said earlier, the relative Chow motive  ${}^{\alpha} \mathcal{V}$  is a direct factor of a Tate twist of  $\pi_{N,*} \mathbb{1}_{\mathbb{B}^N}$ , where  $\pi_N : \mathbb{B}^N \rightarrow M^K$  denotes the  $N$ -fold fibre product of the universal abelian scheme  $\mathbb{B}$  over  $M^K$ .

**Corollary 1.8.** *Assume  $k \geq 1$ . Every element of the Hecke algebra  $\mathfrak{H}(K, G(\mathbb{A}_f))$  acts naturally on the intersection motive  $a_* j_{!*}^{\alpha} \mathcal{V}$ .*

*Proof.* Let  $T \in \mathfrak{H}(K, G(\mathbb{A}_f))$ . According to [Corollary 1.7\(b\)](#), it suffices to show that  $T$  acts on  $\tilde{a}_*^{\alpha} \mathcal{V}$ . To do so, we refer to [\[Wildeshaus 2017, pp. 591–592\]](#).  $\square$

**Corollary 1.9.** *Assume  $k \geq 1$ , and let  $\widetilde{\mathbb{B}}^N$  be any smooth compactification of  $\mathbb{B}^N$ . Then the intersection motive  $a_* j_{!*}^{\alpha} \mathcal{V}$  is a direct factor of a Tate twist of the Chow motive  $b_* \mathbb{1}_{\widetilde{\mathbb{B}}^N}$  ( $b :=$  the structure morphism of the  $\mathbb{Q}$ -scheme  $\widetilde{\mathbb{B}}^N$ ).*

*Proof.* The motive  ${}^{\alpha} \mathcal{V}$  is a direct factor of a Tate twist of  $\pi_{N,*} \mathbb{1}_{\mathbb{B}^N}$ :

$${}^{\alpha} \mathcal{V} \hookrightarrow \pi_{N,*} \mathbb{1}_{\mathbb{B}^N}(\ell)[2\ell] \twoheadrightarrow {}^{\alpha} \mathcal{V},$$

for a suitable integer  $\ell$ . The morphism

$$a_* m : \tilde{a}_! \pi_{N,*} \mathbb{1}_{\mathbb{B}^N} \rightarrow \tilde{a}_* \pi_{N,*} \mathbb{1}_{\mathbb{B}^N}$$

factors through the Chow motive  $b_* \mathbb{1}_{\widetilde{\mathbb{B}}^N}$ , and hence so does

$$a_* m : \tilde{a}_!^{\alpha} \mathcal{V} \rightarrow \tilde{a}_*^{\alpha} \mathcal{V}.$$

Now apply [Corollary 1.7\(c\)](#).  $\square$

**Remark 1.10.** When  $r \geq 0$ , then according to [\[Ancona 2017, Lemma 4.13\]](#), the Chow motive  ${}^{\alpha} \mathcal{V}$  is a direct factor of  $\pi_{N,*} \mathbb{1}_{\mathbb{B}^N}$  (no Tate twist needed). In this context, let us recall [\[Wildeshaus 2018a, Corollary 3.10\]](#): the intersection motive  $a_* j_{!*}^{\alpha} \mathcal{V}$  is canonically dual to the  $e_{\underline{\alpha}}$ -part of the interior motive of  $\mathbb{B}^N$ , where  $e_{\underline{\alpha}}$  is the idempotent endomorphism corresponding to the direct factor  ${}^{\alpha} \mathcal{V}$  of  $\pi_{N,*} \mathbb{1}_{\mathbb{B}^N}$ .

**Corollary 1.11.** *Assume  $k \geq 1$ , i.e., that  $\underline{\alpha}$  is regular. Then for all  $n \in \mathbb{Z}$ , the natural maps*

$$H^n((M^K)^*(\mathbb{C}), j_{!*} \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \rightarrow H^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$$

(in the Hodge theoretic setting) and

$$H^n((M^K)^* \times_{\mathbb{Q}} \overline{\mathbb{Q}}, j_{!*} \mu_{\ell}(V_{\underline{\alpha}})) \rightarrow H^n(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}}))$$

(in the  $\ell$ -adic setting) are injective. Dually,

$$H_c^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \rightarrow H_c^n((M^K)^*(\mathbb{C}), j_{!*} \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$$

and

$$H_c^n(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}})) \rightarrow H_c^n((M^K)^* \times_{\mathbb{Q}} \overline{\mathbb{Q}}, j_{!*} \mu_{\ell}(V_{\underline{\alpha}}))$$

are surjective. In other words, the natural maps from intersection cohomology to cohomology with coefficients in  $\mu_{\mathbf{H}}(V_{\underline{\alpha}})$  and in  $\mu_{\ell}(V_{\underline{\alpha}})$  identify intersection and interior cohomology, respectively.

*Proof.* Write  ${}^{\alpha}\mathcal{V}$  as a direct factor of  $\pi_{N,*} \mathbb{1}_{\mathbb{B}^N}(\ell)[2\ell]$ , for a suitable integer  $\ell$ . Given [Theorem 1.6](#), we may quote [[Wildeshaus 2018a](#), Remark 3.13(a), (b)] (for  $X = (M^K)^*$ ,  $U = M^K$ ,  $C = \mathbb{B}^N$  and  $e = e_{\alpha}$ ).  $\square$

As pointed out in [[Wildeshaus 2018a](#), Remark 3.13(c)], sheaf theoretic considerations alone suffice to show (without any further reference to geometry) that [Theorem 1.6](#) implies [Corollary 1.11](#).

[Corollary 1.11](#) is already known. Indeed, according to [[Mokrane and Tilouine 2002](#), Proposition 1], the result generalizes to Siegel varieties of arbitrary dimension. (However, the proof of [[loc. cit.](#)] is analytic.)

**Remark 1.12.** By [[Wildeshaus 2009](#), Theorem 4.14], control of the reduction of *some* compactification of  $\mathbb{B}^N$  implies control of certain properties of the  $\ell$ -adic realization of the intersection motive  $a_* j_{!*} {}^{\alpha}\mathcal{V}$ . According to [[Faltings and Chai 1990](#), Theorem VI.1.1], there exists a smooth compactification of  $\mathbb{B}^N$  having good reduction at each prime number  $p$  not dividing the level  $n$  of  $K$ .

Theorem 4.14 of [[Wildeshaus 2009](#)] then yields the following:

- (a) for all prime numbers  $p$  not dividing  $n$ , the  $p$ -adic realization of  $a_* j_{!*} {}^{\alpha}\mathcal{V}$  is crystalline;
- (b) if furthermore  $p$  and  $\ell$  are different, then the  $\ell$ -adic realization of  $a_* j_{!*} {}^{\alpha}\mathcal{V}$  is unramified at  $p$ .

**Corollary 1.13.** *Assume  $k \geq 1$ . Let  $p$  be a prime number not dividing the level of  $K$ . Let  $\ell$  be different from  $p$ . Then the characteristic polynomials of the following coincide:*

- (1) the action of Frobenius  $\phi$  on the  $\phi$ -filtered module associated to the (crystalline)  $p$ -adic realization of the intersection motive  $a_* j_{!*} \alpha \mathcal{V}$ ,
- (2) the action of a geometrical Frobenius automorphism at  $p$  on the (unramified)  $\ell$ -adic realization of  $a_* j_{!*} \alpha \mathcal{V}$ .

*Proof.* Fix a smooth compactification  $\widetilde{\mathbf{B}}^N$  of  $\mathbf{B}^N$  with good reduction at  $p$  [Faltings and Chai 1990, Theorem VI.1.1]. Thus the  $\mathbb{Q}_p$ -scheme  $\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \mathbb{Q}_p$  is the generic fibre of a smooth and proper scheme  $\widetilde{\mathcal{B}}^N$  over  $\mathbb{Z}_p$ . Let us denote by  $\widetilde{\mathcal{B}}_{\mathbb{F}_p}^N$  its special fibre.

The  $\phi$ -filtered module associated to  $p$ -adic étale cohomology of  $\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \overline{\mathbb{Q}}$  is first isomorphic to Hyodo–Kato cohomology  $H_{HK}^{\bullet}(\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \overline{\mathbb{Q}}_p)$  [Beilinson 2013, Section 3.2], and this isomorphism can be chosen to be motivic in the sense that it commutes with the action of correspondences in  $\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \widetilde{\mathbf{B}}^N$  [Déglise and Niziol 2018, Section 4.15]. By definition, Hyodo–Kato cohomology is log-crystalline cohomology of a log-smooth model; in our case, given good reduction, such a model is given by  $\widetilde{\mathcal{B}}^N$  (with divisor equal to zero). In other words, Hyodo–Kato cohomology equals crystalline cohomology of  $\widetilde{\mathcal{B}}^N$ . This identification commutes with the action of correspondences in  $\widetilde{\mathcal{B}}^N \times_{\mathbb{Z}_p} \widetilde{\mathcal{B}}^N$ . Finally, crystalline cohomology of  $\widetilde{\mathcal{B}}^N$  equals crystalline cohomology of  $\widetilde{\mathcal{B}}_{\mathbb{F}_p}^N$ .

Altogether, the  $\phi$ -filtered module associated to  $p$ -adic étale cohomology of  $\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \overline{\mathbb{Q}}$  is identified with crystalline cohomology of  $\widetilde{\mathcal{B}}_{\mathbb{F}_p}^N$  in a way compatible with the action of correspondences in  $\widetilde{\mathcal{B}}^N \times_{\mathbb{Z}_p} \widetilde{\mathcal{B}}^N$ . Concretely, this means that given a correspondence  $e$  in  $\widetilde{\mathcal{B}}^N \times_{\mathbb{Z}_p} \widetilde{\mathcal{B}}^N$ , the action of its generic fibre  $e_{\mathbb{Q}_p}$  on  $p$ -adic étale cohomology is identified with the action of its special fibre  $e_{\mathbb{F}_p}$  on crystalline cohomology.

For  $\ell \neq p$ , smooth and proper base change allows us to identify  $\ell$ -adic cohomology of  $\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \overline{\mathbb{Q}}$  and  $\ell$ -adic cohomology of  $\widetilde{\mathcal{B}}_{\mathbb{F}_p}^N \times_{\mathbb{F}_p} \overline{\mathbb{F}}_p$ , again compatibly with correspondences.

According to Corollary 1.9, there is an idempotent endomorphism  $e_{\mathbb{Q}}$  of the Chow motive associated to  $\widetilde{\mathbf{B}}^N$ , or in other words, an idempotent correspondence in  $\widetilde{\mathbf{B}}^N \times_{\mathbb{Q}} \widetilde{\mathbf{B}}^N$ , whose images in the endomorphism rings of the realizations are projections onto the realizations of  $a_* j_{!*} \alpha \mathcal{V}$ . We claim that  $e_{\mathbb{Q}_p} := e_{\mathbb{Q}} \times_{\mathbb{Q}} \mathbb{Q}_p$  can be extended idempotently to  $\widetilde{\mathcal{B}}^N \times_{\mathbb{Z}_p} \widetilde{\mathcal{B}}^N$ . Indeed, according to [O’Sullivan 2011, Proposition 5.1.1], the restriction morphism from the endomorphism ring of the Chow motive associated to  $\widetilde{\mathcal{B}}^N$  to that of the Chow motive associated to  $\widetilde{\mathbf{B}}^N$  is epimorphic, with nilpotent kernel. We now follow a standard line of argument (cf. [Kimura 2005, proof of Corollary 7.8]): let  $\mathfrak{e}$  be any extension of  $e_{\mathbb{Q}_p}$  to  $\widetilde{\mathcal{B}}^N \times_{\mathbb{Z}_p} \widetilde{\mathcal{B}}^N$ . The difference  $\mathfrak{e} - \mathfrak{e}^2$  is nilpotent, say

$$(\mathfrak{e} - \mathfrak{e}^2)^N = 0.$$

But then,

$$e_{\mathbb{Z}_p} := (\text{id}_{\widetilde{\mathcal{B}}^N} - (\text{id}_{\widetilde{\mathcal{B}}^N} - \epsilon)^N)^N$$

equally extends  $e_{\mathbb{Q}_p}$  to  $\widetilde{\mathcal{B}}^N \times_{\mathbb{Z}_p} \widetilde{\mathcal{B}}^N$ , and  $e_{\mathbb{Z}_p}$  is idempotent.

Altogether, there is a smooth and proper scheme  $\widetilde{\mathcal{B}}_{\mathbb{F}_p}^N$  over  $\mathbb{F}_p$ , and an idempotent endomorphism  $e_{\mathbb{F}_p}$  of the Chow motive associated to  $\widetilde{\mathcal{B}}_{\mathbb{F}_p}^N$ , whose images in the endomorphism rings of crystalline and  $\ell$ -adic cohomology, respectively, are projections onto the realizations of  $a_* j_{!*} \alpha \mathcal{V}$ . The claim thus follows from [Katz and Messing 1974, Theorem 2(2)]. □

### 2. Proof of the main result

We keep the notation of the preceding section. In order to prove Theorem 1.6, the idea is to apply the criterion from [Wildeshaus 2018a, Corollary 4.6].

In order to check the hypotheses of [loc. cit.], we first need to fix a finite stratification  $\Phi$  of  $\partial(M^K)^*$  by locally closed subschemes. The canonical choice would be the restriction  $\Phi'$  to  $\partial(M^K)^*$  of the natural (finite) stratification of  $(M^K)^*$  from [Pink 1989, Main Theorem 12.3(c)] — in other words, all the strata of  $(M^K)^*$  except the open one, i.e., except  $M^K$ . According to [Wildeshaus 2017, Lemma 8.2(a)],  $\Phi'$  is *good*, meaning that the closure of every stratum is a union of strata. Furthermore, by [Wildeshaus 2017, Lemma 8.2(b)], all strata, denoted  $i_g(M^{\pi_1(K_1)})$ , are smooth over  $\mathbb{Q}$  (recall that  $K$  is assumed neat, and that  $(G, \mathcal{H})$  satisfies condition (+)), hence regular. The same is therefore true for the following coarser stratification  $\Phi = \{0, 1\}$  of  $\partial(M^K)^*$ : denote by  $i_0 : Z_0 \hookrightarrow \partial(M^K)^*$  the disjoint union of all closed strata of  $\Phi'$ , and by  $i_1 : Z_1 \hookrightarrow \partial(M^K)^*$  the disjoint union of all strata of  $\Phi'$ , which are open in  $\partial(M^K)^*$ . Indeed, according to [Pink 1989, Section 6.3, Example 4.25 (with  $g = 2$ )],

$$\partial(M^K)^* = Z_0 \amalg Z_1;$$

more precisely,  $Z_0$  is of dimension zero, and  $Z_1$  of dimension one (hence so is the whole of  $\partial(M^K)^*$ ). Let us refer to  $Z_0$  as the *Siegel stratum*, and to  $Z_1$  as the *Klingen stratum* of  $\partial(M^K)^*$ . When it is necessary to insist on the structure of stratified scheme of  $\partial(M^K)^*$ , we write  $\partial(M^K)^*(\Phi)$  instead of  $\partial(M^K)^*$ .

**Definition 2.1** [Wildeshaus 2018b, Definitions 3.4 and 3.5]. (a) Let  $S(\mathfrak{S}) = \coprod_{\sigma \in \mathfrak{S}} S_\sigma$  be a good stratification of a scheme  $S(\mathfrak{S})$ . A morphism  $\pi : S(\mathfrak{S}) \rightarrow \partial(M^K)^*(\Phi)$  is said to be a *morphism of good stratifications* if the preimage of any of the strata  $Z_0, Z_1$  of  $\partial(M^K)^*$  is a union of strata  $S_\sigma$ .

(b) A morphism  $\pi : S(\mathfrak{S}) \rightarrow \partial(M^K)^*(\Phi)$  of good stratifications is said to be of *abelian type* if it is proper, and if the following conditions are satisfied.

- (1) All strata  $S_\sigma$ ,  $\sigma \in \mathfrak{S}$ , are nilregular, and for any immersion  $i_\tau : S_\tau \hookrightarrow \overline{S_\sigma}$  of a stratum  $S_\tau$  into the closure  $\overline{S_\sigma}$  of a stratum  $S_\sigma$ , the functor  $i_\tau^!$  maps  $\mathbb{1}_{\overline{S_\sigma}}$  to a Tate motive over  $S_\tau$  [Levine 2010, Section 3.3].
- (2) For all  $\sigma \in \mathfrak{S}$  such that  $S_\sigma$  is a stratum of  $\pi^{-1}(Z_m)$ ,  $m \in \{0, 1\}$ , the morphism  $\pi_\sigma : S_\sigma \rightarrow Z_m$  can be factorized,

$$\pi_\sigma = \pi'_\sigma \circ \pi''_\sigma : S_\sigma \xrightarrow{\pi''_\sigma} B_\sigma \xrightarrow{\pi'_\sigma} Z_m,$$

such that the motive

$$\pi''_{\sigma,*} \mathbb{1}_{S_\sigma} \in DM_{B,c}(B_\sigma)_F$$

belongs to the category  $DMT(B_\sigma)_F$  of Tate motives over  $B_\sigma$ , the morphism  $\pi'_\sigma$  is proper and smooth, and its pull-back to any geometric point of  $Z_m$  lying over a generic point is isomorphic to a finite disjoint union of abelian varieties.

- (c) An object  $V \in DM_{B,c}(\partial(M^K)^*)$  is a  $\Phi$ -constructible motive of abelian type over  $\partial(M^K)^*$  if the following holds: the motive  $V$  belongs to the strict, full, dense,  $\mathbb{Q}$ -linear triangulated subcategory  $DM_{B,c,\Phi}^{Ab}(\partial(M^K)^*)$  generated by the images under  $\pi_*$  of  $\mathfrak{S}$ -constructible Tate motives over  $S(\mathfrak{S})$  [Wildeshaus 2018b, Definition 3.3], where

$$\pi : S(\mathfrak{S}) \rightarrow \partial(M^K)^*(\Phi)$$

runs through the morphisms of abelian type with target equal to  $\partial(M^K)^*(\Phi)$ .

**Theorem 2.2.** Let  $\underline{\alpha} = \alpha(k_1, k_2, r)$ , with  $(k_1, k_2, r) \in \mathbb{Z}^3$  such that

$$r \equiv k_1 + k_2 \pmod{2} \quad \text{and} \quad k_1 \geq k_2 \geq 0,$$

and consider  ${}^\alpha \mathcal{V} = \tilde{\mu}(V_\alpha) \in CHM(M^K)_\mathbb{Q}$ . The motive  $i^* j_* {}^\alpha \mathcal{V}$  belongs to the full subcategory  $DM_{B,c,\Phi}^{Ab}(\partial(M^K)^*)$  of  $DM_{B,c}(\partial(M^K)^*)$ . In other words, it is a  $\Phi$ -constructible motive of abelian type over  $\partial(M^K)^*$ .

*Proof.* As recalled earlier, the relative Chow motive  ${}^\alpha \mathcal{V}$  belongs to the strict, full, dense,  $\mathbb{Q}$ -linear triangulated subcategory

$$\pi_{N,*} DMT(B^N)_\mathbb{Q}^\natural$$

of  $DM_{B,c}(M^K)$  generated by the images under  $\pi_{N,*}$  of the category of Tate motives over  $B^N$ . Here as before,  $\pi_N : B^N \rightarrow M^K$  denotes the  $N$ -fold fibre product of the universal abelian scheme  $B$  over  $M^K$ .

The latter equals the projection from a mixed Shimura variety: indeed [Pink 1989, Example 2.7], the representation  $V$  of  $G$  is of Hodge type  $\{(-1, 0), (0, -1)\}$ . The same is then true for the  $r$ -th power  $V^N$  of  $V$ . By [Pink 1989, Proposition 2.17], this allows for the construction of the unipotent extension  $(P^N, \mathfrak{X}^N)$  of  $(G, \mathcal{H})$

by  $V^N$ . The pair  $(P^N, \mathfrak{X}^N)$  constitute *mixed Shimura data* [Pink 1989, Definition 2.1]. By construction, they come endowed with a morphism of Shimura data  $\pi_N : (P^N, \mathfrak{X}^N) \rightarrow (G, \mathcal{H})$ , identifying  $(G, \mathcal{H})$  with the pure Shimura data underlying  $(P^N, \mathfrak{X}^N)$ . In particular,  $(P^N, \mathfrak{X}^N)$  also satisfies condition (+). Now by [Pink 1989, Theorem 11.18] there is an open compact neat subgroup  $K_N$  of  $P^N(\mathbb{A}_f)$ , whose image under  $\pi_N$  equals  $K$ , such that  $B^N$  is identified with the *mixed Shimura variety*  $M^{K_N} := M^{K_N}(P^N, \mathfrak{X}^N)$ , and such that the morphism  $M^{K_N} \rightarrow M^K$  induced by the morphism  $\pi_N$  of Shimura data is identified with the structure morphism of  $B^N$ .

Choose a smooth *toroidal compactification*  $M^{K_N}(\mathfrak{S}) := M^{K_N}(P^N, \mathfrak{X}^N, \mathfrak{S})$  of  $M^{K_N}$ , associated to a  $K_N$ -admissible complete cone decomposition  $\mathfrak{S}$  [Pink 1989, Section 6.4]. Then by [Pink 1989, proof of Theorem 9.21], modulo a suitable refinement of  $\mathfrak{S}$ , the natural stratification of  $M^{K_N}(\mathfrak{S})$ , also denoted  $\mathfrak{S}$ , satisfies the conclusions of [Wildeshaus 2017, Lemma 8.1], i.e., it is good, and the closures of all strata are regular. Note that the unique open stratum equals  $M^{K_N}$ . According to [Pink 1989, Section 6.24, Main Theorem 12.4(b)], the morphism  $\pi_N : B^N = M^{K_N} \rightarrow M^K$  extends to a proper, surjective morphism  $M^{K_N}(\mathfrak{S}) \rightarrow (M^K)^*$ , still denoted  $\pi_N$ . From the description given in [Pink 1989, Section 7.3], one sees that  $\pi_N$  is a morphism of stratifications.

According to [Wildeshaus 2017, Corollary 4.10(b), Remark 4.7], the category

$$\pi_{N,*}DMT_{\mathfrak{S}}(M^{K_N}(\mathfrak{S}))_{\mathbb{Q}}^{\natural}$$

is obtained by gluing  $\pi_{N,*}DMT(B^N)_{\mathbb{Q}}^{\natural}$  and  $\pi_{N,*}DMT_{\mathfrak{S}}(\pi_N^{-1}(\partial(M^K)^*))_{\mathbb{Q}}^{\natural}$ . In particular,

$$i^*j_*^{\alpha}\mathcal{V} \in \pi_{N,*}DMT_{\mathfrak{S}}(\pi_N^{-1}(\partial(M^K)^*))_{\mathbb{Q}}^{\natural}.$$

But  $\pi_N$  is of abelian type [Wildeshaus 2017, Lemma 8.4]; therefore,

$$\pi_{N,*}DMT_{\mathfrak{S}}(\pi_N^{-1}(\partial(M^K)^*))_{\mathbb{Q}}^{\natural} \subset DM_{B,c,\Phi}^{Ab}(\partial(M^K)^*). \quad \square$$

Next, we collect information on the restriction of  $i^*j_*R_{\ell,M^K}(\alpha)\mathcal{V}$  to the strata  $Z_0$  and  $Z_1$ . The following is essentially due to Lemma [2015, Section 4].

**Theorem 2.3.** *Let  $\ell$  be a prime number.*

(a) *For all integers  $n \leq r + 2$ , the perverse cohomology sheaf*

$$H^n i_0^* i^* j_* R_{\ell,M^K}(\alpha)\mathcal{V}$$

*on  $Z_0$  is of weights  $\leq n - (k_1 - k_2)$ . The perverse cohomology sheaf*

$$H^{r+2} i_0^* i^* j_* R_{\ell,M^K}(\alpha)\mathcal{V}$$

*is nonzero, and pure of weight  $(r + 2) - (k_1 - k_2)$ .*

(b) For all integers  $n \leq r + 2$ , the perverse cohomology sheaf

$$H^n i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V})$$

on  $Z_1$  is of weights  $\leq n - k_2$ . The perverse cohomology sheaf

$$H^{r+2} i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V})$$

is nonzero, and pure of weight  $(r + 2) - k_2$ .

The proof of [Theorem 2.3](#) is given after [Remark 2.6](#). In order to prepare it, recall from [[Pink 1989](#), Example 4.25] that  $Z_0$  and  $Z_1$  correspond bijectively to the  $G(\mathbb{Q})$ -conjugacy classes of proper *rational boundary components* [[Pink 1989](#), Section 4.11] of  $(G, \mathcal{H})$ . Indeed, the group  $G(\mathbb{Q})$  acts transitively on the set of totally isotropic subspaces of  $V$  of a given, strictly positive dimension.

We already fixed a basis  $(e_1, e_2, e_3, e_4)$  of  $V$ , in which our symplectic bilinear form  $J$  acquires the  $4 \times 4$ -matrix

$$\begin{pmatrix} 0 & I_2 \\ -I_2 & 0 \end{pmatrix},$$

which we equally denoted by  $J$ . The subspaces  $V'_0$  and  $V'_1$  generated by  $\{e_1, e_2\}$  and  $\{e_3\}$ , respectively, are both totally isotropic.

Following [[Pink 1989](#), Example 4.25], we put  $Q_m := \text{Stab}_G(V'_m)$ ,  $m = 0, 1$ . Let  $P_m$  denote the normal subgroup of  $Q_m$  underlying the rational boundary component  $(P_m, \mathfrak{X}_m)$  giving rise to  $Z_m$  [[Pink 1989](#), Section 4.7], and  $W_m$  its unipotent radical (which equals the unipotent radical of  $Q_m$ ). Then, still according to [[Pink 1989](#), Example 4.25],

$$\begin{aligned} Q_0 &= \left\{ \begin{pmatrix} q \cdot A & A \cdot M \\ 0 & {}^t A^{-1} \end{pmatrix} : q \in \mathbb{G}_{m, \mathbb{Q}}, A \in \text{GL}_{2, \mathbb{Q}}, {}^t M = M \right\}, \\ P_0 &= \left\{ \begin{pmatrix} q \cdot I_2 & M \\ 0 & I_2 \end{pmatrix} : q \in \mathbb{G}_{m, \mathbb{Q}}, {}^t M = M \right\}, \\ W_0 &= \left\{ \begin{pmatrix} I_2 & M \\ 0 & I_2 \end{pmatrix} : {}^t M = M \right\}, \end{aligned}$$

while

$$Q_1 = \left\{ \begin{pmatrix} a & aq^{-1}(bu+dw) & v & aq^{-1}(cu+ew) \\ 0 & b & w & c \\ 0 & 0 & a^{-1}q & 0 \\ 0 & d & -u & e \end{pmatrix} : a, be - cd = q \in \mathbb{G}_{m, \mathbb{Q}} \right\},$$



$$P_1 = \left\{ \left( \begin{array}{cccc} be - cd & bu + dw & v & cu + ew \\ 0 & b & w & c \\ 0 & 0 & 1 & 0 \\ 0 & d & -u & e \end{array} \right) : be - cd \in \mathbb{G}_{m, \mathbb{Q}} \right\},$$

$$W_1 = \left\{ \left( \begin{array}{cccc} 1 & u & v & w \\ 0 & 1 & w & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -u & 1 \end{array} \right) \right\}.$$

Observe that  $Q_0 \cap Q_1$  equals the Borel subgroup of  $G$  stabilizing the flag  $V'_1 \subset V'_0$ , and that both  $Q_0$  and  $Q_1$  contain the fixed maximal split torus

$$T = \{\text{diag}(a, b, a^{-1}q, b^{-1}q) : a, b, q \in \mathbb{G}_{m, \mathbb{Q}}\}.$$

In particular,  $T$  is canonically identified with a maximal  $\mathbb{Q}$ -split torus of the reductive group  $Q_m/W_m$ , for  $m = 0, 1$ . Given a (representation-theoretic) weight  $\alpha : T \rightarrow \mathbb{G}_{m, \mathbb{Q}}$ , let us denote by  $\alpha_m$  the same application, but with  $T$  seen as a subgroup of  $Q_m/W_m$ ,  $m = 0, 1$ .

Note that

$$R_{\ell, M^K}(\alpha \mathcal{V}) = \mu_{\ell}(V_{\underline{\alpha}})[-r].$$

Recall that we denote by  $\Phi'$  the natural (finite) stratification of  $(M^K)^*$  from [Pink 1989, Main Theorem 12.3(c)], which is finer than  $\Phi$ . In order to determine the classical cohomology objects  $R^n i_m^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}})$ , for  $m = 0, 1$ , and  $n \in \mathbb{Z}$ , one applies the following standard strategy.

(1) By Pink’s theorem [1992, Theorem (5.3.1)], the restriction of  $R^n i_m^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}})$  to any individual stratum  $Z'$  of  $\Phi'$  contributing to  $Z_m$  equals

$$R^n i_m^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}})|_{Z'} = \bigoplus_{p+q=n} \mu_{\ell, Z'}(H^p(H_C/K_W, H^q(\text{Lie}(W_m), V_{\underline{\alpha}}))).$$

Here,  $H_C/K_W$  is an arithmetic subgroup (depending on  $Z'$ ) of  $C_m/W_m$  [Pink 1992, Section (5.2)], where  $C_m$  is the identity component of the Zariski closure of the centralizer in  $Q_m(\mathbb{Q})$  of the rational boundary component  $(P_m, \mathfrak{X}_m)$  [Pink 1992, Section (3.7)], and  $\mu_{\ell, Z'}$  is the canonical construction functor to the category of lisse  $\ell$ -adic sheaves on  $Z'$ .

(2) Apply Kostant’s theorem [Vogan 1981, Theorem 3.2.3], in order to identify  $H^q(\text{Lie}(W_m), V_{\underline{\alpha}})$  as a representation of the reductive group  $Q_m/W_m$ ; this allows us in particular to obtain its weights, and gives potential information concerning cohomology of  $H_C/K_W$  with coefficients in  $H^q(\text{Lie}(W_m), V_{\underline{\alpha}})$ .

The Hodge theoretic analogue of the above strategy yields the cohomology objects of  $i_m^* i^* j_* \mu_{\ell, \mathbf{H}}(V_{\underline{\alpha}})|_{Z'}$ ; this was made explicit in [Lemma 2015, Section 4].

Since steps (2) of the  $\ell$ -adic and the Hodge theoretic strategies are identical, we may use the computations from [loc. cit.] in our setting.

**Proposition 2.4** [Lemma 2015, Section 4.3]. *Let  $\underline{\alpha} = \alpha(k_1, k_2, r)$  with  $(k_1, k_2, r) \in \mathbb{Z}^3$  such that*

$$r \equiv k_1 + k_2 \pmod{2} \quad \text{and} \quad k_1 \geq k_2 \geq 0.$$

(a) *For  $m = 0, 1$ , we have*

$$H^q(\text{Lie}(W_m), V_{\underline{\alpha}}) = 0$$

*whenever  $q < 0$  or  $q > 3$ . If  $0 \leq q \leq 3$ , the  $Q_m/W_m$ -representation  $H^q(\text{Lie}(W_1), V_{\underline{\alpha}})$  is (nonzero and) irreducible.*

(b) *The highest (representation-theoretic) weight of  $H^q(\text{Lie}(W_0), V_{\underline{\alpha}})$ ,  $0 \leq q \leq 3$ , is*

$$\begin{aligned} &\alpha_0(k_1, k_2, r) \quad \text{for } q = 0, \\ &\alpha_0(k_1, -k_2 - 2, r) \quad \text{for } q = 1, \\ &\alpha_0(k_2 - 1, -k_1 - 3, r) \quad \text{for } q = 2, \\ &\alpha_0(-k_2 - 3, -k_1 - 3, r) \quad \text{for } q = 3. \end{aligned}$$

(c) *The highest (representation-theoretic) weight of  $H^q(\text{Lie}(W_1), V_{\underline{\alpha}})$ ,  $0 \leq q \leq 3$ , is*

$$\begin{aligned} &\alpha_1(k_1, k_2, r) \quad \text{for } q = 0, \\ &\alpha_1(k_2 - 1, k_1 + 1, r) \quad \text{for } q = 1, \\ &\alpha_1(-k_2 - 3, k_1 + 1, r) \quad \text{for } q = 2, \\ &\alpha_1(-k_1 - 4, k_2, r) \quad \text{for } q = 3. \end{aligned}$$

*Proof.* Note that given our normalization, we have

$$\alpha(k_1, k_2, r) = \lambda(k_1, k_2, -r)$$

in the notation of [Lemma 2015, top of p. 87].

Part (a) follows from Kostant’s theorem, and from the following fact (see [Lemma 2015, proofs of Lemmas 4.8 and 4.10]), valid for both  $m = 0$  and  $m = 1$ : the set of Weyl representatives for  $Q_m$  contains no element of length  $< 0$  or  $> 3$ , and exactly one element of respective lengths 0, 1, 2 and 3.

As for part (c), we refer to [Lemma 2015, proof of Lemma 4.10].

[Lemma 2015, proof of Lemma 4.8] contains the complete setting for the application of Kostant’s theorem for  $m = 0$ , but makes it explicit only for  $H^2(\text{Lie}(W_0), V_{\underline{\alpha}})$  and  $H^3(\text{Lie}(W_0), V_{\underline{\alpha}})$ . The reader will have no difficulty filling in the missing information needed for part (b).  $\square$

Note that both  $Q_0/W_0$  and  $Q_1/W_1$  are isomorphic to  $\mathbb{G}_{m, \mathbb{Q}} \times_{\mathbb{Q}} \text{GL}_{2, \mathbb{Q}}$ . More precisely,

$$Q_0/W_0 = P_0/W_0 \times_{\mathbb{Q}} \mathrm{GL}_{2,\mathbb{Q}} = \mathbb{G}_{m,\mathbb{Q}} \times_{\mathbb{Q}} \mathrm{GL}_{2,\mathbb{Q}},$$

the identification given by sending the class of a matrix

$$\begin{pmatrix} q \cdot A & A \cdot M \\ 0 & {}^t A^{-1} \end{pmatrix}$$

to the pair  $(q, A)$ , and

$$Q_1/W_1 = P_1/W_1 \times_{\mathbb{Q}} \mathbb{G}_{m,\mathbb{Q}} = \mathrm{GL}_{2,\mathbb{Q}} \times_{\mathbb{Q}} \mathbb{G}_{m,\mathbb{Q}},$$

the identification given by sending the class of a matrix

$$\begin{pmatrix} a & aq^{-1}(bu + dw) & v & aq^{-1}(cu + ew) \\ 0 & b & w & c \\ 0 & 0 & a^{-1}q & 0 \\ 0 & d & -u & e \end{pmatrix}$$

to the pair

$$\left( \begin{pmatrix} b & c \\ d & e \end{pmatrix}, aq^{-1} \right).$$

The restriction of the inverse identification to maximal split tori sends

$$\left( q, \begin{pmatrix} x & 0 \\ 0 & x^{-1}y \end{pmatrix} \right) \in P_0/W_0 \times_{\mathbb{Q}} \mathrm{GL}_{2,\mathbb{Q}}$$

to

$$\mathrm{diag}(qx, qx^{-1}y, x^{-1}, xy^{-1}) \in T \subset Q_0/W_0$$

for  $m = 0$ , and

$$\left( \begin{pmatrix} x & 0 \\ 0 & x^{-1}q \end{pmatrix}, y \right) \in P_1/W_1 \times_{\mathbb{Q}} \mathbb{G}_{m,\mathbb{Q}}$$

to

$$\mathrm{diag}(yq, x, y^{-1}, x^{-1}q) \in T \subset Q_1/W_1$$

for  $m = 1$ .

In the following, the reader should be particularly careful not to confuse two notions of *weight* associated to representations of reductive groups: the highest weights in the sense of representation theory (e.g., those occurring in Kostant’s theorem), when the representation is irreducible, and the weights as determined by the action of the weight cocharacter [Pink 1989, Section 1.3], when the group underlies Shimura data.

**Corollary 2.5.** (a) *The  $Q_0/W_0$ -representations  $H^q(\mathrm{Lie}(W_0), V_{\underline{\alpha}})$ ,  $0 \leq q \leq 2$ , are (irreducible and) regular, except when  $q = 0$  and  $k_1 = k_2$ , in which case  $H^0(\mathrm{Lie}(W_0), V_{\underline{\alpha}})$  factors through the quotient  $\mathbb{G}_{m,\mathbb{Q}} \times_{\mathbb{Q}} \mathbb{G}_{m,\mathbb{Q}}$  of the group*

$$Q_0/W_0 = \mathbb{G}_{m,\mathbb{Q}} \times_{\mathbb{Q}} \mathrm{GL}_{2,\mathbb{Q}}$$

via the determinant on the factor  $\mathrm{GL}_{2,\mathbb{Q}}$ . The restriction to  $\mathrm{SL}_{2,\mathbb{Q}} \subset \mathrm{GL}_{2,\mathbb{Q}}$  of  $H^1(\mathrm{Lie}(W_0), V_\alpha)$  is of highest (representation-theoretic) weight  $k_1 + k_2 + 2$ . The restriction to  $P_0/W_0$  of  $H^0(\mathrm{Lie}(W_0), V_\alpha)$  is of weight  $(r+1) - (k_1 + k_2) - 1$ , and the restriction of  $H^1(\mathrm{Lie}(W_0), V_\alpha)$  is of weight  $(r+2) - (k_1 - k_2)$ .

(b) The restriction to  $P_1/W_1$  of  $H^0(\mathrm{Lie}(W_1), V_\alpha)$  is of weight  $(r+1) - k_1 - 1$ , and the restriction of  $H^1(\mathrm{Lie}(W_1), V_\alpha)$  is of weight  $(r+2) - k_2 - 1$ .

*Proof.* (a): Given the above identifications, the weight  $\alpha_0(n_1, n_2, r)$  on  $T$  maps

$$\left( q, \begin{pmatrix} x & 0 \\ 0 & x^{-1}y \end{pmatrix} \right) \in P_0/W_0 \times_{\mathbb{Q}} \mathrm{GL}_{2,\mathbb{Q}}$$

to

$$\alpha_0(n_1, n_2, r)(\mathrm{diag}(qx, qx^{-1}y, x^{-1}, xy^{-1})) = x^{n_1-n_2}y^{n_2}q^{-(r-n_1-n_2)/2}.$$

In particular, the restriction of  $\alpha_0(n_1, n_2, r)$  to  $T \cap \mathrm{SL}_{2,\mathbb{Q}}$  corresponds to the integer  $n_1 - n_2$ . The first and the second claim thus follow from [Proposition 2.4\(b\)](#).

The weight cocharacter  $\mathbb{G}_{m,\mathbb{Q}} \rightarrow P_0/W_0 = \mathbb{G}_{m,\mathbb{Q}}$  maps  $z$  to  $z^2$  [[Pink 1989](#), Examples 4.25 and 2.8]. Its composition with the inclusion into  $T$ , and with  $\alpha_0(n_1, n_2, r)$  yields

$$\mathbb{G}_{m,\mathbb{Q}} \rightarrow \mathbb{G}_{m,\mathbb{Q}}, \quad z \mapsto z^{-r+n_1+n_2}.$$

The third claim thus follows from [Proposition 2.4\(b\)](#), and from the normalization of weights of representations [[Pink 1989](#), Section 1.3].

(b): The weight cocharacter  $\mathbb{G}_{m,\mathbb{Q}} \rightarrow P_1/W_1 = \mathrm{GL}_{2,\mathbb{Q}}$  maps  $z$  to

$$\begin{pmatrix} z & 0 \\ 0 & z \end{pmatrix}$$

[[Pink 1989](#), Examples 4.25 and 2.8]. Given the above identifications, its composition with the inclusion into  $T$  maps  $z$  to  $\mathrm{diag}(z^2, z, 1, z)$ . Further composition with  $\alpha_1(n_1, n_2, r)$  then yields

$$\mathbb{G}_{m,\mathbb{Q}} \rightarrow \mathbb{G}_{m,\mathbb{Q}}, \quad z \mapsto z^{-r+n_1}.$$

The claim thus follows from [Proposition 2.4\(c\)](#). □

To complete the ingredients needed for the computation of the  $R^n i_m^* i^* j_* \mu_\ell(V_\alpha)$  according to the strategy (1), (2) sketched earlier in this section, observe that the group  $H_C/K_W$  associated to an individual stratum  $Z'$  of  $\Phi'$  contributing to  $Z_m$  is a neat arithmetic subgroup of  $\mathrm{GL}_2(\mathbb{Q})$  for  $m=0$  [[Lemma 2015](#), proof of Lemma 4.8], and hence of  $\mathrm{SL}_2(\mathbb{Q})$ . In particular, it is of cohomological dimension one. For  $m=1$ , the group  $H_C/K_W$ , being a neat arithmetic subgroup of  $\mathbb{G}_m(\mathbb{Q})$ , is trivial [[Lemma 2015](#), proof of Lemma 4.10].

**Remark 2.6.** When  $m = 0$ , let  $V_2$  denote the standard representation of  $\mathrm{SL}_{2,\mathbb{Q}}$ , and  $u \in \mathbb{N}$ . Then  $\mathrm{Sym}^u V_2 \in \mathrm{Rep}(\mathrm{SL}_{2,\mathbb{Q}})$ ; in fact,  $\mathrm{Sym}^u V_2$  is the irreducible representation of highest (representation-theoretic) weight  $u$ . Denote by  $g$  the genus of the quotient of the upper half space by  $H_C/K_W$ , and by  $c \geq 1$  the number of its cusps. (Thus,  $c \geq 3$  if  $g = 0$  since  $H_C/K_W$  is neat.) Then  $H^1(H_C/K_W, \mathrm{Sym}^u V_2)$  is of dimension  $(u + 1)(2g - 2 + c)$  if  $u \geq 1$ , and of dimension  $2g - 1 + c$  if  $u = 0$ . In particular,

$$H^1(H_C/K_W, \mathrm{Sym}^u V_2) \neq 0 \quad \text{for all } u \in \mathbb{N}.$$

*Proof of Theorem 2.3.* (a): According to [Corollary 2.5\(a\)](#) and [Proposition 2.4\(a\)](#),

$$(o) \quad 0 \neq H^0(\mathrm{Lie}(W_0), V_\alpha) \text{ is of weight } (r + 1) - (k_1 + k_2) - 1,$$

$$(i) \quad 0 \neq H^1(\mathrm{Lie}(W_0), V_\alpha) \text{ is of weight } (r + 2) - (k_1 - k_2),$$

and  $H^q(\mathrm{Lie}(W_0), V_\alpha) = 0$  whenever  $q < 0$ . The group  $H_C/K_W$  associated to a stratum  $Z'$  of  $Z_0$  is a neat arithmetic subgroup of  $\mathrm{SL}_2(\mathbb{Q})$ . It is therefore of cohomological dimension one, and admits no nonzero invariants on regular irreducible representations of  $Q_0/W_0 = \mathbb{G}_{m,\mathbb{Q}} \times_{\mathbb{Q}} \mathrm{GL}_{2,\mathbb{Q}}$ .

By [Proposition 2.4\(a\)](#) and [Corollary 2.5\(a\)](#),  $H^q(\mathrm{Lie}(W_0), V_\alpha)$ ,  $0 \leq q \leq 2$ , are irreducible as representations of  $Q_0/W_0$ ; it is regular unless  $q = 0$  and  $k_1 = k_2$ , in which case  $\mathrm{SL}_{2,\mathbb{Q}}$ , so  $H_C/K_W$  acts trivially. Pink's theorem and [[Pink 1992](#), Proposition (5.5.4)] then tell us that

$$(o) \quad R^0 i_0^* i^* j_* \mu_\ell(V_\alpha) \text{ is nonzero if and only if } k_1 = k_2, \text{ in which case it is of weight } r - (k_1 + k_2),$$

$$(i) \quad 0 \neq R^1 i_0^* i^* j_* \mu_\ell(V_\alpha) \text{ is of weight } (r + 1) - (k_1 + k_2) - 1,$$

$$(ii) \quad 0 \neq R^2 i_0^* i^* j_* \mu_\ell(V_\alpha) \text{ is of weight } (r + 2) - (k_1 - k_2),$$

and that  $R^n i_0^* i^* j_* \mu_\ell(V_\alpha) = 0$  whenever  $n < 0$  (for the nonvanishing statements in (i), (ii), see [Remark 2.6](#)).

The scheme  $Z_0$  is of dimension zero; therefore,

$$H^n i_0^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) = H^{n-r} i_0^* i^* j_* \mu_\ell(V_\alpha) = R^{n-r} i_0^* i^* j_* \mu_\ell(V_\alpha).$$

From (o), (i), (ii) and the vanishing of  $R^n i_0^* i^* j_* \mu_\ell(V_\alpha) = 0$  for  $n < 0$ , we conclude that

$$(r) \quad H^r i_0^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) \text{ is zero if } k_1 > k_2, \text{ and nonzero of weight } r - (k_1 + k_2) \text{ if } k_1 = k_2,$$

$$(r+1) \quad 0 \neq H^{r+1} i_0^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) \text{ is of weight } (r + 1) - (k_1 + k_2) - 1,$$

$$(r+2) \quad 0 \neq H^{r+2} i_0^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) \text{ is of weight } (r + 2) - (k_1 - k_2),$$

and that  $H^n i_0^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) = 0$  whenever  $n \leq r - 1$ .

(b): According to [Corollary 2.5\(b\)](#) and [Proposition 2.4\(a\)](#),

- (o)  $0 \neq H^0(\text{Lie}(W_1), V_{\underline{\alpha}})$  is of weight  $(r + 1) - k_1 - 1$ ,
- (i)  $0 \neq H^1(\text{Lie}(W_1), V_{\underline{\alpha}})$  is of weight  $(r + 2) - k_2 - 1$ ,

and  $H^q(\text{Lie}(W_1), V_{\underline{\alpha}}) = 0$  whenever  $q < 0$ . The group  $H_C/K_W$  associated to a stratum  $Z'$  of  $Z_1$  is trivial. Pink's theorem and [Pink 1992, Lemma (5.6.6)] then tell us that

- (o)  $0 \neq R^0 i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}})$  is of weight  $(r + 1) - k_1 - 1$ ,
- (i)  $0 \neq R^1 i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}})$  is of weight  $(r + 2) - k_2 - 1$ ,

and that  $R^n i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}}) = 0$  whenever  $n < 0$ . Furthermore, Pink's theorem tells us that all classical cohomology objects  $R^n i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}})$ ,  $n \in \mathbb{Z}$ , are lisse. The formula

$$H^n i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) = H^{n-r} i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}}) = (R^{n-r-1} i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}}))[1]$$

is valid: the first equation comes from

$$R_{\ell, M^K}(\alpha \mathcal{V}) = \mu_{\ell}(V_{\underline{\alpha}})[-r].$$

As for the second, note that any lisse  $\ell$ -adic sheaf  $\mathcal{F}$  on a one-dimensional regular scheme is a perverse sheaf  $\mathcal{F}'$  up to a shift by  $-1$ :


$$\mathcal{F} = \mathcal{F}'[-1] \quad \text{and} \quad \mathcal{F}' = \mathcal{F}[1].$$

From (o), (i) and the vanishing of  $R^n i_1^* i^* j_* \mu_{\ell}(V_{\underline{\alpha}}) = 0$  for  $n < 0$ , we conclude that

- (r+1)  $0 \neq H^{r+1} i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V})$  is of weight  $(r + 1) - k_1$ ,
- (r+2)  $0 \neq H^{r+2} i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V})$  is of weight  $(r + 2) - k_2$ ,

and that  $H^n i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}) = 0$  whenever  $n \leq r$ . □

For the final step of the proof of Theorem 1.6, the following commutative diagram of immersions will be useful:



Immersiones situated on the same line are complementary to each other (example:  $j''$  and  $i''$ ), the four immersiones marked by “ $\circ$ ” are open (example:  $i_1$ ), and the other four are closed (example:  $i'$ ).

**Remark 2.7.** Denote by  $\tau_{Z_m}^{t \leq \bullet}$  and  $\tau_{Z_m}^{t \geq \bullet}$  the truncation functors with respect to the perverse  $t$ -structure on  $Z_m$ ,  $m = 0, 1$ .

(a) The immersiones  $j'$  and  $i'$  being complementary,

$$(i')^* j'_{!*} \mathcal{F}' = \tau_{Z_1}^{t \leq -1} (i')^* j'_* \mathcal{F}'$$

for any perverse sheaf  $\mathcal{F}'$  on  $M^K$  [Beilinson et al. 1982, Proposition 1.4.23].

(b) The intermediate extension is transitive, i.e.,

$$j!_* = j''_{!*} j'_{!*}$$

[Beilinson et al. 1982, Corollaire 1.4.24]. Application of the functor  $(i'')^* j''_{!*}$  to the exact triangle

$$i'_* \tau_{Z_1}^{t \geq 0} (i')^* j'_*[-1] \rightarrow j'_{!*} \rightarrow j'_* \rightarrow i'_* \tau_{Z_1}^{t \geq 0} (i')^* j'_*$$

of functors on perverse sheaves on  $M^K$  (see (a)) yields the exact triangle

$$i_0^* i_{1,*} \tau_{Z_1}^{t \geq 0} (i')^* j'_*[-1] \rightarrow (i'')^* j''_{!*} j'_{!*} \rightarrow i_0^* i_{1,*} j'_* \rightarrow i_0^* i_{1,*} \tau_{Z_1}^{t \geq 0} (i')^* j'_*.$$

The immersiones  $j''$  and  $i''$  being complementary, we have as in (a)

$$(i'')^* j''_{!*} \mathcal{F}'' = \tau_{Z_0}^{t \leq -1} (i'')^* j''_* \mathcal{F}''$$

for any perverse sheaf  $\mathcal{F}''$  on  $(M^K)^* - Z_0$ . It follows that for any perverse sheaf  $\mathcal{F}'$  on  $M^K$ , there are exact sequences of perverse cohomology objects

$$\begin{aligned} H^{n-1}(i_0^* i_{1,*} \tau_{Z_1}^{t \geq 0} i_1^* i^* j_* \mathcal{F}') &\rightarrow H^n(i_0^* i^* j_* \mathcal{F}') \\ &\rightarrow H^n(i_0^* i^* j_* \mathcal{F}') \rightarrow H^n(i_0^* i_{1,*} \tau_{Z_1}^{t \geq 0} i_1^* i^* j_* \mathcal{F}') \end{aligned}$$

for  $n \leq -1$ , while  $H^n(i_0^* i^* j_* \mathcal{F}') = 0$  for all  $n \geq 0$ .

(c) Recall that  $R_{\ell, M^K}(\alpha \mathcal{V}) = \mu_{\ell}(V_{\alpha})[-r]$ ; the variety  $M^K$  being of dimension three, the complex  $R_{\ell, M^K}(\alpha \mathcal{V})$  is therefore concentrated in perverse degree  $r + 3$ . According to our conventions,  $i_1^* i^* j_{!*} R_{\ell, M^K}(\alpha \mathcal{V}) = (i')^* j'_{!*} R_{\ell, M^K}(\alpha \mathcal{V})$  thus equals

$$((i')^* j'_{!*}(R_{\ell, M^K}(\alpha \mathcal{V})[r + 3]))[-(r + 3)].$$

According to (a), we thus have

$$i_1^* i^* j_{!*} R_{\ell, M^K}(\alpha \mathcal{V}) = \tau_{Z_1}^{t \leq r+2} (i')^* j'_* R_{\ell, M^K}(\alpha \mathcal{V}) = \tau_{Z_1}^{t \leq r+2} i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}).$$

Similarly, following (b),

$$H^n i_0^* i^* j_{!*} R_{\ell, M^K}(\alpha \mathcal{V}) = 0$$

for all  $n \geq r + 3$ , and there are exact sequences

$$\begin{aligned} H^{n-1} i_0^* i_{1,*} \tau_{Z_1}^{t \geq r+3} i_1^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}) &\rightarrow H^n i_0^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}) \\ &\rightarrow H^n i_0^* i_{1,*} \tau_{Z_1}^{t \geq r+3} i_1^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}) \end{aligned}$$

for  $n \leq r + 2$ .

(d) We claim that

$$H^n i_0^* i_{1,*} \tau_{Z_1}^{t \geq r+3} i_1^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}) = 0$$

for all  $n \leq r + 1$ . Equivalently,

$$H^n i_0^* i_{1,*} \tau_{Z_1}^{t \geq 3} i_1^* i^* j_* \mu_{\ell}(V_{\alpha}) = 0$$

for all  $n \leq 1$ . Indeed, by Pink's theorem, the classical cohomology objects of  $i_1^* i^* j_* \mu_{\ell}(V_{\alpha})$  are all lisse. Applying  $\tau_{Z_1}^{t \geq 3}$ , we thus get a complex concentrated in classical degrees  $\geq 2$  (recall that  $Z_1$  is of dimension one). The same is thus true after application of  $i_0^* i_{1,*}$  (recall that inverse images are  $t$ -exact for the classical  $t$ -structure). In other words, the complex

$$i_0^* i_{1,*} \tau_{Z_1}^{t \geq 3} i_1^* i^* j_* \mu_{\ell}(V_{\alpha})$$

has trivial cohomology (classical or perverse; recall that  $Z_0$  is of dimension zero) in degrees  $\leq 1$ .

(e) From (c) and (d), we deduce that

$$H^n i_0^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}) \xrightarrow{\sim} H^n i_0^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V})$$

for  $n \leq r + 1$ , and that  $H^{r+2} i_0^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V})$  equals the kernel of

$$H^{r+2} i_0^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}) \rightarrow H^{r+2} i_0^* i_{1,*} \tau_{Z_1}^{t \geq r+3} i_1^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V}).$$

**Corollary 2.8.** *Let  $\ell$  be a prime number.*

(a) For all  $n \in \mathbb{Z}$ ,

$$H^n i_0^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V})$$

is of weights  $\leq n - (k_1 - k_2)$ .

(b) For all  $n \in \mathbb{Z}$ ,

$$H^n i_1^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V})$$

is of weights  $\leq n - k_2$ . The perverse cohomology sheaf

$$H^{r+2} i_1^* i^* j_* R_{\ell, M^k}(\alpha \mathcal{V})$$

is nonzero, and pure of weight  $(r + 2) - k_2$ .

*Proof.* Part (a) follows from [Remark 2.7\(c\)](#), (e), and from [Theorem 2.3\(a\)](#). Part (b) follows from [Remark 2.7\(c\)](#), and from [Theorem 2.3\(b\)](#).  $\square$



**Corollary 2.8** suffices to prove the part of **Theorem 1.6(b)** asserting that regularity of  $\underline{\alpha}$  is sufficient for weights 0 and 1 to be avoided by  $i^*j_*^\alpha \mathcal{V}$ . In order to prove that it is necessary, we need the following statement.

**Proposition 2.9.** *Let  $\ell$  be a prime number. Then provided that  $k_1 \geq 1$ , the perverse cohomology sheaf*

$$H^{r+2}i_0^*i^*j_{1*}R_{\ell, M^K}(\alpha \mathcal{V})$$

*is nonzero, and pure of weight  $(r + 2) - (k_1 - k_2)$ .*

*Proof.* According to **Remark 2.7(e)**,

$$H^{r+2}i_0^*i^*j_{1*}R_{\ell, M^K}(\alpha \mathcal{V})$$

equals the kernel of

$$\text{ad} : H^{r+2}i_0^*i^*j_*R_{\ell, M^K}(\alpha \mathcal{V}) \rightarrow H^{r+2}i_0^*i_{1,*}\tau_{Z_1}^{t \geq r+3}i_1^*i^*j_*R_{\ell, M^K}(\alpha \mathcal{V})$$

— in particular, it is pure of weight  $(r + 2) - (k_1 - k_2)$  (**Theorem 2.3(a)**) — i.e., it equals the kernel of

$$H^2i_0^*i^*j_*\mu_\ell(V_\alpha) \rightarrow H^2i_0^*i_{1,*}\tau_{Z_1}^{t \geq 3}i_1^*i^*j_*\mu_\ell(V_\alpha).$$

Thanks to Pink’s theorem, the regularity of  $H^2(\text{Lie}(W_0), V_\alpha)$  as a representation of  $Q_0/W_0$  (**Corollary 2.5(a)**), and the fact that the group  $H_C/K_W$  is of cohomological dimension one, locally on  $Z_0$ , the (perverse or classical) sheaf

$$H^2i_0^*i^*j_*\mu_\ell(V_\alpha) = R^2i_0^*i^*j_*\mu_\ell(V_\alpha)$$

equals

$$\mu_{\ell, Z'}(H^1(H_C/K_W, H^1(\text{Lie}(W_0), V_\alpha))),$$

for a stratum  $Z'$  of  $\Phi'$  contributing to  $Z_0$ . Furthermore, by **Corollary 2.5(a)**, the restriction of  $H^1(\text{Lie}(W_0), V_\alpha)$  to  $H_C/K_W$  is isomorphic to the  $(k_1 + k_2 + 2)$ -nd symmetric power of the standard representation of  $\text{SL}_{2, \mathbb{Q}}$ . Therefore, by **Remark 2.6**,  $H^2i_0^*i^*j_*\mu_\ell(V_\alpha)|_{Z'}$  is of constant rank  $(k_1 + k_2 + 3)(2g - 2 + c)$ , where  $g$  denotes the genus of  $H_C/K_W$ , and  $c$  the number of cusps.

We claim that the restriction to the same  $Z'$  of

$$H^2i_0^*i_{1,*}\tau_{Z_1}^{t \geq 3}i_1^*i^*j_*\mu_\ell(V_\alpha)$$

is of constant rank  $c$ . Indeed, according to **Remark 2.7(d)**, the classical cohomology objects of  $i_1^*i^*j_*\mu_\ell(V_\alpha)$  are all lisse. Therefore, perverse truncation above degree three equals classical truncation above degree two (recall that  $Z_1$  is of dimension one). The complex

$$i_0^*i_{1,*}\tau_{Z_1}^{t \geq 3}i_1^*i^*j_*\mu_\ell(V_\alpha)$$

is concentrated in degrees  $\geq 2$ , and we get

$$H^2 i_0^* i_{1,*} \tau_{Z_1}^{t \geq 3} i_1^* i^* j_* \mu_\ell(V_\alpha) = R^0 i_0^* i_{1,*} R^2 i_1^* i^* j_* \mu_\ell(V_\alpha).$$

Restriction to  $Z'$  yields

$$H^2 i_0^* i_{1,*} \tau_{Z_1}^{t \geq 3} i_1^* i^* j_* \mu_\ell(V_\alpha)|_{Z'} = \bigoplus_{Z''} (R^0 i_0^* i_{1,*} (R^2 i_1^* i^* j_* \mu_\ell(V_\alpha)|_{Z''}))|_{Z'},$$

where the direct sum is indexed by all strata  $Z''$  contributing to  $Z_1$ , and containing  $Z'$  in their closure. For every such  $Z''$ ,

$$R^2 i_1^* i^* j_* \mu_\ell(V_\alpha)|_{Z''} = \mu_{\ell, Z''}(H^2(\mathrm{Lie}(W_1), V_\alpha))$$

according to Pink's theorem (since the group  $H_C/K_W$  (for  $m = 1!$ ) is trivial).

Denote by  $j_1 : Z_1 \hookrightarrow Z_1^*$  the Baily–Borel compactification, and by  $i_{01} : \partial Z_1^* \hookrightarrow Z_1^*$  its complement. The immersion  $i_1 : Z_1 \hookrightarrow (M^K)^*$  admits a natural extension  $\bar{i}_1 : Z_1^* \rightarrow (M^K)^*$  [Pink 1989, Main Theorem 12.3(c), Section 7.6], which is finite. The diagram

$$\begin{array}{ccc} Z_1^* & \xleftarrow{i_{01}} & \partial Z_1^* \\ \bar{i}_1 \downarrow & & \downarrow \bar{i}_1 \\ (M^K)^* & \xleftarrow{i_0} & Z_0 \end{array}$$

is cartesian up to nilpotent elements. Proper base change therefore yields the formula

$$R^0 i_0^* i_{1,*} = R^0 \bar{i}_{1,*} i_{0,1}^* j_{1,*}.$$

The functors  $\bar{i}_{1,*}$  and  $i_{0,1}^*$  being exact on sheaves, we have

$$R^0 i_0^* i_{1,*} (R^2 i_1^* i^* j_* \mu_\ell(V_\alpha)|_{Z''}) = \bar{i}_{1,*} i_{0,1}^* R^0 j_{1,*} \mu_{\ell, Z''}(H^2(\mathrm{Lie}(W_1), V_\alpha)).$$

According to Proposition 2.4(a),  $H^2(\mathrm{Lie}(W_1), V_\alpha)$  is irreducible as a representation of  $Q_1/W_1$ , and hence of  $\mathrm{GL}_{2, \mathbb{Q}}$ . Yet another application of Pink's theorem shows that

$$i_{0,1}^* R^0 j_{1,*} \mu_{\ell, Z''}(H^2(\mathrm{Lie}(W_1), V_\alpha))$$

is of constant rank one on the intersection of  $\partial Z_1^*$  with the closure of  $Z''$  in  $(Z_1)^*$ .

Our claim on the rank of

$$H^2 i_0^* i_{1,*} \tau_{Z_1}^{t \geq 3} i_1^* i^* j_* \mu_\ell(V_\alpha)|_{Z'} = \bar{i}_{1,*} \bigoplus_{Z''} (i_{0,1}^* R^0 j_{1,*} \mu_{\ell, Z''}(H^2(\mathrm{Lie}(W_1), V_\alpha)))|_{Z'}$$

is therefore proven as soon as we establish that the number of points in the geometrical fibres of the morphism  $\bar{i}_1 : \partial Z_1^* \rightarrow Z_0$  above  $Z' \subset Z_0$  equals  $c$ . This verification can be done on the level of  $\mathbb{C}$ -valued points, where the adelic description of the situation is at our disposal. More precisely, write  $(G_m, \mathcal{H}_m) := (P_m, \mathfrak{X}_m)/W_m$  [Pink 1989, Proposition 2.9],  $m = 0, 1$ , for the Shimura data contributing to  $\partial(M^K)^*$ , and

$Q_{01}$  for the Borel subgroup  $Q_0 \cap Q_1$  of  $G$ . According to [Pink 1989, Section 6.3], the diagram of  $\mathbb{C}$ -valued points corresponding to the diagram

$$\begin{array}{ccc} Z_1^* & \xleftarrow{i_{01}} & \partial Z_1^* \\ \bar{i}_1 \downarrow & & \downarrow \bar{i}_1 \\ (M^K)^* & \xleftarrow{i_0} & Z_0 \end{array}$$

equals

$$\begin{array}{ccc} Q_1(\mathbb{Q}) \backslash (\mathcal{H}_1^* \times G(\mathbb{A}_f) / K) & \xleftarrow{i_{01}} & Q_{01}(\mathbb{Q}) \backslash (\mathcal{H}_0 \times G(\mathbb{A}_f) / K) \\ \bar{i}_1 \downarrow & & \downarrow \bar{i}_1 \\ G(\mathbb{Q}) \backslash (\mathcal{H}^* \times G(\mathbb{A}_f) / K) & \xleftarrow{i_0} & Q_0(\mathbb{Q}) \backslash (\mathcal{H}_0 \times G(\mathbb{A}_f) / K) \end{array}$$

where all maps are induced by canonical inclusions of groups and spaces. Indeed, the full group  $Q_m(\mathbb{Q})$  (and not only a subgroup of finite index) stabilizes  $\mathcal{H}_m$ ,  $m = 0, 1$ , and two rational boundary components of  $(G_1, \mathcal{H}_1)$  are conjugate under  $G_1(\mathbb{Q})$  if and only if they are conjugate under  $G(\mathbb{Q})$  (by explicit computation, or [Pink 1989, Remark (iii) on p. 91]). The subscheme  $Z' \subset Z_0$  equals the image of a Shimura variety associated to  $(G_0, \mathcal{H}_0)$  under a morphism  $i_g$  associated to an element  $g \in G(\mathbb{A}_f)$  [Pink 1989, Main Theorem 12.3(c)]; given the adelic description of  $i_g$  from [Pink 1989, Section 6.3], we see that under the above identification, any  $z \in Z'(\mathbb{C})$  equals the class  $[h_0, p_0g]$  in

$$Q_0(\mathbb{Q}) \backslash (\mathcal{H}_0 \times G(\mathbb{A}_f) / K)$$

of a pair of the form  $(h_0, p_0g)$ , with  $h_0 \in \mathcal{H}_0$  and  $p_0 \in P_0(\mathbb{A}_f)$ . Put

$$Q_0^+(\mathbb{Q}) := \{q_0 \in Q_0(\mathbb{Q}) : \lambda(q_0) > 0\};$$

this group equals the centralizer in  $Q_0(\mathbb{Q})$  of  $h_0$ , and indeed, of the whole of  $\mathcal{H}_0$ . Putting

$$H'_C := Q_0^+(\mathbb{Q}) \cap p_0gKg^{-1}p_0^{-1},$$

we leave it to the reader to verify that the map

$$Q_{01}(\mathbb{Q}) \backslash Q_0(\mathbb{Q}) / H'_C \rightarrow \bar{i}_1^{-1}(z), \quad [q_0] \mapsto q_0[h_0, p_0g] = [q_0h_0, q_0p_0g]$$

is well-defined, and bijective. By strong approximation,

$$W_0(\mathbb{Q}) \cdot H'_C = Q_0^+(\mathbb{Q}) \cap W_0(\mathbb{A}_f) \cdot p_0gKg^{-1}p_0^{-1}.$$

But

$$Q_0 / W_0 = P_0 / W_0 \times_{\mathbb{Q}} \mathrm{GL}_{2, \mathbb{Q}},$$

meaning that modulo  $W_0$ , elements in  $P_0$  and in  $Q_0$  commute with each other.

Thus,

$$W_0(\mathbb{Q}) \cdot H'_C = Q_0^+(\mathbb{Q}) \cap W_0(\mathbb{A}_f) \cdot gKg^{-1}.$$

The image of  $W_0(\mathbb{Q}) \cdot H'_C$  under the projection  $\pi_0 : Q_0 \rightarrow Q_0/W_0$  coincides with the image of

$$W_0(\mathbb{A}_f) \cdot Q_0^+(\mathbb{Q}) \cap gKg^{-1}$$

(both images equal  $\pi_0(Q_0^+(\mathbb{Q})) \cap \pi(gKg^{-1})$ ). But by definition [Pink 1992, (3.7.4)],  $W_0(\mathbb{A}_f) \cdot Q_0^+(\mathbb{Q}) \cap gKg^{-1}$  equals  $H_C$ . We thus showed that

$$\pi_0(H'_C) = \pi_0(H_C).$$

Now the quotient morphism  $Q_0 \rightarrow Q_0/P_0$ ,  $q_0 \mapsto \bar{q}_0$  induces an isomorphism

$$Q_{01}(\mathbb{Q}) \backslash Q_0(\mathbb{Q})/H'_C \xrightarrow{\sim} \overline{Q_{01}(\mathbb{Q})} \backslash \overline{Q_0(\mathbb{Q})}/\overline{H'_C} = \overline{Q_{01}(\mathbb{Q})} \backslash \overline{Q_0(\mathbb{Q})}/\overline{H_C}.$$

But  $\overline{Q_0(\mathbb{Q})} = \text{GL}_2(\mathbb{Q})$ , and under this identification,  $\overline{Q_{01}(\mathbb{Q})}$  equals the subgroup of upper triangular matrices, while  $\overline{H_C} = H_C/K_W$ . In other words,

$$Q_{01}(\mathbb{Q}) \backslash Q_0(\mathbb{Q})/H'_C$$

is identified with the set up cusps of  $H_C/K_W$ .

The formula

$$(k_1 + k_2 + 3)(2g - 2 + c) \geq 4(2g - 2 + c) > c$$

(recall that  $c$  is greater or equal to 1, and that  $c \geq 3$  if  $g = 0$ ) implies that the rank of the source of  $\text{ad}$  is strictly greater than the rank of its target; the kernel of  $\text{ad}$  is therefore nontrivial.  $\square$

**Remark 2.10.** (a) As the reader may verify,

$$H^{r+2} i_0^* i_{1,*} \tau_{Z_1}^{t \geq r+3} i_1^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V})$$

is pure of weight  $(r + 2) - (k_1 - k_2)$ , i.e., of the same weight as

$$H^{r+2} i_0^* i^* j_* R_{\ell, M^K}(\alpha \mathcal{V}).$$

Weight considerations alone therefore do not imply nontriviality of the kernel of the map  $\text{ad}$  from the proof of Proposition 2.9.

(b) A more conceptual proof of Proposition 2.9 would consist in showing that locally on  $Z_0$ , the map  $\text{ad}$  equals the direct sum over all cusps of  $H_C/K_W$  of the residue maps. Identify  $H^1(H_C/K_W, H^1(\text{Lie}(W_0), V_{\alpha})) \otimes_{\mathbb{Q}} \mathbb{C}$  with the direct sum of the space of modular forms and (the conjugate of) the space of cusp forms for  $H_C/K_W$  of weight  $k_1 + k_2 + 4 \geq 5$ . The kernel of the residues contains the space of cusp forms. Its dimension is computed in [Shimura 1971, Theorems 2.24 and 2.25]; thanks to [Shimura 1971, Proposition 1.40] (always remember that  $H_C/K_W$  is neat), this dimension can be seen to be strictly positive.

(c) On the level of geometry of Baily–Borel compactifications, a “strange duality” seems to be involved in the proof of [Proposition 2.9](#): we need to know how many modular curves in the boundary of  $(M^K)^*$  contain a given cusp  $Z'$  in their closure. The response yields the number of cusps of a “modular curve”, which does not explicitly occur in  $(M^K)^*$ , namely the quotient of the upper half space by  $H_C/K_W$ . It would be interesting to see how this phenomenon generalizes to higher dimensional Siegel varieties.

(d) Our computation of the fibres of the morphism  $\bar{i}_1 : Z_1^* \rightarrow (M^K)^*$  over points of  $Z_0$  is a quantitative version of a classical noninjectivity result of Satake [[1958](#), Exemple on p. 13-06].

**Remark 2.11.** The Hodge theoretic analogues of [Theorem 2.3](#), [Corollary 2.8](#) and [Proposition 2.9](#) hold. The proofs are identical up to the use of Pink’s theorem, which is replaced by [[Burgos and Wildeshaus 2004](#), Theorem 2.9].

*Proof of Theorem 1.6.* According to [Theorem 2.2](#),  $i^* j_*^\alpha \mathcal{V}$  is a  $\Phi$ -constructible motive of abelian type over  $\partial(M^K)^*$ ; this proves part (a) of our claim.

By [[Pink 1989](#), Summary 1.18(d)], there is a perfect pairing

$$V_\alpha \otimes_{\mathbb{Q}} V_\alpha \rightarrow \mathbb{Q}(-r)$$

in  $\text{Rep}(G)$ .

Fix a prime  $\ell$ . Applying  $\mu_\ell$ , we get a perfect pairing

$$\mu_\ell(V_\alpha) \otimes_{\mathbb{Q}_\ell} \mu_\ell(V_\alpha) \rightarrow \mathbb{Q}_\ell(-r)$$

of  $\ell$ -adic lisse sheaves on  $M^K$ . In terms of local duality, the pairing induces an isomorphism

$$\mathbb{D}_{\ell, M^K}(\mu_\ell(V_\alpha)) \cong \mu_\ell(V_\alpha)(r + 3)[6]$$

( $M^K$  is smooth of dimension three). Given  $R_{\ell, M^K}(\alpha\mathcal{V}) = \mu_\ell(V_\alpha)[-r]$ , we find that

$$\mathbb{D}_{\ell, M^K}(R_{\ell, M^K}(\alpha\mathcal{V})) \cong R_{\ell, M^K}(\alpha\mathcal{V})(s)[2s],$$

where  $s = r + 3$ .

[Corollary 2.8](#) tells us that for all  $n \in \mathbb{Z}$ , and  $m = 0, 1$ ,

$$H^n i_m^* i^* j_{!*} R_{\ell, M^K}(\alpha\mathcal{V})$$

is of weights  $\leq n - k$ . According to [[Wildeshaus 2018a](#), Corollary 4.6(b)], the motive  $i^* j_*^\alpha \mathcal{V}$  therefore avoids weights  $-k + 1, -k + 2, \dots, k$ .

In order to conclude the proof of part (b), it remains to show, again thanks to [[Wildeshaus 2018a](#), Corollary 4.6(b)], that for some  $n \in \mathbb{Z}$ , and  $m = 0$  or  $m = 1$ , weight  $n - k$  does occur in

$$H^n i_m^* i^* j_{!*} R_{\ell, M^K}(\alpha\mathcal{V}).$$

We take  $n = r + 2$ , and distinguish two cases. If  $k = k_2$ , i.e.,  $k_2 \leq k_1 - k_2$ , take  $m = 1$ ; the claim then follows from [Corollary 2.8\(b\)](#). Else,  $k_2 > k_1 - k_2$  and  $k = k_1 - k_2$ . Since  $k_1 \geq k_2$ , we necessarily have  $k_1 \geq 1$ . Take  $m = 0$  and apply [Proposition 2.9](#).  $\square$

**Remark 2.12.** (a) An element of  $H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$  is called a *ghost class* if it lies in the image of

$$H^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \rightarrow H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$$

and in the kernel of both restriction maps

$$H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \rightarrow H^n(Z_m(\mathbb{C}), i_m^* i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}})),$$

$m = 0, 1$ . One of the main results of [\[Moya Giusti 2018\]](#) implies that if  $\underline{\alpha}$  is regular, then there are no nonzero ghost classes [\[Moya Giusti 2018, Theorem 3.1\]](#). This result does not formally imply, nor is it implied by, our [Theorem 1.6](#). Nonetheless, it might be worthwhile to note that the weight arguments that occur in the proofs are quite similar. The most relevant information from [Theorem 1.6](#), as far as [\[Moya Giusti 2018, Theorem 3.1\]](#) is concerned, comes from the weight filtration

$$a_* j_{!*}^{\alpha} \mathcal{V} \rightarrow \tilde{a}_*^{\alpha} \mathcal{V} \rightarrow a_* i_* i^! j_{!*}^{\alpha} \mathcal{V}[1] \rightarrow a_* j_{!*}^{\alpha} \mathcal{V}[1]$$

avoiding weights  $1, 2, \dots, k$  ([Corollary 1.7\(a\)](#)), and hence avoiding weight 1 if  $\underline{\alpha}$  is regular, which we assume in the sequel. This implies that any element of  $H^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$  not mapping to zero in  $H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$ , remains nonzero in

$$H^n(\partial(M^K)^*(\mathbb{C}), i^! j_{!*} \mu_{\mathbf{H}}(V_{\underline{\alpha}})[1]) = H^n(\partial(M^K)^*(\mathbb{C}), \tau_{\partial(M^K)^*}^{t \geq 3} i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}})).$$

In other words, a ghost class vanishing in  $H^n(\partial(M^K)^*(\mathbb{C}), \tau_{\partial(M^K)^*}^{t \geq 3} i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$  is zero. The Hodge structure  $H^n(\partial(M^K)^*(\mathbb{C}), \tau_{\partial(M^K)^*}^{t \geq 3} i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$  has weights  $\geq (r + n) + 2$ ; the same type of considerations as those leading to [Corollary 2.8](#) then imply that the direct sum of the restriction maps

$$H^n(\partial(M^K)^*(\mathbb{C}), \tau_{\partial(M^K)^*}^{t \geq 3} i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \rightarrow H^n(Z_m(\mathbb{C}), i_m^* \tau_{\partial(M^K)^*}^{t \geq 3} i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}})),$$

$m = 0, 1$ , is injective.

(b) The above illustrates an observation made by Moya Giusti: *for a class in the cohomology of the boundary whose weight is neither the middle weight nor the middle weight plus one, we can determine exactly whether or not it is in the image of the morphism*

$$H^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \rightarrow H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}})).$$

In fact, it appears amusing to note that the “middle weight” is relevant in another context than the one studied in the present paper. According to [Moya Giusti 2018, p. 2317, second paragraph], the representation  $V_{\underline{\alpha}}$  satisfies the middle weight property if the space of ghost classes in  $H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$  is pure of weight  $r + n$ . In particular, [Moya Giusti 2018, Theorem 3.1] implies that for all  $\underline{\alpha}$  (regular or not), the representation  $V_{\underline{\alpha}}$  does satisfy the middle weight property, while our Theorem 1.6 implies that weights  $\{r + n, r + n + 1\}$  do not occur at all in  $H^n(\partial(M^K)^*(\mathbb{C}), i^* j_* \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$ , as soon as  $\underline{\alpha}$  is regular.

**Remark 2.13.** Saper’s vanishing theorem [2005, Theorem 5] says that if  $\underline{\alpha}$  is regular, then the groups  $H^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$ , and hence (by comparison)

$$H^n(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}})),$$

vanish for  $n < 3 = \dim M^K$ . By duality, one obtains that  $H_c^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) = 0$  and  $H_c^n(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}})) = 0$  for  $n > 3$ . It follows that interior cohomology with coefficients in  $\mu_{\mathbf{H}}(V_{\underline{\alpha}})$ , denoted

$$H_!^n(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})),$$

and interior cohomology with coefficients in  $\mu_{\ell}(V_{\underline{\alpha}})$ , denoted

$$H_!^n(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}})),$$

both vanish for  $n \neq 3$ , provided that  $\underline{\alpha}$  is regular.

### 3. The motive for an automorphic form

This final section contains the analogues for Siegel threefolds of the main results from [Scholl 1990]. Since we do not restrict ourselves to the case of Hecke eigenforms, our notation becomes a little more technical than in [loc. cit.].

We continue to consider the situation of Sections 1 and 2. In particular, we fix a dominant  $\underline{\alpha} = \alpha(k_1, k_2, r)$ , which we assume to be regular, i.e.,  $k_1 > k_2 > 0$ . Consider the intersection motive  $a_* j_{!*}^{\alpha} \mathcal{V} \in CHM(\mathbb{Q})_{\mathbb{Q}}$ , where  $a : (M^K)^* \rightarrow \text{Spec } \mathbb{Q}$  again denotes the structure morphism of  $(M^K)^*$ . According to [Wildeshaus 2018a, Remark 3.13(a)] and Remark 2.13, its Hodge theoretic realization equals

$$H_!^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}}))[-(r + 3)],$$

and its  $\ell$ -adic realization equals

$$H_!^3(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}}))[-(r + 3)].$$

By Corollary 1.8, every element of the Hecke algebra  $\mathfrak{H}(K, G(\mathbb{A}_f))$  acts on  $a_* j_{!*}^{\alpha} \mathcal{V}$ .

**Theorem 3.1** [Harder 2017, Theorem 3.1.1]. *Let  $L$  be any field of characteristic zero. Then the  $\mathfrak{H}(K, G(\mathbb{A}_f)) \otimes_{\mathbb{Q}} L$ -module  $H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L$  is semi-simple.*

Note that [Harder 2017, Section 8.1.6, p. 232] gives a proof of [Theorem 3.1](#), while the statement in [Harder 2017, Theorem 3.1.1] is “nonadelic”. Denote by  $R(\mathfrak{H}) := R(\mathfrak{H}(K, G(\mathbb{A}_f)))$  the image of the Hecke algebra in the endomorphism algebra of  $H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}}))$ .

**Corollary 3.2.** *Let  $L$  be any field of characteristic zero. Then the  $L$ -algebra  $R(\mathfrak{H}) \otimes_{\mathbb{Q}} L$  is semisimple.*

In particular, the isomorphism classes of simple right  $R(\mathfrak{H}) \otimes_{\mathbb{Q}} L$ -modules correspond bijectively to isomorphism classes of minimal right ideals.

Fix  $L$ , and let  $Y_{\pi_f}$  be such a minimal right ideal of  $R(\mathfrak{H}) \otimes_{\mathbb{Q}} L$ . There is a (primitive) idempotent  $e_{\pi_f} \in R(\mathfrak{H}) \otimes_{\mathbb{Q}} L$  generating  $Y_{\pi_f}$ .

**Definition 3.3.** (a) The *Hodge structure*  $W(\pi_f)$  associated to  $Y_{\pi_f}$  is defined as

$$W(\pi_f) := \text{Hom}_{R(\mathfrak{H}) \otimes_{\mathbb{Q}} L}(Y_{\pi_f}, H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L).$$

(b) Let  $\ell$  be a prime number. The *Galois module*  $W(\pi_f)_{\ell}$  associated to  $Y_{\pi_f}$  is defined as

$$W(\pi_f)_{\ell} := \text{Hom}_{R(\mathfrak{H}) \otimes_{\mathbb{Q}} L}(Y_{\pi_f}, H_1^3(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L).$$

[Definition 3.3\(b\)](#) should be compared to [Weissauer 2005, Theorem I].

**Proposition 3.4.** *There is a canonical isomorphism of Hodge structures*

$$W(\pi_f) \xrightarrow{\sim} (H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L) \cdot e_{\pi_f},$$

*and a canonical isomorphism of Galois modules*

$$W(\pi_f)_{\ell} \xrightarrow{\sim} (H_1^3(M^K \times_{\mathbb{Q}} \overline{\mathbb{Q}}, \mu_{\ell}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L) \cdot e_{\pi_f}.$$

*Proof.* We perform the proof for Hodge structures; the one for Galois modules is formally identical. Obviously,

$$\text{Hom}_{R(\mathfrak{H}) \otimes_{\mathbb{Q}} L}(R(\mathfrak{H}) \otimes_{\mathbb{Q}} L, H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L)$$

is canonically identified with

$$H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L$$

by mapping an morphism  $g$  to the image of  $1 = 1_{R(\mathfrak{H})}$  under  $g$ . Inside

$$\text{Hom}_{R(\mathfrak{H}) \otimes_{\mathbb{Q}} L}(R(\mathfrak{H}) \otimes_{\mathbb{Q}} L, H_1^3(M^K(\mathbb{C}), \mu_{\mathbf{H}}(V_{\underline{\alpha}})) \otimes_{\mathbb{Q}} L),$$



the object  $W(\pi_f)$  contains precisely those morphisms  $g$  vanishing on  $1 - e_{\pi_f}$ , or in other words, satisfying the relation  $g(1) = g(e_{\pi_f}) = g(1) \cdot e_{\pi_f}$ .  $\square$

Since we do not know whether the Chow motive  $a_* j_{!*}^{\underline{\alpha}} \mathcal{V}$  is finite dimensional, we cannot apply [Kimura 2005, Corollary 7.8], and therefore do not know whether  $e_{\pi_f}$  can be lifted *idempotently* to the Hecke algebra  $\mathfrak{H}(K, G(\mathbb{A}_f))$ . This is why we need to descend to the level of *Grothendieck motives*. Denote by  $a_* j_{!*}^{\underline{\alpha}} \mathcal{V}'$  the Grothendieck motive underlying  $a_* j_{!*}^{\underline{\alpha}} \mathcal{V}$ .

**Definition 3.5.** Assume  $\underline{\alpha} = \alpha(k_1, k_2, r)$  to be regular. Let  $L$  be a field of characteristic zero, and  $Y_{\pi_f}$  a minimal right ideal of  $R(\mathfrak{H}) \otimes_{\mathbb{Q}} L$ . The motive associated to  $Y_{\pi_f}$  is defined as

$$\mathcal{W}(\pi_f) := a_* j_{!*}^{\underline{\alpha}} \mathcal{V}' \cdot e_{\pi_f}.$$

Definition 3.5 should be compared to [Scholl 1990, Section 4.2.0]. Given our construction, the following is obvious.

**Theorem 3.6.** Assume  $\underline{\alpha} = \alpha(k_1, k_2, r)$  to be regular, i.e.,  $k_1 > k_2 > 0$ . Let  $L$  be a field of characteristic zero, and  $Y_{\pi_f}$  a minimal right ideal of  $R(\mathfrak{H}) \otimes_F L$ . The realizations of the motive  $\mathcal{W}(\pi_f)$  associated to  $Y_{\pi_f}$  are concentrated in the single cohomological degree  $r + 3$ , and they take the values  $W(\pi_f)$  (in the Hodge theoretic setting) and  $W(\pi_f)_{\ell}$  (in the  $\ell$ -adic setting).

A special case occurs when  $Y_{\pi_f}$  is of dimension one over  $L$ , i.e., corresponds to a nontrivial character of  $R(\mathfrak{H})$  with values in  $L$ . The automorphic form is then an eigenform for the Hecke algebra. This is the analogue of the situation considered in [Scholl 1990] for elliptic cusp forms.

The motive  $\mathcal{W}(\pi_f)$  being a direct factor of  $a_* j_{!*}^{\underline{\alpha}} \mathcal{V}'$ , our results on the latter from Section 1 have obvious consequences for the realizations of  $\mathcal{W}(\pi_f)$ .

**Corollary 3.7.** Assume  $\underline{\alpha} = \alpha(k_1, k_2, r)$  to be regular. Let  $L$  be a field of characteristic zero, and  $Y_{\pi_f}$  a minimal right ideal of  $R(\mathfrak{H}) \otimes_{\mathbb{Q}} L$ . Let  $p$  be a prime number not dividing the level of  $K$ . Let  $\ell$  be different from  $p$ .

- (a) The  $p$ -adic realization  $W(\pi_f)_p$  of  $\mathcal{W}(\pi_f)$  is crystalline.
- (b) The  $\ell$ -adic realization  $W(\pi_f)_{\ell}$  of  $\mathcal{W}(\pi_f)$  is unramified at  $p$ .
- (c) The characteristic polynomials of the following coincide: (1) the action of Frobenius  $\phi$  on the  $\phi$ -filtered module associated to  $W(\pi_f)_p$ ; (2) the action of a geometrical Frobenius automorphism at  $p$  on  $W(\pi_f)_{\ell}$ .

*Proof.* Parts (a) and (b) follow from Remark 1.12.

As for (c), in order to apply [Katz and Messing 1974, Theorem 2(2)], use the fact that both realizations are cut out by the *same* cycle from the cohomology of a smooth and proper scheme over the field  $\mathbb{F}_p$  (cf. the proof of Corollary 1.13).  $\square$

Corollary 3.7 should be compared to [Scholl 1990, Theorem 1.2.4].

**Remark 3.8.** Corollary 3.7(c) is already contained in [Urban 2005, Théorème 1].

### Acknowledgments

Part of this work was done during visits to Caltech’s Department of Mathematics (Pasadena), and to the Erwin Schrödinger Institute (Vienna). I am grateful to both institutions. I also wish to thank G. Ancona, J.I. Burgos Gil, M. Cavicchi, F. Déglise, F. Ivorra, F. Lemma, J. Tilouine and A. Vezzani for useful discussions and comments, as well as the referee for observations and suggestions concerning an earlier version of this article.

### References

- [Ancona 2015] G. Ancona, “Décomposition de motifs abéliens”, *Manuscripta Math.* **146**:3-4 (2015), 307–328. [MR](#) [Zbl](#)
- [Ancona 2017] G. Ancona, “Degeneration of Hodge structures over Picard modular surfaces”, *Int. J. Number Theory* **13**:5 (2017), 1145–1164. [MR](#) [Zbl](#)
- [Beilinson 2013] A. Beilinson, “On the crystalline period map”, *Camb. J. Math.* **1**:1 (2013), 1–51. [MR](#) [Zbl](#)
- [Beilinson et al. 1982] A. A. Beilinson, J. Bernstein, and P. Deligne, “Faisceaux pervers”, pp. 5–171 in *Analysis and topology on singular spaces, I* (Luminy, 1981), Astérisque **100**, Société Mathématique de France, Paris, 1982. [MR](#) [Zbl](#)
- [Bondarko 2010] M. V. Bondarko, “Weight structures vs.  $t$ -structures; weight filtrations, spectral sequences, and complexes (for motives and in general)”, *J. K-Theory* **6**:3 (2010), 387–504. [MR](#) [Zbl](#)
- [Burgos and Wildeshaus 2004] J. I. Burgos and J. Wildeshaus, “Hodge modules on Shimura varieties and their higher direct images in the Baily–Borel compactification”, *Ann. Sci. École Norm. Sup.* (4) **37**:3 (2004), 363–413. [MR](#) [Zbl](#)
- [Cisinski and Déglise 2009] D.-C. Cisinski and F. Déglise, “Triangulated categories of mixed motives”, preprint, 2009. [arXiv](#)
- [Cloître 2017] G. Cloître, “On the interior motive of certain Shimura varieties: the case of Picard varieties”, preprint, 2017. [arXiv](#)
- [Déglise and Nizioł 2018] F. Déglise and W. Nizioł, “On  $p$ -adic absolute Hodge cohomology and syntomic coefficients, I”, *Comment. Math. Helv.* **93**:1 (2018), 71–131. [MR](#) [Zbl](#)
- [Deninger and Murre 1991] C. Deninger and J. Murre, “Motivic decomposition of abelian schemes and the Fourier transform”, *J. Reine Angew. Math.* **422** (1991), 201–219. [MR](#) [Zbl](#)
- [Faltings and Chai 1990] G. Faltings and C.-L. Chai, *Degeneration of abelian varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) **22**, Springer, 1990. [MR](#) [Zbl](#)
- [Harder 2017] G. Harder, “Cohomology of arithmetic groups”, preprint, 2017, available at <http://www.math.uni-bonn.de/people/harder/Manuscripts/buch/Volume-III.pdf>.
- [Hébert 2011] D. Hébert, “Structure de poids à la Bondarko sur les motifs de Beilinson”, *Compos. Math.* **147**:5 (2011), 1447–1462. [MR](#) [Zbl](#)
- [Katz and Messing 1974] N. M. Katz and W. Messing, “Some consequences of the Riemann hypothesis for varieties over finite fields”, *Invent. Math.* **23** (1974), 73–77. [MR](#) [Zbl](#)

- [Kimura 2005] S.-I. Kimura, “Chow groups are finite dimensional, in some sense”, *Math. Ann.* **331**:1 (2005), 173–201. [MR](#) [Zbl](#)
- [Lemma 2015] F. Lemma, “On higher regulators of Siegel threefolds, I: The vanishing on the boundary”, *Asian J. Math.* **19**:1 (2015), 83–120. [MR](#) [Zbl](#)
- [Levine 2009] M. Levine, “Smooth motives”, pp. 175–231 in *Motives and algebraic cycles*, edited by R. de Jeu and J. D. Lewis, Fields Inst. Commun. **56**, Amer. Math. Soc., Providence, RI, 2009. [MR](#) [Zbl](#)
- [Levine 2010] M. Levine, “Tate motives and the fundamental group”, pp. 265–392 in *Cycles, motives and Shimura varieties*, edited by V. Srinivas, Tata Inst. Fund. Res. Stud. Math. **21**, Tata Inst. Fund. Res., Mumbai, 2010. [MR](#) [Zbl](#)
- [Mokrane and Tilouine 2002] A. Mokrane and J. Tilouine, “Cohomology of Siegel varieties with  $p$ -adic integral coefficients and applications”, pp. 1–95 in *Cohomology of Siegel varieties*, Astérisque **280**, Société Mathématique de France, Paris, 2002. [MR](#) [Zbl](#)
- [Moya Giusti 2018] M. V. Moya Giusti, “Ghost classes in the cohomology of the Shimura variety associated to  $\mathrm{GSp}_4$ ”, *Proc. Amer. Math. Soc.* **146**:6 (2018), 2315–2325. [MR](#) [Zbl](#)
- [O’Sullivan 2011] P. O’Sullivan, “Algebraic cycles on an abelian variety”, *J. Reine Angew. Math.* **654** (2011), 1–81. [MR](#) [Zbl](#)
- [Pink 1989] R. Pink, *Arithmetical compactification of mixed Shimura varieties*, dissertation, Universität Bonn, 1989, available at <https://people.math.ethz.ch/~pink/dissertation.html>. [MR](#) [Zbl](#)
- [Pink 1992] R. Pink, “On  $l$ -adic sheaves on Shimura varieties and their higher direct images in the Baily–Borel compactification”, *Math. Ann.* **292**:2 (1992), 197–240. [MR](#) [Zbl](#)
- [Saper 2005] L. Saper, “ $\mathcal{L}$ -modules and the conjecture of Rapoport and Goresky–MacPherson”, pp. 319–334 in *Automorphic forms*, vol. I, Astérisque **298**, Société Mathématique de France, Paris, 2005. [MR](#) [Zbl](#)
- [Satake 1958] I. Satake, “Compactification des espaces quotients de Siegel, II”, exposé 13 in *Séminaire Henri Cartan*, 1957/1958, vol. 2, Secr. Math., Paris, 1958.
- [Scholl 1990] A. J. Scholl, “Motives for modular forms”, *Invent. Math.* **100**:2 (1990), 419–430. [MR](#) [Zbl](#)
- [Shimura 1971] G. Shimura, *Introduction to the arithmetic theory of automorphic functions*, Publications of the Mathematical Society of Japan **11**, Iwanami Shoten, Tokyo, 1971. [MR](#) [Zbl](#)
- [Urban 2005] E. Urban, “Sur les représentations  $p$ -adiques associées aux représentations cuspidales de  $\mathrm{GSp}_4/\mathbb{Q}$ ”, pp. 151–176 in *Formes automorphes, II: Le cas du groupe  $\mathrm{GSp}(4)$* , Astérisque **302**, Société Mathématique de France, Paris, 2005. [MR](#) [Zbl](#)
- [Vogan 1981] D. A. Vogan, Jr., *Representations of real reductive Lie groups*, Progress in Math. **15**, Birkhäuser, Boston, 1981. [MR](#) [Zbl](#)
- [Weissauer 2005] R. Weissauer, “Four dimensional Galois representations”, pp. 67–150 in *Formes automorphes, II: Le cas du groupe  $\mathrm{GSp}(4)$* , Astérisque **302**, Société Mathématique de France, Paris, 2005. [MR](#) [Zbl](#)
- [Wildeshaus 1997] J. Wildeshaus, “The canonical construction of mixed sheaves on mixed Shimura varieties”, pp. 77–140 in *Realizations of polylogarithms*, Lecture Notes in Math. **1650**, Springer, 1997. [MR](#) [Zbl](#)
- [Wildeshaus 2007] J. Wildeshaus, “On the boundary motive of a Shimura variety”, *Compos. Math.* **143**:4 (2007), 959–985. [MR](#) [Zbl](#)
- [Wildeshaus 2009] J. Wildeshaus, “Chow motives without projectivity”, *Compos. Math.* **145**:5 (2009), 1196–1226. [MR](#) [Zbl](#)

- [Wildeshaus 2012a] J. Wildeshaus, “[Motivic intersection complex](#)”, pp. 255–276 in *Regulators*, edited by J. I. Burgos Gil et al., Contemp. Math. **571**, Amer. Math. Soc., Providence, RI, 2012. [MR](#) [Zbl](#)
- [Wildeshaus 2012b] J. Wildeshaus, “[On the interior motive of certain Shimura varieties: the case of Hilbert–Blumenthal varieties](#)”, *Int. Math. Res. Not.* **2012**:10 (2012), 2321–2355. [MR](#) [Zbl](#)
- [Wildeshaus 2015] J. Wildeshaus, “[On the interior motive of certain Shimura varieties: the case of Picard surfaces](#)”, *Manuscripta Math.* **148**:3–4 (2015), 351–377. [MR](#) [Zbl](#)
- [Wildeshaus 2017] J. Wildeshaus, “[Intermediate extension of Chow motives of abelian type](#)”, *Adv. Math.* **305** (2017), 515–600. [MR](#) [Zbl](#)
- [Wildeshaus 2018a] J. Wildeshaus, “[Chow motives without projectivity, II](#)”, *Int. Math. Res. Not.* (online publication November 2018).
- [Wildeshaus 2018b] J. Wildeshaus, “[Weights and conservativity](#)”, *Algebr. Geom.* **5**:6 (2018), 686–702. [MR](#) [Zbl](#)

Received 5 Oct 2017. Revised 22 Mar 2019. Accepted 16 Apr 2019.

JÖRG WILDESHAUS: [wildesh@math.univ-paris13.fr](mailto:wildesh@math.univ-paris13.fr)

Laboratoire Analyse, Géométrie et Applications, Université Paris 13, Sorbonne Paris Cité, Villetaneuse, France

# ANNALS OF K-THEORY

[msp.org/akt](http://msp.org/akt)

## EDITORIAL BOARD

Joseph Ayoub	ETH Zürich, Switzerland <a href="mailto:joseph.ayoub@math.uzh.ch">joseph.ayoub@math.uzh.ch</a>
Paul Balmer	University of California, Los Angeles, USA <a href="mailto:balmer@math.ucla.edu">balmer@math.ucla.edu</a>
Guillermo Cortiñas	Universidad de Buenos Aires and CONICET, Argentina <a href="mailto:gcorti@dm.uba.ar">gcorti@dm.uba.ar</a>
Hélène Esnault	Freie Universität Berlin, Germany <a href="mailto:liveesnault@math.fu-berlin.de">liveesnault@math.fu-berlin.de</a>
Eric Friedlander	University of Southern California, USA <a href="mailto:ericmf@usc.edu">ericmf@usc.edu</a>
Max Karoubi	Institut de Mathématiques de Jussieu – Paris Rive Gauche, France <a href="mailto:max.karoubi@imj-prg.fr">max.karoubi@imj-prg.fr</a>
Moritz Kerz	Universität Regensburg, Germany <a href="mailto:moritz.kerz@mathematik.uni-regensburg.de">moritz.kerz@mathematik.uni-regensburg.de</a>
Huaxin Lin	University of Oregon, USA <a href="mailto:livehlin@uoregon.edu">livehlin@uoregon.edu</a>
Alexander Merkurjev	University of California, Los Angeles, USA <a href="mailto:merkurev@math.ucla.edu">merkurev@math.ucla.edu</a>
Birgit Richter	Universität Hamburg, Germany <a href="mailto:birgit.richter@uni-hamburg.de">birgit.richter@uni-hamburg.de</a>
Jonathan Rosenberg	(Managing Editor) University of Maryland, USA <a href="mailto:jmr@math.umd.edu">jmr@math.umd.edu</a>
Marco Schlichting	University of Warwick, UK <a href="mailto:schlichting@warwick.ac.uk">schlichting@warwick.ac.uk</a>
Charles Weibel	(Managing Editor) Rutgers University, USA <a href="mailto:weibel@math.rutgers.edu">weibel@math.rutgers.edu</a>
Guoliang Yu	Texas A&M University, USA <a href="mailto:guoliangyu@math.tamu.edu">guoliangyu@math.tamu.edu</a>

## PRODUCTION

Silvio Levy (Scientific Editor)  
[production@msp.org](mailto:production@msp.org)

Annals of K-Theory is a journal of the [K-Theory Foundation](http://ktheoryfoundation.org) ([ktheoryfoundation.org](http://ktheoryfoundation.org)). The K-Theory Foundation acknowledges the precious support of [Foundation Compositio Mathematica](http://foundationcompositio.com), whose help has been instrumental in the launch of the Annals of K-Theory.

---

See inside back cover or [msp.org/akt](http://msp.org/akt) for submission instructions.

---

The subscription price for 2019 is US \$490/year for the electronic version, and \$550/year (+\$25, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

---

Annals of K-Theory (ISSN 2379-1681 electronic, 2379-1683 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

AKT peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing  
<http://msp.org/>

© 2019 Mathematical Sciences Publishers

# ANNALS OF K-THEORY

2019

vol. 4

no. 4

- On the intersection motive of certain Shimura varieties: the case of Siegel threefolds 525  
JÖRG WILDESHAUS
- A Baum–Connes conjecture for singular foliations 561  
IAKOVOS ANDROULIDAKIS and GEORGES SKANDALIS
- Witt groups of abelian categories and perverse sheaves 621  
JÖRG SCHÜRMAN and JON WOOLF
- A generalized Vaserstein symbol 671  
TARIQ SYED
- A Dolbeault–Hilbert complex for a variety with isolated singular points 707  
JOHN LOTT