

Algebra & Number Theory

Volume 13

2019

No. 3

Fundamental gerbes

Niels Borne and Angelo Vistoli



Fundamental gerbes

Niels Borne and Angelo Vistoli

For a class of affine algebraic groups \mathcal{C} over a field κ , we define the notion of \mathcal{C} -fundamental gerbe of a fibered category, generalizing what we did for finite group schemes in a 2015 paper.

We give necessary and sufficient conditions on \mathcal{C} implying that a fibered category X over κ satisfying mild hypotheses admits a Nori \mathcal{C} -fundamental gerbe. We also give a tannakian interpretation of the gerbe that results by taking as \mathcal{C} the class of virtually unipotent group schemes, under a properness condition on X .

Finally, we prove a general duality result, generalizing the duality between group schemes of multiplicative type and Galois modules, that yields a construction of the multiplicative gerbe of multiplicative type which is independent of the previous theory, and requires weaker hypotheses. This gives a conceptual interpretation of the universal torsor of Colliot-Thélène and Sansuc.

1. Introduction

Previous work. Let X be a reduced proper connected scheme over a field κ , with a rational point $x_0 \in X(\kappa)$. The celebrated result of Nori [1982] says the following:

- (1) There is a profinite group scheme $\pi(X, x_0)$, the *Nori fundamental group scheme*, with a $\pi(X, x_0)$ -torsor $P \rightarrow X$ with a trivialization $P|_{x_0} \simeq \pi(X, x_0)$ such that for every profinite group scheme $G \rightarrow \text{Spec } \kappa$ and every G torsor $Q \rightarrow X$ with a trivialization $\alpha: Q|_{x_0} \simeq G$, there is a unique homomorphism of group schemes $\pi(X, x_0) \rightarrow G$ inducing Q and α .
- (2) There is an equivalence of tannakian categories between representations of the group scheme $\pi(X, x_0)$ and essentially finite locally free sheaves on X .

In our paper [Borne and Vistoli 2015] we extend this result in three ways:

- (1) We relax greatly the hypotheses on X .
- (2) We remove the dependence on the base point, which does not even need to exist, by replacing the fundamental group scheme $\pi(X, x_0)$ with a *fundamental gerbe* $\Pi_{X/\kappa}^N$.
- (3) We give a more general definition of essentially finite locally free sheaf on X .

Borne was supported in part by the Labex CEMPI (ANR-11-LABX-0007-01) and Anr ARIVAF (ANR-10-JCJC 0107). Vistoli was supported in part by research funds from the Scuola Normale Superiore.

MSC2010: primary 14A20; secondary 14H30.

Keywords: fundamental group scheme, Tannaka theory, algebraic stacks.

The fundamental gerbe $\Pi_{X/\kappa}^N$ of a category X fibered in groupoids over the category (Aff/κ) of affine schemes over a fixed base field κ is a profinite gerbe with a morphism $X \rightarrow \Pi_{X/\kappa}^N$ that is universal among morphisms from X to a profinite gerbe.

Nori [1982] also defines a *unipotent fundamental group* scheme $\pi_1^U(X, x_0)$; it is a prounipotent group scheme with a $\pi_1^U(X, x_0)$ -torsor $P \rightarrow X$ that satisfies the analogue of the universal property above for torsors under prounipotent group schemes.

The motivating question. It is a natural question whether one can define a universal prounipotent gerbe $X \rightarrow \Pi_{X/\kappa}^U$.

More generally, suppose that we are given a class \mathcal{C} of affine algebraic groups of finite type defined over extensions of κ , satisfying some natural stability conditions, listed in Definition 5.1. Then one defines a *\mathcal{C} -gerbe over κ* as an affine fpqc gerbe $\Gamma \rightarrow (\text{Aff}/\kappa)$ such that for every extension ℓ of κ and every object ξ of $\Gamma(\ell)$, the group scheme $\underline{\text{Aut}}_\ell \xi$ of automorphisms of ξ is in $\mathcal{C}(\ell)$. A pro- \mathcal{C} -gerbe is a gerbe that is a projective limit of \mathcal{C} -gerbes. If X is a fibered category, we define a *\mathcal{C} -fundamental gerbe* as a pro- \mathcal{C} gerbe $\Pi_{X/\kappa}^\mathcal{C}$ with a morphism $X \rightarrow \Pi_{X/\kappa}^\mathcal{C}$ which is universal among all maps from X to a pro- \mathcal{C} -gerbe. If $x_0 \in X(\kappa)$, ξ is the image of x_0 in $\Pi_{X/\kappa}^\mathcal{C}$, and $\underline{\text{Aut}}_\kappa \xi$ is the automorphism group scheme of ξ over κ , then there exists an $\underline{\text{Aut}}_\kappa \xi$ -torsor $P \rightarrow X$ satisfying the analogue of the universal property above for torsors under projective limits of group schemes in $\mathcal{C}(\kappa)$. (See Section 5 for the rigorous definitions.)

In this paper we answer the following question: under what conditions on \mathcal{C} does $\Pi_{X/\kappa}^\mathcal{C}$ exist for a reasonably large class of fibered categories?

It is certainly not the case that it exists in general. For example, one can show that if \mathcal{C} contains the semidirect product $\mathbb{G}_m \ltimes \mathbb{G}_a$, then $\Pi_{X/\kappa}^\mathcal{C}$ does not exist every time X is a scheme with a line bundle with a nonzero section that vanishes somewhere (Example 5.13).

Existence results for fundamental gerbes. We characterize the classes \mathcal{C} for which $\Pi_{X/\kappa}^\mathcal{C}$ exists for reasonable general X . If G is an affine group scheme of finite type over a field k , we say that G is *virtually nilpotent* if, after passing to the algebraic closure of k , the group G contains a nilpotent subgroup scheme of finite index. *Virtually unipotent* and *virtually abelian* group schemes are defined similarly.

We say that a class \mathcal{C} is *well-founded* when it consists of virtually nilpotent group schemes. Our main examples of well-founded classes are those of virtually abelian and virtually unipotent affine group schemes of finite type.

Our first main result, Theorem 7.1, states that if X satisfies a mild finiteness condition and is geometrically reduced in the sense of Definition 4.3, and $H^0(X, \mathcal{O}) = \kappa$, then $\Pi_{X/\kappa}^\mathcal{C}$ exists for every well-founded class \mathcal{C} . For schemes, the finiteness condition is equivalent to being quasicompact and quasiseparated.

In fact, the condition that the class \mathcal{C} be well-founded turns out to be also necessary (Remark 7.4). In other words, as soon as we admit a group in our class \mathcal{C} that is not virtually nilpotent, then fundamental gerbes $\Pi_{X/\kappa}^\mathcal{C}$ do not exist anymore for a wide class of quasiprojective schemes X satisfying the conditions above.

The proof of [Theorem 7.1](#) is very similar in structure with that of the existence of the fundamental gerbe in [\[Borne and Vistoli 2015\]](#).

There are many examples of well-founded classes, and, correspondingly, many fundamental gerbes, and fundamental group schemes, that one can associate with a fibered category as above. Here are some examples:

- (1) the Nori fundamental gerbe $\Pi_{X/\kappa}^N$, associated with the class of finite group schemes,
- (2) the unipotent fundamental gerbe $\Pi_{X/\kappa}^U$,
- (3) the virtually unipotent fundamental gerbe $\Pi_{X/\kappa}^{VU}$,
- (4) the abelian fundamental gerbe $\Pi_{X/\kappa}^A$,
- (5) the virtually abelian fundamental gerbe $\Pi_{X/\kappa}^{VA}$,
- (6) the fundamental gerbe of multiplicative type $\Pi_{X/\kappa}^{MT}$, associated with the class of group schemes of multiplicative type,
- (7) the virtually nilpotent fundamental gerbe $\Pi_{X/\kappa}^{VN}$; since, by definition, a well-founded class is contained in the class of virtually nilpotent groups, and fundamental gerbes are functorial under inclusion of classes (see [Section 8](#)), the virtually nilpotent group fundamental gerbe $\Pi_{X/\kappa}^{VN}$ dominates all the other $\Pi_{X/\kappa}^{\mathcal{C}}$ (we can call it the one gerbe, in analogy with Tolkien's one ring).

The tannakian interpretations. Of course one would like to have a tannakian interpretation for each of the fundamental gerbes above.

If Γ is an affine gerbe over κ , we denote by $\text{Rep } \Gamma$ the corresponding tannakian category. A morphism $X \rightarrow \Gamma$ induces a pullback $\text{Rep } \Gamma \rightarrow \text{Vect}_X$, where we denote by Vect_X the category of locally free sheaves on X ; in particular for every fibered category satisfying the conditions of [Theorem 7.1](#) and every well-founded class \mathcal{C} , we obtain a functor $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \text{Vect}_X$. The pullbacks $\text{Rep } \Pi_{X/\kappa}^{VN} \rightarrow \text{Vect}_X$, $\text{Rep } \Pi_{X/\kappa}^A \rightarrow \text{Vect}_X$ and $\text{Rep } \Pi_{X/\kappa}^{VA} \rightarrow \text{Vect}_X$ are almost never fully faithful, and we are not able to give a nontautological tannakian interpretation of $\text{Rep } \Pi_{X/\kappa}^A$ and $\text{Rep } \Pi_{X/\kappa}^{VA}$.

In contrast with this, we have that if \mathcal{C} is a well-founded subclass of the class of virtually unipotent group schemes, the pullback $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \text{Vect}_X$ is fully faithful ([Corollary 10.8](#)). In particular, the pullbacks $\text{Rep } \Pi_{X/\kappa}^N \rightarrow \text{Vect}_X$, $\text{Rep } \Pi_{X/\kappa}^U \rightarrow \text{Vect}_X$ and $\text{Rep } \Pi_{X/\kappa}^{VU} \rightarrow \text{Vect}_X$ are fully faithful.

The pullback $\text{Rep } \Pi_{X/\kappa}^N \rightarrow \text{Vect}_X$ induces an equivalence between $\text{Rep } \Pi_{X/\kappa}^N$ and the category of essentially finite bundles on X : this is proved in [\[Borne and Vistoli 2015\]](#).

The pullback $\text{Rep } \Pi_{X/\kappa}^U \rightarrow \text{Vect}_X$ induces an equivalence of $\text{Rep } \Pi_{X/\kappa}^U$ with the class of locally free sheaves that are obtained from successive extensions from trivial bundles ([Theorem 10.5\(1\)](#)). This is a generalization of the tannakian characterization of the unipotent fundamental group scheme due to Nori [\[1982\]](#), and is not at all surprising.

The tannakian interpretation of the virtually unipotent gerbe $\Pi_{X/\kappa}^{VU}$ is somewhat more interesting. S. Otabe [\[2017\]](#) defined *semifinite bundles*: these are locally free sheaves that are obtained as successive

extensions of essentially finite bundles; see [Definition 10.2](#). Our terminology is different, as we call these *extended essentially finite locally free sheaves*.

If $\text{char } \kappa = 0$ we show that the pullback $\text{Rep } \Pi_{X/\kappa}^{\text{VU}} \rightarrow \text{Vect}_X$ gives an equivalence between $\text{Rep } \Pi_{X/\kappa}^{\text{VU}}$ and the category of extended essentially finite locally free sheaves on X ([Theorem 10.5\(2\)](#)). If $\text{char } \kappa > 0$, then $\text{Rep } \Pi_{X/\kappa}^{\text{VU}}$ is equivalent to the category of locally free sheaves that become extended essential finite bundles after pullback by a sufficiently high power of the absolute Frobenius ([Theorem 10.7](#)).

We conclude with a reference to a result, [Theorem 10.9](#), due to Tonini and Zhang. Assume that $\text{char } \kappa > 0$, that X is a pseudoproper geometrically reduced algebraic stack of finite type over κ , and that $H^1(X, E)$ is a finite-dimensional vector space over κ for all locally free sheaves on X . Then $\Pi_{X/\kappa}^{\text{VU}} = \Pi_{X/\kappa}^{\text{SVU}} = \Pi_{X/\kappa}^{\text{N}}$.

Fundamental gerbes of multiplicative type. A particularly interesting fundamental gerbe is the fundamental gerbe $\Pi_{X/\kappa}^{\text{MT}}$ of multiplicative type, as it gives a conceptual interpretation of the universal torsor of Colliot-Thélène and Sansuc [[1976](#); [1987](#)]. Suppose that X is a projective variety over a field κ such that if κ^{sep} is a separable closure of κ , then $\text{Pic}(X_{\kappa^{\text{sep}}})$ is a free abelian group of finite rank, and $x_0 \in X(\kappa)$. Then Colliot-Thélène and Sansuc define a G -torsor on X , where G is the torus associated with the action of the Galois group on $\text{Pic}(X_{\kappa^{\text{sep}}})$. In our language, G is the fundamental group of multiplicative type of (X, x_0) .

In the last section we give a direct construction of $\Pi_{X/\kappa}^{\text{MT}}$, generalizing that of Colliot-Thélène and Sansuc, which is completely independent of the general machinery in the rest of the paper; this works even for fibered categories X satisfying the same mild finiteness condition, with $H^0(X, \mathcal{O}) = \kappa$, but without assuming that X is geometrically reduced. We construct $\Pi_{X/\kappa}^{\text{MT}}$ from the Picard stack $\underline{\text{Pic}}_X$ over the small étale site $\kappa_{\text{ét}}$, which sends every étale κ -algebra A into the groupoid $\underline{\text{Pic}}(X_A)$ of invertible sheaves over X_A ([Theorem 13.11](#)).

Along the way, we prove a very general duality theorem for gerbes of multiplicative type, which extends the well-known duality between groups of multiplicative type and sheaves of abelian groups on $\kappa_{\text{ét}}$ (or, equivalently, abelian groups with a continuous action of the Galois group of $\kappa^{\text{sep}}/\kappa$). More precisely, we prove an equivalence of 2-categories between gerbes of multiplicative type and a certain class of Picard stacks with additional structure ([Theorem 13.10](#)). See also [[Brochard 2014](#); [Braverman and Bezrukavnikov 2007](#), §2.4] for related ideas.

(In this part we do not give the full details of all the proofs, as said details tend to be rather tedious.)

Description of content. The first three sections of the paper aim at fixing the notation, and present some facts about affine gerbes and fibered categories which are undoubtedly known to the experts, but for which we could not find a suitable treatment in the literature.

The real action starts in [Section 5](#), in which we give the general definition of a \mathcal{C} -fundamental gerbe, explain the connection of this with the notion of \mathcal{C} -fundamental group, and give examples to show how fundamental gerbes don't exist in general.

[Section 6](#) contains the definition of a well-founded class, and several technical results on group scheme actions on affine varieties that lead to the characterization of well-founded classes given in [Theorem 6.12](#).

The first main result, the existence of $\Pi_{X/\kappa}^{\mathcal{C}}$ for a well-founded class \mathcal{C} , with appropriate hypotheses on X , is in [Section 7](#).

[Section 8](#) contains a small but very useful result on the relation between $\Pi_{X/\kappa}^{\mathcal{C}}$ and $\Pi_{X/\kappa}^{\mathcal{D}}$, when \mathcal{D} is a subclass of a well-founded class \mathcal{C} .

[Section 9](#) contains a base-change result for $\Pi_{X/\kappa}^{\mathcal{C}}$ under an algebraic extension of κ .

Our main results on the tannakian interpretation of certain fundamental gerbes, [Theorems 10.5 and 10.7](#) are stated in [Section 10](#), together with the result of Tonini and Zhang, [Theorem 10.9](#), mentioned above. In [Section 11](#) we put the problem of giving a tannakian interpretation of fundamental gerbes for a certain fundamental class into a more general framework, and we prove a more general result ([Theorem 11.9](#)) that implies [Theorem 10.5](#). [Section 12](#) contains the proof of [Theorem 10.7](#).

The last section contains our treatment of the duality theorem for gerbes of multiplicative type, and our alternative construction for the universal gerbe of multiplicative type.

2. Notations and conventions

We will fix a base field κ . All schemes and morphisms will be defined over κ . All fibered categories will be fibered in groupoids over the category (Aff/κ) of affine κ -schemes (or, equivalently, over the opposite of the category of κ -algebras). A base-preserving functor between categories fibered in groupoids will be referred to in short as a map, or a morphism. A κ -scheme U will be identified with the category fibered in sets $(\text{Aff}/U) \rightarrow (\text{Aff}/\kappa)$, where (Aff/U) is the category of maps $T \rightarrow U$, where T is an affine scheme.

All group schemes will be affine over extensions ℓ of κ . If G is a group scheme of finite type over ℓ , we will denote by G^0 the connected component of the identity. If ℓ is perfect, G_{red}^0 is a smooth connected subgroup scheme of G .

If a group scheme G over an extension ℓ of κ acts on an ℓ -scheme $X = \text{Spec } A$, we denote by $X//G$ the spectrum of the ℓ -algebra of invariants A^G . We will need the following standard fact, which is, for example, a particular case of Grothendieck's result on the existence of quotients for finite flat groupoids; see [\[SGA 3_I 1970\]](#).

Lemma 2.1. *Assume that ℓ is algebraically closed and G is finite over ℓ . Then the fibers of the function $X(\ell) \rightarrow (X//G)(\ell)$ are precisely the orbits of the action of $G(\ell)$ on $X(\ell)$.*

Let $G \rightarrow \text{Spec } \ell$ be a group scheme and $P \rightarrow \text{Spec } \ell$ a G -torsor. We can use the conjugation action of G and P to define a twisted form of G , which we call, as usual, an *inner form* of G .

3. Generalities on affine gerbes

By *affine gerbe* we will always mean affine fpqc gerbe over the base field κ , that is, an fpqc gerbe over (Aff/κ) with affine diagonal, possessing an affine chart. These admit an obvious description in terms of groupoids, see [\[Borne and Vistoli 2015, §3\]](#), and are called *tannakian gerbes* in [\[Saavedra Rivano 1972, Chapitre III, §2\]](#).

We will denote by $\text{Rep } \Gamma$ the category of vector bundles of finite rank on Γ . This is an example of a *tannakian category* over κ , which by definition is a κ -linear rigid abelian tensor category such that $\text{End}(\mathbf{1}) = \kappa$, that admits moreover a fiber functor over some extension of κ . By [Deligne 1990], the association $\Gamma \mapsto \text{Rep } \Gamma$ establishes a correspondence between affine gerbes and tannakian categories.

We will often consider gerbes of finite type, which by definition are those satisfying the equivalent conditions of the following proposition.

Proposition 3.1. *Let $\Gamma \rightarrow \text{Spec } \kappa$ be an affine gerbe. Then the following are equivalent:*

- (1) Γ is a smooth algebraic stack over κ .
- (2) Γ is an algebraic stack of finite type over κ .
- (3) Γ is an algebraic stack.
- (4) The diagonal of Γ is of finite type.
- (5) If ℓ is an extension of κ and $\xi \in \Gamma(\ell)$, then $\underline{\text{Aut}}_\ell \xi$ is of finite type over ℓ .
- (6) There exists an extension ℓ of κ and an object $\xi \in \Gamma(\ell)$ such that $\underline{\text{Aut}}_\ell \xi$ is of finite type over ℓ .
- (7) If $\{A_i\}_{i \in I}$ is an inductive system of κ -algebras, the natural map

$$\varinjlim_i \Gamma(A_i) \longrightarrow \Gamma(\varinjlim_i A_i)$$

is an equivalence of categories.

- (8) The tannakian category $\text{Rep } \Gamma$ is finitely generated.

Proof. The implications (1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (4) \Rightarrow (5) \Rightarrow (6) are obvious.

The proof of (3) \Rightarrow (1) is given for fppf gerbes in [Bergh 2017, Proposition A.2]; it is also valid for fpqc gerbes.

Here is a sketch of proof that (6) implies (7). Set $G := \underline{\text{Aut}}_\ell \xi$; then $\Gamma_\ell = \mathcal{B}_\ell G$ is an algebraic stack, as it follows from Artin’s theorem [Laumon and Moret-Bailly 2000, Théorème 10.1]. Since (3) \Rightarrow (2) holds, Γ_ℓ is an algebraic stack of finite type over ℓ , or equivalently, of finite presentation. Hence Γ_ℓ preserves filtered colimits; see [Stacks 2005–, Tag 0123]. Set $R_0 := \ell$, $R_1 := \ell \otimes_\kappa \ell$, and $R_2 := \ell \otimes_\kappa \ell \otimes_\kappa \ell$; the natural maps $\iota_1, \iota_2: R_0 \rightarrow R_1$ and ι_{12}, ι_{13} and $\iota_{23}: R_1 \rightarrow R_2$ induce functors $\iota_{1*}, \iota_{2*}: \Gamma(A \otimes_\kappa R_0) \rightarrow \Gamma(A \otimes_\kappa R_1)$ and ι_{12*}, ι_{13*} and $\iota_{23*}: \Gamma(A \otimes_\kappa R_1) \rightarrow \Gamma(A \otimes_\kappa R_2)$ for each κ -algebra A . We call $\Delta \rightarrow (\text{Aff}/\kappa)$ the fibered category of objects of Γ with descent data along the covering $\text{Spec } \ell \rightarrow \text{Spec } \kappa$; if A is a κ -algebra, the objects of $\Delta(A)$ are pairs (ξ, a) , where ξ is an object of $\Gamma(A \otimes_\kappa R_0)$ and a is an isomorphism $a: \iota_{2*}\xi \simeq \iota_{1*}\xi$ satisfying $\iota_{12*}a \circ \iota_{23*}a = \iota_{13*}a$. An arrow $f: (\xi, a) \rightarrow (\eta, b)$ in $\Delta(A)$ is an arrow $f: \xi \rightarrow \eta$ in $\Gamma(A \otimes_\kappa R_0)$, with the property that $b \circ \iota_{2*}f = \iota_{1*}f \circ a$ in $\text{Hom}_{\Gamma(R_1 \otimes_\kappa A)}(\iota_{2*}\xi, \iota_{1*}\eta)$. So the diagram

$$\text{Hom}_{\Delta(A)}((\xi, a), (\eta, b)) \longrightarrow \text{Hom}_{\Gamma(A \otimes_\kappa R_0)}(\xi, \eta) \begin{array}{c} \xrightarrow{f \mapsto b \circ \iota_{2*}f} \\ \xrightarrow{f \mapsto \iota_{1*}f \circ a} \end{array} \text{Hom}_{\Gamma(A \otimes_\kappa R_1)}(\iota_{2*}\xi, \iota_{1*}\eta)$$

is an equalizer.

The obvious functor $\Gamma(A) \rightarrow \Delta(A)$ is an equivalence, because Δ is an fpqc stack. Hence it is enough to prove that for any inductive system of κ -algebras $\{A_i\}$ the functor $\varinjlim_i \Delta(A_i) \rightarrow \Delta(\varinjlim_i A_i)$ is an equivalence.

Let us show that $\varinjlim_i \Delta(A_i) \rightarrow \Delta(\varinjlim_i A_i)$ is fully faithful. For this, notice that if R is an ℓ -algebra, and A a κ -algebra, then the fibered category sending A into $\Gamma(R \otimes_\kappa A) = \Gamma_\ell(R \otimes_\ell (\ell \otimes_\kappa A))$ preserves filtered colimits, because Γ_ℓ does, and tensor products preserve colimits.

Set $A := \varinjlim_i A_i$; we need to prove that the functor $\varinjlim_i \Delta(A_i) \rightarrow \Delta(\varinjlim_i A_i)$ is fully faithful. Take two objects $\{(\xi_i, a_i)\}$ and $\{(\eta_i, b_i)\}$ of $\varinjlim_i \Delta(A_i)$; call (ξ, a) and (η, b) their images in $\Delta(A)$. By definition we have

$$\mathrm{Hom}_{\varinjlim_i \Delta(A_i)}(\{(\xi_i, a_i)\}, \{(\eta_i, b_i)\}) = \varinjlim_i \mathrm{Hom}_{\Delta(A_i)}((\xi_i, a_i), (\eta_i, b_i)).$$

Since filtered colimits preserve equalizers, we have a commutative diagram

$$\begin{array}{ccccc} \varinjlim \mathrm{Hom}((\xi_i, a_i), (\eta_i, b_i)) & \longrightarrow & \varinjlim \mathrm{Hom}(\xi_i, \eta_i) & \rightrightarrows & \varinjlim \mathrm{Hom}(\iota_{2*}\xi_i, \iota_{1*}\eta_i) \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{Hom}((\xi, a), (\eta, b)) & \longrightarrow & \mathrm{Hom}(\xi, \eta) & \rightrightarrows & \mathrm{Hom}(\iota_{2*}\xi, \iota_{1*}\eta) \end{array}$$

in which the rows are equalizers, and the last two columns are bijections. It follows that the first column is also a bijection, which is exactly what we want to show.

The proof of the fact that the functor $\varinjlim_i \Delta(A_i) \rightarrow \Delta(\varinjlim_i A_i)$ essentially surjective is easy, and left to the reader.

It is easy to check that (7) implies (4): this follows from the well-known fact, due to Grothendieck, that an affine scheme over a ring R is finitely presented if and only the functor on R -algebras that it represents preserves inductive limits.

Let us check the stronger result that (7) implies (3). Let ℓ be an extension of κ such that $\Gamma(\ell) \neq \emptyset$. If $\{A_i\}$ is the inductive system of κ -subalgebras of ℓ of finite type over κ ; then $\varinjlim_i A_i = \ell$, so $\varinjlim_i \Gamma(A_i) \simeq \Gamma(\varinjlim_i A_i) \neq \emptyset$. Hence $\Gamma(A_i) \neq \emptyset$ for some i ; by passing to a quotient by a maximal ideal of A_i we see that there is a finite extension k/κ such that $\Gamma(k) \neq \emptyset$. If $\xi \in \Gamma(k)$ and $G := \underline{\mathrm{Aut}}_k \xi$, then G is of finite type over k , because (4) is satisfied; so the map $\mathrm{Spec} k \rightarrow \Gamma_k$ is an fppf cover, and hence the composite $\mathrm{Spec} k \rightarrow \Gamma_k \rightarrow \Gamma$ is an fppf cover. From Artin's theorem [Laumon and Moret-Bailly 2000, Théorème 10.1] we see that Γ is an algebraic stack, as claimed.

The equivalence between (8) and (4) is proved in [Saavedra Rivano 1972, III 3.3.1.1]. □

We will mainly need the following corollary.

Corollary 3.2. *Let Γ be an affine gerbe of finite type over κ . Then there exists a finite separable extension κ'/κ such that $\Gamma(\kappa') \neq \emptyset$.*

Proof. This follows from Proposition 3.1(1). □

In this paper we will use repeatedly the following lemma.

Lemma 3.3. *Let $\phi' : G' \rightarrow G$ and $\phi'' : G'' \rightarrow G$ be homomorphisms of algebraic groups over κ . Then we have an equivalence of fibered categories*

$$\mathcal{B}_\kappa G' \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa G'' \simeq [G/(G' \times G'')],$$

where the action of $G' \times G''$ on G is defined by

$$g \cdot (g', g'') := \phi'(g')^{-1} g \phi''(g'').$$

Proof. An object of the fibered product $\mathcal{B}_\kappa G' \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa G''$ over a scheme T is a triple (P, P'', α) , where $P' \rightarrow T$ and $P'' \rightarrow T$ are, respectively, a G' -torsor and a G'' -torsor, and $\alpha : P' \times^{G'} G \simeq P'' \times^{G''} G$ is an isomorphism of G -torsors. Set $Q := P' \times^{G'} G$, and consider the usual isomorphism $\rho : Q \times G \rightarrow Q \times_T Q$ defined by $(q, g) \rightarrow (q, qg)$; denote by $\pi : Q \times_T Q \rightarrow G$ the composite of ρ^{-1} with the projection $Q \times G \rightarrow G$. Let $u' : P' \rightarrow Q$ be the usual ϕ' -equivariant morphism, and call $u'' : P'' \rightarrow Q$ the composite of the ϕ'' -equivariant morphism $P'' \rightarrow P'' \times^{G''} G$ with α^{-1} .

The composite

$$P' \times_T P'' \xrightarrow{u' \times u''} Q \times_T Q \xrightarrow{\pi} G$$

is easily seen to be $(G' \times G'')$ -equivariant, when the action of $G' \times G''$ on G is the one described above. Of course $P' \times_T P''$ is a $(G' \times G'')$ -torsor.

This defines a base-preserving function from the objects of $\mathcal{B}_\kappa G' \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa G''$ to those of $[G/(G' \times G'')]$; this is immediately seen to extend to a base-preserving functor $\mathcal{B}_\kappa G' \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa G'' \rightarrow [G/(G' \times G'')]$.

To go in the opposite direction, let $P \rightarrow T$ be a $(G' \times G'')$ -torsor and $\theta : P \rightarrow G$ a $(G' \times G'')$ -equivariant morphism. If $P' \rightarrow T$ and $P'' \rightarrow T$ are G' and G'' -torsors associated with P , we have a canonical isomorphism $P \simeq P' \times_T P''$; so we get a $(G' \times G'')$ -equivariant morphism $\theta : P' \times_T P'' \rightarrow G$. From a section $p' \in P'(T)$ we obtain a ϕ'' -equivariant morphism $\theta_{p'} : P'' \rightarrow G$, which in turn yields a G -equivariant morphism $P'' \times^{G''} G \rightarrow G$, which gives a section of $(P'' \times^{G''} G)(T)$. Sending p' into $\theta_{p'}$ gives a ϕ' -equivariant morphism $P' \rightarrow P'' \times^{G''} G$, which extends to an isomorphism of G -torsors $P' \times^{G'} G \simeq P'' \times^{G''} G$. This yields a base-preserving functor $[G/(G' \times G'')] \rightarrow \mathcal{B}_\kappa G' \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa G''$, which is a quasi-inverse to the one above. □

Let $f : \Gamma \rightarrow \Delta$ be a morphism of affine gerbes. Then f is faithful if and only if for some extension ℓ of κ , and some object ξ of $\Delta(\kappa)$, the induced homomorphism of group scheme $\underline{\text{Aut}}_\ell \xi \rightarrow \underline{\text{Aut}}_\ell f(\xi)$ is a monomorphism. Hence, a homomorphism of group schemes $G \rightarrow H$ induces a faithful morphism $\mathcal{B}_\kappa G \rightarrow \mathcal{B}_\kappa H$ if and only if $G \rightarrow H$ is a monomorphism.

If Γ and Δ are of finite type, then f is faithful if and only if it is representable.

Definition 3.4. Let $f : \Gamma \rightarrow \Delta$ be a morphism of affine gerbes over κ . We say that f is *locally full* if for any extension ℓ of κ and any object ξ of $\Gamma(\ell)$, the induced homomorphism of group schemes $\underline{\text{Aut}}_\ell \xi \rightarrow \underline{\text{Aut}}_\ell f(\xi)$ is faithfully flat.


Remark 3.5. If this is true for an extension ℓ and an object ξ of $\Gamma(\ell)$, then it is true for all ℓ and all ξ .

Remark 3.6. If $\phi: G \rightarrow H$ is a homomorphism of affine group schemes, the corresponding morphism $\mathcal{B}_\kappa G \rightarrow \mathcal{B}_\kappa H$ is locally full if and only if ϕ is faithfully flat, or, equivalently, an fpqc cover.

Remark 3.7. A morphism of affine gerbes that is both faithful and locally full is in fact an equivalence.

Definition 3.8. Let $f: \Gamma \rightarrow \Delta$ be a morphism of affine gerbes. A *canonical factorization* of f consists of a factorization $\Gamma \rightarrow \Delta' \rightarrow \Delta$ of f such that $\Gamma \rightarrow \Delta'$ is locally full, and $\Delta' \rightarrow \Delta$ is faithful.

Proposition 3.9. *A morphism of affine gerbes $f: \Gamma \rightarrow \Delta$ has a canonical factorization. Furthermore, if $\Gamma \rightarrow \Delta' \rightarrow \Delta$ and $\Gamma \rightarrow \Delta'' \rightarrow \Delta$ are two canonical factorizations, there exists an equivalence $\phi: \Delta' \rightarrow \Delta''$, and a commutative diagram*



Sketch of proof. For a tannakian proof, see [Tonini and Zhang 2017, Proposition B.4]; here is a direct approach. Suppose that we are given canonical factorization $\Gamma \xrightarrow{g} \Delta' \xrightarrow{h} \Delta$. For each κ -algebra and each $\xi \in \Gamma(A)$, call $K_A(\xi)$ the kernel of the homomorphism of group schemes $\underline{\text{Aut}}_A(\xi) \rightarrow \underline{\text{Aut}}_A(f(\xi))$. If $\xi, \eta \in \Gamma(A)$, then $\underline{\text{Hom}}_A(\xi, \eta)$ is a torsor over $\text{Spec } A$ for the group $\underline{\text{Aut}}_A(\xi)$; thus, by restriction we obtain a free action of $K_A(\xi)$ on $\underline{\text{Hom}}_A(\xi, \eta)$. From the definition of Δ' it follows that the map $\underline{\text{Hom}}_A(\xi, \eta) \rightarrow \underline{\text{Hom}}_A(f(\xi), f(\eta))$ induces an isomorphism of fpqc sheaves

$$\underline{\text{Hom}}_A(\xi, \eta)/K_A(\xi) \simeq \underline{\text{Hom}}_A(f(\xi), f(\eta)).$$

Hence Δ' is the fpqc stackification of the prestack whose objects are the objects of Γ , and whose arrows $\xi \rightarrow \eta$ over a fixed A are sections over $\text{Spec } A$ of the sheaf of sets $\underline{\text{Hom}}_A(\xi, \eta)/K_A(\xi)$. In other words, Δ' is the rigidification of Γ along K_A , as defined in [Abramovich et al. 2008, Appendix A]. This shows the uniqueness of Δ' .

For the existence, we prefer not to use the dubious notion of fpqc stackification, and do the following. Let $U \rightarrow \Gamma$ be an fpqc cover by an affine scheme (for this is it enough that U is a nonempty scheme, for example, the spectrum of a field). Then if we set $R := U \times_\Gamma U$ we get an fpqc groupoid $R \rightrightarrows U$. Set $U \times U = \text{Spec } A$, and call ξ and η the objects of $\Gamma(A)$ corresponding to the composites of the given morphism $U \rightarrow \Gamma$ with the two projections $U \times U \rightarrow U$; then R represents the functor $\underline{\text{Hom}}_A(\xi, \eta)$. Hence there is an action of $K_A(\xi)$ on R , leaving the morphism $R \rightarrow U \times U$ invariant. By passing to the fpqc quotient $R/K_A(\xi)$ we obtain a groupoid $R/K_A(\xi) \rightrightarrows U$, whose stack of torsors is the desired rigidification. There remains to prove that $R/K_A(\xi)$ is an affine scheme, as this shows that the rigidification is an affine gerbe, and ends the proof.

For this, let K be a field extension of κ and let ζ be an object of $\Gamma(K)$. Since Γ is an fpqc gerbe, there exists a faithfully flat extension $A \subseteq B$ such that $K \subseteq B$, such that ξ_B, η_B and ζ_B are all isomorphic. So

the pullback of R to $\text{Spec } B$ is isomorphic to the affine group scheme $\underline{\text{Aut}}_B(\zeta_B)$, and $(R/K_A(\xi))_B$ is the quotient $\underline{\text{Aut}}_B(\zeta_B)/K_B(\zeta_B)$, which is affine. Hence $R/K_A(\xi)$ is affine. \square

Proposition 3.10. *Let $f : \Gamma \rightarrow \Delta$ be a morphism of affine gerbes over κ . The following conditions are equivalent:*

- (1) *The morphism f is locally full.*
- (2) *If S is an affine scheme over κ and ξ and η are two objects of $\Gamma(S)$, the induced morphism of fpqc sheaves $\underline{\text{Isom}}_\ell(\xi, \eta) \rightarrow \underline{\text{Isom}}_\ell(f(\xi), f(\eta))$ is surjective.*
- (3) *The morphism f makes Γ into a relative gerbe over Δ .*
- (4) *If f factors as $\Gamma \rightarrow \Delta' \rightarrow \Delta$, where $\Delta' \rightarrow \Delta$ is a faithful homomorphism of affine gerbes, then $\Delta' \rightarrow \Delta$ is an equivalence.*
- (5) *The pullback homomorphism $\text{Rep } \Delta \rightarrow \text{Rep } \Gamma$ is fully faithful, and any subrepresentation of a representation of Γ in its essential image is also in the essential image.*

Proof. The equivalence between (1), (3) and (5) is also established in [Tonini and Zhang 2017, Proposition B.2(2)]. We give a complete proof for the convenience of the reader.

(2) \implies (1): It follows from the fact that a homomorphism of affine group schemes $G \rightarrow H$ is faithfully flat if and only if it is an fpqc cover.

(1) \implies (2): If $G := \underline{\text{Aut}}_S \xi$ and $H := \underline{\text{Aut}}_S f(\xi)$, then $\underline{\text{Isom}}_\ell(\xi, \eta)$ is a G -torsor and $\underline{\text{Isom}}_\ell(f(\xi), f(\eta))$ is an H -torsor. The map f induces a homomorphism $G \rightarrow H$ of group schemes over S . The map $\underline{\text{Isom}}_\ell(\xi, \eta) \rightarrow \underline{\text{Isom}}_\ell(f(\xi), f(\eta))$ is $(G \rightarrow H)$ -equivariant, so it is enough to show that $G \rightarrow H$ is an fpqc cover.

This follows from the definition if S is the spectrum of a field. In the general case, we may pass to an fpqc cover of S , and assume that there is a morphism $S \rightarrow \text{Spec } \ell$, where ℓ is an extension of κ , and an object ξ_0 of $\Gamma(\ell)$ whose pullback to S is isomorphic to ξ , so the general case follows from the case of a field.

(2) \iff (3): This is straightforward.

(1) \implies (3): We can extend the base field, and assume that $\Gamma = \mathcal{B}_\kappa G$, $\Delta = \mathcal{B}_\kappa H$, and f is induced by a surjective homomorphism of affine group schemes $\phi : G \rightarrow H$.

Then the factorization $\Gamma \rightarrow \Delta' \rightarrow \Delta$ corresponds to a factorization $G \rightarrow H' \rightarrow H$, where $H' \rightarrow H$ is a monomorphism. Since $G \rightarrow H$ is an epimorphism, it follows that $H' \rightarrow H$ is an isomorphism, so that $\Delta' \rightarrow \Delta$ is in fact an equivalence.

(3) \implies (1): Consider the canonical factorization $\Gamma \rightarrow \Delta' \rightarrow \Delta$. Since $\Delta' \rightarrow \Delta$ is faithful, by hypothesis this is an equivalence; hence f is locally full.

(1) \iff (5): See [Saavedra Rivano 1972, III 3.3.2.2]. \square

For the following we need the notion of cofiltered system of affine gerbes, and the projective limit of such a cofiltered system; for this we refer to [Borne and Vistoli 2015, Section 3].

Proposition 3.11. *Let $\{\Delta_i\}$ be a cofiltered system of affine gerbes, $\Gamma \rightarrow \varprojlim \Delta_i$ a morphism of affine gerbes. If each composite $\Gamma \rightarrow \varprojlim \Delta_i \rightarrow \Delta_i$ is locally full, then $\Gamma \rightarrow \varprojlim \Delta_i$ is also locally full.*

Proof. We will use the fact that a homomorphism of affine groups $G \rightarrow H$ over a field k is faithfully flat if and only if the corresponding homomorphism of Hopf algebras $k[H] \rightarrow k[G]$ is injective.

Let ξ be an object of $\Gamma(\ell)$, where ℓ is an extension of κ ; denote by η_i the image of ξ in Δ_i , and by η its image in $\varprojlim \Delta_i$. Set $G := \underline{\text{Aut}}_\ell \xi$, $H_i := \underline{\text{Aut}}_\ell \eta_i$, $H := \underline{\text{Aut}}_\ell \eta$. We need to show that the homomorphism $G \rightarrow H$ is faithfully flat, knowing that the composite $G \rightarrow H \rightarrow H_i$ is. But $\ell[H] = \varinjlim_i \ell[H_i]$; since every homomorphism $\ell[H_i] \rightarrow \ell[G]$ is injective, the conclusion follows. \square

4. Fibered categories

Let $p_X: X \rightarrow (\text{Aff}/\kappa)$ be a category fibered in groupoids.

We will consider X as a site with the fpqc topology inherited from (Aff/κ) : a collection $\{\xi_i \rightarrow \xi\}$ of arrows in X is an fpqc covering if the corresponding maps $p_X \xi_i \rightarrow p_X \xi$ are flat, and $p_X \xi$ is the union of a finite number of images of $p_X \xi_i$.

The fpqc sheaf $\mathcal{O} = \mathcal{O}_X$ sends each object ξ into $\mathcal{O}(U)$, where $U = p_X \xi$. We denote as usual by Vect_X the category of locally free sheaves on X . If κ'/κ is a field extension, denote by $X_{\kappa'}$ the fibered product $(\text{Aff}/\kappa') \times_{(\text{Aff}/\kappa)} X$. There exists an obvious homomorphism of κ -algebras $H^0(X, \mathcal{O}) \rightarrow H^0(X_{\kappa'}, \mathcal{O})$, inducing a homomorphism of κ' -algebras $H^0(X, \mathcal{O}) \otimes_{\kappa} \kappa' \rightarrow H^0(X_{\kappa'}, \mathcal{O})$.

Recall that a quasicompact and quasiseparated morphism, or scheme, or algebraic space, is nowadays called *concentrated*.

Definition 4.1. A fibered category $X \rightarrow (\text{Aff}/\kappa)$ is *concentrated* if there exists an affine scheme U and a representable concentrated faithfully flat morphism $U \rightarrow X$.

Notice that if $U \rightarrow X$ is as above, and we set $R := U \times_X U$, we obtain an fpqc groupoid $R \rightrightarrows U$ in algebraic spaces, in which U and R are concentrated. (If X is an fpqc stack, which we are not assuming, then X is equivalent to the stack of $(R \rightrightarrows U)$ -torsors in the fpqc topology.) From standard arguments in descent theory it follows that we have an exact sequence,

$$0 \longrightarrow H^0(X, \mathcal{O}) \longrightarrow H^0(U, \mathcal{O}) \longrightarrow H^0(R, \mathcal{O}).$$

From this we easily get the following.

Proposition 4.2. *Assume that X is concentrated. For any field extension κ'/κ , the base change homomorphism $H^0(X, \mathcal{O}) \otimes_{\kappa} \kappa' \rightarrow H^0(X_{\kappa'}, \mathcal{O})$ is an isomorphism.*

Definition 4.3. A fibered category X over (Aff/κ) is called *reduced* if every map from X to an algebraic stack Γ over κ factors through the reduced substack $\Gamma_{\text{red}} \subseteq \Gamma$.

It is *geometrically reduced* if the fibered category $X_{\kappa'} \rightarrow (\text{Aff}/\kappa')$ is reduced for any extension κ'/κ .

The following is straightforward.

Proposition 4.4. *Let X be a fibered category. Suppose that there exists a reduced scheme U and a representable faithfully flat map $U \rightarrow X$. Then X is reduced.*

So, for example, an affine gerbe X is reduced, because it has a representable faithfully flat map from the spectrum of a field [Borne and Vistoli 2015, Proposition 3.1(b)]. Since being an affine gerbe is a property that is stable under base change, an affine gerbe is in fact geometrically reduced.

Suppose that G is an affine group scheme over κ , and X is a fibered category. A G -torsor over X is a morphism of fibered categories $X \rightarrow \mathcal{B}_\kappa G$. Morphisms of G -torsors are, of course, base-preserving natural transformations. The resulting category of G -torsors on X will be denoted by $\text{Tors}_G(X)$; it is a groupoid. Of course, if X is a scheme then $\text{Tors}_G(X)$ is equivalent to the categories of classical G -torsors over X .

A homomorphism $\phi: G \rightarrow H$ of affine algebraic group schemes over κ yields a functor $\mathcal{B}_\kappa \phi: \mathcal{B}_\kappa G \rightarrow \mathcal{B}_\kappa H$, sending $E \rightarrow S$ into $E \times^G H \rightarrow S$. Composing with this functor gives a group change functor $\text{Tors}_G(X) \rightarrow \text{Tors}_H(X)$; the image of a torsor E will be denoted by $E \times^G H$.

If $x_0 \in X(\kappa)$, a *pointed torsor* over (X, x_0) will be a pair (E, e_0) , where $E: X \rightarrow \mathcal{B}_\kappa G$ is a G -torsor, and e_0 is a κ -rational point of the G -torsor $E(x_0)$. Pointed G -torsors over (X, x_0) form a category in the obvious way: a morphism $f: (E, e_0) \rightarrow (E', e'_0)$ is a base-preserving natural transformation $f: E \rightarrow E'$ such that $f_{x_0}(e_0) = e'_0$. This yields the groupoid $\text{Tors}_G(X, x_0)$ of pointed G -torsors.

If (E, e_0) is a pointed G -torsor on (X, x_0) and $\phi: G \rightarrow H$ is a homomorphism of affine group schemes, there is a natural map $E(x_0) \rightarrow E(x_0) \times^G H = (E \times^G H)(x_0)$; thus, taking the image of e_0 , the torsor $E \times^G H$ becomes a pointed torsor, which we denote by $(E, e_0) \times^G H$. This gives a group change functor $\text{Tors}_G(X, x_0) \rightarrow \text{Tors}_H(X, x_0)$.

5. Fundamental gerbes

Definition 5.1. Let \mathcal{C} be a class of affine group schemes of finite type over extensions ℓ of κ ; for each ℓ we denote by $\mathcal{C}(\ell)$ the class of group schemes over ℓ that are in \mathcal{C} . We say that \mathcal{C} is *stable* if the following conditions are satisfied:

- (1) Each $\mathcal{C}(\ell)$ is closed under isomorphism of group schemes over ℓ .
- (2) If ℓ is an extension of κ , ℓ' is an extension of ℓ , and G is a group scheme in $\mathcal{C}(\ell)$, then $G_{\ell'}$ is in $\mathcal{C}(\ell')$.
- (3) If G and H are in $\mathcal{C}(\ell)$, then $G \times_\ell H$ is also in $\mathcal{C}(\ell)$.
- (4) Suppose that G is in $\mathcal{C}(\ell)$ and H is an ℓ -subgroup scheme of G . Then H is in $\mathcal{C}(\ell)$.
- (5) Suppose that G is in $\mathcal{C}(\ell)$ and H is a normal ℓ -subgroup scheme of G . Then G/H is in $\mathcal{C}(\ell)$.
- (6) If G in $\mathcal{C}(\ell)$, every inner form of G is in $\mathcal{C}(\ell)$.

Definition 5.2. A stable class \mathcal{C} is said to be *very stable* if whenever ℓ is an extension of κ , ℓ' is a finite extension of ℓ , and G is an affine group scheme of finite type over ℓ , then G is in $\mathcal{C}(\ell)$ if and only if $G_{\ell'}$ is in $\mathcal{C}(\ell')$.

It is called *weakly very stable* if the same is true for all finite separable extensions ℓ'/ℓ .

Definition 5.3. Let \mathcal{C} be a stable class. A *pro- \mathcal{C} -group* over κ is a group scheme that is a projective limit of groups in $\mathcal{C}(\kappa)$.

Definition 5.4. Let \mathcal{C} be a stable class. A *\mathcal{C} -gerbe over κ* is an affine gerbe of finite type over κ such that for any object ξ in $\Gamma(\ell)$, where ℓ is an extension of κ , the group scheme $\underline{\text{Aut}}_\ell \xi$ is in $\mathcal{C}(\ell)$.

A pro- \mathcal{C} -gerbe is a gerbe that is a projective limit of \mathcal{C} -gerbes.

Remark 5.5. It follows from conditions (2) and (6) of Definition 5.1 that if G is a group scheme over κ , then $\mathcal{B}_\kappa G$ is a \mathcal{C} -gerbe if and only if G is in $\mathcal{C}(\kappa)$.

Definition 5.6. Let X be a fibered category over (Aff/κ) . A *\mathcal{C} -fundamental gerbe* $\Pi_{X/\kappa}^\mathcal{C}$ is a pro- \mathcal{C} -gerbe over (Aff/κ) with a morphism of fibered categories $X \rightarrow \Pi_{X/\kappa}^\mathcal{C}$ such that for any other pro- \mathcal{C} -gerbe Γ the induced morphism

$$\text{Hom}_\kappa(\Pi_{X/\kappa}^\mathcal{C}, \Gamma) \longrightarrow \text{Hom}_\kappa(X, \Gamma)$$

is an equivalence of categories.

Remark 5.7. It follows easily from the definition of a projective limit that for $X \rightarrow \Pi_{X/\kappa}^\mathcal{C}$ to be a \mathcal{C} -fundamental gerbe, it is enough to check the condition when Γ is a \mathcal{C} -gerbe.

If $X(\kappa) \neq \emptyset$, the concept of a fundamental gerbe can be recast in the more traditional language of groups and torsors.

Let us fix a stable class \mathcal{C} ; we will consider $\mathcal{C}(\kappa)$ as a full subcategory of the category of affine group schemes over κ .

Definition 5.8. Let x_0 be an element of $X(\kappa)$. The pair (X, x_0) is *\mathcal{C} -rigid* if any pointed G -torsor on (X, x_0) , where $G \in \mathcal{C}(\kappa)$, has trivial automorphism group.

Definition 5.9. Let $x_0 \in X(\kappa)$. A *\mathcal{C} -fundamental group* $\pi_1^\mathcal{C}(X, x_0)$ is a pro- \mathcal{C} -group which prorepresents the functor $\mathcal{C}(\kappa)^{\text{op}} \rightarrow (\text{Set})$ that sends G into the set of isomorphism classes in $\text{Tors}_G(X, x_0)$.

Remark 5.10. Clearly, if $\pi_1^\mathcal{C}(X, x_0)$ is a \mathcal{C} -fundamental group, we have a canonical bijection between homomorphisms $\pi_1^\mathcal{C}(X, x_0) \rightarrow G$ and isomorphism classes of pointed G -torsors on (X, x_0) . This shows that $\pi_1^\mathcal{C}(X, x_0)$ is unique, up to a unique isomorphism.

Proposition 5.11. Let $X \rightarrow (\text{Aff}/\kappa)$ be a fibered category, $x_0 \in X(\kappa)$.

Assume that X has a \mathcal{C} -fundamental gerbe $\rho: X \rightarrow \Pi_{X/\kappa}^\mathcal{C}$, and denote by $\xi \in \Pi_{X/\kappa}^\mathcal{C}(\kappa)$ the image of x_0 . Then (X, x_0) is \mathcal{C} -rigid, and the pro- \mathcal{C} -group scheme $\underline{\text{Aut}}_\kappa \xi$ is a fundamental group scheme for X .

Conversely, assume that X is \mathcal{C} -rigid, and let $\pi_1^\mathcal{C}(X, x_0)$ be a \mathcal{C} -fundamental group scheme. Then there exists a morphism $X \rightarrow \mathcal{B}_\kappa \pi_1^\mathcal{C}(X, x_0)$ making the gerbe $\mathcal{B}_\kappa \pi_1^\mathcal{C}(X, x_0)$ into a \mathcal{C} -fundamental gerbe for X .

Proof. If G and H are group schemes over κ , there is a canonical equivalence between the category of pointed H -torsors on $(\mathcal{B}_\kappa G, e)$, where $e \in \mathcal{B}_\kappa G(\kappa)$ is the trivial torsor, and the set of homomorphisms

$G \rightarrow H$, considered as a category, in which the only arrows are the identities; see [Giraud 1971, Remarque 1.6.7].

Suppose that X has a \mathcal{C} -fundamental gerbe $X \rightarrow \Pi_{X/\kappa}^{\mathcal{C}}$. If we denote by ξ the image of x_0 in $\Pi_{X/\kappa}^{\mathcal{C}}$, we have the equivalence $\mathcal{B}_\kappa \underline{\text{Aut}}_\kappa \xi \simeq \Pi_{X/\kappa}^{\mathcal{C}}$, and an isomorphism between the image of the trivial torsor on $\text{Spec } \kappa$ and ξ . It follows from the definition of a fundamental gerbe that the map $X \rightarrow \Pi_{X/\kappa}^{\mathcal{C}}$ induces an equivalence between pointed G -torsors on (X, x_0) and pointed G -torsors on $(\mathcal{B}_\kappa \underline{\text{Aut}}_\kappa \xi, e)$. Thus (X, x_0) is \mathcal{C} -rigid, because $\text{Tors}_G(X, x_0)$ is equivalent to a set, and we get a bijection between isomorphism classes in $\text{Tors}_G(X, x_0)$ and $\text{Hom}(\underline{\text{Aut}}_\kappa \xi, G)$.

The other direction is proved with similar arguments. □

For most stable classes \mathcal{C} , fundamental gerbes do not exist in any kind of reasonable generality. The point is the following. Suppose that $\Pi_{X/\kappa}^{\mathcal{C}}$ exists, and we are given a 2-commutative diagram

$$\begin{array}{ccc} X & \longrightarrow & \Gamma' \\ \downarrow & & \downarrow \\ \Gamma'' & \longrightarrow & \Gamma \end{array}$$

in which Γ' , Γ'' and Γ are \mathcal{C} -gerbes. Then the induced morphism $X \rightarrow \Gamma' \times_\Gamma \Gamma''$ factors through $\Pi_{X/\kappa}^{\mathcal{C}}$. However, in many cases it is possible to show that $X \rightarrow \Gamma' \times_\Gamma \Gamma''$ cannot factor through a gerbe. Our examples are based on Lemma 3.3.

Example 5.12. Assume that \mathcal{C} contains a reductive nonabelian group. Any such group contains a semisimple nonabelian group G , which in turn contains a parabolic subgroup P . By Lemma 3.3 we have

$$\mathcal{B}_\kappa P \times_{\mathcal{B}_\kappa G} \text{Spec } \kappa \simeq G/P;$$

hence if X is a projective variety with a nonconstant map $X \rightarrow G/P$ (for example \mathbb{P}^1), we have a morphism $X \rightarrow \mathcal{B}_\kappa P \times_{\mathcal{B}_\kappa G} \text{Spec } \kappa$ that does not factor through a gerbe, because any morphism from a gerbe to G/P factors through $\text{Spec } \kappa$, and X cannot have a \mathcal{C} -fundamental gerbe.

Example 5.13. For a more subtle example, let \mathbb{G}_m act on \mathbb{G}_a by multiplication in the usual way. Let \mathcal{C} be a stable class containing $\mathbb{G}_m \times \mathbb{G}_a$. In fact, if $\mathcal{B}_\kappa \mathbb{G}_m \rightarrow \mathcal{B}_\kappa(\mathbb{G}_m \times \mathbb{G}_a)$ is induced by the embedding $\mathbb{G}_m \subseteq (\mathbb{G}_m \times \mathbb{G}_a)$, using the lemma above it is easy to see that

$$\mathcal{B}_\kappa \mathbb{G}_m \times_{\mathcal{B}_\kappa(\mathbb{G}_m \times \mathbb{G}_a)} \mathcal{B}_\kappa \mathbb{G}_m \simeq [\mathbb{G}_a/\mathbb{G}_m].$$

But a morphism $X \rightarrow [\mathbb{G}_a/\mathbb{G}_m]$ corresponds to an invertible sheaf L with a section $s \in H^0(X, L)$; if we take a reduced positive-dimensional projective variety X , this has an invertible sheaf with a section that vanishes at some points, but not everywhere. This defines a morphism $X \rightarrow [\mathbb{G}_a/\mathbb{G}_m]$ that does not factor through a gerbe.

6. Well-founded classes

6A. Well-founded actions. Let G be an affine group scheme of finite type over an algebraically closed field κ , acting on an affine scheme X of finite type over κ .

Definition 6.1. A G -reduced subscheme $V \subseteq X$ is a closed G -invariant subscheme of X with the property that every G -invariant nilpotent sheaf of ideals in \mathcal{O}_V is 0.

Equivalently, a closed G -invariant subscheme $V \subseteq X$ is G -reduced if the quotient stack $[V/G]$ is reduced.

Remark 6.2. If G is smooth, which is automatically the case when $\text{char } \kappa = 0$, a closed G -invariant subscheme $V \subseteq X$ is G -reduced if and only if it is reduced.

Definition 6.3. The action of G on X is *well-founded* if for any G -reduced subscheme $V \subseteq X$ such that $\kappa[V]^G = \kappa$, the action of $G(\kappa)$ on $V(\kappa)$ is transitive.

Remark 6.4. Being well-founded is a property of the quotient stack $[X/G]$: the action is well-founded if for every closed reduced substack $\mathcal{V} \subseteq [X/G]$ with $H^0(\mathcal{V}, \mathcal{O}) = \kappa$, the groupoid $\mathcal{V}(\kappa)$ is transitive (or, equivalently, \mathcal{V} is a gerbe over κ).

Remark 6.5. If the action of G is well-founded, then it has closed orbits. The converse holds if G is reductive, by geometric invariant theory, but not in general (consider the example in which X is a reductive group and G is a parabolic subgroup acting by translation).

It is immediate to give examples of actions that are not well-founded: the action of \mathbb{G}_m on \mathbb{A}^1 by multiplication springs to mind.

A class of examples of well-founded actions comes from the following proposition, which is a variant of the classical Kostant–Rosenlicht theorem. Recall that a linear group scheme G is called *unipotent* if for every nonzero representation $G \rightarrow \text{GL}(V)$ we have $V^G \neq 0$.

Proposition 6.6. *If G is unipotent, the action of G on X is always well-founded.*

Proof. Assume that it is not so. Let $V \subseteq X$ be an invariant closed subset such that $\kappa[V]^G = \kappa$ (we do not need to assume that V is G -reduced). If $v_0 \in V(\kappa)$ is a point whose orbit has minimal dimension, then its scheme-theoretic orbit $\Omega \subseteq V$ is closed. From the lemma below it follows that $\Omega = V$, and the conclusion follows. □

Lemma 6.7. *Let V be an affine κ -scheme with an action of a unipotent κ -group scheme G . Assume that $H^0(V, \mathcal{O})^G = \kappa$. Then the only proper G -invariant closed subscheme of V is \emptyset .*

Proof. Let $W \subseteq V$ be a proper invariant closed subscheme; call $I \subseteq H^0(V, \mathcal{O})$ its ideal. Then G acts rationally on $I \neq 0$; so there exists $f \in I^G \setminus \{0\}$. Since $H^0(V, \mathcal{O})^G = \kappa$ we have $f \in \kappa \setminus \{0\}$. So f is invertible, and therefore $W = \emptyset$. □

In what follows we will use the following notation. Assume $\text{char } \kappa = p > 0$, and $n \in \mathbb{N}$. Let Y be a κ -scheme. We denote by Y_n the scheme Y , considered as a κ -scheme via the composite $Y \rightarrow \text{Spec } \kappa \rightarrow \text{Spec } \kappa$, where the homomorphism $\text{Spec } \kappa \rightarrow \text{Spec } \kappa$ is induced by the ring homomorphism $\kappa \rightarrow \kappa$ defined by $x \mapsto x^{p^{-n}}$. Equivalently we can define Y_n as the fibered product $\text{Spec } \kappa \times_{\text{Spec } \kappa} Y$, where the map $\text{Spec } \kappa \rightarrow \text{Spec } \kappa$ is induced by $x \mapsto x^{p^n}$. We have a relative Frobenius map $F_n: Y_n \rightarrow Y$.

Lemma 6.8. *Suppose that the induced action of G_{red} on X_{red} is well-founded. Then the action of G on X is also well-founded.*

Proof. If $\text{char } \kappa = 0$ then $G = G_{\text{red}}$ is smooth, and the result is obvious, since X_{red} and X have the same closed reduced subschemes.

Assume $\text{char } \kappa = p > 0$; assume that $V \subseteq X$ is a G -reduced subscheme of X with $\kappa[V]^G = \kappa$. It is enough to show that $\kappa[V]^{G_{\text{red}}} = \kappa$, for then the action of $G(\kappa) = G_{\text{red}}(\kappa)$ on $V(\kappa) = V_{\text{red}}(\kappa)$ will be transitive.

Fix a positive integer n : the Frobenius morphism $F_n: X_n \rightarrow X$ is $G_n \rightarrow G$ -equivariant, and carries the closed subscheme $V_n \subseteq X_n$ in V . For $n \gg 0$, the scheme-theoretic images of V_n in V and of G_n in G will be, respectively, V_{red} and G_{red} ; hence $\kappa[V_{\text{red}}]^{G_{\text{red}}} \subseteq \kappa[V_n]^{G_n} = \kappa$, and this completes the proof. \square

Lemma 6.9. *Assume that G is smooth, and that the action of G^0 on X is well-founded. Then the action of G on X is also well-founded.*

Proof. Let $\Gamma := G/G^0$. Let $V \subseteq X$ be a G -reduced subscheme such that $\kappa[V]^G = \kappa$. The action of Γ on the connected components of V is clearly transitive. Let W be a connected component of V , and let $\Delta \subseteq \Gamma$ be the stabilizer of W . Then we have $(\kappa[W]^{G^0})^\Delta = (\kappa[V]^{G^0})^\Gamma = \kappa[V]^G = \kappa$; hence $\kappa[W]^{G^0}$ is an integral extension of κ , so it is contained in the integral closure of κ in $\kappa[W]$, which equals κ . Since the action of G^0 is well-founded by hypothesis, we have that the action of $G^0(\kappa)$ on $W(\kappa)$ is transitive, which implies that the action of $G(\kappa)$ on $V(\kappa)$ is transitive. \square

Lemmas 6.8 and 6.9 imply the following useful fact.

Proposition 6.10. *Suppose that the induced action of G_{red}^0 on X_{red} is well-founded. Then the action of G on X is also well-founded.*

6B. Virtually nilpotent group schemes and well-founded actions. Recall the following facts from the theory of algebraic groups. Assume that κ is algebraically closed, and let G be a smooth connected affine algebraic group over κ :

- (1) G is solvable if and only if G contains no nontrivial semisimple subgroups, and if and only if it is a semidirect product $U \rtimes T$, where T is a torus and U is a smooth unipotent group.
- (2) G is nilpotent if and only if it is the product of a torus and a unipotent group.

Definition 6.11. Let G be an affine group scheme of finite type over a field κ .

We say that G is *virtually nilpotent* if $(G_{\bar{\kappa}})_{\text{red}}^0$ is nilpotent.

Virtually abelian and *virtually unipotent* group schemes are defined similarly.

Our main result in this section is the following characterization of virtually nilpotent group schemes.

Theorem 6.12. *An affine group scheme G of finite type over κ is virtually nilpotent if and only if for any algebraically closed extension ℓ of κ and any two subgroup schemes H and K of G_ℓ , the action of $H \times K$ on G_ℓ defined by $(h, k) \cdot g = h^{-1}gk$ is well-founded.*

The strange-looking condition in [Theorem 6.12](#) is exactly what is needed to make the proof of [Theorem 7.1](#) work.

Proof. For the purpose of the proof we will call an affine group scheme G *well-founded* if it satisfies the condition in the statement. Thus we need to prove that G is well-founded if and only if it is virtually nilpotent.

We will need the following facts.

Lemma 6.13. *Assume that G is an affine group scheme of finite type over κ :*

- (1) *If G is well-founded, subgroups and quotients of G are well-founded.*
- (2) *If ℓ is an extension of κ , then G_ℓ is well-founded if and only if G is well-founded.*
- (3) *G is well-founded if and only if the condition of [Theorem 6.12](#) is satisfied when H and K are smooth and connected.*
- (4) *G is well-founded if and only if $(G_{\bar{\kappa}})_{\text{red}}^0$ is well-founded.*

Proof. Parts (1) and (2) are straightforward.

Part (3) follows from [Proposition 6.10](#).

Let us prove (4). We may assume that κ is algebraically closed. If G is well-founded, so is G_{red}^0 , by part (1).

Assume that G_{red}^0 is well-founded. Let H and K be subgroup schemes of G , and let us check that the action of $H \times K$ on G is well-founded. By [Proposition 6.10](#) we may assume that G is smooth, and H and K are smooth and connected.

Let $V \subseteq G$ be a closed $H \times K$ -invariant subscheme with $\kappa[V]^{H \times K} = \kappa$. Then V must be connected. If $V \subseteq G^0$, then the action of $H \times K$ on V is transitive by hypothesis. Let us reduce to this case.

Let $g \in V(\kappa)$, and set $W := g^{-1}V$. Then W is contained in G^0 . It is also invariant under the action of $(g^{-1}Hg) \times K$, and $\kappa[W]^{(g^{-1}Hg) \times K} \simeq \kappa[V]^{H \times K} = \kappa$. So the action of $(g^{-1}Hg) \times K$ on W is transitive, which implies that the action of $H \times K$ on V is transitive. \square

Let us prove the theorem. By [Lemma 6.13](#), we may assume that κ is algebraically closed, and G is smooth and connected.

Now assume that G is well-founded. We want to show that G is virtually nilpotent.

If G were not solvable, it would contain a nontrivial semisimple group. By [Lemma 6.13\(1\)](#), we may assume that G is nontrivial and semisimple. Let P be a parabolic subgroup of G : by setting $H = P$ and $K = \{1\}$ we see that $\kappa[G]^{H \times K} = \kappa$, but the action of $H \times K$ is clearly not transitive. This is a contradiction.

So G is a semidirect product $U \rtimes T$, where T is a torus and U is a smooth unipotent group. We need to show that the action of T on U is trivial. Take $H = K = T$: the quotient G/T is isomorphic to U , and the corresponding left action on U is given by conjugation. By hypothesis, the orbits of the action of $T \times T$ on G are closed: this implies that the orbits for the action of T on U are closed. Set $U//T := \text{Spec } \kappa[U]^T$: by affine GIT, the fibers of the projection $U \rightarrow U//T$ are, set-theoretically, precisely the orbits of T on U .

But the identity in U forms an orbit: this implies that $\dim U = \dim U//T$, so the generic orbit of T on U is finite. Since T is smooth and connected, this implies that the action is trivial.

For the opposite implication, that virtually nilpotent implies well-founded, by [Lemma 6.13\(4\)](#) it is enough to prove that a product $U \times T$, where U is smooth unipotent and T is a torus, is well-founded.

First, assume that G is abelian. Let H and K be subgroup schemes of G . Then $H \backslash G / K = G / L = G // L$, where L is the subgroup scheme of G generated by H and K , and the statement is easy.

This takes care of the case of the torus. The case that G is unipotent follows from [Proposition 6.6](#).

For the general case, let H be a smooth connected subgroup scheme of G . Since H is also nilpotent, we can split it as a product $H_U \times H_T$ of a unipotent group scheme and a torus: but since all homomorphisms $H_U \rightarrow T$ and $H_T \rightarrow U$ are trivial, we have that $H_U \subseteq U$ and $H_T \subseteq T$.

Let H and K be smooth connected subgroup schemes of G , and let $V \subseteq G$ be a $(H \times K)$ -reduced subscheme of G such that $\kappa[V]^{H \times K} = \kappa$. Let V_U and V_T be the scheme-theoretic images of V in U and T respectively. Then $V \subseteq V_U \times V_T$; furthermore, V_U and V_T are reduced and invariant under the actions of $H_U \times K_U$ and $H_T \times K_T$ respectively. We have

$$\kappa[V_U]^{H_U \times K_U} \subseteq \kappa[V_U]^{H \times K} \subseteq \kappa[V]^{H \times K} = \kappa,$$

so that $\kappa[V_U]^{H_U \times K_U} = \kappa$, and analogously $\kappa[V_T]^{H_T \times K_T} = \kappa$. Since U and T are well-founded, we have that V_U and V_T are orbits for $H_U \times K_U$ and $H_T \times K_T$ respectively. It follows immediately that the action of $H \times T$ on V is transitive. This completes the proof of this implication, and of [Theorem 6.12](#). \square

We will need the following easy consequence of [Theorem 6.12](#), which we record here.

Lemma 6.14. *Let G be a virtually nilpotent group scheme over κ , and let $\phi: H \rightarrow G$ and $\psi: K \rightarrow G$ be two homomorphisms of affine group schemes. Then the action of $H \times K$ on G defined by $g \cdot (h, k) = \phi(h)^{-1} g \psi(k)$ is well-founded.*

Proof. Replace H and K with their images in G , and apply [Theorem 6.12](#). \square

6C. Well-founded classes.

Definition 6.15. A stable class \mathcal{C} is *well-founded* if it consists of virtually nilpotent groups.

The class of all virtually nilpotent groups is stable, and hence well-founded. Obviously, if \mathcal{C} is a well-founded class, any subclass of \mathcal{C} that is closed under isomorphisms, extensions of scalars, and taking products, subgroups, quotients and twisted forms, is also well-founded.

This yields a vast range of examples of well-founded classes:

- (1) all virtually nilpotent group schemes,
- (2) virtually abelian group schemes,
- (3) virtually unipotent group schemes,
- (4) finite group schemes,
- (5) linearly reductive finite group schemes,

- (6) abelian affine group schemes,
- (7) diagonalizable group schemes,
- (8) group schemes of multiplicative type,
- (9) unipotent group schemes.

All these classes are in fact very stable, with the exception of (7).

7. Existence of fundamental gerbes

The following is our first main theorem.

Theorem 7.1. *Let X be a fibered category over (Aff/κ) . Assume that X is concentrated and geometrically reduced, and that $H^0(X, \mathcal{O}) = \kappa$. If \mathcal{C} is a well-founded class, then there exists a fundamental gerbe $X \rightarrow \Pi_{X/\kappa}^{\mathcal{C}}$.*

Corollary 7.2. *Let X be fibered category over (Aff/κ) , and let $x_0 \in X(k)$. Assume that X is concentrated and geometrically reduced, and that $H^0(X, \mathcal{O}) = \kappa$. If \mathcal{C} is a well-founded class, then there exists a \mathcal{C} -fundamental group $\pi_1^{\mathcal{C}}(X, x_0)$.*

Remark 7.3. By definition, the fundamental gerbe $\Pi_{X/\kappa}^{\mathcal{C}}$ only depends on the class of \mathcal{C} -gerbes defined over κ ; there maybe different classes \mathcal{C} for which the class of \mathcal{C} -gerbes defined over κ coincide (for example, when κ is algebraically closed, \mathcal{C} -gerbes over κ are the same when \mathcal{C} is either the class of diagonalizable groups, or of groups schemes of multiplicative type).

Remark 7.4. There is a kind of converse of [Theorem 7.1](#). Suppose that κ is algebraically closed, and that \mathcal{C} is a stable class such that $\Pi_{X/\kappa}^{\mathcal{C}}$ exists for all $X \rightarrow (\text{Aff}/\kappa)$ satisfying the conditions of [Theorem 7.1](#). Let G be in $\mathcal{C}(\kappa)$, and let H and K be subgroup schemes of G . Then we claim that the action of $H \times K$ on G is well-founded.

By [Lemma 3.3](#) we have $\mathcal{B}_\kappa H \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa K = [G/H \times K]$. Let $V \subseteq G$ be an $H \times K$ -reduced subscheme with $\kappa[V]^{H \times K} = \kappa$; then $X := [V/H \times K]$ satisfies the hypotheses of [Theorem 7.1](#). The closed embedding

$$X \subseteq [G/H \times K] = \mathcal{B}_\kappa H \times_{\mathcal{B}_\kappa G} \mathcal{B}_\kappa K$$

must factor through $\Pi_{X/\kappa}^{\mathcal{C}}$, and this clearly implies that $X(\kappa)$ contains a unique isomorphism class, so the action of $(H \times K)(\kappa)$ on $V(\kappa)$ is transitive.

In fact, one can prove that to conclude that the action of $H \times K$ on G is well-founded it is enough to assume that $\Pi_{X/\kappa}^{\mathcal{C}}$ exists for all schemes satisfying the conditions of [Theorem 7.1](#). The proof is somewhat complicated, and we omit it.

The strategy of the proof of the theorem is exactly the same as that in [\[Borne and Vistoli 2015\]](#).

Definition 7.5. Let Γ be a \mathcal{C} -gerbe. A morphism of fibered categories $X \rightarrow \Gamma$ is *Nori-reduced* if for any factorization $X \rightarrow \Gamma' \rightarrow \Gamma$, where Γ' is a \mathcal{C} -gerbe and $\Gamma' \rightarrow \Gamma$ is faithful, then $\Gamma' \rightarrow \Gamma$ is an isomorphism.

Let I be a skeleton of the 2-category of Nori reduced morphisms $X \rightarrow \Gamma$. Thus, an element i of I is a Nori-reduced morphism $X \rightarrow \Gamma_i$, and an arrow $u: j \rightarrow i$ is given by a 2-commutative diagram

$$\begin{array}{ccc} & & \Gamma_j \\ & \nearrow & \downarrow u \\ X & & \Gamma_i \\ & \searrow & \end{array}$$

The fundamental gerbe $\Pi_{X/\kappa}^{\mathcal{C}}$ will be constructed as the limit $\varprojlim_{i \in I} \Gamma_i$; for this we need to show that I is a boolean cofiltered 2-category, and that every morphism from X to a \mathcal{C} -gerbe Γ is bounded by a Nori-reduced morphism $X \rightarrow \Gamma_i$.

Let us recall from [Borne and Vistoli 2015] the notion of *scheme-theoretic image* of a morphism of fibered categories $f: X \rightarrow Y$, where X is a concentrated fibered category and Y is an algebraic stack. We define the scheme-theoretic image Y' of X in Y to be the intersection of all the closed substacks Z of Y such that f factors, necessarily uniquely, through Z . Alternatively, $Y' \subseteq Y$ is the closed substack associated with the kernel of the natural homomorphism $\mathcal{O}_Y \rightarrow f_*\mathcal{O}_X$. The fact that this is quasicoherent is easy when X is a concentrated scheme; the general case reduces to this by using an fpqc cover $U \rightarrow X$.

The key point is the following.

Lemma 7.6. *Let $\Gamma' \rightarrow \Gamma$ and $\Gamma'' \rightarrow \Gamma$ be two morphisms of \mathcal{C} -gerbes. The scheme-theoretic image of any morphism $X \rightarrow \Gamma' \times_{\Gamma} \Gamma''$, where $X \neq \emptyset$, is a \mathcal{C} -gerbe.*

This is the only point in the proof of Theorem 7.1 in which the hypothesis that the class is well-founded is used.

Proof. Call $\Delta \subseteq \Gamma' \times_{\Gamma} \Gamma''$ the scheme-theoretic image of X . We need to show that Δ is a gerbe, and that if η is in $\Delta(\ell)$, where ℓ is an extension of κ , the group scheme $\underline{\text{Aut}}_{\ell} \eta$ is in \mathcal{C} .

For this we can base change to ℓ , and assume that $\kappa = \ell$. Call ξ', ξ'' and ξ the images of η in Γ', Γ'' and Γ respectively, and set $G' := \underline{\text{Aut}}_{\kappa} \xi'$, and do analogously for G'' and G . The morphisms $\Gamma' \rightarrow \Gamma$ and $\Gamma'' \rightarrow \Gamma$ induce homomorphisms $\phi': G' \rightarrow G$ and $\phi'': G'' \rightarrow G$; furthermore we have $\underline{\text{Aut}}_{\kappa} \eta = G' \times_G G''$, and since $G' \times_G G''$ is a subgroup scheme of $G' \times G''$ we have $G' \times_G G'' \in \mathcal{C}$.

Let us check that Δ is a gerbe. We can make a further extension of κ , and assume that κ is algebraically closed. There are isomorphisms $\Gamma \simeq \mathcal{B}_{\kappa} G$, $\Gamma' \simeq \mathcal{B}_{\kappa} G'$ and $\Gamma'' \simeq \mathcal{B}_{\kappa} G''$, and the morphisms $\Gamma' \rightarrow \Gamma$ and $\Gamma'' \rightarrow \Gamma$ are induced by ϕ' and ϕ'' . By Lemma 3.3 we have an isomorphism

$$\mathcal{B}_{\kappa} G' \times_{\mathcal{B}_{\kappa} G} \mathcal{B}_{\kappa} G'' \simeq [G/(G' \times G'')],$$

where the action of $G' \times G''$ on G is defined by

$$g \cdot (g', g'') := \phi'(g')^{-1} g \phi''(g'').$$

The scheme-theoretic image of X in $[G/(G' \times G'')]$ is of the form $[V/(G' \times G'')]$, where V is a $(G' \times G'')$ -stable subscheme of G . Call A the κ -algebra corresponding to V , so that $V/(G' \times G'') = \text{Spec } A^{G' \times G''}$. Since $H^0(X, \mathcal{O}) = \kappa$, we have that the composite $X \rightarrow [V/(G' \times G'')] \rightarrow V/(G' \times G'')$ factors through

$\text{Spec } \kappa$; but since the scheme-theoretic image of X in $[V/(G' \times G'')]$ is $[V/(G' \times G'')]$ itself, we see that $A^{G' \times G''} = \kappa$. Then it follows that the action of $H := G' \times G''$ on V has only one orbit (this is immediately seen by replacing G' and G'' with their images in G). Choose a point $v_0 \in V(\kappa)$, and call $K \subseteq H$ the stabilizer of v_0 ; we have a closed embedding $\Omega := H/K \subseteq V$. Call I the sheaf of ideals of $[\Omega/H]$ in $[V/H]$; the inverse image of I in \mathcal{O}_V is nilpotent, so the inverse image of I in X is 0, because X is reduced. Hence $X \rightarrow [V/H]$ factors through $[\Omega/H] = [(H/K)/H] = \mathcal{B}_\kappa K$. So $[V/H] = \mathcal{B}_\kappa K$, and this ends the proof of the lemma. \square

Lemma 7.7. *If Δ is a \mathcal{C} -gerbe, every morphism $X \rightarrow \Delta$ factors as $X \rightarrow \Gamma$, where Γ is a \mathcal{C} -gerbe, $X \rightarrow \Gamma$ is Nori-reduced and $\Gamma \rightarrow \Delta$ is representable.*

Proof. If $X \rightarrow \Delta$ is not Nori-reduced, choose a factorization $X \rightarrow \Delta_1 \rightarrow \Delta$, where $\Delta_1 \rightarrow \Delta$ is representable. If $X \rightarrow \Delta_1$ is not Nori-reduced, let us choose an analogous factorization $X \rightarrow \Delta_2 \rightarrow \Delta_1$. This process cannot continue ad infinitum, as is shown in the following lemma. \square

Lemma 7.8. *If Δ is an affine gerbe, and*

$$\dots \longrightarrow \Delta_i \longrightarrow \Delta_{i-1} \longrightarrow \dots \longrightarrow \Delta_1 \longrightarrow \Delta$$

is an infinite sequence of representable maps of affine gerbes, there exists a positive integer m such that $\Delta_i \rightarrow \Delta_{i-1}$ is an equivalence for all $i \geq m$.

Proof. We can extend the base field κ , and assume that κ is algebraically closed. If ℓ is an algebraically closed extension of κ such that $\Delta_i(\ell) \neq \emptyset$, choose an object ξ of $\Delta_i(\ell)$. Call d_i the dimension of $\text{Aut}_\ell \xi$, and e_i the degree of the finite scheme $\text{Aut}_\ell \xi / (\text{Aut}_\ell \xi)_{\text{red}}^0$. It is immediate to show that d_i and e_i are independent of ξ and ℓ . If η denotes the image of ξ in $\Delta_{i-1}(\ell)$, the morphism $\text{Aut}_\ell \xi \rightarrow \text{Aut}_\ell \eta$ induced by the map $\Delta_i \rightarrow \Delta_{i-1}$ is a monomorphism, and an isomorphism if and only if $\Delta_i \rightarrow \Delta_{i-1}$ is an equivalence.

Clearly $d_i \leq d_{i-1}$, and if $d_i = d_{i-1}$ then $e_i \leq e_{i-1}$. Furthermore, $d_i = d_{i-1}$ and $e_i = e_{i-1}$ if and only if $\text{Aut}_\ell \xi \rightarrow \text{Aut}_\ell \eta$ is an isomorphism, and the statement follows from this. \square

Let us show that I is a cofiltered category; we need to show that given two arrows $j \rightarrow i$ and $k \rightarrow i$ in I , there exists a commutative diagram

$$\begin{array}{ccc} & & j \\ & \nearrow & \searrow \\ l & & i \\ & \searrow & \nearrow \\ & & k \end{array}$$

and, that given two arrows $u, v : j \rightarrow i$, there exists a unique 2-arrow $u \rightarrow v$.

For the first point, the two arrows $j \rightarrow i$ and $k \rightarrow i$ correspond to a 2-commutative diagram

$$\begin{array}{ccc} & & \Gamma_j \\ & \nearrow & \searrow \\ X & & \Gamma_i \\ & \searrow & \nearrow \\ & & \Gamma_k \end{array}$$

inducing a morphism $X \rightarrow \Gamma_j \times_{\Gamma_k} \Gamma_i$. By [Lemma 7.6](#) this factors through a \mathcal{C} -subgerbe $\Delta \subseteq \Gamma_j \times_{\Gamma_k} \Gamma_i$, and by [Lemma 7.7](#) it factors through a Nori-reduced morphism $X \rightarrow \Gamma_l$, proving what we want.

The second fact follows from the analogue of [\[Borne and Vistoli 2015, Lemma 5.13\]](#).

Lemma 7.9. *Let $f: X \rightarrow \Gamma$ and $g: X \rightarrow \Delta$ be morphisms, where Γ and Δ are \mathcal{C} -gerbes and f is Nori-reduced. Suppose that $u, v: \Gamma \rightarrow \Delta$ are morphisms of fibered categories, and $\alpha: u \circ f \simeq g$ and $\beta: v \circ f \simeq g$ are isomorphisms. Then there exists a unique isomorphism $\gamma: u \simeq v$ such that $\beta \circ (\gamma * \text{id}_f) = \alpha$.*

This can be expressed by saying that, given two 2-commutative diagrams

$$\begin{array}{ccc}
 X & \begin{array}{c} \xrightarrow{f} \\ \searrow g \end{array} & \begin{array}{c} \Gamma \\ \downarrow u \\ \Delta \end{array}
 \end{array}
 \quad \text{and} \quad
 \begin{array}{ccc}
 X & \begin{array}{c} \xrightarrow{f} \\ \searrow g \end{array} & \begin{array}{c} \Gamma \\ \downarrow v \\ \Delta \end{array}
 \end{array}$$

in which f is Nori-reduced, there exists a unique isomorphism $u \simeq v$ making the diagram

$$\begin{array}{ccc}
 X & \begin{array}{c} \xrightarrow{f} \\ \searrow g \end{array} & \begin{array}{c} \Gamma \\ \downarrow u \\ \Delta \end{array} \\
 & & \begin{array}{c} \downarrow v \\ \Delta \end{array}
 \end{array}$$

2-commutative.

Proof. This is virtually identical to the proof of [\[Borne and Vistoli 2015, Lemma 5.13\]](#).

Consider the category $\Gamma' \rightarrow \Gamma$ fibered in sets over Γ , whose objects over a κ -scheme T are pairs (ξ, ρ) , where ξ is an object of $\Gamma(T)$ and ρ is an isomorphism of $u(\xi)$ with $v(\xi)$ in $\Delta(T)$. This can be written as a fibered product

$$\begin{array}{ccc}
 \Gamma' & \longrightarrow & \Gamma \\
 \downarrow & & \downarrow \langle u, v \rangle \\
 \Delta & \xrightarrow{\delta} & \Delta \times \Delta
 \end{array}$$

where $\delta: \Delta \rightarrow \Delta \times \Delta$ is the diagonal.

An isomorphism $u \simeq v$ corresponds to a section of the projection $\Gamma' \rightarrow \Gamma$, or, again, to a substack $\Gamma'' \subseteq \Gamma'$ such that the restriction $\Gamma'' \rightarrow \Gamma$ of the projection is an isomorphism. The composite isomorphism $u \circ f \xrightarrow{\alpha} g \xrightarrow{\beta^{-1}} v \circ f$ yields a lifting $X \rightarrow \Gamma'$ of $f: X \rightarrow \Gamma$; the thesis can be translated into the condition that there exists a unique substack $\Gamma'' \subseteq \Gamma'$ as above, such that $X \rightarrow \Gamma'$ factors through Γ'' . By [Lemma 7.6](#), there is a unique closed substack Γ'' of Γ' that is a gerbe, such that $X \rightarrow \Gamma'$ factors through Γ'' . However, $\Gamma'' \rightarrow \Gamma$ is representable, because $\Gamma' \rightarrow \Gamma$ is, so $\Gamma'' \rightarrow \Gamma$ is an equivalence, since f is Nori-reduced. \square

Now we set $\Pi_{X/\kappa}^{\mathcal{C}} := \varprojlim_{i \in I} \Gamma_i$; the morphisms $X \rightarrow \Gamma_i$ yield a morphism $X \rightarrow \Pi_{X/\kappa}^{\mathcal{C}}$. We need to show that if Δ is a \mathcal{C} -gerbe, the induced functor

$$\text{Hom}_{\kappa}(\Pi_{X/\kappa}^{\mathcal{C}}, \Delta) \longrightarrow \text{Hom}_{\kappa}(X, \Delta)$$

is an equivalence. From [Lemma 7.7](#) it follows that it is essentially surjective.

Now, the natural functor

$$\varinjlim_{i \in I} \text{Hom}_\kappa(\Gamma_i, \Delta) \longrightarrow \text{Hom}_\kappa(\Pi_{X/\kappa}^\mathcal{C}, \Delta)$$

is an equivalence, by [Borne and Vistoli 2015, Proposition 3.7]; hence to prove that the above functor is fully faithful it is enough to show that for every $i \in I$ the induced functor $\text{Hom}(\Gamma_i, \Delta) \rightarrow \text{Hom}(X, \Delta)$ is fully faithful.

Call $f : X \rightarrow \Gamma_i$ the canonical morphism. Consider two morphisms $u, v : \Gamma_i \rightarrow \Delta$, and an isomorphism of functors $\beta : u \circ f \simeq v \circ f$. We need to check that there exists a unique isomorphism $\gamma : u \simeq v$ such that $\gamma * \text{id}_f = \beta$. This follows immediately from Lemma 7.6.

8. Change of class

Let \mathcal{C} be a well-founded class of group schemes over κ , and let Γ be an affine gerbe. It follows from Proposition 3.10 that a morphism $\Gamma \rightarrow \Delta$ is Nori-reduced if and only if it is locally full. From this, and from Proposition 3.11 we deduce that the morphism $\Gamma \rightarrow \Pi_{\Gamma/\kappa}^\mathcal{C}$ is locally full.

From now on we use the notation $\Gamma^\mathcal{C}$ for $\Pi_{\Gamma/\kappa}^\mathcal{C}$. Another way of thinking about $\Gamma^\mathcal{C}$ is the following. We can write Γ as the projective limit $\varprojlim \Gamma_i$, where the limit is taken over all the locally full morphisms $\Gamma \rightarrow \Gamma_i$. Let $\{\Gamma_j\}_{j \in J}$ denote the full subcategory of I consisting of those Γ_i that are \mathcal{C} -gerbes. This is a cofiltered full subcategory of I . Every morphism from Γ to a \mathcal{C} -gerbe factors uniquely through $\varprojlim \Gamma_j$; hence $\Gamma^\mathcal{C} = \varprojlim \Delta_i$.

Suppose that \mathcal{D} is a stable subclass of \mathcal{C} . Since a pro- \mathcal{D} -gerbe is also a pro- \mathcal{C} -gerbe, the morphism $X \rightarrow \Pi_{X/\kappa}^\mathcal{D}$ induces a morphism $\Pi_{X/\kappa}^\mathcal{C} \rightarrow \Pi_{X/\kappa}^\mathcal{D}$.

Proposition 8.1. *The induced morphism $\Pi_{X/\kappa}^\mathcal{C} \rightarrow \Pi_{X/\kappa}^\mathcal{D}$ is locally full.*

Proof. Clearly the morphism $X \rightarrow \Pi_{X/\kappa}^\mathcal{C}$ induces an equivalence between $\Pi_{X/\kappa}^\mathcal{D}$ and $\Pi_{\Pi_{X/\kappa}^\mathcal{C}/\kappa}^\mathcal{D}$, so the result follows from the previous discussion. □

Corollary 8.2. *If $\text{Rep } \Pi_{X/\kappa}^\mathcal{C} \rightarrow \text{Vect } X$ is fully faithful, so is $\text{Rep } \Pi_{X/\kappa}^\mathcal{D} \rightarrow \text{Vect } X$.*

Proof. This follows from Propositions 8.1 and 3.10. □

Corollary 8.3. *If \mathcal{C} is a stable class of finite group schemes, the pullback functor $\text{Rep } \Pi_{X/\kappa}^\mathcal{C} \rightarrow \text{Vect}_X$ is fully faithful.*

Proof. If \mathcal{C} is the whole class of finite group schemes, this is in [Borne and Vistoli 2015, Theorem 7.9]. The general case follows from Corollary 8.2. □

9. Weil restriction and change of base

If A is a finite κ -algebra, and X is a quasiprojective A -scheme, we have the Weil restriction $R_{A/\kappa} X$, which is a κ -scheme defined by $R_{A/\kappa} X(R) = X(R \otimes_\kappa A)$, for each κ -algebra R . There is an obvious generalization when $X \rightarrow (\text{Aff}/A)$ is a fibered category: we have the Weil restriction $R_{A/\kappa} X \rightarrow (\text{Aff}/\kappa)$; for the definition and the basic properties we refer to [Borne and Vistoli 2015, Section 6].

We will use the following fact.

Lemma 9.1. *Let G be an affine group scheme over a finite extension ℓ of κ . Then the Weil restriction $\mathbf{R}_{\ell/\kappa}(\mathcal{B}_\ell G)$ is canonically equivalent to $\mathcal{B}_\kappa(\mathbf{R}_{\ell/\kappa} G)$.*

Proof. Let T be an affine κ -scheme, $T \rightarrow \mathbf{R}_{\ell/\kappa}(\mathcal{B}_\ell G)$ a morphism, corresponding to a morphism $T_\ell \rightarrow \mathcal{B}_\ell G$, which in turn corresponds to a G -torsor $E \rightarrow T_\ell$. By applying the Weil restriction functor we obtain a morphism $\mathbf{R}_{\ell/\kappa} E \rightarrow \mathbf{R}_{\ell/\kappa}(T_\ell)$; since the Weil restriction commutes with fibered products, from the action of G on E we get an action $\mathbf{R}_{\ell/\kappa} E \times_\ell \mathbf{R}_{\ell/\kappa} G = \mathbf{R}_{\ell/\kappa}(E \times G) \rightarrow \mathbf{R}_{\ell/\kappa} E$, which leaves the morphism $\mathbf{R}_{\ell/\kappa} E \rightarrow \mathbf{R}_{\ell/\kappa}(T_\ell)$ invariant. By pulling back along the unit morphism $T \rightarrow \mathbf{R}_{\ell/\kappa}(T_\ell)$ we obtain a morphism $F \rightarrow T$, with an action of $\mathbf{R}_{\ell/\kappa} G$ on F leaving it invariant. It is easy to see that $F \rightarrow T$ is a $\mathbf{R}_{\ell/\kappa} G$ -torsor: when $E = T_\ell \times G$ this follows from the fact that $\mathbf{R}_{\ell/\kappa}$ preserves products. In general, if $T' \rightarrow T$ is a faithfully flat morphism of affine schemes such that $T' \times_T E \simeq T' \times G$, we get a diagram

$$\begin{array}{ccc} T' \times \mathbf{R}_{\ell/\kappa} G & \longrightarrow & F \\ \downarrow & & \downarrow \\ T' & \longrightarrow & T \end{array}$$

which is easily checked to be cartesian.

Thus we get a functor $\mathbf{R}_{\ell/\kappa}(\mathcal{B}_\ell G) \rightarrow \mathcal{B}_\kappa(\mathbf{R}_{\ell/\kappa} G)$. Let us produce a functor in the opposite direction.

Let T be an affine κ -scheme, $T \rightarrow \mathcal{B}_\kappa(\mathbf{R}_{\ell/\kappa} G)$ a morphism, corresponding to a $\mathbf{R}_{\ell/\kappa} G$ -torsor $E \rightarrow T$. Then the pullback $E_\ell \rightarrow T_\ell$ is a $(\mathbf{R}_{\ell/\kappa} G)_\ell$ -torsor; with a change of group along the unit morphism $(\mathbf{R}_{\ell/\kappa})_\ell \rightarrow G$ we obtain a G -torsor $E \rightarrow T_\ell$, which yields a morphism $T \rightarrow \mathbf{R}_{\ell/\kappa}(\mathcal{B}_\ell G)$.

We leave it to the reader to check that the two functors above are in fact quasi-inverses. □

Proposition 9.2. *Let \mathcal{C} be a stable class of groups over κ . Let ℓ/κ be a finite extension, and Γ a \mathcal{C} -gerbe over ℓ . Assume that either*

- (1) \mathcal{C} is weakly very stable and ℓ/κ is separable, or
- (2) \mathcal{C} is a very stable class of groups over κ , and every extension of a group in \mathcal{C} by a product of copies of \mathbb{G}_a is still in \mathcal{C} .

Then $\mathbf{R}_{\ell/\kappa} \Gamma$ is a \mathcal{C} -gerbe.

Proof. Suppose that ℓ/κ is separable of degree d , and let κ' be a Galois closure of ℓ/κ ; call v_1, \dots, v_d the embeddings of ℓ in κ' . Denote by Γ_i the fibered product $\Gamma_{\kappa'}$ induced by $v_i: \ell \rightarrow \kappa'$.

We have that $\ell \otimes_\kappa \kappa'$ is isomorphic to the κ' -algebra $(\kappa')^d$, where the i -th projection $\ell \otimes_\kappa \kappa' \simeq (\kappa')^d \rightarrow \kappa'$ corresponds to v_i . But

$$(\mathbf{R}_{\ell/\kappa} \Gamma)_{\kappa'} \simeq \mathbf{R}_{\ell \otimes_\kappa \kappa' / \kappa'} \Gamma_{\ell \otimes_\kappa \kappa'} \simeq \mathbf{R}_{\ell \otimes_\kappa \kappa' / \kappa'} \prod_{i=1}^d \Gamma_i \simeq \prod_{i=1}^d \Gamma_i,$$

and each of the Γ_i is a \mathcal{C} -gerbe; since \mathcal{C} is weakly very stable, it follows that $\mathbf{R}_{\ell/\kappa} \Gamma$ is a \mathcal{C} -gerbe. This proves (1).

For (2), we may replace κ with its separable closure in ℓ , and assume that ℓ/κ is purely inseparable. Call p the characteristic of κ ; since the Weil restriction is functorial, we may assume $\ell = \kappa(\sqrt[p]{a})$.

Let κ' be a finite separable extension of κ , $\ell' := \kappa' \otimes_{\kappa} \ell$. Then $\ell' = \kappa'(\sqrt[p]{a})$; it is enough to show that $(\mathbf{R}_{\ell/\kappa} \Gamma)_{\kappa'} = \mathbf{R}_{\ell'/\kappa'}(\Gamma_{\ell'})$ is a \mathcal{C} -gerbe. Since every finite separable extension of ℓ is of the type above, we may assume that $\Gamma(\ell) \neq \emptyset$, so that $\Gamma = \mathcal{B}_{\ell} G$ for some \mathcal{C} -group G over ℓ . By Lemma 9.1, we need to show that $\mathbf{R}_{\ell/\kappa} G$ is a \mathcal{C} -group. Since the class \mathcal{C} is by hypothesis very stable, it is enough to show that $(\mathbf{R}_{\ell/\kappa} G)_{\ell}$ is a \mathcal{C} -group.

For each nonnegative integer n , denote by A_n the ℓ -algebra $\ell[x]/(x^{n+1})$. Then $\ell \otimes_{\kappa} \ell \simeq A_{p-1}$, and $(\mathbf{R}_{\ell/\kappa} G)_{\ell}$ is the group scheme $\underline{\mathrm{Hom}}_{\ell}(\mathrm{Spec} A_{p-1}, G)$.

Let us show by induction on n that the group scheme $G_n := \underline{\mathrm{Hom}}_{\ell}(\mathrm{Spec} A_n, G)$ is in \mathcal{C} . In fact $G_0 = G \in \mathcal{C}$; the embedding $\mathrm{Spec} A_{n-1} \subseteq \mathrm{Spec} A_n$ induces a homomorphism $G_n \rightarrow G_{n-1}$, whose image is a subgroup scheme of G_{n-1} , which is in \mathcal{C} , by induction hypothesis. It is easy to see that its kernel is isomorphic to the Lie algebra of G , which is a product of copies of \mathbb{G}_a ; hence G_n is in \mathcal{C} . □

Let $X \rightarrow (\mathrm{Aff}/\kappa)$ be a concentrated geometrically reduced fibered category and \mathcal{C} a well-founded class. Let κ'/κ be a field extension. Consider the morphism $X_{\kappa'} \rightarrow (\Pi_{X/\kappa}^{\mathcal{C}})_{\kappa'}$ obtained by base change from the morphism $X \rightarrow \Pi_{X/\kappa}^{\mathcal{C}}$; since $(\Pi_{X/\kappa}^{\mathcal{C}})_{\kappa'}$ is a pro- \mathcal{C} -gerbe, it will factor through $\Pi_{X_{\kappa'}/\kappa'}^{\mathcal{C}}$; so we obtain a morphism of κ' -gerbes $\Pi_{X_{\kappa'}/\kappa'}^{\mathcal{C}} \rightarrow (\Pi_{X/\kappa}^{\mathcal{C}})_{\kappa'}$.

Theorem 9.3. *Assume one of the following hypotheses:*

- (1) *The class \mathcal{C} is weakly very stable, and κ'/κ is an algebraic separable extension.*
- (2) *The class \mathcal{C} is very stable, every extension of a group in \mathcal{C} by a product of copies of \mathbb{G}_a is still in \mathcal{C} , and κ'/κ is an algebraic extension.*

Then the map $\Pi_{X_{\kappa'}/\kappa'}^{\mathcal{C}} \rightarrow (\Pi_{X/\kappa}^{\mathcal{C}})_{\kappa'}$ is an equivalence.

Proof. The proof is virtually identical to the proof of [Borne and Vistoli 2015, Proposition 6.1], using Proposition 9.2. □

10. The tannakian interpretations of the unipotent and virtually unipotent fundamental gerbes

Let X be a concentrated fibered category over (Aff/κ) with $H^0(X, \mathcal{O}) = \kappa$, and \mathcal{C} a well-founded class. The gerbe $\Pi_{X/\kappa}^{\mathcal{C}}$ is tannakian; hence the category of representations $\mathrm{Rep} \Pi_{X/\kappa}^{\mathcal{C}}$ is a tannakian category. For any \mathcal{C} , one can ask if it is possible to give a direct description of $\mathrm{Rep} \Pi_{X/\kappa}^{\mathcal{C}}$ in terms of X . Of course there is a pullback map $\mathrm{Rep} \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \mathrm{Vect}_X$ into the category of locally free sheaves on X ; however, this is in general not fully faithful. For example, when \mathcal{C} is the class of abelian group schemes it is immediate to see that the pullback functor is not fully faithful when there are maps between invertible sheaves on X that are neither zero nor isomorphisms. And in fact we don't have a candidate for such a description of $\mathrm{Rep} \Pi_{X/\kappa}^{\mathcal{C}}$ when \mathcal{C} is the class of abelian groups.

However when \mathcal{C} is the category of unipotent, or virtually unipotent, group schemes, the functor is fully faithful; and in these cases there is a good description of $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}}$.

In what follows we will assume the following conditions on X .

Conditions 10.1. (1) X is concentrated.

(2) X is geometrically reduced.

(3) $H^0(X, \mathcal{O}) = \kappa$.

(4) X is pseudoproper, in the sense of [Borne and Vistoli 2015].

Recall that X is pseudoproper if for any locally free sheaf E on X , the κ -vector space $H^0(X, E)$ is finite-dimensional.

For example, an affine gerbe always satisfies these conditions.

We can't think of an example in which (1), (2) and (3) are satisfied but (4) is not, but we have no doubt that this is for lack of trying.

Notice that if X satisfies the conditions above and κ' is a finite extension of κ , the fibered category $X_{\kappa'} \rightarrow (\text{Aff}/\kappa')$ also satisfies them.

Definition 10.2. Let E be a locally free sheaf on X :

(1) We say that E is *unipotent* if it admits a filtration

$$0 = E_{r+1} \subseteq E_r \subseteq E_{r-1} \subseteq \cdots \subseteq E_1 \subseteq E_0 = E$$

in which all the quotients E_i/E_{i+1} are free.

(2) We say that E is an *extended essentially finite sheaf* if there is a filtration as above in which all the quotients E_i/E_{i+1} are essentially finite.

Unipotent bundles were introduced by M. Nori [1982] under the name *nilpotent bundles*. The second class of bundles was introduced by S. Otabe [2017]; he calls them *semifinite bundles*.

Both classes form a tannakian category, and have a natural interpretation in our language.

Definition 10.3. A group scheme $G \rightarrow \text{Spec } \ell$ is *strongly virtually unipotent* if it has a normal unipotent subgroup $H \subseteq G$ such that G/H is finite.

A strongly virtually unipotent group scheme is clearly virtually unipotent. If G is smooth, then the converse holds; hence if $\text{char } \kappa = 0$, then the converse holds. This is not true in positive characteristic, as the following examples show.

Examples 10.4. Let us give two examples of two group schemes in positive characteristic that are virtually unipotent, but not strongly virtually unipotent. The first one is abelian and defined over a nonperfect field; the second one is not abelian, but is defined over an arbitrary field of positive characteristic. The first example also tells us that the class of strongly virtually unipotent groups, which is weakly very stable, is not very stable.

(1) Let k be a nonperfect field of characteristic $p > 0$ and ℓ/k a purely inseparable extension of degree p . Denote by G the Weil transfer $R_{\ell/k}\mu_p$. Then we claim that G is virtually unipotent, but not strongly virtually unipotent.

We have $\ell \otimes_k \ell \simeq \ell[\varepsilon] := \ell[x]/(x^p)$; hence G_ℓ is the Weil transfer $R_{\ell[\varepsilon]/\ell}\mu_p$. This can be described as the group scheme

$$H := \underline{\text{Hom}}_\ell(\text{Spec } \ell[\varepsilon], \mu_p)$$

whose sections over an ℓ -algebra A are the homomorphisms of A -algebras

$$A[t]/((t - 1)^p) \longrightarrow A[\varepsilon] = A[x]/(x^p).$$

These are uniquely determined by the image of t in $A[\varepsilon]$, which is an element $a_0 + a_1\varepsilon + \dots + a_{p-1}\varepsilon^{p-1}$ with $a_0^p = 1$; the product structure is given by the product in $A[\varepsilon]$. There is a projection $H \rightarrow \mu_p$, defined by sending $a_0 + a_1\varepsilon + \dots + a_{p-1}\varepsilon^{p-1}$ into a_0 , whose kernel is easily seen to be unipotent. Hence G_ℓ is an extension of μ_p by unipotent group scheme, so G is virtually unipotent, and has dimension $p - 1$.

On the other hand, a homomorphism from a unipotent group scheme U on k to G corresponds, by adjunction, to a homomorphism $U_\ell \rightarrow \mu_p$, which must be trivial; so G does not contain any nontrivial unipotent subgroups, and is not strongly virtually unipotent.

(2) Here k can be an arbitrary field of positive characteristic, and n an integer with $n > 1$. Call $\text{GL}_n^{(1)}$ the Frobenius kernel in GL_n , and $U_n \subseteq \text{GL}_n$ the subgroup consisting of strictly upper triangular matrices. Let $G := U_n \ltimes \text{GL}_n^{(1)}$, where the action of U_n on $\text{GL}_n^{(1)}$ is by conjugation. Clearly $G_{\text{red}}^0 = U_n$, so G is virtually unipotent.

Suppose that $H \subseteq G$ is a normal unipotent subgroup such that G/H is finite. Then $U_n \subseteq H$, and $K := H \cap \text{GL}_n^{(1)}$. Then K is nontrivial, because the action of U_n on $\text{GL}_n^{(1)}$ is nontrivial, so U_n is not normal in G . Since K is unipotent we have that the invariant subspace $(k^n)^K$ in k^n is proper and nontrivial; but K is normal in $\text{GL}_n^{(1)}$, so $(k^n)^K$ is $\text{GL}_n^{(1)}$ -invariant. But obviously $\text{GL}_n^{(1)}$ is not contained in any proper parabolic subgroup of GL_n , and this gives a contradiction, showing that H cannot exist, and that G is not strongly virtually unipotent.

In what follows we will denote by $\Pi_{X/\kappa}^U$, $\Pi_{X/\kappa}^{\text{VU}}$ and $\Pi_{X/\kappa}^{\text{SVU}}$ the fundamental gerbe $\Pi_{X/\kappa}^{\mathcal{C}}$, when \mathcal{C} is, respectively, the class of unipotent, virtually unipotent, or strongly virtually unipotent groups.

Theorem 10.5. (1) *The pullback $\text{Rep } \Pi_{X/\kappa}^U \rightarrow \text{Vect}_X$ induces an equivalence of $\text{Rep } \Pi_{X/\kappa}^U$ with the full subcategory of $\text{Vect } X$ whose objects are unipotent locally free sheaves.*

(2) *The pullback $\text{Rep } \Pi_{X/\kappa}^{\text{SVU}} \rightarrow \text{Vect}_X$ induces an equivalence of $\text{Rep } \Pi_{X/\kappa}^{\text{SVU}}$ with the full subcategory of $\text{Vect } X$ whose objects are extended essentially finite sheaves.*

So, if $\text{char } \kappa = 0$, this gives a tannakian interpretation of $\Pi_{X/\kappa}^{\text{VU}} = \Pi_{X/\kappa}^{\text{SVU}}$.

[Theorem 10.5](#) is a particular case of the more general [Theorem 11.9](#) in the next section.

There also a tannakian interpretation of $\Pi_{X/\kappa}^{\text{VU}}$ in positive characteristic, at least with a weak additional assumption on X . Assume that $\text{char } \kappa = p > 0$.

If U is a scheme over κ , denote by $\text{Frob}_U : U \rightarrow U$ the absolute Frobenius map of U .

Denote by $\text{Frob}_X : X \rightarrow X$ the functor sending an object $\xi \in X(T)$, where T is an affine scheme over κ to the pullback $\text{Frob}_T^* \xi$; this is a morphism of fibered categories over $(\text{Aff}/\mathbb{F}_p)$, not over (Aff/κ) . Notice that this definition involves the choice of a cleavage for X ; but the resulting functor is unique, up to a unique isomorphism.

If U is an affine scheme, then $\text{Frob}_{(\text{Aff}/U)} : (\text{Aff}/U) \rightarrow (\text{Aff}/U)$ is the functor corresponding to the morphism $\text{Frob}_U : U \rightarrow U$; we will also denote it by Frob_U . If $U = \text{Spec } A$ we will also use the notation Frob_A .

Clearly, the diagram

$$\begin{array}{ccc} X & \xrightarrow{\text{Frob}_X} & X \\ \downarrow & & \downarrow \\ (\text{Aff}/\kappa) & \xrightarrow{\text{Frob}_\kappa} & (\text{Aff}/\kappa) \end{array}$$

is strictly commutative. If $F : X \rightarrow Y$ is a morphism of fibered categories over (Aff/κ) , then we have an obvious commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\text{Frob}_X} & X \\ \downarrow F & & \downarrow F \\ Y & \xrightarrow{\text{Frob}_Y} & Y \end{array}$$

Definition 10.6. A locally free sheaf E on X is *virtually unipotent* if there exists a positive integer n such that $(\text{Frob}_X^n)^* E$ is an extended essentially finite sheaf.

Theorem 10.7. Assume that X has an fpqc cover $U \rightarrow X$, where U is a noetherian reduced scheme. Then the pullback $\text{Rep } \Pi_{X/\kappa}^{\text{VU}} \rightarrow \text{Vect}_X$ induces an equivalence of the tannakian category $\text{Rep } \Pi_{X/\kappa}^{\text{VU}}$ with the full subcategory of $\text{Vect } X$ consisting of virtually unipotent sheaves.

We do not know whether the (rather weak) condition on the existence of a cover $U \rightarrow X$ as above is necessary for the conclusion to hold. It is certainly satisfied when X is an affine gerbe, because any morphism from the spectrum of a field to an affine gerbe is an fpqc cover.

Notice that from [Theorem 10.7](#) and [Corollary 8.2](#) we obtain the following.

Corollary 10.8. If \mathcal{C} is a stable subclass of the class of virtually unipotent group schemes, then the pullback $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \text{Vect}_X$ is fully faithful.

The following is due to Tonini and Zhang.

Theorem 10.9 [[Tonini and Zhang 2018](#), Corollary II]. Assume that $\text{char } \kappa > 0$, that X is a pseudoproper geometrically reduced algebraic stack of finite type over κ , and that $H^1(X, E)$ is a finite-dimensional vector space over κ for all locally free sheaves on X . Then

$$\Pi_{X/\kappa}^{\text{VU}} = \Pi_{X/\kappa}^{\text{SVU}} = \Pi_{X/\kappa}^{\text{N}}.$$

This is clearly false without the hypothesis on H^1 (for example, take $X = \mathcal{B}_\kappa \mathbb{G}_a$).

11. Unipotent saturations

Suppose that \mathcal{V} is a class of locally free sheaves on fibered categories satisfying [Conditions 10.1](#); for each such fibered category X we denote by $\mathcal{V}(X)$ the class of locally free sheaves on X that are in \mathcal{V} , and also the corresponding full subcategory of Vect_X .

Definition 11.1. Let \mathcal{V} be a class of locally free sheaves on fibered categories satisfying [Conditions 10.1](#). We say that \mathcal{V} is a *tannakian class* if it satisfies the following conditions (where X and Y are arbitrary fibered categories satisfying [Conditions 10.1](#) and Γ is an affine gerbe):

- (1) For each X , the subcategory $\mathcal{V}(X) \subseteq \text{Vect}_X$ is a tannakian subcategory.
- (2) If $f: Y \rightarrow X$ is a morphism and E in $\mathcal{V}(X)$, then f^*E is in $\mathcal{V}(Y)$.
- (3) Let $f: X \rightarrow \Gamma$ be a morphism such that the pullback $f^*: \text{Rep } \Gamma \rightarrow \text{Vect}_X$ induces an equivalence between $\text{Rep } \Gamma$ and $\mathcal{V}(X)$. Then $\mathcal{V}(\Gamma) = \text{Rep } \Gamma$.

Here by a *tannakian subcategory* $\mathcal{V}(X) \subseteq \text{Vect}_X$ we mean that it is a monoidal subcategory closed under isomorphisms and taking duals, that is, tannakian with respect to the induced rigid monoidal structure, and that kernels and cokernels in $\mathcal{V}(X)$ are also kernels and cokernels as homomorphism of sheaves of \mathcal{O}_X -modules.

Remark 11.2. Condition (3) of the definition above may look strange; it has been introduced because it is essential for the proof of [Lemma 11.4](#). We should point out that we don't know of examples in which (1) and (2) hold, but (3) does not.

There are many examples of tannakian classes, for example, the class of free locally free sheaves and that of essentially finite locally free sheaves. Many more examples are provided by the following proposition.

Proposition 11.3. *Let \mathcal{C} be a well-founded class of group schemes; assume that for each X satisfying [Conditions 10.1](#) the functor $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \text{Vect}_X$ is fully faithful. Denote by $\mathcal{V}(X)$ its essential image. Then \mathcal{V} is a tannakian class.*

Proof. The proof is straightforward. □

We will call this \mathcal{V} the *tannakian realization* of the class \mathcal{C} . Classes \mathcal{C} satisfying the condition of [Proposition 11.3](#) will be called *realizable*. Every stable class of finite group schemes is realizable, because of [Corollary 8.3](#).

The following gives a criterion to check that a tannakian class \mathcal{V} is the tannakian realization of a fundamental class \mathcal{C} .

Lemma 11.4. *Let \mathcal{V} be a tannakian class and \mathcal{C} a fundamental class of group schemes. Assume that for every affine gerbe Γ over κ , a representation of Γ is in $\mathcal{V}(\Gamma)$ if and only if it is the pullback of a representation of $\Pi_{\Gamma/\kappa}^{\mathcal{C}}$.*

Then \mathcal{V} is the tannakian realization of \mathcal{C} .

Proof. If Γ is an affine gerbe, we claim that Γ is a pro- \mathcal{C} -gerbe if and only if $\mathcal{V}(\Gamma) = \text{Rep } \Gamma$. In fact, Γ is a pro- \mathcal{C} -gerbe if and only if $\Gamma \rightarrow \Pi_{\Gamma/\kappa}^{\mathcal{C}}$ is an equivalence, that is, by Tannaka duality, if and only if the pullback $\text{Rep } \Pi_{\Gamma/\kappa}^{\mathcal{C}} \rightarrow \text{Rep } \Gamma$ is an equivalence. But $\text{Rep } \Pi_{\Gamma/\kappa}^{\mathcal{C}} \rightarrow \text{Rep } \Gamma$ is fully faithful, by [Proposition 8.1](#), so Γ is a pro- \mathcal{C} -gerbe if and only if $\text{Rep } \Pi_{\Gamma/\kappa}^{\mathcal{C}} \rightarrow \text{Rep } \Gamma$ is essentially surjective. But the condition in the lemma tells us that the essential image of $\text{Rep } \Pi_{\Gamma/\kappa}^{\mathcal{C}} \rightarrow \text{Rep } \Gamma$ is precisely $\mathcal{V}(\Gamma)$.

Let Π be the affine gerbe corresponding to the tannakian category $\mathcal{V}(X)$; this is a pro- \mathcal{C} -gerbe, because of what we just showed. By Deligne’s theorem [[1990](#), Théorème 1.12] we obtain a map $X \rightarrow \Pi$ such that the pullback $\text{Rep } \Pi \rightarrow \text{Vect}_X$ induces an equivalence $\text{Rep } \Pi \simeq \mathcal{V}(X)$. Hence $\mathcal{V}(\Pi) = \text{Rep } \Pi$, because of part (3) of [Definition 11.1](#), so Π is a pro- \mathcal{C} -gerbe, because of the result above. Consider the factorization $X \rightarrow \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \Pi$, which induces a factorization $\text{Rep } \Pi \rightarrow \text{Rep } \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \text{Vect } X$. But $\text{Rep } \Pi$ and $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}}$ have the same essential image $\mathcal{V}(X)$, $\text{Rep } \Pi_{X/\kappa}^{\mathcal{C}} \rightarrow \text{Vect}_X$ is faithful, and $\text{Rep } \Pi \rightarrow \text{Vect}_X$ is fully faithful, so $\text{Rep } \Pi \rightarrow \text{Rep } \Pi_{X/\kappa}^{\mathcal{C}}$ is an equivalence. □

Definition 11.5. Let \mathcal{V} be a tannakian class. The *unipotent saturation* $\overline{\mathcal{V}}$ is defined as follows. Let E be a locally free sheaf on X . We say that E is in $\overline{\mathcal{V}}$ if it admits a filtration

$$0 = E_{r+1} \subseteq E_r \subseteq E_{r-1} \subseteq \cdots \subseteq E_1 \subseteq E_0 = E$$

in which all the quotients E_i/E_{i+1} are in \mathcal{V} .

Thus, the unipotent saturation of the class of free sheaves is the class of unipotent sheaves, while that of the class of essentially free sheaves is the class of extended essentially finite sheaves.

Proposition 11.6. *The unipotent saturation of a tannakian class is a tannakian class.*

Proof. Let us check that the three conditions of [Definition 11.1](#) are satisfied. This is trivial for (2).

Condition (1) is easily proved by adapting the proof in [[Nori 1982](#), Chapter IV.1] for unipotent locally free sheaves.

For condition (3), let $f: X \rightarrow \Gamma$ be a morphism such that $f^*: \text{Rep } \Gamma \rightarrow \text{Vect}_X$ induces an equivalence between $\text{Rep } \Gamma$ and $\overline{\mathcal{V}}(X)$. Let Δ be the affine gerbe corresponding to the tannakian category $\mathcal{V}(X)$. The embedding $\mathcal{V}(X) \subseteq \overline{\mathcal{V}}(X)$ induces a morphism $\phi: \Gamma \rightarrow \Delta$, and a commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & \Gamma \\ & \searrow g & \downarrow \phi \\ & & \Delta \end{array}$$

such that $g^*: \text{Rep } \Delta \rightarrow \text{Vect}_X$ induces an equivalence between $\text{Rep } \Delta$ and $\mathcal{V}(X)$. Since \mathcal{V} is a tannakian class, we have $\mathcal{V}(\Delta) = \text{Rep } \Delta$.

Let E be a nonzero representation of Γ ; we need to show that E is in $\overline{\mathcal{V}}(\Gamma)$. We proceed by induction on $\text{rk } E$, and assume that all representations of Γ of rank less than $\text{rk } E$ are in $\overline{\mathcal{V}}(\Gamma)$. If f^*E is in $\mathcal{V}(X)$, then it is the pullback of a representation $G_0 \in \text{Rep } \Delta = \mathcal{V}(\Delta)$, so $E \simeq \phi^*G_0$ is in $\mathcal{V}(\Gamma)$, and we are done.

By hypothesis we have $f^*E \in \overline{\mathcal{V}}(X)$; hence there exists a subsheaf $0 \neq F_0 \subseteq f^*E$ with $F_0 \in \mathcal{V}(X)$. Let E_0 be a representation of Γ with $f^*E_0 \simeq F_0$; by the argument above, $E_0 \in \mathcal{V}(\Gamma)$. Since f^* is fully faithful we get an embedding $E_0 \subseteq E$. By induction hypothesis E/E_0 is in $\overline{\mathcal{V}}(X)$, so $E \in \overline{\mathcal{V}}(X)$. \square

In general, the unipotent saturation of the tannakian realization of a fundamental class of group schemes is not the tannakian realization of a fundamental class; for example, one can show that the unipotent saturation of the tannakian realization of the class of abelian group schemes is not a tannakian realization.

The main result of this section is that the tannakian realization of the unipotent saturation of a very stable fundamental class of finite group schemes is again a tannakian realization.

Definition 11.7. Let \mathcal{D} be a very stable class of finite group schemes. The *unipotent saturation* $\overline{\mathcal{D}}$ of \mathcal{D} is the class of affine algebraic groups G over extensions ℓ of κ , with the property that there exists a normal unipotent subgroup scheme H such that G/H is in \mathcal{D} .

Proposition 11.8. *The unipotent saturation of a very stable class of finite group schemes is well-founded.*

Proof. Let \mathcal{D} be a very stable class of finite group schemes. Since $\overline{\mathcal{D}}$ is a subclass of the class of all virtually finite group schemes, which is well-founded, it is enough to show that $\overline{\mathcal{D}}$ is stable. This is straightforward. \square

Theorem 11.9. *Let \mathcal{D} be a very stable class of finite group schemes and \mathcal{V} its tannakian realization. Then the tannakian realization of the unipotent saturation $\overline{\mathcal{D}}$ is the unipotent saturation $\overline{\mathcal{V}}$ of \mathcal{V} .*

When applied to the class \mathcal{D} consisting of trivial groups, and to the class of all finite group schemes, this immediately implies [Theorem 10.5](#).

The proof of [Theorem 11.9](#) will occupy the rest of this section. We use [Proposition 11.6](#) and [Lemma 11.4](#); we only have to check that the condition of [Lemma 11.4](#) is satisfied. This is the content of the following proposition. Recall that we use the notation $\Gamma^{\mathcal{D}}$ for $\Pi_{\Gamma/k}^{\mathcal{D}}$.

Proposition 11.10. *If Γ is an affine gerbe, a representation E is in $\overline{\mathcal{V}}(\Gamma)$ if and only if it is a pullback from $\Gamma^{\mathcal{D}}$.*

For the proof of [Proposition 11.10](#) we need the following fact.

Let $\Gamma \rightarrow \Delta$ be a morphism of gerbes over κ . Let us assume that this is locally full, or, equivalently, that Γ is a gerbe over Δ . Then we say that Γ is *unipotent over Δ* if for any morphism $\text{Spec } \ell \rightarrow \Delta$, where ℓ is a field, the fibered product $\text{Spec } \ell \times_{\Delta} \Gamma$ is unipotent.

It is easy to see that if $\text{Spec } \ell \times_{\Delta} \Gamma$ is unipotent for some morphism $\text{Spec } \ell \rightarrow \Delta$, then it is unipotent for all such morphisms.

Proposition 11.11. *Let Γ be a gerbe of finite type over κ ; then Γ is a $\overline{\mathcal{D}}$ -gerbe if and only if Γ is unipotent over $\Gamma^{\mathcal{D}}$.*

Proof of [Proposition 11.10](#), assuming [Proposition 11.11](#). Let Γ be an affine gerbe over κ .

Let us prove that every representation in $\overline{\mathcal{V}}(\Gamma)$ comes from $\Gamma^{\overline{\mathcal{D}}}$. This is obviously true for representations in $\mathcal{V}(\Gamma)$, by definition; hence, it is enough to show that given an extension

$$0 \longrightarrow E_1 \longrightarrow E \longrightarrow E_2 \longrightarrow 0$$

in which both E_1 and E_2 come from $\Gamma^{\overline{\mathcal{D}}}$, the representation E also comes from $\Gamma^{\overline{\mathcal{D}}}$. Let Γ' be a $\overline{\mathcal{D}}$ -gerbe of finite type with a map $\Gamma \rightarrow \Gamma'$ such that E_1 and E_2 come from representations F_1 and F_2 of Γ' .

Consider the fibered category Δ over (Aff/κ) defined as follows. Given an object $\xi \in \Gamma'(A)$, where A is a κ -algebra, we denote by $(E_1)_\xi$ and $(E_2)_\xi$ the pullbacks to $\text{Spec } A$ obtained from the morphism $\text{Spec } A \rightarrow \Gamma'$ corresponding to ξ . An object of Δ over a κ -algebra A is an extension

$$0 \longrightarrow (E_1)_\xi \longrightarrow F \longrightarrow (E_2)_\xi \longrightarrow 0$$

of sheaves of \mathcal{O} -modules over $\text{Spec } A$.

The arrows in the fiber category $\Gamma(A)$ are given by homomorphisms of sheaves $F \rightarrow F'$ of sheaves of \mathcal{O} -modules over $\text{Spec } A$ fitting into a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & (F_1)_\xi & \longrightarrow & F & \longrightarrow & (F_2)_\xi \longrightarrow 0 \\ & & \parallel & & \downarrow & & \parallel \\ 0 & \longrightarrow & (F_1)_\xi & \longrightarrow & F' & \longrightarrow & (F_2)_\xi \longrightarrow 0 \end{array}$$

It is immediate to check that Δ is a gerbe over Γ' . Furthermore, if ℓ is an extension of κ and η is an object of $\Delta(\ell)$ mapping to ξ in $\Gamma'(\ell)$, the kernel of the natural homomorphism of group schemes $\underline{\text{Aut}}_\ell \eta \rightarrow \underline{\text{Aut}}_\ell \xi$ is the vector space $\text{Hom}_\ell((E_2)_\xi, (E_1)_\xi)$; hence it is unipotent. Clearly, an extension of a $\overline{\mathcal{D}}$ -group with unipotent kernel is again in $\overline{\mathcal{D}}$; hence $\underline{\text{Aut}}_\ell \eta$ is in $\overline{\mathcal{D}}$, and Δ is a $\overline{\mathcal{D}}$ -gerbe.

The extension

$$0 \longrightarrow E_1 \longrightarrow E \longrightarrow E_2 \longrightarrow 0$$

gives a lifting $\Gamma \rightarrow \Delta$ of the given morphism $\Gamma \rightarrow \Gamma'$; since Δ is a $\overline{\mathcal{D}}$ -gerbe, the morphism $X \rightarrow \Delta$ factors through $\Gamma^{\overline{\mathcal{D}}}$, and E come from $\Gamma^{\overline{\mathcal{D}}}$, as claimed.

In the other direction, we can replace Γ with $\Gamma^{\overline{\mathcal{D}}}$; so it enough to show that every representation of a $\overline{\mathcal{D}}$ -gerbe Γ is in $\mathcal{V}(\Gamma)$. This follows immediately from the following lemma, by an obvious induction on the rank.

Lemma 11.12. *Let Γ be a $\overline{\mathcal{D}}$ -gerbe over κ ; denote by $\pi : \Gamma \rightarrow \Gamma^{\overline{\mathcal{D}}}$ the projection. Let E be a nonzero representation of Γ . Then $\pi_* E \neq 0$, and the counit homomorphism $\pi^* \pi_* E \rightarrow E$ is injective.*

Proof. Let ℓ be an extension of κ such that $\Gamma(\ell) \neq 0$. By Proposition 11.11, the pullback $\text{Spec } \ell \times_{\Gamma^{\overline{\mathcal{D}}}} \Gamma$ is of the form $\mathcal{B}_\ell G$, where G is a unipotent group scheme over ℓ , and we have a cartesian diagram

$$\begin{array}{ccc} \mathcal{B}_\ell G & \xrightarrow{\psi} & \Gamma \\ \downarrow \rho & & \downarrow \pi \\ \text{Spec } \ell & \xrightarrow{\phi} & \Gamma^{\overline{\mathcal{D}}} \end{array}$$

Set $V := \psi^*E$. Since ϕ is faithfully flat, so is ψ . Furthermore, the formation of π_* commutes with base change, so it is enough to show that $\psi^*\pi^*\pi_*E = \rho^*\rho_*V$ injects in V . But thinking of V as a representation of G we have $\rho^*\rho_*V = V^G$, and the statement is clear. \square

Remark 11.13. The argument in the proof of [Lemma 11.12](#) proves the following fact, which will be used later.

Let $\pi : \Gamma \rightarrow \Delta$ be a locally full morphism of affine gerbes. If E is a representation of Γ , the counit homomorphism $\pi^*\pi_*E \rightarrow E$ is injective.

This ends the proof of [Proposition 11.10](#), assuming [Proposition 11.11](#). \square

Proof of [Proposition 11.11](#). Assume that Γ is unipotent on $\Gamma^{\mathcal{D}}$, and let ξ be an object of $\Gamma(\ell)$ for some extension ℓ/κ . Let η be the image of ξ in $\Gamma^{\mathcal{D}}(\ell)$; we need to show that $\underline{\text{Aut}}_{\ell} \xi$ is a $\overline{\mathcal{D}}$ -group. Let η be the image of ξ in $\Gamma^{\mathcal{D}}(\ell)$; the kernel of the natural surjective map $\underline{\text{Aut}}_{\ell} \xi \rightarrow \underline{\text{Aut}}_{\ell} \eta$ is the automorphism of the object of $(\text{Spec } \ell \times_{\Gamma^{\mathcal{D}}} \Gamma)(\ell)$ coming from ξ , so it is a unipotent group, while $\underline{\text{Aut}}_{\ell} \eta$ is in \mathcal{D} .

For the other implication, let Γ be a $\overline{\mathcal{D}}$ -gerbe. By [Corollary 3.2](#), there exists a finite separable extension ℓ/κ such that $\Gamma(\ell) \neq \emptyset$. Let ξ be an object of $\Gamma(\ell)$; consider the fibered product $\Delta := \text{Spec } \ell \times_{\Gamma^{\mathcal{D}}} \Gamma$. We need to show that Δ is a unipotent gerbe. By [Theorem 9.3\(1\)](#) we see that the formation of Δ commutes with base change along the morphism $\text{Spec } \ell \rightarrow \text{Spec } \kappa$; hence, by base changing we may assume that $\Delta(\kappa) \neq \emptyset$. Let η be the image of $\xi \in \Delta(\kappa)$ in $\Gamma(\kappa)$, and ζ its image in $\Gamma^{\mathcal{D}}(\kappa)$. Set $H := \underline{\text{Aut}}_{\kappa} \xi$, $G := \underline{\text{Aut}}_{\kappa} \eta$ and $L := \underline{\text{Aut}}_{\kappa} \zeta$; then H is a normal subgroup of G , and $L = G/H$. We need to show that H is unipotent.

By hypothesis G is in $\overline{\mathcal{D}}(\kappa)$; hence there exists a normal unipotent subgroup $G' \subseteq G$ such that $G'' := G/G'$ is in $\mathcal{D}(\kappa)$. The corresponding morphism $\mathcal{B}_{\kappa} G \rightarrow \mathcal{B}_{\kappa} G''$ factors through $\mathcal{B}_{\kappa} G \rightarrow (\mathcal{B}_{\kappa} G)^{\mathcal{D}} = \mathcal{B}_{\kappa} L$; this gives a factorization $G \rightarrow L \rightarrow G''$, and shows that H is included in G' . Since G' is unipotent, so is H , as claimed. \square

This ends the proof of [Theorem 11.9](#).

12. The proof of [Theorem 10.7](#)

By [Lemma 11.4](#), to prove [Theorem 10.7](#) it is enough to show the following two results.

Proposition 12.1. *Assume that X has an fpqc cover $U \rightarrow X$, where U is a noetherian reduced scheme. Then the virtually unipotent locally free sheaves form a tannakian class.*

Proposition 12.2. *Let Γ be an affine gerbe. Then a representation of Γ is virtually unipotent if and only if it is isomorphic to a pullback from $\text{Rep } \Gamma^{\text{VU}}$.*

Proof of [Proposition 12.1](#). Let us check again that the three conditions of [Definition 11.1](#) are satisfied. This is clear for (2).

For condition (1), the only nonobvious thing to prove is that virtually unipotent locally free sheaves form an abelian subcategory; for this we need to show that the cokernel of a homomorphism of virtually unipotent locally free sheaves is again locally free and virtually unipotent.

Let $\phi: F \rightarrow G$ be a homomorphism of virtually unipotent sheaves, and let Q be its cokernel. Choose an integer n such that $(\text{Frob}_X^n)^*F$ and $(\text{Frob}_X^n)^*G$ are extended essentially finite sheaves on X ; then $(\text{Frob}_X^n)^*Q$ is the cokernel of the pullback homomorphism $(\text{Frob}_X^n)^*\phi$, so it is an extended essentially finite sheaf on X , from [Theorem 10.5\(2\)](#).

Choose an fpqc cover $\pi: U \rightarrow X$, where U is a noetherian reduced scheme. We have a commutative diagram

$$\begin{array}{ccc} U & \xrightarrow{\text{Frob}_U^n} & U \\ \downarrow \pi & & \downarrow \pi \\ X & \xrightarrow{\text{Frob}_X^n} & X \end{array}$$

Now, $\pi^*(\text{Frob}_X^n)^*Q = (\text{Frob}_U^n)^*\pi^*Q$ is locally free on U ; since Frob_U is a homeomorphism, we have that the function $U \rightarrow \mathbb{N}$ which sends $p \in U$ into $\dim_{k(p)}(\pi^*Q \otimes_{\mathcal{O}_U} k(p))$ is locally constant. We deduce that π^*Q is locally free on X ; hence Q is a virtually unipotent locally free sheaf on X , and this concludes the proof.

Let us now check condition (3). Let $f: X \rightarrow \Gamma$ be a morphism such that $f^*: \text{Rep } \Gamma \rightarrow \text{Vect}_X$ induces an equivalence between $\text{Rep } \Gamma$ and virtually unipotent locally free sheaves on X . Let E be a representation of Γ ; we need to show that E is virtually unipotent. We use induction on the rank r of E , the result being clear for $r = 0$. Assume $r > 0$, and fix an integer n such that $f^*(\text{Frob}_\Gamma^n)^*E = (\text{Frob}_X^n)^*f^*E$ is a unipotent sheaf on X . So there exists a positive integer s and a monomorphism $\mathcal{O}_X^{\oplus s} \rightarrow f^*(\text{Frob}_\Gamma^n)^*E$. Since f is fully faithful, this lifts into a monomorphism $\mathcal{O}_\Gamma^{\oplus s} \rightarrow (\text{Frob}_\Gamma^n)^*E$. Because the rank of $(\text{Frob}_\Gamma^n)^*E/\mathcal{O}_\Gamma^{\oplus s}$ is less than r , it is virtually unipotent. Since virtually unipotent sheaves are stable by extensions, we conclude that $(\text{Frob}_\Gamma^n)^*E$ is virtually unipotent, which implies in turn that E itself is virtually unipotent. \square

Proof of Proposition 12.2. The proof is somewhat long, so we split it into three steps.

Step 1: reduction to the case that Γ is of finite type.

Let us assume that the proposition holds for affine gerbes of finite type; let Γ be an arbitrary affine gerbe over κ , and let V be a representation of Γ .

If V comes from Γ^{vU} , we need to show that V is virtually unipotent. Since the pullback of a virtually unipotent representation is virtually unipotent, we can replace Γ with Γ^{vU} , and assume that Γ is virtually unipotent. Choose a locally full morphism $\Gamma \rightarrow \Delta$, where Δ is a gerbe of finite type, and a representation W of Δ whose pullback to Γ is isomorphic to V . Then Δ is virtually unipotent; hence W is virtually unipotent, and so V is virtually unipotent.

Conversely, let V be a virtually unipotent representation of Γ ; we need to show that V is a pullback from Γ^{vU} . Once again, choose a locally full morphism $\Gamma \rightarrow \Delta$, where Δ is a gerbe of finite type, and a representation W of Δ whose pullback to Γ is isomorphic to V . It is enough to show that there exists a factorization $\Gamma \rightarrow \Delta_1 \rightarrow \Delta$, where Δ_1 is again an affine gerbe of finite type, such that the pullback W_1 of W to Δ_1 is virtually unipotent. In fact, this implies that W_1 is a pullback from Δ_1^{vU} , and, since the composite $\Gamma \rightarrow \Delta_1^{\text{vU}}$ factors through Γ^{vU} , the conclusion follows.

Assume that V is an extended essentially finite sheaf. In this case we know from [Proposition 11.10](#) that V comes from Γ^{SVU} ; choose a locally full morphism $\Gamma^{\text{SVU}} \rightarrow \Theta$ and a representation Z of Θ , whose pullback to Γ is isomorphic to V . We can choose a locally full morphism $\Gamma \rightarrow \Delta_1$, where Δ_1 is an affine gerbe of finite type, such that both $\Gamma \rightarrow \Delta$ and $\Gamma \rightarrow \Theta$ factor through Δ_1 . If Z_1 denotes the pullback of Z to Δ_1 , we have that Z_1 is an extended essentially finite sheaf. The pullback of W_1 and Z_1 to Γ are both isomorphic to V ; since the pullback $\text{Rep } \Delta_1 \rightarrow \text{Rep } \Gamma$ is fully faithful, it follows that W_1 and Z_1 are isomorphic, so W_1 is an extended essentially finite sheaf.

In the general case, choose a positive integer such that $(\text{Frob}_\Gamma^m)^* V$ is an extended essentially finite sheaf; choose a factorization $\Gamma \rightarrow \Delta_1 \rightarrow \Delta$ of the desired type, such that the pullback of $(\text{Frob}_\Delta^m)^* W$ is an extended essentially finite sheaf; but this pullback is $(\text{Frob}_{\Delta_1}^m)^* W_1$, and the conclusion follows.

Step 2: reduction to the case that $\Gamma(\kappa) \neq \emptyset$.

Since by [Corollary 3.2](#) there exists a finite Galois extension κ' of κ with $\Gamma(\kappa') \neq \emptyset$, the following lemma allows us to assume that $\Gamma(\kappa) \neq \emptyset$, so that $\Gamma = \mathcal{B}_\kappa G$ for some affine group G of finite type over κ .

Lemma 12.3. *Let κ' be a finite Galois extension of κ . Let V be a representation of Γ .*

- (1) *The representation V is a pullback from Γ^{VU} if and only if the pullback $V_{\kappa'}$ of V to $\Gamma_{\kappa'}$ is a pullback from $(\Gamma_{\kappa'})^{\text{VU}}$.*
- (2) *The representation V is virtually unipotent if and only if the pullback of V to $\Gamma_{\kappa'}$ is virtually unipotent.*

Proof. Call G the Galois group of κ'/κ . By descent theory, we have an equivalence between $\text{Rep } \Gamma$ and the category of G -equivariant locally free sheaves on $\Gamma_{\kappa'}$.

For part (1), assume that $V_{\kappa'}$ comes from a representation W of $(\Gamma_{\kappa'})^{\text{VU}}$. Notice that by [Theorem 9.3](#) we have $(\Gamma_{\kappa'})^{\text{VU}} = (\Gamma^{\text{VU}})_{\kappa'}$. For each $g \in G$ we denote by $g: \Gamma_{\kappa'} \rightarrow \Gamma_{\kappa'}$ and $g: (\Gamma^{\text{VU}})_{\kappa'} \rightarrow (\Gamma^{\text{VU}})_{\kappa'}$ the induced morphisms. The G -equivariant structure of $V_{\kappa'}$ gives a collection of isomorphisms $V_{\kappa'} \simeq g^* V_{\kappa'}$ of locally free sheaves over $\Gamma_{\kappa'}$; since the pullback $\text{Rep}(\Gamma^{\text{VU}})_{\kappa'} \rightarrow \text{Rep } \Gamma_{\kappa'}$ is fully faithful, these give isomorphisms $g^*: W \simeq g^* W$ that give W a structure of G -equivariant representation on $(\Gamma^{\text{VU}})_{\kappa'}$. Thus W descends to a representation of Γ^{VU} whose pullback to Γ is isomorphic to V .

For part (2), first of all notice that the argument above also applies to the Γ^{SVU} fundamental gerbe; since the representations coming from the Γ^{SVU} are the extended essentially finite representations, this shows that if $V_{\kappa'}$ is an extended essentially finite representation, then so is V .

Now fix V a representation of Γ so that $V_{\kappa'}$ is virtually unipotent. Since the pullback $(\text{Frob}_{\Gamma_{\kappa'}}^m)^* V_{\kappa'}$ is isomorphic to $((\text{Frob}_\Gamma^m)^* V)_{\kappa'}$, we may assume that $V_{\kappa'}$ is an extended essentially finite sheaf, and the conclusion follows. □

Step 3: the conclusion.

Let us show that every locally free sheaf on $\Gamma = \mathcal{B}_\kappa G$ coming from Γ^{VU} is virtually unipotent; since being virtually unipotent is a property that is stable under pullback, we may assume that $\Gamma = \Gamma^{\text{VU}}$, so that G is virtually unipotent. Let V be a representation of G ; for $n \gg 0$ we have that the Frobenius morphism

$\text{Frob}_G^n : G \rightarrow G$ factors through G_{red} . Since G_{red} is strongly virtually unipotent, we have that the pullback $(\text{Frob}_{\mathcal{B}_\kappa G}^n)^* V$ is strongly virtually unipotent, so V is virtually unipotent.

Conversely, assume that V is a virtually unipotent representation of G . We can replace G with its image in $\text{GL}(V)$, and assume that V is faithful. Denote by $V^{(n)}$ the pullback of V under the Frobenius map $\text{Frob}_{\mathcal{B}_\kappa G}^n$; then for $n \gg 0$ the representation $V^{(n)}$ is strongly virtually unipotent. The kernel of $G \rightarrow \text{GL}(V^{(n)})$ is clearly finite; call H the image of G in $\text{GL}(V^{(n)})$; then $V^{(n)}$ is strongly virtually unipotent as a representation of H . From [Proposition 11.10](#) it follows that $V^{(n)}$ is a pullback of a representation of a strongly virtually unipotent quotient of H ; but $V^{(n)}$ is a faithful representation, so H is strongly virtually unipotent. The following lemma allows us to conclude.

Lemma 12.4. *Let G be an affine algebraic group of finite type over κ . Assume that there exists a normal finite subgroup scheme $K \subseteq G$ such that G/K is virtually unipotent. Then G is virtually unipotent.*

Proof. We can base change to the algebraic closure of κ , and assume that κ is algebraically closed. Furthermore, if $H = G/K$, then $H_{\text{red}}^0 = G_{\text{red}}^0 / (G_{\text{red}}^0 \cap K)$; so we may assume that G , and therefore H , is smooth and connected. It follows that H is unipotent; we will show that G is also unipotent.

If K is infinitesimal and we denote by $G^{[n]}$ the pullback under the isomorphism $\text{Frob}_\kappa^n : \text{Spec } \kappa \rightarrow \text{Spec } \kappa$; then the relative Frobenius homomorphism $G \rightarrow G^{[n]}$ factors through G/K for $n \gg 0$. Hence $G^{[n]}$ is a quotient of G/K , so it is unipotent; it follows that G is unipotent.

In the general case, by the previous case we can replace G by G/K^0 , and assume that K is étale. Every smooth unipotent group scheme contains a normal subgroup scheme which is isomorphic to \mathbb{G}_a ; let $H' \subseteq H = G/K$ be such a subgroup scheme. Call G' the connected component of the identity in the inverse image of H' in G , and K' the inverse image of G' in K . Then G/G' is an extension of H/H' by K/K' ; by induction on the dimension of G we may assume that G/G' is unipotent. Since every extension of unipotent groups is unipotent, it is enough to prove that G' is unipotent. However, G' is a smooth connected 1-dimensional affine group scheme, so it is isomorphic to \mathbb{G}_a or to \mathbb{G}_m . Since G/K is isomorphic to \mathbb{G}_a , G cannot be \mathbb{G}_m , so it is \mathbb{G}_a . □

This ends the proof of [Proposition 12.2](#), and of [Theorem 10.7](#). □

13. Gerbes of multiplicative type and Picard stacks

13A. Groups and gerbes of multiplicative type. Recall the following definitions.

Given an abelian group A , we consider the functor $D(A) : (\text{Aff}/\kappa)^{\text{op}} \rightarrow (\text{Ab})$ that sends a κ -algebra R into the group of homomorphisms of abelian groups $A \rightarrow R^\times$. Then $D(A)$ is an affine group scheme; if A is finitely generated, then $D(A)$ is of finite type. A group scheme over κ is called *diagonalizable* if it is isomorphic to some $D(A)$.

A group scheme G over κ is called *of multiplicative type* if it satisfies one the following equivalent conditions:

- (1) $G_{\kappa^{\text{sep}}}$ is diagonalizable.

- (2) $G_{\bar{\kappa}}$ is diagonalizable.
- (3) G_{ℓ} is diagonalizable for some extension ℓ of κ .
- (4) G is commutative, and all representations of G are semisimple.

The category of groups of multiplicative type is closed under taking subgroups, quotients and projective limits. Hence it coincides with the class of pro- \mathcal{C} -groups, where \mathcal{C} is the class of multiplicative groups of finite type over κ .

Let us set up some notation.

If R is a commutative ring, we will denote by $R_{\text{ét}}$ the small étale site of $\text{Spec } R$, which we think of as the dual of the category of étale R -algebras.

If R is a κ -algebra, we denote by $\mathcal{O}_{R_{\text{ét}}}^{\times} : R_{\text{ét}} \rightarrow (\text{Ab})$ the sheaf $A \rightarrow A^{\times}$. We shorten $\mathcal{O}_{\kappa_{\text{ét}}}^{\times}$ to \mathcal{O}^{\times} .

We denote by $\tilde{\mathcal{O}}_{R_{\text{ét}}} : \kappa_{\text{ét}} \rightarrow (\text{Ab})$ the pushforward of $\mathcal{O}_{R_{\text{ét}}}^{\times}$ to $\kappa_{\text{ét}}$ via the morphism $R_{\text{ét}} \rightarrow \kappa_{\text{ét}}$ induced by the homomorphism $\kappa \rightarrow R$; this is the sheaf that sends an étale κ -algebra A into the group $(A \otimes_{\kappa} R)^{\times}$. A homomorphism of κ -algebras $R \rightarrow S$ induces a homomorphism of sheaves $\tilde{\mathcal{O}}_{R_{\text{ét}}} \rightarrow \tilde{\mathcal{O}}_{S_{\text{ét}}}$.

Recall that the category of groups of multiplicative type over κ is antiequivalent to the category of sheaves of abelian groups over the small étale site $\kappa_{\text{ét}}$ of $\text{Spec } \kappa$; we will mostly think of $\kappa_{\text{ét}}$ as the dual of the category of étale κ -algebras. With a group of multiplicative type G we associate the sheaf of characters $\widehat{G} : \kappa_{\text{ét}} \rightarrow (\text{Ab})$ defined as the functor sending each étale κ -algebra A to the group of characters $\text{Hom}_A(G_A, \mathcal{O}_A^{\times})$.

In the other direction, given a sheaf F on $\kappa_{\text{ét}}$ we define a functor

$$D(F) : (\text{Aff}/\kappa)^{\text{op}} \longrightarrow (\text{Ab})$$

by sending a κ -algebra R into the group $\text{Hom}(F, \tilde{\mathcal{O}}_{R_{\text{ét}}})$.

Another way of stating this equivalence is the following. Let \mathcal{G} be the Galois group of κ^{sep} over κ . The abelian group $M := \text{Hom}(G_{\kappa^{\text{sep}}}, \mathbb{G}_{\text{m}, \kappa^{\text{sep}}})$ has a natural continuous action of \mathcal{G} ; sending G into M gives an equivalence between the opposite of the category of group schemes of multiplicative type over κ with that of abelian groups with a continuous action of \mathcal{G} .

We will call a pro- \mathcal{C} -gerbe, where \mathcal{C} is the class of groups of multiplicative type and finite type, a *gerbe of multiplicative type*. Equivalently, a gerbe of multiplicative type is an affine gerbe banded by a group of multiplicative type, not necessarily of finite type.

The aim of this section is give a description of gerbes of multiplicative type that is very similar in spirit to that of groups of multiplicative type given above.

13B. Picard stacks. In this subsection we recall some known facts about Picard stacks.

By a *Picard stack* P over κ we will mean, as in [SGA 4₃ 1973, Exposé XVIII], a stack in strictly commutative monoidal groupoids $P \rightarrow \kappa_{\text{ét}}$, whose operation $P \times_{\kappa_{\text{ét}}} P \rightarrow P$ is denoted by $(\xi, \eta) \mapsto \xi \otimes \eta$, such that every object of P is invertible. We will denote by $\varepsilon : \kappa_{\text{ét}} \rightarrow P$ the section corresponding to the identity; if $A \in \kappa_{\text{ét}}$, the image of A in $P(A)$ will be denoted by ε_A . If $\phi : A \rightarrow B$ is a homomorphism of étale κ -algebras, the image of the corresponding arrow $\text{Spec } B \rightarrow \text{Spec } A$ in $\kappa_{\text{ét}}$ will be denoted by $\varepsilon_{\phi} : \varepsilon_B \rightarrow \varepsilon_A$.

We denote by $\pi_0(P)$ the sheafification of the presheaf on $\kappa_{\text{ét}}$ sending A into the abelian group of isomorphism classes in $P(A)$. By $\pi_1(P)$ we denote the sheaf of automorphisms of the identity section $\varepsilon : \kappa_{\text{ét}} \rightarrow P$. Because of the monoidal structure of P , the sheaf of groups $\pi_1(P)$ is abelian. The inverse image of the identity in $\pi_0(P)$ in P is equivalent to the classifying stack $\mathcal{B}_{\kappa_{\text{ét}}}\pi_1(P)$.

Furthermore, if $\xi \in P(A)$ we have a homomorphism of groups

$$\pi_1(P)(A) = \text{Aut}_A(\varepsilon_A) \longrightarrow \text{Aut}_A(\varepsilon_A \otimes \xi)$$

sending α into $\alpha \otimes \text{id}_\xi$; by using the given isomorphism $\varepsilon_A \otimes \xi \simeq \xi$ we obtain a group homomorphism $\pi_1(P)(A) \rightarrow \text{Aut}_A \xi$, which is in fact an isomorphism. This gives an equivalence of Picard stacks between the inertia I_P and the product $\pi_1(P) \times P$.

The rigidification $P // \pi_1(P)$ (in the sense of [Abramovich et al. 2008, Appendix A]) is a sheaf, isomorphic to $\pi_0(P)$. Thus, P can be thought of as a central extension of the sheaf $\pi_0(P)$ by the classifying stack $\mathcal{B}\pi_1(P)$.

A homomorphism $P \rightarrow Q$ of Picard stacks is an equivalence if and only if the induced homomorphisms of sheaves $\pi_1(P) \rightarrow \pi_1(Q)$ and $\pi_0(P) \rightarrow \pi_0(Q)$ are isomorphisms.

Picard stacks over $\kappa_{\text{ét}}$ form a strict 2-category, whose 1-arrows are symmetric monoidal base-preserving functors; we call these *homomorphisms of Picard stacks*.

In [SGA 4₃ 1973, Exposé XVIII], Deligne also showed how to describe Picard stacks in terms of complexes of sheaves of abelian groups: a Picard stack is a quotient $[L^0/L^{-1}]$ where L^\bullet is a complex of sheaves on $\kappa_{\text{ét}}$ concentrated in degrees -1 and 0 , and L^{-1} is an injective sheaf. Such complexes form a 2-category, the 2-arrows being given by homotopies. Sending L^\bullet into $[L^0/L^{-1}]$ gives an equivalence between the 2-category of complexes of this type, and the 2-category of Picard stacks.

If we fix two complexes of sheaves G_0 and G_1 , consider the 2-category of 2-extensions of G_0 by G_1 . An object of this 2-category is a Picard stack P , with a fixed isomorphism $\pi_0(P) \simeq G_0$ and $\pi_1(P) \simeq G_1$. As a corollary of Deligne’s result, we have that equivalence classes of such extensions are parametrized by $\text{Ext}_{\kappa_{\text{ét}}}^2(G_0, G_1)$.

13C. \mathcal{O}^\times -stacks. A simple example of a Picard stack is the classifying stack

$$\mathcal{B}_{\kappa_{\text{ét}}}\mathcal{O}^\times \longrightarrow \kappa_{\text{ét}};$$

it is the Picard stack in which every $\mathcal{B}_{\kappa_{\text{ét}}}\mathcal{O}^\times(A)$ has a unique object, whose automorphism group is A^\times . Notice that it is a stack in the étale topology, because every étale κ -algebra has trivial Picard group.

We will often shorten $\mathcal{B}_{\kappa_{\text{ét}}}\mathcal{O}^\times$ to $\mathcal{B}\mathcal{O}^\times$.

Definition 13.1. An \mathcal{O}^\times -stack over κ is a Picard stack P over κ , together with a homomorphism of sheaves of groups $\rho_P : \mathcal{O}^\times \rightarrow \pi_1(P)$.

An \mathcal{O}^\times -stack is *rigid* if $\rho_P : \mathcal{O}^\times_{\kappa_{\text{ét}}} \rightarrow \pi_1(P)$ is an isomorphism.

Equivalently, an \mathcal{O}^\times -stack is a Picard stack with a homomorphism of Picard stacks $\mathcal{B}_{\kappa_{\text{ét}}}\mathcal{O}^\times \rightarrow P$.

There is an obvious 2-category of \mathcal{O}^\times -stacks over κ : a homomorphism of \mathcal{O}^\times -stacks $\phi: P \rightarrow Q$ is a homomorphism of Picard stacks, such that the composite $\mathcal{O}^\times \xrightarrow{\rho_P} \pi_1(P) \xrightarrow{\pi_1(\phi)} \pi_1(Q)$ equals ρ_Q .

A homomorphism $P \rightarrow Q$ of rigid \mathcal{O}^\times -stacks is an equivalence if and only if the induced homomorphism $\pi_0(P) \rightarrow \pi_0(Q)$ is an isomorphism.

Every rigid \mathcal{O}^\times -stack is a quotient stack $[L^0/L^{-1}]$, with a fixed isomorphism $\ker(L^{-1} \rightarrow L^0) \simeq \mathcal{O}^\times$. We will call complexes of this type \mathcal{O}^\times -complexes. They form a 2-category; the 2-arrows are given by homotopies. We have a functor from the 2-category of \mathcal{O}^\times -complexes into \mathcal{O}^\times -stacks that sends L^\bullet to $[L^0/L^{-1}]$; restricting this functor to \mathcal{O}^\times -complexes L^\bullet with L^{-1} injective gives an equivalence of 2-categories.

This implies that given a sheaf F on $\kappa_{\text{ét}}$, equivalence classes of rigid \mathcal{O}^\times -stacks P with a fixed isomorphism $\pi_0(P) \simeq F$ are classified by $\text{Ext}^2(F, \mathcal{O}^\times)$, where the Ext^2 is taken in the category of sheaves on $\kappa_{\text{ét}}$.

13D. From gerbes to rigid \mathcal{O}^\times -stacks. Let $X \rightarrow (\text{Aff}/\kappa)$ be a fibered category. If A is an étale κ algebra, the composite $X_A \rightarrow (\text{Aff}/A) \rightarrow (\text{Aff}/\kappa)$ makes X_A into a category fibered over (Aff/κ) .

The Picard stack $\underline{\text{Pic}}_{X/\kappa} = \underline{\text{Pic}}_X \rightarrow \kappa_{\text{ét}}$ is the fibered category corresponding to the pseudofunctor on $\kappa_{\text{ét}}$ that sends A into the groupoid $\underline{\text{Pic}}(X_A)$ of invertible sheaves on X_A . It can be conveniently defined as the stack of morphism $X_A \rightarrow \mathcal{B}_\kappa \mathbb{G}_m$.

The stack $\underline{\text{Pic}}_X$ has a canonical structure of \mathcal{O}^\times -stack. Furthermore, if X is concentrated and $H^0(X, \mathcal{O}) = \kappa$, then $\underline{\text{Pic}}_X$ is rigid.

In particular, if Γ is a gerbe of multiplicative type over κ , its Picard stack $\underline{\text{Pic}}_\Gamma \rightarrow \kappa_{\text{ét}}$ is a rigid \mathcal{O}^\times -stack. Notice that $\underline{\text{Pic}}_{(\text{Aff}/\kappa)}$ is equivalent to $\mathcal{B}\mathcal{O}^\times$.

A morphism of fibered categories $X \rightarrow Y$ over (Aff/κ) yields a homomorphism of \mathcal{O}^\times -stacks $\underline{\text{Pic}}_Y \rightarrow \underline{\text{Pic}}_X$; this defines a strict 2-functor from the 2-category of fibered categories over (Aff/κ) to the 2-category of \mathcal{O}^\times -stacks over κ .

If R is a κ algebra, we set $\underline{\text{Pic}}_R := \underline{\text{Pic}}_{\text{Spec } R}$; then $\pi_1(\underline{\text{Pic}}_R) = \tilde{\mathcal{O}}_{R_{\text{ét}}}$.

Proposition 13.2. *Let $G \rightarrow \text{Spec } \kappa$ be a group of multiplicative type. Then*

$$\underline{\text{Pic}}_{\mathcal{B}_\kappa G} \simeq \mathcal{B}_{\kappa_{\text{ét}}} \mathcal{O}^\times \times \widehat{G}.$$

Proof. There is an equivalence between invertible sheaves on $\mathcal{B}_A G_A = (\mathcal{B}_\kappa G)_A$ and morphisms $\mathcal{B}_A G_A \rightarrow \mathcal{B}_A \mathbb{G}_{m,A}$ over (Aff/A) . If A is an étale κ -algebra, by [Giraud 1971, III, Remarque 1.6.7] the category of such morphisms is equivalent to the category of global objects of the stack

$$\mathcal{B}_\kappa \mathbb{G}_m(A) \times \widehat{G}(A) \simeq (\mathcal{B}_{\kappa_{\text{ét}}} \mathcal{O}^\times \times \widehat{G})(A),$$

and this gives the desired equivalence. □

For a general gerbe we have the following.

Proposition 13.3. *Let Γ be a gerbe of multiplicative type; call G its band. Then $\pi_0(\underline{\text{Pic}}_\Gamma)$ is canonically isomorphic to \widehat{G} .*

Proof. Suppose that $\Gamma(\kappa) \neq \emptyset$; then the choice of an object ξ of $\Gamma(\kappa)$ gives an isomorphism between $\pi_0(\underline{\text{Pic}}_\Gamma)$ and \widehat{G} , where $G = \underline{\text{Aut}}_\kappa \xi$. This is independent of the choice of ξ ; it is also functorial under finite separable extensions of the base field.

In the general case, for each separable extension κ'/κ such that $G(\kappa') \neq \emptyset$ we obtain an isomorphism between the band of $\Gamma_{\kappa'}$, which is the restriction of the band of Γ , and $\widehat{G}_{\kappa'}$. These isomorphisms are canonical, and descend to the desired isomorphism between the band of Γ and \widehat{G} . \square

Consider a cofiltered system $\{\Gamma_i\}_{i \in I}$ of affine gerbes over κ , and set $\Gamma := \varprojlim_i \Gamma_i$. For each arrow $j \rightarrow i$ in I we have a morphism $\Gamma_j \rightarrow \Gamma_i$, and hence a homomorphism of \mathcal{O}^\times -stacks $\underline{\text{Pic}}_{\Gamma_i} \rightarrow \underline{\text{Pic}}_{\Gamma_j}$. This gives a strict 2-functor from I^{op} to Picard stacks. The projection $\varprojlim_i \Gamma_i \rightarrow \Gamma_i$ induces a homomorphism of Picard stacks $\underline{\text{Pic}}_{\Gamma_i} \rightarrow \underline{\text{Pic}}_{\varprojlim_i \Gamma_i}$, and consequently a homomorphism $\varinjlim_i \underline{\text{Pic}}_{\Gamma_i} \rightarrow \underline{\text{Pic}}_{\varprojlim_i \Gamma_i}$.

Proposition 13.4. *The homomorphism $\varinjlim_i \underline{\text{Pic}}_{\Gamma_i} \rightarrow \underline{\text{Pic}}_{\varprojlim_i \Gamma_i}$ above is an equivalence.*

Proof. This follows from [Borne and Vistoli 2015, Proposition 3.8]. \square

13E. From rigid \mathcal{O}^\times -stacks to gerbes. We can also go from \mathcal{O}^\times -stacks P to fibered categories over (Aff/κ) in the following way. Let P be an \mathcal{O}^\times -stack. Let us define a fibered category $\underline{\text{Ger}}_P \rightarrow (\text{Aff}/\kappa)$ as follows. If R is a κ -algebra, an object of $\underline{\text{Ger}}_P(R)$ is a homomorphism of \mathcal{O}^\times -stacks $P \rightarrow \underline{\text{Pic}}_R$. A morphism $\text{Spec } S \rightarrow \text{Spec } R$ in (Aff/κ) induces a pullback functor $\underline{\text{Pic}}_R \rightarrow \underline{\text{Pic}}_S$. Composing with this gives the function $R \mapsto \underline{\text{Ger}}_P(R)$ the structure of a pseudofunctor; we define $\underline{\text{Ger}}_P$ to be the associated fibered category. (The notation here is a little improper, as $\underline{\text{Ger}}_P$ is not necessarily a gerbe if P is not rigid.)

This construction gives a strict 2-functor from the 2-category of \mathcal{O}^\times -stacks to that of stacks in groupoids on (Aff/κ) .

Proposition 13.5. *Assume that P is a rigid \mathcal{O}^\times -stack; then the fibered category $\underline{\text{Ger}}_P$ is an affine gerbe. Its band is the group scheme of multiplicative type $\text{D}(\pi_0(P))$ corresponding to $\pi_0(P)$.*

Proof. We will divide the proof into several steps.

Step 1: $\underline{\text{Ger}}_P$ is an fpqc stack.

This is a straightforward exercise in descent theory.

Step 2: formation of $\underline{\text{Ger}}_P$ commutes with separable algebraic extensions of the base field.

Suppose that ℓ is a separable algebraic extension of κ . The obvious functor $\kappa_{\text{ét}} \rightarrow \ell_{\text{ét}}$ sending A into $A \otimes_\kappa \ell$ induces a morphism of sites $\phi: \ell_{\text{ét}} \rightarrow \kappa_{\text{ét}}$, which gives a pullback functor ϕ^{-1} from sheaves, or stacks, over $\kappa_{\text{ét}}$ to sheaves, or stacks, over $\ell_{\text{ét}}$. Since $\phi^{-1} \mathcal{O}_{\kappa_{\text{ét}}} = \mathcal{O}_{\ell_{\text{ét}}}$, we have that $P_\ell := \text{Spec } \ell \times_{\text{Spec } \kappa} P$ is an \mathcal{O}^\times -stack over ℓ . It is immediate to conclude that $\underline{\text{Ger}}_{P_\ell} = \text{Spec } \ell \times_{\text{Spec } \kappa} \underline{\text{Ger}}_P$.

Step 3: describing $\underline{\text{Ger}}_P$ in terms of complexes of sheaves.

Let R be a κ -algebra; call $\phi: R_{\text{ét}} \rightarrow \kappa_{\text{ét}}$ the morphism of sites induced by the homomorphism $\kappa \rightarrow R$. Fix an embedding $\mathcal{O}_{R_{\text{ét}}}^\times \subseteq I^{-1}$ into an injective sheaf, set $I^0 := I^{-1}/\mathcal{O}_{R_{\text{ét}}}^\times$, and extend the projection

$I^{-1} \rightarrow I^0$ into a complex I^\bullet . Then, under Deligne’s correspondence the Picard stack $\underline{\text{Pic}}_{R/\kappa}$ corresponds to the complex $\phi_* I^\bullet$.

Now, represent P as a quotient $[L^0/L^{-1}]$, where L^\bullet is an \mathcal{O}^\times -complex. From the adjunction between pushforward and pullback, we obtain that $\underline{\text{Ger}}_P(R)$ is equivalent to the category that has the following description:

- (a) An object is a homomorphism $\xi : \phi^{-1} L^\bullet \rightarrow I^\bullet$ of sheaves on $\kappa_{\text{ét}}$ such that $H^{-1}(\xi) : \phi^{-1} \mathcal{O}_{\kappa_{\text{ét}}}^\times \rightarrow \mathcal{O}_{R_{\text{ét}}}^\times$ is the canonical homomorphism.
- (b) An arrow $\xi \rightarrow \eta$ is a homomorphism of graded sheaves $a : \phi^{-1} L^\bullet \rightarrow I^\bullet[-1]$ such that

$$\eta - \xi = \delta_{I^\bullet} \circ a + a \circ \delta_{\phi^{-1} L^\bullet}.$$

Step 4: identifying the sheaf of automorphisms of an object of $\underline{\text{Ger}}_P(R)$.

From the description above it is immediate to check the automorphism group of an object ξ is the group

$$\begin{aligned} \underline{\text{Hom}}_{R_{\text{ét}}}(\text{coker}(\phi^{-1} L^{-1} \rightarrow \phi^{-1} L^0), \mathcal{O}_R^\times) &= \underline{\text{Hom}}_{R_{\text{ét}}}(\phi^{-1} \pi_0(P), \mathcal{O}_R^\times) \\ &= \underline{\text{Hom}}_{\kappa_{\text{ét}}}(\pi_0(P), \tilde{\mathcal{O}}_{R_{\text{ét}}}) \\ &= \text{D}(\pi_0(P)) \end{aligned}$$

as claimed.

Step 5: $\underline{\text{Ger}}_P$ is nonempty.

Let us show that $\underline{\text{Ger}}_P(\kappa^{\text{sep}}) \neq \emptyset$, where κ^{sep} is a separable closure of κ . By Step 2 we can base-change to κ^{sep} , and assume that κ is separably closed. In this case $\text{Ext}^2(\pi_0(P), \mathcal{O}^\times) = 0$, so that $P \simeq \mathcal{B}\mathcal{O}^\times \times \pi_0(P)$, and composing the projection $P \rightarrow \mathcal{B}\mathcal{O}^\times$ with the natural homomorphism of Picard stacks $\mathcal{B}\mathcal{O}^\times \rightarrow \underline{\text{Pic}}_{\text{Spec } \kappa}$, we obtain an object of $\underline{\text{Ger}}_P(\kappa)$.

Step 6: two objects of $\underline{\text{Ger}}_P$ are fpqc locally isomorphic.

Once again, we may assume that κ is separably closed, so that P is of the form $\mathcal{B}\mathcal{O}^\times \times \pi_0(P)$. Let R be a κ -algebra, ξ and η two objects of $\underline{\text{Ger}}_P(R)$.

Lemma 13.6. *For any κ -algebra R , there exists a faithfully flat extension $R \subseteq S$ such that*

- (a) *every faithfully flat étale ring homomorphism $S \rightarrow T$ has a retraction $T \rightarrow S$, and*
- (b) *the abelian group S^\times is divisible.*

Proof. For part (a) we will switch to the language of affine schemes, as this seems to make the proof somewhat more intuitive. Set $X := \text{Spec } R$. First of all, let us show there exists an faithfully flat map $Y \rightarrow X$, where Y is an affine scheme with the property that every étale surjective map $Z \rightarrow Y$, where Z is an affine scheme, has a section $Y \rightarrow Z$.

The construction is straightforward, and mirrors the standard construction of the strict henselization at a point. The existence of S also follows from [Bhatt and Scholze 2015, Corollary 2.2.14]; for the

convenience of the reader, we give a direct proof. For each $p \in X$ choose a separable closure $\kappa(p)^{\text{sep}}$, and set $X_0 := \bigsqcup_{p \in X} \text{Spec}(\kappa(p)^{\text{sep}})$. Let us define a category I whose objects (U, f, a) are commutative diagrams

$$\begin{array}{ccc} & & U \\ & \nearrow a & \downarrow f \\ X_0 & \longrightarrow & X \end{array}$$

in which the bottom morphism is the obvious map, and the vertical one is an étale map with U affine. A map from (U, f, a) to (V, g, b) is, naturally, a morphism $\phi: U \rightarrow V$ with $f = g\phi$ and $b = \phi a$. This category has fibered products and a terminal object, so it is cofiltered. Define $Y = \varprojlim_I U$; the natural map $Y \rightarrow X$ is faithfully flat. If $Z \rightarrow Y$ is an étale surjective map with Z affine, then Z is a pullback $W \times_U Y$ of an étale surjective map $\phi: W \rightarrow U$ for some object (U, f, a) of I [Stacks 2005–, Tag 00U2]. The map $a: X_0 \rightarrow U$ lifts to a map $b: X_0 \rightarrow W$; so ϕ is an arrow $(W, f\phi, b) \rightarrow (U, f, a)$. Hence $Y \rightarrow U$ lifts to $Y \rightarrow W$, and this gives a section $Y \rightarrow Z$.

Now, set $S := \mathcal{O}(Y)$; this S has property (a). If $\text{char } \kappa = 0$ and $f \in S^\times$, the extension $S \subseteq S[x]/(x^n - f)$ is étale; hence it has a section $S[x]/(x^n - f) \rightarrow S$, which means exactly that f has an n -th root, so S also has property (b).

Assume that $\text{char } \kappa = p > 0$; in this case the argument above only works when p does not divide n , and we need to extend S further, so that it has p^n -th roots of all the elements of S^\times for all n . Let us start by replacing R with S , and assume that R has property (a).

Consider the set J of finite subsets $S \subseteq R^\times$, ordered by inclusion. For each $S \in J$ define an R -algebra $R[S]$ by adding an indeterminate x_f for each $f \in S$, and dividing by the ideal generated by the polynomials $x_f^p - f$ for all $f \in S$. The algebra $R[S]$ is clearly faithfully flat, and the induced morphism $\text{Spec } R[S] \rightarrow \text{Spec } R$ is a universal homeomorphism.

If $S \subseteq T$ there is an induced homomorphism $R[S] \rightarrow R[T]$ defined by sending $[x_f] \in R[S]$ into $[x_f] \in R[T]$ for each $f \in S$. Set

$$R_1 := \varinjlim_{S \in J} R[S];$$

from the construction it is clear that every $f \in R^\times$ has a p -th root in R_1 . Also, R_1 is faithfully flat over R , and the induced morphism $\text{Spec } R_1 \rightarrow \text{Spec } R$ is a universal homeomorphism.

Let us iterate this construction: for each $n > 0$ define $R_n := (R_{n-1})_1$. We get a sequence of faithfully flat extensions $R \subseteq R_1 \subseteq \dots \subseteq R_n \subseteq \dots$ such that every induced morphism $\text{Spec } R \rightarrow \text{Spec } R_{n-1}$ is a universal homeomorphism, and every element of R_{n-1}^\times has p -th root in R_n . Set $S := \varinjlim_n R_n$; it follows that S^\times is p -divisible.

Furthermore, S is faithfully flat over R , and $\text{Spec } S \rightarrow \text{Spec } R$ is a universal homeomorphism; from this it follows that pullback induces an equivalence between affine étale surjective maps to $\text{Spec } R$ and affine étale surjective maps to $\text{Spec } S$. Since R has property (a), it follows that S has it too. But this implies that S^\times is n -divisible for all n not divisible by p . □

So, we may assume that $R = S$. Consider the morphism of sites $\phi: R_{\text{ét}} \rightarrow \kappa_{\text{ét}}$ given by the homomorphism $\kappa \rightarrow R$. Condition (a) of Lemma 13.6 implies that sending a sheaf F into its group of global sections is an exact functor; in other words, the pushforward ϕ_* from sheaves on $R_{\text{ét}}$ to $\kappa_{\text{ét}}$ is exact.

Choose an injective sheaf I^{-1} containing $\mathcal{O}_{R_{\text{ét}}}^\times$, and set $I^0 := I^{-1}/\mathcal{O}_{R_{\text{ét}}}^\times$, as in Step 3; hence $\phi_* I^\bullet$ is an injective resolution of R^\times . Now, P is represented by the complex $\mathcal{O}^\times \xrightarrow{0} \pi_0(P)$, where $\pi_0(P)$ is in degree 0; hence the difference $\eta - \xi$ gives a homomorphism of sheaves from $\mathcal{O}^\times \xrightarrow{0} \pi_0(P)$ to $\phi_* I^\bullet$, which is 0 in degree -1 . We need to show that this is homotopic to 0, which is clear, because the sequence

$$0 \longrightarrow R^\times \longrightarrow \phi_* I^{-1} \longrightarrow \phi_* I^0 \longrightarrow 0$$

is split exact, since, by condition (b) of Lemma 13.6, the abelian group R^\times is injective.

This ends the proof of Proposition 13.5. □

Thus we get a functor from the 2-category of rigid \mathcal{O}^\times -stacks to the category of gerbes of multiplicative type sending P into $\underline{\text{Ger}}_P$.

As a particular case of Proposition 13.5, we obtain the following description of the gerbe associated with a split \mathcal{O}^\times stack $\mathcal{B}\mathcal{O}^\times \times A$.

Corollary 13.7. *Let A be a sheaf of abelian groups in $\kappa_{\text{ét}}$. Then $\underline{\text{Ger}}_{\mathcal{B}\mathcal{O}^\times \times A}$ is canonically equivalent to $\mathcal{B}_\kappa \text{D}(A)$.*

Proof. The projection $\mathcal{B}\mathcal{O}^\times \times A \rightarrow \mathcal{B}\mathcal{O}^\times = \underline{\text{Pic}}_\kappa$ gives an object $\xi \in \underline{\text{Ger}}_{\mathcal{B}\mathcal{O}^\times \times A}(\kappa)$, and hence an equivalence $\underline{\text{Ger}}_{\mathcal{B}\mathcal{O}^\times \times A} \simeq \mathcal{B} \underline{\text{Aut}}_\kappa \xi$. But according to Proposition 13.5 $\underline{\text{Aut}}_\kappa \xi$ is canonically isomorphic to $\text{D}(A)$. □

The main result. If X is a concentrated fibered category, it is easy to construct a functor $\Phi_X: X \rightarrow \underline{\text{Ger}}_{\underline{\text{Pic}}_X}$. Suppose that we are given an object ξ of $X(R)$, corresponding to a morphism $\text{Spec } R \rightarrow X$, which in turn gives a symmetric monoidal functor $\underline{\text{Pic}}_X \rightarrow \underline{\text{Pic}}_R$, which is, by definition, an object of $\underline{\text{Ger}}_{\underline{\text{Pic}}_X}(R)$. This gives a function on the objects, that extends easily to a base-preserving functor $\Phi_X: X \rightarrow \underline{\text{Ger}}_{\underline{\text{Pic}}_X}$.

A morphism of fibered categories $X \rightarrow Y$ gives morphisms of fibered categories $\underline{\text{Pic}}_Y \rightarrow \underline{\text{Pic}}_X$ and $\underline{\text{Ger}}_{\underline{\text{Pic}}_X} \rightarrow \underline{\text{Ger}}_{\underline{\text{Pic}}_Y}$; it is straightforward to show that Φ is a natural transformation of 2-functors from the identity to $\underline{\text{Ger}}_{\underline{\text{Pic}}_\gamma}$.

Proposition 13.8. *If Γ is a gerbe of multiplicative type, the morphism $\Phi_\Gamma: \Gamma \rightarrow \underline{\text{Ger}}_{\underline{\text{Pic}}_\Gamma}$ is an equivalence.*

Proof. We split the proof into three steps.

Step 1: Assume that $\Gamma(\kappa) \neq \emptyset$, so that $\Gamma \simeq \mathcal{B}_\kappa G$ for some group scheme of multiplicative type G over κ . By Proposition 13.2 we have a canonical equivalence $\underline{\text{Pic}}_{\mathcal{B}_\kappa} \simeq \mathcal{B}\mathcal{O}^\times \times \widehat{G}$; by Corollary 13.7 we have $\underline{\text{Ger}}_{\underline{\text{Pic}}_{\mathcal{B}_\kappa}} \simeq \mathcal{B}_\kappa \text{D}(\widehat{G})$. The composite $\mathcal{B}_\kappa G \rightarrow \mathcal{B}_\kappa \text{D}(\widehat{G})$ is easily seen to be isomorphic to the functor induced by the canonical map $G \rightarrow \text{D}(\widehat{G})$, which is an isomorphism; this proves the result.

Step 2: Assume that Γ is of finite type over κ . Then, by Proposition 3.1 we have that Γ is a smooth algebraic stack over κ ; hence there exists a finite separable extension κ'/κ such that $\Gamma(\kappa') \neq \emptyset$.

Since Γ and $\underline{\text{Ger}}_{\underline{\text{Pic}}_\Gamma}$ are fpqc stacks, is enough to show that the morphism $(\Phi_X)_{\kappa'} : \Gamma_{\kappa'} \rightarrow (\underline{\text{Ger}}_{\underline{\text{Pic}}_\Gamma})_{\kappa'}$ of gerbes over κ' is an equivalence. But it is obvious that $\underline{\text{Pic}}_{\Gamma_{\kappa'}}$ is the restriction of $\underline{\text{Pic}}_\Gamma$ to $\kappa'_{\text{ét}}$; hence $(\underline{\text{Ger}}_{\underline{\text{Pic}}_\Gamma})_{\kappa'} = \underline{\text{Ger}}_{\underline{\text{Pic}}_{\Gamma_{\kappa'}}}$. So, since $\Gamma_{\kappa'}(\kappa') \neq \emptyset$, the result follows from the first step.

Step 3: In the general case, write Γ as a projective limit $\varprojlim_i \Gamma_i$ of affine gerbes of finite type over κ . Then by Proposition 13.4 we have that $\underline{\text{Pic}}_\Gamma \simeq \varinjlim_i \underline{\text{Pic}}_{\Gamma_i}$, and it is easy to see that $\underline{\text{Ger}}_{\varinjlim_i \underline{\text{Pic}}_{\Gamma_i}} \simeq \varprojlim_i \underline{\text{Ger}}_{\underline{\text{Pic}}_{\Gamma_i}}$. \square

If P is an \mathcal{O}^\times -stack, let us construct a homomorphism of \mathcal{O}^\times -stacks $\Psi_P : P \rightarrow \underline{\text{Pic}}_{\underline{\text{Ger}}_P}$. Suppose that η is an object of $P(A)$ for some étale κ -algebra A ; we need to define an object $\Psi_P(\eta) \in \underline{\text{Pic}}_{\underline{\text{Ger}}_P}(A)$, that is, a morphism of fibered categories $\Psi_P(\eta) : (\underline{\text{Ger}}_P)_A \rightarrow \mathcal{B}_\kappa \mathbb{G}_m$.

Assume that we have an object ϕ of $(\underline{\text{Ger}}_P)_A(R)$; this consists of an A -algebra structure on the κ -algebra R , and an object of ϕ of $\underline{\text{Ger}}_P(R)$, that is, a symmetric monoidal functor $\phi : P \rightarrow \underline{\text{Pic}}_R$. By applying ϕ to η we obtain an element $\phi(\eta)$ of $\underline{\text{Pic}}_R(A) = \underline{\text{Pic}}(A \otimes R)$; we define $\Psi_P(\eta)(\phi) := \phi(\eta)$ as the image in $\underline{\text{Pic}}(R)$ of $\phi(\eta)$, via the functor $\underline{\text{Pic}}(A \otimes R) \rightarrow \underline{\text{Pic}}(R)$ induced by the product homomorphism $A \otimes R \rightarrow R$ (the one that gives the A -algebra structure on R). This defines $\Psi_P(\eta)$ at the level of objects. This is easily extended to a morphism of fibered categories.

We leave to the reader the straightforward, but dull, task of defining Ψ_P as a symmetric monoidal functor, and to check that this makes Ψ into a natural transformation of 2-functors from the identity to $\underline{\text{Pic}}_{\underline{\text{Ger}}_P}$.

Proposition 13.9. *If P is rigid, the homomorphism $\Psi_P : P \rightarrow \underline{\text{Pic}}_{\underline{\text{Ger}}_P}$ is an equivalence.*

Proof. We need to check that the homomorphism $\pi_0(P) \rightarrow \pi_0(\underline{\text{Pic}}_{\underline{\text{Ger}}_P})$ induced by Ψ_P is an isomorphism. It follows from Propositions 13.3 and 13.5 that $\pi_0(\underline{\text{Pic}}_{\underline{\text{Ger}}_P})$ is canonically isomorphic to $\widehat{D(\pi_0(P))}$, and one checks that the homomorphism $\pi_0(P) \rightarrow \widehat{D(\pi_0(P))}$ induced by Ψ_P is the biduality map, which is an isomorphism. \square

Propositions 13.8 and 13.9 imply the following.

Theorem 13.10. *Sending a gerbe Γ of multiplicative type into $\underline{\text{Pic}}_\Gamma$, and a rigid \mathcal{O}^\times -stack P into $\underline{\text{Ger}}_P$, gives an equivalence between the 2-category of gerbes of multiplicative type and the opposite of the 2-category of rigid \mathcal{O}^\times -stacks.*

Here is the main result of this section.

Theorem 13.11. *Let $X \rightarrow (\text{Aff}/\kappa)$ be a concentrated fibered category such that $H^0(X, \mathcal{O}) = \kappa$. Then the morphism $\Phi_X : X \rightarrow \underline{\text{Ger}}_{\underline{\text{Pic}}_X}$ makes $\underline{\text{Ger}}_{\underline{\text{Pic}}_X}$ into the fundamental gerbe of X for the class of group schemes of multiplicative type.*

Proof. Let $\Gamma \rightarrow (\text{Aff}/\kappa)$ be a gerbe of multiplicative type; we need to show that the functor

$$\text{Hom}(\underline{\text{Ger}}_{\underline{\text{Pic}}_X}, \Gamma) \longrightarrow \text{Hom}(X, \Gamma)$$

induced by composition with $\Phi_X : X \rightarrow \underline{\text{Ger}}_{\underline{\text{Pic}}_X}$ is an equivalence.

Set $P := \underline{\text{Pic}}_\Gamma$; by [Theorem 13.10](#) we have $\Gamma \simeq \underline{\text{Ger}}_P$. There is a 2-commutative diagram of functors

$$\begin{array}{ccc} \text{Hom}(\underline{\text{Ger}}_{\underline{\text{Pic}}_X}, \Gamma) & \xrightarrow{\quad\quad\quad} & \text{Hom}(X, \Gamma) \\ & \searrow \quad \swarrow & \\ & \text{Hom}(P, \underline{\text{Pic}}_X)^{\text{op}} & \end{array}$$

The leftmost arrow is an equivalence, by [Theorem 13.10](#); hence to prove the theorem it is enough to show that the rightmost arrow is an equivalence. This holds for any concentrated stack, as the following lemma states; this completes the proof of the theorem.

Lemma 13.12. *Let X be a concentrated stack and P a rigid \mathcal{O}^\times -stack. Then the natural functor*

$$\text{Hom}(X, \underline{\text{Ger}}_P) \longrightarrow \text{Hom}(P, \underline{\text{Pic}}_X)$$

is an equivalence.

Proof. If $X = \text{Spec } R$, this follows from the definition of $\underline{\text{Ger}}_P$, and Yoneda’s lemma. The general case follows easily from this. □

14. Acknowledgments

We are grateful to Sylvain Brochard and Lei Zhang for very useful discussions. We heartily thank Marta Pieropan, who pointed out to us the possible connection of our theory of fundamental gerbes with the theory of the universal torsor of Colliot-Thélène and Sansuc.

We are especially indebted to Fabio Tonini for several helpful remarks.

The fact that [Theorem 6.12](#) should be true was pointed out to us by Andrei Okounkov and Johan De Jong, to whom we express our appreciation.

Finally, we also would like to thank the anonymous referee for his constructive comments.

References

[Abramovich et al. 2008] D. Abramovich, M. Olsson, and A. Vistoli, “Tame stacks in positive characteristic”, *Ann. Inst. Fourier (Grenoble)* **58**:4 (2008), 1057–1091. [MR](#) [Zbl](#)

[Bergh 2017] D. Bergh, “Functorial destackification of tame stacks with abelian stabilisers”, *Compos. Math.* **153**:6 (2017), 1257–1315. [MR](#) [Zbl](#)

[Bhatt and Scholze 2015] B. Bhatt and P. Scholze, “The pro-étale topology for schemes”, pp. 99–201 in *Astérisque* **369**, Soc. Math. France, Paris, 2015. [MR](#) [Zbl](#)

[Borne and Vistoli 2015] N. Borne and A. Vistoli, “The Nori fundamental gerbe of a fibered category”, *J. Algebraic Geom.* **24**:2 (2015), 311–353. [MR](#) [Zbl](#)

[Braverman and Bezrukavnikov 2007] A. Braverman and R. Bezrukavnikov, “Geometric Langlands correspondence for \mathcal{D} -modules in prime characteristic: the $GL(n)$ case”, *Pure Appl. Math. Q.* **3**:1 (2007), 153–179. [MR](#) [Zbl](#)

[Brochard 2014] S. Brochard, “Duality for commutative group stacks”, preprint, 2014. [arXiv](#)

[Colliot-Thélène and Sansuc 1976] J.-L. Colliot-Thélène and J.-J. Sansuc, “Torseurs sous des groupes de type multiplicatif; applications à l’étude des points rationnels de certaines variétés algébriques”, *C. R. Acad. Sci. Paris Sér. A-B* **282**:18 (1976), A1113–A1116. [MR](#) [Zbl](#)

- [Colliot-Thélène and Sansuc 1987] J.-L. Colliot-Thélène and J.-J. Sansuc, “La descente sur les variétés rationnelles, II”, *Duke Math. J.* **54**:2 (1987), 375–492. [MR](#) [Zbl](#)
- [Deligne 1990] P. Deligne, “Catégories tannakiennes”, pp. 111–195 in *The Grothendieck Festschrift, II*, edited by P. Cartier et al., Progr. Math. **87**, Birkhäuser, Boston, 1990. [MR](#) [Zbl](#)
- [Giraud 1971] J. Giraud, *Cohomologie non abélienne*, Grundlehren der Math. Wissenschaften **179**, Springer, 1971. [MR](#) [Zbl](#)
- [Laumon and Moret-Bailly 2000] G. Laumon and L. Moret-Bailly, *Champs algébriques*, Ergebnisse der Mathematik (3) **39**, Springer, 2000. [MR](#) [Zbl](#)
- [Nori 1982] M. V. Nori, “The fundamental group-scheme”, *Proc. Indian Acad. Sci. Math. Sci.* **91**:2 (1982), 73–122. [MR](#) [Zbl](#)
- [Otabe 2017] S. Otabe, “An extension of Nori fundamental group”, *Comm. Algebra* **45**:8 (2017), 3422–3448. [MR](#) [Zbl](#)
- [Saavedra Rivano 1972] N. Saavedra Rivano, *Catégories tannakiennes*, Lecture Notes in Math. **265**, Springer, 1972. [MR](#) [Zbl](#)
- [SGA 3₁ 1970] P. Gabriel, “Construction de préschémas quotient”, pp. 251–286 in *Schémas en groupes, Tome I: Propriétés générales des schémas en groupes, Exposés I–VII* (Séminaire de Géométrie Algébrique du Bois Marie 1962–1964), edited by M. Demazure and A. Grothendieck, Lecture Notes in Math. **151**, Springer, 1970. [MR](#)
- [SGA 4₃ 1973] M. Artin, A. Grothendieck, and J. L. Verdier, *Théorie des topos et cohomologie étale des schémas, Tome 3: Exposés IX–XIX* (Séminaire de Géométrie Algébrique du Bois Marie 1963–1964), Lecture Notes in Math. **305**, Springer, 1973. [MR](#) [Zbl](#)
- [Stacks 2005–] P. Belmans, A. J. de Jong, et al., “The Stacks project”, electronic reference, 2005–, available at <http://stacks.math.columbia.edu>.
- [Tonini and Zhang 2017] F. Tonini and L. Zhang, “Algebraic and Nori fundamental gerbes”, *J. Inst. Math. Jussieu* (online publication July 2017).
- [Tonini and Zhang 2018] F. Tonini and L. Zhang, “ F -divided sheaves trivialized by dominant maps are essentially finite”, *Trans. Amer. Math. Soc.* (online publication August 2018).

Communicated by Martin Olsson

Received 2017-09-18 Revised 2018-08-28 Accepted 2019-01-21

niels.borne@math.univ-lille1.fr

Laboratoire Paul Painlevé, Université de Lille, U.M.R. CNRS 8524,
U.F.R. de Mathématiques, Villeneuve-d’Ascq, France

angelo.vistoli@sns.it

Department of Mathematics, Scuola Normale Superiore, Pisa, Italy

Algebra & Number Theory

msp.org/ant

EDITORS

MANAGING EDITOR

Bjorn Poonen
Massachusetts Institute of Technology
Cambridge, USA

EDITORIAL BOARD CHAIR

David Eisenbud
University of California
Berkeley, USA

BOARD OF EDITORS

Richard E. Borcherds	University of California, Berkeley, USA	Raman Parimala	Emory University, USA
Antoine Chambert-Loir	Université Paris-Diderot, France	Jonathan Pila	University of Oxford, UK
J-L. Colliot-Thélène	CNRS, Université Paris-Sud, France	Anand Pillay	University of Notre Dame, USA
Brian D. Conrad	Stanford University, USA	Michael Rapoport	Universität Bonn, Germany
Samit Dasgupta	University of California, Santa Cruz, USA	Victor Reiner	University of Minnesota, USA
Hélène Esnault	Freie Universität Berlin, Germany	Peter Sarnak	Princeton University, USA
Gavril Farkas	Humboldt Universität zu Berlin, Germany	Joseph H. Silverman	Brown University, USA
Hubert Flenner	Ruhr-Universität, Germany	Michael Singer	North Carolina State University, USA
Sergey Fomin	University of Michigan, USA	Christopher Skinner	Princeton University, USA
Edward Frenkel	University of California, Berkeley, USA	Vasudevan Srinivas	Tata Inst. of Fund. Research, India
Andrew Granville	Université de Montréal, Canada	J. Toby Stafford	University of Michigan, USA
Joseph Gubeladze	San Francisco State University, USA	Pham Huu Tiep	University of Arizona, USA
Roger Heath-Brown	Oxford University, UK	Ravi Vakil	Stanford University, USA
Craig Huneke	University of Virginia, USA	Michel van den Bergh	Hasselt University, Belgium
Kiran S. Kedlaya	Univ. of California, San Diego, USA	Akshay Venkatesh	Institute for Advanced Study, USA
János Kollár	Princeton University, USA	Marie-France Vignéras	Université Paris VII, France
Philippe Michel	École Polytechnique Fédérale de Lausanne	Kei-Ichi Watanabe	Nihon University, Japan
Susan Montgomery	University of Southern California, USA	Melanie Matchett Wood	University of Wisconsin, Madison, USA
Shigefumi Mori	RIMS, Kyoto University, Japan	Shou-Wu Zhang	Princeton University, USA
Martin Olsson	University of California, Berkeley, USA		

PRODUCTION

production@msp.org

Silvio Levy, Scientific Editor


See inside back cover or msp.org/ant for submission instructions.

The subscription price for 2019 is US \$385/year for the electronic version, and \$590/year (+\$60, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Algebra & Number Theory (ISSN 1944-7833 electronic, 1937-0652 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

ANT peer review and production are managed by EditFLOW® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

Algebra & Number Theory

Volume 13 No. 3 2019

Fundamental gerbes	531
NIELS BORNE and ANGELO VISTOLI	
A new proof of the Waldspurger formula, I	577
RAHUL KRISHNA	
Algebraic independence for values of integral curves	643
TIAGO J. FONSECA	
Generically split octonion algebras and \mathbb{A}^1-homotopy theory	695
ARAVIND ASOK, MARC HOYOIS and MATTHIAS WENDT	