

The pillowcase and perturbations of traceless representations of knot groups

MATTHEW HEDDEN
CHRISTOPHER M HERALD
PAUL KIRK

We introduce explicit holonomy perturbations of the Chern–Simons functional on a 3–ball containing a pair of unknotted arcs. These perturbations give us a concrete local method for making the moduli spaces of flat singular $SO(3)$ connections relevant to Kronheimer and Mrowka’s singular instanton knot homology nondegenerate. The mechanism for this study is a (Lagrangian) intersection diagram which arises, through restriction of representations, from a tangle decomposition of a knot. When one of the tangles is trivial, our perturbations allow us to study isolated intersections of two Lagrangians to produce minimal generating sets for singular instanton knot homology. The (symplectic) manifold where this intersection occurs corresponds to the traceless character variety of the four-punctured 2–sphere, which we identify with the familiar pillowcase. We investigate the image in this pillowcase of the traceless representations of tangles obtained by removing a trivial tangle from 2–bridge knots and torus knots. Using this, we compute the singular instanton homology of a variety of torus knots. In many cases, our computations allow us to understand nontrivial differentials in the spectral sequence from Khovanov homology to singular instanton homology.

[57M27](#), [57R58](#), [57M25](#); [81T13](#)

1 Introduction

Kronheimer and Mrowka have recently developed a variant of instanton homology for knots in 3–manifolds [[26](#); [25](#); [21](#)] which they call *singular instanton knot homology*. In [[25](#)], they construct a filtered chain complex whose total homology is the singular instanton homology of a knot $K \subset S^3$, and whose spectral sequence has E_2 page the Khovanov homology of K . Inspired by early observations of similarities between their theory and Khovanov homology, Lewallen [[28](#)] showed that for 2–bridge knots the Khovanov homology is isomorphic to the homology of the variety of $SU(2)$ representations of the fundamental group of the knot complement sending the meridian to

a traceless matrix. For alternating knots, he showed that the Khovanov homology is isomorphic to the homology of the subvariety of binary dihedral representations.

In fact for alternating knots the Khovanov and singular instanton homology groups are isomorphic (with the Khovanov bigrading appropriately collapsed to a $\mathbb{Z}/4$ grading), a fact implied by the collapse of Kronheimer and Mrowka's spectral sequence at the E_2 page. In contrast, they show that there are nontrivial higher differentials in the spectral sequence associated to the $(4, 5)$ torus knot [21, Section 11], and hence Khovanov and instanton homology do not have the same rank, in general. (Rasmussen noticed that the Khovanov homology of the $(4, 5)$ torus knot also has larger rank than its Heegaard knot Floer homology groups. It is conjectured that there is a similar spectral sequence in that context.) Zentner [41] showed that for some alternating pretzel knots there are nonbinary dihedral traceless representations (in contrast to 2-bridge knots), so that for these families one expects there to be nontrivial differentials in the singular instanton chain complex.

To make sense of this expectation, we should recall that the chain complex from which the instanton homology is computed is the Morse complex associated to a perturbation of a particular Chern–Simons functional. This chain complex is $\mathbb{Z}/4$ graded, and is generated by certain gauge equivalence classes of perturbed-flat connections. These connections live on an $SO(3)$ bundle on the complement of the link formed by adding a small meridional circle H to K . The second Stiefel–Whitney class of the bundle is Poincaré dual to an arc W connecting K and H , as in Figure 1. The unperturbed flat moduli space can be identified with the conjugacy classes of $SU(2)$ representations of the fundamental group of $S^3 \setminus (K \cup H \cup W)$, which take the meridians of K and H to traceless matrices in $SU(2)$, and the meridian of W to the nontrivial central element -1 . A feature of these representation spaces is that, with the exception of the unknot in S^3 , they are never nondegenerate; the corresponding (unperturbed) Chern–Simons functional is not Morse. Thus it is necessary to perturb the functional to identify the generators of the instanton complex. A common method for perturbing the Chern–Simons functional is through the use of so-called *holonomy perturbations*, and such perturbations can be constructed quite generally in instanton Floer theories. The use of such perturbations, however, obscures the connection between generators of the instanton chain complex and representations of the fundamental group. In particular, it would be desirable to be able to effectively estimate the rank of a reduced instanton chain complex from a presentation of the fundamental group of the knot or link complement. A general perturbation of the Chern–Simons functional will make this estimation impossible.

The purpose of this article is to identify a certain perturbation explicitly and to use it to identify generators of the reduced instanton chain complex. The main conceptual step




Figure 1: A small segment of a knot K and the meridional circle H and arc W used in the construction of the reduced singular instanton homology.

is to split a 3-manifold containing a knot along a *Conway sphere*; that is, a 2-sphere which intersects the knot in four points. The sphere decomposes the knot into two tangles, and determining generators of the instanton complex becomes an intersection problem for two Lagrangians in the relevant character variety of the 4-punctured sphere. The key to our result lies in picking a Conway sphere for which one of the associated tangles is trivial, and constructing concrete local (and hence universal) perturbations in this trivial piece.

It turns out ([Proposition 3.1](#)) that the relevant character variety for a 2-sphere with four marked points, $R(S^2, \{a, b, c, d\})$, is a *pillowcase*, ie a topological 2-sphere with four singular points, understood as the quotient of the torus by the hyperelliptic involution or, equivalently, as the quotient of \mathbb{R}^2 :

$$R(S^2, \{a, b, c, d\}) \cong \mathbb{R}^2 / \sim, \quad (\gamma, \theta) \sim (-\gamma, -\theta) \sim (\gamma + 2\pi m, \theta + 2\pi n).$$

We describe paths in $R(S^2, \{a, b, c, d\})$ by giving formulas for their lifts to \mathbb{R}^2 , eg $t \mapsto (\gamma(t), \theta(t))$. In terms of the pillowcase, our main results, [Theorem 7.1](#) and [Corollary 7.2](#), state the following.

Theorem 1 *Let $A_1 \cup A_2 \subset B^3$ be a pair of unknotted arcs in the 3-ball, H a small meridian of A_1 , and W an arc connecting H and A_1 . Let $P \subset B^3 \setminus (A_1 \cup A_2 \cup H \cup W)$ be the perturbation curve illustrated in [Figure 9](#).*

Given small perturbation data $\pi = (P, \epsilon, f)$ (where $\epsilon > 0$ and f is an odd, 2π -periodic function, eg $f(\beta) = \sin(\beta)$), the perturbed reduced moduli space $R_\pi^h(B^2, A_1 \cup A_2)$ is

homeomorphic to a circle, and the image of the restriction map to the pillowcase

$$\rho: R_{\pi}^{\sharp}(B^2, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$$

is an immersion, given by

$$\rho(\beta) = (\gamma(\beta), \theta(\beta)) = (\beta + \epsilon f(\beta) + \frac{\pi}{2}, \beta - \epsilon f(\beta) + \frac{\pi}{2}), \beta \in \mathbb{R}/(2\pi\mathbb{Z})$$

as illustrated in Figure 10. As $\epsilon \rightarrow 0$, this immersion limits to a generically two-to-one map onto the diagonal arc $\gamma = \theta$ in the pillowcase.

A more precise statement can be found in the body of the paper. For all the results in this article it suffices to take $f(\beta) = \sin(\beta)$ in the perturbation data.

In Section 9 we treat the case corresponding to the unreduced instanton homology $I^{\sharp}(Y, K)$. The counterpart of Theorem 7.1 in this context is Theorem 9.1, which states in part, the following.

Theorem 2 *With perturbation data $\pi = (P_1 \cup P_2, \epsilon, \sin(\beta))$ supported on the two curves P_1 and P_2 in Figure 11, the perturbed unreduced moduli space $R_{\pi}^{\sharp}(B^3, A_1 \cup A_2)$ is a disjoint union of two circles, parameterized by $\beta \in \mathbb{R}/(2\pi\mathbb{Z})$ and $i = 1, 2$. The restriction map of $R_{\pi}^{\sharp}(B^3, A_1 \cup A_2)$ to the pillowcase is an immersion, given by*

$$\rho_i(\beta) = (-\tau_i(\beta) + \frac{\pi}{2} + \beta + \epsilon \sin \beta, -\tau_i(\beta) + \frac{\pi}{2} + \beta - \epsilon \sin \beta), \quad i = 1, 2,$$


where $\tau_1(\beta) = \arcsin(-\frac{1}{2} \sin \beta)$ and $\tau_2(\beta) = \pi - \tau_1(\beta)$. As $\epsilon \rightarrow 0$, this immersion limits to a generically four-to-one map onto the diagonal arc $\gamma = \theta$ in the pillowcase.

Figure 12 illustrates the images in the pillowcase of the three curves, ρ_1, ρ_2 of Theorem 2, and ρ of Theorem 1.

As alluded to above, a useful implication of these results is that it reduces the problem of identifying generators of the singular instanton complexes for a knot K in a 3-manifold Y to an intersection problem in the pillowcase. More precisely, consider a 3-ball which intersects (Y, K) in a pair of unknotted arcs $A_1 \cup A_2$. Setting $(Y_0, K_0) := (Y \setminus B^3, K \setminus (A_1 \cup A_2))$, we obtain a tangle decomposition

$$(1) \quad (Y, K) = (Y_0, K_0) \cup_{(S^2, \{a, b, c, d\})} (B^3, A_1 \cup A_2).$$

This decomposition yields, upon passing to the appropriate moduli spaces, an intersection diagram:



The intersection $R_{\pi}^{\natural}(Y, K)$ parameterizes the generators of the chain complex defining the instanton homology $I^{\natural}(Y, K)$, provided this is a nondegenerate set. The unperturbed space $R^{\natural}(Y, K)$ is never nondegenerate, except for the unknot in S^3 , in contrast to other examples of this method in low-dimensional topology such as Casson’s invariant; see Akbulut and McCarthy [1].

In the decomposition (1), the nontrivial part of the $SO(3)$ bundle and the perturbation π needed to make the set $R(Y, K)$ nondegenerate has been placed entirely inside the simple space B^3 . Theorem 7.1 states that $R_{\pi}^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ is a smooth immersion of a circle, and identifies the image precisely.

It follows that the problem of describing the set $R^{\natural}(Y, K)$ of generators of the instanton homology chain complex is reduced to understanding the space $R(Y_0, K_0)$ and its restriction to the pillowcase, a problem that involves only the fundamental group of the 2–stranded tangle complement $Y_0 \setminus K_0$ and its peripheral structure. Indeed, for simple knots like 2–bridge knots and torus knots, no further perturbations are needed. For general knots only perturbations in the knot complement (which have been studied in detail for a long time, see eg Herald [12]) are required.

With this understanding in place, we turn our attention to the problem of describing $R(Y_0, K_0)$ and its image in the pillowcase $R(S^2, \{a, b, c, d\})$ for 2–bridge knots and torus knots. More precisely, given a knot K in S^3 and a 3–ball B^3 meeting K in a pair of unknotted arcs $A_1 \cup A_2$, the complement Y_0 of this 3–ball is again a 3–ball and contains the two-component tangle $K_0 = K \setminus (A_1 \cup A_2)$. The moduli space $R(Y_0, K_0)$ is identified with the space of conjugacy classes of $SU(2)$ representations of $\pi_1(Y_0 \setminus K_0)$ which send meridians of K_0 to traceless matrices. This space, and its restriction to the pillowcase, turns out to be a very interesting tangle invariant. In Section 10 we identify $R(Y_0, K_0)$ and its image in the pillowcase for rational tangles (the tangles which glue with the trivial tangle to produce 2–bridge knots) and prove the following theorem.

Theorem 3 For the 2–bridge knot $K = K(p/q)$ and the 3–ball $B^3 \subset S^3$ meeting K in a pair of unknotted arcs as in [Figure 14](#), the space $R(Y_0, K_0)$ is an arc, and the restriction $R(Y_0, K_0) \rightarrow R(S^2, \{a, b, c, d\})$ is the embedding

$$(\gamma, \theta) = (qt, (q - p)t), \quad t \in [0, \pi].$$

Using [Theorem 7.1](#) and [Corollary 7.2](#) it follows that the set $R_\pi^{\natural}(S^3, K)$ is a union of pairs of isolated nondegenerate points x_{ℓ_1}, x_{ℓ_2} , $\ell = 1, 2, \dots, (p - 1)/2$ and one additional point α' . In particular, we have that the chain complex $CI^{\natural}(S^3, K)$ for the reduced instanton homology of $K(p/q)$ is generated by these $2((p - 1)/2) + 1 = p$ points. From [\[25\]](#) we know that all differentials are zero so that these points generate the instanton homology. While the instanton homology of 2–bridge knots is easily determined by the spectral sequence of [\[25\]](#), we find it interesting to be able to produce an explicit complex with trivial differential (in contrast to the complex coming from the spectral sequence which has many more generators than the rank of homology, or the highly degenerate unperturbed character variety studied in [\[28\]](#)).

In [Section 11](#) we analyze the corresponding representation spaces for tangles arising from torus knots. These are more complicated than the spaces associated to 2–bridge knots. For a particular tangle decomposition $(Y_0, K_0) \cup (B^3, A_1 \cup A_2)$ of the (p, q) torus knot, the space $R(Y_0, K_0)$ can be identified with a certain singular semialgebraic curve in \mathbb{R}^2 , cut out by a 2–variable polynomial determined entirely by the integers p, q and r, s , where $pr + qs = 1$. We obtain several results ([Theorems 11.1](#) and [11.3](#)) which give different descriptions of $R(Y_0, K_0)$ for torus knots. See [Figure 19](#) for the explicit example of the $(3, 4)$ torus knot, a knot whose instanton chain complex (for all small enough perturbations) necessarily has a nontrivial differential. Our results show that for torus knots a reduced singular instanton chain complex can be found with $|\sigma(K)| + 1$ generators, where $\sigma(K)$ denotes the signature of K . Moreover, for small generic holonomy perturbations this is the fewest possible generators, even if the instanton homology has smaller rank.

The pillowcase arises in another context when studying the $SU(2)$ representation spaces of knot complements: as the character variety $\chi(T)$ of all $SU(2)$ representations of the fundamental group of a torus. We will have occasion to use both incarnations, $R(S^2, \{a, b, c, d\})$ and $\chi(T)$, of the pillowcase in this article. Indeed, our analysis of $R^{\natural}(Y, K)$ calls on the examination of the restriction from $R(Y_0, K_0)$ to the pillowcase $R(S^2, \{a, b, c, d\})$, as well as the restriction of the full character variety $\chi(Y, K)$ (the space of conjugacy classes of all representations of $\pi_1(Y \setminus K)$) to the pillowcase $\chi(T)$ associated to the peripheral torus.

This latter context is familiar and can be found in many places in the literature, starting with Klassen's article [20], and, in the enlarged context of $SL(2, \mathbb{C})$ representation, as the variety defined by the A -polynomial of a knot; see Cooper, Culler, Gillet, Long and Shalen [7]. The interplay between these two manifestations of the pillowcase is exploited in Section 12. We combine our results with the foundational theorems of Kronheimer and Mrowka to calculate and tabulate the reduced instanton complex, the Khovanov homology and the instanton homology for various families of torus knots. In particular we give new examples of torus knots for which there are nontrivial differentials in the instanton chain complex, examples for which there are many higher nontrivial differentials in the spectral sequence, and non-2-bridge examples for which all differentials in the chain complex are zero. For example, the spectral sequence for the $(5, 7)$ torus knot drops in rank from 29 to 17 after the E_2 page, and the spectral sequence for the $(13, 28)$ torus knot collapses at Khovanov homology.

It is worth contrasting our approach with the one taken in Jacobsson and Rubinsztein [15], where a knot in S^3 is described by a closed braid, and a Lagrangian intersection picture is obtained by cutting along a $2n$ -punctured 2-sphere which separates the braid from a trivial braid used to close it. The purpose of that article, however, is to explore the symplectogeometric aspects of their setup, whereas our emphasis is on singular instanton homology and its efficient computation.

Readers interested in a quick geometric overview of the contents of this article are encouraged to examine Figures 9 and 10, which encapsulate the statement of Theorem 1. Figures 11 and 12 illustrate Theorem 2. The reader is also encouraged to compare Figures 4 and 15 corresponding to the trefoil knot, Figures 20 and 19 corresponding to the $(3, 4)$ torus knot and Figures 6 and 17 corresponding to the knot 7_2 . These figures illustrate how to determine generators of the singular instanton chain complexes of these knots from the intersection of arcs of traceless and perturbed traceless representations in the pillowcase.

Acknowledgements The first author gratefully acknowledges support from NSF grant DMS-0906258, NSF CAREER grant DMS-1150872, and an Alfred P. Sloan Research Fellowship. The third author gratefully acknowledges support from NSF grant DMS-1007196.

2 Unit quaternions

In this section we establish notation and recall some basic facts about $SU(2)$. Identify $SU(2)$ with the unit quaternions,

$$SU(2) = \{a + bi + cj + dk \mid a, b, c, d \in \mathbb{R}, a^2 + b^2 + c^2 + d^2 = 1\}.$$

The inverse of a unit quaternion $q = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$ is equal to its conjugate $\bar{q} = a - b\mathbf{i} - c\mathbf{j} - d\mathbf{k}$. The Lie algebra of $SU(2)$ is identified with the *purely imaginary quaternions*

$$su(2) = \{b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \mid b, c, d \in \mathbb{R}\}$$

with invariant inner product $v \cdot w = -\text{Re}(vw)$. We denote the exponential map $su(2) \rightarrow SU(2)$ by $q \mapsto e^q$. Let $C(\mathbf{i}) \subset SU(2)$ denote the conjugacy class of \mathbf{i} : this is the 2–sphere of *purely imaginary unit quaternions*

$$C(\mathbf{i}) = \{b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \mid b, c, d \in \mathbb{R}, b^2 + c^2 + d^2 = 1\}.$$

We also call $C(\mathbf{i})$ the *traceless unit quaternions*, since they correspond to the traceless 2×2 matrices in the usual description of $SU(2)$.

Notice that $C(\mathbf{i})$ lies in the Lie algebra $su(2)$, and that as a subset of $SU(2)$ it is the set of unit quaternions $q \in SU(2)$ satisfying $\text{Re}(q) = 0$. Furthermore,

$$e^{vQ} = \cos(v) + \sin(v)Q \text{ for } v \in \mathbb{R}, \quad Q^2 = -1 \text{ for } Q \in C(\mathbf{i}).$$

In particular, every nonzero vector in $su(2)$ can be uniquely written in the form tQ for some $t \in \mathbb{R}_{>0}$, $Q \in C(\mathbf{i})$. Every element of $SU(2)$ can be written in the form e^{tQ} for some $t \in \mathbb{R}$, $Q \in C(\mathbf{i})$; this representation is unique for $SU(2) \setminus \{\pm 1\}$ if we choose $0 < t < \pi$.

We summarize a few well known and easily verified properties of the conjugation action of $SU(2)$ on itself in the following proposition.

Proposition 2.1 *Consider the action of $SU(2)$ on itself by conjugation.*

- (i) *The stabilizer of any subgroup $G \subset SU(2)$,*

$$\text{Stab}(G) = \{a \in SU(2) \mid ag\bar{a} = g \text{ for all } g \in G\},$$

can be one of the three types: $\{\pm 1\}$, $\{e^{tQ}\}$ for some $Q \in C(\mathbf{i})$, or $SU(2)$ according to whether G is nonabelian, G is abelian but noncentral, or G is contained in the center $\{\pm 1\}$.

- (ii) *Given a traceless unit quaternion $Q \in C(\mathbf{i})$ and $t \in \mathbb{R}$, the conjugation action of e^{tQ} on the 2–sphere $C(\mathbf{i})$ of traceless unit quaternions is rotation about the axis through Q of angle $2t$.*
- (iii) *If q is a unit quaternion and $Q \in C(\mathbf{i})$ is a traceless unit quaternion which together satisfy $-Q = qQ\bar{q}$, then q is itself a traceless unit quaternion, and q and Q are perpendicular, ie, $\text{Re}(qQ) = 0$.*

(iv) If $Q_1, Q_2 \in C(\mathbf{i})$ and $t_1, t_2 \in \mathbb{R}$, then

$$\operatorname{Re}(e^{t_1 Q_1} e^{t_2 Q_2}) = \cos t_1 \cos t_2 - \sin t_1 \sin t_2 \cos \nu,$$

where ν denotes angle (in $su(2) = \mathbb{R}^3$) between Q_1 and Q_2 .

(v) If e^{tQ_1} and e^{sQ_2} commute, with $Q_i \in C(\mathbf{i})$ and $e^{tQ_1}, e^{sQ_2} \neq \pm 1$, then $Q_1 = \pm Q_2$. □

3 The pillowcase

In this section we introduce the pillowcase as a quotient space of \mathbb{R}^2 , and describe how it arises as an $SU(2)$ character variety in two different ways.

3.1 The pillowcase as quotient of \mathbb{R}^2

Let G denote the split extension of \mathbb{Z}^2 by $\mathbb{Z}/2$ where the generator $\tau \in \mathbb{Z}/2$ acts on $(m, n) \in \mathbb{Z}^2$ by $\tau \cdot (m, n) = -(m, n)$. Then G acts affinely on the plane \mathbb{R}^2 by

$$(m, n) \cdot (x, y) = (x + 2\pi m, y + 2\pi n), \quad \tau \cdot (x, y) = (-x, -y).$$

The quotient \mathbb{R}^2/G is called *the pillowcase*. The quotient map

$$(3) \quad \mathbb{R}^2 \rightarrow \mathbb{R}^2/G$$

is a branched cover, branched over four points with preimage the lattice $(\mathbb{Z}\pi)^2 \subset \mathbb{R}^2$ of points with nontrivial isotropy. The pillowcase is homeomorphic to a 2-sphere.

A fundamental domain for the action is the rectangle $[0, \pi] \times [0, 2\pi]$ (see [Figure 2](#)), and the identifications along its boundary are

$$(4) \quad (x, 0) \sim (x, 2\pi), \quad (0, y) \sim (0, 2\pi - y), \quad (\pi, y) \sim (\pi, 2\pi - y).$$

Hence the moniker “pillowcase.” Taking the quotient in two steps, first by \mathbb{Z}^2 and then by $\mathbb{Z}/2$, exhibits the pillowcase as the quotient of the torus by the hyperelliptic involution.

We will often describe a curve in the pillowcase in terms of a lift to \mathbb{R}^2 . A curve may have different lifts, for example the straight line segments $t \mapsto (2t, 3t)$ and $t \mapsto (-2t, -3t + 4\pi)$, $t \in [0, \frac{\pi}{2}]$ define the same curve in the pillowcase.




Figure 2: The pillowcase: on the left is a fundamental domain for the action of $\mathbb{Z}^2 \rtimes \mathbb{Z}/2$ on \mathbb{R}^2 ; on the right is the pillowcase, obtained by performing the identifications on the left. Topologically, the pillowcase is a 2–sphere.

3.2 The pillowcase as the $SU(2)$ character variety of the torus

The $SU(2)$ character variety of the 2–dimensional torus T is the space of conjugacy classes of representations:

$$\chi(T) = \{\rho: \pi_1(T) \rightarrow SU(2)\} / \text{conjugation}.$$

If $\mu, \lambda \in \pi_1(T)$ denote generators, then to any pair $(x, y) \in \mathbb{R}^2$ of real numbers one can assign the conjugacy class of the representation

$$\mu \mapsto e^{xi}, \quad \lambda \mapsto e^{yi},$$

in $\chi(T)$. The resulting map $\mathbb{R}^2 \rightarrow \chi(T)$ factors through the branched cover of Equation (3) and induces a homeomorphism of the pillowcase with $\chi(T)$. Note that the identification depends on the choice of generators μ, λ .

The representations which send μ to a traceless matrix, ie to $C(i)$, correspond exactly to the line $x = \frac{\pi}{2}$, since $e^{\pi i/2} = i$. The line $\{x = \frac{\pi}{2}\} \subset \mathbb{R}^2$ is sent to the circle in the pillowcase:

$$(5) \quad S(i) = \{\rho: \pi_1(T) \rightarrow SU(2) \mid \rho(\mu) = i, \rho(\lambda) = e^{yi}\}$$

(For more details, see eg Kirk and Klassen [18].)

3.3 The pillowcase as the traceless $SU(2)$ character variety of a four-punctured 2-sphere

Consider a 2-sphere with four marked points, labelled a, b, c , and d , respectively, so that the fundamental group $\pi_1(S^2 \setminus \{a, b, c, d\})$ is presented (by abuse of notation) as

$$\pi_1(S^2 \setminus \{a, b, c, d\}) = \langle a, b, c, d \mid ba = cd \rangle$$

as indicated in Figure 3.

We denote by $R(S^2, \{a, b, c, d\})$ the space of conjugacy classes of homomorphisms which take the loops at each puncture to a traceless quaternion:

$$R(S^2, \{a, b, c, d\}) = \{ \rho: \langle a, b, c, d \mid ba = cd \rangle \rightarrow SU(2) \mid \rho(a), \rho(b), \rho(c), \rho(d) \in C(\mathbf{i}) \} / \text{conjugation}.$$

The space $R(S^2, \{a, b, c, d\})$ is known to be a topological 2-sphere with four singular points; see eg Lin [30, Lemma 2.1] and Heusener and Kroll [14, Lemma 4.1]. The following proposition gives an explicit identification with the pillowcase, considered as a quotient of \mathbb{R}^2 .

Proposition 3.1 *There is a surjective quotient map*

$$\psi: \mathbb{R}^2 \rightarrow R(S^2, \{a, b, c, d\})$$

given by

$$\psi(\gamma, \theta): \quad a \mapsto \mathbf{i}, \quad b \mapsto e^{\gamma \mathbf{k}} \mathbf{i}, \quad c \mapsto e^{\theta \mathbf{k}} \mathbf{i}, \quad d \mapsto e^{(\theta - \gamma) \mathbf{k}} \mathbf{i}.$$

The map ψ factors through the branched cover of Equation (3) and induces a homeomorphism of the pillowcase with $R(S^2, \{a, b, c, d\})$. The four corner points correspond to reducible noncentral representations, and all other points correspond to irreducible representations.

Proof Given any $(\gamma, \theta) \in \mathbb{R}^2$, the assignment $a \mapsto \mathbf{i}$, $b \mapsto e^{\gamma \mathbf{k}} \mathbf{i}$, $c \mapsto e^{\theta \mathbf{k}} \mathbf{i}$, $d \mapsto e^{(\theta - \gamma) \mathbf{k}} \mathbf{i}$ satisfies $ba = cd$, and hence ψ maps into $R(S^2, \{a, b, c, d\})$.

Any traceless representation $\rho: \pi_1(S^2 \setminus \{a, b, c, d\}) \rightarrow SU(2)$ can be conjugated so that $\rho(a) = \mathbf{i}$. Since $e^{-t/2 \mathbf{i}} (e^{t \mathbf{i}} \mathbf{j}) e^{t/2 \mathbf{i}} = \mathbf{j}$, ρ can be further conjugated (fixing $\rho(a)$) so that $\rho(b) = \cos(\gamma) \mathbf{i} + \sin(\gamma) \mathbf{j} = e^{\gamma \mathbf{k}} \mathbf{i}$ for some $\gamma \in [0, \pi]$.

Suppose that $\rho(c) = c_1 \mathbf{i} + c_2 \mathbf{j} + c_3 \mathbf{k}$. Since $q \in SU(2)$ lies in $C(\mathbf{i})$ if and only if $\text{Re}(q) = 0$, the equation

$$0 = \text{Re}(\rho(d)) = \text{Re}(\rho(c^{-1}ba)) = \text{Re}(\rho(c^{-1})(e^{\gamma \mathbf{k}} \mathbf{i}) \mathbf{i}) = -c_3 \sin \gamma$$




Figure 3: A four-punctured sphere with boundary curves added that generate its fundamental group (after choosing a base point near the barycenter of the curves and arcs to the boundary).

implies that either $\sin \gamma = 0$ or else $c_3 = 0$. If $\sin \gamma = 0$, then $\rho(a) = \mathbf{i}$ and $\rho(b) = \pm \mathbf{i}$, and hence ρ may be further conjugated by $e^{-t/2\mathbf{i}}$, fixing $\rho(a)$ and $\rho(b)$, so that $c_3 = 0$. So we may assume by conjugating that $c_3 = 0$ in either case. Hence $\rho(c) = c_1 \mathbf{i} + c_2 \mathbf{j} = e^{\theta \mathbf{k}} \mathbf{i}$ for some $\theta \in [0, 2\pi]$.

Therefore the map $\psi: \mathbb{R}^2 \rightarrow R(S^2, \{a, b, c, d\})$ is onto. It is easy to check that $\psi(\gamma_1, \theta_1) = \psi(\gamma_2, \theta_2)$ if and only if $(\gamma_2, \theta_2) = \pm(\gamma_1, \theta_1) + (2\pi m, 2\pi n)$, so that ψ passes to a homeomorphism $\mathbb{R}^2/G \rightarrow R(S^2, \{a, b, c, d\})$. \square

4 Representation spaces of knots

To a knot K in a 3-manifold Y with tubular neighborhood $N(K)$ we will assign several spaces of conjugacy classes of $SU(2)$ representations. The notation “ χ ” will be used when we consider all conjugacy classes of representations of $\pi_1(Y \setminus N(K))$, and “ R ” will be used when we restrict to those representations which send the meridians of K into the conjugacy class of traceless unit quaternions $C(\mathbf{i})$.

4.1 All representations

First, let $\chi(Y, K)$ denote the space of conjugacy classes of all representations of $\pi_1(Y \setminus N(K))$ into $SU(2)$:

$$(6) \quad \chi(Y, K) = \{\rho: \pi_1(Y \setminus N(K)) \rightarrow SU(2)\} / \text{conjugation}.$$

Restricting to the boundary of $N(K)$ yields a map

$$\chi(Y, K) \rightarrow \chi(\partial N(K)).$$

Choosing generators μ_K, λ_K for $\pi_1(\partial N(K))$ gives an identification of $\chi(\partial N(K))$ with the pillowcase. In particular, if Y is a homology sphere we take the generators μ_K to be the canonical (isotopy class of) meridian and λ_K the longitude arising as the boundary of an oriented Seifert surface.

If Z is any 3-manifold containing a torus $T = S^1 \times S^1 \subset Z$, restricting representations yields a map from the character variety $\chi(Z)$ of (all) $SU(2)$ representations of $\pi_1(Z)$ to the pillowcase:

$$(7) \quad \chi(Z) \rightarrow \chi(T).$$

4.2 Traceless representations

Next, denote by $R(Y, K) \subset \chi(Y, K)$ the subset of conjugacy classes of representations sending every meridian of K into $C(\mathbf{i})$. Note that all meridians of K are conjugate since we are assuming that K is a knot rather than a link, and so any representation sending a particular meridian $\mu_K \in \pi_1(Y \setminus N(K))$ into $C(\mathbf{i})$ takes all meridians into $C(\mathbf{i})$. Hence

$$(8) \quad R(Y, K) = \{\rho: \pi_1(Y \setminus N(K)) \rightarrow SU(2) \mid \rho(\mu_K) \in C(\mathbf{i})\} / \text{conjugation}.$$

Given a pair μ_K, λ_K of generators of $\pi_1(\partial N(K))$ with μ_K a meridian, then $R(Y, K)$ can be described as the preimage under the restriction $\chi(Y, K) \rightarrow \chi(\partial N(K))$ of the circle $S(\mathbf{i})$ of Equation (5).

If $S^2 \subset Y$ is an embedded 2-sphere intersecting K transversely in four points a, b, c, d , we can restrict representations to S^2 to obtain a map

$$(9) \quad R(Y, K) \rightarrow R(S^2, \{a, b, c, d\}),$$

which we consider as the traceless analogue of (7).

4.3 Adding an earring to avoid reducibles

The spaces $\chi(Y, K)$ and $R(Y, K)$ are stratified according to the three possible stabilizers of the constituent $SU(2)$ representations: $\{\pm 1\}$, S^1 or $SU(2)$. To avoid singularities in moduli spaces arising from reducible connections, Kronheimer and Mrowka [25] introduce an auxiliary construction which ensures that only the center $\{\pm 1\}$ appears as a stabilizer (actually their construction requires the use of connections in a nontrivial $SO(3)$ bundle, with w_2 dual to the arc W described below). Let us recall their construction.

Fix a base point on K and identify $N(K)$ with the unit normal (disk) bundle of K . Let H be the circle of radius $\frac{1}{2}$ in the normal disk fiber containing the base point.

Denote the link $K \cup H$ by K^{\natural} . Let $W \subset Y$ be a radial arc in this normal disk connecting K and H ; see [Figure 1](#).

The boundary of a small tubular neighborhood of H is a torus which is punctured once by the arc W . Thus for the appropriate choices of basings and orientation of loops, the meridian μ_H and longitude λ_H of this torus satisfy $[\lambda_H, \mu_H] = \mu_W$. Moreover, since H is a small circle linking K , the longitude of H equals the meridian of K , ie $\lambda_H = \mu_K$. Thus $[\mu_K, \mu_H] = \mu_W$ in $\pi_1(Y \setminus (K^{\natural} \cup W))$.

We will use $R^{\natural}(Y, K)$ to denote the space of conjugacy classes of representations $\rho: \pi_1(Y \setminus (K^{\natural} \cup W)) \rightarrow \text{SU}(2)$ which send the meridians μ_K, μ_H of K and H to $C(\mathbf{i})$ and the meridian μ_W of the arc W to -1 :

$$(10) \quad R^{\natural}(Y, K) = \{ \rho: \pi_1(Y \setminus (K^{\natural} \cup W)) \rightarrow \text{SU}(2) \mid \rho(\mu_K), \rho(\mu_H) \in C(\mathbf{i}), \rho(\mu_W) = -1 \} / \text{conjugation}$$

When $Y = S^3$ (or any homology sphere), there is a distinguished conjugacy class $\alpha \in R^{\natural}(Y, K)$ which is characterized completely by the requirement that its restriction to the complement of a (large) tubular neighborhood $N(K)$ of K containing $H \cup W$ is abelian. The representation α can be uniquely conjugated to satisfy

$$(11) \quad \alpha(\mu_K) = \mathbf{i}, \quad \alpha(\mu_H) = \mathbf{j}, \quad \alpha(\mu_W) = -1, \quad \alpha(\lambda_K) = 1.$$

Since the set of all conjugates of μ_K by loops in $Y \setminus N(K)$ generate $\pi_1(Y \setminus N(K))$, α sends each of these conjugates to \mathbf{i} . The representation α and its restriction to $\pi_1(Y \setminus N(K))$, which will be denoted $\alpha_{\pi/2}$, appear frequently below.

If $Z \subset Y$ is a codimension-zero submanifold which contains $K^{\natural} \cup W$, then there is a restriction map $R^{\natural}(Y, K) \rightarrow R^{\natural}(Z, K)$. For example, one can take $Z = N(K)$, a tubular neighborhood of K large enough to contain $H \cup W$ to get $R^{\natural}(N(K), K)$.

Proposition 4.1 *The space $R^{\natural}(N(K), K)$ is homeomorphic to a circle, and every representation in $R^{\natural}(N(K), K)$ is nonabelian. Moreover, the restriction map*

$$R^{\natural}(N(K), K) \rightarrow \chi(\partial N(K))$$

is injective, with image the vertical circle $S(\mathbf{i})$ of [Equation \(5\)](#).

Proof Any representation in $R^{\natural}(N(K), K)$ may be conjugated so that $\rho(\mu_K) = \mathbf{i}$. Since $\rho(\lambda_K)$ commutes with $\rho(\mu_K)$ it follows that $\rho(\lambda_K) = e^{\ell \mathbf{i}}$ for some ℓ .

The relation $[\rho(\mu_K), \rho(\mu_H)] = \rho(\mu_W) = -1$ implies ρ is nonabelian and that $\rho(\mu_H)$ is perpendicular to \mathbf{i} ([Proposition 2.1](#)). Thus $\rho(\mu_H) = \cos \theta \mathbf{j} + \sin \theta \mathbf{k} = e^{\theta \mathbf{i}} \mathbf{j}$ for

some θ . Further conjugation by $e^{-\theta \mathbf{i}/2}$ fixes $\rho(\mu_K) = \mathbf{i}$ and $\rho(\lambda_K) = e^{\ell \mathbf{i}}$ but rotates so that $\rho(\mu_H) = \mathbf{j}$.

Conversely, any choice of $e^{\ell \mathbf{i}} \in S^1$ defines a unique conjugacy class in $R^{\natural}(N(K), K)$ by sending μ_K to \mathbf{i} , λ_K to $e^{\ell \mathbf{i}}$, μ_H to \mathbf{j} , and μ_W to -1 . \square

Note that we can think of K^{\natural} as obtained from K by forming the connected sum of K with one component of a Hopf link whose components are spanned by the arc W . Kronheimer and Mrowka also introduce a variant of this construction, which they denote by K^{\sharp} , obtained by taking the disjoint union of K with a Hopf link rather than the connected sum. This leads to a different representation space $R^{\sharp}(Y, K)$; see Section 9 below.

4.4 Gluing representations and the relation between the spaces $R^{\natural}(Y, K)$ and $R(Y, K)$

The spaces $R(Y, K)$ are better known than $R^{\natural}(Y, K)$. The relationship between the two is that $R^{\natural}(Y, K)$ maps to $R(Y, K)$ with fibers either a circle or a point depending on the reducibility of the image. We make this precise in Proposition 4.3 below.

We first present a folklore description of the character variety of a free product with amalgamation in terms of the character varieties of its pieces. To describe it, fix a compact Lie group G and (for this section only) let $\mathcal{H}(-)$ denote the functor which takes a discrete group H to the space $\text{Hom}(H, G)$ and let $\chi(-)$ denote the functor that takes H to $\text{Hom}(H, G)/\text{conjugation}$.


Suppose that the finitely presented group H is given a decomposition as a free product with amalgamation $H = A *_S B$. Then clearly

$$\mathcal{H}(H) \cong \mathcal{H}(A) \times_{\mathcal{H}(S)} \mathcal{H}(B) := \{(\rho_A, \rho_B) \in \mathcal{H}(A) \times \mathcal{H}(B) \mid \rho_A|_S = \rho_B|_S\}.$$

The Lie group G acts diagonally by conjugation on $\mathcal{H}(H)$ with quotient $\chi(H)$, and similarly for the subgroups A, B , and S . There is a surjective map

$$\mathcal{H}(A) \times_{\mathcal{H}(S)} \mathcal{H}(B) \rightarrow \chi(A) \times_{\chi(S)} \chi(B) := \{(c_A, c_B) \in \chi(A) \times \chi(B) \mid c_A|_S = c_B|_S\}$$

which factors through $\chi(H)$. These form a diagram:



Let $\text{Stab}(\rho) \subset G$ denote the centralizer of the image of a representation ρ :

$$(12) \quad \text{Stab}(\rho) = \{g \in G \mid \rho(x) = g\rho(x)g^{-1} \text{ for all } x\}$$

Lemma 4.2 *Let $(c_A, c_B) \in \chi(A) \times_{\chi(S)} \chi(B)$ and choose $(\rho_A, \rho_B) \in p_3^{-1}(c_A, c_B)$ (so that $\rho_A|_S = \rho_B|_S$). Then the fiber $p_2^{-1}(c_A, c_B)$ is homeomorphic to the double coset space*

$$\text{Stab}(\rho_A) \backslash \text{Stab}(\rho_A|_S) / \text{Stab}(\rho_B).$$

Remark Lemma 4.2 can be applied in situations when we consider representation spaces and character varieties that place conjugation-invariant conditions on the value that the representations take on specified elements in the groups H , A , B , and S . For our purposes we will be interested in the traceless representations which arise in the definition of R and R^{\natural} .

Proof of Lemma 4.2 Denote by ρ_S the restriction of ρ_A to S . We first identify the fiber $p_3^{-1}(c_A, c_B)$ with a quotient of $G \times \text{Stab}(\rho_S)$. If $(\rho'_A, \rho'_B) \in p_3^{-1}(c_A, c_B)$ then there are $g_1, g_2 \in G$ so that $(\rho'_A, \rho'_B) = (g_1\rho_A g_1^{-1}, g_2\rho_B g_2^{-1})$. Since ρ'_A and ρ'_B agree on S , (as do ρ_A and ρ_B) it follows that $g_1^{-1}g_2 \in \text{Stab}(\rho_S)$. Thus the map $G \times \text{Stab}(\rho_S) \rightarrow p_3^{-1}(c_A, c_B)$ taking (g, t) to $(g\rho_A g^{-1}, gt\rho_B(gt)^{-1})$ is surjective.

Since p_1 is surjective, it maps $p_3^{-1}(c_A, c_B)$ surjectively to $p_2^{-1}(c_A, c_B)$. Suppose that $(g, t), (g', t') \in G \times \text{Stab}(\rho_S)$. Then

$$p_1(g'\rho_A g'^{-1}, g't'\rho_B(g't')^{-1}) = p_1(g\rho_A g^{-1}, gt\rho_B(gt)^{-1})$$

if and only if there is an $h \in G$ so that

$$(g'\rho_A g'^{-1}, g't'\rho_B(g't')^{-1}) = (hg\rho_A(hg)^{-1}, hgt\rho_B(hgt)^{-1}).$$

This is equivalent to $g'^{-1}hg \in \text{Stab}(\rho_A)$ and $(g't')^{-1}hgt \in \text{Stab}(\rho_B)$. Writing $a = g'^{-1}hg$ and $b = (g't')^{-1}hgt$ we see that $h = g'ag^{-1}$ and $t'^{-1}at = b$. It follows that (g, t) and (g', t') correspond to the same element of $p_2^{-1}(c_A, c_B)$ if and only if there exist $a \in \text{Stab}(\rho_A)$ and $b \in \text{Stab}(\rho_B)$ so that $t' = atb^{-1}$. \square

We now use Lemma 4.2 to compare $R(Y, K)$ and $R^{\natural}(Y, K)$. To do this, write

$$(13) \quad (Y, K^{\natural} \cup W) = (Y \setminus N(K), \emptyset) \cup_{\partial N(K)} (N(K), K^{\natural} \cup W).$$

Suppose $\rho_A \in \chi(Y, K)$ and $\rho_B \in R^{\natural}(N(K), K)$ map to the same point $\rho_S \in \chi(\partial N(K))$. Conjugate ρ_B so that the restrictions of ρ_A and ρ_B to $\partial N(K)$ agree. Proposition 4.1 shows $\rho_B \in R^{\natural}(N(K), K)$ has nonabelian image. Moreover, since $\rho_B(\mu_K) \in C(\mathfrak{i})$, then the restriction ρ_A to $Y \setminus N(K)$ of any representation $\rho \in R^{\natural}(Y, K)$ lies in

$R(Y, K) \subset \chi(Y, K)$. Furthermore, the restriction ρ_S to the separating torus $\partial N(K)$ is abelian but noncentral, since it sends the meridian μ_K into $C(\mathbf{i})$. Thus $\text{Stab}(\rho_S) \cong S^1$.

From Lemma 4.2 we conclude that the fiber of the restriction map

$$(14) \quad R^{\natural}(Y, K) \rightarrow R(Y, K) \times_{\chi(\partial N(K))} R^{\natural}(N(K), K)$$

over $\rho_A * \rho_B$ is

$$\text{Stab}(\rho_A) \backslash S^1 / \{\pm 1\}.$$

This is a single point if the restriction ρ_A of ρ to $Y \setminus N(K)$ has abelian image, and a circle if the restriction has nonabelian image.

Proposition 4.1 asserts that the restriction map $R^{\natural}(N(K), K) \rightarrow \chi(\partial N(K))$ is injective. This allows us to identify the restriction map of Equation (14) with the surjective map

$$(15) \quad R^{\natural}(Y, K) \rightarrow R(Y, K).$$

In summary, we have the following.

Proposition 4.3 *Every representation in $R^{\natural}(Y, K)$ has nonabelian image. The forgetful map $R^{\natural}(Y, K) \rightarrow R(Y, K)$ is a surjection. The fiber over a conjugacy class ρ is either a circle or a point, and this depends on whether ρ has nonabelian or abelian image. \square*

Lemma 4.2 is also useful when studying $R^{\natural}(Y, K)$ via a decomposition of the pair (Y, K) along a four-punctured 2-sphere. Suppose that $Y = Y_1 \cup_{S^2} Y_2$ is a decomposition of Y along a 2-sphere $S^2 \subset Y$ which intersects K in four points a, b, c, d . We assume that $H \cup W$ lies in the interior of Y_2 . Then Lemma 4.2 allows us to identify $R^{\natural}(Y, K)$ as a fiber product of $R(Y_1, K_1)$ and $R^{\natural}(Y_2, K_2)$ over the second pillowcase $R(S^2, \{a, b, c, d\})$ (where each $K_i = K \cap Y_i$ is a union of two arcs). Precisely, there is a restriction map

$$(16) \quad R^{\natural}(Y, K) \rightarrow R(Y_1, K_1) \times_{R(S^2, \{a, b, c, d\})} R^{\natural}(Y_2, K_2)$$

whose fiber over $\rho_1 * \rho_2$ is

$$\text{Stab}(\rho_1) \backslash \text{Stab}(\rho_{S^2}) / \text{Stab}(\rho_2).$$

Now $\text{Stab}(\rho_2) = \pm 1$, since $H \cup W$ is contained in Y_2 . The stabilizer of any nonsingular point in the pillowcase $R(S^2, \{a, b, c, d\})$ is just the center ± 1 , so the fiber of the map in Equation (16) over a pair (ρ_1, ρ_2) is a single point if the restriction to S^2 is not one of the four abelian conjugacy classes in $R(S^2, \{a, b, c, d\})$.

However, at one of the four abelian conjugacy classes, the stabilizer is S^1 , and hence the fiber of the map in Equation (16) is

$$\text{Stab}(\rho_1) \backslash S^1,$$

which is a circle if ρ_1 is nonabelian and a single point if ρ_1 is abelian. In summary, we have the following.

Proposition 4.4 *Suppose that $S^2 \subset Y$ is a separating 2–sphere meeting the knot K transversely in four points a, b, c, d and that $H \cup W$ lies in Y_2 . Restricting to the two pieces in the decomposition $Y = Y_1 \cup_{S^2} Y_2$ gives a surjection*

$$R^{\natural}(Y, K) \rightarrow R(Y_1, K_1) \times_{R(S^2, \{a, b, c, d\})} R^{\natural}(Y_2, K_2)$$

whose fibers are single points, with the exception of the fibers over the nonabelian representations in $R(Y_1, K_1)$ which restrict to one of the four abelian (corner) points in $R(S^2, \{a, b, c, d\})$. The fiber above these latter points is a circle. \square

In the examples below we will take Y_2 to be a 3–ball intersecting K in a pair of unknotted arcs. We will explain, for torus and 2–bridge knots, how to choose the 3–ball judiciously so that the restriction of every nonabelian representation in $R(Y_1, K_1)$ to $R(S^2, \{a, b, c, d\})$ avoids the corner points. The map of Proposition 4.4 then has point fibers. Then a perturbation will be applied to make $R^{\natural}(Y_2, K_2)$ generic, in the sense that the fiber product in Proposition 4.4 has finitely many points, corresponding to a finite intersection of $R(Y_1, K_1)$ and $R^{\natural}(Y_2, K_2)$ in $R(S^2, \{a, b, c, d\})$. These points will provide a finite generating set for the reduced singular instanton chain complex.

5 Knots with simple representation varieties

Much is known about the spaces $\chi(Y, K)$ for various (Y, K) , starting with Klassen’s influential article [20]. The image of $\chi(S^3, K)$ in the pillowcase $\chi(\partial N(K))$ is also a well studied space: for example, it appears as part of the real locus of the A –polynomial of [7]. The identification of the algebraic count of the intersections of this image with the circle $S(\mathbf{i})$ of Equation (5), corresponding to the line $\{x = \frac{\pi}{2}\}$ (and more generally circles corresponding to the line $\{x = \theta\}$ for $\theta \in [0, \pi]$) with Levine–Tristram knot signatures is explained in [30; 14] and Herald [13].

Definition 5.1 We say a knot K in S^3 has a *simple representation variety*, or $\chi(S^3, K)$ is *generic* if $\chi(S^3, K)$ is homeomorphic to a 1-complex made out of:

(i) An arc of abelian representations, parameterized by

$$t \in [0, \pi] \mapsto \alpha_t: \pi_1(S^3 \setminus N(K)) \rightarrow H_1(S^3 \setminus N(K)) \rightarrow \text{SU}(2), \quad \alpha_t(\mu_K) = e^{ti}.$$

(ii) A finite number of smooth arcs of representations

$$n_i: [0, 1] \rightarrow \text{Hom}(\pi_1(S^3 \setminus N(K)), \text{SU}(2)), \quad i = 1, \dots, k,$$

whose interior points are nonabelian and whose endpoints $n_i(0)$ and $n_i(1)$ equal $\alpha_{s_{i,0}}$ and $\alpha_{s_{i,1}}$ for a pairwise distinct set of points $s_{1,0}, s_{1,1}, s_{2,0}, s_{2,1}, \dots, s_{k,0}, s_{k,1} \in [0, \pi]$.

(iii) A finite number of disjoint smooth circles

$$c_i: S^1 \subset \chi(S^3, K), \quad i = 1, \dots, m,$$

of nonabelian representations disjoint from the arcs α_t and n_i .

(iv) The restriction map $\chi(S^3, K) \rightarrow \chi(\partial N(K))$ restricts to an immersion on each arc α_t, n_i and each circle c_i . This immersion is transverse to the circle $S(i)$ of representations which are traceless on the meridian (Equation (5)).

The points $s_{1,0}, s_{1,1}, \dots, s_{k,0}, s_{k,1}$ are called *bifurcation points*, since they correspond to places where the irreducible representations bifurcate from the abelian representations. This implies that the roots of the Alexander polynomial of K which lie on the unit circle are those of the form $e^{2s_{i,j}i}$, and are simple and distinct [20; 13].

Not all knots have simple representation varieties, but the results of [12; 13] imply that perturbation data (as explained below) π can be found so that the perturbed moduli space $\chi_\pi(S^3, K)$ has the properties listed above. For small perturbations, the distinct points $s_{1,0}, s_{1,1}, \dots, s_{k,0}, s_{k,1}$ are close, but not necessarily equal to, the set of roots of the Alexander polynomial.

5.1 Torus knots

All torus knots have simple representation varieties. In fact $\chi(S^3, T_{p,q})$ is a union of an arc α_t of abelian representations as described above, and $(p-1)(q-1)/2$ arcs of nonabelian representations that limit to abelian representations at roots of the Alexander polynomial [20]. We review some of the details.

The (p, q) torus knot group has presentation $\langle x, y \mid x^p = y^q \rangle$. In this presentation $\mu_K = x^s y^r$ and $\lambda_K = x^p (x^s y^r)^{-pq}$ where $pr + qs = 1$ (see eg Burde [5, Proposition 3.28]). The arc of conjugacy classes of abelian representations is parameterized by

$\alpha_t(\mu_K) = e^{ti}$, $t \in [0, \pi]$. In terms of the generators x and y we have $\alpha_t(x) = e^{qti}$ and $\alpha_t(y) = e^{pti}$.

Since $x^p = y^q$ is central, then any nonabelian representation ρ will send $x^p = y^q$ to ± 1 . Hence x is sent to a p^{th} root of ± 1 in $SU(2)$ and y is sent to a q^{th} root of ± 1 . If ρ is conjugated so that $\rho(\mu_K) = e^{mi}$ for some $m \in \mathbb{R}$, then we have that $\rho(\lambda_K) = \rho(x^p \mu_K^{-pq}) = \pm e^{-pqmi}$. Hence the image of the nonabelian part of $\chi(S^3, K)$ in the pillowcase lies in the straight lines $\ell = -pqm + k\pi$ (via the quotient $\mathbb{R}^2 \rightarrow \chi(T)$, taking (m, ℓ) to the conjugacy class of $\mu_K \mapsto e^{mi}$, $\lambda_K \mapsto e^{\ell i}$). In fact the images in the pillowcase are embedded arcs that start and end at α_{s_i} where $\Delta_K(e^{2s_i i}) = 0$ (see [20, Theorem 19] and the discussion which follows its proof).

Each arc of nonabelian representations is completely determined by the choice of p^{th} and q^{th} root of ± 1 . Explicitly, given a pair a, b of integers of the same parity, then the assignment

$$(17) \quad x \mapsto e^{a\pi i/p}, \quad y \mapsto \cos(b\pi/q) + \sin(b\pi/q)(\cos(u)\mathbf{i} + \sin(u)\mathbf{j}), \quad u \in [0, \pi]$$

defines an arc of representations whose projection to the space of conjugacy classes is one-to-one. Conversely, given any representation ρ , $\rho(x)$ can be conjugated to $e^{a\pi i/p}$ and $\rho(y)$ can be conjugated to $e^{b\pi i/q}$ for some choice of integers of the same parity, and some choice of conjugating elements. It follows that any representation can be conjugated to a unique point on the corresponding path of Equation (17): we first conjugate ρ so that $\rho(x)$ has the desired image, and then further conjugate by an element in the circle through $\rho(x)$ so that y has the form of (17). This sets up a bijection between arcs of nonabelian representations and pairs (a, b) of the same parity satisfying $0 < a < p$ and $0 < b < q$. For more details see [20, Theorem 1].

The endpoints of this arc are abelian, and hence lie on the arc α_t . These endpoints are determined by computing where they send μ_K . When $u = 0$ the pair (μ_K, λ_K) is sent to $(e^{(as/p+br/q)\pi i}, e^{pr(a-b)\pi i})$. Hence the endpoint with $u = 0$ equals α_{s_0} where $(s_0, 0)$ and $((as/p + br/q)\pi, pr(a-b)\pi)$ in \mathbb{R}^2 map to the same point in the pillowcase. Similarly one determines where the endpoint with $u = \pi$ is sent.

These considerations suffice to show that all torus knots have simple representation varieties and to completely determine $\chi(S^3, K)$ and its image in the pillowcase. We illustrate two examples.

The simplest example is the trefoil knot, ie the $(2, 3)$ torus knot. It has one arc of nonabelian representations, attached to the arc of abelian representations at the points $s_0 = \pi/6$ and $s_1 = 5\pi/6$, corresponding to the fact that $\Delta_K(e^{2s_i}) = 0$, $i = 0, 1$. This arc of nonabelian representations is sent to an arc of slope -6 in the pillowcase. Notice that the map $\chi(S^3, K) \rightarrow \chi(\partial N(K))$ fails to be injective at precisely the two

points in $\chi(S^3, K)$ sent to the circle $S(i)$ in the pillowcase (see Equation (5)). In particular, $R(S^3, K)$ consists of exactly two points, the abelian representation $\alpha_{\pi/2}$, and the nonabelian representation corresponding to $u = \pi/2$.




Figure 4: The character variety $\chi(S^3, T_{2,3})$ and its image in the pillowcase

Next we consider the $(3, 5)$ torus knot. This knot has Alexander polynomial $\Delta_K(t) = t^8 - t^7 + t^5 - t^4 + t^3 - t + 1$. Thus $\Delta_K(e^{2s_i}) = 0$ when $s_i = \pi/15, 2\pi/15, 4\pi/15, 7\pi/15, 8\pi/15, 11\pi/15, 13\pi/15$ and $14\pi/15$. The corresponding α_{s_i} form the endpoints of the four arcs of nonabelian representations, determined by the four possible choices $(a, b) = (1, 1), (1, 3), (2, 2)$ and $(2, 4)$. One calculates that the first arc has endpoints $\{\alpha_{\pi/15}, \alpha_{11\pi/15}\}$, the second $\{\alpha_{7\pi/15}, \alpha_{13\pi/15}\}$, the third $\{\alpha_{2\pi/15}, \alpha_{8\pi/15}\}$, and the fourth $\{\alpha_{4\pi/15}, \alpha_{14\pi/15}\}$. The restriction $\chi(S^3, K) \rightarrow \chi(\partial N(K))$ is far from injective, although it is injective when restricted to each arc of nonabelian representations. In this example, the space $R(S^3, K)$ consists of five points: the abelian $\alpha_{\pi/2}$, and four other points, one interior to each of the nonabelian arcs. Two of these four points are sent to the point in the pillowcase with (\mathbb{R}^2) coordinates $(\pi/2, 3\pi/2)$; the other two are sent to the point with coordinates $(\pi/2, \pi/2)$. In particular, this implies that these traceless representations are not binary dihedral, since the longitude is not sent to ± 1 . We illustrate this in Figure 5, where we have only drawn the image of the first arc in

the pillowcase for clarity. It intersects the circle $S(i)$ (in the figure, the vertical line $\gamma = \pi/2$) in the point with coordinates $(\pi/2, 3\pi/2)$.




Figure 5: The character variety $\chi(S^3, T_{3,5})$ and part of its image in the pillowcase

The representation variety $\chi(S^3, K)$ for K a general (p, q) torus knot consists of an arc α_t of abelian representations and $(p - 1)(q - 1)/2$ arcs of nonabelian representations with distinct endpoints α_{s_i} on α_t , parameterized by pairs of integers (a_i, b_i) in Equation (17) above.

For the purposes of describing the image of each arc in the pillowcase, it is more convenient to parameterize the paths by their endpoints, rather than the pairs (a_i, b_i) . Therefore $\chi(S^3, K)$ for K any (p, q) torus knot and its image in the pillowcase $\chi(\partial N(K))$ is completely described by the data consisting of the pair (p, q) and an additional $(p - 1)(q - 1)/2$ pairs (c_i, d_i) (for $i = 1, \dots, (p - 1)(q - 1)/2$ with $c_i < d_i$) of integers determined by the requirement that the endpoints of the i^{th} arc of nonabelian representations are $\frac{c_i \pi}{pq}$ and $\frac{d_i \pi}{pq}$; these satisfy $\Delta_K(e^{(2c_i \pi)/(pq)}) = 0 = \Delta_K(e^{(2d_i \pi)/(pq)})$. The pairs (c_i, d_i) determine and are determined by the (a_i, b_i) , but the relationship is awkward to describe explicitly, and so we will use (c_i, d_i) to prescribe each arc of nonabelians.

The subvariety $R(S^3, K)$ of $\chi(S^3, K)$ contains $|\sigma(K)|/2 + 1$ points where $\sigma(K)$ denotes the signature of K [30; 13; 14]. One of these points is $\alpha_{\pi/2}$, the rest correspond

to intersections of some of the arcs of nonabelian representations with the circle $S(i)$ of Equation (5).

The image of the arc of abelians α_t in the pillowcase has slope 0, and each arc of nonabelian representations maps to a line of slope $-pq$. Hence, once we know their endpoints, we know how they map to the pillowcase. In particular, there is always one abelian traceless representation $\alpha_{\pi/2}$, and one nonabelian traceless representation for each pair (c_i, d_i) as above satisfying $c_i < \frac{pq}{2} < d_i$.

For example, for the trefoil $T_{2,3}$, the data (c_i, d_i) is just $(1, 5)$ and so $R(S^3, T_{2,3})$ has one abelian and one nonabelian traceless representation. For $K = T_{3,5}$, the data are

$$(1, 11), \quad (7, 13), \quad (2, 15), \quad (4, 14),$$

and so $R(S^3, T_{3,5})$ has one abelian and four nonabelian traceless representations, since each corresponding interval contains $\frac{15}{2}$. The data clearly determines the information in Figure 5.

In general, not every arc of nonabelian representations contains a point of $R(S^3, K)$: the inequality $|\sigma(K)|/2 \leq (p-1)(q-1)/2$ may be strict. For example, the data for the $(3, 7)$ -torus knot are

$$(1, 13), \quad (11, 17), \quad (5, 19), \quad (2, 16), \quad (4, 10), \quad (8, 20),$$

and so the arcs determined by the pairs $(11, 17)$ and $(4, 10)$ do not contain traceless representations. This corresponds to the fact that $\sigma/2 = 4$. Similarly, the data for the $(4, 9)$ -torus knot are

$$(1, 17), \quad (15, 33), \quad (23, 31), \quad (7, 25), \quad (2, 34), \quad (14, 22), \quad (6, 30), \\ (10, 26), \quad (19, 35), \quad (3, 21), \quad (5, 13), \quad (11, 29),$$

and $\sigma/2 = 8$. It is known that the signature of a nontrivial torus knot is always nonzero (see eg Kirk and Livingston [19]) from which it follows that $R(S^3, K)$ always contains at least one point on a nonabelian arc. A much deeper result is the theorem of Kronheimer and Mrowka [22] that every nontrivial knot in S^3 admits a nonabelian traceless representation.

Applying Proposition 4.3 we conclude that for K the (p, q) torus knot, $R^{\natural}(S^3, K)$ consists of one isolated point α , corresponding to $\alpha_{\pi/2}$, and $|\sigma(K)|/2$ circles, one for each irreducible traceless representation of $\pi_1(S^3 \setminus K)$.

5.2 2-bridge knots

We recall some facts about $\chi(S^3, K)$ when K is a 2-bridge knot. Klassen [20] identified the spaces $\chi(S^3, K)$ for the twists knots. Building on work of Riley [37]

(who considered $\mathrm{SL}(2, \mathbb{C})$ representations), Burde [5] determined $\chi(S^3, K)$ for all 2–bridge knots. In contrast to torus knots, the image of $\chi(S^3, K)$ in the pillowcase is not given by linear equations, but rather by more complicated polynomial equations. However, we will give explicit (and linear) equations which determine the *traceless* representation varieties $R(S^3, K)$ and $R^{\natural}(S^3, K)$ for any 2–bridge knot in Section 10 below.

For the m –twist knot (this is the 2–bridge knot corresponding to the continued fraction expansion $[1, 1, m]$), $\chi(S^3, K)$ is a union of the arc α_t of abelian representations, $[\frac{m}{2}]$ circles of nonabelian representations, and, if m is odd, one arc of nonabelian representations with endpoints on the arc α_t corresponding to the two roots of the Alexander polynomial on the unit circle. Each circle contributes two points to $R(S^3, K)$, and the arc of nonabelian representations (when m is odd) meets $R(S^3, K)$ in one point. As for all knots, the abelian representation $\alpha_{\pi/2}$ lies in $R(S^3, K)$.

Thus, for K the m –twist knot, $R(S^3, K)$ consists of one abelian representation, and $2[\frac{m}{2}] = m$ nonabelian representations if m is even, and $2[\frac{m}{2}] + 1 = m$ nonabelian representations if m is odd. Using Proposition 4.3 it follows that $R^{\natural}(S^3, K)$ consists of one isolated point α and m circles.

For example, when K is the figure 8 knot, corresponding to $m = 2$, $\chi(S^3, K)$ consists of the arc α_t of abelian representations and a disjoint circle of nonabelian representations. A construction of this circle and arc, together with an explicit description of the image of $\chi(S^3, K) \rightarrow \chi(\partial N)$ can be found in [18, Proposition 5.4]. From this one concludes that $R(S^3, K)$ consists of exactly three points: the arc α_t contributes the point $\alpha_{\pi/2}$ to $R(S^3, K)$, and the circle contributes two points to $R(S^3, K)$. This identifies $R^{\natural}(S^3, K)$ as the union of an isolated point α and two circles.

We illustrate the more complicated example of the m twisted double of the unknot when $m = 5$, this is the knot 7_2 in the knot tables. The space $\chi(S^3, K)$ consists of two circles and one arc of nonabelians, and the arc α_t of abelians. The Alexander polynomial is $3 - 5t + 3t^3$, with roots $(5 \pm \sqrt{-11})/6 \approx e^{(0.1)2\pi i}$. The arc of nonabelians is embedded, but the two circles of nonabelians are immersed in the pillowcase with one transverse double point at $(m, \ell) = (\frac{\pi}{2}, 0)$. The proof of these facts can be found in Burde’s article [5].

In particular, the space $R(S^3, K)$ contains six points: the abelian point $\alpha_{\pi/2}$, the midpoint of the arc of nonabelians, and two points on each of the two circles of nonabelians. Each of these six points is mapped to the point $(\frac{\pi}{2}, 0)$ in the pillowcase. The space $R^{\natural}(S^3, K)$ thus consists of one isolated point α corresponding to $\alpha_{\pi/2}$ via Proposition 4.3 and five circles corresponding to the five nonabelian representations in $R(S^3, K)$.




Figure 6: $\chi(S^3, 7_2)$ and its image in the pillowcase

6 Traceless representations of certain tangles

Having established the presence of circles in $R^{\mathbb{H}}(Y, K)$ in Proposition 4.3, we are faced with the problem that this set is never finite (except for the unknot) and yet should give a generating set for the reduced instanton homology of (Y, K) , since it appears as the critical set of the relevant Chern–Simons functional. The circles arise via the mechanism of Lemma 4.2 because of the presence of the torus (ie the boundary of the tubular neighborhood of K which contains $H \cup W$) along which we can bend a representation. Holonomy perturbations of the Chern–Simons functional (described below) are used to correct this problem. We will show below that the circles can be eliminated by using a holonomy perturbation that lies in a 3–ball intersecting K in two unknotted arcs and containing $H \cup W$. Thus we turn our attention to the representation spaces corresponding to a pair of arcs in a ball.

6.1 The space $R(B^3, A_1 \cup A_2)$

Consider the space of conjugacy classes of $SU(2)$ representations $R(B^3, A_1 \cup A_2)$ of the complement of a pair of unknotted arcs in a 3–ball which send their meridians to elements in $C(\mathbf{i})$, as illustrated in Figure 7. This corresponds exactly to the subspace of $R(S^2, \{a, b, c, d\})$ consisting of those representations which satisfy $\rho(a) = \rho(d)$ and $\rho(b) = \rho(c)$.

We have the following simple observation.

Proposition 6.1 *The space $R(B^3, A_1 \cup A_2)$ can be identified with the arc $[0, \pi]$ via*

$$a \mapsto \mathbf{i}, \quad b \mapsto e^{\gamma k} \mathbf{i}, \quad c \mapsto e^{\gamma k} \mathbf{i}, \quad d \mapsto \mathbf{i},$$

for $\gamma \in [0, \pi]$. The restriction map

$$R(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$$

is injective, with image the diagonal arc $\psi(\gamma, \gamma)$, $\gamma \in [0, \pi]$. □




Figure 7: Two arcs in a 3-ball, $(B^3, A_1 \cup A_2)$

Proposition 6.1 is illustrated in Figure 8, where $R(S^2, \{a, b, c, d\})$ is represented as an identification space obtained from the rectangle $[0, \pi] \times [0, 2\pi]$.

We now consider the space $R^{\natural}(B^3, A_1 \cup A_2)$ consisting of conjugacy classes of $SU(2)$ representations which send a, b, c, d and h to $C(\mathbf{i})$ and w to -1 , where h denotes the meridian of H and w the meridian to the arc W . As before, we may conjugate so that (abusing notation)

$$a = d = \mathbf{i}, \quad b = c = e^{\gamma k} \mathbf{i}, \quad w = -1,$$

with $\gamma \in [0, \pi]$. The relation $[\bar{a}, \bar{h}] = w = -1$ implies that

$$h = e^{\tau \mathbf{i}} \mathbf{j}$$

for some τ .

Hence the image of the restriction $R^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ is equal to the image $R(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$. Both are given by the arc $\psi(\gamma, \gamma)$, $\gamma \in [0, \pi]$, as illustrated above. In the case of $R^{\natural}(B^3, A_1 \cup A_2)$, however, the restriction is not injective. In fact, there is a circle action on $R^{\natural}(B^3, A_1 \cup A_2)$ defined by the parameter τ . This action is free away from the two fixed points, which occur when $\sin \gamma = 0$. The restriction map $R^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ factors through the orbit map, which is just the map $R^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(B^3, A_1 \cup A_2)$ that forgets h and w . Summarizing, we have the following.




Figure 8: The image of $R(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ is the arc $\theta = \gamma$.

Proposition 6.2 $R^{\natural}(B^3, A_1 \cup A_2)$ is a 2-sphere, corresponding to

$$a \mapsto \mathbf{i}, \quad b \mapsto e^{\gamma \mathbf{k}} \mathbf{i}, \quad c \mapsto e^{\gamma \mathbf{k}} \mathbf{i}, \quad d \mapsto \mathbf{i}, \quad h \mapsto e^{\tau \mathbf{i}} \mathbf{j}, \quad w \mapsto -1,$$

for $(\gamma, \tau) \in [0, \pi] \times [0, 2\pi]$, cylindrical coordinates on S^2 . The restriction to the pillowcase

$$R^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$$

has image the diagonal arc $\psi(\gamma, \gamma)$, $\gamma \in [0, \pi]$, and the fibers are circles parameterized by τ over the interior of the arc and a single point over each endpoint. \square

7 Perturbations

7.1 Holonomy perturbations of the Chern–Simons functional and its critical points

We recall some aspects of the definition and basic properties of instanton homology and the related holonomy perturbations that we need. We provide the minimal description needed for our purposes, and refer to the series of articles [26; 25; 21] for details.

To the triple (Y, K^{\natural}, W) , Kronheimer and Mrowka assign *singular bundle data* \mathbf{P} which consists of an $\mathrm{SO}(3)$ bundle over $Y \setminus K^{\natural}$ whose second Stiefel–Whitney class is Poincaré dual to W , as well as a certain kind of $O(2)$ reduction near $K \cup H$. The singular bundle data in turn give rise to an affine space of $\mathrm{SO}(3)$ connections $\mathcal{C}(Y, K; \mathbf{P})$ which have a prescribed singularity near K^{\natural} , with tangent spaces identified with spaces of bundle-valued 1-forms with appropriate behavior near K . The determinant one gauge group $\mathcal{G}(Y, K; \mathbf{P})$ acts on $\mathcal{C}(Y, K; \mathbf{P})$.

The Chern–Simons functional $\mathrm{CS}: \mathcal{C}(Y, K; \mathbf{P}) \rightarrow \mathbb{R}$ has the property that its gradient vector field with respect to the L^2 inner product on 1-forms is

$$(\mathrm{grad} \mathrm{CS})_B = - * F_B.$$

The critical points of the Chern–Simons functional therefore consist of *flat* singular $\mathrm{SO}(3)$ connections. The gauge group $\mathcal{G}(Y, K; \mathbf{P})$ preserves the set of flat connections. As explained in [25], the set of gauge equivalence classes of critical points, $\mathfrak{C}(Y, K; \mathbf{P})$, is identified, via the holonomy, with $R^{\natural}(Y, K)$.

Any critical point A of CS restricts to an honest flat connection on $Y - N(K)$. The bundle is trivial over $Y \setminus N(K)$ since $W \subset N(K)$ and hence the holonomy of A lifts uniquely to $\mathrm{SU}(2)$, giving a representation $\rho_A: \pi_1(Y \setminus N(K)) \rightarrow \mathrm{SU}(2)$ which sends the meridians to traceless unit quaternions, ie $\rho_A \in R(Y, K)$.

The reduced instanton homology $I^{\natural}(Y, K)$ is the Morse–Floer homology of a $\mathbb{Z}/4$ graded-chain complex $CI^{\natural}(Y, K)$ associated to CS. The chain complex $CI^{\natural}(Y, K)$ should therefore be generated by the set of gauge equivalence classes $\mathfrak{C}(Y, K; \mathbf{P})$ of critical points of CS, which are identified with $R^{\natural}(Y, K)$. For any nontrivial knot K in S^3 , however, Kronheimer and Mrowka have proved [22] that there exist nonabelian traceless $\mathrm{SU}(2)$ representations. Thus Proposition 4.3 shows that $R^{\natural}(S^3, K)$ always contains circles. Since the Morse complex of CS must be finitely generated in order to define a sensible theory, we see that we must perturb CS to ensure that its critical points are isolated and finite in number.

This is achieved by adding to CS a *holonomy perturbation* $h_{\pi}: \mathcal{C}(Y, K; \mathbf{P}) \rightarrow \mathbb{R}$. The function h_{π} is constructed from data consisting of a collection of embeddings $e_i: S^1 \times D^2 \rightarrow Y \setminus (K^{\natural} \cup W)$ and some conjugation invariant functions $g_i: \mathrm{SU}(2) \rightarrow \mathbb{R}$. Following [26; 25], $\pi = \{e_i, g_i\}$ is called the *perturbation data*, and determines the function $h_{\pi}: \mathcal{C}(Y, K; \mathbf{P}) \rightarrow \mathbb{R}$ by the formula

$$h_{\pi}(A) = \sum_i \int_{D^2} g_i(\mathrm{hol}_{S^1 \times \{x\}}(A)) \eta(x) d^2x,$$

where $\eta(x)$ is a (fixed) radially symmetric smooth cutoff function on D^2 , and, given $x \in D^2$, $\text{hol}_{S^1 \times \{x\}}(A)$ denotes the holonomy of A around the loop $t \mapsto e_i(e^{it}, x)$, $t \in [0, 2\pi]$.

Kronheimer and Mrowka consider more general perturbation functions, but these functions suffice for our purposes. In fact, we will only require one embedding $e: S^1 \times D^2 \rightarrow Y \setminus (K^{\natural} \cup W)$, and the function g will be taken to be $g(q) = \epsilon \text{Re}(q)$ for some small $\epsilon > 0$.

Denote by P_j the image of e_j , a solid torus. We will abuse terminology and call P_j a *perturbation curve*. Denote by p_j the meridian of P_j , ie the (suitably based) loop $e_j(* \times \partial D^2)$, and by ℓ_j a choice of longitude, eg $e_j(S^1 \times *)$. Let $P = \bigsqcup_j P_j$. Denote by μ_K, μ_H, μ_W (suitably based) meridians of K, H, W respectively.

Any critical point of $\text{CS} + h_\pi$ is the gauge equivalence class of a connection which is flat outside the image P of the embeddings $e_i: S^1 \times D^2 \rightarrow Y \setminus (K^{\natural} \cup W)$. Therefore, critical points of the perturbed functional $\text{CS} + h_\pi$ are identified with conjugacy classes of representations $\pi_1(Y \setminus (K \cup H \cup W \cup P)) \rightarrow \text{SU}(2)$ which send the meridians μ_K, μ_H to $C(\mathbf{i})$, and μ_W to -1 , and which also satisfy a certain constraint determined by g_j when restricted to the meridian and longitude of the j^{th} boundary torus of P .

Lemma 61 of [12] identifies the constraint and shows that given any list of smooth functions $f_j: \mathbb{R} \rightarrow \mathbb{R}$ satisfying $f_j(-x) = -f_j(x)$ and f_j is 2π periodic (for example, $f_j(x) = \sin(x)$), there exist g_j as above so that the constraint on the j^{th} boundary torus is given by the *perturbation condition*:

If the representation $\rho: \pi_1(Y \setminus (K \cup H \cup W \cup P)) \rightarrow \text{SU}(2)$ takes the meridian and longitude p_j, ℓ_j of the j^{th} component of P to $e^{v_j} Q_j$ and $e^{\beta_j} Q_j$ respectively, for some $Q_j \in C(\mathbf{i})$, then $v_j = f_j(\beta_j)$.

We denote the space of conjugacy classes of such representations by $R_\pi^{\natural}(Y, K)$. Explicitly, $R_\pi^{\natural}(Y, K)$ is the space of $\text{SU}(2)$ -conjugacy classes of representations

$$\rho: \pi_1(Y \setminus (K \cup H \cup W \cup P)) \rightarrow \text{SU}(2)$$

satisfying

$$(18) \quad \begin{aligned} &\rho(\mu_K), \rho(\mu_H) \in C(\mathbf{i}), \quad \rho(\mu_W) = -1, \\ &v_j = f_j(\beta_j) \quad \text{when } \rho(p_j) = e^{v_j} Q_j \text{ and } \rho(\ell_j) = e^{\beta_j} Q_j. \end{aligned}$$

We can apply similar constructions to $\chi(Y, K)$, the space of conjugacy classes of all representations $\pi_1(Y \setminus N(K)) \rightarrow \text{SU}(2)$. One can perturb using embeddings $e_i: S^1 \times D^2 \rightarrow Y \setminus N(K)$; the resulting critical set is denoted by $\chi_\pi(Y, K)$, and

is identified with those representations $\pi_1(Y \setminus (K \cup P)) \rightarrow \text{SU}(2)$ satisfying the constraints

$$(19) \quad v_j = f_j(\beta_j) \quad \text{when } \rho(p_j) = e^{v_j} Q_j \quad \text{and } \rho(\ell_j) = e^{\beta_j} Q_j$$

on each ∂P_j .

When P lies outside the tubular neighborhood $N(K)$ of K containing $K^{\natural} \cup W$, restriction defines a map $R_{\pi}^{\natural}(Y, K) \rightarrow \chi_{\pi}(Y, K)$ with image which we denote by $R_{\pi}(Y, K)$. Just as in the unperturbed case, the fibers of the restriction map

$$R_{\pi}^{\natural}(Y, K) \rightarrow R_{\pi}(Y, K)$$

are circles over every representation with nonabelian image, and a point over every representation with abelian image.

In particular, any choice of perturbation data $\pi = \{e_i, g_i\}$ which make $R_{\pi}^{\natural}(Y, K)$ a finite union of isolated points must include at least one perturbation curve $P \subset Y \setminus (K^{\natural} \cup W)$ which intersects the separating torus $\partial N(K)$ essentially and hence must link $H \cup W$ in some way. This observation motivates using the perturbation curve P in [Figure 9](#). However, before analyzing the effect of perturbing along P to turn circles into pairs of isolated points, a discussion concerning the *nondegeneracy* of the space $R_{\pi}^{\natural}(Y, K)$ for appropriate perturbations is in order.

7.2 Nondegeneracy

In Floer-type theories, the chain complexes are generated by critical points of a functional defined on a configuration space, and the boundary operator is defined by counting integral curves for the gradient of the functional that connect the critical points (so-called “gradient flowlines”). The critical points of the functional form a moduli space. In order to have a finitely generated complex, then, one must ensure that the number of points in this space is finite. This can be achieved by the analogue of a slight perturbation of a real-valued function on a finite-dimensional manifold to ensure that its critical points are nondegenerate, and hence isolated (and finite in number if the manifold is compact). In the infinite-dimensional setting, we likewise must perturb the functional defining the Floer theory to achieve nondegeneracy of the Hessian. This can be viewed as a first step in constructing a Floer theory.

To ensure that differentials are defined, and that the resulting homology is well defined, however, requires more. Namely, we must have some form of transversality for the moduli spaces of gradient flow lines connecting critical points of the functional. Precise conditions depend on the context, and are usually subsumed under the terms

“regularity” or “transversality.” For example, in finite dimensions a function $f: M \rightarrow \mathbb{R}$ is nondegenerate if it is a Morse function, and regularity adds the requirement that the stable and unstable manifolds intersect transversally (which is usually referred to as the Morse–Smale condition). In Floer theory for Lagrangian intersections, nondegeneracy is typically achieved by a Hamiltonian isotopy of the Lagrangian submanifolds so that they intersect transversally. Regularity is achieved by perturbing the almost complex structure which defines the J -holomorphic curve equation satisfied by gradient flowlines in the space of paths connecting the two Lagrangians. In the context of instanton homology, nondegeneracy is expressed by the condition that the perturbed Chern–Simons functional is Morse in a suitable infinite-dimensional sense.

A Lagrangian–Floer theory in the context of traceless character varieties of knot complements in 3-manifolds has not yet been worked out and identified with singular instanton knot homology. Providing such an identification is an instance of the *Atiyah–Floer Conjecture*, which has been established in several other settings. In all known cases critical points of the Chern–Simons functional are nondegenerate (ie Morse) exactly when the corresponding Lagrangian intersections are transverse. This is proved by identifying the kernel of the Hessian of the Chern–Simons functional at a critical point with some form of de Rham cohomology, which is then related to transversality of the Lagrangian submanifolds via the de Rham theorem and the Mayer–Vietoris sequence.

To carry this out carefully below in our context would take us too far afield from the intent of this article. We offer instead the following as a notion of nondegeneracy in this article, which is adapted from Weil’s observation [40] that $H^1(\pi; \mathfrak{g})$ is identified with the tangent space at a smooth point of the character variety $\chi(\pi, G)$.

Given perturbation data π with the f_j real analytic, the space $R_\pi^{\mathfrak{h}}(Y, K)$ is the orbit space of the free $SU(2)/\pm 1 = SO(3)$ conjugation action on a real analytic variety. This variety is constructed as follows.

Consider a finite presentation $\langle G \mid R \rangle$ of $\pi_1(Y \setminus (K \cup H \cup W \cup P))$. The presentation defines a polynomial map $F_1: SU(2)^G \rightarrow SU(2)^R$ with preimage $F_1^{-1}(1, 1, \dots, 1)$ parameterizing all homomorphisms $\pi_1(Y \setminus (K \cup H \cup W \cup P)) \rightarrow SU(2)$, in the usual way.

Fixing three words in the generators representing the meridians μ_K, μ_H and μ_W , a map $F_2: SU(2)^G \rightarrow SU(2)$ is defined by sending a G -tuple to the image of μ_W . Then we have that $F_2^{-1}(-1)$ meets $F_1^{-1}(1, 1, \dots, 1)$ in those homomorphisms sending μ_K, μ_H to perpendicular elements of $C(\mathfrak{i})$ and μ_W to -1 . (We are assuming K is a knot, not a link. In the case when K has n components and H links K_1 , define

$F_2: \text{SU}(2)^G \rightarrow \text{SU}(2) \times \mathbb{R}^{n-1}$ by taking the last factors to be the value $\text{Re}(\mu_{K_i})$, $i = 2, \dots, n$.)

Finally, for each perturbation curve P_j , we will let $G_j: \text{SU}(2) \rightarrow \text{SU}(2)$ be the map $G_j(e^{\nu}Q) = e^{f_j(\nu)}Q$. The properties of f_j imply that G_j is well defined. The meridian and longitude p_j, ℓ_j of P_j , expressed as words in the generators G , give rise to a map $F_{3,j}: \text{SU}(2)^G \rightarrow \text{SU}(2)$ by $F_{3,j} = p_j^{-1}G_j(\ell_j)$. The preimage of 1 precisely captures the perturbation constraints (Equation (19)).

The product

$$F = F_2 \times F_1 \times \prod_j F_{3,j}: \text{SU}(2)^G \rightarrow \text{SU}(2) \times \text{SU}(2)^R \times \prod_j \text{SU}(2)$$

defines $R_\pi^{\natural}(Y, K)$ in the sense that

$$R_\pi^{\natural}(Y, K) = F^{-1}(1, 1, \dots, 1)/\text{SU}(2).$$

Thus $R_\pi^{\natural}(Y, K)$ is finite if and only if $F^{-1}(1, 1, \dots, 1)$ is a finite union of copies of $\text{SO}(3)$ with $\text{SU}(2)/\pm 1$ acting freely.

A representation ρ whose conjugacy class lies in $R_\pi^{\natural}(Y, K)$ determines a point (which we denote again by ρ) in $F^{-1}(1, 1, \dots, 1)$, by evaluating ρ on the elements of G . We call ρ *nondegenerate* if it is isolated in $R_\pi^{\natural}(Y, K)$ and the kernel of the differential dF_ρ is 3-dimensional, ie it is the tangent space to the orbit under the diagonal conjugation action of $\text{SU}(2)$ on $\text{SU}(2)^G$. (The fact that, for any G -tuple in the preimage, μ_K, μ_H are sent to noncommuting elements of $\text{SU}(2)$ implies that the stabilizer of the G -tuple is $\{\pm 1\}$ and the orbit is 3-dimensional.) We then call $R_\pi^{\natural}(Y, K)$ *nondegenerate* if it is finite and all its points are nondegenerate.

Standard arguments show that the notion of nondegeneracy is unchanged by changing the presentation. Indeed, any two finite have a common stabilization obtained by adding generators and relations to each, and stabilization preserves nondegeneracy.

It will be clear from our constructions below that as a space of representations, $R_\pi^{\natural}(Y, K)$ is a transverse intersection of two smooth 1-dimensional manifolds in a 2-manifold, in two different ways corresponding to the two different pillowcase pictures. One can show that transversality in these two contexts is equivalent, and equivalent to nondegeneracy as defined above, and we will explore this issue in subsequent work. In this sense we produce in Theorem 7.1 below (in concert with the results of [12]) a perturbation which makes $R_\pi^{\natural}(Y, K)$ nondegenerate.

A more sophisticated approach is to recast this in the context of group cohomology, which identifies the kernel of dF_ρ with certain 1-cocycles and the tangents

to the conjugation orbits with coboundaries, at least when the perturbation is trivial. Then nondegeneracy is equivalent to the vanishing of $H^1(Y \setminus (K \cup H); so(3)_\rho)$ (or, if K has n components, the vanishing of the kernel of $H^1(Y \setminus (K \cup H); so(3)_\rho) \rightarrow \bigoplus_{i=1}^n H^1(\mu_{K_i}; so(3)_\rho)$; see [26, Lemma 3.13] and also Gerard [11, Proposition 2.10]. For nontrivial perturbations similar conditions apply. The references show that nondegeneracy in this cohomological sense is equivalent to the perturbed Chern–Simons functional $CS + h_\pi$ having a nondegenerate Hessian at its critical points, and therefore its critical points serve as generators for the instanton Floer complex defining $I^{\natural}(Y, K)$.

To illustrate these ideas, we show that the distinguished representation $\alpha \in R^{\natural}(S^3, K)$, defined in Equation (11), is nondegenerate.

Decompose $S^3 \setminus (K \cup H \cup W)$ into $S^3 \setminus N(K)$ and $N(K) \setminus (K \cup H \cup W)$ along a torus $T = \partial N(K)$ as in Equation (13). Let μ_K, μ_H, μ_W denote the meridians of K, H, W and λ_K the longitude of K . Then

$$\pi_1(N(K) \setminus (K \cup H \cup W)) = \langle \mu_K, \mu_H, \mu_W, \lambda_K \mid \mu_W = [\mu_K, \mu_H], 1 = [\mu_K, \lambda_K] \rangle.$$

The representation α restricts to the unique abelian representation on $S^3 \setminus N(K)$ sending the meridian μ_K (which generates $H_1(S^3 \setminus N(K))$) to \mathbf{i} . The longitude λ_K is sent to 1 since it maps to zero in $H_1(S^3 \setminus N(K))$. Finally, α restricts to the nonabelian representation

$$\alpha(\mu_K) = \mathbf{i}, \quad \alpha(\mu_H) = \mathbf{j}, \quad \alpha(\mu_W) = -1, \quad \alpha(\lambda_K) = 1,$$

on $N(K) \setminus (K \cup H \cup W)$.

Hence (taking coefficients in $su(2)$ twisted by α),

$$H^0(S^3 \setminus N(K)) = \mathbb{R} = H^0(T), \quad H^0(N(K) \setminus (K \cup H \cup W)) = 0.$$

Since $\alpha(\mu_K) = e^{i\pi/2}$ satisfies $\Delta_K(\alpha(\mu_K)^2) \neq 0$, $H^1(S^3 \setminus N(K)) = \mathbb{R}$ (see [20]) and $H^1(T) = \mathbb{R}^2$.

A straightforward calculation yields $H^1(N(K) \setminus (K \cup H \cup W)) = \mathbb{R}^4$ and that the restriction to $H^1(T) = \mathbb{R}^2$ is surjective. Indeed, given any pair of unit quaternions (q_1, q_2) near (\mathbf{i}, \mathbf{j}) and a third unit quaternion q_3 in the unique circle subgroup through q_1 , the assignment

$$(20) \quad \mu_K \mapsto q_1, \quad \mu_H \mapsto q_2, \quad \mu_W \mapsto [q_1, q_2], \quad \lambda_K \mapsto q_3,$$

gives a smooth 7–dimensional family of irreducible representations near α on which conjugation acts freely modulo ± 1 , so that $H^1(N(K) \setminus (K \cup H \cup W))$ is 4–dimensional and maps onto $H^1(T)$.

The Mayer–Vietoris sequence then shows that $H^1(S^3 \setminus (K \cup H \cup W)) = \mathbb{R}^3$. Since $H^1(S^3 \setminus (K \cup H \cup W)) \cong \ker dF_1/B_1$, where B_1 denotes the tangent space to the 3–dimensional orbit through α of the conjugation action (see eg [40]), it follows that $\ker dF_1 \cong \mathbb{R}^6$. The map $SU(2) \times SU(2) \rightarrow SU(2)$ taking a pair to their commutator has -1 as a regular value, and hence the map F_2 is a submersion near α because the quaternions q_1, q_2 of Equation (20) can be chosen arbitrarily near \mathbf{i}, \mathbf{j} . This implies that the kernel of $d(F_1 \times F_2)$ at α is $6 - 3 = 3$ dimensional, so that α is nondegenerate. In particular, α remains nondegenerate under small perturbations.

7.3 Perturbation in a 3–ball

Place a loop P inside B^3 , linking A_2 and H as illustrated in Figure 9. Use the standard meridian-longitude framing of P to think of P as the image of an embedding of a solid torus $e: S^1 \times D^2 \rightarrow B^3$. Label the generators of $\pi_1(B^3 \setminus (A_1 \cup A_2 \cup H \cup W \cup P))$ by a, b, c, d, h, w , and p as indicated in the figure.




Figure 9: The local picture in the 3–ball where the perturbation occurs: the holonomy perturbation occurs in the neighborhood of the blue circle.

Fix a smooth function $f: \mathbb{R} \rightarrow \mathbb{R}$ satisfying $f(0) = 0, |f(x)| \leq 1, f(-x) = -f(x), f$ is 2π periodic and $f(x) \neq 0$ when x is not a multiple of π (it suffices for our purposes to take $f(x) = \sin(x)$). Note that $f(n\pi) = 0$ for integers n .

Fix an $\epsilon \geq 0$ and denote the data (P, f, ϵ) by π . Define the π –perturbed moduli space $R_\pi^h(B^3, A_1 \cup A_2)$ to be the space of conjugacy classes of $SU(2)$ representations which send a, b, c, d and h to $C(\mathbf{i}), w$ to -1 , and which satisfy the perturbation condition. Thus if the longitude $\lambda_P = bh$ of the component P is sent to $e^{\beta Q}$ and the meridian p to $e^{\nu Q}$ for some $Q \in C(\mathbf{i})$, then

$$\nu = \epsilon f(\beta).$$

In particular, when $\epsilon = 0$, $R_{\pi}^{\natural}(B^3, A_1 \cup A_2) = R^{\natural}(B^3, A_1 \cup A_2)$.

The following is our main result. Its statement is perhaps best understood by examining Figure 10.

Theorem 7.1 *For all $\epsilon > 0$ small enough, the space $R_{\pi}^{\natural}(B^3, A_1 \cup A_2)$ is homeomorphic to a circle, parameterized by $\beta \in \mathbb{R}/2\pi\mathbb{Z}$ by the assignment*

$$\begin{aligned} \rho(\beta): \quad a &\mapsto \mathbf{i}, & b &\mapsto e^{(\pi/2+\beta+\epsilon f(\beta))\mathbf{k}} \mathbf{i}, & c &\mapsto e^{(\pi/2+\beta-\epsilon f(\beta))\mathbf{k}} \mathbf{i}, \\ d &\mapsto e^{-2\epsilon f(\beta)\mathbf{k}} \mathbf{i}, & h &\mapsto -\mathbf{j} e^{-\epsilon f(\beta)\mathbf{k}}, & p &\mapsto e^{\epsilon f(\beta)\mathbf{k}}, & w &\mapsto -1. \end{aligned}$$

Proof A straightforward calculation using the Seifert–Van Kampen theorem shows that

$$\pi_1(B^3 \setminus (A_1 \cup A_2 \cup P \cup H \cup W))$$

is presented with generators a, b, c, d, w, h, p , and relations

$$(21) \quad c = \bar{p}bp, \quad d = \bar{c}ba, \quad [bh, p] = 1, \quad [a\bar{p}, h] = (ha)w(\bar{a}\bar{h}).$$

(The two commutator relations come from the tori which form the boundaries of the tubular neighborhoods of P and H .)

Let $R_{\pi}^{\natural}(B^3 \setminus P, A_1 \cup A_2)$ denote the space of conjugacy classes of $SU(2)$ representations which send a, b, c, d, h to $C(\mathbf{i})$ and w to -1 , with no restrictions on where p is sent.

Since a, b, c, d satisfy the same relation $d = \bar{c}ba$ in $\pi_1(B^3 \setminus (A_1 \cup A_2 \cup P \cup H \cup W))$ that is the defining relation in $\pi_1(S^2 \setminus \{a, b, c, d\})$, Proposition 3.1 implies that we may assume up to conjugation that any representation satisfies (abusing notation slightly to identify generators with their image in $SU(2)$)

$$a = \mathbf{i}, \quad b = e^{\gamma\mathbf{k}} \mathbf{i}, \quad c = e^{\theta\mathbf{k}} \mathbf{i}, \quad d = e^{(\theta-\gamma)\mathbf{k}} \mathbf{i}$$

for $(\gamma, \theta) \in [0, \pi] \times \mathbb{R}/(2\pi\mathbb{Z})$. It will be convenient to relax this condition and assume

$$(\gamma, \theta) \in \mathbb{R}/(2\pi\mathbb{Z}) \times \mathbb{R}/(2\pi\mathbb{Z}) = S^1 \times S^1$$

for most of this proof, and then determine which pairs of representations are equivalent at the end of the argument.

Denote by Q_H the image of h , and let $Q_P \in C(\mathbf{i})$ be a traceless unit quaternion so that $\lambda_P = bh$ is sent to $e^{\beta Q_P}$ for some $\beta \in [0, 2\pi)$. Since p commutes with λ_P , it follows that the representation sends p to $e^{\nu Q_P}$ for some angle ν . The perturbation condition implies that $\nu = \epsilon f(\beta)$, but we'll impose this condition last. For the moment, we

assume ϵ is small, so that $|\nu| < \frac{\pi}{2}$. Note that Q_P is determined up to ± 1 , unless λ_P is sent to ± 1 , in which case we may take Q_P to be any element of $C(\mathbf{i})$.

Summarizing what we have accomplished so far: any representation whose conjugacy class lies in $R_{\pi}^{\natural}(B^3, A_1 \cup A_2)$ can be conjugated so that

$$(22) \quad \begin{aligned} a &= \mathbf{i}, & b &= e^{\gamma \mathbf{k}} \mathbf{i}, & c &= e^{\theta \mathbf{k}} \mathbf{i}, & d &= e^{(\theta - \gamma) \mathbf{k}} \mathbf{i}, \\ h &= Q_H, & p &= e^{\nu Q_P}, & w &= -1, & \lambda_P &= e^{\beta Q_P}, \end{aligned}$$

for some 6-tuple

$$(23) \quad (\gamma, \theta, Q_H, Q_P, \beta, \nu) \in [0, 2\pi) \times [0, 2\pi) \times C(\mathbf{i}) \times C(\mathbf{i}) \times [0, 2\pi) \times (-\frac{\pi}{2}, \frac{\pi}{2}).$$

One obvious redundancy in this description is that we have (ν, β, Q_P) is equivalent to $(-\nu, \beta + \pi, -Q_P)$.

The relation $\bar{p}bp = c$ implies that

$$(24) \quad e^{-\nu Q_P} e^{\gamma \mathbf{k}} \mathbf{i} e^{\nu Q_P} = e^{\theta \mathbf{k}} \mathbf{i}.$$

Recall from Proposition 2.1 that the conjugation action of e^{tQ} on the 2-sphere $C(\mathbf{i})$ is rotation about the axis through $\pm Q$ of angle $2t$.

If $e^{\gamma \mathbf{k}} \neq e^{\theta \mathbf{k}}$, then Equation (24) implies that Q_P lies on the great circle in $C(\mathbf{i})$ through \mathbf{k} and $e^{(\gamma + \theta)/2 \mathbf{k}} \mathbf{i}$. This is because the only axes $\pm Q$ for which the orbits of rotation pass through both $e^{\gamma \mathbf{k}} \mathbf{i}$ and $e^{\theta \mathbf{k}} \mathbf{i}$ lie on this great circle.

If $e^{\gamma \mathbf{k}} = e^{\theta \mathbf{k}}$, then $e^{\nu Q_P}$ stabilizes $e^{\gamma \mathbf{k}} \mathbf{i}$. When $\nu \neq 0$, this implies that $Q_P = \pm e^{\gamma \mathbf{k}} \mathbf{i}$, so we have that either $\nu = 0$ (and Q_P is not constrained by this relation) or that $Q_P = \pm e^{\gamma \mathbf{k}} \mathbf{i} = \pm \sin(\frac{\pi}{2}) e^{(\gamma + \theta)/2 \mathbf{k}} \mathbf{i}$.

Hence, in any case,

$$(25) \quad \nu = 0 \text{ and } \gamma = \theta, \quad \text{or } Q_P = \cos t \mathbf{k} + \sin t e^{(\gamma + \theta)/2 \mathbf{k}} \mathbf{i}$$

for some $t \in [0, 2\pi)$.

The relation $h = \bar{b} \lambda_P$ implies that Q_H is determined by γ, β and Q_P by the equation

$$Q_H = -\mathbf{i} e^{-\gamma \mathbf{k}} e^{\beta Q_P}.$$

Since $\text{Re}(Q_H) = 0$, using Equation (25) we see that when $v \neq 0$,

$$\begin{aligned} 0 &= \text{Re}(Q_H) = \text{Re}(-i e^{-\gamma k} e^{\beta Q_P}) \\ &= \text{Re}(-i e^{-\gamma k} (\cos \beta + \sin \beta (\cos t k + \sin t e^{(\gamma+\theta)/2k} i))) \\ &= \sin \beta \text{Re}(-i e^{-\gamma k} (\cos t k + \sin t e^{(\gamma+\theta)/2k} i)) \\ &= \sin \beta \sin t \text{Re}(-i e^{-\gamma k} e^{(\gamma+\theta)/2k} i) \\ &= \sin \beta \sin t \cos\left(\frac{\theta-\gamma}{2}\right). \end{aligned}$$

Hence

$$(26) \quad 0 = \sin \beta \sin t \cos\left(\frac{\theta-\gamma}{2}\right)$$

Next, consider the relation $[a\bar{p}, h] = haw\bar{a}\bar{h} = -1$, or, in terms of our chosen coordinates,

$$[i e^{-v Q_P}, Q_H] = -1.$$

This equation can be rewritten, using the fact that if $Q \in C(i)$, then $Q^{-1} = -Q$, as

$$i e^{-v Q_P} Q_H (i e^{-v Q_P})^{-1} = -Q_H.$$

Part (iii) of Proposition 2.1 shows that $i e^{-v Q_P}$ is itself a traceless unit quaternion, and is perpendicular to Q_H . Hence

$$\begin{aligned} (27) \quad 0 &= \text{Re}(i e^{-v Q_P}) = -\sin v \text{Re}(i Q_P) \\ &= -\sin v \text{Re}(i (\cos t k + \sin t e^{(\gamma+\theta)/2k} i)) \\ &= \sin v \sin t \cos\left(\frac{\gamma+\theta}{2}\right) \end{aligned}$$

So far we have shown all $SU(2)$ representations of $\pi_1(B^3 \setminus (A_1 \cup A_2 \cup P \cup H \cup W))$ sending a, b, c, d, h into $C(i)$, and $w \mapsto -1$ can be conjugated to have the form in (22), for a 6-tuple in (23); furthermore, either $v = 0$ and $\gamma = \theta$, or we can express Q_P as

$$Q_P = \cos t k + \sin t e^{((\gamma+\theta)/2)k} i.$$

In the latter case, equations (26) and (27) must also hold.

Now assume that such a representation corresponds to the holonomy of a perturbed flat connection on $B^3 \setminus (A_1 \cup A_2 \cup H \cup W)$. Then there is the additional condition that $v = \epsilon f(\beta)$, where f is an odd, 2π -periodic, function, whose zeroes occur only at multiples of π . We will examine what this additional restriction implies about the representations in $R_\pi^{\natural}(B^3, A_1 \cup A_2)$.

To begin, note that $\nu = 0$ if and only if $\sin \beta = 0$. If $\nu = \sin \beta = 0$, then the representation is independent of $Q_P \in C(\mathbf{i})$; in particular, in this case, we can assume $Q_P = \pm \mathbf{k}$. We next examine the case that $\nu \neq 0$ and $\sin \beta \neq 0$. In this case, either

$$\sin t = 0, \quad \text{or} \quad \cos\left(\frac{\theta+\gamma}{2}\right) = 0, \quad \cos\left(\frac{\theta-\gamma}{2}\right) = 0.$$

(Note we are using the fact that $\nu = 0$ if and only if $\sin \nu = 0$, since $|\nu| < \frac{\pi}{2}$.)

Suppose $\cos\left(\frac{\theta+\gamma}{2}\right) = 0$ and $\cos\left(\frac{\theta-\gamma}{2}\right) = 0$. Since $\gamma \in [0, 2\pi)$ and $\theta \in [0, 2\pi)$, there are two solutions:

$$(\gamma, \theta) = (0, \pi) \quad \text{or} \quad (\pi, 0).$$

If (γ, θ) is equal to $(\pi, 0)$, then Equation (22) implies that $a = \mathbf{i}, b = -\mathbf{i}, c = \mathbf{i}$. If $(\gamma, \theta) = (0, \pi)$, then $a = \mathbf{i}, b = \mathbf{i}, c = -\mathbf{i}$. The relation $\bar{p}bp = c$ implies in either case that $e^{-\nu Q_P} \mathbf{i} e^{\nu Q_P} = -\mathbf{i}$. Proposition 2.1 then implies that $e^{-\nu Q_P}$ is a traceless unit quaternion, which is impossible since $|\nu| < \frac{\pi}{2}$. Thus one of $\cos\left(\frac{\theta+\gamma}{2}\right)$ and $\cos\left(\frac{\theta-\gamma}{2}\right)$ must be nonzero. It follows that if $\sin \beta \neq 0$, then $\sin t = 0$. In particular, either $\sin \beta \neq 0$ and $\sin t$ must equal zero, or $\sin \beta = 0$ and we can assume $\sin t = 0$ without changing the representation. Hence we may assume that $Q_P = \sigma \mathbf{k}$ for some $\sigma \in \{\pm 1\}$.

We've now seen that any representation whose conjugacy class lies in $R_{\pi}^{\natural}(B^3, A_1 \cup A_2)$ can be conjugated so that

$$(28) \quad \begin{aligned} a &= \mathbf{i}, & b &= e^{\gamma \mathbf{k}} \mathbf{i}, & c &= e^{\theta \mathbf{k}} \mathbf{i}, & d &= e^{(\theta-\gamma) \mathbf{k}} \mathbf{i}, \\ h &= -\mathbf{i} e^{(\beta\sigma-\gamma) \mathbf{k}}, & p &= e^{\epsilon f(\beta) \sigma \mathbf{k}}, & w &= -1, & \lambda_P &= e^{\beta \sigma \mathbf{k}}, \end{aligned}$$

for some 4-tuple

$$(29) \quad (\gamma, \theta, \beta, \sigma) \in [0, 2\pi) \times [0, 2\pi) \times [0, 2\pi) \times \{\pm 1\}.$$

Since $f(\beta + \pi) = -f(\beta)$, then we have that $(\gamma, \theta, \beta, \sigma)$ gives the same representation as $(\gamma, \theta, 2\pi - \beta, -\sigma)$ when $\beta > 0$. When $\beta = 0$, then $\beta = -\beta$. Hence we may assume that $\sigma = 1$, or more precisely, any representation whose conjugacy class lies in $R_{\pi}^{\natural}(B^3, A_1 \cup A_2)$ can be conjugated so that

$$(30) \quad \begin{aligned} a &= \mathbf{i}, & b &= e^{\gamma \mathbf{k}} \mathbf{i}, & c &= e^{\theta \mathbf{k}} \mathbf{i}, & d &= e^{(\theta-\gamma) \mathbf{k}} \mathbf{i}, \\ h &= -\mathbf{i} e^{(\beta-\gamma) \mathbf{k}}, & p &= e^{\epsilon f(\beta) \mathbf{k}}, & w &= -1, & \lambda_P &= e^{\beta \mathbf{k}}, \end{aligned}$$

where so far, the angles γ, θ, β could lie anywhere in $[0, 2\pi)$.

Finally, we determine what relations between the angles are necessary to satisfy the perturbed flat equation.

The relation $\bar{p}bp = c$ implies that $e^{-\epsilon f(\beta)\mathbf{k}} e^{\gamma\mathbf{k}} \mathbf{i} e^{\epsilon f(\beta)\mathbf{k}} = e^{\theta\mathbf{k}} \mathbf{i}$, which implies $e^{(\gamma-2\epsilon f(\beta))\mathbf{k}} = e^{\theta\mathbf{k}}$, so

$$(31) \quad \theta \equiv \gamma - 2\epsilon f(\beta) \pmod{2\pi}.$$

The relation $[a\bar{p}, h] = -1$ gives

$$[\mathbf{i} e^{-\epsilon f(\beta)\mathbf{k}}, -\mathbf{i} e^{(\beta-\gamma)\mathbf{k}}] = -1$$

and so

$$-1 = \mathbf{i} e^{-\epsilon f(\beta)\mathbf{k}} (-\mathbf{i}) e^{(\beta-\gamma)\mathbf{k}} e^{\epsilon f(\beta)\mathbf{k}} (-\mathbf{i}) e^{-(\beta-\gamma)\mathbf{k}} \mathbf{i} = e^{2(\epsilon f(\beta)+\beta-\gamma)\mathbf{k}}.$$

Hence

$$(32) \quad \gamma \equiv \beta + \epsilon f(\beta) + \frac{\pi}{2} \pmod{\pi}.$$

In other words, we either have

$$(33) \quad \begin{aligned} \gamma &\equiv \beta + \epsilon f(\beta) + \frac{\pi}{2} \pmod{2\pi}, \\ \theta &\equiv \beta - \epsilon f(\beta) + \frac{\pi}{2} \pmod{2\pi}, \end{aligned}$$

or else

$$(34) \quad \begin{aligned} \gamma &\equiv \beta + \epsilon f(\beta) + \frac{3\pi}{2} \pmod{2\pi}, \\ \theta &\equiv \beta - \epsilon f(\beta) + \frac{3\pi}{2} \pmod{2\pi}. \end{aligned}$$

These formulas give us two families of representations indexed by $\beta \in [0, 2\pi)$ whose union maps surjectively to $R_{\pi}^{\natural}(B^3, A_1 \cup A_2)$. These define parameterizations of two smooth circles:

$$(35) \quad \begin{array}{ll} \rho(\beta): a \mapsto \mathbf{i} & \rho'(\beta): a \mapsto \mathbf{i} \\ b \mapsto e^{(\beta+\epsilon f(\beta))\mathbf{k}} \mathbf{j} & b \mapsto -e^{(\beta+\epsilon f(\beta))\mathbf{k}} \mathbf{j} \\ c \mapsto e^{(\beta-\epsilon f(\beta))\mathbf{k}} \mathbf{j} & c \mapsto -e^{(\beta-\epsilon f(\beta))\mathbf{k}} \mathbf{j} \\ d \mapsto e^{-2\epsilon f(\beta)\mathbf{k}} \mathbf{i} & d \mapsto e^{-2\epsilon f(\beta)\mathbf{k}} \mathbf{i} \\ h \mapsto -\mathbf{j} e^{(-\epsilon f(\beta))\mathbf{k}} & h \mapsto \mathbf{j} e^{(-\epsilon f(\beta))\mathbf{k}} \\ p \mapsto e^{\epsilon f(\beta)\mathbf{k}} & p \mapsto e^{\epsilon f(\beta)\mathbf{k}} \\ w \mapsto -1 & w \mapsto -1 \end{array}$$

We next observe that the second circle of representations, ρ' , is simply a conjugate of a reparametrization of the first one, ρ . Indeed, from the odd symmetry of the function f , and the fact that conjugation by \mathbf{i} sends $\mathbf{j} \mapsto -\mathbf{j}$ and $\mathbf{k} \mapsto -\mathbf{k}$, a straightforward calculation using (35) shows that \mathbf{i} conjugates $\rho'(2\pi - \beta)$ to $\rho(\beta)$.

Finally, we show that in the circle ρ , distinct values $\beta_1, \beta_2 \in [0, 2\pi)$ never give conjugate representations (which includes *equal representations*). Suppose that conjugation by some $g \in \text{SU}(2)$ sends $\rho(\beta_1)$ to $\rho(\beta_2)$. Then conjugation by g fixes $\mathbf{i} = \rho(\beta_i)(a)$, so $g = e^{\tau \mathbf{i}}$ for some τ .

Considering the real part of the condition that $g(\rho(\beta_1)(p))g^{-1} = \rho(\beta_2)(p)$, we see that $f(\beta_1) = \pm f(\beta_2)$. The equation $f(\beta_1) = f(\beta_2) = 0$ only occurs when $\{\beta_1, \beta_2\} = \{0, \pi\}$, by our assumptions about the function f . Then consideration of the image of b shows $g \mathbf{j} g^{-1} = -\mathbf{j}$ (so $g = \pm \mathbf{i}$), but the image of h gives the contradictory condition that $g \mathbf{j} g^{-1} = \mathbf{j}$.

For any other pair of β values, $g(\rho(\beta_1)(p))g^{-1} = \rho(\beta_2)(p)$ is impossible unless $g \in \{\pm 1, \pm \mathbf{i}\}$. The cases $g = \pm 1$ imply $\rho(\beta_1) = \rho(\beta_2)$, which is easy to rule out by considering the images of p and b .

If $g = \pm \mathbf{i}$, then $\rho(\beta_2)(p) = g(\rho(\beta_1)(p))g^{-1}$ implies that $f(\beta_2) = -f(\beta_1)$. But then $\rho(\beta_2)(h) = g(\rho(\beta_1)(h))g^{-1}$ implies that $\epsilon f(\beta_2) = -\epsilon f(\beta_1) + \pi \pmod{2\pi}$, which is impossible given that $\epsilon|f(x)| < \frac{\pi}{2}$. This shows that $\rho(\beta)$, $\beta \in [0, 2\pi)$, represent distinct conjugacy classes of representations. \square




Figure 10: The restriction to the pillowcase of the circle of perturbed traceless representations of the 3-ball containing a trivial tangle: the black arc in the figure on the left is the image of the unperturbed traceless representations, and the brown circle is the image of the perturbed traceless representations.

Theorem 7.1 allows us to determine the image $R_{\pi}^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$, as well as the properties of the limit of $R_{\pi}^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ as

$\epsilon \rightarrow 0$. To emphasize the dependence on the parameter ϵ , we write

$$R_{(P,f,\epsilon)}^{\natural}(B^3, A_1 \cup A_2) \cong \{\rho^\epsilon(\beta) \mid \beta \in \mathbb{R}/2\pi\mathbb{Z}\},$$

where ρ^ϵ denotes the circle of representations of Equation (35).

Theorem 7.1 shows that the circle fibers of the restriction $R^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ have been perturbed away to pairs of points in the restriction $R_{\pi}^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$. We make this more precise in the following corollary.

Corollary 7.2 *Given perturbation data $\pi = (P, f, \epsilon)$ with P as in Figure 9, the restriction map to the pillowcase*

$$R_{(P,f,\epsilon)}^{\natural}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$$

is an immersion of the circle $\{\rho(\beta) \mid \beta \in \mathbb{R}/2\pi\mathbb{Z}\}$ for $\epsilon > 0$ small. Its image is the image of the curve in \mathbb{R}^2 , $\beta \mapsto (\beta + \epsilon f(\beta) + \frac{\pi}{2}, \beta - \epsilon f(\beta) + \frac{\pi}{2})$ under the canonical projection of \mathbb{R}^2 to the pillowcase.

The limit

$$\lim_{\epsilon \rightarrow 0} R_{(P,f,\epsilon)}^{\natural}(B^3, A_1 \cup A_2) \subset R^{\natural}(B^3, A_1 \cup A_2)$$

is a circle consisting of those representations which send h to $\pm \mathbf{j}$, ie those points with $\tau = \pm \frac{\pi}{2}$ in Proposition 6.2. Its projection to the pillowcase is a map from the circle onto the diagonal $\beta \mapsto (\beta + \frac{\pi}{2}, \beta + \frac{\pi}{2})$, which is a two-to-one immersion except at the corners when $\beta = \frac{\pi}{2}$ and $\frac{3\pi}{2}$.

Proof The statements all follow immediately from the formulas of Theorem 7.1 except possibly the fact that the circle $\{\rho^\epsilon(\beta)\}$ is immersed in the pillowcase when $\epsilon > 0$ is small enough.

This follows from the fact that the smooth embedding

$$\mathbb{R} \rightarrow \mathbb{R}^2, \quad \beta \mapsto (\beta + \epsilon f(\beta) + \frac{\pi}{2}, \beta - \epsilon f(\beta) + \frac{\pi}{2})$$

avoids the branch points $(\pi\mathbb{Z}) \times (\pi\mathbb{Z})$ in the branched cover $\mathbb{R}^2 \rightarrow R(S^2, \{a, b, c, d\})$ of the pillowcase.

To see why this is true, recall that the branch points are the points $\{(\pi k, \pi \ell) \mid k, \ell \in \mathbb{Z}\}$. Suppose $(\beta + \epsilon f(\beta) + \frac{\pi}{2}, \beta - \epsilon f(\beta) + \frac{\pi}{2}) = (\pi k, \pi \ell)$, then $\epsilon f(\beta) = \frac{\pi}{2}(k - \ell)$, and hence $k = \ell$ and $f(\beta) = 0$. But we also have $\beta + \frac{\pi}{2} = \frac{\pi}{2}(k + \ell)$, so that $\beta = \frac{\pi}{2}(2\ell - 1)$ and hence $f(\beta) \neq 0$, since f only vanishes for multiples of π . \square

Figure 10 illustrates the situation when $f(x) = \sin(x)$ and $\epsilon = 0.2$. The immersed brown circle (which by abuse of notation we also denote ρ) has one double point, corresponding to the parameter values $\beta = 0, \pi$. The black diagonal arc corresponds to the image of the unperturbed moduli space, each point in the interior of the black arc corresponding to a latitudinal circle in the 2–sphere $R^{\natural}(B^3, A_1 \cup A_2)$ and the endpoints corresponding to the poles. As ϵ shrinks towards 0, the circle approaches the black curve, with pairs of points collapsing to a single point along the interior of the arc. In the limit, the pair of distinct representations on each circle fiber over the black curve are the two unperturbed representations of Proposition 6.2 sending h to j and $-j$ and which project to the given point on the black arc in the pillowcase.

Notice that the restriction map $R^{\natural}_{\pi}(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{a, b, c, d\})$ is not an embedding and encloses zero signed area in the pillowcase. This can presumably be explained by the fact that the image has to lift to a Legendrian circle with respect to a natural connection in a Chern–Simons $U(1)$ bundle over $R(S^2, \{a, b, c, d\})$ as in the situation for closed surfaces; see Ramadas, Singer and Weitsman [35] and [12].

8 The intersection picture

The following corollary summarizes the results of Theorem 7.1 and Corollary 7.2 in a statement that suggests an intersection picture corresponding to Diagram (2). Perturbing along the unknotted curve P in a 3–ball corrects, in a manner which is independent of the pair (Y, K) , for the problem that the unperturbed Chern–Simons functional is never Morse.

Corollary 8.1 *Suppose that $K \subset Y$ is a knot and $B^3 \subset Y$ is a 3–ball intersecting K transversally in two trivial arcs $A = A_1 \cup A_2$. Let $(Y_0, K_0) := (Y \setminus B^3, K \setminus A)$.*

Assume that $R(Y_0, K_0)$ is a smooth 1–manifold away from a finite number of points and that the restriction to the pillowcase

$$R(Y_0, K_0) \rightarrow R(S^2, \{a, b, c, d\})$$

is an immersion transverse to the arc $\theta = \gamma$ on the manifold points, and takes the nonmanifold points outside a neighborhood of the arc $\theta = \gamma$. Then for a small enough choice of perturbation, the intersection in the pillowcase of $R(Y_0, K_0)$ and the circle ρ is transverse, and hence the set $R^{\natural}_{\pi}(Y, K)$ is nondegenerate (and finite). \square

In complete generality, achieving the conditions on $R(Y_0, K_0)$ will also involve suitable perturbations along curves in $Y_0 \setminus K_0$. One of our primary goals is to “combinatorialize” the instanton homology of a knot in terms of the intersection picture given

by Diagram (2) and Corollary 8.1. In later work we will explore the calculation of gradings, and explore differentials in terms of this picture. The reader should look at Figures 15 and 16 below for an illustration of Corollary 8.1.

9 The unreduced case

Kronheimer and Mrowka construct two versions of their singular instanton homology for (Y, K) . The first, *reduced* instanton homology $I^{\natural}(Y, K)$, corresponds to taking the connected sum K^{\natural} of K with a Hopf link and working with an $\text{SO}(3)$ bundle which is nontrivial on the torus which separates the components of the Hopf link. For this version of their theory, the critical set of the Chern–Simons functional is identified with $R^{\natural}(Y, K)$. For knots in S^3 with simple representation varieties (or for all knots, after applying a perturbation outside a neighborhood of K which contains H and W), $R^{\natural}(S^3, K)$ consists of one circle for each nonabelian point in the space $R(S^3, K)$ of traceless representations, and one nondegenerate isolated point corresponding to the abelian representation in $R(S^3, K)$. Theorem 7.1 then shows how to perturb along one curve P to turn each circle into a nondegenerate pair of isolated points.

The second version, *unreduced* instanton homology $I^{\sharp}(Y, K)$, corresponds to taking the disjoint union K^{\sharp} of K with a Hopf link $H_1 \cup H_2$ rather than the connected sum. The critical set of the corresponding Chern–Simons functional is $R^{\sharp}(Y, K) = R^{\natural}(Y, K \cup H_1)$. For knots in S^3 with simple representation varieties (or for all knots, after further perturbation), $R^{\sharp}(S^3, K)$ consists of a copy of $\text{SO}(3)$ for each nonabelian point in the space $R(S^3, K)$ of traceless representations, and one 2–sphere corresponding to the abelian representation in $R(S^3, K)$. This follows quickly from Lemma 4.2 by decomposing along the 2–sphere separating the Hopf link from K , by similar but easier versions of Propositions 4.3 and 4.4. In this section we prove a counterpart to Theorem 7.1 for $R^{\sharp}(Y, K)$ by using two perturbation curves.

Given a knot (or link) K in a 3–manifold Y , consider a 3–ball $B^3 \subset Y$ intersecting K in two unknotted arcs $A_1 \cup A_2$. Place the Hopf link H inside B^3 and place an arc W spanning the two components of the Hopf link, as shown by the black curves in Figure 11.

As Kronheimer and Mrowka observe in [26; 25], every conjugacy class of representations which take the meridians of K and H to traceless matrices and the meridian of W to -1 can be uniquely conjugated so that the two meridians of H are sent to i and j respectively. This shows that the corresponding unperturbed representation space $R^{\sharp}(Y, K)$ is homeomorphic to the space $\tilde{R}(Y, K)$ of all traceless representations of $\pi_1(Y \setminus K)$ (not conjugacy classes). Since the orbits of the $\text{SU}(2)$ conjugation

action on $\tilde{R}(Y, K)$ have dimension greater than 1, $R^\sharp(Y, K)$ is never finite. For knots in S^3 with simple representation varieties, $R^\sharp(Y, K)$ is a disjoint union of a 2–sphere corresponding to the distinguished representation α and a number of copies of $SO(3) = SU(2)/\pm 1$.

Place two smaller 3–balls B_1^3 and B_2^3 in the interior of B^3 as illustrated in Figure 11. Denote their boundaries by $S_i^2 = \partial B_i^3$. Place a perturbation curve P_1 inside B_1^3 and a second perturbation curve P_2 inside B_2^3 , as indicated. Label the various meridians a, b, c, d, m, n , and p_2 , as indicated.




Figure 11: The “ \sharp ” construction and two perturbation curves: the black Hopf link in the center and the arc connecting its components represents the additional data defining the \sharp construction relevant to unreduced singular instanton homology; the holonomy perturbation takes place in the neighborhood of the blue circles P_1 and P_2 .

Fix $\epsilon > 0$, and choose the perturbation function for the curve P_1 to be $\epsilon \sin(x)$. For the curve P_2 , take the perturbation function $2\epsilon \sin(x)$. With this perturbation data π in place, we obtain a space $R_\pi^\sharp(B^3, A_1 \cup A_2)$ and a restriction map to the pillowcase:

$$R_\pi^\sharp(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{m, b, c, n\})$$

Let $\arcsin: [-1, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$ denote the inverse sine function and define two smooth functions

$$(36) \quad \begin{aligned} \tau_1: S^1 &\rightarrow [-\frac{\pi}{6}, \frac{\pi}{6}], & \tau_1(\beta) &= \arcsin(-\frac{1}{2} \sin(\beta)), \\ \tau_2: S^1 &\rightarrow [\frac{5\pi}{6}, \frac{7\pi}{6}], & \tau_2(\beta) &= \pi - \tau_1(\beta). \end{aligned}$$

The set $\{(\beta, \tau) \in S^1 \times S^1 \mid \sin \beta = -2 \sin \tau\}$ is precisely the disjoint union of the graphs of $\tau_1(\beta)$ and $\tau_2(\beta)$.

Theorem 9.1 *The space $R_\pi^\sharp(B^3, A_1 \cup A_2)$ is the disjoint union of two circles, parameterized by $\beta \in \mathbb{R}/(2\pi\mathbb{Z})$ and $i = 1, 2$. These satisfy*

$$\begin{aligned} m &\mapsto \mathbf{i}, & b &\mapsto e^{-\tau_i(\beta)\mathbf{k}} e^{(\pi/2+\beta+\epsilon \sin \beta)\mathbf{k}} \mathbf{i}, & c &\mapsto e^{-\tau_i(\beta)\mathbf{k}} e^{(\pi/2+\beta-\epsilon \sin \beta)\mathbf{k}} \mathbf{i}, \\ n &\mapsto e^{-2\epsilon \sin(\beta)\mathbf{k}} \mathbf{i}, & a &\mapsto e^{-\tau_i(\beta)\mathbf{k}} \mathbf{i}, & d &\mapsto e^{(-\tau_i(\beta)-2\epsilon \sin(\beta))\mathbf{k}} \mathbf{i}. \end{aligned}$$

In particular the restriction map $R_\pi^\sharp(B^3, A_1 \cup A_2) \rightarrow R(S^2, \{m, b, c, n\})$ is given by

$$\rho_i(\beta) = (-\tau_i(\beta) + \frac{\pi}{2} + \beta + \epsilon \sin \beta, -\tau_i(\beta) + \frac{\pi}{2} + \beta - \epsilon \sin \beta), \quad i = 1, 2.$$

Proof Choose $\rho \in R_\pi^\sharp(B^3, A_1 \cup A_2)$. [Theorem 7.1](#) applied to the sphere S^2_1 implies that ρ may be uniquely conjugated to a representation satisfying

$$a \mapsto \mathbf{i}, \quad b \mapsto e^{(\pi/2+\beta+\epsilon \sin(\beta))\mathbf{k}} \mathbf{i}, \quad c \mapsto e^{(\pi/2+\beta-\epsilon \sin(\beta))\mathbf{k}} \mathbf{i}, \quad d \mapsto e^{-2\epsilon \sin(\beta)\mathbf{k}} \mathbf{i},$$

for some $\beta \in [0, 2\pi)$. [Proposition 3.1](#), applied to S^2 (or S^2_2), then implies that there exists a $\tau \in S^1$ so that

$$m \mapsto e^{\tau\mathbf{k}} \mathbf{i}, \quad n \mapsto e^{(\tau-2\epsilon \sin \beta)\mathbf{k}} \mathbf{i}.$$

The fundamental group $\pi_1(B^3_2 \setminus (K \cup H \cup P_2))$ is generated by a, d, p_2, m, n subject to the relations $d = \bar{p}_2 a p_2, n = \bar{p}_2 m p_2$, and $[\bar{a}m, p_2] = 1$. The longitude of P_2 is equal to $\bar{a}m$, which is sent to

$$-\mathbf{i} e^{\tau\mathbf{k}} \mathbf{i} = e^{-\tau\mathbf{k}}.$$

The perturbation condition for P_2 then says that the meridian p_2 is sent to $e^{-2\epsilon \sin(\tau)\mathbf{k}}$. The relation $d = \bar{p}_2 a p_2$ then implies

$$e^{-2\epsilon \sin(\beta)\mathbf{k}} \mathbf{i} = e^{2\epsilon \sin(\tau)\mathbf{k}} \mathbf{i} e^{-2\epsilon \sin(\tau)\mathbf{k}} = e^{4\epsilon \sin(\tau)\mathbf{k}} \mathbf{i}$$

so that (since ϵ is small) $\sin \beta = -2 \sin \tau$. The relation $n = \bar{p}_2 m p_2$ places the same restriction $\sin \beta = -2 \sin \tau$. Thus $\tau = \tau_1(\beta)$ or $\tau = \tau_2(\beta)$.

Conversely, given any τ satisfying $\sin \beta = -2 \sin \tau$, the assignment

$$(37) \quad \begin{aligned} a &\mapsto \mathbf{i}, & b &\mapsto e^{(\pi/2+\beta+\epsilon \sin(\beta))\mathbf{k}} \mathbf{i}, & c &\mapsto e^{(\pi/2+\beta-\epsilon \sin(\beta))\mathbf{k}} \mathbf{i}, \\ d &\mapsto e^{-2\epsilon \sin(\beta)\mathbf{k}} \mathbf{i}, & m &\mapsto e^{\tau\mathbf{k}} \mathbf{i}, & n &\mapsto e^{(\tau-2\epsilon \sin \beta)\mathbf{k}} \mathbf{i}, & p_2 &\mapsto e^{-2\epsilon \sin(\tau)\mathbf{k}}, \end{aligned}$$

uniquely defines a representation whose conjugacy class lies in $R_\pi^\sharp(B^3, A_1 \cup A_2 \cup H)$.

Conjugating by $e^{-\tau/2\mathbf{k}}$ completes the proof. □

Figure 12 illustrates the image of the two circles of Theorem 9.1, as well as the circle of Theorem 7.1. The red and green circles correspond to ρ_1 and ρ_2 , respectively, from Theorem 9.1. The brown circle corresponds to the circle ρ of Theorem 7.1 and Corollary 7.2. In this figure we used a moderately sized perturbation ($\epsilon = .4$) to highlight the fact that these three circles map to three distinct (but close) immersed circles in the pillowcase. It is straightforward to check that (just as in the case of ρ) for $i = 1, 2$, the map $\rho_i: S^1 \rightarrow R(S^2, \{m, b, c, n\})$ is an immersion with a single double point corresponding to $\rho_i(0) = \rho_i(\pi)$. As $\epsilon \rightarrow 0$, each circle limits to a generically two-to-one map onto the diagonal arc $\gamma = \theta$.




Figure 12: Illustrating Theorem 9.1: The theorem analyzes the perturbed traceless representations of the ball from Figure 11. There are two circles, ρ_1 and ρ_2 , of such representations, and the figure shows their restriction to the pillowcase. These are the red and green circles. Also shown (in brown) is the image of the circle of representations ρ , from Theorem 7.1.

In particular, Corollary 8.1 has the following unreduced counterpart.

Corollary 9.2 *Suppose that $K \subset Y$ is a knot and $B^3 \subset Y$ is a 3–ball intersecting K transversally in two trivial arcs $A = A_1 \cup A_2$.*

Assume that $R(Y_0, K_0) = R(Y \setminus B^3, K \setminus A)$ is a smooth 1–manifold away from a finite number of points and that the restriction to the pillowcase

$$R(Y_0, K_0) \rightarrow R(S^2, \{a, b, c, d\})$$

is an immersion transverse to the arc $\theta = \gamma$ on the manifold points, and takes the non-manifold points outside a neighborhood of the arc $\theta = \gamma$ (or that a suitable perturbation in $Y_0 \setminus K_0$ has been applied to achieve these conditions).

Then for a small enough choice of perturbation, the intersection in the pillowcase of $R(Y_0, K_0)$ and each circle ρ_1, ρ_2 is transverse, and hence the set $R_\pi^\sharp(Y, K)$ is nondegenerate (and finite), and contains two points for each point of $R_\pi^\sharp(Y, K)$. \square

In particular, for knots in the 3–sphere with simple representation varieties, the 2–sphere corresponding to the representation $\alpha_{\pi/2}$ is perturbed into a pair of isolated points and each $\text{SO}(3)$ component is perturbed into four isolated points. This is the unreduced counterpart to the fact that in the reduced case, the isolated representation α perturbs to a nondegenerate isolated point and each circle perturbs to a pair of isolated points.

For convenience, we will restrict our calculations in Section 12 below to the reduced instanton chain complex $CI^\sharp(Y, K)$. By using Theorem 9.1, each calculation has its unreduced counterpart, and in each case $CI^\sharp(Y, K)$ has twice as many generators as $CI^\natural(Y, K)$.

10 Examples: 2–bridge knots

Now that we have worked out the perturbation picture inside the 3–ball, completing the analysis of Diagram (2) is reduced to understanding the restriction map from $R(Y_0, K_0) = R(Y \setminus B^3, K \setminus (A_1 \cup A_2))$ to the pillowcase for various (Y, K) . We explore this in detail for 2–bridge knots and torus knots in this and the following section. In contrast to the analogous question for knot complements and the image of their full representation varieties in the pillowcase as the character variety of the torus, the situation is much simpler (in fact linear) for 2–bridge knots and complicated for torus knots.

Consider the arcs $A_1(n) \cup A_2(n) \subset B^3$ indicated in Figure 13, where n refers to the number of half-twists (positive or negative according to the sign of n). Orient these arcs arbitrarily and let a, b, c, d denote their oriented meridians as indicated.




Figure 13: A box labeled “ n ” indicates n positive or negative crossings, according to whether $n > 0$ or $n < 0$.

It is straightforward to check that if $\pi_1(B^3 \setminus (A_1(n) \cup A_2(n))) \rightarrow \text{SU}(2)$ is a representation given by

$$b \mapsto e^{xk} \mathbf{i}, \quad a \mapsto e^{yk} \mathbf{i}, \quad c \mapsto e^{zk} \mathbf{i}, \quad d \mapsto e^{wk} \mathbf{i},$$

then

$$(38) \quad \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1+n & -n \\ n & 1-n \end{pmatrix} \begin{pmatrix} z \\ w \end{pmatrix}.$$

Note that this formula holds with any choice of orientations of the arcs and choice of sign of n .

Let a_1, a_2, \dots, a_m be integers, with m odd, and define the rational number p/q with p and q relatively prime by the continued fraction expansion

$$(39) \quad \frac{p}{q} = a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}.$$

Consider the diagram of the 2–bridge knot $K = K(p/q)$ associated to this sequence (see Burde and Zieschang [6]), with a 3–ball intersecting in unknotted arcs as indicated in Figure 14. We assume p is odd so that $K(p/q)$ is a knot, not a link.

Our notational convention is that of Burde and Zieschang [6] and is consistent with the convention that the 2–fold branched cover of $K(p/q)$ is $L(p, q)$, where $L(p, q)$ is oriented as the quotient of S^3 , in other words $L(p, q)$ is $-p/q$ surgery on the unknot. Some other authors use other conventions.

The complement $S^3 \setminus (K \cup B^3)$ is homeomorphic to a 3–ball with two unknotted arcs removed, and hence Proposition 6.1 implies that $R(S^3 \setminus B^3, K \setminus (A_1 \cup A_2))$ is an arc

$$\tau_t: \pi_1(S^3 \setminus (K \cup B^3)) \rightarrow \text{SU}(2), \quad t \in [0, \pi]$$

determined by

$$\tau_t(a') = \mathbf{i}, \quad \tau_t(b') = \mathbf{i}, \quad \tau_t(c') = e^{t\mathbf{k}}\mathbf{i}, \quad \tau_t(d') = e^{t\mathbf{k}}\mathbf{i},$$

where a', b', c', d' are illustrated in Figure 14.

Then $\tau_t(a) = \mathbf{i}$, and writing

$$\tau_t(b) = e^{\gamma\mathbf{k}}\mathbf{i}, \quad \tau_t(c) = e^{\theta\mathbf{k}}\mathbf{i}, \quad \tau_t(d) = e^{(\theta-\gamma)\mathbf{k}}\mathbf{i},$$

Equation (38) implies that

$$(40) \quad \begin{pmatrix} \gamma(t) \\ \theta(t) \\ \theta(t) - \gamma(t) \end{pmatrix} = M(-a_1)N(a_2)M(-a_3)\cdots N(a_{m-1})M(-a_m) \begin{pmatrix} t \\ t \\ 0 \end{pmatrix} \pmod{2\pi},$$

where

$$M(a) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1+a & -a \\ 0 & a & 1-a \end{pmatrix}, \quad N(a) = \begin{pmatrix} 1+a & -a & 0 \\ a & 1-a & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$




Figure 14: A tangle decomposition for a 2-bridge knot

Thus the path τ_t will restrict to a linear path in the pillowcase of the form

$$(\gamma(t), \theta(t)) = (mt, nt), t \in [0, \pi]$$

for some integers m, n . More precisely, this linear path in \mathbb{R}^2 projects to a path in the pillowcase via the branched cover $\mathbb{R}^2 \rightarrow R(S^2, \{a, b, c, d\})$. We next show that $(m, n) = (q, q - p)$.

Lemma 10.1 *Let p, q be relatively prime nonzero integers and a_1, a_2, \dots, a_m a sequence giving the continued fraction expansion for p/q . Then the solution to Equation (40) is*

$$(\gamma(t), \theta(t)) = \pm(qt, (q - p)t).$$

Proof Note that

$$M(-a_m) \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 1 - a_m \\ -a_m \end{pmatrix}.$$

Suppose by induction that

$$M(-a_3)N(a_4) \cdots N(a_{m-1})M(-a_m) \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = \begin{pmatrix} s \\ s-r \\ -r \end{pmatrix}$$

for some relatively prime pair of integers r, s so that $\frac{s}{r}$ has continued fraction expansion given by a_3, a_4, \dots, a_m . Then

$$\begin{aligned} M(-a_1)N(a_2) \cdots M(-a_m) \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} &= M(-a_1)N(a_2) \begin{pmatrix} s \\ s-r \\ -r \end{pmatrix} \\ &= \begin{pmatrix} s+a_2r \\ s-r+a_2r-a_1s-a_1a_2r \\ -a_1s-a_1a_2r-r \end{pmatrix}. \end{aligned}$$

Setting $p_0 = a_1s + a_1a_2r + r$ and $q_0 = s + a_2r$ it is easy to check that p_0 and q_0 are relatively prime and

$$\frac{p_0}{q_0} = a_1 + \frac{1}{a_2 + \frac{s}{r}}.$$

In particular, p_0/q_0 has continued fraction expansion given by a_1, \dots, a_m so that $(p_0, q_0) = \pm(p, q)$, and

$$M(-a_1)N(a_2) \cdots M(-a_m) \begin{pmatrix} t \\ t \\ 0 \end{pmatrix} = \pm t \begin{pmatrix} q \\ q-p \\ -p \end{pmatrix}. \quad \square$$

The two paths

$$t \mapsto (qt, (q-p)t), \quad t \mapsto (-qt, -(q-p)t), \quad t \in [0, \pi],$$

are identical as maps to the pillowcase $R(S^2, \{a, b, c, d\}) = \mathbb{R}^2 / \sim$. Hence the sign ambiguity in Lemma 10.1 does not affect the image in the pillowcase.

The intersection of the curve $(\gamma(t), \theta(t)) = (qt, (q-p)t)$, $t \in [0, \pi]$ with the curve $\gamma = \theta$ in $R(S^2, \{a, b, c, d\})$ occurs at the $\frac{p+1}{2}$ points

$$(41) \quad x_\ell = (q \frac{2\pi\ell}{p}, (q-p) \frac{2\pi\ell}{p}), \quad \ell = 0, 1, \dots, \frac{p-1}{2}.$$

Note that the intersection point x_0 corresponds to the distinguished representation α of Equation (11).

Combining this observation with Proposition 6.2 and Lemma 4.2 one immediately concludes with the following.

Theorem 10.2 For the 2-bridge knot $K = K(p/q)$, the space $R^{\sharp}(S^3, K)$ is a union of circles and one isolated point α , one circle for each intersection point x_{ℓ} , $\ell = 1, 2, \dots, \frac{p-1}{2}$ with α corresponding to x_0 .

For perturbation data π as above, the space $R_{\pi}^{\sharp}(S^3, K)$ is a union of pairs of isolated nondegenerate points $x_{\ell,1}, x_{\ell,2}$, $\ell = 1, 2, \dots, \frac{p-1}{2}$, and one additional nondegenerate point α' , corresponding to the intersections in $R(S^2, \{a, b, c, d\}) = [0, \pi] \times [0, 2\pi] / \sim$ of the curve $(qt, (q-p)t)$, $t \in [0, \pi]$ with the circle $\{\rho^{\epsilon}(\beta)\}$.

Proof The first assertion follows from Proposition 6.2 and Lemma 4.2. The second then follows similarly from Corollary 7.2 and Lemma 4.2. \square

Note that if we consider the space $\tilde{R}(S^3, K)$ of traceless $SU(2)$ representations of K (not modulo conjugation) then the same reasoning gives that for $K = K(p/q)$, $\tilde{R}(S^3, K)$ is homeomorphic to the union of a 2-sphere (the conjugacy class of α) and $\frac{p-1}{2}$ copies of $SO(3) = SU(2)/\pm 1$ (the conjugacy classes of each x_{ℓ}), giving a different argument for the result of Lewallen [28, Theorem 2.4]. Since the spaces $\tilde{R}(S^3, K)$ and $R^{\sharp}(S^3, K)$ are homeomorphic, Theorem 9.1 and Corollary 9.2 implies that after perturbing along the two curves P_1, P_2 , $R_{\pi}^{\sharp}(S^3, K)$ is the union of 4ℓ points $x_{\ell,1}, x_{\ell,2}, x_{\ell,3}, x_{\ell,4}$, $\ell = 1, 2, \dots, \frac{p-1}{2}$ and two additional points α'_1, α'_2 .




Figure 15: R^{\sharp} and R_{π}^{\sharp} for the trefoil




Figure 16: R^{\sharp} and R_{π}^{\sharp} for the figure 8 knot

We illustrate [Theorem 10.2](#) in a few examples. Consider first the $(2, n)$ torus knot $T_{2,n}$, corresponding to $p/q = -n/1$. Thus the restriction of $(\gamma(t), \theta(t))$ to the pillowcase is parameterized by

$$t \in [0, \pi] \mapsto (t, (n + 1)t) \in [0, \pi] \times [0, 2\pi].$$

[Figure 15](#) illustrates the case of the right handed trefoil knot, $T_{2,3} = K(-3/1)$, with corresponding curve $t \mapsto (t, 4t)$. The unperturbed moduli space $R^{\natural}(S^3, T_{2,3})$ consists of a isolated point α and a circle c . The perturbed moduli space $R^{\natural}_{\pi}(S^3, T_{2,3})$ with $\pi = (P, \sin(x), \epsilon)$, consists of three isolated representations, α', x_1, x_2 . As $\epsilon \rightarrow 0$, $\alpha' \rightarrow \alpha$ and x_1, x_2 converge to a pair of antipodal points on the circle c . These three points are the generators of the instanton knot homology chain complex of the trefoil.

As a further example [Figure 16](#) illustrates the situation for K the figure 8 knot, which can be expressed as $K(-5/3)$. This gives the curve $t \mapsto (3t, 8t)$. We conclude that the unperturbed space $R^{\natural}(S^3, K)$ consists of a pair of circles c_x, c_y and the isolated point α . The perturbed moduli space (generating the instanton chain complex) $R^{\natural}_{\pi}(S^3, K)$ contains five points, $\alpha', x_1, x_2, y_1, y_2$.




Figure 17: R^{\natural}_{π} for the knot 7_2

It is worth recalling that the 2-fold branched cover of $K(p/q)$ is $L(p, q)$, and hence the fraction p/q does not uniquely determine $K = K(p/q)$. For example the figure 8 knot can also be described as $K(5/2)$, and hence its character variety $R(S^3, K)$ can also be described as the intersection of $(\gamma, \theta) = (2t, -3t)$ with $\{\gamma = \theta\}$. This corresponds to the fact that one can choose different (up to isotopy) 3-balls intersecting K in two unknotted arcs, for the same reason that 3-manifolds admit different Heegaard splittings.

Finally, Figure 17 illustrates the case when $K = 7_2 = K(-11/5)$. The restriction $R(S^3 \setminus B^3, K \setminus (A_1 \cup A_2)) \rightarrow R(S^2, \{a, b, c, d\})$ takes the arc to $(\gamma(t), \theta(t)) = (5t, 16t)$. The perturbed moduli space $R_{\pi}^h(S^3, K)$ contains 11 points,

$$\alpha', \quad x_1, \quad x_2, \quad y_1, \quad y_2, \quad z_1, \quad z_2, \quad w_1, \quad w_2, \quad v_1, \quad v_2.$$

11 Examples: Torus knots

In contrast to the situation for the full character varieties $\chi(S^3, T_{p,q})$, the varieties of traceless representations $R(S^3, T_{p,q})$ for torus knots are more complicated than those for 2-bridge knots. We explore the situation in enough detail to establish that a 3-ball intersecting a torus knot in 2 unknotted arcs can be found so that the restriction to the pillowcase of any nonabelian representation in $R(S^3, T_{p,q})$ avoids the corners. We then give a method to describe a 2-variable polynomial which cuts out the traceless character variety of the complement of a 3-ball meeting a torus knot in two arcs.

Note that counting the points of $R(S^3, T_{p,q})$ can be done by looking at the intersections of $\chi(S^3, T_{p,q})$ with the circle $S(i)$ of Equation (5), or equivalently by computing the signature of a torus knot. We establish that perturbing in a ball using Theorem 7.1 gives rise to a set of $\sigma(T_{p,q}) + 1$ generators for the instanton complex $CI^h(S^3, T_{p,q})$.

Figure 18 illustrates a (p, q) torus knot $T_{p,q}$ in S^3 . We view S^3 as $\frac{q}{r}$ and $-\frac{s}{p}$ Dehn surgery on the two components of a Hopf link, where $pr + qs = 1$. The knot $T_{p,q}$ is isotopic to a curve parallel to the first component, a fact which can be verified by identifying the parallel curve with a regular fiber in a Seifert fibering of S^3 with singular fibers of order p and q . In the figure, $T_{p,q}$ has been isotoped so that it meets a 3-ball in a pair of trivial arcs $A_1 \cup A_2$.

We wish to identify the space

$$R(Y_0, K_0) := R(S^3 \setminus B^3, T_{p,q} \setminus (A_1 \cup A_2))$$

and its image in the pillowcase. Note that

$$(S^3 \setminus B^3) \setminus (T_{p,q} \setminus (A_1 \cup A_2)) = S^3 \setminus (B^3 \cup T_{p,q}).$$




Figure 18: A tangle decomposition for the (p, q) torus knot

A straightforward calculation using the Wirtinger presentation shows $\pi_1(S^3 \setminus (B^3 \cup T_{p,q}))$ has the presentation

$$\langle x, y, a, b, c, d \mid c = \bar{x}ax, ad\bar{a} = yxb\bar{x}\bar{y}, \\ 1 = [y, xb] = [x, d\bar{a}y] = y^s(xb)^{-p} = x^q(d\bar{a}y)^r \rangle$$

with x, y, a, b, c, d the generators illustrated in Figure 18.

A priori, it is clear that $\pi_1(S^3 \setminus (B^3 \cup T_{p,q}))$ is a free group on two generators: up to homotopy $S^3 \setminus (B^3 \cup T_{p,q})$ is obtained by gluing two solid tori along a disk, just as a torus knot complement is obtained by gluing two solid tori along an annulus. Word manipulation in the presentation above, however, provides an explicit identification with a free group on generators

$$A = (xb)^q y^r, \quad B = (d\bar{a}y)^{-s} x^p.$$

Manipulating words shows that

$$A^s = xb, \quad A^p = y, \quad B^r = x, \quad B^{-q} = d\bar{a}y.$$

Solving for a, b, c, d (using the relation $d\bar{a} = \bar{c}b$) yields

$$(42) \quad \begin{aligned} a &= A^{s+p} B^{q-r}, & b &= B^{-r} A^s, & c &= B^{-r} A^{s+p} B^q = B^{-r} a B^r, \\ d &= B^{-q} A^s B^{q-r} = B^{-(q-r)} b B^{q-r}. \end{aligned}$$

Since $\pi_1(S^3 \setminus (B^3 \cup T_{p,q}))$ is free on A and B , the assignment $A \mapsto M, B \mapsto N$ gives a representation for any pair $M, N \in \text{SU}(2)$. A general such assignment will not yield a traceless representation. However, writing the pair (M, N) as

$$M = e^{uQ}, N = e^{vR}, \quad Q, R \in C(\mathbf{i}), \quad u, v \in [0, \pi],$$

then the corresponding representation determines a point in $R(Y_0, K_0)$ if and only if the images of a and b

$$(43) \quad M^{s+p} N^{q-r}, \quad N^{-r} M^s,$$

are traceless, or, equivalently, if and only if

$$(44) \quad 0 = \text{Re}(e^{(s+p)uQ} e^{(q-r)vR}), \quad 0 = \text{Re}(e^{-rvR} e^{suQ}).$$

Define

$$V_{p,q,r,s} = \{(M, N) \in \text{SU}(2) \times \text{SU}(2) \mid M^{s+p} N^{q-r} = \mathbf{i} \text{ and } N^{-r} M^s = e^{\gamma \mathbf{k}} \mathbf{i}, \gamma \in [0, \pi]\}.$$

An analysis using Equation (43) leads to the following result.

Theorem 11.1 *The assignment $A \mapsto M, B \mapsto N$ induces a homeomorphism*

$$V_{p,q,r,s} \cong R(S^3 \setminus B^3, T_{p,q} \setminus (A_1 \cup A_2)) = R(Y_0, K_0).$$

For $(M, N) \in V_{p,q,r,s}$, define $\gamma \in [0, \pi]$ and $\theta \in [0, 2\pi)$ by

$$N^{-r} M^s = e^{\gamma \mathbf{k}} \mathbf{i}, \quad N^{-r} M^{s+p} N^q = e^{\theta \mathbf{k}} \mathbf{i}.$$

Then $e^{(\gamma-\theta)\mathbf{k}} = M^p N^q$ and the space $R(S^3, T_{p,q})$ is homeomorphic to the subset of $V_{p,q,r,s}$ consisting of those pairs (M, N) satisfying $M^p N^q = 1$.

The restriction to the pillowcase $R(S^3, T_{p,q}) \rightarrow R(S^2, \{a, b, c, d\})$ takes every non-abelian representation to a nonabelian representation.

Finally, for suitable small perturbations, $R_\pi^{\natural}(S^3, T_{p,q})$ contains $|\sigma(T_{p,q})| + 1$ nondegenerate points.

Proof Let I denote the interval of traceless unit quaternions of the form $e^{\gamma \mathbf{k}} \mathbf{i}$ with $\gamma \in [0, \pi]$ and define $F: \text{SU}(2) \times \text{SU}(2) \rightarrow \text{SU}(2) \times \text{SU}(2)$ by the formula $F(M, N) = (M^{s+p} N^{q-r}, N^{-r} M^s)$ so that $V_{p,q,r,s} = F^{-1}(\{\mathbf{i}\} \times I)$.

The pair $(M, N) \in V_{p,q,r,s}$ determines a representation which sends a to \mathbf{i} and b to a unit quaternion, say $e^{\gamma\mathbf{k}}\mathbf{i}$ in I . Then c is sent to $N^{-r}aN^r$ and d is sent to $N^{q-r}bN^{q-r}$, so that c and d are sent to the conjugacy class $C(\mathbf{i})$ of traceless unit quaternions. This shows $V_{p,q,r,s}$ maps into $R(Y_0, K_0)$.

The map is surjective since any representation in $R(Y_0, K_0)$ can be conjugated to send a to \mathbf{i} and b into I . If A and B are sent to noncommuting quaternions then the resulting conjugacy class is uniquely determined.

If A, B are sent to commuting elements M, N , then $(M^{s+p}N^{q-r}, N^{-r}M^s) = (\mathbf{i}, \pm\mathbf{i})$, and hence M and N lie in the circle $\{e^{\mu\mathbf{i}}\}$, say $M = e^{\mu_1\mathbf{i}}, N = e^{\mu_2\mathbf{i}}$. Therefore

$$(\mathbf{i}, \pm\mathbf{i}) = (e^{((s+p)\mu_1+(q+r)\mu_2)\mathbf{i}}, e^{(s\mu_1-r\mu_2)\mathbf{i}})$$

so that $((s+p)\mu_1+(q+r)\mu_2) = \frac{\pi}{2}$ and $s\mu_1-r\mu_2 = \pm\frac{\pi}{2}$ modulo 2π . Since $pr+qs=1$, these equations uniquely determine μ_1 and μ_2 modulo 2π , and hence M and N are again uniquely determined, and so the map is a homeomorphism.

One computes

$$e^{(\gamma-\theta)\mathbf{k}} = -e^{\gamma\mathbf{k}}\mathbf{i}e^{\theta\mathbf{k}}\mathbf{i} = (N^{-r}M^s)^{-1}N^{-r}M^{s+p}N^q = M^pN^q.$$

The space $R(S^3, T_{p,q})$ corresponds to those representations in $R(Y_0, K_0)$ satisfying $a = d$ and $b = c$. From Equation (42) we see that if A, B are sent to M, N , then $a = d$ and $b = c$ precisely when $M^pN^q = 1$.

Suppose a representation in $R(S^3, T_{p,q})$ restricts to an abelian representation in the pillowcase $R(S^2, \{a, b, c, d\})$. The representation is given by a pair $(M, N) \in V_{p,q,r,s}$ satisfying $M^pN^q = 1$, and, since its restriction is abelian, Equation (42) gives

$$(45) \quad \begin{aligned} 2\mathbf{i} &= M^{s+p}N^{q-r}, & \sigma\mathbf{i} &= N^{-r}M^s, \\ \sigma\mathbf{i} &= N^{-r}\mathbf{i}N^r, & \mathbf{i} &= N^{-(q-r)}(\sigma\mathbf{i})N^{q-r} \end{aligned}$$

for some choice of sign $\sigma = \pm 1$.

Hence

$$N^{-1}\mathbf{i}N = N^{-r p - q s}\mathbf{i}N^{r p + q s} = (N^{-r})^{p+s}(N^{-(q-r)})^s\mathbf{i}(N^{(q-r)})^s(N^r)^{p+s} = \sigma^{2s+p}\mathbf{i}.$$

When $\sigma^p = 1$, $N = e^{\mu\mathbf{i}}$ for some μ . Then $M = M^{pr+qs} = N^{-qr}(N^r\sigma\mathbf{i})^q$ and so M and N commute, so that the representation is abelian.

We will show that $\sigma^p = -1$ is impossible. Suppose to the contrary that $\sigma^p = -1$, so that $\sigma = -1$ and p is odd. The equation $N^{-1}\mathbf{i}N = -\mathbf{i}$ implies that N is a traceless

unit quaternion, and hence $N^2 = -1$. Then $-i = N^{-r} i N^r = N^{-(q-r)} i N^{q-r}$ imply that r is odd and q is even. Since $M^p = N^{-q}$, it follows that $M^{2p} = 1$.

From Equation (45) we obtain

$$\begin{aligned} 1 = (-i)i &= N^{-r} M^s M^{s+p} N^{q-r} = N^{-r} M^{2s} M^p N^q N^{-r} \\ &= N^{-r} M^{2s} N^{-r} = -N^{-r} M^{2s} N^r \end{aligned}$$

so that $M^{2s} = -1$. But then $M^2 = M^{2(pr+qs)} = M^{2qs} = (-1)^q = 1$, a contradiction. Hence $\sigma^p \neq -1$.

It follows from [13; 14] that $R(S^3, T_{p,q})$ contains $|\sigma(T_{p,q})|/2$ nonabelian representations and one abelian representation. Since the restriction of any nonabelian representation to the pillowcase avoids the corners, it corresponds to an intersection point of the image $V_{p,q,r,s} \rightarrow R(S^2, \{a, b, c, d\})$ with the interior of the arc $\gamma = \theta$. Applying Propositions 4.3, 4.4 and Theorem 7.1 we see that each nonabelian representation in $R(S^3, T_{p,q})$ gives rise to two nondegenerate points of $R^1(S^3, T_{p,q})$ and the representation α gives one more nondegenerate point. \square

Theorem 11.1 does not give an explicit description of $R(Y_0, K_0) \cong V_{p,q,r,s}$ or its image in the pillowcase in the same sense as for 2-bridge knots above. The space $V_{p,q,r,s}$ is complicated; in fact the map $F: \text{SU}(2) \times \text{SU}(2) \rightarrow \text{SU}(2) \times \text{SU}(2)$ is not transverse to the interval $\{i\} \times I$; we will see an example below where $V_{p,q,r,s}$ is a singular variety.

We give a more explicit description of $V_{p,q,r,s}$ in two different ways in Proposition 11.2 and Theorem 11.3 below. In Proposition 11.2 we prove that, for any p, q, r , and s , $V_{p,q,r,s}$ contains an arc which maps to a straight line segment in the pillowcase. In Theorem 11.3 we show that $V_{p,q,r,s}$ is a semialgebraic set in \mathbb{R}^2 contained in the zero set of a polynomial determined explicitly by the integers p, q, r, s . The method of Theorem 11.3 lends itself easily to computer calculation and makes it easy describe $V_{p,q,r,s}$ in particular examples.

Write I for $\{i\} \times I$, so that $V_{p,q,r,s} = F^{-1}(I)$. The following result shows how to construct a splitting of $V_{p,q,r,s} \rightarrow I$, yielding a curve in $V_{p,q,r,s}$ whose image in the pillowcase is a straight line.

Proposition 11.2 *There is a cross section $s: I \rightarrow V_{p,q,r,s}$ of the map $F: V_{p,q,r,s} \rightarrow I$ whose image consists of binary dihedral representations. The image of $s(I)$ in the pillowcase $R(S^2, \{a, b, c, d\})$ is a straight line segment.*

If p and q are odd, or if p is even and $q-2r = \pm 1$, then the initial point $s(i) = s(e^{0k} i)$ is the restriction to $(S^3 \setminus B^3, T_{p,q} \setminus (A_1 \cup A_2))$ of the unique abelian traceless representation of $\pi_1(S^3 \setminus T_{p,q})$.

Proof Assume that q is odd by interchanging p and q if necessary. Fix $\gamma \in [0, \pi]$. We construct $s(e^{\gamma\mathbf{k}}\mathbf{i})$. We consider three cases:

- (1) p, r both odd
- (2) p odd, r even
- (3) p even

For the first case, p, q, r , and $p + s$ are odd, and s and $q - r$ are even. Set

$$s(e^{\gamma\mathbf{k}}\mathbf{i}) = ((-1)^{(p+s+q-r-1)/2}\mathbf{i}, (-1)^{(s-r-1)/2}e^{\gamma\mathbf{k}}\mathbf{i}).$$

For the second case, p, q, s , and $q - r$ are odd, and $r, p + s$ are even. Set

$$s(e^{\gamma\mathbf{k}}\mathbf{i}) = ((-1)^{(r-s+1)/2}e^{\gamma\mathbf{k}}\mathbf{i}, (-1)^{(p+s+q-r-1)/2}\mathbf{i}).$$

For the third case, q, s , and $p + s$ are odd. Set

$$s(e^{\gamma\mathbf{k}}\mathbf{i}) = (e^{\tau\mathbf{k}}\mathbf{i}, e^{\psi\mathbf{k}}),$$

where

$$\tau = \frac{q-r}{q-2r}\gamma + \frac{\pi(rp-qs+2rs+q-2r)}{2(q-2r)}, \quad \psi = \frac{1}{q-2r}\gamma + \frac{\pi p}{2(q-2r)}.$$

A calculation shows that in each of the three cases, $F(s(e^{\gamma\mathbf{k}}\mathbf{i})) = (\mathbf{i}, e^{\gamma\mathbf{k}}\mathbf{i})$. Note that in each case the generators are sent to the binary dihedral subgroup $\{e^{\theta\mathbf{k}}\} \cup \{e^{\theta\mathbf{k}}\mathbf{i}\}$ of $SU(2)$.

Denote $s(e^{\gamma\mathbf{k}}\mathbf{i})$ by $(M(\gamma), N(\gamma))$. Then $M(0)$ and $N(0)$ commute so that the corresponding representation is abelian. Moreover, when p and q are both odd, or if p is even and $q - 2r = \pm 1$, $M(0)^p N(0)^q = 1$, so that this abelian representation extends to a traceless abelian representation of $\pi_1(S^3 \setminus T_{p,q})$.

Theorem 11.1 implies that $e^{(\gamma-\theta)\mathbf{k}} = M^p(\gamma)N^q(\gamma)$. For the three cases, we have

$$e^{(\gamma-\theta)\mathbf{k}} = M^p(\gamma)N^q(\gamma) = \begin{cases} e^{-\gamma\mathbf{k}} & \text{case (1),} \\ e^{\gamma\mathbf{k}} & \text{case (2),} \\ e^{q/(q-2r)\gamma\mathbf{k}} e^{(\pi p(2q-2r))/(2(q-2r))\mathbf{k}} & \text{case (3),} \end{cases}$$

so that in each case θ is a linear function of $\gamma \in [0, \pi]$. □

We turn now to a different description of $R(Y_0, K_0) \cong V_{p,q,r,s}$ in terms of Chebyshev polynomials. **Theorem 11.3** below roughly says that if $(M, N) = (e^{uQ}, e^{vR}) \in V_{p,q,r,s}$, then there exists a polynomial $p(x, y)$ so that $p(\cos u, \cos v) = 0$, and that conversely, the zero set of this polynomial, subject to some inequalities (essentially $|x|, |y| \leq 1$), parameterizes $V_{p,q,r,s}$.

First, we return to Equation (44). Using Proposition 2.1 we may rewrite this as

$$(46) \quad \begin{aligned} 0 &= \cos((s+p)u) \cos((q-r)v) - \sin((s+p)u) \sin((q-r)v) Q \cdot R, \\ 0 &= \cos(su) \cos(-rv) - \sin(su) \sin(-rv) Q \cdot R, \end{aligned}$$

where $Q \cdot R$ denotes the dot product of Q and R .

Thus, if we define $\tilde{V}_{p,q,r,s}$ to be the set of pairs $(M, N) = (e^{uQ}, e^{vR}) \in \text{SU}(2) \times \text{SU}(2)$ satisfying (46), then $\tilde{V}_{p,q,r,s}$ is in bijective correspondence with all representations (not conjugacy classes) of traceless representations. Note that $V_{p,q,r,s} \subset \tilde{V}_{p,q,r,s}$.

For each integer n , there exist (Chebyshev) polynomials $T_n(x)$ and $S_n(x)$ so that

$$(47) \quad \cos(nu) = T_n(\cos u), \quad \sin(nu) = \sin u S_n(\cos u).$$

Hence Equation (46) can be rewritten as

$$(48) \quad \begin{aligned} 0 &= T_{s+p}(\cos u) T_{q-r}(\cos v) - \sin(u) \sin(v) S_{s+p}(\cos u) S_{q-r}(\cos v) Q \cdot R, \\ 0 &= T_s(\cos u) T_{-r}(\cos v) - \sin(u) \sin(v) S_s(\cos u) S_{-r}(\cos v) Q \cdot R. \end{aligned}$$

Substituting $x = \cos u, y = \cos v$, multiplying the first equation by $S_s(x)S_{-r}(y)$, the second by $S_{s+p}(x)S_{q-r}(y)$ and subtracting yields the polynomial equation

$$(49) \quad \begin{aligned} p_{p,q,r,s}(x, y) \\ := T_{s+p}(x)T_{q-r}(y)S_s(x)S_{-r}(y) - S_{s+p}(x)S_{q-r}(y)T_s(x)T_{-r}(y) = 0. \end{aligned}$$

Thus the map

$$\tilde{V}_{p,q,r,s} \rightarrow \mathbb{R}^2, \quad (e^{uQ}, e^{vR}) \mapsto (\cos u, \cos v),$$

takes its image in the zero set of the polynomial $p_{p,q,r,s}(x, y)$ of Equation (49).

Denote by Z the zero set of $p_{p,q,r,s}$. For $(x, y) \in Z$, the two ratios

$$(50) \quad \frac{T_{s+p}(x)T_{q-r}(y)}{\sqrt{(1-x^2)(1-y^2)}S_{s+p}(x)S_{q-r}(y)}, \quad \frac{T_s(x)T_{-r}(y)}{\sqrt{(1-x^2)(1-y^2)}S_s(x)S_{-r}(y)},$$

are equal if neither denominator vanishes.

Denote by $Z_0 \subset Z$ the subset containing those points so that at least one of the denominators in the ratios of (50) is nonzero, and if the other is zero, so is its numerator. Define the function $\tau(x, y)$ on Z_0 to be one of these two ratios, so that

$$\tau(x, y) = \frac{T_{s+p}(x)T_{q-r}(y)}{\sqrt{(1-x^2)(1-y^2)}S_{s+p}(x)S_{q-r}(y)} = \frac{T_s(x)T_{-r}(y)}{\sqrt{(1-x^2)(1-y^2)}S_s(x)S_{-r}(y)}$$

for $(x, y) \in Z_0$ (and at least one of these ratios is defined).

Denote by $Z_1 \subset Z$ the subset containing those points so that both denominators and both numerators in the ratios of (50) are zero.

Call two pairs of quaternions (M, N) and (M', N') conjugate if there exists $g \in \text{SU}(2)$ so that $(M', N') = (gMg^{-1}, gNg^{-1})$.

Theorem 11.3 *Let $p_{p,q,r,s}(x, y)$ be the polynomial defined above in Equation (49), and let Z denote its zero set in \mathbb{R}^2 , and $Z_0, Z_1 \subset Z$ the subsets defined above.*

If $(M, N) = (e^{uQ}, e^{vR}) \in V_{p,q,r,s}$ then $p_{p,q,r,s}(\cos u, \cos v) = 0$, and the fiber of the map

$$V_{p,q,r,s} \rightarrow Z, \quad (e^{uQ}, e^{vR}) \mapsto (\cos u, \cos v),$$

over a point $(x, y) \in Z$ is given as follows.

(i) *If $|x| < 1, |y| < 1, (x, y) \in Z_0$, and $|\tau(x, y)| \leq 1$, then the fiber over (x, y) is a single point, conjugate to (e^{uQ}, e^{vR}) , where $u = \arccos x, v = \arccos y$, and*

$$Q = i, \quad R = e^{tk}i, \quad t = \arccos(\tau(x, y)).$$

The corresponding point (γ, θ) in the pillowcase $R(S^2, \{a, b, c, d\})$ satisfies

$$\cos \gamma = -T_{2s+p}(x)T_{q-2r}(y) + \sqrt{(1-x^2)(1-y^2)}S_{2s+p}(x)S_{q-2r}(y)\tau(x, y),$$

$$\cos(\gamma - \theta) = T_p(x)T_q(y) - \sqrt{(1-x^2)(1-y^2)}S_p(x)S_q(y)\tau(x, y).$$

(ii) *If $|x| < 1, |y| < 1$, and $(x, y) \in Z_1$, then the fiber over (x, y) is an arc conjugate to the arc $t \mapsto (e^{ui}, e^{ve^{tk}i})$, $t \in [0, \pi]$, where $u = \arccos x, v = \arccos y$.*

The image $(\gamma(t), \theta(t))$ of this path in the pillowcase $R(S^2, \{a, b, c, d\})$ satisfies

$$\cos \gamma = -T_{2s+p}(x)T_{q-2r}(y) + \sqrt{(1-x^2)(1-y^2)}S_{2s+p}(x)S_{q-2r}(y) \cos t,$$

$$\cos(\gamma - \theta) = T_p(x)T_q(y) - \sqrt{(1-x^2)(1-y^2)}S_p(x)S_q(y) \cos t.$$

(iii) *If $|x| \leq 1, |y| \leq 1$, one of $|x|, |y|$ equals 1, and $(x, y) \in Z_1$, then the fiber over (x, y) is a single point.*

The corresponding point (γ, θ) in the pillowcase $R(S^2, \{a, b, c, d\})$ satisfies

$$\begin{aligned} \cos \gamma &= -T_{2s+p}(x)T_{q-2r}(y), \\ \cos(\gamma - \theta) &= T_p(x)T_q(y). \end{aligned}$$

(iv) *In all other cases, the fiber is empty, ie (x, y) is not in the image.*

Proof If $(M, N) = (e^{uQ}, e^{vR}) \in V_{p,q,r,s}$, then (u, v, Q, R) satisfy (48), and hence satisfy $p_{p,q,r,s}(x, y) = 0$, where $x = \cos v, y = \cos u$. In particular, $|x| \leq 1, |y| \leq 1$.

If neither of the denominators in Equation (50) vanishes, then $(x, y) \in Z_0, |x| < 1, |y| < 1$, and $\tau(x, y) = Q \cdot R$ and hence $|\tau(x, y)| \leq 1$. If exactly one of the denominators in Equation (50) vanishes, then (48) show its numerator also vanishes, thus $(x, y) \in Z_0$ and $|\tau(x, y)| = |Q \cdot R| \leq 1$. If $(x, y) \notin Z_0$, then (48) show that both numerators are zero, so that $(x, y) \in Z_1$. Moreover, $Q \cdot R$ can be any number in $[-1, 1]$, ie $Q \cdot R = \cos t$ for some $t \in [0, \pi]$. By conjugating the pair (Q, R) we may assume $Q = \mathbf{i}$ and $R = e^{tk} \mathbf{i}$. Thus the image of $V_{p,q,r,s} \rightarrow Z$ lies in $Z_0 \cup Z_1$ and we have established (iii).

Conversely, suppose that $(x, y) \in Z_0, |x| < 1, |y| < 1$, and $|\tau(x, y)| \leq 1$. Let

$$u = \arccos x, \quad v = \arccos y, \quad t = \arccos(\tau(x, y)), \quad Q = \mathbf{i}, \quad R = e^{tk} \mathbf{i}.$$

Then (u, v, Q, R) satisfy (48). Hence the pair $(M', N') = (e^{uQ}, e^{vR})$ satisfies Equation (44), thus defining a traceless representation, which is uniquely conjugate to a pair $(M, N) \in V_{p,q,r,s}$ by Theorem 11.1. Conjugation does not change u, v nor x, y . Proposition 3.1 implies that there exists a unit quaternion which conjugates the triple

$$(e^{(p+s)uQ} e^{(q-r)vR}, e^{-rvR} e^{suQ}, e^{puQ} e^{qvR}) \quad \text{to} \quad (\mathbf{i}, e^{\gamma k} \mathbf{i}, e^{(\gamma-\theta)k}).$$

Therefore,

$$-\cos \gamma = \text{Re}(e^{-rvR} e^{suQ} e^{(p+s)uQ} e^{(q-r)vR}), \quad \cos(\gamma - \theta) = \text{Re}(e^{puQ} e^{qvR}).$$

Using conjugation invariance and part (iv) of Proposition 2.1, these equations can be rewritten as

$$\begin{aligned} (51) \quad & -\cos \gamma = \cos((2s + p)u) \cos((q - 2r)v) \\ & -\sin((2s + p)u) \sin((q - 2r)v) Q \cdot R, \\ & \cos(\gamma - \theta) = \cos(pu) \cos(qv) - \sin(pu) \sin(qv) Q \cdot R. \end{aligned}$$

Substituting x and y transforms these to

$$\begin{aligned} -\cos \gamma &= T_{2s+p}(x) T_{q-2r}(y) - \sqrt{(1-x^2)(1-y^2)} S_{2s+p}(x) S_{q-2r}(y) \tau(x, y), \\ \cos(\gamma - \theta) &= T_p(x) T_q(y) - \sqrt{(1-x^2)(1-y^2)} S_p(x) S_q(y) \tau(x, y), \end{aligned}$$

establishing (i).

Suppose that $(x, y) \in Z_1$ and $|x| \leq 1, |y| \leq 1$. Then, for any $t \in [0, \pi]$, define

$$u = \arccos x, \quad v = \arccos y, \quad Q = \mathbf{i}, \quad R = e^{tk} \mathbf{i}.$$

Then (u, v, Q, R) satisfy (48). Hence $(M', N') = (e^{uQ}, e^{vR})$ satisfies Equation (44), defining a traceless representation which is conjugate to a pair $(M, N) \in V_{p,q,r,s}$ by Theorem 11.1. Conjugation does not change $u, v, Q \cdot R$, nor x, y .

If $|x| = 1$, then $u = 0$ or π , so that $(M', N') = (\pm 1, e^{vR})$, which is conjugate to $(\pm 1, e^{vi}) \in V_{p,q,r,s}$, and in particular independent of t . Therefore the fiber of $V_{p,q,r,s} \rightarrow Z_1 \cap \{|x| = 1\}$ is a single point. If $|y| = 1$, $(M', N') = (e^{ui}, \pm 1) = (M, N)$ and so again the fiber of $V_{p,q,r,s} \rightarrow Z_1 \cap \{|y| = 1\}$ is a single point.

When $|x| < 1$ and $|y| < 1$, (M, N) is uniquely determined by (M', N') . The assertions about $\cos \gamma$ and $\cos(\gamma - \theta)$ follow from Equation (51). Thus we have established (ii) and (iii). □

The statement of Theorem 11.3 is unfortunately somewhat technical, and does not easily reveal the structure of $R(Y_0, K_0)$ and its image in the pillowcase. However, the polynomials $p_{p,q,r,s}(x, y)$ can be computed and their zero sets graphed using computer algebra software. We present a few examples.

The $(2, 2n + 1)$ torus knots are particularly simple to understand in this context. Take $p = 2, q = 2n + 1, r = n + 1, s = -1$. Then $p_{2,2n+1,n+1,-1}(x, y) = x$, and $\tau(x, y) = 0$ along the arc $x = 0, y \in [-1, 1]$. Theorem 11.3 then says that $V_{2,2n+1,n+1,-1}$ is an arc, and gives a parameterization by (e^{uQ}, e^{vR}) where

$$u = \arccos(0) = t = \frac{\pi}{2}, \quad v = \arccos(y) \in [0, \pi], \quad Q = i, \quad R = e^{tk} i = j.$$

Conjugating $(e^{uQ}, e^{vR}) = (i, e^{vj})$ by $e^{\pi/4i} e^{(\pi nv)/2j}$ yields $(M, N) = (e^{nvk} i, e^{vk})$ which satisfies $F(M, N) = (i, e^{(\pi-v)k} i)$, and hence parametrizes $V_{2,2n+1,n+1,-1}$. Computing $M^2 N^{2n+1}$ yields $e^{(\pi+(2n+1)v)k}$ and from Theorem 11.1 we see that $\gamma = \pi - v$ and $\gamma - \theta = \pi + (2n + 1)v$, so that $\theta = (2n + 2)\gamma \pmod{2\pi}$. This is the same arc identified in the examination of 2-bridge knots in Section 10, and is also the same arc produced by the cross section of Proposition 11.2 (note that $q - 2r = -1$). Theorem 11.3 gives $\cos \gamma = -\cos v$ and $\cos(\gamma - \theta) = -\cos((2n + 1)v)$. The first equation implies $\gamma = \pi - v$ and the second that $\gamma - \theta = \pi \pm (2n + 1)v$, which implies $(1 \pm (2n + 1))\gamma = \theta$. So these equations are not quite sharp enough to give $\theta = (2n + 2)\gamma$.

We turn to the $(3, 4)$ torus knot. This knot is interesting because $R(Y_0, K_0)$ is singular and, as we shall see below, the instanton complex $CI^{\natural}(S^3, T_{3,4})$ has a nontrivial differential.

Take $p = 3, q = 4, r = 3, s = -2$. The polynomial $p_{3,4,3,-2}(x, y)$ of Equation (49) is computed, using Equation (47), as

$$p_{3,4,3,-2}(x, y) = y(4x^2 + 4y^2 - 3).$$

We show that $V_{3,4,3,-2}$ is the union of an arc and a circle that meet in two points.

The zero set of $p_{3,4,3,-2}$ meets the set $Z_0 \cup Z_1$ of [Theorem 11.3](#) in union of the arc $(x, 0), x \in [-1, 1]$ and the circle $x^2 + y^2 = \frac{3}{4}$. The endpoints $(\pm 1, 0)$ of the arc lie in Z_1 and fall under case (iii) of [Theorem 11.3](#), and the rest of the points lie in Z_0 . Moreover, $\tau(x, y) = xy/\sqrt{(1-x^2)(1-y^2)}$, which is less than 1 on Z_0 , so that the map

$$V_{3,4,3,-2} \rightarrow Z \cap \{|x| \leq 1, |y| \leq 1\}$$

is a homeomorphism.

Applying [Theorem 11.3](#) we see that the arc $(x, 0), x \in [-1, 1]$ has $y = 0$ and $\tau(x, y) = 0$, so that $v = t = \frac{\pi}{2}$ and hence gives the arc $(e^{uQ}, e^{vR}) = (e^{ui}, \mathbf{j}), u \in [0, \pi]$ in $\tilde{V}_{3,4,3,-2}$. Conjugating by $e^{-\pi/4i} e^{-\pi/4k} e^{-u/2i}$ yields the arc $(e^{uk}, e^{-uk} \mathbf{j})$. This arc lies in $V_{3,4,3,-2}$, since it is sent by F to the arc $(\mathbf{i}, e^{uk} \mathbf{i})$. This is the arc identified in [Proposition 11.2](#) (after a change of notation, since we are taking p odd and q even here; the condition $q - 2r = \pm 1$ transforms to $p + 2s = \pm 1$).

Since $y = \cos v = 0$ along this arc and $\tau(x, 0) = 0$, [Theorem 11.3](#) gives

$$\begin{pmatrix} \cos \gamma \\ \cos(\gamma - \theta) \end{pmatrix} = \begin{pmatrix} -T_1(x)T_{-2}(y) \\ T_3(x)T_4(y) \end{pmatrix} = \begin{pmatrix} \cos u \\ \cos(3u) \end{pmatrix}$$

and so $\gamma = u$ and $\gamma - \theta = \pm 3u$. Thus $\theta = -2\gamma$ or $\theta = 4\gamma$. At the point $x = 0, u = \frac{\pi}{2}$ and so $(e^{uk}, e^{-uk} \mathbf{j}) = (\mathbf{k}, -\mathbf{j})$. Hence $N^{-r} M^s = \mathbf{j} = e^{\pi/2k} \mathbf{j}$ so that $\gamma = \frac{\pi}{2}$, and $M^p N^q = -\mathbf{k} = e^{-\pi/2k} \mathbf{k}$ so that $\gamma - \theta = -\frac{\pi}{2}$.

Hence at this point (and by continuity along the entire arc) $\theta = -2\gamma$ rather than 4γ . We denote this arc in $V_{3,4,3,-2}$ by I_0 , and consider it parameterized by $\gamma = u \in [0, \pi]$.

The circle $4x^2 + 4y^2 - 3 = 0$ lies entirely in Z_0 , since $\tau(x, y) < 1$ on this circle. It intersects the arc I_0 in the points $(x, y) = (\pm\sqrt{3}/2, 0)$ and since $x = \cos(u)$ and $u = \gamma$, the intersection points occur when $\gamma = \frac{\pi}{6}$ and $\frac{5\pi}{6}$. In particular $V_{3,4,3,-2}$ is singular, and so the map $F: \text{SU}(2) \times \text{SU}(2) \rightarrow \text{SU}(2) \times \text{SU}(2)$ of [Theorem 11.1](#) is not transverse to $\mathbf{i} \times \mathbf{I}$, in contrast to the case for $(2, n)$ torus knots.

We will now use [Theorem 11.3](#) to find the image of the circle in the pillowcase. When $4x^2 + 4y^2 = 3$, then one calculates $\cos \gamma = x$ and $\cos(\gamma - \theta) = -4x^3 + 3x$. Since $\cos(3\gamma) = 4 \cos^2 \gamma + 3 \cos \gamma$, this yields

$$\cos(\gamma - \theta) = -\cos 3\gamma,$$

so that θ equals $\pi + 4\gamma$ or $\pi - 2\gamma$. Since $\cos \gamma = x = \cos u$ and $\gamma, u \in [\frac{\pi}{6}, \frac{5\pi}{6}]$, it follows that $u = \gamma$. From this one computes that at the points $(x, y) = (0, \pm\sqrt{3}/2)$,

$\gamma = \frac{\pi}{2}$ and $\theta = \pi$, so that

$$\theta = \pi + 4\gamma.$$

The map of the circle to the pillowcase takes the pairs (x, y) and $(x, -y)$ in this circle to the same point. Thus we have proved the following.

Proposition 11.4 *For the $(3, 4)$ torus knot, the space $V_{3,4,3,-2} \cong R(Y_0, K_0)$ is homeomorphic to the union of three arcs,*

$$I_0: [0, \pi] \rightarrow V_{3,4,3,-2}, \quad I_{\pm}: [\frac{\pi}{6}, \frac{5\pi}{6}] \rightarrow V_{3,4,3,-2},$$

where

$$I_{\pm}(\frac{\pi}{6}) = I_0(\frac{\pi}{6}), \quad I_{\pm}(\frac{5\pi}{6}) = I_0(\frac{5\pi}{6}).$$

The arc I_0 maps to $\theta = -2\gamma$ and each arc I_{\pm} maps to the arc $\theta = 4\gamma + \pi$ in the pillowcase. □

The following (Figure 19) illustrates both the space $V_{3,4,3,-1}$ and its image in the pillowcase $R(S^2, \{a, b, c, d\})$.




Figure 19: $R(Y_0, T_0)$ and its image in the pillowcase for the $(3,4)$ torus knot

Notice that the images of I_+ , and I_- in the pillowcase each intersect the arc $\theta = \gamma$ (transversely) in one interior point, and I_0 intersects $\theta = \gamma$ in two points, the distinguished representation α at one corner and one interior point. This gives the four traceless representations of the $(3, 4)$ torus knot. Perturbing as above we conclude that the instanton chain complex for $T_{3,4}$ is generated by seven generators, $\alpha', x_1, x_2, y_1, y_2, z_1, z_2$ of the reduced instanton complex. Note that x_i and z_i are mapped to the same point in the pillowcase. The representation labelled y lies on I_0 , and hence by Proposition 11.2 corresponds to a binary dihedral representation.

The perturbation illustrated by the brown circle in Figure 19 corresponds to a small choice of perturbation parameter ϵ . Increasing ϵ eventually moves the intersection point labelled y_2 past the singular point where I_+ and I_- bifurcate from I_0 . This also creates two new intersection points, one each on I_+ and I_- . This presumably corresponds to an elementary expansion of the reduced instanton complex.

The reduced Khovanov homology of the $(3, 4)$ torus knot has rank 5, and since it forms the E_2 page of a spectral sequence converging to $I^{\natural}(S^3, K)$ (see Section 12), one differential in $CI^{\natural}(S^3, T_{3,4})$ is nontrivial. Arguments involving the gradings as in [25] suggest that the differential involves the generator coming from the abelian representation.

We briefly list some other calculations. For the $(4, 5)$ torus knot, taking $p = 4, q = 5, r = 4, s = -3$, we have

$$p_{4,5,4,-3}(x, y) = x(16y^4 + 16x^2y^2 - 20y^2 - 4x^2 + 3),$$

$$\tau(x, y) = \frac{xy}{\sqrt{1-x^2}\sqrt{1-y^2}}.$$

The intersection of the zero set of $p_{4,5,4,-3}(x, y)$ with $Z_0 \cup Z_1$ is the union of an arc (corresponding to $x = 0$) and a circle which meets the arc in two points. The zero set of $p_{4,5,4,-3}(x, y)$ contains two other components which lie outside the square $|x| \leq 1, |y| \leq 1$ and hence do not contribute to $V_{4,5,4,-3}$, and so $V_{4,5,4,-3}$ is the union of a circle and an arc, and is homeomorphic to $V_{3,4,3,-2}$.

For the $(3, 5)$ torus knot, $p_{3,5,2,-1}(x, y) = -8y^4 + 6y^2 - 2x^2$. Its zero set is a figure 8 curve, ie an immersed circle with one double point, located at $(0, 0)$. Moreover, in the notation of Theorem 11.3, $(0, 0) \in Z_1$ so that $V_{3,5,2,-1} \cong R(Y_0, T_0)$ is again a union of a circle and an arc intersecting in two points, and is homeomorphic to $V_{3,4,3,-2}$.

It is not clear from our description whether the image of $R(Y_0, K_0)$ in the pillowcase is always contained in the union of straight lines for all (p, q) torus knots, as is the case

for (p, q) equal to $(2, n)$ and $(3, 4)$. Further calculations reveal that the polynomials $p_{p,q,r,s}$ can be quite complicated and their zero sets highly singular. This topic is explored further in Fukumoto, Kirk and Pinzón-Caicedo [10], where examples are given with nonlinear image.

12 Calculations of $CI^{\natural}(S^3, K)$

In this section, we use our results to carry out some calculations of instanton homology. To set the stage, observe that for a knot in a homology sphere with a simple representation variety, we have arranged that $R_{\pi}^{\natural}(Y, K)$ has $2k + 1$ points, which we label as α' and $\beta_{i,1}, \beta_{i,2}$, $i = 1, \dots, k$. Here α' corresponds to the perturbation of the distinguished isolated point $\alpha \in R^{\natural}(Y, K)$ that restricts to the abelian representation $\alpha_{\pi/2}$ on $Y \setminus N(K)$. For each $i = 1, \dots, k$, the two points $\beta_{i,j} \in R_{\pi}^{\natural}(Y, K)$ are those that result from perturbing the circle of nonabelian traceless representations in $R^{\natural}(Y, K)$ coming from $\beta_i \in R(Y, K)$.

12.1 Summary of the results of Kronheimer and Mrowka

The generators of $CI^{\natural}(Y, K)$ are the points of $R_{\pi}^{\natural}(Y, K)$ for some generic perturbation data π . The differential on $CI^{\natural}(Y, K)$ is defined using moduli spaces of singular instantons on the cylinder $Y \times \mathbb{R}$. Though we suppress it from the notation, it is important to emphasize that the chain group $CI^{\natural}(Y, K)$ depends on the choice of perturbation data π , and that the differential depends on additional perturbation data on the cylinder. Generators of $CI^{\natural}(Y, K)$ come equipped with a relative $\mathbb{Z}/4$ -grading. This grading is determined by the spectral flow of the family of Hessians of the Chern–Simons functional along a path of connections joining a pair of generators. A standard argument shows that the gradings of $\beta_{i,1}$ and $\beta_{i,2}$ differ by 1 (after relabeling if necessary). The relative grading is promoted to an absolute grading by defining a grading difference associated to paths of connections on a cobordism of pairs from (Y, K) to (S^3, U) , where U is the unknot, and normalizing the grading of the unique point of $R^{\natural}(S^3, U)$ [25, Proposition 4.4]. One can use a splitting theorem for spectral flow (eg Atiyah, Patodi and Singer [2], Nicolaescu [32] and Daniel and Kirk [8]) as in Boden, Herald, Kirk and Klassen [4] to compute the relative grading between a pair generators that lie on the same path component in $\chi(Y, K)$. This is implicit in the discussion in [21, Section 11].

In [25], Kronheimer and Mrowka define a different $\mathbb{Z}/4$ -graded chain complex for knots in S^3 whose homology is $I^{\natural}(S^3, K)$. To distinguish it from $CI^{\natural}(S^3, K)$ we denote it by $FCI^{\natural}(S^3, K)$. This chain complex is filtered, and the associated spectral

sequence has E_2 page isomorphic to the reduced Khovanov homology of the mirror image K^m of K , $E_2 \cong \text{Kh}^{\text{red}}(K^m)$. The spectral sequence is $\mathbb{Z}/4$ -graded, and the bigrading on Khovanov homology determines the modulo 4 grading on the E_2 page. Explicitly, a generator of $\text{Kh}_{i,j}^{\text{red}}(K^m)$ with quantum grading i and homological grading j inherits the grading $i - j + 1 \pmod 4$ in the E_2 page [25, Section 8.1]. Note that in that reference it is shown that a generator of *unreduced* Khovanov homology in bigrading (i, j) determines a generator of unreduced instanton homology in bigrading $i - j - 1$; a shift of 2 occurs when passing to reduced homology. Henceforth, the Khovanov groups expressed with a single grading (eg Kh_i^{red}) will refer to this modulo 4 grading.

The chain complex $FCT^{\natural}(S^3, K)$ is built from the hypercube of complete unoriented resolutions of a diagram of K . Its construction relies on the fact that the singular instanton homology groups of knots which differ by the unoriented skein relation fit into an exact triangle. In general, this means that the rank of $FCT^{\natural}(S^3, K)$ will be exponentially greater than the rank of any complex $CT^{\natural}(S^3, K)$ obtained by a nondegenerate perturbation of the Chern–Simons functional. These considerations also show that for each $i \in \mathbb{Z}/4$,

$$(52) \quad \text{rank } FCT_i^{\natural}(S^3, K) \geq \text{rank } \text{Kh}_i^{\text{red}}(K^m) \geq \text{rank } I^{\natural}(S^3, K)_i.$$

Since the homology of $CT^{\natural}(S^3, K)$ equals $I^{\natural}(S^3, K)$, we have the obvious inequality

$$(53) \quad \text{rank } CT_i^{\natural}(S^3, K) \geq \text{rank } I^{\natural}(S^3, K)_i, \quad \text{for each } i \in \mathbb{Z}/4.$$

Kronheimer and Mrowka use excision for instanton homology to show that $I^{\natural}(S^3, K)$ is isomorphic, as a $\mathbb{Z}/4$ graded group, to the sutured instanton Floer homology $KHI(K)$; see [25, Proposition 1.4]. This latter invariant was defined by Floer in [9] and revisited in Kronheimer and Mrowka [24, Section 7]. It has the advantage of possessing an additional \mathbb{Z} grading, and the graded Euler characteristic with respect to this grading equals the Alexander polynomial; see Kronheimer, and Mrowka [23] and Lim [29]. It follows that if $\Delta_K(t) = \sum_i a_i t^i$ denotes the Alexander polynomial, then

$$(54) \quad \text{rank } I^{\natural}(S^3, K) \geq \sum_i |a_i| \geq \left| \sum_i a_i (-1)^i \right| = |\det(K)|.$$

On the other hand, the Euler characteristic of Khovanov homology equals the Jones polynomial [17]

$$J_K(q) = \sum_{i,j} (-1)^j q^{-i} \text{rank } \text{Kh}_{i,j}^{\text{red}}(K).$$

Thus, if we let $J_K(q) = \sum_i b_i q^i$, then we have

$$(55) \quad \text{rank Kh}^{\text{red}}(K) \geq \sum_i |b_i| \geq |J_K(-1)| = |\det(K)|.$$

These inequalities are obviously useful for understanding the behavior of Kronheimer and Mrowka's spectral sequence. For instance, they immediately show that spectral sequence collapses for all 2-bridge knots. This is because $|\det(K)|$ equals the rank of $\text{Kh}^{\text{red}}(K)$ for 2-bridge knots. More generally, the spectral sequence collapses for the same reason for all alternating and quasialternating knots [17], Lee [27] and Manolescu and Ozsváth [31]. Hence Equations (52), (54) and (55) imply that for these knots, $I^{\natural}(S^3, K) = \text{Kh}^{\text{red}}(K^m)$.

Similar facts hold for the unreduced theory $I^{\sharp}(S^3, K)$. In particular the corresponding spectral sequence has E_2 term the unreduced Khovanov homology of K with its bigrading appropriately reduced to a modulo 4 grading. Thus the calculations we give below can be modified to handle the case of unreduced Khovanov homology and $I^{\sharp}(S^3, K)$. The corresponding chain complex $CI^{\sharp}(S^3, K)$ is generated by $R_{\pi}^{\sharp}(S^3, K)$, which, using the perturbations described in Theorems 7.1 and 9.1, has twice as many points as $R_{\pi}^{\natural}(S^3, K)$.

12.2 Remarks on calculations

The calculations of reduced Khovanov homology we give below were obtained using Dror Bar-Natan's Knot theory Mathematica workbook [3]. We work over \mathbb{Q} for the remainder of the article.

Use the notation (a, b, c, d) for the $\mathbb{Z}/4$ graded vector space $(\mathbb{Q}^a, \mathbb{Q}^b, \mathbb{Q}^c, \mathbb{Q}^d)$, so, eg the rank in grading 2 is equal to c and in grading 3 is equal to d . More generally, let $(a, b, c, d)_e$ denote the result of shifting (a, b, c, d) to the right e slots, so eg $(0, 1, 2, 3)_3 = (1, 2, 3, 0) = (\mathbb{Q}, \mathbb{Q}^2, \mathbb{Q}^3, 0)$.

12.2.1 2-bridge knots Theorem 10.2 says that if $K(p/q)$ is a 2-bridge knot, $CI^{\natural}(S^3, K(p/q))$ is generated by $|p| = |\det(K)|$ points. Thus for 2-bridge knots,

$$CI^{\natural}(S^3, K(p/q)) = I^{\natural}(S^3, K) = \text{Kh}^{\text{red}}(K^m)$$

and all differentials in $CI^{\natural}(S^3, K(p/q))$ are zero, as are all higher differentials in the spectral sequence from $\text{Kh}^{\text{red}}(K^m)$ to $I^{\natural}(S^3, K)$. While these facts could be deduced without the use of Theorem 10.2, it would be interesting to use these examples to investigate what grading and differential information can be gleaned from the intersection diagram (2).

For example, Figures 15, 16 and 17 illustrate the trefoil $= K(-3/1)$, with $\text{Kh}^{\text{red}}(K^m) = I^{\natural}(K) = (1, 0, 1, 1)$, the $(2, 5)$ torus knot $= K(-5, 1)$, with $\text{Kh}^{\text{red}}(K^m) = I^{\natural}(K) = (2, 1, 1, 1)$, the figure 8 knot $= K(-5/3)$, with $\text{Kh}^{\text{red}}(K^m) = I^{\natural}(K) = (1, 1, 2, 1)$, and the knot $7_2 = K(-11/5)$, with $\text{Kh}^{\text{red}}(K^m) = I^{\natural}(K) = (3, 2, 3, 3)$.

12.2.2 Knots with simple representation varieties We illustrate one sample calculation; all the data below were obtained by the same method, which is the method described in [21].

12.2.3 The $(3, 4)$ torus knot The bigraded reduced Khovanov homology has Poincaré polynomial $\text{Kh}^{\text{red}}(T_{3,4}^m) = q^{-7} + q^{-17}t^{-5} + q^{-13}t^{-4} + q^{-13}t^{-3} + q^{-11}t^{-2}$. This means that there is a generator of homology in bidegree $(-7, 0)$, one in $(-17, -5)$ etc. The induced modulo 4 graded group has a generator in degree $-7 - 0 + 1 = -6 \equiv 2 \pmod 4$, one in degree $-17 - (-5) + 1 \equiv 1$, etc. Thus the $\mathbb{Z}/4$ graded reduced Khovanov homology has ranks equalling $(2, 1, 1, 1)$.

The character variety $\chi(S^3, T_{3,4})$ is illustrated in Figure 20, and is determined by the data (see Section 5.1)

$$(1, 7), \quad (5, 11), \quad (2, 10).$$

There are four points in $R(S^3, T_{3,4})$, labelled α, x, y, z in the figure.




Figure 20: $\chi(S^3, T_{3,4})$ and its image in the pillowcase

In terms of $R(S^3, T_{3,4})$ and its image in the other pillowcase $R(S^2, \{a, b, c, d\})$, Proposition 11.2 shows that representations which lie on I_0 send N^q to 1, since $q = 4$. Thus the two traceless representations labelled α and y in Figure 20 correspond to the two intersections of $\{\theta = \gamma\}$ with the arc I_0 in Figure 19: the intersection at the corner is α and the nonabelian intersection in the interior of I_0 is y . The remaining two points, x and z lie on the intersection of I_+ and I_- with $\{\theta = \gamma\}$.

From [Theorem 7.1](#) and [Corollary 7.2](#) we see that x, y and z each perturb to give two generators for $CI^{\natural}(S^3, T_{3,4})$, which we label $\{x_1, x_2, y_1, y_2, z_1, z_2\}$. These arise by perturbing a Morse–Bott critical circle, and hence the gradings satisfy $\text{gr}(x_1) - \text{gr}(x_2) = \text{gr}(y_1) - \text{gr}(y_2) = \text{gr}(z_1) - \text{gr}(z_2) = 1$ (after perhaps reindexing).

The last generator α' (the perturbation of α) contributes $(1, 0, 0, 0)_a$ to $CI^{\natural}(S^3, T_{3,4})$. Thus we see that $CI^{\natural}(S^3, T_{3,4})$ is a direct sum of graded groups

$$CI^{\natural}(S^3, T_{3,4}) = (1, 0, 0, 0)_a \oplus (1, 1, 0, 0)_{b(x)} \oplus (1, 1, 0, 0)_{b(y)} \oplus (1, 1, 0, 0)_{b(z)}.$$

Using splitting theorems for spectral flow, such as those of [\[4\]](#) (see also the discussion in [\[21\]](#)) one can see from [Figure 20](#) that $b(y) = b(x) + 2$ and $b(z) = b(x) + 4 \equiv b(x)$. Thus

$$CI^{\natural}(S^3, T_{3,4}) = (1, 0, 0, 0)_a \oplus (2, 2, 1, 1)_b,$$

where $b = b(x)$. The Alexander polynomial $\Delta_K(t) = t^3 + 1/t^3 - t^2 - 1/t^2 + 1$. The sum of the absolute value of its coefficients tells us the rank of $I^{\natural}(S^3, T_{3,4})$ is at least 5, and hence $I^{\natural}(S^3, T_{3,4}) = \text{Kh}^{\text{red}}(T_{3,4}^m) = (2, 1, 1, 1)$. Thus the rank of the differential on the complex $CI^{\natural}(S^3, T_{3,4})$ is one, and the spectral sequence collapses after the E_2 page.

In this example, the representation labeled y is the only nonabelian binary dihedral representations, and hence corresponds to the representation y of [Figure 19](#).

12.3 The $(2, n)$ torus knots

For the $(2, 2k + 1)$ torus knots, there are k nonabelian arcs in $\chi(S^3, T_{2,2k+1})$, and these are nested. Each arc contains a traceless representation, and the signature of $T_{2,2k+1}$ equals $-2k$.

Therefore $R_{\pi}^{\natural}(S^3, K)$ consists of $2k + 1$ points which correspondingly generate $CI^{\natural}(S^3, T_{2,2k+1})$. Spectral flow considerations show that since the nonabelian arcs in $\chi(S^3, T_{2,2k+1})$ are nested, the grading difference from α to each nonabelian point in $R(S^3, K)$ is successively $b, b + 2, b + 4, \dots$ modulo 4 for some integer b . Hence

$$\begin{aligned} I^{\natural}(S^3, K) &= CI^{\natural}(S^3, T_{2,2k+1}) \\ &= (1, 0, 0, 0)_{a(k)} \oplus \bigoplus_{j=1}^k (1, 1, 0, 0)_{b+2j} \\ &= (1, 0, 0, 0)_{a(k)} \oplus \begin{cases} \left(\frac{k+1}{2}, \frac{k+1}{2}, \frac{k-1}{2}, \frac{k-1}{2}\right)_{b(k)} & \text{if } k \text{ is odd,} \\ \left(\frac{k}{2}, \frac{k}{2}, \frac{k}{2}, \frac{k}{2}\right) & \text{if } k \text{ is even.} \end{cases} \end{aligned}$$

where $a(k)$ denotes the grading of the generator α' , and $b(k)$ is some integer.

On the other hand, one can easily compute $\text{Kh}^{\text{red}}(T_{2,2k+1}^m)$ (eg, by the method of Khovanov [16, Section 6.2]) to find

$$\text{Kh}^{\text{red}}(T_{2,2k+1}^m) = (1, 0, 0, 0) \oplus \begin{cases} \left(\frac{k-1}{2}, \frac{k-1}{2}, \frac{k+1}{2}, \frac{k+1}{2}\right) & \text{if } k \text{ is odd,} \\ \left(\frac{k}{2}, \frac{k}{2}, \frac{k}{2}, \frac{k}{2}\right) & \text{if } k \text{ is even.} \end{cases}$$

One sees that these values are consistent with the possibility that $a(k) = \sigma(T_{2,2k+1}) \pmod 4$ and $b(k) = 3$ for all k .

12.4 The (3, n) torus knots

For the (3, n) torus knots with $n \leq 38$, we have

$$CI^{\natural}(S^3, T_{3,n}) = (1, 0, 0, 0)_{a(n)} \oplus \begin{cases} (2k, 2k, 2k, 2k) & \text{if } n = 6k + 1, \\ (2k + 1, 2k + 1, 2k, 2k)_{b(n)} & \text{if } n = 6k + 2, \\ (2k + 2, 2k + 2, 2k + 1, 2k + 1)_{b(n)} & \text{if } n = 6k + 4, \\ (2k + 2, 2k + 2, 2k + 2, 2k + 2) & \text{if } n = 6k + 5, \end{cases}$$

$$\text{Kh}^{\text{red}}(T_{3,n}^m) = \begin{cases} (2k + 1, 2k, 2k, 2k) & \text{if } n = 6k + 1, \\ (2k + 1, 2k, 2k + 1, 2k + 1) & \text{if } n = 6k + 2, \\ (2k + 2, 2k + 1, 2k + 1, 2k + 1) & \text{if } n = 6k + 4, \\ (2k + 2, 2k + 1, 2k + 2, 2k + 2) & \text{if } n = 6k + 5. \end{cases}$$

For these knots, the sum of the absolute values of the coefficients of the Alexander polynomial $|\Delta|$ equals the rank of the reduced Khovanov homology. The absolute value of the signature $|\sigma|$ satisfies

$$|\sigma(T_{3,n})| + 1 = \begin{cases} |\Delta(T_{3,n})| & \text{if } n = 6k + 1, 6k + 2, \\ |\Delta(T_{3,n})| + 2 & \text{if } n = 6k + 4, 6k + 5. \end{cases}$$

Thus $\text{Kh}^{\text{red}}(T_{3,n}^m) = I^{\natural}(S^3, K_{3,n})$, and when $n = 6k + 1, 6k + 2$, there are no nonzero differentials in the complex $CI^{\natural}(S^3, T_{3,n})$. When $n = 6k + 4, 6k + 5$, the rank of the differential is one. In either case there are no nonzero differentials after the E_2 page in the Kronheimer–Mrowka spectral sequence. The data are consistent with the guess $a(n) = \sigma(T_{3,n}) \pmod 4$ and $b(n) = 3$. The unreduced Khovanov homology is known for all (3, n) torus knots; see Turner [38]. Presumably similar calculations could verify the formulae above for all n .

12.5 Other torus knots

Patterns for more complicated torus knots are not as obvious. But for fixed m , the rank $|\sigma| + 1$ of $CI^{\natural}(S^3, T_{m,n})$ and the lower bound $|\Delta|$ grow linearly in n , whereas the rank of $\text{Kh}^{\text{red}}(T_{m,n})$ seems to be growing more quickly for $m \geq 4$.

The computations of CI^{\natural} for torus knots are done by computing their data in the sense of Section 5.1, using the MAPLE computer algebra package. Some examples were listed in Section 5.1. Table 1 includes the signature σ and the sum of the absolute values of the coefficients of the Alexander polynomial $|\Delta|$. We write A_a for $(1, 0, 0, 0)_a$. Whenever $CI^{\natural}(K)$ and $I^{\natural}(K)$ differ the chain complex $CI^{\natural}(K)$ has nontrivial differential. Whenever $\text{Kh}^{\text{red}}(K)$ and $I^{\natural}(K)$ differ the Kronheimer–Mrowka spectral sequence has nontrivial higher differentials.

A torus knot which exhibits interesting CI^{\natural} is the $(4, 5)$ torus knot, with (c_i, d_i) equal to

$$(1, 9), \quad (7, 17), \quad (2, 18), \quad (6, 14), \quad (11, 19), \quad (3, 13).$$

The first and fourth arc do not contain traceless representations, and hence do not contribute to $CI^{\natural}(S^3, T_{4,5})$. The remaining four arcs contain traceless representations, and hence $CI^{\natural}(S^3, T_{4,5})$ is generated by $8 + 1 = 9$ elements. As explained in [21], this example is remarkable because although the Khovanov homology also has 9 generators, the gradings ($CI^{\natural} = (3, 2, 2, 2)_a$ and $\text{Kh}^{\text{red}} = (2, 1, 3, 3)$) are incompatible with a spectral sequence with no higher differentials. Since $|\Delta| = 7$, it follows that the differential on $CI^{\natural}(S^3, T_{4,5})$ has rank one. As mentioned in [21], this is not quite enough to compute $I^{\natural}(S^3, T_{4,5})$, as both $(2, 1, 2, 2)$ and $(1, 1, 3, 2)$ are compatible. In [21] they establish that $I^{\natural}(S^3, T_{4,5}) = (2, 1, 2, 2)$ using the results of [23; 29] which identify the coefficients of the Alexander polynomial with a kind of Euler characteristic associated to the generalized eigenspaces of an operator $\mu: KHI(K) \rightarrow KHI(K)$. If, as seems likely, the grading of the generator α occurs in degree $\sigma(K) \bmod 4$ (see below) then this would also show that $I^{\natural}(S^3, T_{4,5}) = (2, 1, 2, 2)$.

In the spectral sequence for $T_{4,9}$, the rank from the E_2 page to the limit drops by at least 8. For $T_{5,7}$, all differentials in $CI^{\natural}(S^3, T_{5,7})$ are zero, but in the spectral sequence, the rank drops from 29 for the E_2 page to 17 in the limit.

For $T_{5,6}$, the ranks of $CI^{\natural}(S^3, T_{5,6})$ and $\text{Kh}^{\text{red}}(T_{5,6}^m)$ are equal, but the gradings are different so that the spectral sequence necessarily has higher differentials, and hence $CI^{\natural}(S^3, T_{5,6})$ also has nontrivial differentials. For this knot, the knot Floer homology group has rank equal to 9, and hence further examination of this knot may shed light on the relationship between the three knot invariants.

For the larger values of (p, q) in Table 1, examples were chosen so that the Alexander polynomial sufficed to conclude that $CI^{\natural}(S^3, K)$ has no higher differentials. This is because $|\sigma(K)| + 1$ equals $|\Delta|$.

12.6 Speculation

The data calculated above are consistent with the conjecture that the generator α of $CI^{\natural}(K)$ has grading equal to the signature $\sigma(K)$ modulo 4. This is likely true, and a proof should follow from the following outline. Push a Seifert surface for K into the 4–ball and surger along half a symplectic basis. This yields a 4–manifold X with boundary in which K bounds a disk D , such that α extends to $\pi_1(X_0)$ where $X_0 = X \setminus N(D)$. The signature of (X_0, α) gives the signature of K . Now glue in $(S^1 \times D^2) \times [0, 1]$, containing the surfaces $(S^1 \sqcup H) \times [0, 1]$ together with the singular bundle data over $W \times [0, 1]$ (ie cross Figure 1 with an interval and glue it to X_0 along an annulus in D times an interval). This gives a flat singular cobordism from (S^3, K) to (S^3, U) , where U denotes the unknot. The Fredholm index of this flat cobordism (in the sense of [25, Proposition 4.4]) should be the signature of K by an excision argument.

Note that it has been conjectured by Kronheimer and Mrowka [24, page 302] that the sutured instanton homology groups of a knot defined in [24] are isomorphic to the Heegaard knot Floer homology groups of Ozsváth and Szabó [33] and Rasmussen [36]. Since the sutured instanton homology groups have the same rank as $I^{\natural}(K)$, this conjecture would imply that $I^{\natural}(K)$ has the same rank as the Heegaard knot Floer homology groups. Denoted $\widehat{HFK}(K)$, these latter groups are known to satisfy

$$\text{rk}\widehat{HFK}(K) = |\Delta|$$

in the special case when K is a torus knot (or, more generally, when K is a so-called L –space knot; see Ozsváth and Szabó [34, Theorem 1.2]). This would imply that all the lower bounds for $I^{\natural}(K)$ in Table 1 are attained, and hence the boundary operator acting on our chain complex should have large rank, in general.

Looking at Table 1, one could also conjecture that $b = 3 = -1 \pmod 4$. This corresponds to a spectral flow along an arc of flat connections in $\chi(S^3, K)$ starting at the abelian flat connection α traveling towards the trivial connection (ie α_t as t decreases from $t = \frac{1}{2}$ in the notation of Definition 5.1) and changing branches into the first irreducible arc encountered which meets $S(\mathbf{i})$. For general knots with simple representation varieties b might be ∓ 1 , according to the sign of the change in Levine–Tristram signatures.

Torus knot	σ	$ \Delta $	$CI^\natural(K)$	$\text{Kh}^{\text{red}}(K^m)$	$I^\natural(K)$
(4,5)	-8	7	$(3, 2, 2, 2)_a$	(2, 1, 3, 3)	
(4,7)	-14	11	$A_a \oplus (4, 4, 3, 3)_b$,	(4, 4, 5, 4)	$11 \leq \text{rank} \leq 15$
(4,9)	-16	13	$(5, 4, 4, 4)_a$	(7, 6, 6, 6)	$13 \leq \text{rank} \leq 17$
(4,11)	-22	17	$A_a \oplus (6, 6, 5, 5)_b$	(10, 9, 9, 9)	$17 \leq \text{rank} \leq 23$
(4,13)	-24	19	$(7, 6, 6, 6)_a$	(12, 11, 13, 13)	$19 \leq \text{rank} \leq 25$
(4,15)	-30	23	$A_a \oplus (8, 8, 7, 7)_b$	(16, 16, 17, 16)	$23 \leq \text{rank} \leq 31$
(4,17)	-32	25	$A_a \oplus (8, 8, 8, 8)$	(21, 20, 20, 20)	$25 \leq \text{rank} \leq 33$
(4,19)	-38	29	$A_a \oplus (10, 10, 9, 9)_b$	(26, 25, 25, 25)	$29 \leq \text{rank} \leq 39$
(4,21)	-40	31	$A_a \oplus (10, 10, 10, 10)$	(30, 29, 31, 31)	$31 \leq \text{rank} \leq 41$
(4,23)	-46	35	$A_a \oplus (12, 12, 11, 11)_b$	(36, 36, 37, 36)	$35 \leq \text{rank} \leq 47$
(4,25)	-48	37	$A_a \oplus (12, 12, 12, 12)$	(43, 42, 43, 42)	$37 \leq \text{rank} \leq 49$
(5,6)	-16	9	$(5, 4, 4, 4)_a$	(5, 3, 4, 5)	$9 \leq \text{rank} \leq 15$
(5,7)	-16	17	$(5, 4, 4, 4)_a$	(8, 6, 7, 8)	$(5, 4, 4, 4)_a$
(5,8)	-20	19	$(6, 5, 5, 5)_a$	(9, 8, 9, 9)	$19 \leq \text{rank} \leq 21$
(5,9)	-24	15	$(7, 6, 6, 6)_a$	(10, 10, 11, 10)	$15 \leq \text{rank} \leq 25$
(5,11)	-24	17	$(7, 6, 6, 6)_a$	(15, 14, 14, 14)	$17 \leq \text{rank} \leq 25$
(5,12)	-28	29	$(8, 7, 7, 7)_a$	(20, 19, 19, 19)	$(8, 7, 7, 7)_a$
(5,17)	-40	41	$(11, 10, 10, 10)_a$	(38, 36, 37, 38)	$(11, 10, 10, 10)_a$
(5,22)	-52	53	$(14, 13, 13, 13)_a$	(62, 61, 61, 61)	$(14, 13, 13, 13)_a$
(5,117)	-280	281	$(71, 70, 70, 70)_a$?	$(71, 70, 70, 70)_a$
(6,7)	-18	11	$A_a \oplus (5, 5, 4, 4)_b$	(7, 7, 9, 8)	$11 \leq \text{rank} \leq 19$
(7,16)	-54	55	$A_a \oplus (14, 14, 13, 13)_b$?	$A_a \oplus (14, 14, 13, 13)_b$
(7,30)	-102	103	$A_a \oplus (26, 26, 25, 25)_b$?	$A_a \oplus (26, 26, 25, 25)_b$
(9,11)	-48	49	$(13, 12, 12, 12)_a$?	$(13, 12, 12, 12)_a$
(9,25)	-112	111	$(29, 28, 28, 28)_a$?	$111 \leq \text{rank} \leq 113$
(9,29)	-128	129	$(33, 32, 32, 32)_a$?	$(33, 32, 32, 32)_a$
(11,24)	-130	131	$A_a \oplus (33, 33, 32, 32)_b$?	$A_a \oplus (33, 33, 32, 32)_b$
(11,31)	-168	169	$(43, 42, 42, 42)_a$?	$(43, 42, 42, 42)_a$
(13,15)	-96	97	$(25, 24, 24, 24)_a$?	$(25, 24, 24, 24)_a$
(13,28)	-180	181	$(46, 45, 45, 45)_a$?	$(46, 45, 45, 45)_a$

Table 1: Calculations for torus knots

Much more ambitiously, one could hope that an Atiyah–Floer conjecture holds in this context, which would describe how to calculate gradings and differentials from intersection diagrams in the pillowcase (and perhaps additional data internal to $R(Y_0, K_0)$). Exploring this topic provides motivation for the problem of describing the spaces $R(Y_0, K_0)$ and their image in the pillowcase for more general tangles than those that arise from 2–bridge and torus knots. It would be interesting to relate our examples and calculations to the approach of Wehrheim and Woodward as alluded to in [39].

References

- [1] **S Akbulut, JD McCarthy**, *Casson’s invariant for oriented homology 3–spheres*, Mathematical Notes 36, Princeton Univ. Press (1990) [MR1030042](#)
- [2] **MF Atiyah, V K Patodi, IM Singer**, *Spectral asymmetry and Riemannian geometry, I*, Math. Proc. Cambridge Philos. Soc. 77 (1975) 43–69 [MR0397797](#)
- [3] **D Bar-Natan**, *The Mathematica package KnotTheory* Available at http://katlas.math.toronto.edu/wiki/The_Mathematica_Package_KnotTheory
- [4] **H U Boden, C M Herald, P A Kirk, E P Klassen**, *Gauge theoretic invariants of Dehn surgeries on knots*, Geom. Topol. 5 (2001) 143–226 [MR1825661](#)
- [5] **G Burde**, *SU(2)–representation spaces for two-bridge knot groups*, Math. Ann. 288 (1990) 103–119 [MR1070927](#)
- [6] **G Burde, H Zieschang**, *Knots*, de Gruyter Studies in Mathematics 5, de Gruyter, Berlin (1985) [MR808776](#)
- [7] **D Cooper, M Culler, H Gillet, DD Long, P B Shalen**, *Plane curves associated to character varieties of 3–manifolds*, Invent. Math. 118 (1994) 47–84 [MR1288467](#)
- [8] **M Daniel, P A Kirk**, *A general splitting formula for the spectral flow*, Michigan Math. J. 46 (1999) 589–617 [MR1721567](#) With an appendix by K. P. Wojciechowski
- [9] **A Floer**, *Instanton homology, surgery, and knots*, from: “Geometry of low-dimensional manifolds, 1”, (S K Donaldson, C B Thomas, editors), London Math. Soc. Lecture Note Ser. 150, Cambridge Univ. Press (1990) 97–114 [MR1171893](#)
- [10] **Y Fukumoto, P A Kirk, J Pinzón-Caicedo**, *Traceless SU(2) representations of 2–stranded tangles* [arXiv:1305.6042](#)
- [11] **B Gerard**, *Singular connections on 3–manifolds and manifolds with cylindrical ends* [arXiv:math/9805036](#)
- [12] **CM Herald**, *Legendrian cobordism and Chern–Simons theory on 3–manifolds with boundary*, Comm. Anal. Geom. 2 (1994) 337–413 [MR1305710](#)
- [13] **CM Herald**, *Flat connections, the Alexander invariant, and Casson’s invariant*, Comm. Anal. Geom. 5 (1997) 93–120 [MR1456309](#)

- [14] **M Heusener, J Kroll**, *Deforming abelian $SU(2)$ -representations of knot groups*, Comment. Math. Helv. 73 (1998) 480–498 [MR1633375](#)
- [15] **M Jacobsson, R L Rubinsztein**, *Symplectic topology of $SU(2)$ -representation varieties and link homology, I: Symplectic braid action and the first Chern class* [arXiv:0806.2902](#)
- [16] **M Khovanov**, *A categorification of the Jones polynomial*, Duke Math. J. 101 (2000) 359–426 [MR1740682](#)
- [17] **M Khovanov**, *Patterns in knot cohomology, I*, Experiment. Math. 12 (2003) 365–374 [MR2034399](#)
- [18] **P A Kirk, E P Klassen**, *Chern–Simons invariants of 3-manifolds and representation spaces of knot groups*, Math. Ann. 287 (1990) 343–367 [MR1054574](#)
- [19] **P A Kirk, C Livingston**, *Concordance and mutation*, Geom. Topol. 5 (2001) 831–883 [MR1871406](#)
- [20] **E P Klassen**, *Representations of knot groups in $SU(2)$* , Trans. Amer. Math. Soc. 326 (1991) 795–828 [MR1008696](#)
- [21] **P B Kronheimer, T S Mrowka**, *Filtrations on instanton homology* [arXiv 1110.1290](#)
- [22] **P B Kronheimer, T S Mrowka**, *Dehn surgery, the fundamental group and $SU(2)$* , Math. Res. Lett. 11 (2004) 741–754 [MR2106239](#)
- [23] **P B Kronheimer, T S Mrowka**, *Instanton Floer homology and the Alexander polynomial*, Algebr. Geom. Topol. 10 (2010) 1715–1738 [MR2683750](#)
- [24] **P B Kronheimer, T S Mrowka**, *Knots, sutures, and excision*, J. Differential Geom. 84 (2010) 301–364 [MR2652464](#)
- [25] **P B Kronheimer, T S Mrowka**, *Khovanov homology is an unknot-detector*, Publ. Math. Inst. Hautes Études Sci. (2011) 97–208 [MR2805599](#)
- [26] **P B Kronheimer, T S Mrowka**, *Knot homology groups from instantons*, J. Topol. 4 (2011) 835–918 [MR2860345](#)
- [27] **E S Lee**, *An endomorphism of the Khovanov invariant*, Adv. Math. 197 (2005) 554–586 [MR2173845](#)
- [28] **S Lewallen**, *Khovanov homology of alternating links and $SU(2)$ representations of the link group* [arXiv:0910.5047](#)
- [29] **Y Lim**, *Instanton homology and the Alexander polynomial*, Proc. Amer. Math. Soc. 138 (2010) 3759–3768 [MR2661575](#)
- [30] **X-S Lin**, *A knot invariant via representation spaces*, J. Differential Geom. 35 (1992) 337–357 [MR1158339](#)

- [31] **C Manolescu, P Ozsváth**, *On the Khovanov and knot Floer homologies of quasi-alternating links*, from: “Proceedings of Gökova Geometry–Topology Conference 2007”, GGT, Gökova (2008) 60–81 [MR2509750](#)
- [32] **LI Nicolaescu**, *The Maslov index, the spectral flow, and decompositions of manifolds*, Duke Math. J. 80 (1995) 485–533 [MR1369400](#)
- [33] **P Ozsváth, Z Szabó**, *Holomorphic disks and knot invariants*, Adv. Math. 186 (2004) 58–116 [MR2065507](#)
- [34] **P Ozsváth, Z Szabó**, *On knot Floer homology and lens space surgeries*, Topology 44 (2005) 1281–1300 [MR2168576](#)
- [35] **T R Ramadas, I M Singer, J Weitsman**, *Some comments on Chern–Simons gauge theory*, Comm. Math. Phys. 126 (1989) 409–420 [MR1027504](#)
- [36] **J A Rasmussen**, *Floer homology and knot complements*, PhD Thesis, Harvard University (2003) Available at <http://search.proquest.com/docview/305332635>
- [37] **R Riley**, *Nonabelian representations of 2–bridge knot groups*, Quart. J. Math. Oxford Ser. 35 (1984) 191–208 [MR745421](#)
- [38] **P Turner**, *A spectral sequence for Khovanov homology with an application to $(3, q)$ –torus links*, Algebr. Geom. Topol. 8 (2008) 869–884 [MR2443099](#)
- [39] **K Wehrheim, C T Woodward**, *Functoriality for Lagrangian correspondences in Floer theory*, Quantum Topol. 1 (2010) 129–170 [MR2657646](#)
- [40] **A Weil**, *Remarks on the cohomology of groups*, Ann. of Math. 80 (1964) 149–157 [MR0169956](#)
- [41] **R Zentner**, *Representation spaces of pretzel knots*, Algebr. Geom. Topol. 11 (2011) 2941–2970 [MR2869448](#)

Department of Mathematics, Michigan State University
A338 WH, East Lansing, MI 48824, USA

Department of Mathematics, University of Nevada
1664 N. Virginia Street, Reno, NV 89557, USA

Department of Mathematics, Indiana University
Rawles Hall, 831 East 3 Street, Bloomington, IN 47405, USA

mhedden@math.msu.edu, herald@unr.edu, pkirk@indiana.edu

Proposed: Ronald Fintushel

Received: 5 January 2013

Seconded: Simon Donaldson, Ronald Stern

Accepted: 13 July 2013