

# involve

a journal of mathematics

On the size of the resonant set for the products of  $2 \times 2$   
matrices

Jeffrey Allen, Benjamin Seeger and Deborah Unger

 mathematical sciences publishers

2011

vol. 4, no. 2



# On the size of the resonant set for the products of $2 \times 2$ matrices

Jeffrey Allen, Benjamin Seeger and Deborah Unger

(Communicated by Chi-Kwong Li)

For  $\theta \in [0, 2\pi)$  and  $\lambda > 1$ , consider the matrix  $h = \begin{pmatrix} \lambda & 0 \\ 0 & 0 \end{pmatrix}$  and the rotation matrix  $R_\theta$ . Let  $W_n(\theta)$  denote some product of  $m$  instances of  $R_\theta$  and  $n$  of  $h$ , with the condition  $m \leq \epsilon n$  ( $0 < \epsilon < 1$ ). We analyze the measure of the set of  $\theta$  for which  $\|W_n(\theta)\| \geq \lambda^{\delta n}$  ( $0 < \delta < 1$ ). This can be regarded as a model problem for the Bochi–Fayad conjecture.

## 1. Introduction

Avila and Roblin [2009] considered the following problem. Take the two matrices

$$H = \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} \quad (1)$$

and

$$R_\theta = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}, \quad \theta \in [0, 2\pi).$$

Fix  $\lambda > 1$  and let  $m, n \in \mathbb{N}$ . Consider words of the form

$$W_n(\theta) = H^{i_1} R_\theta^{j_1} \dots H^{i_k} R_\theta^{j_k},$$

where  $k$  is arbitrary and  $i_1, \dots, i_k, j_1, \dots, j_k \in \mathbb{N} \cup \{0\}$  are such that

$$i_1 + \dots + i_k = n, \quad j_1 + \dots + j_k = m.$$

Assume that  $m$  is much smaller than  $n$  and take a “generic” angle  $\theta$ . It is not unreasonable to conjecture that  $\|W_n\|$  grows geometrically with  $n$  regardless of the combinatorics of the word. Avila and Roblin proved the following theorem, where the norm is given by  $\|W\| = |a| + |b| + |c| + |d|$  if  $W = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ .

*MSC2010:* primary 37H15; secondary 37H05, 37C85.

*Keywords:* Bochi–Fayad conjecture, resonant set, measure, rotation matrix, Fayad, Krikorian, exponential growth.

**Theorem 1.** *Assume that  $0 < \delta < 1$  is fixed. Then there is an  $n$ -independent set  $\Omega$  such that  $|\Omega| = 2\pi$  and for any  $\theta \in \Omega$  there is  $\epsilon > 0$  so that*

$$\min_{W_n} \|W_n(\theta)\| > \lambda^{\delta n}$$

provided  $m < \epsilon n (\ln n \ln \ln n)^{-1}$ .

Here the minimum is over all words  $W_n$  for  $n$  fixed and  $m$  as given. This theorem improved earlier results by Fayad and Krikorian [2008]. The special case of the Bochi–Fayad conjecture [Avila and Roblin 2009; Fayad and Krikorian 2008] deals with the similar situation when  $m < \epsilon n$  and  $\epsilon$  is small. One might expect that  $|\Omega| \rightarrow 2\pi$  as  $\epsilon \rightarrow 0$  in this case. Proving it seems to be quite hard. We investigate a simpler case. In (1), consider the matrix  $H$  when  $\lambda$  is large. Then  $\lambda^{-1} \rightarrow 0$  as  $\lambda \rightarrow \infty$  and one might wonder what happens if  $\lambda^{-1}$  is dropped. Thus we consider  $h = \begin{pmatrix} \lambda & 0 \\ 0 & 0 \end{pmatrix} \notin \mathrm{SL}(2, \mathbb{R})$  instead of  $H$ . It turns out that a very precise analysis can be performed for this simpler model problem, as we shall see in the next section. Section 3 provides some numerical evidence and comparison of the model case with the real problem.

## 2. The model problem

In the previous setting, take  $h = \begin{pmatrix} \lambda & 0 \\ 0 & 0 \end{pmatrix}$  instead of  $H$  and fix  $\epsilon \in (0, 1)$ . Given  $n$ , set

$$f_n(\theta) = \min_{W_n} \|W_n(\theta)\|,$$

where the norm is again given by the sum of absolute values of matrix entries and the minimum is taken over all  $W_n$  with  $m \leq \epsilon n$ . Note that we can take the minimum because for a given  $n$  there are only finitely many possibilities for  $W_n$ .

Finally, we fix  $0 < \delta < 1$  and define the *resonant set*  $\mathcal{R}$  thus:  $\theta \in \mathcal{R}$  if there exists some  $n$  such that  $f_n(\theta) < \lambda^{\delta n}$ . We claim that  $|\mathcal{R}| < C\lambda^{-(1-\delta)/\epsilon}$ , where  $C$  is some constant that can be explicitly computed and  $|\mathcal{R}|$  denotes the Lebesgue measure of the set  $\mathcal{R}$ .

We now make the convention that *there are no zero exponents in the expression of  $W_n(\theta)$* . Then, for words having precisely  $k$  blocks of rotation matrices, there are four possibilities, differing in which matrix ( $h$  or  $R_\theta$ ) begins the word and which matrix ends it:

$$W_n(\theta) = h^{i_1} R_\theta^{j_1} \cdots h^{i_k} R_\theta^{j_k}, \quad (2)$$

$$W_n(\theta) = R_\theta^{j_1} h^{i_1} \cdots R_\theta^{j_k} h^{i_k}, \quad (3)$$

$$W_n(\theta) = R_\theta^{j_1} h^{i_1} \cdots R_\theta^{j_{k-1}} h^{i_{k-1}} R_\theta^{j_k}, \quad (4)$$

$$W_n(\theta) = h^{i_1} R_\theta^{j_1} \cdots h^{i_k} R_\theta^{j_k} h^{i_{k+1}}. \quad (5)$$

For the word in (2), the product has this explicit form:

$$\begin{aligned}
 W_n(\theta) &= h^{i_1} R_\theta^{j_1} \dots h^{i_k} R_\theta^{j_k} \\
 &= \begin{pmatrix} \lambda^{i_1} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos j_1\theta & -\sin j_1\theta \\ \sin j_1\theta & \cos j_1\theta \end{pmatrix} \dots \begin{pmatrix} \lambda^{i_k} & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} \cos j_k\theta & -\sin j_k\theta \\ \sin j_k\theta & \cos j_k\theta \end{pmatrix} \\
 &= \begin{pmatrix} \lambda^{i_1} \cos j_1\theta & -\lambda^{i_1} \sin j_1\theta \\ 0 & 0 \end{pmatrix} \dots \begin{pmatrix} \lambda^{i_k} \cos j_k\theta & -\lambda^{i_k} \sin j_k\theta \\ 0 & 0 \end{pmatrix} \\
 &= \begin{pmatrix} \lambda^n \cos j_1\theta \dots \cos j_k\theta & \lambda^n \cos j_1\theta \dots \cos j_{k-1}\theta \sin j_k\theta \\ 0 & 0 \end{pmatrix}. \tag{6}
 \end{aligned}$$

Likewise, for (3), we obtain

$$\begin{aligned}
 W_n(\theta) &= R_\theta^{j_1} h^{i_1} \dots R_\theta^{j_k} h^{i_k} \\
 &= \begin{pmatrix} \cos j_1\theta & -\sin j_1\theta \\ \sin j_1\theta & \cos j_1\theta \end{pmatrix} \begin{pmatrix} \lambda^{i_1} & 0 \\ 0 & 0 \end{pmatrix} \dots \begin{pmatrix} \cos j_k\theta & -\sin j_k\theta \\ \sin j_k\theta & \cos j_k\theta \end{pmatrix} \begin{pmatrix} \lambda^{i_k} & 0 \\ 0 & 0 \end{pmatrix} \\
 &= \begin{pmatrix} \lambda^{i_1} \cos j_1\theta & 0 \\ \lambda^{i_1} \sin j_1\theta & 0 \end{pmatrix} \dots \begin{pmatrix} \lambda^{i_k} \cos j_k\theta & 0 \\ \lambda^{i_k} \sin j_k\theta & 0 \end{pmatrix} \\
 &= \begin{pmatrix} \lambda^n \cos j_1\theta \dots \cos j_k\theta & 0 \\ \lambda^n \sin j_1\theta \cos j_2\theta \dots \cos j_k\theta & 0 \end{pmatrix}. \tag{7}
 \end{aligned}$$

Using the result in (7), we get for the word (4)

$$\begin{aligned}
 W_n(\theta) &= (R_\theta^{j_1} h^{i_1} \dots R_\theta^{j_{k-1}} h^{i_{k-1}}) R_\theta^{j_k} \\
 &= \begin{pmatrix} \lambda^{i_1+\dots+i_{k-1}} \cos j_1\theta \cos j_2\theta \dots \cos j_{k-1}\theta & 0 \\ \lambda^{i_1+\dots+i_{k-1}} \sin j_1\theta \cos j_2\theta \dots \cos j_{k-1}\theta & 0 \end{pmatrix} \begin{pmatrix} \cos j_k\theta & -\sin j_k\theta \\ \sin j_k\theta & \cos j_k\theta \end{pmatrix} \\
 &= \begin{pmatrix} \lambda^n \cos j_1\theta \cos j_2\theta \dots \cos j_k\theta & -\lambda^n \cos j_1\theta \cos j_2\theta \dots \cos j_{k-1}\theta \sin j_k\theta \\ \lambda^n \sin j_1\theta \cos j_2\theta \dots \cos j_k\theta & -\lambda^n \sin j_1\theta \cos j_2\theta \dots \cos j_{k-1}\theta \sin j_k\theta \end{pmatrix}. \tag{8}
 \end{aligned}$$

Finally, using (6), we get for the word (5) simply

$$\begin{aligned}
 W_n(\theta) &= (h^{i_1} R_\theta^{j_1} \dots h^{i_{k-1}} R_\theta^{j_{k-1}} h^{i_k} R_\theta^{j_k}) h^{i_{k+1}} \\
 &= \begin{pmatrix} \lambda^{i_1+\dots+i_k} \cos j_1\theta \dots \cos j_k\theta & \lambda^{i_1+\dots+i_k} \cos j_1\theta \dots \cos j_{k-1}\theta \sin j_k\theta \\ 0 & 0 \end{pmatrix} \\
 &\hspace{25em} \times \begin{pmatrix} \lambda^{i_{k+1}} & 0 \\ 0 & 0 \end{pmatrix} \\
 &= \begin{pmatrix} \lambda^n \cos j_1\theta \dots \cos j_k\theta & 0 \\ 0 & 0 \end{pmatrix}. \tag{9}
 \end{aligned}$$

Therefore  $\|W_n(\theta)\|$  is given by

$$\begin{aligned} \lambda^n |\cos j_1 \theta \cdots \cos j_{k-1} \theta| (|\cos j_k \theta| + |\sin j_k \theta|) & \quad \text{for } W_n \text{ of type (2),} \\ \lambda^n |\cos j_2 \theta \cdots \cos j_k \theta| (|\cos j_1 \theta| + |\sin j_1 \theta|) & \quad \text{for } W_n \text{ of type (3),} \\ \lambda^n |\cos j_2 \theta \cdots \cos j_{k-1} \theta| (|\cos j_1 \theta| + |\sin j_1 \theta|) \\ & \quad \times (|\cos j_k \theta| + |\sin j_k \theta|) \quad \text{for } W_n \text{ of type (4),} \\ \lambda^n |\cos j_1 \theta \cdots \cos j_k \theta| & \quad \text{for } W_n \text{ of type (5).} \end{aligned}$$

**Remark.** This shows that, among words with  $k$  rotation blocks,  $\min_{W_n} \|W_n(\theta)\|$  is reached by words of type (5).

**Theorem 2.** *Let*

$$\begin{aligned} S_\alpha &= \{ \theta \in [0, 2\pi) \mid |\cos \alpha \theta| < \lambda^{-(1-\delta)\alpha/\epsilon-1} \}, \\ \tilde{S}_\alpha &= \{ \theta \in [0, 2\pi) \mid |\cos \alpha \theta| < \lambda^{-(1-\delta)\alpha/\epsilon} \}. \end{aligned}$$

*Then the resonant set  $\mathcal{R}$  satisfies*

$$\bigcup_{\alpha \in \mathbb{N}} S_\alpha \subseteq \mathcal{R} \subseteq \bigcup_{\alpha \in \mathbb{N}} \tilde{S}_\alpha.$$

*Proof.* Suppose  $\theta \in \bigcup_{\alpha \in \mathbb{N}} S_\alpha$ . Then  $\theta \in S_\alpha$  for some  $\alpha \in \mathbb{N}$  and

$$|\cos \alpha \theta| < \lambda^{-(1-\delta)\alpha/\epsilon-1}.$$

Let  $n = [\alpha/\epsilon] + 1$ . Then,  $n - 1 \leq \alpha/\epsilon < n$  and  $\alpha < \epsilon n$ . Consider the word  $\omega_n(\theta) = h^{i_1} R_\theta^\alpha h^{i_2}$  where  $i_1 + i_2 = n$ . Since  $m = \alpha$ , we have  $m \leq \epsilon n$ . Then

$$f_n(\theta) = \min_{W_n(\theta)} \|W_n(\theta)\| \leq \|\omega_n(\theta)\| = \lambda^n |\cos \alpha \theta| < \lambda^n \cdot \lambda^{-(1-\delta)\alpha/\epsilon-1} \leq \lambda^{\delta n}.$$

Therefore  $\theta \in \mathcal{R}$ .

Now suppose  $\theta \notin \bigcup_{\alpha \in \mathbb{N}} \tilde{S}_\alpha$ . Then  $|\cos \alpha \theta| \geq \lambda^{-(1-\delta)\alpha/\epsilon}$  for all  $\alpha \in \mathbb{N}$ . Choose an arbitrary  $n \in \mathbb{N}$ . Then

$$\begin{aligned} f_n(\theta) &= \min_{W_n(\theta)} \|W_n(\theta)\| = \|\omega_n(\theta)\| & \quad (\text{for some word } \omega_n(\theta)) \\ &= \lambda^n |\cos j_1 \theta \cdots \cos j_k \theta| & \quad (\text{by the remark above}) \\ &= \lambda^n |\cos \alpha_1 \theta^{m_1} \cdots \cos \alpha_l \theta^{m_l}|, \end{aligned}$$

where  $\alpha_1 < \cdots < \alpha_l$  and  $m_1 \alpha_1 + \cdots + m_l \alpha_l = m \leq \epsilon n$ . Then

$$\begin{aligned} f_n(\theta) &= \lambda^n |\cos \alpha_1 \theta^{m_1} \cdots \cos \alpha_l \theta^{m_l}| \\ &\geq \lambda^n \cdot \lambda^{-(1-\delta)(m_1 \alpha_1 + \cdots + m_l \alpha_l)/\epsilon} = \lambda^n \cdot \lambda^{-m(1-\delta)/\epsilon} \geq \lambda^{\delta n}, \end{aligned}$$

and therefore  $\theta \notin \mathcal{R}$ . □

We claim that  $\mathcal{R}$  is a dense open set. To show that  $\mathcal{R}$  is open, we show that for each  $n$ ,  $f_n$  is continuous. For each  $n$ ,

$$R_n = \{\theta \in [0, 2\pi) \mid f_n(\theta) < \lambda^{\delta n}\} = f_n^{-1}((-\infty, \lambda^{\delta n})),$$

which is open as the preimage of a continuous function of an open set. Note that  $\mathcal{R} = \bigcup_{n=1}^{\infty} R_n$ , a union of open sets, so  $\mathcal{R}$  is open.

To show that  $f_n$  is continuous, we note that  $f_n$  is the minimum of a finite number of continuous functions (the norms of a finite number of words). Denote these functions by  $F_1, F_2, \dots, F_M$ ,  $M \in \mathbb{N}$ . Fix arbitrary  $\theta \in [0, 2\pi)$ , fix  $\zeta > 0$ , and let  $\eta > 0$  be such that whenever  $|\theta - \tilde{\theta}| < \eta$ ,  $|F_k(\theta) - F_k(\tilde{\theta})| < \zeta$  for all  $k = 1, \dots, M$ . Consider arbitrary  $\tilde{\theta} \in (\theta - \eta, \theta + \eta)$ . For some  $i, j$ ,

$$f_n(\theta) = F_i(\theta) \quad \text{and} \quad f_n(\tilde{\theta}) = F_j(\tilde{\theta}).$$

By the definition of  $f_n$ ,

$$F_i(\theta) \leq F_j(\theta) \quad \text{and} \quad F_j(\tilde{\theta}) \leq F_i(\tilde{\theta}).$$

Notice that if  $F_i(\theta) = F_j(\tilde{\theta})$ , then  $|f_n(\theta) - f_n(\tilde{\theta})| = 0 < \zeta$  and we are done. Suppose that  $F_i(\theta) > F_j(\tilde{\theta})$ . Then  $|f_n(\theta) - f_n(\tilde{\theta})| = F_i(\theta) - F_j(\tilde{\theta}) \leq F_j(\theta) - F_j(\tilde{\theta}) < \zeta$ . Otherwise, if  $F_i(\theta) < F_j(\tilde{\theta})$ , then

$$|f_n(\theta) - f_n(\tilde{\theta})| = F_j(\tilde{\theta}) - F_i(\theta) \leq F_i(\tilde{\theta}) - F_i(\theta) < \zeta.$$

To see that  $\mathcal{R}$  is dense, let  $I$  be any open interval in  $[0, 2\pi)$ . The collection of points  $R_\alpha = \{\pi/2\alpha + (\pi/\alpha)k : k \in \{1, \dots, 2\alpha - 1\}\}$  is in  $S_\alpha$ ; indeed, for any  $\phi \in R_\alpha$ ,  $\cos \alpha\phi = 0 < \lambda^{-(1-\delta)\alpha/\epsilon-1}$ . If we choose  $\alpha > |I|/\pi$ , then there must be some element  $\phi$  of  $R_\alpha$  in  $I$ . Since  $\phi \in \bigcup_{\alpha \in \mathbb{N}} S_\alpha \subseteq \mathcal{R} \subseteq \bigcup_{\alpha \in \mathbb{N}} \tilde{S}_\alpha$ , we see that every open interval in  $[0, 2\pi)$  contains a point in  $\mathcal{R}$ .

Now we are ready to estimate the size of  $\mathcal{R}$ . Consider  $\tilde{S}_\alpha$  for arbitrary  $\alpha \in \mathbb{N}$ . The measure of this set is

$$\begin{aligned} |\tilde{S}_\alpha| &= 4\alpha \left( \frac{\pi}{2\alpha} - \frac{1}{\alpha} \cos^{-1}(\lambda^{-(1-\delta)\alpha/\epsilon}) \right) \\ &= 2\pi - 4 \cos^{-1}(\lambda^{-(1-\delta)\alpha/\epsilon}) \\ &\approx 2\pi - 2\pi + 4\lambda^{-(1-\delta)\alpha/\epsilon} \\ &= 4\lambda^{-(1-\delta)\alpha/\epsilon}. \end{aligned}$$

Then our estimate for the size of  $\mathcal{R}$  is

$$|\mathcal{R}| \leq \left| \bigcup_{\alpha \in \mathbb{N}} \tilde{S}_\alpha \right| \leq \sum_{\alpha=1}^{\infty} |\tilde{S}_\alpha| \approx 4 \sum_{\alpha=1}^{\infty} \lambda^{-(1-\delta)\alpha/\epsilon} = \frac{4\lambda^{-(1-\delta)/\epsilon}}{1 - \lambda^{-(1-\delta)/\epsilon}} \approx 4\lambda^{-(1-\delta)/\epsilon},$$


as  $\epsilon \sim 0$  and  $\lambda, \delta$  are fixed.

### 3. Some numerical evidence


We provide some numerical and graphical evidence of what was proved. We see how the graphs of the model case and the real case compare for fixed  $n$  and  $m$ . In addition, it is shown graphically that changing the multiplicities of  $H$  affects the word's norm, but in the model case the word's norm is invariant under these changes. The graphs in this section were plotted with Maple 14.

Based on the similarities between the pictures of the real case and the model case, we conjecture that the resonant set in the model case is, in some sense, the limiting set of the resonant set in the real case as  $\lambda$  grows large, since the  $\lambda^{-1}$  term goes to 0 as  $\lambda$  goes to infinity. Of course, since here we take  $\lambda$  relatively small ( $\lambda = 2$ ) for graphing convenience, this is a rough conjecture; in fact, proving it seems to be rather difficult.

Figure 1 shows a red curve and a blue curve. The blue curve is the graph of  $\|h^{i_1} R_\theta^2 h^{i_2} R_\theta^3 h^{i_3}\|$  where  $i_1, i_2, i_3 \in \mathbb{N}$  and  $i_1 + i_2 + i_3 = 15$ . Recall that  $h$  is the matrix we use in the model case, where  $\lambda^{-1}$  is replaced by 0. With these combinatorics, varying  $i_1, i_2, i_3$  does not change the graph, as long as their sum is 15. Specifically, Figure 1 is the model case of  $n = 15, m = 5$  ( $j_1 = 2, j_2 = 3$ ), and  $\lambda = 2$ .



**Figure 1.** Graphs of the functions  $\|H^5 R_\theta^2 H^5 R_\theta^3 H^5\|$  (red curve) and  $\|h^5 R_\theta^2 h^5 R_\theta^3 h^5\|$  (blue curve) when  $\lambda = 2$ . (The curves have been slightly offset horizontally; otherwise they would coincide at this resolution.) The thin black curve on the bottom right is the difference between the first and second functions, with the y-axis expanded 1000 times.



**Figure 2.** Graphs of the functions  $\|H^5 R_\theta^2 H^9 R_\theta^3 H^1\|$  (red curve) and  $\|h^5 R_\theta^2 h^9 R_\theta^3 h^1\|$  (blue curve) when  $\lambda = 2$ .

The red curve in [Figure 1](#) shows the case where we replace  $h$  with  $H$  and set  $i_1 = i_2 = i_3 = 5$ . Explicitly, we are graphing  $\|H^5 R_\theta^2 H^5 R_\theta^3 H^5\|$ .


In [Figure 2](#) we change only two parameters relative to [Figure 1](#): the lengths of the last two blocks of  $H$ 's (or  $h$ 's) are now  $i_2 = 9$ , and  $i_3 = 1$ . As already seen, the  $h$  curve (blue) remains the same, but the  $H$  curve (red)—that is, the graph of the function  $\|H^5 R_\theta^2 H^9 R_\theta^3 H^1\|$ —changes significantly as a result of changing the order of multiplication in the word.

By comparing the red curves in [Figures 1](#) and [2](#), we observe that a greater disparity between the multiplicities of  $H$  (the  $i_k$ 's) is correlated with a smaller resonant set (the set of points  $\theta$  between 0 and  $2\pi$  such that the norm of the word is within a certain distance of zero). The slope of the word's norm is steeper in [Figure 2](#) than it is in [Figure 1](#) and the peaks in [Figure 2](#) are associated with larger values of the word's norm than in the case depicted by [Figure 1](#). Both conditions lead to fewer points  $\theta$  that are mapped to a norm of the word that is close to zero.

[Figure 3](#) shows the graph of  $\|H^1 R_\theta^2 H^1 R_\theta^3 H^{13}\|$ . Comparing this graph with [Figure 2](#) provides further evidence that a greater disparity between the multiplicities of  $H$  results in a smaller resonant set.

To further justify our use of the model case, consider the [Figure 4](#), which treats the case of a word of the form (4). Specifically, the blue curve shows the function  $\|R_\theta h^{i_1} R_\theta h^{i_2} R_\theta\|$ , where  $i_1 + i_2 = 15$  and  $\lambda = 2$ . We take  $i_1 = 7$  and  $i_2 = 8$  and replace





**Figure 3.** Graphs of the functions  $\|H^1 R_\theta^2 H^1 R_\theta^3 H^{13}\|$  (red curve) and  $\|h^1 R_\theta^2 h^1 R_\theta^3 h^{13}\|$  (blue curve) when  $\lambda = 2$ .

$h$  by  $H$  to obtain the red curve. As in Figure 1, the two curves are indistinguishable to within the plot’s resolution.

Note that the blue curve in Figure 4 is not comparable to that of Figures 1–3. Both show the model case, but with different combinatorics on the word: expression (5) for the earlier figures, and (4) for Figure 4. Both graphs still have a small resonant set.


Figure 5 shows the graphs of  $\|R_\theta H^{14} R_\theta H R_\theta\|$  (red) and of  $\|R_\theta h^{14} R_\theta h R_\theta\|$  (blue); the latter of course is the same as the blue curve of Figure 4. Comparing Figure 4 with Figure 5, again we see that a greater disparity between the multiplicities of  $H$  results in a smaller resonant set.

### 4. Conclusion


We hope that our model problem is a viable approximation for what happens when the matrix  $H$  is used. The next step might be to express the the Bochi–Fayad problem in terms of the model problem. One way to do this might be to write  $H = h + e$ , where  $e$  is the matrix

$$e = \begin{pmatrix} 0 & 0 \\ 0 & \lambda^{-1} \end{pmatrix},$$

and then express a product of  $H$ ’s and  $R_\theta$ ’s in terms of a product of  $h$ ’s and  $R_\theta$ ’s, and some other, hopefully small, error terms. The numerical evidence above suggests



**Figure 4.** Graphs of the functions  $\|R_\theta H^7 R_\theta H^8 R_\theta\|$  (red curve) and  $\|R_\theta h^7 R_\theta h^8 R_\theta\|$  (blue curve) when  $\lambda = 2$ . (The curves have been slightly offset horizontally; otherwise they would coincide at this resolution.) The thin black curve on the bottom right is the difference between the first and second functions, with the y-axis expanded 2500 times.



**Figure 5.** Graphs of  $\|R_\theta h^{14} R_\theta h R_\theta\|$  and  $\|R_\theta H^{14} R_\theta H R_\theta\|$  when  $\lambda = 2$ .

that the resonant set for words using the matrix  $H$  would actually be smaller than that for words using the matrix  $h$ , especially if the distribution of  $H$ 's is in some sense irregular. This behavior might become more apparent when  $\lambda$  is taken much larger, and  $\epsilon$  much smaller, than in the experiments above.

### Acknowledgements

We would like to thank Professors Serguei Denisov and Alexander Kiselev for their constant support throughout the research and writing processes. We are grateful to them for suggesting the topic and providing us with the opportunity to apply the concepts we have learned throughout our mathematical careers. We thank them both for their excellent advice and encouragement.

### References

- [Avila and Roblin 2009] A. Avila and T. Roblin, “Uniform exponential growth for some  $SL(2, \mathbb{R})$  matrix products”, *J. Mod. Dyn.* **3**:4 (2009), 549–554. [MR 2011f:37099](#) [Zbl 1189.37060](#)
- [Fayad and Krikorian 2008] B. Fayad and R. Krikorian, “Exponential growth of product of matrices in  $SL(2, \mathbb{R})$ ”, *Nonlinearity* **21**:2 (2008), 319–323. [MR 2009d:37041](#) [Zbl 1142.37024](#)

Received: 2010-12-10

Revised: 2011-02-17

Accepted: 2011-04-03

[jhallen2@wisc.edu](mailto:jhallen2@wisc.edu)

*University Of Wisconsin – Madison, Madison, WI 53706,  
United States*

[bseeger@wisc.edu](mailto:bseeger@wisc.edu)

*University Of Wisconsin – Madison, Madison, WI 53706,  
United States*

[dunger@wisc.edu](mailto:dunger@wisc.edu)

*University Of Wisconsin – Madison, Madison, WI 53715,  
United States*

## EDITORS

### MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, [berenhks@wfu.edu](mailto:berenhks@wfu.edu)

### BOARD OF EDITORS

John V. Baxley	Wake Forest University, NC, USA <a href="mailto:baxley@wfu.edu">baxley@wfu.edu</a>	Chi-Kwong Li	College of William and Mary, USA <a href="mailto:ckli@math.wm.edu">ckli@math.wm.edu</a>
Arthur T. Benjamin	Harvey Mudd College, USA <a href="mailto:benjamin@hmc.edu">benjamin@hmc.edu</a>	Robert B. Lund	Clemson University, USA <a href="mailto:lund@clemson.edu">lund@clemson.edu</a>
Martin Bohner	Missouri U of Science and Technology, USA <a href="mailto:bohner@mst.edu">bohner@mst.edu</a>	Gaven J. Martin	Massey University, New Zealand <a href="mailto:g.j.martin@massey.ac.nz">g.j.martin@massey.ac.nz</a>
Nigel Boston	University of Wisconsin, USA <a href="mailto:boston@math.wisc.edu">boston@math.wisc.edu</a>	Mary Meyer	Colorado State University, USA <a href="mailto:meyer@stat.colostate.edu">meyer@stat.colostate.edu</a>
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA <a href="mailto:budhiraj@email.unc.edu">budhiraj@email.unc.edu</a>	Emil Minchev	Ruse, Bulgaria <a href="mailto:eminchev@hotmail.com">eminchev@hotmail.com</a>
Pietro Cerone	Victoria University, Australia <a href="mailto:pietro.cerone@vu.edu.au">pietro.cerone@vu.edu.au</a>	Frank Morgan	Williams College, USA <a href="mailto:frank.morgan@williams.edu">frank.morgan@williams.edu</a>
Scott Chapman	Sam Houston State University, USA <a href="mailto:scott.chapman@shsu.edu">scott.chapman@shsu.edu</a>	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran <a href="mailto:moslehian@ferdowsi.um.ac.ir">moslehian@ferdowsi.um.ac.ir</a>
Jem N. Corcoran	University of Colorado, USA <a href="mailto:corcoran@colorado.edu">corcoran@colorado.edu</a>	Zuhair Nashed	University of Central Florida, USA <a href="mailto:znashed@mail.ucf.edu">znashed@mail.ucf.edu</a>
Michael Dorff	Brigham Young University, USA <a href="mailto:mdorff@math.byu.edu">mdorff@math.byu.edu</a>	Ken Ono	University of Wisconsin, USA <a href="mailto:ono@math.wisc.edu">ono@math.wisc.edu</a>
Sever S. Dragomir	Victoria University, Australia <a href="mailto:sever@matilda.vu.edu.au">sever@matilda.vu.edu.au</a>	Joseph O'Rourke	Smith College, USA <a href="mailto:orourke@cs.smith.edu">orourke@cs.smith.edu</a>
Behrouz Emamizadeh	The Petroleum Institute, UAE <a href="mailto:bemamizadeh@pi.ac.ae">bemamizadeh@pi.ac.ae</a>	Yuval Peres	Microsoft Research, USA <a href="mailto:peres@microsoft.com">peres@microsoft.com</a>
Errin W. Fulp	Wake Forest University, USA <a href="mailto:fulp@wfu.edu">fulp@wfu.edu</a>	Y.-F. S. Pétermann	Université de Genève, Switzerland <a href="mailto:petermann@math.unige.ch">petermann@math.unige.ch</a>
Andrew Granville	Université Montréal, Canada <a href="mailto:andrew@dms.umontreal.ca">andrew@dms.umontreal.ca</a>	Robert J. Plemmons	Wake Forest University, USA <a href="mailto:rplemmons@wfu.edu">rplemmons@wfu.edu</a>
Jerrold Griggs	University of South Carolina, USA <a href="mailto:griggs@math.sc.edu">griggs@math.sc.edu</a>	Carl B. Pomerance	Dartmouth College, USA <a href="mailto:carl.pomerance@dartmouth.edu">carl.pomerance@dartmouth.edu</a>
Ron Gould	Emory University, USA <a href="mailto:rg@mathcs.emory.edu">rg@mathcs.emory.edu</a>	Bjorn Poonen	UC Berkeley, USA <a href="mailto:poonen@math.berkeley.edu">poonen@math.berkeley.edu</a>
Sat Gupta	U of North Carolina, Greensboro, USA <a href="mailto:sgupta@uncg.edu">sgupta@uncg.edu</a>	James Propp	U Mass Lowell, USA <a href="mailto:jpropp@cs.uml.edu">jpropp@cs.uml.edu</a>
Jim Haglund	University of Pennsylvania, USA <a href="mailto:jhaglund@math.upenn.edu">jhaglund@math.upenn.edu</a>	József H. Przytycki	George Washington University, USA <a href="mailto:przytyck@gwu.edu">przytyck@gwu.edu</a>
Johnny Henderson	Baylor University, USA <a href="mailto:johnny_henderson@baylor.edu">johnny_henderson@baylor.edu</a>	Richard Rebarber	University of Nebraska, USA <a href="mailto:rrebarbe@math.unl.edu">rrebarbe@math.unl.edu</a>
Natalia Hritonenko	Prairie View A&M University, USA <a href="mailto:nahritonenko@pvamu.edu">nahritonenko@pvamu.edu</a>	Robert W. Robinson	University of Georgia, USA <a href="mailto:rwr@cs.uga.edu">rwr@cs.uga.edu</a>
Charles R. Johnson	College of William and Mary, USA <a href="mailto:crjohnso@math.wm.edu">crjohnso@math.wm.edu</a>	Filip Saidak	U of North Carolina, Greensboro, USA <a href="mailto:f_saidak@uncg.edu">f_saidak@uncg.edu</a>
Karen Kafadar	University of Colorado, USA <a href="mailto:karen.kafadar@cudenver.edu">karen.kafadar@cudenver.edu</a>	Andrew J. Sterge	Honorary Editor <a href="mailto:andy@ajsterge.com">andy@ajsterge.com</a>
K. B. Kulasekera	Clemson University, USA <a href="mailto:kk@ces.clemson.edu">kk@ces.clemson.edu</a>	Ann Trenk	Wellesley College, USA <a href="mailto:atrenk@wellesley.edu">atrenk@wellesley.edu</a>
Gerry Ladas	University of Rhode Island, USA <a href="mailto:gladas@math.uri.edu">gladas@math.uri.edu</a>	Ravi Vakil	Stanford University, USA <a href="mailto:vakil@math.stanford.edu">vakil@math.stanford.edu</a>
David Larson	Texas A&M University, USA <a href="mailto:larson@math.tamu.edu">larson@math.tamu.edu</a>	Ram U. Verma	University of Toledo, USA <a href="mailto:verma99@msn.com">verma99@msn.com</a>
Suzanne Lenhart	University of Tennessee, USA <a href="mailto:lenhart@math.utk.edu">lenhart@math.utk.edu</a>	John C. Wierman	Johns Hopkins University, USA <a href="mailto:wierman@jhu.edu">wierman@jhu.edu</a>

## PRODUCTION

Silvio Levy, Scientific Editor

Sheila Newbery, Senior Production Editor

Cover design: ©2008 Alex Scorpan

See inside back cover or <http://pjm.math.berkeley.edu/involve> for submission instructions.

The subscription price for 2011 is US \$100/year for the electronic version, and \$130/year (+\$35 shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94704-3840, USA.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, Department of Mathematics, University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW™ from Mathematical Sciences Publishers.

PUBLISHED BY  
 **mathematical sciences publishers**  
<http://msp.org/>

A NON-PROFIT CORPORATION

Typeset in L<sup>A</sup>T<sub>E</sub>X

Copyright ©2011 by Mathematical Sciences Publishers

# involve

2011

vol. 4

no. 2

<a href="#">The visual boundary of <math>\mathbb{Z}^2</math></a>	103
KYLE KITZMILLER AND MATT RATHBUN	
<a href="#">An observation on generating functions with an application to a sum of secant powers</a>	117
JEFFREY MUDROCK	
<a href="#">Clique-relaxed graph coloring</a>	127
CHARLES DUNN, JENNIFER FIRKINS NORDSTROM, CASSANDRA NAYMIE, ERIN PITNEY, WILLIAM SEHORN AND CHARLIE SUER	
<a href="#">Cost-conscious voters in referendum elections</a>	139
KYLE GOLENBIEWSKI, JONATHAN K. HODGE AND LISA MOATS	
<a href="#">On the size of the resonant set for the products of <math>2 \times 2</math> matrices</a>	157
JEFFREY ALLEN, BENJAMIN SEEGER AND DEBORAH UNGER	
<a href="#">Continuous <math>p</math>-Bessel mappings and continuous <math>p</math>-frames in Banach spaces</a>	167
MOHAMMAD HASAN FAROUGH AND ELNAZ OSGOOEI	
<a href="#">The multidimensional Frobenius problem</a>	187
JEFFREY AMOS, IULIANA PASCU, VADIM PONOMARENKO, ENRIQUE TREVIÑO AND YAN ZHANG	
<a href="#">The Gauss–Bonnet formula on surfaces with densities</a>	199
IVAN CORWIN AND FRANK MORGAN	