

involve

a journal of mathematics

Matrix coefficients of depth-zero supercuspidal
representations of $GL(2)$

Andrew Knightly and Carl Ragsdale



Matrix coefficients of depth-zero supercuspidal representations of $GL(2)$

Andrew Knightly and Carl Ragsdale

(Communicated by Michael E. Zieve)

We give explicit formulas for matrix coefficients of the depth-zero supercuspidal representations of $GL(2)$ over a nonarchimedean local field, highlighting the case where the test vector is a unit new vector. We also describe the partition of the set of such representations according to central character, and compute sums of matrix coefficients over all representations in a given class.

Introduction

Let F be a nonarchimedean local field with integer ring \mathfrak{o} , maximal ideal $\mathfrak{p} = \varpi \mathfrak{o}$, and residue field $k = \mathfrak{o}/\mathfrak{p}$ of cardinality q . The supercuspidal representations of $GL_2(F)$ are precisely those irreducible admissible representations which do not arise as constituents of parabolic induction. They are characterized by having matrix coefficients which are compactly supported modulo the center.

In this paper we explicitly compute the matrix coefficients of depth-zero supercuspidal representations. These are the supercuspidals with the smallest possible conductor exponent, namely 2. First discovered by Mautner [1964, Section 9], they arise by compact induction from the $(q - 1)$ -dimensional representations of $GL_2(\mathfrak{o})$ inflated from the cuspidal series of the finite group $GL_2(k)$.

In the first section, we show that the matrix coefficients of any supercuspidal representation are expressible in terms of those of the finite-dimensional inducing representation. Thus, the task at hand essentially reduces to a computation of the matrix coefficients of the cuspidal representations of $GL_2(k)$. The latter is achieved in [Theorem 2.7](#) using the explicit model from [Piatetski-Shapiro 1983].

With global applications in mind, in [Section 3](#) we single out the case where the test vector in the supercuspidal matrix coefficient is a unit new vector. The resulting function, given in [\(3-6\)](#) and [Theorem 3.2](#), may be used to define an integral operator on the global automorphic spectrum of GL_2 which isolates those cuspidal

MSC2010: 22E50.

Keywords: supercuspidal, matrix coefficients, regular characters.

newforms with depth-zero supercuspidal local type (see [Section 3.4](#)). Possible applications include various trace formulas involving newforms of level p^2 which are supercuspidal (as opposed to principal series or special) at p . For a recent example involving the supercuspidal representations of conductor p^3 , see [\[Knightly and Li 2012\]](#).

It is often desirable to organize representations according to central character. For example, there are exactly $2(q - 1)$ supercuspidal representations of $\mathrm{GL}_2(F)$ of conductor p^3 with a given central character (see [\[Bushnell and Henniart 2014, Remark 2.2\]](#)). By contrast, there are $(q/2)(q - 1)$ distinct cuspidal representations of $\mathrm{GL}_2(k)$ and $(q - 1)$ possible central characters, but obviously there cannot be $q/2$ of each kind if q is odd. We sort this out in [Proposition 2.3](#), and use it to give a formula [\(4-1\)](#) for the number of supercuspidals of conductor p^2 with a given central character. It depends on the parity of q and the order of the central character. We then give formulas for various sums of matrix coefficients over the set of depth-zero supercuspidal representations with a given central character. These computations rely on sum formulas for primitive characters, derived in [Proposition 2.4](#). We close in the final section with some simple examples.

1. Matrix coefficients of supercuspidal representations

In this section let $G = \mathrm{GL}_n(F)$, or more generally, any unimodular locally profinite group [\[Bushnell and Henniart 2006\]](#) with center Z . Let $H \subset G$ be an open and closed subgroup containing Z with H/Z compact, and let (ρ, V) be an irreducible smooth representation of H . Consider the compact induction $\pi = \mathrm{c}\text{-Ind}_H^G(\rho)$. It consists of the functions $\phi : G \rightarrow V$ with compact support (mod Z) for which $\phi(hg) = \rho(h)\phi(g)$ for all $h \in H, g \in G$, with G acting on the space by right translation. Here we show that, as observed by Mautner [\[1964\]](#), the matrix coefficients of π are essentially those of ρ . These matrix coefficients are compactly supported (modulo Z), so if π is irreducible and admissible, it is supercuspidal. Conversely, it is conjectured that all supercuspidal representations arise in this way. This was proven by Bushnell and Kutzko [\[1993\]](#) for $G = \mathrm{GL}_n(F)$, and more recently in great (but not complete) generality in [\[Stevens 2008; Kim 2007\]](#).

We assume for simplicity that ρ has unitary central character, so that by the fact that H/Z is compact, ρ is unitarizable. Let $\langle v, w \rangle_V$ denote an H -equivariant inner product on V . Then the inner product on $\mathrm{c}\text{-Ind}_H^G(\rho)$ given by

$$\langle \phi, \psi \rangle = \sum_{x \in H \backslash G} \langle \phi(x), \psi(x) \rangle_V$$

is convergent (in fact a finite sum) and well-defined. Further, for any $g \in G$,

$$\langle \pi(g)\phi, \pi(g)\psi \rangle = \sum_{x \in H \backslash G} \langle \phi(xg), \psi(xg) \rangle_V = \sum_{x \in H \backslash G} \langle \phi(x), \psi(x) \rangle_V = \langle \phi, \psi \rangle.$$

Thus π is unitary relative to this inner product.

For $v \in V$ and $y \in G$, define a function $f_{y,v} \in \mathfrak{c}\text{-Ind}_H^G(\rho)$ by

$$f_{y,v}(g) = \begin{cases} \rho(h)v & \text{if } g = hy \in Hy, \\ 0 & \text{if } g \notin Hy. \end{cases}$$

Then the set $\{f_{y,v} \mid y \in H \backslash G, v \in V\}$ spans the space $\mathfrak{c}\text{-Ind}_H^G(\rho)$. (Note that $f_{hy,v} = f_{y,\rho(h^{-1}v)}$.)

Proposition 1.1. *For $y, z \in G$ and $v, w \in V$,*

$$\langle \pi(g)f_{y,v}, f_{z,w} \rangle = \begin{cases} \langle \rho(h)v, w \rangle_V & \text{if } g = z^{-1}hy \in z^{-1}Hy, \\ 0 & \text{if } g \notin z^{-1}Hy. \end{cases}$$

Proof. By definition of the inner product,

$$\begin{aligned} \langle \pi(g)f_{y,v}, f_{z,w} \rangle &= \sum_{x \in H \backslash G} \langle \pi(g)f_{y,v}(x), f_{z,w}(x) \rangle_V \\ &= \sum_{x \in H \backslash G} \langle f_{y,v}(xg), f_{z,w}(x) \rangle_V \\ &= \langle f_{y,v}(zg), w \rangle_V, \end{aligned}$$

since $f_{z,w}(x)$ vanishes unless $x \in Hz$. If $g = z^{-1}hy \in z^{-1}Hy$, then the above is equal to

$$\langle f_{y,v}(hy), w \rangle_V = \langle \rho(h)v, w \rangle_V,$$

as needed. If $g \notin z^{-1}Hy$, then $zg \notin Hy$, so $f_{y,v}(zg) = 0$ and the inner product vanishes. □

If we let $\bar{G} = G/Z$, then the formal degree d_π of π is a positive constant satisfying

$$\int_{\bar{G}} |\langle \pi(g)f, f \rangle|^2 dg = \frac{\|f\|^4}{d_\pi} \tag{1-1}$$

for all $f \in \mathfrak{c}\text{-Ind}_H^G(\rho)$. It depends on the choice of Haar measure on \bar{G} . (The existence of d_π is due to Godement; see, for example, [Knightly and Li 2006, Proposition 10.4].)

Proposition 1.2. *For any choice of Haar measure on $\bar{G} = G/Z$, the associated formal degree of π is given by*

$$d_\pi = \frac{\dim \rho}{\text{meas}(\bar{H})},$$

where \bar{H} is the (open) image of H in \bar{G} .

Proof. Let $v \in V$ be a unit vector, and consider the function $f_{1,v}$. By [Proposition 1.1](#),

$$\langle \pi(g) f_{1,v}, f_{1,v} \rangle = \begin{cases} \langle \rho(g)v, v \rangle_V & \text{if } g \in H, \\ 0 & \text{if } g \notin H. \end{cases}$$

Therefore,

$$\begin{aligned} \int_{\bar{G}} |\langle \pi(g) f_{1,v}, f_{1,v} \rangle|^2 dg &= \int_{\bar{H}} |\langle \rho(g)v, v \rangle_V|^2 dg \\ &= \frac{\|v\|^4}{\dim(\rho)} \text{meas}(\bar{H}) = \frac{\text{meas}(\bar{H})}{\dim(\rho)}, \end{aligned}$$

by the Schur orthogonality relations for irreducible representations of compact groups. By [\(1-1\)](#),

$$\frac{\text{meas}(\bar{H})}{\dim(\rho)} = \frac{\|f_{1,v}\|^4}{d_\pi}.$$

Therefore, it suffices to show that $\|f_{1,v}\| = 1$. This can be done via a direct computation:

$$\begin{aligned} \|f_{1,v}\|^2 = \langle f_{1,v}, f_{1,v} \rangle &= \sum_{x \in H \setminus G} \langle f_{1,v}(x), f_{1,v}(x) \rangle_V \\ &= \langle f_{1,v}(1), f_{1,v}(1) \rangle_V = \langle v, v \rangle = 1, \end{aligned}$$

as needed. □

2. Cuspidal representations of $\text{GL}_2(k)$

Let q be a prime power, let k be the finite field with q elements, let L be the unique quadratic extension of k , and let $G = \text{GL}_2(k)$. Define the subgroups

$$U = \left\{ \begin{pmatrix} 1 & b \\ 0 & 1 \end{pmatrix} \in G \right\}, \quad Z = \left\{ \begin{pmatrix} a & \\ & a \end{pmatrix} \in G \right\}, \quad B = \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \in G \right\}.$$

Note that Z is the center of G . Recall that the cuspidal representations of G are those that do not contain the trivial character of the unipotent subgroup U . These are precisely the irreducible representations that do not arise via parabolic induction. They have dimension $q - 1$, and are parametrized by the Galois orbits of the primitive characters of L^* , defined below.

2.1. Primitive characters of L^* . For any finite abelian group H , we write \widehat{H} for the dual group, consisting of the characters $\chi : H \rightarrow \mathbb{C}^*$. We recall that

$$H \cong \widehat{\widehat{H}}.$$

Thus $\widehat{L^*}$ is a cyclic group of order $q^2 - 1$. A character $\nu \in \widehat{L^*}$ is *primitive* if $\nu^q \neq \nu$. Otherwise, ν is *imprimitive*. Letting $\bar{\alpha} = \alpha^q$ denote the Frobenius map, the norm

map $N : L \rightarrow k$ is given by

$$N(\alpha) = \alpha\bar{\alpha} = \alpha^{q+1}.$$

Proposition 2.1. *Let $\nu : L^* \rightarrow \mathbb{C}^*$ be a character of L^* . Then the following are equivalent:*

- (i) ν is imprimitive; that is, $\nu^q = \nu$.
- (ii) $\nu = \chi \circ N$ for some $\chi \in \widehat{k^*}$.
- (iii) $L^1 \subset \ker(\nu)$, where L^1 is the subgroup of norm 1 elements of L^* .

Proof. Let θ be a generator of the cyclic group L^* . Then θ^{q+1} is a generator of k^* . If $\nu^q = \nu$, then $\nu(\theta)$ is a $(q - 1)$ -st root of unity, so we may define $\chi \in \widehat{k^*}$ by $\chi(\theta^{q+1}) = \nu(\theta)$. Then $\nu = \chi \circ N$, so (i) implies (ii). It is clear that (ii) implies (iii). On the other hand, for any $x \in L^*$, $N(x^{q-1}) = N(x)^{q-1} = 1$, so that $x^{q-1} \in L^1$. Therefore if (iii) holds, $\nu^q(x) = \nu(x^q) = \nu(x^{q-1}x) = \nu(x^{q-1})\nu(x) = \nu(x)$. Hence (iii) implies (i). □

The imprimitive characters thus correspond bijectively with the characters of k^* , so there are $q - 1$ of them. It follows that there are $(q^2 - 1) - (q - 1) = q^2 - q$ primitive characters of L^* .

Lemma 2.2. *Let ν be a primitive character of L^* . Then, for all $\alpha \in k^*$,*

$$\sum_{\substack{x \in L^* \\ N(x) = \alpha}} \nu(x) = 0.$$

Proof. By Proposition 2.1, there exists $\lambda \in L^1$ such that $\nu(\lambda) \neq 1$. Thus,

$$\sum_{N(x) = \alpha} \nu(x) = \sum_{N(x) = \alpha} \nu(\lambda x) = \nu(\lambda) \sum_{N(x) = \alpha} \nu(x).$$

It follows that $\sum_{N(x) = \alpha} \nu(x) = 0$. □

Next, we examine the partition of primitive characters into classes according to their restrictions to k^* . This will allow us to count the number of depth-zero supercuspidal representations with a given central character (see (4-1)).

Proposition 2.3. *Suppose ω is a given character of k^* . Let P_ω denote the number of primitive characters ν of L^* for which $\nu|_{k^*} = \omega$. Then*

$$P_\omega = \begin{cases} q - 1 & \text{if } q \text{ is odd and } \omega^{(q-1)/2} \text{ is trivial,} \\ q + 1 & \text{if } q \text{ is odd and } \omega^{(q-1)/2} \text{ is nontrivial,} \\ q & \text{if } q \text{ is even.} \end{cases}$$

Proof. Let ξ be a generator of the cyclic group $\widehat{L^*}$. Let $\nu_0 = \xi^{q-1}$. Note that for $\alpha \in k^*$,

$$\nu_0(\alpha) = \xi(\alpha^{q-1}) = \xi(1) = 1.$$

In fact, ν_0 is a generator of the order $q + 1$ subgroup $\widehat{L^*/k^*}$ of $\widehat{L^*}$. Let us consider two characters of L^* to be equivalent if they have the same restriction to k^* . Then the equivalence class of a given character ν is the set

$$\{\nu, \nu\nu_0, \nu\nu_0^2, \dots, \nu\nu_0^q\}. \quad (2-1)$$

If $\omega = \nu|_{k^*}$, then $P_\omega = q + 1 - A_\omega$, where A_ω is the number of imprimitive elements of the above set. Write $\nu = \xi^b$. Without loss of generality (replacing ν by some $\nu\nu_0^m$), we may assume that $0 \leq b < q - 1$. A character ξ^a is imprimitive if and only if $\xi^{qa} = \xi^a$, or equivalently, $(q + 1) \mid a$. Suppose $\nu\nu_0^s$ and $\nu\nu_0^k$ are both imprimitive for $0 \leq s < k < q$. Then $b + (q - 1)s$ and $b + (q - 1)k$ are both divisible by $q + 1$ and strictly less than $q^2 - 1$. Their difference $(k - s)(q - 1) \geq 0$ also has these properties, and furthermore it is divisible by

$$\text{lcm}(q - 1, q + 1) = \begin{cases} (q^2 - 1)/2 & \text{if } q \text{ is odd,} \\ q^2 - 1 & \text{if } q \text{ is even.} \end{cases}$$

It follows that $k - s = 0$ if q is even, and $k - s \in \{0, (q + 1)/2\}$ if q is odd. This means that $A_\omega \leq 1$ if q is even, and $A_\omega \leq 2$ if q is odd.

Suppose q is odd and b is even. Then there are two imprimitive elements, namely

$$b + \frac{1}{2}b(q - 1) = \frac{1}{2}b(q + 1), \quad (2-2)$$

giving $\nu\nu_0^{b/2} = \xi^{b/2} \circ N$, and

$$b + \frac{b + q + 1}{2}(q - 1) = \frac{b + q - 1}{2}(q + 1), \quad (2-3)$$

giving $\nu\nu_0^{(b+q+1)/2} = \xi^{(b+q-1)/2} \circ N$. Hence $A_\omega = 2$ in this case. Noting that b is even if and only if $\omega^{(q-1)/2} = 1$, we obtain the first claim of the proposition: $P_\omega = q + 1 - A_\omega = q - 1$.

Suppose q is odd and b is odd. Then for all k , $b + k(q - 1)$ is odd, and hence it cannot be divisible by the even number $q + 1$. So $A_\omega = 0$, and $P_\omega = q + 1$, proving the second claim of the proposition.

If q is even, then as shown above, $A_\omega \leq 1$. If b is even then (2-2) is a solution, and if b is odd, (2-3) is a solution. Either way, this shows that $A_\omega \geq 1$, and hence $A_\omega = 1$, proving the final claim that $P_\omega = q + 1 - A_\omega = q$ when q is even. \square

The character sums in the next proposition will be used in Section 4 when we sum matrix coefficients over all representations with a given central character.

Proposition 2.4. *Let ω be a character of k^* , and let $[\omega]$ denote the set of primitive characters of L^* extending ω .*

Suppose q is odd and $\omega^{(q-1)/2}$ is nontrivial. Then for $\alpha \in L^$,*

$$\sum_{\nu \in [\omega]} \nu(\alpha) = \begin{cases} (q+1)\omega(\alpha) & \text{if } \alpha \in k^*, \\ 0 & \text{if } \alpha \notin k^*. \end{cases} \tag{2-4}$$

Suppose q is odd and $\omega^{(q-1)/2}$ is trivial. Then for $\alpha \in L^$,*

$$\sum_{\nu \in [\omega]} \nu(\alpha) = \begin{cases} (q-1)\omega(\alpha) & \text{if } \alpha \in k^*, \\ -2\omega(\alpha^{(q+1)/2}) & \text{if } \alpha \notin k^*, \alpha^{(q^2-1)/2} = 1, \\ 0 & \text{if } \alpha^{(q^2-1)/2} = -1. \end{cases} \tag{2-5}$$

(Note that necessarily $\alpha \notin k^$ if $\alpha^{(q^2-1)/2} \neq 1$.)*

Suppose q is even. Then for $\alpha \in L^$,*

$$\sum_{\nu \in [\omega]} \nu(\alpha) = \begin{cases} q\omega(\alpha) & \text{if } \alpha \in k^*, \\ -\omega(N(\alpha)^{1/2}) & \text{if } \alpha \notin k^*. \end{cases} \tag{2-6}$$

Here, we note that the square root is unique in k^ , since the square function is a bijection when q is even.*

Proof. If $\alpha \in k^*$, then the sum is equal to $P_\omega\omega(\alpha)$ and the assertions follow from the previous proposition. So we may assume that $\alpha \notin k^*$. We use the notation from the previous proof. Suppose q is odd and $\omega^{(q-1)/2}$ is nontrivial. By the proof of the previous proposition,

$$\sum_{\nu \in [\omega]} \nu(\alpha) = \nu(\alpha) \sum_{m=0}^q \nu_0^m(\alpha),$$

where on the right-hand side ν is any fixed element of $[\omega]$. Noting that

$$\sum_{m=0}^q \nu_0^m(\alpha) = \sum_{\chi \in \widehat{L^*/k^*}} \chi(\alpha) = \begin{cases} q+1 & \text{if } \alpha \in k^*, \\ 0 & \text{if } \alpha \notin k^*, \end{cases} \tag{2-7}$$

(2-4) follows.

Now suppose q is odd and $\omega^{(q-1)/2}$ is trivial. By the proof of the previous proposition,

$$\sum_{\nu \in [\omega]} \nu(\alpha) = \xi^b(\alpha) \left(\sum_{m=0}^q \nu_0^m(\alpha) - \nu_0(\alpha)^{b/2} - \nu_0(\alpha)^{(b+q+1)/2} \right).$$

Since $\alpha \notin k^*$, by (2-7) this is equal to

$$-\xi^b(\alpha) [\nu_0(\alpha)^{b/2} + \nu_0(\alpha)^{b/2} \nu_0(\alpha)^{(q+1)/2}].$$

Recalling that $\nu_0 = \xi^{q-1}$ and writing $\xi^b = \nu$, this is

$$= -\nu(\alpha^{1+((q-1)/2)})[1 + \xi(\alpha^{(q^2-1)/2})] = -\nu(\alpha^{(q+1)/2})[1 + \xi(\alpha^{(q^2-1)/2})]. \tag{2-8}$$

Observe that $\alpha^{(q^2-1)/2} = \pm 1$ since its square is 1. If it is equal to +1, then $\alpha^{(q+1)/2} \in k^*$ since its $(q-1)$ -st power is 1, and we immediately obtain the middle line of (2-5). Otherwise $\xi(\alpha^{(q^2-1)/2}) = \xi(-1) = -1$ and (2-8) vanishes.

When q is even, there exists a choice of ξ for which $\omega = \xi^b$ with b even. Then using (2-2), we find that when $\alpha \notin k^*$,

$$\sum_{\nu \in [\omega]} \nu(\alpha) = -\xi^{b/2}(N(\alpha)) = -\omega(N(\alpha)^{1/2}). \quad \square$$

2.2. Model for cuspidal representations. There are various ways to construct the cuspidal representation ρ_ν attached to a primitive character ν . The action of L^* on the k -vector space $L \cong k^2$ by multiplication gives an identification

$$L^* \cong T \tag{2-9}$$

of L^* with a nonsplit torus $T \subset G$, with $k^* \subset L^*$ mapping onto $Z \subset T$. The characteristic polynomial of an element $g \in G$ is irreducible over k if and only if g is conjugate to an element of $T - Z$.

Fix a nontrivial character of the additive group

$$\psi : k \longrightarrow \mathbb{C}^*,$$

viewed in the obvious way as a character of U . Then one may define ρ_ν implicitly by

$$\text{Ind}_{ZU}^G(\nu \otimes \psi) = \rho_\nu \oplus \text{Ind}_T^G \nu; \tag{2-10}$$

see [Bushnell and Henniart 2006, Theorem 6.4]. Although (2-10) allows for computation of the trace of ρ_ν (see (2-20) below), it is not convenient for computing the matrix coefficients. For this purpose we shall use the explicit model for ρ_ν defined in [Piatetski-Shapiro 1983, Section 13] as follows.¹

Given a primitive character ν of L^* and ψ as above, let

$$V = \mathbb{C}[k^*]$$

¹There is a minus sign missing from the definition of $j(x)$ in Equation (4) of Section 13 of [Piatetski-Shapiro 1983] (otherwise his identity (6) will not hold). Likewise a minus sign is missing from (16) on page 40. The expression four lines above (16) is correct (except K should be K^*).

be the vector space of functions $f : k^* \rightarrow \mathbb{C}$. We define a representation ρ_ν of G on V as follows. For any $\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in B$, $f \in V$, let

$$\left[\rho_\nu \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} f \right] (x) = \nu(d) \psi(bd^{-1}x) f(ad^{-1}x) \quad (x \in k^*), \tag{2-11}$$

and for $g \in G - B$, define

$$(\rho_\nu(g)f)(x) = \sum_{y \in k^*} \phi(x, y; g) f(y) \quad (x \in k^*), \tag{2-12}$$

where, for $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G - B$,

$$\phi(x, y; g) = -\frac{1}{q} \psi \left[\frac{ax + dy}{c} \right] \sum_{\substack{t \in L^* \\ N(t) = xy^{-1} \det g}} \psi \left(-\frac{y}{c} (t + \bar{t}) \right) \nu(t). \tag{2-13}$$

Theorem 2.5. *If ν is a primitive character of L^* , (2-11) and (2-12) give a well-defined representation (ρ_ν, V) which is cuspidal. Furthermore, every cuspidal representation is isomorphic to some ρ_ν , and $\rho_\nu \cong \rho_{\nu'}$ if and only if $\nu' \in \{\nu, \nu^q\}$. In particular, there are $(q^2 - q)/2$ distinct cuspidal representations.*

Proof. See [Piatetski-Shapiro 1983, Section 13–14], where it is assumed that $q > 2$ throughout. When $q = 2$, G is isomorphic to the symmetric group S_3 . The unique cuspidal representation is the character sending each permutation to its sign. It is readily checked that the above construction defines this character as well, so the theorem remains valid when $q = 2$. □

Define an inner product on V by

$$\langle f_1, f_2 \rangle = \sum_{x \in k^*} f_1(x) \overline{f_2(x)}. \tag{2-14}$$

We will work with the orthonormal basis

$$\mathfrak{B} = \{f_r\}_{r \in k^*} \quad \text{for } f_r(x) = \begin{cases} 1 & \text{if } x = r, \\ 0 & \text{if } x \neq r. \end{cases}$$

Proposition 2.6. *Let ν be a primitive character of L^* , and let ρ_ν be the associated cuspidal representation of G . Then ρ_ν is unitary with respect to the inner product (2-14).*

Proof. By linearity, it suffices to prove that for all $f_r, f_s \in \mathfrak{B}$ and $g \in G$,

$$\langle \rho_\nu(g)f_r, \rho_\nu(g)f_s \rangle = \langle f_r, f_s \rangle. \tag{2-15}$$

By the Bruhat decomposition $G = B \cup Bw'U$ for $w' = \begin{pmatrix} & 1 \\ -1 & \end{pmatrix}$ and the fact that ρ_ν is a homomorphism, we only need to consider $g \in B$ and $g = w'$.

Suppose first that $g = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in B$. Then by (2-11),

$$\langle \rho_\nu(g) f_r, \rho_\nu(g) f_s \rangle = \sum_{x \in k^*} \nu(d) \psi(bd^{-1}x) f_r(ad^{-1}x) \overline{\nu(d) \psi(bd^{-1}x) f_s(ad^{-1}x)}.$$

Using the fact that ν and ψ are unitary, and replacing x by $a^{-1}dx$, we see that this expression equals

$$\sum_{x \in k^*} f_r(x) \overline{f_s(x)} = \langle f_r, f_s \rangle,$$

as needed.

It remains to prove (2-15) for $g = w' = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. By (2-12),

$$\begin{aligned} \rho_\nu(w') f(x) &= \sum_{y \in k^*} \phi(x, y; w') f(y) = -\frac{1}{q} \sum_{y \in k^*} \sum_{\substack{t \in L^* \\ N(t) = xy^{-1}}} \psi(y(t + \bar{t})) \nu(t) f(y) \\ &= -\frac{1}{q} \sum_{y \in k^*} \sum_{\substack{u \in L^* \\ N(u) = xy}} \psi(u + \bar{u}) \nu(u) \nu(y^{-1}) f(y) = \sum_{y \in k^*} \nu(y^{-1}) j(xy) f(y), \end{aligned}$$

where

$$j(t) = -\frac{1}{q} \sum_{\substack{u \in L^* \\ N(u) = t}} \psi(u + \bar{u}) \nu(u). \quad (2-16)$$

Hence

$$\rho_\nu(w') f_r(x) = \sum_{y \in k^*} \nu(y^{-1}) j(xy) f_r(y) = \nu(r^{-1}) j(rx).$$

We now see that

$$\begin{aligned} \langle \rho_\nu(w') f_r, \rho_\nu(w') f_s \rangle &= \sum_{x \in k^*} \nu(r^{-1}) j(rx) \overline{\nu(s^{-1}) j(sx)} = \nu(sr^{-1}) \sum_{x \in k^*} j(rx) \overline{j(sx)} \\ &= \nu(sr^{-1}) \sum_{x \in k^*} j(rs^{-1}x) \overline{j(x)}. \end{aligned}$$

Taking $r' = rs^{-1}$, it suffices to prove that

$$\sum_{x \in k^*} j(r'x) \overline{j(x)} = \begin{cases} 0 & \text{if } r' \neq 1, \\ 1 & \text{if } r' = 1. \end{cases} \quad (2-17)$$

From the definition (2-16), we have

$$\begin{aligned} \sum_{x \in k^*} j(r'x) \overline{j(x)} &= \frac{1}{q^2} \sum_{x \in k^*} \sum_{N(\alpha) = r'x} \sum_{N(\beta) = x} \psi(\alpha + \bar{\alpha}) \nu(\alpha) \psi(-\beta - \bar{\beta}) \nu(\beta^{-1}) \\ &= \frac{1}{q^2} \sum_{\beta \in L^*} \sum_{N(\alpha) = r'N(\beta)} \psi(\alpha + \bar{\alpha} - \beta - \bar{\beta}) \nu(\alpha \beta^{-1}). \end{aligned}$$

Since the norm map is surjective, there exists $z \in L^*$ such that $N(z) = r'$. Then $\alpha = z\beta u$ for some $u \in L^1$. This allows us to rewrite the above sum as

$$\begin{aligned} \sum_{x \in k^*} j(r'x)\overline{j(x)} &= \frac{1}{q^2} \sum_{\beta \in L^*} \sum_{u \in L^1} \psi(z\beta u + \overline{z\beta u} - \beta - \bar{\beta})v(zu) \\ &= \frac{v(z)}{q^2} \sum_{u \in L^1} v(u) \sum_{\beta \in L^*} \psi(\text{tr}[(zu - 1)\beta]). \end{aligned} \tag{2-18}$$

Generally, for $c \in L$, the map $R(\beta) = \psi(\text{tr}[c\beta])$ is a homomorphism from L to \mathbb{C}^* . If $c \neq 0$, then R is nontrivial since the trace map from L to k is surjective. It follows that

$$\sum_{\beta \in L^*} \psi(\text{tr}[c\beta]) = \begin{cases} -1 & \text{if } c \neq 0, \\ q^2 - 1 & \text{if } c = 0. \end{cases} \tag{2-19}$$

Suppose $r' \neq 1$. Then $N(zu) = N(z) = r' \neq 1$, so in particular $zu \neq 1$. Therefore (2-18) becomes

$$\sum_{x \in k^*} j(r'x)\overline{j(x)} = -\frac{v(z)}{q^2} \sum_{u \in L^1} v(u) = 0,$$

where we have used the fact (Proposition 2.1) that v is a nontrivial character of L^1 since v is primitive.

Now suppose $r' = 1$. Then we can take $z = 1$, so by (2-18) and (2-19),

$$\begin{aligned} \sum_{x \in k^*} j(x)\overline{j(x)} &= \frac{1}{q^2} \sum_{u \in L^1} v(u) \sum_{\beta \in L^*} \psi(\text{tr}[(u - 1)\beta]) \\ &= \frac{q^2 - 1}{q^2} - \frac{1}{q^2} \sum_{\substack{u \in L^1 \\ u \neq 1}} v(u) = \left(1 - \frac{1}{q^2}\right) + \frac{1}{q^2} = 1, \end{aligned}$$

since $\sum_{\substack{u \in L^1 \\ u \neq 1}} v(u) = -1$, again because v is nontrivial on L^1 . This proves (2-17). \square

2.3. Matrix coefficients of cuspidal representations. Let v be a primitive character of L^* . Using (2-10), one finds that

$$\text{tr } \rho_v(x) = \begin{cases} (q - 1)v(x) & \text{if } x \in Z, \\ -v(z) & \text{if } x = zu, z \in Z, u \in U, u \neq 1, \\ -v(x) - v^q(x) & \text{if } x \in T, x \notin Z, \\ 0 & \text{if no conjugate of } x \text{ is in } T \cup ZU \end{cases} \tag{2-20}$$

(see [Bushnell and Henniart 2006, (6.4.1)]). This is a sum of matrix coefficients. For the coefficients themselves, we use the model given in the previous section to prove the following (which can also be used to derive (2-20)).

Theorem 2.7. *Let $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G$ and $f_r, f_s \in \mathfrak{B}$. Let ρ_ν be a cuspidal representation of G . If $g \in B$, then*

$$\langle \rho_\nu(g) f_r, f_s \rangle = \begin{cases} \nu(d)\psi(bd^{-1}s) & \text{if } r = ad^{-1}s, \\ 0 & \text{if } r \neq ad^{-1}s. \end{cases}$$

If $g \notin B$, then

$$\langle \rho_\nu(g) f_r, f_s \rangle = \phi(s, r; g),$$

where ϕ is defined in (2-13).

Proof. First suppose $g \in B$ (i.e., $c = 0$). Then

$$\begin{aligned} \langle \rho_\nu(g) f_r, f_s \rangle &= \sum_{x \in k^*} [\rho_\nu(g) f_r](x) \overline{f_s(x)} = [\rho_\nu(g) f_r](s) \\ &= \nu(d)\psi(bd^{-1}s) f_r(ad^{-1}s) = \begin{cases} \nu(d)\psi(bd^{-1}s) & \text{if } r = ad^{-1}s, \\ 0 & \text{if } r \neq ad^{-1}s. \end{cases} \end{aligned}$$

Now, suppose $g \notin B$. Then

$$[\rho_\nu(g) f_r](x) = \sum_{y \in k^*} \phi(x, y; g) f_r(y) = \phi(x, r; g).$$

Therefore

$$\langle \rho_\nu f_r, f_s \rangle = \sum_{x \in k^*} [\rho_\nu(g) f_r](x) \overline{f_s(x)} = [\rho_\nu(g) f_r](s) = \phi(s, r; g),$$

as needed. □

3. Depth-zero supercuspidal representations of $GL_2(F)$

We move now to the p -adic setting. When no field is specified, G, Z, B, U , etc., will henceforth denote the corresponding subgroups of $GL_2(F)$ rather than $GL_2(k)$. Fix a primitive character ν , and let ρ_ν be the associated cuspidal representation of $GL_2(k)$. We view ρ_ν as a representation of $K = GL_2(\mathfrak{o})$ via reduction modulo \mathfrak{p} :

$$K \longrightarrow GL_2(k) \longrightarrow GL(V).$$

The central character of this representation is given by $z \mapsto \nu(z(1 + \mathfrak{p}))$ for $z \in \mathfrak{o}^*$. Extend this character of \mathfrak{o}^* to $Z \cong F^* = \bigcup_{n \in \mathbb{Z}} \varpi^n \mathfrak{o}^*$ by choosing a complex number $\nu(\varpi)$ of absolute value 1. We denote this character of F^* by ν . This allows us to view ρ_ν as a unitary representation of the group ZK . Let

$$\pi_\nu = \text{c-Ind}_{ZK}^{GL_2(F)}(\rho_\nu)$$

be the representation of G compactly induced from ρ_ν . This representation is irreducible and supercuspidal (see, for example, [Bump 1997, Theorem 4.8.1]).

3.1. Matrix coefficients. Define an inner product on $\text{c-Ind}_{ZK}^G(\rho_v)$ by

$$\langle f_1, f_2 \rangle = \sum_{x \in ZK \setminus G} \langle f_1(x), f_2(x) \rangle_V, \tag{3-1}$$

where $\langle \cdot, \cdot \rangle_V$ denotes the inner product on V defined in (2-14). As in Section 1, this inner product is G -equivariant.

The matrix coefficients of π_v can now be computed explicitly by using Proposition 1.1 in conjunction with Theorem 2.7. Likewise, by Proposition 1.2, if we normalize so that $\text{meas}(\overline{K}) = 1$, then

$$d_{\pi_v} = \dim \rho_v = q - 1. \tag{3-2}$$

Define a function $\phi_v : G \rightarrow \mathbb{C}$ by

$$\phi_v(x) = \begin{cases} \overline{\text{tr } \rho_v(x)} & \text{if } x \in ZK, \\ 0 & \text{otherwise.} \end{cases} \tag{3-3}$$

Then this is a *pseudocoefficient* of π_v in the sense that for any irreducible tempered representation π of G with central character ω ,

$$\text{tr } \pi(\phi_v) = \begin{cases} 1 & \text{if } \pi \cong \pi_v, \\ 0 & \text{otherwise} \end{cases}$$

(see [Palm 2012, Section 9.4.1]). The function ϕ_v may be computed explicitly using (2-20).

3.2. New vectors. For an integer $n \geq 0$, define the congruence subgroup

$$K_1(\mathfrak{p}^n) = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in K \mid c, (d - 1) \in \mathfrak{p}^n \right\}.$$

If π is a representation of G , we let $\pi^{K_1(\mathfrak{p}^n)}$ denote the space of vectors fixed by $K_1(\mathfrak{p}^n)$. By a result of Casselman [1973], for any irreducible admissible representation π of G , there exists a unique ideal \mathfrak{p}^n (the *conductor* of π) for which $\dim \pi^{K_1(\mathfrak{p}^n)} = 1$ and $\dim \pi^{K_1(\mathfrak{p}^{n-1})} = 0$. A nonzero vector fixed by $K_1(\mathfrak{p}^n)$ is called a *new vector*. The supercuspidal representations constructed above have conductor \mathfrak{p}^2 . We shall give an elementary proof below, and exhibit a new vector. More generally, the new vectors for depth-zero supercuspidal representations of $\text{GL}_n(F)$ were identified by Reeder [1991, Example (2.3)].

Proposition 3.1. *The supercuspidal representation π_v defined above has conductor \mathfrak{p}^2 . If we let $w \in \mathbb{C}[k^*]$ denote the constant function 1, that is,*

$$w = \sum_{r \in k^*} f_r, \tag{3-4}$$

then the function $f = f_{(\varpi \ 1),w}$ supported on the coset

$$ZK \begin{pmatrix} \varpi & 0 \\ 0 & 1 \end{pmatrix} = ZK \begin{pmatrix} \varpi & 0 \\ 0 & 1 \end{pmatrix} K_1(\mathfrak{p}^2),$$

and defined by

$$f \left(zk \begin{pmatrix} \varpi & 0 \\ 0 & 1 \end{pmatrix} \right) = \rho_v(zk)w,$$

is a new vector of π_v .

Proof. To see that f is $K_1(\mathfrak{p}^2)$ -invariant, it suffices to show that

$$f \left(\begin{pmatrix} \varpi & \\ & 1 \end{pmatrix} k \right) = w \quad \text{for all } k \in K_1(\mathfrak{p}^2).$$

Writing $k = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $c \in \mathfrak{p}^2$ and $d \in 1 + \mathfrak{p}^2$, we have

$$\begin{aligned} f \left(\begin{pmatrix} \varpi & \\ & 1 \end{pmatrix} k \right) &= f \left(\begin{pmatrix} \varpi & \\ & 1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \varpi^{-1} & \\ & 1 \end{pmatrix} \begin{pmatrix} \varpi & \\ & 1 \end{pmatrix} \right) \\ &= \rho_v \left(\begin{pmatrix} a & \varpi b \\ \varpi^{-1}c & d \end{pmatrix} \right) w = \rho_v \left(\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \right) w \\ &= \sum_{r \in k^*} \rho_v \left(\begin{pmatrix} a & \\ & 1 \end{pmatrix} \right) f_r = \sum_{r \in k^*} f_{a^{-1}r} = w, \end{aligned}$$

as needed. This shows that $\pi_v^{K_1(\mathfrak{p}^2)} \neq 0$, so the conductor divides \mathfrak{p}^2 . There are various ways to see that the conductor is exactly \mathfrak{p}^2 . When $n > 1$, it is straightforward to show that a continuous irreducible n -dimensional complex representation of the Weil group of F has Artin conductor of exponent at least n (see, for example, [Gross and Reeder 2010, Equation (1)]). So by the local Langlands correspondence, the conductor of any supercuspidal representation of $\text{GL}_n(F)$ is divisible by \mathfrak{p}^n , giving the desired conclusion here when $n = 2$. For an elementary proof in the present situation, one can observe that a function $f \in \text{c-Ind}_{ZK}^G(\rho_v)$ supported on a coset ZKx is $K_1(\mathfrak{p})$ -invariant if and only if ρ_v is trivial on $K \cap xK_1(\mathfrak{p})x^{-1}$. Using the double coset decomposition

$$G = \bigcup_{n \geq 0} ZK \begin{pmatrix} \varpi^n & \\ & 1 \end{pmatrix} K = \bigcup_{n \geq 0} \bigcup_{\delta \in \overline{K/K_1(\mathfrak{p})}} ZK \begin{pmatrix} \varpi^n & \\ & 1 \end{pmatrix} \delta K_1(\mathfrak{p})$$

(we may use the representatives $\delta \in \{1\} \cup \{(y \ 1) \mid y \in \mathfrak{o}/\mathfrak{p}\}$; see, for example, [Knightly and Li 2006, Lemma 13.1]), it suffices to consider $x = \begin{pmatrix} \varpi^n & \\ & 1 \end{pmatrix} \delta$, and one checks that in each case ρ_v is *not* trivial on $K \cap xK_1(\mathfrak{p})x^{-1}$, so $\pi_v^{K_1(\mathfrak{p})} = \{0\}$. \square

3.3. Matrix coefficient of the new vector. Generally, if π is a supercuspidal representation with unit new vector v and formal degree d_π , the function

$$g \mapsto d_\pi \overline{\langle \pi(g)v, v \rangle}$$

can be used to define a projection operator onto $\mathbb{C}v$.

In the present context, if f is the new vector defined in Proposition 3.1, one finds easily that $\|f\|^2 = (q - 1)$, so with the standard normalization $\text{meas}(\overline{K}) = 1$, by (3-2) we have

$$\Phi_v(g) \stackrel{\text{def}}{=} d_{\pi_v} \overline{\left\langle \pi_v(g) \frac{f}{\|f\|}, \frac{f}{\|f\|} \right\rangle} = \overline{\langle \pi_v(g)f, f \rangle}. \tag{3-5}$$

By Proposition 1.1,

$$\text{Supp}(\Phi_v) = \begin{pmatrix} \varpi^{-1} & \\ & 1 \end{pmatrix} ZK \begin{pmatrix} \varpi & \\ & 1 \end{pmatrix},$$

and for $g = \begin{pmatrix} \varpi^{-1} & \\ & 1 \end{pmatrix} h \begin{pmatrix} \varpi & \\ & 1 \end{pmatrix} \in \text{Supp}(\Phi_v)$,

$$\Phi_v(g) = \overline{\langle \rho_v(h)w, w \rangle}_V \tag{3-6}$$

for w as in (3-4). This is computed as follows.

Theorem 3.2. Let $h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(k) = \text{GL}_2(k)$, and let $w \in V$ be the function defined in (3-4). Then

$$\langle \rho_v(h)w, w \rangle_V = \begin{cases} (q - 1)v(d) & \text{if } b = c = 0, \\ -v(d) & \text{if } c = 0, b \neq 0, \\ - \sum_{\substack{\alpha \in L^* \\ \alpha + \bar{\alpha} = \frac{aN(\alpha)}{\det h} + d}} v(\alpha) & \text{if } c \neq 0. \end{cases}$$

Remark. The sum may be evaluated using Proposition 3.3 below.

Proof. To ease notation, we drop the subscript V from the inner product. Suppose $h = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in B(k)$. Then applying Theorem 2.7,

$$\langle \rho_v(h)w, w \rangle = \sum_{r, s \in k^*} \langle \rho_v(h) f_r, f_s \rangle = \sum_{s \in k^*} \langle \rho_v(h) f_{ad^{-1}s}, f_s \rangle = v(d) \sum_{s \in k^*} \psi(bd^{-1}s).$$

This gives

$$\langle \rho_v(h)w, w \rangle = \begin{cases} (q - 1)v(d) & \text{if } b = 0, \\ -v(d) & \text{if } b \neq 0. \end{cases}$$

Now suppose $h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(k) - B(k)$. Then by [Theorem 2.7](#),

$$\begin{aligned} \langle \rho_v(h)w, w \rangle &= \sum_{r,s \in k^*} \phi(s, r; h) \\ &= -\frac{1}{q} \sum_{r,s \in k^*} \psi(c^{-1}(sa + rd)) \sum_{\substack{\alpha \in L^* \\ N(\alpha) = sr^{-1} \det h}} \psi(-rc^{-1}(\alpha + \bar{\alpha}))v(\alpha). \end{aligned}$$

Let $l = sr^{-1}$, so $s = rl$. From the previous display we have

$$\begin{aligned} \langle \rho_v(h)w, w \rangle &= -\frac{1}{q} \sum_{r \in k^*} \sum_{l \in k^*} \psi(c^{-1}(rla + rd)) \sum_{N(\alpha) = l \det h} \psi(-rc^{-1}(\alpha + \bar{\alpha}))v(\alpha) \\ &= -\frac{1}{q} \sum_{l \in k^*} \sum_{N(\alpha) = l \det h} v(\alpha) \sum_{r \in k^*} \psi(rc^{-1}(al + d - (\alpha + \bar{\alpha}))) \\ &= -\frac{1}{q} \sum_{l \in k^*} \sum_{N(\alpha) = l \det h} v(\alpha) \sum_{r \in k^*} \psi(r(al + d - (\alpha + \bar{\alpha}))). \end{aligned}$$

There are two cases for the inner sum:

$$\sum_{r \in k^*} \psi(r(al + d - (\alpha + \bar{\alpha}))) = \begin{cases} -1 & \text{if } al + d - (\alpha + \bar{\alpha}) \neq 0, \\ q - 1 & \text{if } al + d - (\alpha + \bar{\alpha}) = 0. \end{cases}$$

Therefore,

$$\begin{aligned} \langle \rho_v(h)w, w \rangle &= -\frac{1}{q} \sum_{l \in k^*} \left(\sum_{\substack{N(\alpha) = l \det h \\ \alpha + \bar{\alpha} \neq al + d}} -v(\alpha) + \sum_{\substack{N(\alpha) = l \det h \\ \alpha + \bar{\alpha} = al + d}} (q - 1)v(\alpha) \right) \\ &= -\frac{1}{q} \sum_{l \in k^*} \left(- \sum_{N(\alpha) = l \det h} v(\alpha) + q \sum_{\substack{N(\alpha) = l \det h \\ \alpha + \bar{\alpha} = al + d}} v(\alpha) \right). \end{aligned}$$

By [Lemma 2.2](#), the first sum in the big parentheses vanishes. So

$$\langle \rho_v(h)w, w \rangle = -\frac{1}{q} \sum_{l \in k^*} q \sum_{\substack{N(\alpha) = l \det h \\ \alpha + \bar{\alpha} = al + d}} v(\alpha) = - \sum_{\substack{\alpha \in L^* \\ \alpha + \bar{\alpha} = \frac{aN(\alpha)}{\det h} + d}} v(\alpha), \tag{3-7}$$

as claimed. □

This sum can be refined as follows.

Proposition 3.3. *For $l \in k^*$ and $h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(k)$ with $c \neq 0$, define*

$$p_{h,l}(X) = X^2 - \left(\frac{al}{\det h} + d \right) X + l \in k[X]. \tag{3-8}$$

Then

$$\langle \rho_\nu(h)w, w \rangle = - \sum_{\substack{l \in k^* \\ p_{h,l}(X) \text{ irred. over } k}} \sum_{\substack{\text{roots } \alpha \text{ of} \\ p_{h,l}(X) \text{ in } L^*}} \nu(\alpha) - \sum_{\substack{l \in k^* \\ p_{h,l}(X) = (X-\alpha)^2 \text{ in } k[X]}} \nu(\alpha). \quad (3-9)$$

Proof. If $\alpha \in L^*$ contributes to the sum (3-7), then it is a root of

$$X^2 - \left(\frac{aN(\alpha)}{\det h} + d \right) X + N(\alpha).$$

Conversely, given $l \in k^*$, a root α_l of $p_{h,l}(X)$ contributes to (3-7) if and only if $p_{h,l}(X) = (X - \alpha_l)(X - \bar{\alpha}_l)$ in $L[X]$, which is the case if and only if either $p_{h,l}(X)$ is irreducible or $p_{h,l}(X) = (X - \alpha_l)^2$ with $\alpha_l \in k^*$. The proposition now follows. \square

3.4. Motivation. Although specific global applications of the above formulas are beyond the scope of this article, perhaps a few words of motivation will be helpful. The two functions ϕ_ν and Φ_ν of (3-3) and (3-5) serve slightly different purposes. The former is simpler and hence easier to work with. Taken as a local component of a global test function, it is well suited for use in the Arthur–Selberg trace formula, for example if one is interested in detecting those automorphic representations $\pi = \bigotimes_w \pi_w$ of the adelic group $\text{GL}_2(\mathbb{A}_\mathbb{Q})$ with the local condition $\pi_w \cong \pi_\nu$ at a given finite place w (e.g., to obtain a dimension formula for the associated space of classical newforms). This method was treated in detail recently by Palm [2012]. On the other hand, as mentioned in the Introduction, the matrix coefficient Φ_ν gives rise to an operator projecting onto the span of the newforms attached to the global representations π as above (see [Knightly and Li 2012, Section 2.5]). It can be used in variants of the trace formula to extract finer information, like Fourier coefficients or L -values of these newforms. Of course, the utility of Φ_ν in explicit computation is limited by the complexity of the sum in Theorem 3.2.

4. Consideration of central character

The supercuspidal representations which have a given central character ω occur naturally together as irreducible subrepresentations of the right regular representation of $G(F)$ on the space of L^2 functions $f : G(F) \rightarrow \mathbb{C}$ that transform under the center by $\bar{\omega}$. It is often the case in number theory that one can achieve a certain amount of simplification by simultaneously treating all objects in a family via averaging. Here we sum the trace functions ϕ_ν from (3-3) over all isomorphism classes of depth-zero supercuspidal π_ν with a given central character, and similarly for the new vector matrix coefficients Φ_ν of (3-5).

Let ω be a unitary character of F^* , and let S_ω denote the set of isomorphism classes of depth-zero supercuspidal representations of $\text{GL}_2(F)$ with central character

ω . In order for S_ω to be nonempty, ω must be trivial on $1 + \mathfrak{p}$. In fact we have

$$|S_\omega| = \begin{cases} P_\omega/2 & \text{if } \omega|_{(1+\mathfrak{p})} = 1, \\ 0 & \text{otherwise,} \end{cases} \tag{4-1}$$

for P_ω as in Proposition 2.3. (Note that ν and ν^q have the same restriction to k^* and $\rho_\nu \cong \rho_{\nu^q}$.)

Assuming $\omega|_{(1+\mathfrak{p})}$ is trivial, consider the sum of the trace functions ϕ_ν defined in (3-3). In view of the fact that $\phi_\nu = \phi_{\nu^q}$, we define

$$\phi_\omega = \frac{1}{2} \sum_{\nu \in [\omega]} \phi_\nu, \tag{4-2}$$

with notation as in Proposition 2.4. We can make it explicit with the following.

Theorem 4.1. *Suppose q is odd and $\omega^{(q-1)/2}$ is nontrivial. Then for $x \in G(k)$,*

$$\sum_{\nu \in [\omega]} \text{tr } \rho_\nu(x) = \begin{cases} (q^2 - 1)\omega(x) & \text{if } x \in Z, \\ -(q + 1)\omega(z) & \text{if } x = zu, z \in Z, u \in U, u \neq 1, \\ 0 & \text{if no conjugate of } x \text{ is in } ZU. \end{cases}$$

Suppose q is odd and $\omega^{(q-1)/2}$ is trivial. Then with T as in (2-9),

$$\sum_{\nu \in [\omega]} \text{tr } \rho_\nu(x) = \begin{cases} (q - 1)^2\omega(x) & \text{if } x \in Z, \\ -(q - 1)\omega(z) & \text{if } x = zu, z \in Z, u \in U, u \neq 1, \\ 4\omega(x^{(q+1)/2}) & \text{if } x \in T - Z, x^{(q^2-1)/2} = 1, \\ 0 & \text{if } x \in T - Z, x^{(q^2-1)/2} = -1, \text{ or if} \\ & \text{no conjugate of } x \text{ is in } T \cup ZU. \end{cases}$$

Suppose q is even. Then

$$\sum_{\nu \in [\omega]} \text{tr } \rho_\nu(x) = \begin{cases} q(q - 1)\omega(x) & \text{if } x \in Z, \\ -q\omega(z) & \text{if } x = zu, z \in Z, u \in U, u \neq 1, \\ 2\omega(N(x)^{1/2}) & \text{if } x \in T - Z, \\ 0 & \text{if no conjugate of } x \text{ is in } T \cup ZU. \end{cases}$$

Proof. This follows immediately by examining the various cases using (2-20) and Proposition 2.4. □

Likewise, for a depth-zero supercuspidal representation $\pi = \pi_\nu$, let $\Phi_\pi = \Phi_\nu$ be the matrix coefficient defined in (3-5). Define a function Φ_ω on G by

$$\Phi_\omega(g) = \sum_{\pi \in S_\omega} \Phi_\pi(g) = \frac{1}{2} \sum_{\nu \in [\omega]} \overline{\langle \rho_\nu(h)w, w \rangle}_\nu \tag{4-3}$$

for $g = \begin{pmatrix} \varpi^{-1} & \\ & 1 \end{pmatrix} h \begin{pmatrix} \varpi & \\ & 1 \end{pmatrix}$ with $h \in ZK$. In principle, this function can be used to define an operator that projects the automorphic spectrum of $\text{GL}_2(\mathbf{A}_\mathbb{Q})$ onto the span

of those newforms of a given weight and level p^2 that correspond to automorphic representations which are unramified away from p and are supercuspidal (as opposed to special or principal series) at p .

One can evaluate Φ_ω via the following.

Theorem 4.2. *Suppose $h = \begin{pmatrix} a & \\ & d \end{pmatrix} \in G(k)$ is diagonal. Then*

$$\sum_{v \in [\omega]} \langle \rho_v(h)w, w \rangle_V = \begin{cases} (q^2 - 1)\omega(d) & \text{if } q \text{ is odd and } \omega^{(q-1)/2} \text{ is nontrivial,} \\ (q - 1)^2\omega(d) & \text{if } q \text{ is odd and } \omega^{(q-1)/2} \text{ is trivial,} \\ q(q - 1)\omega(d) & \text{if } q \text{ is even.} \end{cases} \quad (4-4)$$

If $h = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \in B(k)$ with $b \neq 0$, then

$$\sum_{v \in [\omega]} \langle \rho_v(h)w, w \rangle_V = \begin{cases} -(q + 1)\omega(d) & \text{if } q \text{ is odd and } \omega^{(q-1)/2} \text{ is nontrivial,} \\ -(q - 1)\omega(d) & \text{if } q \text{ is odd and } \omega^{(q-1)/2} \text{ is trivial,} \\ -q\omega(d) & \text{if } q \text{ is even.} \end{cases} \quad (4-5)$$

If $g \in G(k) - B(k)$, then the sum is given by (4-6) below.

Proof. Suppose $h = \begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$ is diagonal. Then by [Theorem 3.2](#), we can write

$$\sum_{v \in [\omega]} \langle \rho_v(h)w, w \rangle = (q - 1) \sum_{v \in [\omega]} v(d).$$

Applying [Proposition 2.4](#) now gives (4-4), using the fact that $d \in k^*$.

Similarly, if $h = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$ with $b \neq 0$, then applying [Theorem 3.2](#),

$$\sum_{v \in [\omega]} \langle \rho_v(h)w, w \rangle = - \sum_{v \in [\omega]} v(d).$$

Using [Proposition 2.4](#), this gives (4-5).

Now suppose $h = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G(k) - B(k)$. By [Proposition 3.3](#),

$$\sum_{v \in [\omega]} \langle \rho_v(h)w, w \rangle = - \sum_{\substack{l \in k^* \\ p_{h,l}(X) \text{ irred.}}} \sum_{\substack{\text{roots } \alpha \text{ of} \\ p_{h,l}(X) \text{ in } L^*}} \sum_{v \in [\omega]} v(\alpha) - \sum_{\substack{l \in k^* \\ p_{h,l}(X) = (X - \alpha)^2}} \sum_{v \in [\omega]} v(\alpha),$$

where $p_{h,l}(X) = X^2 - ((al/\det h) + d)X + l \in k[X]$. If we fix one root $\alpha_l \in L^*$ of $p_{h,l}(X)$ for each l , then the above is

$$= -2 \sum_{\substack{l \in k^* \\ p_{h,l}(X) \text{ irred.}}} \sum_{v \in [\omega]} v(\alpha_l) - P_\omega \sum_{\substack{l \in k^* \\ p_{h,l}(X) = (X - \alpha_l)^2}} \omega(\alpha_l), \quad (4-6)$$

where $P_\omega = |[\omega]|$ as in [Proposition 2.3](#). We have used the fact that

$$\sum_{v \in [\omega]} v(\alpha_l) = \sum_{v \in [\omega]} v(\bar{\alpha}_l),$$

since ν and ν^q both belong to $[\omega]$. Once again, (4-6) can be evaluated on a case-by-case basis using Proposition 2.4.

For instance, suppose q is odd. If $\omega^{(q-1)/2}$ is nontrivial, the first term of (4-6) vanishes. If $(al + d \det h) = 0$, then l makes no contribution to the second term. In particular, the term vanishes if $a = d = 0$. Generally, at most two l contribute to the second term when q is odd since $2\alpha_l = ((al / \det h) + d)$ implies that l satisfies the quadratic equation $l = \alpha_l^2 = \frac{1}{4}((al / \det h) + d)^2$.

On the other hand, if q is even, a given $l \in k^*$ contributes to the second term of (4-6) if and only if $(al + d \det h) = 0$ since $(X - \alpha_l)^2 = X^2 - \alpha_l^2$, and as remarked earlier every l is a square when q is even. □

5. Examples

Take $q = 2$. By (4-1), there is a unique supercuspidal representation π of $\text{GL}_2(\mathbb{Q}_2)$ of conductor 2^2 . As mentioned before, the cuspidal representation of $\text{GL}_2(\mathbb{F}_2) \cong S_3$ is the character ρ sending a matrix g to $(-1)^{|g|+1}$, where $|g|$ is the order of g in the finite group. Explicitly, ρ sends $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ to $1, -1, -1, -1, 1, 1$ respectively. This one-dimensional representation of course coincides with its matrix coefficient:

$$\langle \rho(h)w, w \rangle = \rho(h)\langle w, w \rangle = \rho(h).$$

Indeed, one may verify that Theorem 3.2 recovers ρ when $q = 2$. For example, consider $h = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$. The polynomial (3-8) becomes $p_{h,1}(X) = X^2 + X + 1$. If $\theta \in \mathbb{F}_4^*$ is a root, then $\nu(\theta) = \exp(2\pi i / 3)$ defines a primitive character. By (3-9),

$$\rho(h) = -(\nu(\theta) - \nu(\theta^2)) = 1.$$

By (3-6), the matrix coefficient Φ attached to the new vector of π has the simple expression

$$\Phi(g) = \begin{cases} \rho(h) & \text{if } g = z \begin{pmatrix} 2^{-1} & \\ & 1 \end{pmatrix} h \begin{pmatrix} 2 & \\ & 1 \end{pmatrix} \in Z \begin{pmatrix} 2^{-1} & \\ & 1 \end{pmatrix} K \begin{pmatrix} 2 & \\ & 1 \end{pmatrix}, \\ 0 & \text{otherwise.} \end{cases} \tag{5-1}$$

Now consider $k = \mathbb{F}_5$ and $L = \mathbb{F}_{25}$. Then $L = k[\theta]$, where $\theta^2 = 2$. One finds that $1 + 2\theta$ generates the cyclic group L^* , so the characters of L^* are the maps

$$\nu_n(1 + 2\theta) = \zeta^n \quad (n \in \mathbb{Z}/24\mathbb{Z}),$$

where $\zeta = \exp(2\pi i / 24)$. Note that ν_n is primitive if and only if $6 \nmid n$. There are exactly four characters of k^* , given by

$$\omega_n(3) = i^n \quad (n \in \mathbb{Z}/4\mathbb{Z}).$$

Noting that $v_n(3) = v_n((1 + 2\theta)^6) = \zeta^{6n} = i^n$, we see that $v_n|_{k^*} = \omega_n$. So the equivalence classes of primitive characters of L^* are as follows:

$$[\omega_0] = \{\nu_4, \nu_{20}, \nu_8, \nu_{16}\}, \quad [\omega_1] = \{\nu_1, \nu_5, \nu_9, \nu_{21}, \nu_{13}, \nu_{17}\},$$

$$[\omega_2] = \{\nu_2, \nu_{10}, \nu_{14}, \nu_{22}\}, \quad [\omega_3] = \{\nu_3, \nu_{15}, \nu_7, \nu_{11}, \nu_{19}, \nu_{23}\},$$

where each primitive character is listed alongside its conjugate. The above illustrates [Proposition 2.3](#). As a simple illustration of [Theorem 4.2](#), we now show that

$$\sum_{\nu \in [\omega_0]} \left\langle \rho_\nu \left(\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right) w, w \right\rangle = 0.$$

For $h = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ the polynomial [\(3-8\)](#) becomes $p_{h,l}(X) = X^2 + l$. The second term in [\(4-6\)](#) vanishes. Thus an element $l \in k^*$ contributes to [\(4-6\)](#) only if $-l$ is a quadratic nonresidue in k . The roots of $X^2 + 2$ are

$$\alpha_2 = (1 + 2\theta)^3 \quad \text{and} \quad \bar{\alpha}_2 = (1 + 2\theta)^{15}.$$

The roots of $X^2 + 3$ are

$$\alpha_3 = \theta = (1 + 2\theta)^9 \quad \text{and} \quad \bar{\theta} = (1 + 2\theta)^{21}.$$

For any $\nu \in [\omega_0]$, $\nu(\alpha_j)$ is a power of $\zeta^{12} = -1$ when $4|n$. Hence $\nu^5(\alpha_j) = \nu(\alpha_j)$. Thus in this case [\(4-6\)](#) equals

$$-2[2\nu_4(\alpha_2) + 2\nu_8(\alpha_2) + 2\nu_4(\alpha_3) + 2\nu_8(\alpha_3)] = -4[-1 + 1 - 1 + 1] = 0.$$

Acknowledgements

This work is based on Ragsdale’s Master’s thesis at the University of Maine. We thank the referee for carefully reading the manuscript and making several suggestions and corrections which have greatly improved the exposition. Ragsdale thanks the Department of Mathematics and Statistics of the University of Maine for summer support during the final stages of writing this paper. Both authors were supported by NSF grant DMS 0902145.

References

[Bump 1997] D. Bump, *Automorphic forms and representations*, Cambridge Studies in Advanced Mathematics **55**, Cambridge University Press, 1997. [MR 97k:11080](#) [Zbl 0868.11022](#)

[Bushnell and Henniart 2006] C. J. Bushnell and G. Henniart, *The local Langlands conjecture for $GL(2)$* , Grundlehren der Math. Wiss. **335**, Springer, Berlin, 2006. [MR 2007m:22013](#) [Zbl 1100.11041](#)

[Bushnell and Henniart 2014] C. J. Bushnell and G. Henniart, “Langlands parameters for epipelagic representations of GL_n ”, *Math. Ann.* **358**:1–2 (2014), 433–463. [arXiv 1302.4304](#)

[Bushnell and Kutzko 1993] C. J. Bushnell and P. C. Kutzko, *The admissible dual of $GL(N)$ via compact open subgroups*, Annals of Mathematics Studies **129**, Princeton University Press, 1993. [MR 94h:22007](#) [Zbl 0787.22016](#)

- [Casselman 1973] W. Casselman, “On some results of Atkin and Lehner”, *Math. Ann.* **201** (1973), 301–314. [MR 49 #2558](#) [Zbl 0239.10015](#)
- [Gross and Reeder 2010] B. H. Gross and M. Reeder, “Arithmetic invariants of discrete Langlands parameters”, *Duke Math. J.* **154**:3 (2010), 431–508. [MR 2012c:11252](#) [Zbl 1207.11111](#)
- [Kim 2007] J.-L. Kim, “Supercuspidal representations: an exhaustion theorem”, *J. Amer. Math. Soc.* **20**:2 (2007), 273–320. [MR 2008c:22014](#) [Zbl 1111.22015](#)
- [Knightly and Li 2006] A. Knightly and C. Li, *Traces of Hecke operators*, Mathematical Surveys and Monographs **133**, American Mathematical Society, Providence, RI, 2006. [MR 2008g:11090](#) [Zbl 1120.11024](#)
- [Knightly and Li 2012] A. Knightly and C. Li, “Modular L -values of cubic level”, *Pacific J. Math.* **260**:2 (2012), 527–563. [MR 3001804](#) [Zbl 06136442](#)
- [Mautner 1964] F. I. Mautner, “Spherical functions over b -adic fields, II”, *Amer. J. Math.* **86** (1964), 171–200. [MR 29 #3582](#) [Zbl 0135.17204](#)
- [Palm 2012] M. Palm, *Explicit $GL(2)$ trace formulas and uniform, mixed Weyl laws*, Ph.D. thesis, Göttingen, 2012. [arXiv 1212.4282](#)
- [Piatetski-Shapiro 1983] I. Piatetski-Shapiro, *Complex representations of $GL(2, K)$ for finite fields K* , Contemporary Mathematics **16**, American Mathematical Society, Providence, RI, 1983. [MR 84m:20046](#) [Zbl 0513.20026](#)
- [Reeder 1991] M. Reeder, “Old forms on GL_n ”, *Amer. J. Math.* **113**:5 (1991), 911–930. [MR 92i:22018](#) [Zbl 0758.11027](#)
- [Stevens 2008] S. Stevens, “The supercuspidal representations of p -adic classical groups”, *Invent. Math.* **172**:2 (2008), 289–352. [MR 2010e:22008](#) [Zbl 1140.22016](#)

Received: 2013-08-14

Revised: 2013-11-12

Accepted: 2013-11-16

knightly@math.umaine.edu

*Department of Mathematics and Statistics,
University of Maine, 5752 Neville Hall, Room 333,
Orono, ME 04469-5752, United States*

cwragsda@syrr.edu

*Department of Mathematics and Statistics,
University of Maine, 5752 Neville Hall, Room 333,
Orono, ME 04469-5752, United States*

EDITORS

MANAGING EDITOR

Kenneth S. Berenhaut, Wake Forest University, USA, berenhks@wfu.edu

BOARD OF EDITORS

Colin Adams	Williams College, USA colin.c.adams@williams.edu	David Larson	Texas A&M University, USA larson@math.tamu.edu
John V. Baxley	Wake Forest University, NC, USA baxley@wfu.edu	Suzanne Lenhart	University of Tennessee, USA lenhart@math.utk.edu
Arthur T. Benjamin	Harvey Mudd College, USA benjamin@hmc.edu	Chi-Kwong Li	College of William and Mary, USA ckli@math.wm.edu
Martin Bohner	Missouri U of Science and Technology, USA bohner@mst.edu	Robert B. Lund	Clemson University, USA lund@clemson.edu
Nigel Boston	University of Wisconsin, USA boston@math.wisc.edu	Gaven J. Martin	Massey University, New Zealand g.j.martin@massey.ac.nz
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA budhiraj@email.unc.edu	Mary Meyer	Colorado State University, USA meyer@stat.colostate.edu
Pietro Cerone	La Trobe University, Australia P.Cerone@latrobe.edu.au	Emil Minchev	Ruse, Bulgaria eminchev@hotmail.com
Scott Chapman	Sam Houston State University, USA scott.chapman@shsu.edu	Frank Morgan	Williams College, USA frank.morgan@williams.edu
Joshua N. Cooper	University of South Carolina, USA cooper@math.sc.edu	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran moslehian@ferdowsi.um.ac.ir
Jem N. Corcoran	University of Colorado, USA corcoran@colorado.edu	Zuhair Nashed	University of Central Florida, USA znashed@mail.ucf.edu
Toka Diagana	Howard University, USA tdiagana@howard.edu	Ken Ono	Emory University, USA ono@mathcs.emory.edu
Michael Dorff	Brigham Young University, USA mdorff@math.byu.edu	Timothy E. O'Brien	Loyola University Chicago, USA tbriell@luc.edu
Sever S. Dragomir	Victoria University, Australia sever@matilda.vu.edu.au	Joseph O'Rourke	Smith College, USA orourke@cs.smith.edu
Behrouz Emamizadeh	The Petroleum Institute, UAE bemamizadeh@pi.ac.ae	Yuval Peres	Microsoft Research, USA peres@microsoft.com
Joel Foisy	SUNY Potsdam foisyjs@potsdam.edu	Y.-F. S. Pétermann	Université de Genève, Switzerland petermann@math.unige.ch
Errin W. Fulp	Wake Forest University, USA fulp@wfu.edu	Robert J. Plemmons	Wake Forest University, USA rplemmons@wfu.edu
Joseph Gallian	University of Minnesota Duluth, USA kgallian@d.umn.edu	Carl B. Pomerance	Dartmouth College, USA carl.pomerance@dartmouth.edu
Stephan R. Garcia	Pomona College, USA stephan.garcia@pomona.edu	Vadim Ponomarenko	San Diego State University, USA vadim@sciences.sdsu.edu
Anant Godbole	East Tennessee State University, USA godbole@etsu.edu	Bjorn Poonen	UC Berkeley, USA poonen@math.berkeley.edu
Ron Gould	Emory University, USA rg@mathcs.emory.edu	James Propp	U Mass Lowell, USA jpropp@cs.uml.edu
Andrew Granville	Université Montréal, Canada andrew.andrew@dms.umontreal.ca	József H. Przytycki	George Washington University, USA przytyck@gwu.edu
Jerrold Griggs	University of South Carolina, USA griggs@math.sc.edu	Richard Rebarber	University of Nebraska, USA rrebarbe@math.unl.edu
Sat Gupta	U of North Carolina, Greensboro, USA sgupta@uncg.edu	Robert W. Robinson	University of Georgia, USA rwr@cs.uga.edu
Jim Haglund	University of Pennsylvania, USA jhaglund@math.upenn.edu	Filip Saidak	U of North Carolina, Greensboro, USA f_saidak@uncg.edu
Johnny Henderson	Baylor University, USA johnny_henderson@baylor.edu	James A. Sellers	Penn State University, USA sellersj@math.psu.edu
Jim Hoste	Pitzer College jhoste@pitzer.edu	Andrew J. Sterge	Honorary Editor andy@ajsterge.com
Natalia Hritonenko	Prairie View A&M University, USA nahritonenko@pvamu.edu	Ann Trenk	Wellesley College, USA atrenk@wellesley.edu
Glenn H. Hurlbert	Arizona State University, USA hurlbert@asu.edu	Ravi Vakil	Stanford University, USA vakill@math.stanford.edu
Charles R. Johnson	College of William and Mary, USA crjohnso@math.wm.edu	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy antonia.vecchio@cnr.it
K. B. Kulasekera	Clemson University, USA kk@ces.clemson.edu	Ram U. Verma	University of Toledo, USA verma99@msn.com
Gerry Ladas	University of Rhode Island, USA gladas@math.uri.edu	John C. Wierman	Johns Hopkins University, USA wierman@jhu.edu
		Michael E. Zieve	University of Michigan, USA zieve@umich.edu

PRODUCTION

Silvio Levy, Scientific Editor

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2014 is US \$120/year for the electronic version, and \$165/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscribers address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2014 Mathematical Sciences Publishers

involve

2014

vol. 7

no. 5

Infinite cardinalities in the Hausdorff metric geometry	585
ALEXANDER ZUPAN	
Computing positive semidefinite minimum rank for small graphs	595
STEVEN OSBORNE AND NATHAN WARNBERG	
The complement of Fermat curves in the plane	611
SETH DUTTER, MELISSA HAIRE AND ARIEL SETNIKER	
Quadratic forms representing all primes	619
JUSTIN DEBENEDETTO	
Counting matrices over a finite field with all eigenvalues in the field	627
LISA KAYLOR AND DAVID OFFNER	
A not-so-simple Lie bracket expansion	647
JULIE BEIER AND MCCABE OLSEN	
On the omega values of generators of embedding dimension-three numerical monoids generated by an interval	657
SCOTT T. CHAPMAN, WALTER PUCKETT AND KATY SHOUR	
Matrix coefficients of depth-zero supercuspidal representations of $GL(2)$	669
ANDREW KNIGHTLY AND CARL RAGSDALE	
The sock matching problem	691
SARAH GILLIAND, CHARLES JOHNSON, SAM RUSH AND DEBORAH WOOD	
Superlinear convergence via mixed generalized quasilinearization method and generalized monotone method	699
VINCHENCIA ANDERSON, COURTNEY BETTIS, SHALA BROWN, JACQKIS DAVIS, NAEEM TULL-WALKER, VINODH CHELLAMUTHU AND AGHALAYA S. VATSALA	