

involve

a journal of mathematics

A solution to a problem of Frechette and Locus

Chenthuran Abeyakaran



A solution to a problem of Frechette and Locus

Chenthuran Abeyakaran

(Communicated by Ken Ono)

In a recent paper, Frechette and Locus examined and found expressions for the infinite product $D_m(q) := \prod_{t=1}^{\infty} (1 - q^{mt}) / (1 - q^t)$ in terms of products of q -series of the Rogers–Ramanujan type coming from Hall–Littlewood polynomials, when $m \equiv 0, 1, 2 \pmod{4}$. These q -series were originally discovered in 2014 by Griffin, Ono, and Warnaar in their work on the framework of the Rogers–Ramanujan identities. Extending this framework, Rains and Warnaar also recently discovered more q -series and their corresponding infinite products. Frechette and Locus left open the case where $m \equiv 3 \pmod{4}$. Here we find such an expression for the infinite products for $m \equiv 3 \pmod{4}$ by making use of the new q -series obtained by Rains and Warnaar.

1. Introduction

The Rogers–Ramanujan identities [Andrews 1971]

$$G(q) := \sum_{n=0}^{\infty} \frac{q^{n^2}}{(1-q)(1-q^2)\dots(1-q^n)} = \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})}, \quad (1-1)$$

$$H(q) := \sum_{n=0}^{\infty} \frac{q^{n^2+n}}{(1-q)(1-q^2)\dots(1-q^n)} = \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+2})(1-q^{5n+3})}, \quad (1-2)$$

have inspired research and discoveries in many areas of mathematics and physics, such as modular forms and elliptic curves, conformal field theory, knot theory, probability, and statistical mechanics. (See next citation for some discussion.) Given the importance of these identities, it had been an open problem for nearly a century to build a theory suggested by these two Rogers–Ramanujan identities. In 2014 Griffin, Ono, and Warnaar [Griffin et al. 2016] discovered¹ a more general framework for identities similar to that of Rogers–Ramanujan, where an infinite sum, defined using Hall–Littlewood polynomials $P_{\lambda}(x; q)$, is equal to an infinite product with periodic exponents.

MSC2010: 11P84.

Keywords: Rogers–Ramanujan Identities.

¹Their work was named the 15th top story in science in 2014 by *Discover* magazine.

In order to define the Hall–Littlewood polynomials, we recall the definition of an integer partition and the following notation. A *partition* is a nonincreasing sequence of nonnegative integers with finitely many nonzero terms. For a partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$, we define the *weight* of the partition to be $|\lambda| := \lambda_1 + \lambda_2 + \dots + \lambda_n$ and the *length* of the partition λ to be n . In addition, we let $2\lambda := (2\lambda_1, 2\lambda_2, \dots, 2\lambda_n)$. Let m_i denote the multiplicity of size i parts. Also, let $(q)_k = (q; q)_k$ denote the q -Pochhammer symbol, which is defined as follows:

$$(a)_k := (a; q)_k = \begin{cases} (1 - a)(1 - aq)(1 - aq^2) \cdots (1 - aq^{k-1}) & \text{if } k \geq 0, \\ \prod_{n=0}^{\infty} (1 - aq^n) & \text{if } k = \infty. \end{cases}$$

If λ has length n , the Hall–Littlewood polynomial is a symmetric function in n variables, namely x_1, x_2, \dots, x_n , defined as

$$P_\lambda(x; q) = \frac{1}{v_\lambda(q)} \sum_{w \in S_n} w \left(x^\lambda \prod_{i < j} \frac{x_i - qx_j}{x_i - x_j} \right), \tag{1-3}$$

where $x^\lambda := x_1^{\lambda_1} x_2^{\lambda_2} x_3^{\lambda_3} \dots x_n^{\lambda_n}$, $v_\lambda(q) := \prod_{i=0}^n (q)_{m_i} / (1 - q)^{m_i}$, and the symmetric group S_n acts on x by permuting all the x_i .

For ordered pairs $\nu = (c, d) \in \{(1, -1), (2, -1), (1, 0), (2, -2)\}$ and arbitrary $a, b \geq 1$, Griffin, Ono, and Warnaar [Griffin et al. 2016], and more recently Rains and Warnaar [2015], defined the q -series

$$R_\nu(a, b; q) = \sum_{\lambda, \lambda_1 \leq a} q^{c|\lambda|} P_{2\lambda}(1, q, q^2, \dots; q^{2b+d}), \tag{1-4}$$

$$S(a, b; q) = \sum_{\lambda, \lambda_1 \leq a} q^{|\lambda|/2} P_\lambda(1, q, q^2, \dots; q^{2b}), \tag{1-5}$$

and

$$T(a, b; q) = \sum_{\lambda, \lambda_1 \leq a} q^{|\lambda|} \left(\prod_{i=1}^{a-1} (-q^{b-1/2}; q^{b-1/2})_{m_i(\lambda)} \right) P_\lambda(1, q, q^2, \dots; q^{2b-1}). \tag{1-6}$$

Here the $P_\lambda(1, q, q^2, \dots; q^T)$ are Hall–Littlewood q -series in infinitely many parameters. To define these Hall–Littlewood q -series, we must first express the Hall–Littlewood polynomial $P_\lambda(x_1, x_2, \dots, x_n; q^T)$ in terms of the r -th power sum symmetric function, $x_1^r + x_2^r + \dots + x_n^r$. This is possible due to a well-known fact in abstract algebra which states that every symmetric polynomial can be written as a sum of products of r -th power sum symmetric functions with rational coefficients [Macdonald 1995]. Now we obtain the Hall–Littlewood polynomial $P_\lambda(1, q, q^2, \dots; q^T)$ by replacing $(x_1^r + x_2^r + \dots + x_n^r)$ with $1^r + q^r + q^{2r} + \dots = 1/(1 - q^r)$.

To motivate our work, consider another interesting property of the Roger–Ramanujan identities. When we take the product of both Rogers–Ramanujan

identities, we can see that

$$\begin{aligned} G(q) \cdot H(q) &= \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})} \cdot \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+2})(1-q^{5n+3})} \\ &= \prod_{n=0}^{\infty} \frac{1}{(1-q^{5n+1})(1-q^{5n+4})(1-q^{5n+2})(1-q^{5n+3})} \\ &= \prod_{n=1}^{\infty} \frac{(1-q^{5n})}{(1-q^n)}. \end{aligned}$$

Inspired by this emergence of this infinite product, Frechette and Locus [2016] explored the natural question of more generalized products of q -series of the Rogers–Ramanujan type that result in an infinite product of the form

$$D_m(q) := \prod_{t=1}^{\infty} \frac{(1-q^{mt})}{(1-q^t)}. \tag{1-7}$$

Making use of the Rogers–Ramanujan framework of [Griffin et al. 2016], Frechette and Locus obtained explicit formulas for $D_m(q)$ when $m \equiv 0, 1, 2 \pmod{4}$. When $m \geq 8$ and is even, they found that

$$D_m(q) = \frac{R_{(1,0)}\left(2, \frac{m}{2} - 3; q\right) \cdot R_{(2,-2)}\left(\frac{m}{2}, 2; q\right)}{R_{(2,-2)}\left(\frac{m}{2} - 3, 3; q\right)}. \tag{1-8}$$

They also found for $m \equiv 1 \pmod{4}$ and $m > 1$,

$$D_m(q) = \frac{R_{(1,-1)}\left(\frac{m-1}{2} - 1, 1; q\right) \cdot R_{(2,-1)}\left(\frac{m-1}{4}, \frac{m-1}{4}; q\right)}{R_{(2,-1)}\left(\frac{m-1}{4} + 1, \frac{m-1}{4} - 1; q\right)}. \tag{1-9}$$

However, they were unable to construct $D_m(q)$ for positive integers $m \equiv 3 \pmod{4}$. Their difficulty arose from the fact that the Rogers–Ramanujan framework of [Griffin et al. 2016]. Rains and Warnaar [2015] recently found this extension. In this paper, we address this case and provide such a formula.

Theorem 1.1. *If $m \equiv 3 \pmod{4}$ and $m \geq 7$, we have*

$$D_m(q) = \frac{T\left(\frac{m+1}{2}, \frac{m+1}{4}; q\right) S\left(\frac{m+1}{2}, \frac{m+1}{4} - 1; q\right) R_{(1,-1)}\left(\frac{m-1}{2} - 1, 1; q\right)}{T\left(\frac{m+1}{2} + 2, \frac{m+1}{4} - 1; q\right) S\left(\frac{m+1}{2} - 2, \frac{m+1}{4}; q\right)}.$$

In Section 2, we cover preliminaries on q -series and state the results of [Griffin et al. 2016; Rains and Warnaar 2015; Frechette and Locus 2016]. In Section 3, we use these results to prove Theorem 1.1.

2. Preliminaries

In order to simplify the products involved in writing $R_\nu(a, b; q)$, $S(a, b; q)$, and $T(a, b; q)$, we use a modified theta function

$$\theta(a; q) := (a; q)_\infty (q/a; q)_\infty, \tag{2-1}$$

where $(a; q)_\infty$ denotes the q -Pochhammer symbol we previously defined. We then define

$$\theta(a_1, a_2, \dots, a_n; q) = \theta(a_1; q) \cdot \theta(a_2; q) \cdots \theta(a_n; q). \tag{2-2}$$

Theorem 2.1 [Griffin et al. 2016, Theorem 1.1]. *If a and b are positive integers and $\kappa := 2a + 2b + 1$, we have*

$$\begin{aligned} R_{(1,-1)}(a, b; q) &:= \sum_{\lambda, \lambda_1 \leq a} q^{|\lambda|} P_{2\lambda}(1, q, q^2, \dots; q^{2b-1}) \\ &= \frac{(q^\kappa; q^\kappa)_\infty^b}{(q)_\infty^b} \cdot \prod_{i=1}^b \theta(q^{i+a}; q^\kappa) \prod_{1 \leq i < j \leq b} \theta(q^{j-i}, q^{i+j-1}; q^\kappa) \\ &= \frac{(q^\kappa; q^\kappa)_\infty^a}{(q)_\infty^a} \cdot \prod_{i=1}^a \theta(q^{i+1}; q^\kappa) \prod_{1 \leq i < j \leq a} \theta(q^{j-i}, q^{i+j+1}; q^\kappa). \end{aligned}$$

This result has generalizations to the functions S and T :

Theorem 2.2 [Rains and Warnaar 2015, Theorem 5.10]. *If a and b are positive integers and $\kappa := a + 2b + 1$, we have*

$$\begin{aligned} S(a, b; q) &:= \sum_{\lambda, \lambda_1 \leq a} q^{|\lambda|/2} P_\lambda(1, q, q^2, \dots; q^{2b}) \\ &= \frac{(q^\kappa; q^\kappa)_\infty^{b-1} (q^{\kappa/2}; q^{\kappa/2})_\infty}{(q)_\infty^{b-1} (q^{1/2}; q^{1/2})_\infty} \prod_{i=1}^b \theta(q^i; q^{\kappa/2}) \prod_{1 \leq i < j \leq b} \theta(q^{j-i}, q^{i+j}; q^\kappa). \end{aligned}$$

Theorem 2.3 [Rains and Warnaar 2015, Remark 5.13]. *If a and b are positive integers and $\kappa := a + 2b - 1$, we have*

$$\begin{aligned} T(a, b; q) &:= \sum_{\lambda, \lambda_1 \leq a} q^{|\lambda|} \left(\prod_{i=1}^{a-1} (-q^{b-1/2}; q^{b-1/2})_{m_i(\lambda)} \right) P_\lambda(1, q, q^2, \dots; q^{2b-1}) \\ &= \frac{(q^\kappa; q^\kappa)_\infty^b}{(q)_\infty^{b-1} (q^{1/2}; q^{1/2})_\infty} \prod_{i=1}^b \theta(q^{i-1/2}; q^\kappa) \prod_{1 \leq i < j \leq b} \theta(q^{j-i}, q^{i+j-1}; q^\kappa). \end{aligned}$$

To prove our result, we combine **Theorem 2.3** with the following proposition.

Proposition 2.4. *If $i \in \mathbb{Z}^+$, we have*

$$\theta(q^i; q^{m/2}) = \theta(q^i; q^m)\theta(q^{m/2-i}; q^m).$$

Proof. Using the definition of the modified theta function, we have

$$\begin{aligned} \theta(q^i; q^{m/2}) &= (q^i; q^{m/2})_\infty (q^{m/2-i}; q^{m/2})_\infty \\ &= \prod_{n=0}^\infty (1 - q^i \cdot q^{mn/2}) \prod_{n=0}^\infty (1 - q^{m/2-i} \cdot q^{mn/2}). \end{aligned}$$

Separating terms in the infinite product on the right-hand side based on the parity of n , we have

$$\begin{aligned} \theta(q^i; q^{m/2}) &= \prod_{n=0}^\infty (1 - q^i \cdot q^{mn}) (1 - q^{i+m/2} \cdot q^{mn}) \prod_{n=0}^\infty (1 - q^{m/2-i} \cdot q^{mn}) (1 - q^{m-i} \cdot q^{mn}) \\ &= \prod_{n=0}^\infty (1 - q^i \cdot q^{mn}) (1 - q^{m-i} \cdot q^{mn}) (1 - q^{m/2-i} \cdot q^{mn}) (1 - q^{i+m/2} \cdot q^{mn}) \\ &= (q^i; q^m)_\infty (q^{m-i}; q^m)_\infty (q^{m/2-i}; q^m)_\infty (q^{m/2+i}; q^m)_\infty \\ &= \theta(q^i; q^m)\theta(q^{m/2-i}; q^m). \quad \square \end{aligned}$$

3. Proof of Theorem 1.1

We shall now prove Theorem 1.1. By Theorem 2.3, when $m = a + 2b - 1$, we have

$$\begin{aligned} T\left(\frac{m+1}{2}, \frac{m+1}{4}; q\right) &= \frac{(q^m; q^m)_{\infty}^{(m+1)/4}}{(q)_{\infty}^{(m+1)/4-1} (q^{1/2}; q^{1/2})_{\infty}} \prod_{i=1}^{(m+1)/4} \theta(q^{i-1/2}; q^m) \\ &\quad \prod_{1 \leq i < j \leq (m+1)/4} \theta(q^{j-i}, q^{i+j-1}; q^m) \quad (3-1) \end{aligned}$$

and

$$\begin{aligned} T\left(\frac{m+1}{2} + 2, \frac{m+1}{4} - 1; q\right) &= \frac{(q^m; q^m)_{\infty}^{(m-3)/4}}{(q)_{\infty}^{(m-7)/4} (q^{1/2}; q^{1/2})_{\infty}} \prod_{i=1}^{(m-3)/4} \theta(q^{i-1/2}; q^m) \\ &\quad \prod_{1 \leq i < j \leq (m-3)/4} \theta(q^{j-i}, q^{i+j-1}; q^m), \quad (3-2) \end{aligned}$$

which gives

$$\begin{aligned}
 & \frac{T\left(\frac{m+1}{2}, \frac{m+1}{4}; q\right)}{T\left(\frac{m+1}{2} + 2, \frac{m+1}{4} - 1; q\right)} \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m-1)/4}; q^m) \prod_{i=1}^{(m+1)/4-1} \theta(q^{(m+1)/4-i}, q^{(m+1)/4-1+i}; q^m) \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m-1)/4}; q^m) \prod_{i=1}^{(m+1)/4-1} \theta(q^i; q^m) \prod_{i=(m+1)/4}^{(m+1)/2-2} \theta(q^i; q^m) \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m-1)/4}; q^m) \prod_{i=1}^{(m-1)/2} \theta(q^i; q^m). \tag{3-3}
 \end{aligned}$$

Using [Theorem 2.2](#) and [Proposition 2.4](#), we have

$$\begin{aligned}
 & S\left(\frac{m+1}{2} - 2, \frac{m+1}{4}; q\right) \\
 &= \frac{(q^m; q^m)_\infty^{(m+1)/4-1} (q^{m/2}; q^{m/2})_\infty}{(q)_\infty^{(m+1)/4-1} (q^{1/2}; q^{1/2})_\infty} \prod_{i=1}^{(m+1)/4} \theta(q^i; q^m) \theta(q^{m/2-i}; q^m) \\
 & \quad \prod_{1 \leq i < j \leq (m+1)/4} \theta(q^{j-i}, q^{i+j}; q^m), \tag{3-4}
 \end{aligned}$$

and

$$\begin{aligned}
 & S\left(\frac{m+1}{2}, \frac{m+1}{4} - 1; q\right) \\
 &= \frac{(q^m; q^m)_\infty^{(m+1)/4-2} (q^{m/2}; q^{m/2})_\infty}{(q)_\infty^{(m+1)/4-2} (q^{1/2}; q^{1/2})_\infty} \prod_{i=1}^{(m+1)/4-1} \theta(q^i; q^m) \theta(q^{m/2-i}; q^{m/2}) \\
 & \quad \prod_{1 \leq i < j \leq (m+1)/4-1} \theta(q^{j-i}, q^{i+j}; q^m). \tag{3-5}
 \end{aligned}$$

Now, we evaluate the following quotient:

$$\begin{aligned}
 & \frac{S\left(\frac{m+1}{2} - 2, \frac{(m+1)}{4}; q\right)}{S\left(\frac{m+1}{2}, \frac{m+1}{4} - 1; q\right)} \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m+1)/4}; q^m) \theta(q^{(m-1)/4}; q^m) \\
 & \quad \prod_{i=1}^{(m+1)/4-1} \theta(q^{(m+1)/4-i}, q^{(m+1)/4+i}; q^m)
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m+1)/4}; q^m) \theta(q^{(m-1)/4}; q^m) \\
 &\quad \prod_{i=1}^{(m+1)/4-1} \theta(q^i; q^m) \prod_{i=(m+1)/4+1}^{(m+1)/2-1} \theta(q^i; q^m) \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m-1)/4}; q^m) \prod_{i=1}^{(m-1)/2} \theta(q^i; q^m). \tag{3-6}
 \end{aligned}$$

Dividing (3-3) by (3-6), we obtain

$$\begin{aligned}
 &\frac{T\left(\frac{m+1}{2}, \frac{m+1}{4}; q\right) S\left(\frac{m+1}{2}, \frac{m+1}{4} - 1; q\right)}{T\left(\frac{m+1}{2} + 2, \frac{m+1}{4} - 1; q\right) S\left(\frac{m+1}{2} - 2, \frac{m+1}{4}; q\right)} \\
 &= \frac{(q^m; q^m)_\infty / (q)_\infty \theta(q^{(m-1)/4}; q^m) \prod_{i=1}^{(m-3)/2} \theta(q^i; q^m)}{(q^m; q^m)_\infty / (q)_\infty \theta(q^{(m-1)/4}; q^m) \prod_{i=1}^{(m-1)/2} \theta(q^i; q^m)} \tag{3-7} \\
 &= \frac{1}{\theta(q^{(m-1)/2}; q^m)}.
 \end{aligned}$$

By Theorem 2.1, we have the identity

$$R_{(1,-1)}\left(\frac{m-1}{2} - 1, 1; q\right) = \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m-1)/2}; q^m). \tag{3-8}$$

Multiplying (3-8) and (3-7) gives us our desired result:

$$\begin{aligned}
 &\frac{T\left(\frac{m+1}{2} + 1, \frac{m+1}{4}; q\right) S\left(\frac{m+1}{2}, \frac{m+1}{4} - 1; q\right) R_{(1,-1)}\left(\frac{m-1}{2} - 1, 1; q\right)}{T\left(\frac{m+1}{2} + 3, \frac{m+1}{4} - 1; q\right) S\left(\frac{m+1}{2} - 2, \frac{m+1}{4}; q\right)} \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} \theta(q^{(m-1)/2}; q^m) \cdot \frac{1}{\theta(q^{(m-1)/2}; q^m)} \\
 &= \frac{(q^m; q^m)_\infty}{(q)_\infty} = \prod_{t=1}^{\infty} \frac{(1-q^{mt})}{(1-q^t)} = D_m(q). \quad \square
 \end{aligned}$$

Acknowledgements

I would like to thank Ken Ono for introducing me to this problem and for his guidance throughout this project. I would also like to thank Tessa Cotron, Sarah Fleming, and Robert Dicks for proofreading this paper and checking my work.

References

[Andrews 1971] G. E. Andrews, *Number theory*, Saunders, Philadelphia, 1971. Corrected reprint by Dover, New York, 1994. [MR](#) [Zbl](#)

- [Frechette and Locus 2016] C. Frechette and M. Locus, “Combinatorial properties of Rogers–Ramanujan type identities arising from Hall–Littlewood polynomials”, *Ann. Comb.* **20**:2 (2016), 345–360. [MR](#) [Zbl](#)
- [Griffin et al. 2016] M. J. Griffin, K. Ono, and S. O. Warnaar, “A framework of Rogers–Ramanujan identities and their arithmetic properties”, *Duke Math. J.* **165**:8 (2016), 1475–1527. [MR](#) [Zbl](#)
- [Macdonald 1995] I. G. Macdonald, *Symmetric functions and Hall polynomials*, 2nd ed., Oxford University Press, New York, 1995. [MR](#) [Zbl](#)
- [Rains and Warnaar 2015] E. M. Rains and S. O. Warnaar, “Bounded Littlewood identities”, preprint, 2015. [arXiv](#)

Received: 2016-07-29

Revised: 2016-08-18

Accepted: 2016-08-21

ChenthuranJA@gmail.com

*Chamblee Charter High School, 3688 Chamblee Dunwoody Rd,
Chamblee, GA 30341, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor


Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

involve

2017

vol. 10

no. 5

Algorithms for finding knight's tours on Aztec diamonds	721
SAMANTHA DAVIES, CHENXIAO XUE AND CARL R. YERGER	
Optimal aggression in kleptoparasitic interactions	735
DAVID G. SYKES AND JAN RYCHTÁŘ	
Domination with decay in triangular matchstick arrangement graphs	749
JILL COCHRAN, TERRY HENDERSON, AARON OSTRANDER AND RON TAYLOR	
On the tree cover number of a graph	767
CHASSIDY BOZEMAN, MINERVA CATRAL, BRENDAN COOK, OSCAR E. GONZÁLEZ AND CAROLYN REINHART	
Matrix completions for linear matrix equations	781
GEOFFREY BUHL, ELIJAH CRONK, ROSA MORENO, KIRSTEN MORRIS, DIANNE PEDROZA AND JACK RYAN	
The Hamiltonian problem and t -path traceable graphs	801
KASHIF BARI AND MICHAEL E. O'SULLIVAN	
Relations between the conditions of admitting cycles in Boolean and ODE network systems	813
YUNJIAO WANG, BAMIDELE OMIDIRAN, FRANKLIN KIGWE AND KIRAN CHILAKAMARRI	
Weak and strong solutions to the inverse-square brachistochrone problem on circular and annular domains	833
CHRISTOPHER GRIMM AND JOHN A. GEMMER	
Numerical existence and stability of steady state solutions to the distributed spruce budworm model	857
HALA AL-KHALIL, CATHERINE BRENNAN, ROBERT DECKER, ASLIHAN DEMIRKAYA AND JAMIE NAGODE	
Integer solutions to $x^2 + y^2 = z^2 - k$ for a fixed integer value k	881
WANDA BOYER, GARY MACGILLIVRAY, LAURA MORRISON, C. M. (KIEKA) MYNHARDT AND SHAHLA NASSERASR	
A solution to a problem of Frechette and Locus	893
CHENTHURAN ABEYAKARAN	