

ON INTERTWINING BY AN OPERATOR HAVING A DENSE RANGE

EITOKU GOYA AND TEISHIRÔ SAITÔ

(Received February 28, 1980)

1. Throughout the paper, by an operator we mean a bounded linear transformation acting on a Hilbert space H . The algebra of all operators on H is denoted by $B(H)$.

We formulate an algebraic version of generalized Putnam-Fuglede theorem [3; Theorem 1], and we show that a paranormal contraction T is unitary, if S is a coisometry, if W is an operator having a dense range and if $TW = WS$. This is a generalization of a result due to Okubo [1].

Let $T \in B(H)$. T is *hyponormal* (resp. *cohyponormal*) if $T^*T - TT^* \geq 0$ (resp. $TT^* - T^*T \geq 0$). T is *dominant* if $\text{range}(T - \lambda) \subset \text{range}(T - \lambda)^*$ for all $\lambda \in \sigma(T)$, the spectrum of T . This condition is equivalent to the existence of a constant M_λ for each $\lambda \in \sigma(T)$ such that

$$\|(T - \lambda)^*x\| \leq M_\lambda \|(T - \lambda)x\|$$

for all $x \in H$. Thus every hyponormal operator is dominant. T is *paranormal* if

$$\|Tx\|^2 \leq \|T^2x\| \|x\|$$

for all $x \in H$.

2. The following theorem is a version of [3; Theorem 1]. The proof of [3] applies to this version. We include it for completeness.

THEOREM 1. *Let T, S , and $W \in B(H)$, where W has a dense range. Assume that $TW = WS$ and $T^*W = WS^*$. Then*

- (i) *T is hyponormal (resp. cohyponormal), if so is S .*
- (ii) *T is isometric (resp. coisometric), if so is S . In particular, T is unitary, if so is S .*
- (iii) *T is normal, if so is S .*

PROOF. Let $W^* = V^*B$ be the polar decomposition of W^* . Since W has a dense range, W^* is injective. Thus $B^2 = WW^*$ is injective, and V is coisometric. From equations $TW = WS$ and $T^*W = WS^*$, we have

$$TWW^* = WSW^*, \quad WW^*T = WSW^*.$$

Thus, WW^* commutes with T , and so B commutes with T . Hence we have

$$BTV = TBV = TW = WS = BVS,$$

which implies that $TV = VS$ because B is injective. Since V is coisometric, we obtain

$$T = TVV^* = VSV^*.$$

From the equations $W^*T = SW^*$ and $TB = BT$, we have

$$V^*TB = V^*BT = W^*T = SW^* = SV^*B,$$

which implies that $V^*T = SV^*$. Hence

$$V^*VS = V^*TV = SV^*V.$$

First we assume that S is normal. Since $S^*S = SS^*$, we obtain

$$\begin{aligned} T^*T &= (VSV^*)^*(VSV^*) = VS^*V^*VSV^* = VS^*SV^*VV^* = VS^*SV^* \\ &= VSS^*V^* = VV^*VSS^*V^* = (VSV^*)(VSV^*)^* = TT^*, \end{aligned}$$

whence T is normal.

To prove (i), assume that S is hyponormal (resp. cohyponormal). Since $S^*S \geq SS^*$ (resp. $SS^* \geq S^*S$), the above computation implies that

$$\begin{aligned} T^*T &= VS^*SV^* \geq VSS^*V^* = TT^*, \\ (\text{resp. } TT^* &= VSS^*V^* \geq VS^*SV^* = T^*T), \end{aligned}$$

and the assertion of (i) follows.

To prove (ii), assume that S is isometric (resp. coisometric). Again, by the above computation,

$$T^*T = VS^*SV^* = VV^* = I, \quad (\text{resp. } TT^* = VSS^*V^* = VV^* = I),$$

whence T is isometric (resp. coisometric).

The rest of the theorem is obvious.

REMARK. In Theorem 1, if W is injective and has a dense range, V is a unitary operator which implements the unitary equivalence of S and T .

The next theorem is a generalization of [1; Proposition 1].

THEOREM 2. *Let T , V , and $W \in B(H)$, where T is a paranormal contraction, V is a coisometry and W has a dense range. Assume that $TW = WV$. Then T is a unitary operator. In particular, if W is injective and has a dense range, then V is also a unitary operator.*

PROOF. Let $x \in H$ such that $Wx \neq 0$, and define

$$y_n = WV^{*n}x \quad (n = 0, 1, 2, \dots).$$

Then we have

$$Ty_{n+1} = TWV^{*(n+1)}x = WV^{*(n+1)}x = WV^{*n}x = y_n.$$

Since T is a contraction,

$$\|y_n\| = \|Ty_{n+1}\| \leq \|y_{n+1}\| = \|WV^{*(n+1)}x\| \leq \|W\| \|x\|$$

and hence $\{\|y_{n+1}\|\}$ is a monotone increasing convergent sequence. By the paranormality of T , we have

$$\|y_n\|^2 = \|Ty_{n+1}\|^2 \leq \|T^2y_{n+1}\| \|y_{n+1}\| = \|y_{n-1}\| \|y_{n+1}\|$$

and

$$1 \geq \frac{\|y_0\|}{\|y_1\|} \geq \frac{\|y_1\|}{\|y_2\|} \geq \dots \geq \frac{\|y_{n-1}\|}{\|y_n\|} \rightarrow 1 \quad (n \rightarrow \infty).$$

In particular, $\|y_0\| = \|y_1\|$, that is,

$$\|Wx\| = \|WV^*x\|.$$

Thus

$$\|WV^*x\| = \|Wx\| = \|WV^{*2}x\| = \|TWV^*x\| \leq \|WV^*x\|,$$

and so

$$\|WV^*x\| = \|Wx\| = \|TWV^*x\|.$$

Note that these equalities are valid for $x \in H$ such that $Wx \neq 0$. Hence

$$\begin{aligned} & \|T^*Wx - WV^*x\|^2 \\ &= \|T^*Wx\|^2 + \|WV^*x\|^2 - (T^*Wx, WV^*x) - (WV^*x, T^*Wx) \\ &\leq 2\|Wx\|^2 - (Wx, TWV^*x) - (TWV^*x, Wx) \\ &= 2\|Wx\|^2 - (Wx, WV^{*2}x) - (WV^{*2}x, Wx) \\ &= 2\|Wx\|^2 - 2\|Wx\|^2 = 0 \end{aligned}$$

for all $x \in H$, and $TW^* = WV^*$. It follows from Theorem 1 that T is a coisometry. Since T is paranormal, T is unitary by [2; Lemma 3]. The rest is clear by the remark after Theorem 1.

REMARK. Our proof of Theorem 2 is a modification of the argument due to Okubo [1]. He proved Theorem 2 under the hypothesis that V is unitary.

COROLLARY 3. Let $T \in B(H)$ be a paranormal contraction. Let $TW = WV$, where $V \in B(H)$ is a coisometry and $W \in B(H)$ is any non-

zero operator. Then T has a nontrivial invariant subspace.

PROOF. Let \mathfrak{M} be the closure of $\text{range } W$. If W does not have the dense range, \mathfrak{M} is a nontrivial invariant subspace of T . If W has the dense range, then T is unitary by Theorem 2, and T has a nontrivial invariant subspace.

3. As an application of Theorems 1 and 2, we give an alternative proof to the following theorem.

THEOREM 4. Let $T \in B(H)$ be a contraction. Let

$$\mathfrak{M} = \{x \in H \mid \|T^{*n}x\| \rightarrow 0 \ (n \rightarrow \infty)\}.$$

If T is dominant or paranormal, then \mathfrak{M} is a reducing subspace for T such that $T|_{\mathfrak{M}^\perp}$ is unitary and $T|_{\mathfrak{M}}$ is completely non-unitary (i.e., $T|_{\mathfrak{M}}$ has no nontrivial reducing subspace on which $T|_{\mathfrak{M}}$ is unitary).

This theorem was first proved for dominant operators in [4] and for paranormal operators in [1]. Note that the statements in [4; Theorem 2] contain a slip, because $\{x \in H \mid \|T^{*n}x\| \geq \varepsilon_x > 0\}$ is not a linear subspace of H .

To prove Theorem 4, we need the following simple lemma.

LEMMA 5. Let $T \in B(H)$ be a contraction. Let $\mathfrak{M} \subset H$ be an invariant subspace for T . If $T|_{\mathfrak{M}}$ is a coisometry, then \mathfrak{M} reduces T .

PROOF. Let $S = T|_{\mathfrak{M}}$, and let $x \in \mathfrak{M}$. Then, since S^* is isometric

$$\begin{aligned} \|S^*x - T^*x\|^2 &= \|S^*x\|^2 + \|T^*x\|^2 - (S^*x, T^*x) - (T^*x, S^*x) \\ &\leq \|x\|^2 + \|x\|^2 - (TS^*x, x) - (x, TS^*x) \\ &= 2\|x\|^2 - 2\|S^*x\|^2 = 2\|x\|^2 - 2\|x\|^2 = 0. \end{aligned}$$

Thus, $T^*x = (T|_{\mathfrak{M}})^*x \in \mathfrak{M}$ for all $x \in \mathfrak{M}$, which implies that \mathfrak{M} is invariant under T^* .

PROOF OF THEOREM 4. Since $\|T\| \leq 1$, the sequence $\{T^n T^{*n}\}$ converges strongly to a positive contraction. Let

$$A = (\lim_{n \rightarrow \infty} T^n T^{*n})^{1/2}.$$

Then, $\mathfrak{M} = \ker A$ and $TA^2T^* = A^2$. Since

$$\|AT^*x\|^2 = (TA^2T^*x, x) = (A^2x, x) = \|Ax\|^2$$

for all $x \in H$, there exists a partial isometry $W \in B(H)$ such that

$$AT^* = WA, \quad W|_{\mathfrak{M}} = 0.$$

It is easy to see that \mathfrak{M}^\perp is invariant under T . Let us write the equa-

tion $AT^* = WA$ in matrix form on $H = \mathfrak{M} \oplus \mathfrak{M}^\perp$. Then

$$\begin{bmatrix} 0 & 0 \\ 0 & A_1 \end{bmatrix} \begin{bmatrix} S_1 & S_2 \\ 0 & S_3 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & W_1 \end{bmatrix} \begin{bmatrix} 0 & 0 \\ 0 & A_1 \end{bmatrix},$$

whence $A_1 S_3 = W_1 A_1$, or $S_3^* A_1 = A_1 W_1^*$. Note that $A_1 = A|_{\mathfrak{M}^\perp}$ is injective and has a dense range, and $W_1 = W|_{\mathfrak{M}^\perp}$ is an isometry.

Case 1. Assume that T is dominant. Since S_3^* is dominant and W_1^* is coisometric, S_3^* and W_1^* are unitarily equivalent normal operators by [4; Theorem 1] and the remark after Theorem 1. Thus \mathfrak{M}^\perp reduces T by [3; Lemma 2]. Since W_1 is normal and isometric, W_1 is unitary and so is S_3 .

Case 2. Assume that T is paranormal. Since $S_3^* = T|_{\mathfrak{M}^\perp}$ is paranormal, S_3^* is unitary by Theorem 2. Thus \mathfrak{M}^\perp reduces T by Lemma 5.

It is clear that $T|_{\mathfrak{M}}$ is completely non-unitary in each case.

REMARK. In Theorem 4, A is the projection onto \mathfrak{M}^\perp . This was proved in [1] for a paranormal contraction.

COROLLARY 6. *Let $T \in B(H)$ be a dominant or paranormal contraction. If there exists a vector $x_0 \in H$ such that $\|T^{*n}x_0\| \geq \epsilon > 0$ for $n = 1, 2, 3, \dots$, then T has a non-trivial invariant subspace.*

PROOF. Let $\mathfrak{M} = \{x \in H \mid \|T^{*n}x\| \rightarrow 0 \ (n \rightarrow \infty)\}$. By hypothesis, $\mathfrak{M} \neq H$ or $\mathfrak{M}^\perp \neq \{0\}$. By Theorem 4, $T = T_1 \oplus U$, where $U = T|_{\mathfrak{M}^\perp}$ is unitary, and thus T has a non-trivial invariant subspace.

REFERENCES

[1] K. OKUBO, The unitary part of paranormal operators, *Hokkaido Math. J.* 6 (1977), 273-275.
 [2] T. SAITÔ, On a theorem by S. M. Patel, *Rev. Roumaine Math. pures et appl.* 21 (1976), 1407-1409.
 [3] J. G. STAMPFLI AND B. L. WADHWA, An asymmetric Putnam-Fuglede theorem for dominant operators, *Indiana Univ. Math. J.* 25 (1976), 359-365.
 [4] J. G. STAMPFLI AND B. L. WADHWA, On dominant operators, *Monatshefte für Math.* 84 (1977), 143-153.
 [5] T. YOSHINO, On the unitary part of paranormal contractions, preprint.

FACULTY OF EDUCATION	AND	COLLEGE OF GENERAL EDUCATION
RYUKYU UNIVERSITY		TÔHOKU UNIVERSITY
NAHA, 903 JAPAN		SENDAI, 980 JAPAN

