

FINITE TIME DEAD-CORE RATE FOR THE HEAT EQUATION WITH A STRONG ABSORPTION

JONG-SHENQ GUO AND CHIN-CHIN WU

(Received October 10, 2006, revised May 1, 2007)

Abstract. We study the solution of the heat equation with a strong absorption. It is well-known that the solution develops a dead-core in finite time for a large class of initial data. It is also known that the exact dead-core rate is faster than the corresponding self-similar rate. By using the idea of matching, we formally derive the exact dead-core rates under a dynamical theory assumption. Moreover, we also construct some special solutions for the corresponding Cauchy problem satisfying this dynamical theory assumption. These solutions provide some examples with certain given polynomial rates.

1. Introduction. In this paper, we study the following initial boundary value problem (P) for the heat equation with a strong absorption:

$$(1.1) \quad u_t = u_{xx} - u^p, \quad 0 < x < 1, \quad t > 0,$$

$$(1.2) \quad u_x(0, t) = 0, \quad u(1, t) = k, \quad t > 0,$$

$$(1.3) \quad u(x, 0) = u_0(x), \quad 0 \leq x \leq 1,$$

where $p \in (0, 1)$, k is a positive constant, and u_0 is a smooth positive function defined on $[0, 1]$ such that

$$(1.4) \quad u'_0(0) = 0, \quad u_0(1) = k, \quad u'_0(x) \geq 0 \quad \text{for } x \in [0, 1].$$

Problem (P) has been studied extensively for past years. It arises in the modeling of an isothermal reaction-diffusion process [2, 11] in which the solution u of (P) represents the concentration of the reactant which is injected with a fixed amount on the boundary $x = \pm 1$ (by a symmetric reflection), and p is the order of reaction. It also arises in the modeling of a description of thermal energy transport in plasma [8, 7]. For more references, we refer the reader to a recent work of Guo-Souplet [4] and the references cited therein.

In the literature, the region where $u = 0$ is called the *dead-core* and the first time when u reaches zero is called the *dead-core time*.

2000 *Mathematics Subject Classification.* Primary 35K57; Secondary 35K55, 34B16.

Key words and phrases. dead-core, dead-core rate, heat equation, absorption.

This work is partially supported by the National Science Council of the Republic of China under the grant NSC 94-2115-M-003-005. We would like to thank Professor Noriko Mizoguchi for some valuable discussions and the referee for some helpful comments.

It is shown in [2] that for a large class of initial data u_0 the solution of (P) develops a dead-core in a finite time, say, T . By the following transformation

$$(1.5) \quad u(x, t) = (T - t)^\alpha w(y, s), \quad \alpha := 1/(1 - p),$$

$$(1.6) \quad y = x/\sqrt{T - t}, \quad s = -\ln(T - t),$$

we see that u is a solution of (P) if and only if w is a solution of the following initial and boundary value problem (Q):

$$(1.7) \quad w_s = w_{yy} - w^p + \alpha w - \frac{1}{2}yw_y, \quad 0 < y < R(s), \quad s > s_0,$$

$$(1.8) \quad w_y(0, s) = 0, \quad w(R(s), s) = ke^{\alpha s}, \quad s > s_0,$$

$$(1.9) \quad w(y, s_0) = w_0(y) := u_0(y\sqrt{T})/T^\alpha, \quad 0 \leq y \leq 1/\sqrt{T},$$

where $R(s) := e^{s/2}$ and $s_0 := -\ln T$. Furthermore, it is shown in [4] that, as $s \rightarrow \infty$, $w(y, s) \rightarrow U(y)$ uniformly on compact subsets. Here $U(y) := k_p y^{2\alpha}$, $k_p := [2\alpha(2\alpha - 1)]^{-\alpha}$. In particular, $w(0, s) \rightarrow 0$ as $s \rightarrow \infty$. Therefore, the exact convergence rate is still not determined.

The main purpose of this paper is to find the exact convergence rate of $w(0, s)$. For the same question to different problems, we refer to the reader to the recent works of Dold-Galaktionov-Lacey-Vazquez [3] and Souplet-Vazquez [10]. The main idea of these two works is to use a matching of the inner and outer expansions.

For the inner expansion, we enlarge the inner region near $y = 0$ by a re-scaling. Then the inner expansion is derived by studying a stabilization problem as the time goes to infinity.

For the outer expansion, we first study the linearized operator of the right-hand side of (1.7) around the singular steady state U . Then, from the dynamical point of view, we assume that there exist an integer $l \geq 1$ and positive constants ε, K, s_1 with ε sufficiently small such that

$$(1.10) \quad |w(y, s) - U(y) - \hat{c}_l e^{-(l-1/2)s} \phi_l(y)| \leq \varepsilon e^{-(l-1/2)s} y^{2\alpha-1}$$

for $y \in [Ke^{-(l-1/2)s}, 1]$ and $s \geq s_1$ for some nonzero constant \hat{c}_l , where ϕ_l is the l -th eigenfunction of the linearized operator.

Then, by a matching, the rate of convergence of $w(0, s)$ can be formally derived as

$$(1.11) \quad \{\ln[w(0, s)]/s\} = -2\alpha(l - 1/2) + O(1/s) \quad \text{as } s \rightarrow \infty.$$

Note that the estimate (1.11) implies that

$$u(0, t) \sim (T - t)^{\alpha+2\alpha(l-1/2)} \quad \text{as } t \uparrow T^-.$$

Hence, under the assumption (1.10), the dead-core rate is at most polynomially. But, it is faster than the so-called self-similar rate.

Although we are unable to verify the assumption (1.10) for general solutions, motivated by the works of Herrero-Velazquez [6] and Mizoguchi [9], we can construct some special solutions for the corresponding Cauchy problem such that the assumption (1.10) is satisfied by these solutions for any odd integer l .

We note that the dead-core rate of the solution of (P) should depend on the initial data u_0 . We observe from the exact expression of ϕ_l (see Section 4 below) that there are exact l intersections of $w(y, s)$ (constructed in Theorem 6.1 below) and $U(y)$ in $(0, \infty)$ for any $s \geq s_1$. Notice that the number of intersections of $w(y, s)$ and $U(y)$ is the same as the number of intersections of $u(x, t)$ and $U(x)$ due to the scaling invariance of U under the scaling (1.6). Also, as $t \uparrow T^-$ (or $s \rightarrow \infty$), the y -domain of w tends to the whole real line. It is nature to expect that (1.10) is satisfied with an integer l which is related to the number of intersections of $u(x, t)$ and $U(x)$ in $(0, 1)$. This gives a connection between the initial data and the assumption (1.10) on the solution.

The paper is organized as follows. In Section 2, we study the structure of steady states of (P). The inner expansion is given in Section 3. In Section 4, we first study the spectrum of the linearized operator around the singular steady state U . With this information on the spectrum, we then give a formal outer expansion. Then, in Section 5, we formally derive the exact convergence rate of $w(0, s)$ under the dynamical theory assumption (1.10). Finally, to illustrate the plausibility of the assumption (1.10), we construct some special solutions for the corresponding Cauchy problem with certain given rates in Section 6. These solutions satisfy the dynamical theory assumption (1.10). The proof of a key lemma in this construction is given in Section 7. This involves a quite heavy analysis.

2. Steady states. In this section, we shall study the steady states of (P). For any $\eta \geq 0$, let U_η be the solution of

$$(2.1) \quad u'' = u^p, \quad u > 0 \quad \text{for any } y > 0; \quad u(0) = \eta, \quad u'(0) = 0.$$

In particular, $U_0(y) = U(y) = k_p y^{2\alpha}$ for $y \geq 0$. Note that, by a re-scaling, we have

$$(2.2) \quad U_\eta(y) = \eta U_1(\eta^{(p-1)/2} y) \quad \text{for any } \eta > 0.$$

Also, by a simple comparison, we have $U_{\eta_1} > U_{\eta_2}$ if $\eta_1 > \eta_2 \geq 0$. Moreover, $U_\eta \rightarrow U_0$ as $\eta \rightarrow 0^+$.

REMARK 1. For $\eta = 0$, there are non-negative solutions in the form

$$U_0^\varepsilon(y) := k_p (y - \varepsilon)_+^{2\alpha}$$

for any $\varepsilon > 0$. Also, these give all the possible non-negative non-trivial solutions of (2.1).

Concerning the asymptotic behavior of U_η as $\eta \rightarrow 0^+$, we have

LEMMA 2.1. As $\eta \rightarrow 0^+$,

$$(2.3) \quad U_\eta(x) = U_0(x) + k(\eta)x^{2\alpha-1}(1 + o(1))$$

for any $x > 0$, where $k(\eta) := a\eta^{(1-p)/2}$ for some $a > 0$.

PROOF. First, we study the asymptotic behavior of $U_1(y)$ as $y \rightarrow \infty$. For this, we write $U_1 = U_0 + v$. Then v satisfies the equation

$$v'' = by^{-2}v + c_2y^{-2-2\alpha}v^2 + c_3y^{-2-4\alpha}v^3 + \dots$$

for some constants c_i , $i \geq 2$, where $b := (2\alpha - 1)(2\alpha - 2)$. Assume that $v(y) \sim y^\gamma$ as $y \rightarrow \infty$ for some $\gamma > 0$. Then we must have

$$\gamma(\gamma - 1) = b.$$

By writing $\gamma = 2\alpha - \delta$, we obtain that either $\delta = 1$ or $\delta = 4\alpha - 2 > 2\alpha$, which is impossible. Hence we obtain that

$$(2.4) \quad U_1(y) = U_0(y) + ay^{2\alpha-1}(1 + o(1)) \quad \text{as } y \rightarrow \infty$$

for some constant a . The constant a is positive, since $U_1 > U_0$.

Now, for any $x > 0$, from (2.2) and (2.4) it follows that

$$U_\eta(x) = \eta U_1(\eta^{(p-1)/2}x) = U_0(x) + a\eta^{(1-p)/2}x^{2\alpha-1}(1 + o(1)) \quad \text{as } \eta \rightarrow 0^+.$$

The lemma is proved. \square

3. Inner expansion. In the sequel, for convenience we denote $\sigma(s) := w(0, s)$. To derive the inner expansion, we make the following transformations

$$(3.1) \quad w(y, s) := \sigma(s)\theta(\xi, \tau), \quad \xi := \sigma(s)^{(p-1)/2}y, \quad \tau = \tau(s) := \int_{s_0}^s \sigma(r)^{p-1}dr.$$

Then it is easy to check that θ satisfies the equation

$$(3.2) \quad \theta_\tau = \theta_{\xi\xi} - \theta^p + [\sigma^{1-p}(s) + g(\tau)]\left(\alpha\theta - \frac{1}{2}\xi\theta_\xi\right),$$

where

$$(3.3) \quad g(\tau) := -(1-p)\sigma'(s)\sigma(s)^{-p}.$$

Also, $\theta(0, \tau) = 1$ and $\theta_\xi(0, \tau) = 0$ for all $\tau > 0$. We shall study the stabilization of the solution θ of (3.2).

First, recall from (3.7) of [4] that

$$0 \leq u_x \leq Cu^{(p+1)/2} \quad \text{for any } x \in [0, 1], \quad T/2 \leq t \leq T,$$

where C is a positive constant. Hence we have

$$(3.4) \quad 0 \leq w_y \leq Cw^{(p+1)/2} \quad \text{for any } y \in [0, R(s)], \quad s \geq -\ln(T/2) := s_1,$$

for some positive constant C . Consequently, by an integration, we deduce from (3.4) that

$$(3.5) \quad w(y, s) \leq [\sigma(s)^{(1-p)/2} + cy]^{2\alpha} \quad \text{for any } y \in [0, R(s)], \quad s \geq s_1,$$

for some positive constant c . Using (3.5), (3.1) and $w_y = \sigma^{(1+p)/2}\theta_\xi$, we obtain the following estimate

$$(3.6) \quad 0 < \theta(\xi, \tau) \leq (1 + c\xi)^{2\alpha}, \quad 0 \leq \xi\theta_\xi(\xi, \tau) \leq C(1 + \xi)^{2\alpha}$$

for all $\xi \in [0, R(s)\sigma^{(p-1)/2}(s)]$, $\tau \geq \tau_1 := \tau(s_1)$, for some positive constants c and C .

On the other hand, recall from (1.7) of [4] that

$$(3.7) \quad u(x, t) \geq [u(0, t)^{1-p} + cx^2]^\alpha \quad \text{for any } x \in [0, 1], \quad T/2 \leq t \leq T,$$

for some positive constant c . Hence $w(y, s)$ grows at least as fast as $y^{2\alpha}$ for $y \gg 1$ and $s \gg 1$.

Next, it follows from Hopf's Lemma that $w_{yy}(0, s) > 0$ and so $w_s(0, s) > -w^p(0, s)$ by (1.7). Hence $g(\tau) < 1 - p$ for all $\tau > 0$. In the sequel, we assume, in addition to (1.4), that u_0 satisfies the condition

$$(3.8) \quad u_0'' - u_0^p \leq 0 \quad \text{in } [0, 1].$$

It follows from the maximum principle that $u_t \leq 0$ in $[0, 1] \times [0, T]$. From the relation

$$u_t(x, t) = (T - t)^{\alpha-1} [w_s(y, s) - \alpha w(y, s) + y w_y(y, s)/2],$$

it follows that $w_s(0, s) \leq \alpha w(0, s)$ for all $s > s_0$. Hence $g(\tau) \geq -w^{1-p}(0, s)$ for all $s > s_0$. Therefore, g is bounded and $\liminf_{\tau \rightarrow \infty} g(\tau) \geq 0$.

Note that

$$\int_0^\infty g(\tau) d\tau = -(1-p) \int_{s_0}^\infty \sigma^{-1}(s) \sigma'(s) ds = \infty.$$

Hence either $\limsup_{\tau \rightarrow \infty} g(\tau) = 0$ or $\limsup_{\tau \rightarrow \infty} g(\tau) > 0$. Indeed, the first case holds as shown in the following useful lemma.

LEMMA 3.1. *There holds $\lim_{\tau \rightarrow \infty} g(\tau) = 0$.*

PROOF. Otherwise, there is a sequence $\{\tau_n\} \rightarrow \infty$ such that $g(\tau_n) \rightarrow \gamma$ as $n \rightarrow \infty$ for some constant $\gamma > 0$. By using (3.6) and the standard regularity theory, we can show that there is a subsequence, still denoted by $\{\tau_n\}$, such that

$$\theta(\xi, \tau + \tau_n) \rightarrow \tilde{\theta}(\xi, \tau) \quad \text{as } n \rightarrow \infty$$

uniformly on any compact subsets, where $\tilde{\theta}$ solves the equation

$$\tilde{\theta}_\tau = \tilde{\theta}_{\xi\xi} - \tilde{\theta}^p + \gamma \left(\alpha \tilde{\theta} - \frac{1}{2} \xi \tilde{\theta}_\xi \right), \quad \xi > 0, \quad \tau > 0,$$

with $\tilde{\theta}(0, \tau) = 1$ and $\tilde{\theta}_\xi(0, \tau) = 0$. Moreover, it is easy to check that $\tilde{\theta}_\xi \geq 0$ and $\tilde{\theta}(\xi, \tau)$ grows at most polynomially as $\xi \rightarrow \infty$ for $\tau \gg 1$.

Now, it follows from the so-called energy argument (cf. the proof of Proposition 3.1 in [4]) that $\tilde{\theta}(\xi, \tau) \rightarrow V(\xi)$ as $\tau \rightarrow \infty$ for some V satisfying

$$\begin{aligned} V'' - V^p + \gamma \left(\alpha V - \frac{1}{2} \xi V' \right) &= 0, \quad \xi > 0, \\ V'(0) &= 0, \quad V(0) = 1. \end{aligned}$$

Note that $V' \geq 0$ and V grows at most polynomially. Set

$$W(y) := \gamma^\alpha V(y/\sqrt{\gamma}).$$

Then W satisfies

$$\begin{aligned} W'' - W^p + \alpha W - \frac{1}{2} y W' &= 0, \quad y > 0, \\ W'(0) &= 0, \quad W(0) = \gamma^\alpha. \end{aligned}$$

Since $W > 0$, $W' \geq 0$ for $y > 0$, and taking into account of the polynomial boundedness of W , it follows from Proposition 3.3 of [4] that either $W = U$ or $W \equiv \alpha^{-\alpha}$. The first case is impossible, since $U(0) = 0$. The second case is also impossible, since θ is unbounded by (3.7). Hence the lemma follows. \square

Again, by the standard limiting process with the estimate (3.6) and Lemma 3.1, for any given sequence $\{\tau_n\} \rightarrow \infty$ we can show that there is a limit $\tilde{\theta}$ satisfying

$$\begin{aligned}\tilde{\theta}_\tau &= \tilde{\theta}_{\xi\xi} - \tilde{\theta}^p, \quad \xi > 0, \quad \tau \in \mathbf{R}, \\ \tilde{\theta}(0, \tau) &= 1, \quad \tilde{\theta}_\xi(0, \tau) = 0,\end{aligned}$$

such that $\theta(\xi, \tau + \tau_n) \rightarrow \tilde{\theta}(\xi, \tau)$ as $n \rightarrow \infty$ uniformly on compact subsets of $(0, \infty) \times \mathbf{R}$. Since we also have

$$0 = U_1'' - U_1^p, \quad \tilde{\theta}(0, \tau) = U_1(0), \quad \tilde{\theta}_\xi(0, \tau) = (U_1)_\xi(0),$$

Hopf's Lemma and the real analyticity of $\tilde{\theta} - U_1$ imply that $\tilde{\theta} \equiv U_1$. Indeed, suppose on the contrary that $\tilde{\theta} \not\equiv U_1$. Taking any finite τ_0 , it follows from the real analyticity of $v(\xi) := \tilde{\theta}(\xi, \tau_0) - U_1(\xi)$ in ξ that the zeros of v are isolated. Assume that the smallest positive zero of v is ξ_0 ($\xi_0 := \infty$, if there is no finite zero). Without loss of generality we may assume that $v < 0$ in $(0, \xi_0)$. In the connected component

$$Q := \{(\xi, \tau) \mid \xi > 0, \tau \geq \tau_0, \tilde{\theta} - U_1 < 0\}$$

containing $(0, \xi_0) \times \{\tau_0\}$, any point $(0, \tau)$ with $\tau > \tau_0$ is a maximum point. Then Hopf's Lemma implies that $(\tilde{\theta} - U_1)_\xi(0, \tau) < 0$, a contradiction. Therefore, we must have $\tilde{\theta} \equiv U_1$.

Since this limit is independent of the given sequence $\{\tau_n\}$, we see that $\theta(\xi, \tau) \rightarrow U_1(\xi)$ as $\tau \rightarrow \infty$ uniformly on any compact subsets. Returning to the original variables and using the relation (2.2), we thus have proved the following so-called inner expansion.

THEOREM 3.2. *As $s \rightarrow \infty$, we have*

$$w(y, s) = U_{\sigma(s)}(y)(1 + o(1))$$

uniformly in $\{0 \leq \sigma^{(p-1)/2}(s)y \leq C\}$ for any positive constant C .

4. Outer expansion. In the matching process, we need to study the following linearized operator

$$\mathcal{L}v := -v'' + \frac{y}{2}v' + \frac{b}{y^2}v, \quad b := (2\alpha - 1)(2\alpha - 2)$$

which comes from the linearization of (1.7) around the singular steady state U .

Consider the eigenvalue problem

$$(4.1) \quad \mathcal{L}\phi = \lambda\phi, \quad y > 0; \quad \phi'(0) = 0, \quad \phi(0) = 0.$$

Set $\rho(y) := \exp(-y^2/4)$. We introduce the following weighted Hilbert spaces:

$$L_\rho^2 := \left\{ \phi \text{ is measurable in } \mathbf{R} \mid \phi(y) = \phi(|y|), \int_0^\infty \phi^2(y)\rho(y)dy < \infty \right\},$$

$$\mathbf{H} := \left\{ \phi \in L_\rho^2 \mid \int_0^\infty \phi_y^2(y)\rho(y)dy < \infty, \int_0^\infty \frac{\phi^2(y)}{y^2}\rho(y)dy < \infty \right\}.$$

Also, we set

$$J(\phi) := \int_0^\infty \phi_y^2(y)\rho(y)dy + b \int_0^\infty \frac{\phi^2(y)}{y^2}\rho(y)dy,$$

$$I(\phi) := \int_0^\infty \phi^2(y)\rho(y)dy.$$

Then the principal eigenvalue λ_0 of (4.1) can be characterized by

$$(4.2) \quad \lambda_0 := \inf\{J(\phi)/I(\phi) \mid \phi \in \mathbf{H}, I(\phi) > 0\}.$$

It is easy to see that $\lambda_0 > 0$. Also, by taking a minimization sequence, we can show that this λ_0 can be attained by a function $\phi_0 \in \mathbf{H}$ which is the eigenfunction of (4.1) such that

$$\phi_0 > 0 \quad \text{in } (0, \infty), \quad \int_0^\infty (\phi_0(y))^2 \rho(y) dy = 1.$$

Moreover, from the standard theory on eigenvalue problem, there is a sequence of eigenpairs $\{(\lambda_n, \phi_n)\}_{n \geq 0}$ of (4.1) with $0 < \lambda_0 < \lambda_1 \leq \lambda_2 \leq \dots$ and $\lambda_n \rightarrow \infty$ as $n \rightarrow \infty$. Since $\phi_n \not\equiv 0$, without loss of generality we may assume that $\phi_n > 0$ for $y > 0$ and small. Also, we take ϕ_n so that $\int_0^\infty (\phi_n(y))^2 \rho(y) dy = 1$. It is also easy to see that

$$(4.3) \quad \phi_n(y) = d_n y^{\gamma_+} (1 + o(1)) \quad \text{as } y \rightarrow 0^+$$

for some positive constant d_n , where $\gamma_+ := 2\alpha - 1$.

To compute the eigenvalues, we set

$$\phi(y) = |y|^{\gamma_+} H(\eta), \quad \eta = y^2/4.$$

Then ϕ satisfies (4.1) if and only if H satisfies

$$(4.4) \quad \eta H''(\eta) + (\hat{b} - \eta)H'(\eta) - \hat{a}H(\eta) = 0,$$

where $\hat{a} := \gamma_+/2 - \lambda$ and $\hat{b} := \gamma_+ + 1/2$. This is Kummer's Equation (cf. [1]) and its general solution is given by

$$c_1 M(\hat{a}, \hat{b}, \eta) + c_2 U(\hat{a}, \hat{b}, \eta)$$

for some constants c_1 and c_2 , where

$$U(\hat{a}, \hat{b}, \eta) := \frac{\pi}{\sin \hat{b}\pi} \left[\frac{M(\hat{a}, \hat{b}, \eta)}{\Gamma(1 + \hat{a} - \hat{b})\Gamma(\hat{b})} - \eta^{1-\hat{b}} \frac{M(1 + \hat{a} - \hat{b}, 2 - \hat{b}, \eta)}{\Gamma(\hat{a})\Gamma(2 - \hat{b})} \right]$$

with M Kummer's function and Γ the Gamma function.

Since $H(0)$ is finite, $1 - \hat{b} = 3/2 - 2\alpha < 0$, and $M(\cdot, \cdot, 0) = 1$, it follows that the solution of (4.4) is given by

$$H(\eta) = cM(\hat{a}, \hat{b}, \eta)$$

for some constant c .

Since $\hat{b} > 0$, $H(\eta)$ is always well-defined. If $-\hat{a} \in \mathcal{N} \cup \{0\}$, then $M(\hat{a}, \hat{b}, \eta)$ is a polynomial of degree $-\hat{a}$ in η (cf. [1]). Indeed, $M \equiv 1$ if $\hat{a} = 0$. For $n \in \mathcal{N}$, we have

$$M(-n, \hat{b}, \eta) = 1 + \sum_{m=1}^n (-1)^m \frac{n!}{(n-m)!m!} \frac{1}{\hat{b} \cdots (\hat{b} + m - 1)} \eta^m.$$

Otherwise, if $-\hat{a} \notin \mathcal{N} \cup \{0\}$, then

$$M\left(\hat{a}, \hat{b}, \frac{y^2}{4}\right) \sim \frac{\Gamma(\hat{b})}{\Gamma(\hat{a})} \left(\frac{y^2}{4}\right)^{-q} \exp\left(\frac{y^2}{4}\right)$$

as $y \rightarrow \infty$, where $q := \gamma_+/2 + 1/2 + \lambda = \alpha + \lambda$. Since $\phi \in \mathbf{H}$, we conclude that the eigenvalues of \mathcal{L} are given by $\lambda_0 = \alpha - 1/2$ and $\lambda_n = \gamma_+/2 + n = \alpha + n - 1/2$ for $n \in \mathcal{N}$.

Note that $\phi_0(y) = c_0|y|^{2\alpha-1}$ for some positive constant c_0 . Since $\phi(y) = y^{\gamma_+} H(\eta)$, we have

$$(4.5) \quad \phi_n(y) = c_n y^{2\alpha-1} (1 + o(1)) \quad \text{as } y \rightarrow 0^+,$$

$$(4.6) \quad \phi_n(y) = \tilde{c}_n y^{2\lambda_n} (1 + o(1)) \quad \text{as } y \rightarrow \infty,$$

for some constants $c_n > 0$ and $(-1)^n \tilde{c}_n > 0$ for all $n \in \mathcal{N}$.

In this paper, we are unable to derive a good outer expansion rigorously. Since $w \rightarrow U$ as $s \rightarrow \infty$, from the dynamical theory point of view, we assume instead that there exist an integer $l \geq 1$ and positive constants ε, K, s_1 with ε sufficiently small such that

$$(4.7) \quad |w(y, s) - U(y) - \hat{c}_l e^{-(\lambda_l - \alpha)s} \phi_l(y)| \leq \varepsilon e^{-(\lambda_l - \alpha)s} y^{2\alpha-1}$$

for $y \in [K e^{-(\lambda_l - \alpha)s}, 1]$ and $s \geq s_1$ for some nonzero constant \hat{c}_l .

5. Rate of convergence. In this section, we shall use the idea of matching to derive formally the exact convergence rate of $\sigma(s) := w(0, s)$ to zero as $s \rightarrow \infty$.

We recall from Lemma 2.1 and Theorem 3.2 that for any $y \in (0, C\sigma^{(1-p)/2}(s)]$, $C > 0$:

$$(5.1) \quad w(y, s) - U(y) = a\sigma^{(1-p)/2}(s) y^{2\alpha-1} (1 + o(1)) \quad \text{as } s \rightarrow \infty.$$

On the other hand, by (4.7) and (4.3), we have, for $y(s) := K e^{-(\lambda_l - \alpha)s}$,

$$(5.2) \quad \begin{aligned} (\hat{c}_l d_l - \varepsilon) e^{-(l-1/2)s} y(s)^{2\alpha-1} &\leq w(y(s), s) - U(y(s)) \\ &\leq (\hat{c}_l d_l + \varepsilon) e^{-(l-1/2)s} y(s)^{2\alpha-1} \end{aligned}$$

for all $s \geq s_1$ for some $l \geq 1$. Consequently, (1.11) can be formally derived under the dynamical theory assumption (4.7).

6. Construction of some special solutions. In this section, we shall construct some special solutions for the problem (CP):

$$(6.1) \quad w_s = w_{yy} - \frac{y}{2}w_y + \alpha w - w^p \quad \text{for } y > 0, \quad s > s_1,$$

$$(6.2) \quad w_y(0, s) = 0 \quad \text{for } s > s_1,$$

$$(6.3) \quad w(y, s_1) = w_0(y) > 0 \quad \text{for } y \in [0, \infty).$$

By setting $v := w - U$, (6.1) is equivalent to

$$(6.4) \quad v_s = -\mathcal{A}v + f(v),$$

where

$$\begin{aligned} \mathcal{A}v &:= -v_{yy} + \frac{y}{2}v_y + \frac{b}{y^2}v - \alpha v, \quad b := (2\alpha - 1)(2\alpha - 2), \\ f(v) &:= U^p - (v + U)^p + bv/y^2. \end{aligned}$$

First, we fix some notation. For each $l \in \mathbb{N}$, let c_l be the constant in (4.5) with $n = l$. Let $k(\eta)$ be the constant in (2.3). We choose (for a fix l) two positive constants η_1 and η_2 so that $k(\eta_1) < c_l < k(\eta_2)$.

THEOREM 6.1. *For each odd $l \in \mathbb{N}$, there exists a positive solution w of (CP) such that, for some positive constants $s_1, K, \sigma, \varepsilon$ with $(\lambda_l - \alpha)/(2\lambda_l - 2\alpha + 1) < \sigma < 1/2, s_1 \gg 1, K \gg 1$, and $\varepsilon \ll 1$,*

$$(6.5) \quad e^{-\gamma s} U_{\eta_1}(e^{\beta s} y) < w(y, s) < e^{-\gamma s} U_{\eta_2}(e^{\beta s} y)$$

for $y \in [0, K e^{-\beta s}]$ and $s \geq s_1$, and

$$(6.6) \quad |w(y, s) - U(y) - e^{-\beta s} \phi_l(y)| \leq \varepsilon e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l})$$

for $y \in [K e^{-\beta s}, e^{\sigma s}]$ and $s \geq s_1$, where $\beta = \beta_l := \lambda_l - \alpha$ and $\gamma = \gamma_l := 2\alpha\beta$.

Hereafter l is a fixed odd positive integer so that $\beta = \lambda_l - \alpha$ and $\gamma = 2\alpha\beta$ are fixed.

For a given $d := (d_0, \dots, d_{l-1}) \in \mathbf{R}^l$ with the property

$$(D) \quad \sum_{n=0}^{l-1} |d_n| < \varepsilon e^{-\beta s_1},$$

we let $w(y, s; d)$ be the solution of (6.1)–(6.3) with the initial data $w_0(y) = v(y, s_1) + U(y)$, where

$$(6.7) \quad v(y, s_1) = v(y, s_1; d) := \sum_{n=0}^{l-1} d_n \phi_n(y) + e^{-\beta s_1} \tilde{\phi}_l(y)$$

satisfying, for some (fixed) constants $\tilde{\sigma}$ and \tilde{K} with $\tilde{\sigma} \in ((\lambda_l - \alpha)/(2\lambda_l - 2\alpha + 1), 1/2)$ and $\tilde{K} \gg 1$,

$$(V1) \quad \tilde{\phi}_l(y) := -e^{\beta s_1} \left\{ U(y) - e^{-\gamma s_1} U_*(e^{\beta s_1} y) + \sum_{n=0}^{l-1} d_n \phi_n(y) \right\}, \quad y \in [0, \tilde{K} e^{-\beta s_1}],$$

where U_* is the solution of (2.1) so that $k(\eta) = c_l$ in (2.3);

- (V2) $\tilde{\phi}_l(y) := \phi_l(y)$, $y \in [\tilde{K}e^{-\beta s_1}, e^{\tilde{\sigma} s_1}]$;
(V3) $\tilde{\phi}_l(y)$ is smooth for $y \geq \tilde{K}e^{-\beta s_1}$ and satisfies the bound

$$\tilde{\phi}_l(y) \leq \min\{\phi_l(y), U(y)\} \quad \text{for } y \geq e^{\tilde{\sigma} s_1};$$

- (V4) $v(y, s_1) + U(y) > 0$ in $(0, \infty)$.

Fix a positive constant $\varepsilon \ll 1$. Take $\sigma \in ((\lambda_l - \alpha)/(2\lambda_l - 2\alpha + 1), \tilde{\sigma})$, $s_1 \gg 1$, and $K \gg \tilde{K}$ such that $e^{\beta s_1/8} < K < e^{\beta s_1/2}$. For $s_2 \geq s_1 > 0$ and $\theta \in (0, 1]$, let W_{s_1, s_2}^θ be the set of (symmetric) continuous functions w on $[0, \infty)$ satisfying

$$(6.8) \quad \begin{aligned} |w(y, s) - U(y) - e^{-\beta s} \phi_l(y)| &\leq \theta \varepsilon e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}) \\ &\text{if } y \in [Ke^{-\beta s}, e^{\sigma s}], s \in [s_1, s_2]. \end{aligned}$$

Define U_{s_1, s_2} to be the set of all $d \in \mathbf{R}^l$ with the property (D) such that $w(y, s; d) \in W_{s_1, s_2}^1$. Also, we define \bar{U}_{s_1, s_2} to be the closure of U_{s_1, s_2} . Hereafter the constant σ is fixed.

PROPOSITION 6.1. *Let $s_1 \gg 1$ and $K \gg \tilde{K}$. If $d \in \bar{U}_{s_1, s_2}$ with some $s_2 > s_1$, then there exists a positive constant $\delta_0 \in (0, 1)$ such that*

$$(6.9) \quad (1 + \delta_0)e^{-\gamma s} U_{\eta_1}(e^{\beta s} y) < w(y, s) < (1 - \delta_0)e^{-\gamma s} U_{\eta_2}(e^{\beta s} y)$$

for $y \in [0, Ke^{-\beta s}]$ and $s \in [s_1, s_2]$, where $w(y, s) := w(y, s; d)$.

PROOF. First, we recall from the proof of Lemma 2.1 that, as $\eta^{(p-1)/2} e^{\beta s} y \rightarrow \infty$,

$$(6.10) \quad e^{-\gamma s} U_\eta(e^{\beta s} y) = U(y) + k(\eta)e^{-\beta s} y^{2\alpha-1} (1 + o(1)).$$

By assumption, $w(y, s) \in W_{s_1, s_2}^1$, since $d \in \bar{U}_{s_1, s_2}$. By (6.8) and the choices of $\eta_1, \eta_2, \varepsilon$, there exists a positive constant $\delta_1 \in (0, 1)$ such that

$$(6.11) \quad (1 + \delta_1)e^{-\gamma s} U_{\eta_1}(e^{\beta s} y) < w(y, s) < (1 - \delta_1)e^{-\gamma s} U_{\eta_2}(e^{\beta s} y)$$

in $D_1 := \{y = Ke^{-\beta s}, s \in [s_1, s_2]\}$, if $K \gg 1$. From (D), (V1) and (V2), by choosing $\delta_2 \in (0, \delta_1)$ small enough, we have

$$(6.12) \quad (1 + \delta_2)e^{-\gamma s} U_{\eta_1}(e^{\beta s_1} y) < w(y, s_1) < (1 - \delta_2)e^{-\gamma s_1} U_{\eta_2}(e^{\beta s_1} y)$$

in $D_2 := \{y \in [0, Ke^{-\beta s_1}], s = s_1\}$, if $s_1 \gg 1$ and $K \gg \tilde{K}$. Set $\delta_0 := \delta_2$. From (6.11) and (6.12), it follows that (6.9) holds in $D_1 \cup D_2$, if $K \gg \tilde{K}$ and $s_1 \gg 1$.

Now, we define

$$\begin{aligned} \tilde{w}(y, s) &= (1 - \delta_0)e^{-\gamma s} U_{\eta_2}(e^{\beta s} y), \\ \hat{w}(y, s) &= (1 + \delta_0)e^{-\gamma s} U_{\eta_1}(e^{\beta s} y). \end{aligned}$$

Then from a direct calculation we get, for $y \in [0, Ke^{-\beta s}]$ and $s \in [s_1, s_2]$,

$$\begin{aligned} \tilde{w}_s - \tilde{w}_{yy} + \frac{1}{2}y\tilde{w}_y - \alpha\tilde{w} + \tilde{w}^p \\ = (1 - \delta_0)e^{(-\gamma+2\beta)s} \{[-1 + (1 - \delta_0)^{p-1}]U_{\eta_2}^p + e^{-2\beta s}B_2\} > 0 \quad \text{if } s_1 \gg 1, \\ \hat{w}_s - \hat{w}_{yy} + \frac{1}{2}y\hat{w}_y - \alpha\hat{w} + \hat{w}^p \\ = (1 + \delta_0)e^{(-\gamma+2\beta)s} \{[-1 + (1 + \delta_0)^{p-1}]U_{\eta_1}^p + e^{-2\beta s}B_1\} < 0 \quad \text{if } s_1 \gg 1, \end{aligned}$$

where $B_i := -(\gamma + \alpha)U_{\eta_i}(e^{\beta s}y) + (\beta + 1/2)(e^{\beta s}y)U'_{\eta_i}(e^{\beta s}y)$, $i = 1, 2$. Since $w_y(0, s) = 0 = U'_{\eta_i}(0)$, the proposition follows by a comparison principle. \square

We next derive the estimate in the region $\{y \geq e^{\sigma s}\}$ as follows.

PROPOSITION 6.2. *Let $s_1 \gg 1$ and $K \gg \tilde{K}$. If $d \in \bar{U}_{s_1, s_2}$ with some $s_2 > s_1$, then*

$$(6.13) \quad 0 < w(y, s) \leq U(y)$$

for $y \geq e^{\sigma s}$ and $s \in [s_1, s_2]$, where $w(y, s) := w(y, s; d)$.

PROOF. Note that, by (V4), it follows from the maximum principle that $w > 0$. Since $d \in \bar{U}_{s_1, s_2}$, by (6.8) and (4.6), we have

$$w(y, s) \leq U(y) + e^{-\beta s}(\tilde{c}_l - 2\varepsilon)y^{2\lambda_l} < U(y) \quad \text{if } \varepsilon \ll 1 \text{ and } s_1 \gg 1,$$

for $y = e^{\sigma s}$ and $s \in [s_1, s_2]$. Here l is assumed to be odd and so $\tilde{c}_l < 0$. Clearly, (6.13) holds for $y \geq e^{\sigma s}$ and $s = s_1$, if $s_1 \gg 1$, since $v(y, s_1) < 0$ for $y \geq e^{\sigma s_1}$ by using (D), (V2) and (V3). The proposition follows by a comparison principle. \square

Next, we define the operator $P(\cdot; s_1, s_2)$ from \bar{U}_{s_1, s_2} to \mathbf{R}^l by

$$P(d; s_1, s_2) = (p_0, \dots, p_{l-1}), \quad p_n := \langle v(y, s_2; d), \phi_n(y) \rangle, \quad n = 0, \dots, l-1,$$

where $v(y, s; d) := w(y, s; d) - U(y)$ and $\langle f, g \rangle := \int_0^\infty f(y)g(y)\rho(y)dy$.

PROPOSITION 6.3. *Let $s_1 \gg 1$ and $K \gg \tilde{K}$. If there is $d \in \bar{U}_{s_1, s_2}$ for some $s_2 > s_1$ such that $P(d; s_1, s_2) = 0$, then $w(y, s; d) \in W_{s_1, s_2}^\theta$ for some $\theta \in (0, 1)$.*

To prove this key proposition, we shall apply an idea of Herrero and Velazquez [6], which was modified by Mizoguchi [9]. Since it involves rather complicated computations, we shall postpone its proof at the end of this paper. We continue to prove Theorem 6.1.

PROPOSITION 6.4. *Let $s_1 \gg 1$. If $\bar{U}_{s_1, s_2} \neq \emptyset$ for some $s_2 > s_1$, then there exists $d \in \bar{U}_{s_1, s_2}$ such that $P(d; s_1, s_2) = 0$.*

PROOF. We shall apply the degree theory to prove that

$$\deg(P(d; s_1, s_2), 0, U_{s_1, s_2}) = 1,$$

where $\deg(P(d; s_1, s_2), 0, U_{s_1, s_2})$ denotes the degree of $P(\cdot; s_1, s_2)$ with respect to 0 in the set U_{s_1, s_2} . First, we claim that $\deg(P(d; s_1, s_1), 0, U_{s_1, s_1}) = 1$. Note that $p_n(d; s_1, s_1) =$

$d_n + e^{-\beta s_1} \langle \tilde{\phi}_l, \phi_n \rangle$. Set

$$h^t(d) = (1-t)I(d) + tP(d; s_1, s_1),$$

where $I(d)$ is the identity mapping in \mathbf{R}^l and $t \in [0, 1]$. Through the choice of $\tilde{\phi}_l$, for $s_1 \gg 1$ we have $h^t(\partial U_{s_1, s_1}) \neq \emptyset$ for $t \in [0, 1]$. By the standard degree theory, $\deg(h^t(d), 0, U_{s_1, s_1})$ is independent of $t \in [0, 1]$. Hence we obtain that

$$\deg(P(d; s_1, s_1), 0, U_{s_1, s_1}) = \deg(I(d), 0, U_{s_1, s_1}) = 1.$$

Moreover, by Proposition 6.3 and Lemma 7.3, we have $P(d; s_1, s_2) \neq 0$ for all $d \in \partial U_{s_1, s_2}$, if $s_1 \gg 1$. Therefore, it follows from the homotopy invariance of degree theory that $\deg(P(d; s_1, s_2), 0, U_{s_1, s_2}) = 1$. This proves the proposition.

PROPOSITION 6.5. *Let $s_1 \gg 1$. Then $U_{s_1, s_2} \neq \emptyset$ for all $s_2 > s_1$.*

PROOF. Set $s^* := \sup\{s \geq s_1 \mid U_{s_1, s} \neq \emptyset\}$. From the theory of continuous dependence on initial data, we see that $U_{s_1, s} \neq \emptyset$ for $s > s_1$ with $s - s_1 \ll 1$. Hence $s^* > s_1$. Claim that $s^* = \infty$. If not, then there exists a sequence $\{s_j\}$ with $s_j < s^* < \infty$ such that $s_j \rightarrow s^*$ as $j \rightarrow \infty$. By Proposition 6.4, there exists $d_j \in \bar{U}_{s_1, s_j}$ such that $P(d_j; s_1, s_j) = 0$ for each j . Since $\{d_j\}$ is bounded, without loss of generality, we may assume $d_j \rightarrow d^*$ as $j \rightarrow \infty$ for some $d^* \in \mathbf{R}^l$. Moreover, we have $d^* \in \bar{U}_{s_1, s^*}$ and $P(d^*; s_1, s^*) = 0$ from the continuity of P . From Proposition 6.3, there exists $\theta \in (0, 1)$ such that $w(y, s; d^*) \in W_{s_1, s^*}^\theta$. Hence we get $w(y, s; d^*) \in U_{s_1, s^* + \eta}$ for some $\eta > 0$. This is a contradiction to the definition of s^* and the proposition is proved.

Now, we are ready to prove Theorem 6.1.

PROOF OF THEOREM 6.1. From Proposition 6.5 it follows that there exists a strictly increasing sequence $\{s_j\} \subset (s_1, \infty)$ with $s_j \rightarrow \infty$ as $j \rightarrow \infty$ such that $U_{s_1, s_j} \neq \emptyset$. By Proposition 6.4, there exists $d_j \in \bar{U}_{s_1, s_j}$ such that $P(d_j; s_1, s_j) = 0$ for each j . Hence, from Proposition 6.3, we have $w(y, s; d_j) \in W_{s_1, s_j}^1$ for each j . Since $\{d_j\}$ is bounded, without loss of generality we may assume that $d_j \rightarrow \bar{d}$ as $j \rightarrow \infty$ for some $\bar{d} \in \mathbf{R}^l$. Let $w(y, s; \bar{d})$ be the solution of (CP) with initial value $w(y, s_1; \bar{d}) = v(y, s_1; \bar{d}) + U(y)$. Since $d_k \in \bar{U}_{s_1, s_k} \subseteq \bar{U}_{s_1, s_j}$ for all $k > j$ for any given j , we have $\bar{d} \in \bar{U}_{s_1, s_j}$ for all j . Therefore, we conclude that $w(y, s; \bar{d}) \in W_{s_1, \infty}^1$. \square

7. Proof of Proposition 6.3. In the sequel, C denotes a (universal) positive constant, which may be different from one line to another, depending on p and l .

Recall that $f(v) := U^p - (v + U)^p + bv/y^2$. Note that, by (V4), $w = v + U > 0$ and so f is well-defined. From

$$v_s = -\mathcal{A}v + f(v) \quad \text{in } \mathbf{R} \times (s_1, \infty),$$

it follows that (cf. [5])

$$(7.1) \quad v(y, s) = e^{-\mathcal{A}(s-s_1)}v(y, s_1) + \int_{s_1}^s e^{-\mathcal{A}(s-\tau)}f(v(y, \tau))d\tau$$

for $s \geq s_1$. Hereafter s_1 is a very large constant.

LEMMA 7.1. *We have*

$$(7.2) \quad 0 \leq f(v) \leq CU^{p-2}v^2$$

for some $C > 0$.

PROOF. First, since

$$f(v) = \frac{1}{2}p(1-p)(U + \hat{v})^{p-2}v^2$$

for some \hat{v} between 0 and v , we see that $f(v) \geq 0$.

To prove the upper bound, we divide our proof into two cases.

Case 1: $-U/2 \leq v$. Applying the mean value theorem, we have

$$\begin{aligned} f(v) &= U^p - (v+U)^p + pU^{p-1}v \\ &= -p(U + \theta_1 v)^{p-1}v + pU^{p-1}v = -p[(U + \theta_1 v)^{p-1} - U^{p-1}]v \\ &= \theta_1 p(1-p)(U + \theta_1 \theta_2 v)^{p-2}v^2 \\ &\leq \theta_1 p(1-p) \left(U - \frac{1}{2}\theta_1 \theta_2 U \right)^{p-2} v^2 \\ &\leq \theta_1 p(1-p) \left(\frac{1}{2}U \right)^{p-2} v^2 \leq CU^{p-2}v^2, \end{aligned}$$

where the constants $\theta_i \in (0, 1)$, $i = 1, 2$.

Case 2: $-U \leq v \leq -U/2$. Since $-U \leq v \leq -U/2$, we have $-2 \leq U/v \leq -1$. Hence

$$\begin{aligned} f(v) &= U^p - (v+U)^p + pU^{p-1}v \\ &\leq U^p - (v+U)^p \\ &= U^{p-2}v^2 \left[\left(\frac{U}{v} \right)^2 - \left(\frac{v}{U} + 1 \right)^p \left(\frac{U}{v} \right)^2 \right] \\ &\leq 2U^{p-2}v^2 \left(\frac{U}{v} \right)^2 \leq CU^{p-2}v^2. \end{aligned}$$

Combining Case 1 with Case 2, we have proved the lemma.

LEMMA 7.2. *Let $\delta > 1$. If $d \in \bar{U}_{s_1, s_2}$ for some $s_2 > s_1$, then*

$$(7.3) \quad 0 \leq f(v) \leq CK^{2\alpha}e^{-2\alpha\beta s}y^{-2} \quad \text{for } y \in [0, Ke^{-\beta s}],$$

$$(7.4) \quad 0 \leq f(v) \leq Ce^{-2\beta s}y^{2\alpha-4} \quad \text{for } y \in [Ke^{-\beta s}, \delta],$$

$$(7.5) \quad 0 \leq f(v) \leq Ce^{-2\beta s}y^{4\lambda_1-2\alpha-2} \quad \text{for } y \in [\delta, e^{\sigma s}],$$

$$(7.6) \quad 0 \leq f(v) \leq Cy^{2\alpha-2} \quad \text{for } y \geq e^{\sigma s}$$

for each $s \in [s_1, s_2]$, where $v := v(y, s; d) = w(y, s; d) - U(y)$.

PROOF. Given any $s \in [s_1, s_2]$. From (6.9), there exists a constant $C > 0$ such that

$$|v(y, s)| \leq CK^{2\alpha} e^{-2\alpha\beta s}$$

for $y \in [0, Ke^{-\beta s}]$. Then (7.3) follows from the definition of f . From (6.8), (4.5) and (4.6), we have

$$\begin{aligned} |v(y, s)| &\leq Ce^{-\beta s} y^{2\alpha-1} \quad \text{if } y \in [Ke^{-\beta s}, \delta]; \\ |v(y, s)| &\leq Ce^{-\beta s} y^{2\lambda_l} \quad \text{if } y \in [\delta, e^{\sigma s}]. \end{aligned}$$

Moreover, from (6.13) it follows that $|v(y, s)| \leq Cy^{2\alpha}$ for $y \geq e^{\sigma s}$. Therefore, (7.4) through (7.6) follow by using (7.2). \square

LEMMA 7.3. Suppose that $d \in \tilde{U}_{s_1, s_2}$ such that $P(d; s_1, s_2) = 0$. Then for any $v \in (0, 1)$ there exists $s_1 \gg 1$ such that

$$(7.7) \quad \sum_{n=0}^{l-1} |d_n| \leq v e^{-\beta s_1}.$$

PROOF. Since $P(d; s_1, s_2) = 0$, we have $\langle v(y, s_2), \phi_n \rangle = 0$ for $n = 0, \dots, l-1$. Fix $n \in \{0, \dots, l-1\}$. Then we get from (7.1) and (6.7) that $|d_n| \leq I_{n,1} + I_{n,2}$, where

$$\begin{aligned} I_{n,1} &:= e^{-\beta s_1} |\langle \tilde{\phi}_l, \phi_n \rangle|, \\ I_{n,2} &:= e^{-(\lambda_n - \alpha)s_1} \int_{s_1}^{s_2} e^{(\lambda_n - \alpha)\tau} |\langle f, \phi_n \rangle| d\tau. \end{aligned}$$

Since $\tilde{\phi}_l \rightarrow \phi_l$ in L^2_ρ as $s_1 \rightarrow \infty$, $|\langle \tilde{\phi}_l, \phi_n \rangle| \rightarrow 0$ as $s_1 \rightarrow \infty$. Thus for any $v \in (0, 1)$ there exists $s_1 \gg 1$ such that

$$I_{n,1} \leq \frac{v}{2l} e^{-\beta s_1}.$$

Now, we shall apply Lemma 7.2 to estimate $|\langle f, \phi_n \rangle|$ for each $\tau \in [s_1, s_2]$. Recall that $K < e^{\beta s_1/2}$. First, from (7.3) it follows that

$$(7.8) \quad \left| \int_0^{Ke^{-\beta\tau}} f(v(y, \tau)) \phi_n(y) \rho(y) dy \right| \leq CK^{2\alpha} e^{-2\alpha\beta\tau} \leq Ce^{(1-\alpha)\beta\tau} e^{-\beta\tau}.$$

Next, by (7.4) and (7.5), we have

$$(7.9) \quad \left| \left\{ \int_{Ke^{-\beta\tau}}^\delta + \int_\delta^{e^{\sigma\tau}} \right\} f(v(y, \tau)) \phi_n(y) \rho(y) dy \right| \leq Ce^{-2\beta\tau}.$$

Finally, by (7.6), we have

$$(7.10) \quad \left| \int_{e^{\sigma\tau}}^\infty f(v(y, \tau)) \phi_n(y) \rho(y) dy \right| \leq Ce^{-e^{2\sigma\tau}/8} \int_{e^{\sigma\tau}}^\infty y^{2\lambda_l+2\alpha-2} e^{-y^2/8} dy \leq Ce^{-e^{2\sigma\tau}/8}.$$

Combining (7.8), (7.9) and (7.10), we could find a constant $\kappa > 0$ such that

$$(7.11) \quad |\langle f(v(y, \tau)), \phi_n \rangle| \leq Ce^{-(\beta+\kappa)\tau} \quad \text{if } s_1 \gg 1.$$

Hence

$$I_{n,2} \leq \frac{v}{2l} e^{-\beta s_1} \quad \text{if } s_1 \gg 1.$$

This completes the proof of the lemma.

Throughout the following, we suppose that $d \in \bar{U}_{s_1, s_2}$ with $1 \ll s_1 < s_2$ such that $P(d; s_1, s_2) = 0$.

7.1. Short time. $s_1 \leq s \leq s_1 + 1$.

Since $v = \sum_{n=0}^{\infty} \langle v, \phi_n \rangle \phi_n$, we may rewrite $v(y, s_1)$ as

$$v(y, s_1) = \sum_{n=0}^{l-1} (d_n + e^{-\beta s_1} \langle \tilde{\phi}_l, \phi_n \rangle) \phi_n + e^{-\beta s_1} \langle \tilde{\phi}_l, \phi_l \rangle \phi_l + \sum_{n=l+1}^{\infty} e^{-\beta s_1} \langle \tilde{\phi}_l, \phi_n \rangle \phi_n.$$

Thus from (7.1) we may write, for $s \geq s_1$,

$$v(y, s) = S_1(y, s) + S_2(y, s) + S_3(y, s),$$

where

$$(7.12) \quad S_1(y, s) := e^{-\beta s} \langle \tilde{\phi}_l, \phi_l \rangle \phi_l,$$

$$(7.13) \quad S_2(y, s) := \sum_{n \neq l} e^{-(\lambda_n - \alpha)(s - s_1)} \langle \tilde{\phi}_l, \phi_n \rangle e^{-\beta s_1} \phi_n + \sum_{n=0}^{l-1} d_n e^{-(\lambda_n - \alpha)(s - s_1)} \phi_n,$$

$$(7.14) \quad S_3(y, s) := \int_{s_1}^s e^{-\mathcal{A}(s-\tau)} f(v(y, \tau)) d\tau.$$

LEMMA 7.4. For any $v \in (0, 1)$, there exist $s_1 \gg 1$ and $K \gg \tilde{K}$ such that

$$|S_2(y, s)| \leq v e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}),$$

if $K e^{-\beta s} \leq y \leq e^{\sigma s}$ and $s_1 \leq s \leq s_1 + 1$, where $\sigma \in (0, \tilde{\sigma})$.

PROOF. Given any $v \in (0, 1)$. Note that $S_2(y, s)$ satisfies

$$(S_2)_s = (S_2)_{yy} - \frac{1}{2} y (S_2)_y + \alpha S_2 - \frac{b}{y^2} S_2.$$

Set $S_2(y, s) = y^{2\alpha-1} V(y, s)$. We have

$$V_s = V_{yy} + \left(\frac{2(2\alpha-1)}{y} - \frac{y}{2} \right) V_y + \frac{1}{2} V.$$

Moreover, by setting $\bar{V} = e^{-s/2} V$, we obtain that

$$\bar{V}_s = \bar{V}_{yy} + \left(\frac{2(2\alpha-1)}{y} - \frac{y}{2} \right) \bar{V}_y.$$

From Proposition 6.1 of [1], it follows that

$$\begin{aligned} \bar{V}(y, s) &= c_\alpha \frac{\exp((4\alpha-3)(s-s_1)/4)}{1 - e^{-(s-s_1)}} \int_0^\infty I_{(4\alpha-3)/2} \left(\frac{e^{-(s-s_1)/2} r y}{2(1 - e^{-(s-s_1)})} \right) \\ &\quad \cdot r^{(4\alpha-1)/2} y^{-(4\alpha-3)/2} \exp\left(-\frac{y^2 e^{-(s-s_1)} + r^2}{4(1 - e^{-(s-s_1)})}\right) \bar{V}(r, s_1) dr, \end{aligned}$$

where $I_\mu(z)$ is the modified Bessel function of order μ and c_α is a positive constant depending on α . Hence

$$(7.15) \quad S_2(y, s) = Cy^{2\alpha-1} \frac{\exp((4\alpha-1)(s-s_1)/4)}{1-e^{-(s-s_1)}} \int_0^\infty H(r, y; s-s_1) r^{1/2} v_0 dr,$$

where $v_0(y, s_1) := v(y, s_1) - e^{-\beta s_1} \langle \tilde{\phi}_l, \phi_l \rangle \phi_l(y)$ and

$$(7.16) \quad H(r, y; s-s_1) := I_{(4\alpha-3)/2} \left(\frac{e^{-(s-s_1)/2} r y}{2(1-e^{-(s-s_1)})} \right) \exp \left(-\frac{y^2 e^{-(s-s_1)} + r^2}{4(1-e^{-(s-s_1)})} \right) y^{-(4\alpha-3)/2}.$$

Since

$$|I_\mu(z)| \leq \frac{Cz^\mu e^z}{(1+z)^{\mu+1/2}},$$

from (7.15) we have for $s_1 \leq s \leq s_1 + 1$

$$|S_2(y, s)| \leq \frac{Cy^{2\alpha-1}}{1-e^{-(s-s_1)}} \int_0^\infty T(r, y; s-s_1) r^{1/2} |v_0| dr := S_{2,1} + S_{2,2},$$

where

$$\begin{aligned} S_{2,1} &:= \frac{Cy^{2\alpha-1}}{1-e^{-(s-s_1)}} \int_0^{\tilde{K}e^{-\beta s_1}} T(r, y; s-s_1) r^{1/2} |v_0| dr, \\ S_{2,2} &:= \frac{Cy^{2\alpha-1}}{1-e^{-(s-s_1)}} \int_{\tilde{K}e^{-\beta s_1}}^\infty T(r, y; s-s_1) r^{1/2} |v_0| dr, \\ T(r, y; s-s_1) &:= \frac{(e^{-(s-s_1)/2} r y / 2(1-e^{-(s-s_1)}))^{(4\alpha-3)/2}}{(1+e^{-(s-s_1)/2} r y / 2(1-e^{-(s-s_1)}))^{2\alpha-1}} \\ &\quad \cdot \exp \left(-\frac{|y e^{-(s-s_1)/2} - r|^2}{4(1-e^{-(s-s_1)})} \right) y^{-2\alpha+3/2}. \end{aligned}$$

Firstly, we consider $S_{2,1}$. Using (V1) and (6.10), for $s_1 \gg 1$ we have

$$(7.17) \quad |v_0(y, s_1)| \leq C\tilde{K}^{2\alpha} e^{-2\alpha\beta s_1} \quad \text{for } y \in [0, \tilde{K}e^{-\beta s_1}].$$

From (7.17), we have

$$S_{2,1}(y, s) \leq \frac{C\tilde{K}^{2\alpha} e^{-2\alpha\beta s_1} y^{2\alpha-1}}{1-e^{-(s-s_1)}} \int_0^{\tilde{K}e^{-\beta s_1}} T(r, y; s-s_1) r^{1/2} dr.$$

For $y \geq Ke^{-\beta s}$ and $r \leq \tilde{K}e^{-\beta s_1}$, we have (using $s_1 \leq s \leq s_1 + 1$)

$$y e^{-(s-s_1)/2} - r \geq r \quad \text{if } K \geq 2\tilde{K}e^{\beta+1/2}.$$

Hence we obtain

$$S_{2,1}(y, s) \leq \frac{C\tilde{K}^{2\alpha} e^{-2\alpha\beta s_1}}{(1-e^{-(s-s_1)})^{1/2}} \int_0^{\tilde{K}e^{-\beta s_1}} \exp \left(-\frac{r^2}{4(1-e^{-(s-s_1)})} \right) dr.$$

Set $t = r/(1 - e^{-(s-s_1)})^{1/2}$. We then have

$$\begin{aligned} S_{2,1}(y, s) &\leq C \tilde{K}^{2\alpha} e^{-2\alpha\beta s_1} \int_0^\infty \exp\left(-\frac{t^2}{4}\right) dt \\ &\leq C \tilde{K}^{2\alpha} e^{-2\alpha\beta s_1} y^{2\alpha-1} (K e^{-\beta s})^{-2\alpha+1} \\ &\leq \nu e^{-\beta s} y^{2\alpha-1} \quad \text{if } K \gg \tilde{K}. \end{aligned}$$

Secondly, we consider $S_{2,2}(y, s)$. From Lemma 7.3, (4.5) and (4.6), for the given $\nu \in (0, 1)$ there exists $s_1 \gg 1$ such that

$$(7.18) \quad |v_0(y, s_1)| \leq \nu e^{-\beta s_1} (y^{2\alpha-1} + y^{2\lambda_l}) \quad \text{for } \tilde{K} e^{-\beta s_1} \leq y \leq e^{\tilde{\sigma} s_1}.$$

Moreover, it also holds

$$(7.19) \quad |v_0(y, s_1)| \leq C y^{2\lambda_l} \quad \text{for } y \geq e^{\tilde{\sigma} s_1}.$$

From (7.18) and (7.19), for $s_1 \gg 1$ we can estimate $S_{2,2}$ by

$$S_{2,2} \leq S_{2,2}^1 + S_{2,2}^2 + S_{2,2}^3,$$

where

$$\begin{aligned} S_{2,2}^1 &= \frac{C y^{2\alpha-1}}{1 - e^{-(s-s_1)}} \int_{e^{\tilde{\sigma} s_1}}^\infty T(r, y; s - s_1) r^{1/2+2\lambda_l} dr, \\ S_{2,2}^2 &= \frac{C \nu e^{-\beta s_1} y^{2\alpha-1}}{1 - e^{-(s-s_1)}} \int_{2y e^{-(s-s_1)/2}}^\infty T(r, y; s - s_1) r^{2\alpha-1/2} (1 + r^{2\lambda_l-2\alpha+1}) dr, \\ S_{2,2}^3 &= \frac{C \nu e^{-\beta s_1} y^{2\alpha-1}}{1 - e^{-(s-s_1)}} \int_0^{2y e^{-(s-s_1)/2}} T(r, y; s - s_1) r^{2\alpha-1/2} (1 + r^{2\lambda_l-2\alpha+1}) dr. \end{aligned}$$

Since

$$r - y e^{-(s-s_1)/2} \geq r(1 - e^{-s_1(\tilde{\sigma}-\sigma)}) \geq r/2 \quad \text{if } r \geq e^{\tilde{\sigma} s_1}, y \leq e^{\sigma s} \quad \text{and } s_1 \gg 1,$$

we have

$$\begin{aligned} S_{2,2}^1 &\leq \frac{C y^{2\alpha-1}}{1 - e^{-(s-s_1)}} \int_{e^{\tilde{\sigma} s_1}}^\infty \left(\frac{r}{2(1 - e^{-(s-s_1)})}\right)^{(4\alpha-3)/2} \exp\left(-\frac{r^2}{16(1 - e^{-(s-s_1)})}\right) r^{1/2+2\lambda_l} dr \\ &\leq \frac{C y^{2\alpha-1}}{1 - e^{-(s-s_1)}} e^{-e^{2\tilde{\sigma} s_1}/32} \int_{e^{\tilde{\sigma} s_1}}^\infty \left(\frac{r}{2(1 - e^{-(s-s_1)})}\right)^{(4\alpha-3)/2} \\ &\quad \cdot \exp\left(-\frac{r^2}{32(1 - e^{-(s-s_1)})}\right) r^{1/2+2\lambda_l} dr. \end{aligned}$$

Set $t = r/(1 - e^{-(s-s_1)})^{1/2}$. Then

$$\begin{aligned} S_{2,2}^1 &\leq C y^{2\alpha-1} e^{-C e^{2\tilde{\sigma} s_1}} \int_0^\infty t^{2\alpha+2\lambda_l-1} e^{-t^2/32} dt \\ &\leq \nu e^{-\beta s} y^{2\alpha-1} \quad \text{if } s_1 \gg 1. \end{aligned}$$

For $S_{2,2}^2$, since

$$r - ye^{-(s-s_1)/2} \geq r/2 \quad \text{if } r \geq 2ye^{-(s-s_1)/2},$$

we have

$$\begin{aligned} S_{2,2}^2 &\leq \frac{Cve^{-\beta s_1} y^{2\alpha-1}}{1 - e^{-(s-s_1)}} \int_{2ye^{-(s-s_1)/2}}^{\infty} \left(\frac{r}{2(1 - e^{-(s-s_1)})} \right)^{(4\alpha-3)/2} \\ &\quad \cdot \exp\left(\frac{-r^2}{16(1 - e^{-(s-s_1)})} \right) r^{2\alpha-1/2} (1 + r^{2\lambda_l - 2\alpha + 1}) dr. \end{aligned}$$

Set $t = r/(1 - e^{-(s-s_1)})^{1/2}$. Then

$$\begin{aligned} S_{2,2}^2 &\leq Cve^{-\beta s_1} y^{2\alpha-1} \int_0^{\infty} t^{4\alpha-2} e^{-t^2/16} (1 + t^{2\lambda_l - 2\alpha + 1}) dt \\ &\leq Cve^{-\beta s} y^{2\alpha-1} \quad \text{if } s_1 \gg 1. \end{aligned}$$

Finally, since

$$e^{-(s-s_1)/2} ry \geq r^2/2 \quad \text{if } r \leq 2ye^{-(s-s_1)/2},$$

we have

$$\begin{aligned} S_{2,2}^3 &\leq \frac{Cve^{-\beta s_1} y^{2\alpha-1}}{1 - e^{-(s-s_1)}} \int_0^{2ye^{-(s-s_1)/2}} \left(\frac{r}{2(1 - e^{-(s-s_1)})} \right)^{(4\alpha-3)/2} \left(\frac{r^2}{4(1 - e^{-(s-s_1)})} \right)^{1-2\alpha} \\ &\quad \cdot \exp\left(-\frac{|ye^{-(s-s_1)/2} - r|^2}{4(1 - e^{-(s-s_1)})} \right) r^{2\alpha-1/2} (1 + r^{2\lambda_l - 2\alpha + 1}) dr \\ &\leq \frac{Cve^{-\beta s_1} y^{2\alpha-1}}{(1 - e^{-(s-s_1)})^{1/2}} (1 + y^{2\lambda_l - 2\alpha + 1}) \int_0^{\infty} \exp\left(-\frac{|ye^{-(s-s_1)/2} - r|^2}{4(1 - e^{-(s-s_1)})} \right) dr \\ &\leq Cve^{-\beta s_1} (y^{2\alpha-1} + y^{2\lambda_l}) \quad \text{if } s_1 \gg 1. \end{aligned}$$

This completes the proof.

LEMMA 7.5. *For any $v \in (0, 1)$, there exist $s_1 \gg 1$ such that*

$$|S_3(y, s)| \leq ve^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}),$$

if $Ke^{-\beta s} \leq y \leq e^{\sigma s}$ and $s_1 \leq s \leq s_1 + 1$, where $\sigma \in ((\lambda_l - \alpha)/(2\lambda_l - 2\alpha + 1), \tilde{\sigma})$.

PROOF. Let $Z(y, s) = e^{-\mathcal{A}(s-\tau)} f(v(y, \tau))$, $s_1 < \tau < s$. Then

$$Z_s = -\mathcal{A}Z = Z_{yy} - \frac{y}{2}Z_y - \frac{b}{y^2}Z + \alpha Z.$$

Following the same reasoning as in the proof of Lemma 7.4, we have

$$Z(y, s) = Cy^{2\alpha-1} \frac{\exp((4\alpha-1)(s-\tau)/4)}{1 - e^{-(s-\tau)}} \int_0^{\infty} H(r, y; s-\tau) r^{1/2} f(v(r, \tau)) dr,$$

where H is defined in (7.16). Therefore, we obtain

$$S_3(y, s) = Cy^{2\alpha-1} \int_{s_1}^s \frac{\exp((4\alpha-1)(s-\tau)/4)}{1 - e^{-(s-\tau)}} \int_0^{\infty} H(r, y; s-\tau) r^{1/2} f(v(r, \tau)) dr d\tau.$$

Thus

$$\begin{aligned} |S_3(y, s)| &\leq Cy^{2\alpha-1} \int_{s_1}^s \frac{1}{(1 - e^{-(s-\tau)})^{(4\alpha-1)/2}} \int_0^\infty \left(1 + \frac{e^{-(s-\tau)/2} ry}{2(1 - e^{-(s-\tau)})}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-\frac{|ye^{-(s-\tau)/2} - r|^2}{4(1 - e^{-(s-\tau)})}\right) r^{2\alpha-1} |f(v(r, \tau))| dr d\tau \\ &\leq S_{3,1} + S_{3,2} + S_{3,3}, \end{aligned}$$

where

$$\begin{aligned} S_{3,1} &:= Cy^{2\alpha-1} \int_{s_1}^s (s - \tau)^{-2\alpha+1/2} \int_0^{Le^{-\beta\tau}} \left(1 + \frac{Cry}{s - \tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s - \tau}\right) r^{2\alpha-1} |f(v(r, \tau))| dr d\tau, \\ S_{3,2} &:= Cy^{2\alpha-1} \int_{s_1}^s (s - \tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{e^{\sigma\tau}} \left(1 + \frac{Cry}{s - \tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s - \tau}\right) r^{2\alpha-1} |f(v(r, \tau))| dr d\tau, \\ S_{3,3} &:= Cy^{2\alpha-1} \int_{s_1}^s (s - \tau)^{-2\alpha+1/2} \int_{e^{\sigma\tau}}^\infty \left(1 + \frac{Cry}{s - \tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s - \tau}\right) r^{2\alpha-1} |f(v(r, \tau))| dr d\tau, \end{aligned}$$

by using $(s - \tau)/e \leq 1 - e^{-(s-\tau)} \leq (s - \tau)$ for $s - \tau \in [0, 1]$. Here $1 \ll L \ll K$.

I. Estimate of $S_{3,1}$. By the same way of estimating (7.3), we have

$$\begin{aligned} S_{3,1} &\leq CL^{2\alpha} e^{-2\alpha\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s - \tau)^{-2\alpha+1/2} \\ &\quad \cdot \int_0^{Le^{-\beta\tau}} \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s - \tau}\right) r^{2\alpha-3} dr d\tau. \end{aligned}$$

Since $y \geq Ke^{-\beta s}$ and $r \leq Le^{-\beta\tau}$, we get

$$\exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s - \tau}\right) \leq \exp\left(\frac{-C(y^2 + r^2)}{s - \tau}\right).$$

Set $u = r/(s - \tau)^{1/2}$. Then we obtain

$$S_{3,1} \leq CL^{2\alpha} e^{-2\alpha\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s - \tau)^{-\alpha-1/2} e^{-Cy^2/(s-\tau)} \int_0^{Le^{-\beta\tau}/\sqrt{s-\tau}} e^{-Cu^2} u^{2\alpha-3} du d\tau.$$

Set $t = y^2/(s - \tau)$. Then

$$S_{3,1} \leq CL^{2\alpha} e^{-2\alpha\beta s_1} y^{2\alpha-1} \int_{y^2/(s-s_1)}^\infty t^{\alpha-3/2} e^{-Ct} y^{-2\alpha+1} \int_0^\infty e^{-Cu^2} u^{2\alpha-3} du dt.$$

Since $y \geq Ke^{-\beta s}$, we obtain that

$$S_{3,1} \leq CL^{2\alpha} e^{-2\alpha\beta s_1} y^{2\alpha-1} (Ke^{-\beta s})^{-2\alpha+1} \leq ve^{-\beta s_1} y^{2\alpha-1} \quad \text{if } K \gg L.$$

II. Estimate of $S_{3,2}$. From (7.4), (7.5) and Proposition 6.1, we have $S_{3,2} \leq S_{3,2}^1 + S_{3,2}^2$, where

$$\begin{aligned} S_{3,2}^1 &:= Ce^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{e^{\sigma\tau}} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-5} dr d\tau, \\ S_{3,2}^2 &:= Ce^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{e^{\sigma\tau}} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\lambda_l-3} dr d\tau. \end{aligned}$$

Firstly, we consider $S_{3,2}^1$. Note that $S_{3,2}^1 \leq S_{3,2}^{1,1} + S_{3,2}^{1,2}$, where

$$\begin{aligned} S_{3,2}^{1,1} &:= Ce^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{4y} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-5} dr d\tau, \\ S_{3,2}^{1,2} &:= Ce^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{4y}^{\infty} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-5} dr d\tau. \end{aligned}$$

Consider $S_{3,2}^{1,1}$. Since $ry \geq r^2/4$, if $r \leq 4y$, we have

$$\begin{aligned} S_{3,2}^{1,1} &\leq Ce^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{4y} \left(1 + \frac{Cr^2}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-5} dr d\tau. \end{aligned}$$

Set $u = r/(s-\tau)^{1/2}$. Then

$$\begin{aligned} S_{3,2}^{1,1} &\leq Ce^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-3/2} \int_{(Le^{-\beta\tau})/\sqrt{s-\tau}}^{4y/\sqrt{s-\tau}} (1 + Cu^2)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right|^2\right) u^{4\alpha-5} du d\tau. \end{aligned}$$

To estimate $S_{3,2}^{1,1}$, we divide it into two cases and define

$$D_1 := \left\{ u \in \left[\frac{Le^{-\beta\tau}}{\sqrt{s-\tau}}, \frac{4y}{\sqrt{s-\tau}} \right]; \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right| \geq \frac{ye^{-(s-\tau)/2}}{2\sqrt{s-\tau}} \right\},$$

$$D_2 := \left\{ u \in \left[\frac{Le^{-\beta\tau}}{\sqrt{s-\tau}}, \frac{4y}{\sqrt{s-\tau}} \right]; \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right| \leq \frac{ye^{-(s-\tau)/2}}{2\sqrt{s-\tau}} \right\}.$$

Since

$$\begin{aligned} & \int_{D_1} (1 + Cu^2)^{-2\alpha+1} \exp\left(-C \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right|^2\right) u^{4\alpha-5} du \\ & \leq \int_0^{4y/\sqrt{s-\tau}} e^{-Cy^2/(s-\tau)} u^{4\alpha-5} du \leq C e^{-Cy^2/(s-\tau)} (y^2/(s-\tau))^{2\alpha-2}, \end{aligned}$$

by setting $t = y^2/(s-\tau)$, we have

$$\begin{aligned} S_{3,2}^{1,1,1} & := C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-3/2} \int_{D_1} (1 + Cu^2)^{-2\alpha+1} \\ & \quad \cdot \exp\left(-C \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right|^2\right) u^{4\alpha-5} dud\tau \\ & \leq C e^{-2\beta s_1} y^{2\alpha-2} \int_0^\infty t^{2\alpha-5/2} e^{-Ct} dt \\ & \leq C e^{-2\beta s_1} y^{2\alpha-1} (Ke^{-\beta s})^{-1} \\ & \leq \nu e^{-\beta s_1} y^{2\alpha-1} \quad \text{if } K \gg 1. \end{aligned}$$

Moreover, for any $a \in (0, 1/8)$, we have

$$\begin{aligned} & \int_{D_2} (1 + Cu^2)^{-2\alpha+1} \exp\left(-C \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right|^2\right) u^{4\alpha-5} du \\ & \leq \int_{D_2} u^{4\alpha-3+a} (1 + Cu^2)^{-2\alpha+3/2-a/2} (1 + Cu^2)^{-1/2+a/2} u^{-2-a} du \\ & \leq \int_{D_2} u^{-2-a} du \leq Cy^{-1-a} (s-\tau)^{1/2+a/2}, \end{aligned}$$

since

$$\frac{ye^{-(s-\tau)/2}}{2\sqrt{s-\tau}} \leq u \leq \frac{3ye^{-(s-\tau)/2}}{2\sqrt{s-\tau}} \quad \text{if } u \in D_2.$$

Hence, by noting that $K \in (e^{\beta s_1/8}, e^{\beta s_1/2})$,

$$\begin{aligned} S_{3,2}^{1,1,2} &:= C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-3/2} \int_{D_2} (1+Cu^2)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \left| \frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u \right|^2\right) u^{4\alpha-5} dud\tau \\ &\leq C e^{-2\beta s_1} y^{2\alpha-1} y^{-1-a} \int_{s_1}^s (s-\tau)^{-1+a/2} d\tau \\ &\leq C e^{-2\beta s_1} y^{2\alpha-1} (K e^{-\beta s})^{-1-a} \\ &\leq \nu e^{-\beta s_1} y^{2\alpha-1} \quad \text{if } s_1 \gg 1. \end{aligned}$$

We conclude that

$$S_{3,2}^{1,1} \leq \nu e^{-\beta s_1} y^{2\alpha-1} \quad \text{if } s_1 \gg 1.$$

Consider $S_{3,2}^{1,2}$. Since

$$|ye^{-(s-\tau)/2} - r|^2 \geq y^2 e^{-1} + r^2/2 \quad \text{if } r \geq 4y,$$

we have

$$S_{3,2}^{1,2} \leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{4y}^{\infty} e^{-Cy^2/(s-\tau)} e^{-Cr^2/(s-\tau)} r^{4\alpha-5} dr d\tau.$$

Set $u = r/(s-\tau)^{1/2}$. Then

$$S_{3,2}^{1,2} \leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-3/2} \int_{4y/\sqrt{s-\tau}}^{\infty} e^{-Cy^2/(s-\tau)} e^{-Cu^2} u^{4\alpha-5} dud\tau.$$

Set $t = y^2/(s-\tau)$. Then we get

$$\begin{aligned} S_{3,2}^{1,2} &\leq C e^{-2\beta s_1} y^{2\alpha-1} y^{-1} \int_{y^2/(s-s_1)}^{\infty} t^{-1/2} e^{-Ct} \int_{4\sqrt{t}}^{\infty} e^{-Cu^2} u^{4\alpha-5} dudt \\ &\leq C e^{-2\beta s_1} y^{2\alpha-1} (K e^{-\beta s})^{-1} \leq \nu e^{-\beta s_1} y^{2\alpha-1} \quad \text{if } K \gg 1. \end{aligned}$$

Secondly, we consider $S_{3,2}^2$. Note that $S_{3,2}^2 \leq S_{3,2}^{2,1} + S_{3,2}^{2,2}$, where

$$\begin{aligned} S_{3,2}^{2,1} &:= C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{4y} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\lambda_l-3} dr d\tau, \\ S_{3,2}^{2,2} &:= C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{4y}^{\infty} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\lambda_l-3} dr d\tau. \end{aligned}$$

Consider $S_{3,2}^{2,1}$. Since $ry \geq r^2/4$, if $r \leq 4y$, we have

$$\begin{aligned} S_{3,2}^{2,1} &\leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{Le^{-\beta\tau}}^{4y} \left(1 + \frac{Cr^2}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\lambda_l-3} dr d\tau. \end{aligned}$$

Set $u = r/(s-\tau)^{1/2}$. Then we have

$$\begin{aligned} S_{3,2}^{2,1} &\leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{2\lambda_l-2\alpha-1/2} \\ &\quad \cdot \int_{Le^{-\beta\tau}/\sqrt{s-\tau}}^{4y/\sqrt{s-\tau}} \exp\left(-C \left|\frac{ye^{-(s-\tau)/2}}{\sqrt{s-\tau}} - u\right|^2\right) u^{4\lambda_l-4\alpha-1} du d\tau \\ &\leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{2\lambda_l-2\alpha-1/2} \left(\frac{4y}{\sqrt{s-\tau}}\right)^{4\lambda_l-4\alpha} d\tau \\ &\leq C e^{-2\beta s_1} y^{4\lambda_l-2\alpha-1} \leq C e^{-\beta s_1} y^{2\lambda_l} (e^{-\beta s_1} e^{\sigma(2\lambda_l-2\alpha-1)s}) \\ &\leq \nu e^{-\beta s_1} y^{2\lambda_l} \quad \text{if } s_1 \gg 1. \end{aligned}$$

Consider $S_{3,2}^{2,2}$. Since

$$|ye^{-(s-\tau)/2} - r|^2 \geq y^2 e^{-1} + r^2/2 \quad \text{if } r \geq 4y,$$

we have

$$S_{3,2}^{2,2} \leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{4y}^{\infty} e^{-Cy^2/(s-\tau)} e^{-Cr^2/(s-\tau)} r^{4\lambda_l-3} dr d\tau.$$

Set $u = r/\sqrt{s-\tau}$. Then

$$\begin{aligned} S_{3,2}^{2,2} &\leq C e^{-2\beta s_1} y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{2\lambda_l-2\alpha-1/2} e^{-Cy^2/(s-\tau)} \int_{4y/\sqrt{s-\tau}}^{\infty} e^{-Cu^2} u^{4\lambda_l-3} du d\tau \\ &\leq C e^{-2\beta s_1} y^{2\alpha-1} \leq \nu e^{-\beta s_1} y^{2\alpha-1} \quad \text{if } s_1 \gg 1. \end{aligned}$$

III. Estimate of $S_{3,3}$. From (7.6), we have

$$\begin{aligned} S_{3,3} &\leq C y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{e^{\sigma\tau}}^{\infty} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-3} dr d\tau. \end{aligned}$$

To estimate $S_{3,3}$, we divide it into two cases: $y \leq e^{\sigma s}/4$ and $y \geq e^{\sigma s}/4$.

Case 1: $y \leq e^{\sigma s}/4$. Since

$$\exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) \leq e^{-Ce^{2\sigma\tau}} e^{-Cr^2/(s-\tau)} \quad \text{if } r \geq e^{\sigma\tau} \quad \text{and } y \leq e^{\sigma s}/4,$$

we have

$$S_{3,3} \leq Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{e^{\sigma\tau}}^{\infty} e^{-Ce^{2\sigma\tau}} e^{-Cr^2/(s-\tau)} r^{4\alpha-3} dr d\tau.$$

Set $u = r/\sqrt{s-\tau}$. Then we get

$$\begin{aligned} S_{3,3} &\leq Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-1/2} \int_{e^{\sigma\tau}/\sqrt{s-\tau}}^{\infty} e^{-Ce^{2\sigma\tau}} e^{-Cu^2} u^{4\alpha-3} du d\tau \\ &\leq Ce^{-Ce^{2\sigma s_1}} y^{2\alpha-1} \leq ve^{-\beta s_1} y^{2\alpha-1} \quad \text{if } s_1 \gg 1. \end{aligned}$$

Case 2: $y \geq e^{\sigma s}/4$. Note that $S_{3,3} = S_{3,3}^1 + S_{3,3}^2$, where

$$\begin{aligned} S_{3,3}^1 &:= Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{e^{\sigma\tau}}^{4y} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-3} dr d\tau, \\ S_{3,3}^2 &:= Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{4y}^{\infty} \left(1 + \frac{Cry}{s-\tau}\right)^{-2\alpha+1} \\ &\quad \cdot \exp\left(-C \frac{|ye^{-(s-\tau)/2} - r|^2}{s-\tau}\right) r^{4\alpha-3} dr d\tau. \end{aligned}$$

Consider $S_{3,3}^1$. Since $ry \geq r^2/4$, if $r \leq 4y$, we have

$$S_{3,3}^1 \leq Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{e^{\sigma\tau}}^{4y} \left(1 + \frac{Cr^2}{s-\tau}\right)^{-2\alpha+1} r^{4\alpha-3} dr d\tau.$$

Set $u = r/\sqrt{s-\tau}$. Then we have

$$\begin{aligned} S_{3,3}^1 &\leq Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-1/2} \int_{e^{\sigma\tau}/\sqrt{s-\tau}}^{4y/\sqrt{s-\tau}} u^{-1} du d\tau \\ &\leq Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-1/2} \left(\frac{e^{\sigma\tau}}{\sqrt{s-\tau}}\right)^{-1} \left(\frac{4y}{\sqrt{s-\tau}}\right) d\tau \\ &\leq Cy^{2\alpha} e^{-\sigma s_1} \\ &\leq Ce^{-\beta s_1} y^{2\lambda_l} e^{\beta s_1 + \sigma(-2\lambda_l + 2\alpha - 1)s_1} \\ &\leq ve^{-\beta s_1} y^{2\lambda_l} \quad \text{if } s_1 \gg 1, \end{aligned}$$

where the assumption $\sigma > (\lambda_l - \alpha)/(2\lambda_l - 2\alpha + 1)$ is used.

Consider $S_{3,3}^2$. Since

$$|ye^{-(s-\tau)/2} - r|^2 \geq y^2 e^{-1} + r^2/2 \quad \text{if } r \geq 4y,$$

we have

$$S_{3,3}^2 \leq Cy^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-2\alpha+1/2} \int_{4y}^{\infty} e^{-Cr^2/(s-\tau)} e^{-Cy^2/(s-\tau)} r^{4\alpha-3} dr d\tau.$$

Set $u = r/\sqrt{s-\tau}$. Then we get

$$\begin{aligned} S_{3,3}^2 &\leq C y^{2\alpha-1} \int_{s_1}^s (s-\tau)^{-1/2} \int_{4y/\sqrt{s-\tau}}^{\infty} e^{-Cu^2} u^{4\alpha-3} du d\tau \\ &\leq C y^{2\alpha-1} \leq C e^{-\beta s_1} y^{2\lambda_l} e^{\beta s_1 + \sigma(-2\lambda_l + 2\alpha - 1)s_1} \\ &\leq \nu e^{-\beta s_1} y^{2\lambda_l} \quad \text{if } s_1 \gg 1, \end{aligned}$$

where the assumption $\sigma > (\lambda_l - \alpha)/(2\lambda_l - 2\alpha + 1)$ is used.

Putting together all the above estimates, the lemma is proved.

7.2. Long time: $s_2 > s_1 + 1$.

Since $P(d; s_1, s_2) = 0$, we have

$$d_n = -e^{-\beta s_1} \langle \tilde{\phi}_l, \phi_n \rangle - \int_{s_1}^{s_2} e^{-(\lambda_n - \alpha)(s_1 - \tau)} \langle f(v(\tau)), \phi_n \rangle d\tau$$

for $n = 0, 1, \dots, l-1$. From (7.1) we may write

$$v(y, s) = I_1(y, s) + I_2(y, s) + I_3(y, s) + I_4(y, s),$$

where

$$(7.20) \quad I_1 := e^{-\beta s} \langle \tilde{\phi}_l, \phi_l \rangle \phi_l,$$

$$(7.21) \quad I_2 := \sum_{n=l+1}^{\infty} e^{-(\lambda_n - \alpha)(s - s_1)} \langle \tilde{\phi}_l, \phi_n \rangle e^{-\beta s_1} \phi_n,$$

$$(7.22) \quad I_3 := \sum_{n=0}^{\infty} \int_{s_1}^s e^{-(\lambda_n - \alpha)(s - \tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau,$$

$$(7.23) \quad I_4 := - \sum_{n=0}^{l-1} \int_{s_1}^{s_2} e^{-(\lambda_n - \alpha)(s - \tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau.$$

Notice that $I_3 = S_3$ and $S_2 = I_2 + I_4$.

LEMMA 7.6. *For any $\nu \in (0, 1)$, there exists $s_1 \gg 1$ such that*

$$|I_4(y, s)| \leq \nu e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l})$$

for all $y \geq 0$ and $s_1 + 1 \leq s \leq s_2$.

PROOF. From (4.5), (4.6) and (7.11), we get

$$\begin{aligned} |I_4| &\leq C (y^{2\alpha-1} + y^{2\lambda_l}) \int_{s_1}^{s_2} e^{-(\lambda_n - \alpha)(s - \tau)} e^{-(\beta + \kappa)\tau} d\tau \\ &= C e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}) \int_{s_1}^{s_2} e^{(\lambda_l - \lambda_n)(s - \tau)} e^{-\kappa\tau} d\tau \\ &\leq C e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}) e^{-\kappa s_1} \\ &\leq \nu e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}) \quad \text{if } s_1 \gg 1, \end{aligned}$$

for any $y \geq 0$ and $s \in [s_1 + 1, s_2]$. □

From now on, we shall fix $R > 1$.

LEMMA 7.7. *For any $\nu \in (0, 1)$, there exists $s_1 \gg 1$ such that*

$$|I_2(y, s)| \leq \nu e^{-\beta s} y^{2\alpha-1}$$

for $Ke^{-\beta s} \leq y \leq R$ and $s_1 + 1 \leq s \leq s_2$.

PROOF. Fix $s \geq s_1 + 1$. Since $\tilde{\phi}_l \rightarrow \phi_l$ in L^2_ρ as $s_1 \rightarrow \infty$, $|\langle \tilde{\phi}_l, \phi_n \rangle| \rightarrow 0$ as $s_1 \rightarrow \infty$ uniformly on n for $n \geq l + 1$. Moreover, we have $e^{-(\lambda_n - \alpha)(s - s_1)} e^{-\beta s_1} = e^{-\beta s} e^{-(\lambda_n - \lambda_l)(s - s_1)}$. Hence for a given $\nu \in (0, 1)$ there exists $s_1 \gg 1$ such that

$$|I_2(y, s)| \leq \nu e^{-\beta s} \sum_{n=l+1}^{\infty} e^{-(n-l)} |\phi_n|.$$

Since $\int_0^\infty |\phi_n(y)|^2 \rho(y) dy = 1$ and $\phi_n(y) = c_n y^{2\alpha-1} M(-n, 2\alpha - 1/2; y^2/4)$, we have

$$c_n^2 \cdot 2^{4\alpha-2} \int_0^\infty \xi^{2\alpha-3/2} e^{-\xi} (M(-n, 2\alpha - 1/2; \xi))^2 d\xi = 1.$$

Making use of

$$\int_0^\infty x^{t-1} e^{-x} (M(-n, \hat{\alpha}; \xi))^2 dx = \frac{[\Gamma(\hat{\alpha})]^2 \Gamma(n+1)}{\Gamma(\hat{\alpha} + n)},$$

we have

$$\int_0^\infty \xi^{2\alpha-3/2} e^{-\xi} (M(-n, 2\alpha - 1/2; \xi))^2 d\xi = \frac{[\Gamma(2\alpha - 1/2)]^2 \Gamma(n+1)}{\Gamma(2\alpha + n - 1/2)}.$$

Hence

$$c_n^2 = \frac{\Gamma(2\alpha + n - 1/2)}{2^{4\alpha-2} [\Gamma(2\alpha - 1/2)]^2 \Gamma(n+1)}.$$

Since we know that $\Gamma(x) \sim \sqrt{2\pi} e^{-x} x^{x-1/2}$ as $x \rightarrow \infty$, we get

$$\left(\frac{\Gamma(2\alpha + n - 1/2)}{\Gamma(n+1)} \right)^{1/2} \sim C n^{\alpha-3/4} \quad \text{for } n \gg 1.$$

Thus

$$(7.24) \quad c_n \sim C n^{\alpha-3/4} \quad \text{for } n \gg 1.$$

Moreover, since

$$\begin{aligned} M(-n, 2\alpha - 1/2; y^2/4) &= \pi^{-1/2} \Gamma(2\alpha - 1/2) e^{y^2/4} [(\alpha + n - 1/4) y^2/4]^{1/2-\alpha} \\ &\cdot \cos\left(\frac{\sqrt{4\alpha + 4n - 1}}{2} y - (\alpha - 1/2)\pi\right) (1 + o(1)) \quad \text{as } n \rightarrow \infty \end{aligned}$$

uniformly for $y \in [0, R]$, there exists a positive constant $C = C(R)$ such that

$$(7.25) \quad |M(-n, 2\alpha - 1/2; y^2/4)| \leq C n^{1/2-\alpha} \quad \text{for } y \in [0, R] \text{ and } n \gg 1.$$

From (7.24) and (7.25), we obtain

$$(7.26) \quad |\phi_n| \leq C y^{2\alpha-1} n^{-1/4} \quad \text{for } y \in [0, R] \text{ and } n \gg 1.$$

Since

$$\sum_{n=l+1}^{\infty} e^{-n} n^{-1/4} < \infty,$$

it follows that there exists $s_1 \gg 1$ such that

$$|I_2(y, s)| \leq \nu e^{-\beta s} y^{2\alpha-1}.$$

This proves the lemma.

LEMMA 7.8. *For any $\nu \in (0, 1)$, there exists $s_1 \gg 1$ such that*

$$|I_3(y, s)| \leq \nu e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l})$$

for $Ke^{-\beta s} \leq y \leq R$ and $s_1 + 1 \leq s \leq s_2$.

PROOF. Given any $\nu \in (0, 1)$, note that $I_3 = I_{3,1} + I_{3,2}$, where

$$I_{3,1} := \sum_{n=1}^{l-1} \int_{s_1}^s e^{-(\lambda_n - \alpha)(s-\tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau,$$

$$I_{3,2} := \sum_{n=l}^{\infty} \int_{s_1}^s e^{-(\lambda_n - \alpha)(s-\tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau.$$

By the same reasoning as the proof of Lemma 7.6, it follows immediately that

$$|I_{3,1}| \leq \nu e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}) \quad \text{if } s_1 \gg 1.$$

We write $I_{3,2} = I_{3,2}^1 + I_{3,2}^2$, where

$$I_{3,2}^1 := \sum_{n=l}^{\infty} \int_{s-1}^s e^{-(\lambda_n - \alpha)(s-\tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau,$$

$$I_{3,2}^2 := \sum_{n=l}^{\infty} \int_{s_1}^{s-1} e^{-(\lambda_n - \alpha)(s-\tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau.$$

Since

$$I_{3,2}^1 = \int_{s-1}^s e^{-\mathcal{A}(s-\tau)} f(v(\tau)) d\tau - \sum_{n=0}^{l-1} \int_{s-1}^s e^{-(\lambda_n - \alpha)(s-\tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau,$$

by the same arguments as the proofs of Lemma 7.5 and the estimate of $I_{3,1}$, we get

$$I_{3,2}^1 \leq \nu e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l}) \quad \text{if } s_1 \gg 1.$$

For $I_{3,2}^2$, we have

$$I_{3,2}^2 = \int_{s_1}^{s-1} e^{-\beta(s-\tau)} \sum_{n=l}^{\infty} e^{-(\lambda_n - \lambda_l)(s-\tau)} \langle f(v(\tau)), \phi_n \rangle \phi_n d\tau.$$

Thus

$$|I_{3,2}^2| \leq \int_{s_1}^{s-1} e^{-\beta(s-\tau)} \left(\sum_{n=l}^{\infty} e^{-2(\lambda_n-\lambda_l)(s-\tau)} |\phi_n|^2 \lambda_n^2 \right)^{1/2} \left(\sum_{n=l}^{\infty} \frac{|(f(v(\tau)), \phi_n)|^2}{\lambda_n^2} \right)^{1/2} d\tau.$$

From (7.26), we have for $n \gg 1$ and $\tau \in [s_1, s-1]$

$$e^{-2(\lambda_n-\lambda_l)(s-\tau)} |\phi_n|^2 \lambda_n^2 \leq C e^{-2n} n^{3/2} y^{4\alpha-2}.$$

Since $\sum_{n=l}^{\infty} e^{-2n} n^{3/2} < \infty$, we obtain

$$\left(\sum_{n=l}^{\infty} e^{-2(\lambda_n-\lambda_l)(s-\tau)} |\phi_n|^2 \lambda_n^2 \right)^{1/2} \leq C y^{2\alpha-1}.$$

Moreover, from (7.26)

$$\begin{aligned} \left(\sum_{n=l}^{\infty} \frac{|(f(v(r, \tau)), \phi_n)|^2}{\lambda_n^2} \right)^{1/2} &\leq \sum_{n=l}^{\infty} \frac{|(f(v(r, \tau)), \phi_n)|}{\lambda_n} \\ &\leq \sum_{n=l}^{\infty} \frac{\int_0^{\infty} |f(v(r, \tau))| |\phi_n| e^{-r^2/4} dr}{\lambda_n} \\ &\leq C \sum_{n=l}^{\infty} \frac{n^{-1/4}}{\alpha - 1/2 + n} \int_0^{\infty} |f(v(r, \tau))| r^{2\alpha-1} e^{-r^2/4} dr \\ &\leq C \int_0^{\infty} |f(v(r, \tau))| r^{2\alpha-1} e^{-r^2/4} dr, \end{aligned}$$

since $\sum_{n=l}^{\infty} n^{-5/4} < \infty$. Hence we have

$$(7.27) \quad |I_{3,2}^2| \leq C y^{2\alpha-1} \int_{s_1}^{s-1} e^{-\beta(s-\tau)} \int_0^{\infty} |f(v(r, \tau))| r^{2\alpha-1} e^{-r^2/4} dr d\tau.$$

From Lemma 7.2, we have the estimate

$$\int_0^{\infty} |f(v(r, \tau))| r^{2\alpha-1} e^{-r^2/4} dr \leq J_1 + J_2 + J_3,$$

where

$$\begin{aligned} J_1 &:= C K^{2\alpha} e^{-2\alpha\beta\tau} \int_0^{K e^{-\beta\tau}} r^{2\alpha-3} e^{-r^2/4} dr, \\ J_2 &:= C e^{-2\beta\tau} \int_{K e^{-\beta\tau}}^{e^{\sigma\tau}} (r^{2\alpha-4} + r^{4\lambda_l-2\alpha-2}) r^{2\alpha-1} e^{-r^2/4} dr, \\ J_3 &:= C e^{-2\beta\tau} \int_{e^{\sigma\tau}}^{\infty} r^{4\alpha-3} e^{-r^2/4} dr. \end{aligned}$$

By a similar argument to the proof of Lemma 7.3, we have

$$\int_0^{\infty} |f(v(r, \tau))| r^{2\alpha-1} e^{-r^2/4} dr \ll e^{-\beta\tau} \quad \text{if } s_1 \gg 1.$$

Plugging into (7.27), we obtain

$$|I_{3,2}^2| \leq \nu e^{-\beta s} y^{2\alpha-1} \quad \text{if } s_1 \gg 1.$$

This proves the lemma.

LEMMA 7.9. *For any $\nu \in (0, 1)$, there exists $s_1 \gg 1$ such that*

$$|I_2(y, s)| \leq \nu e^{-\beta s} (y^{2\alpha-1} + y^{2\lambda_l})$$

for $y \in \{K e^{-\beta s} \leq y \leq e^{\sigma s}\} \cap \{y \geq e^{(s-s_1)/2}\}$ and $s_1 + 1 \leq s \leq s_2$.

PROOF. Since $I_2 = S_2 - I_4$, by Lemma 7.6, we only need to estimate S_2 .

From (7.17), (7.18) and (7.19), we could easily find that for a given $\nu \in (0, 1)$ there exists $s_1 \gg 1$ such that

$$(7.28) \quad |v_0(y, s_1)| \leq \nu e^{-\beta s_1} (y^{2\alpha-1} + y^{2\lambda_l}) \quad \text{if } K e^{-\beta s_1} \leq y \leq e^{\sigma s_1},$$

$$(7.29) \quad |v_0(y, s_1)| \leq C K^{2\alpha} e^{-2\alpha\beta s_1} \quad \text{if } y \leq K e^{-\beta s_1},$$

$$(7.30) \quad |v_0(y, s_1)| \leq C y^{2\lambda_l} \quad \text{if } y \geq e^{\sigma s_1}.$$

Moreover, from (7.15) and Lemma 7.2, we have $|S_2| \leq S_2^1 + S_2^2 + S_2^3$, where

$$\begin{aligned} S_2^1 &:= C K^{2\alpha} e^{-2\alpha\beta s_1} y^{1/2} e^{(4\alpha-1)(s-s_1)/4} \int_0^{K e^{-\beta s_1}} r^{1/2} dr, \\ S_2^2 &:= C \nu e^{-\beta s_1} y^{2\alpha-1} e^{(s-s_1)/2} \int_{K e^{-\beta s_1}}^{e^{\sigma s}} (1 + C e^{-(s-s_1)/2} r y)^{-2\alpha+1} \\ &\quad \cdot \exp(-C |y e^{-(s-s_1)/2} - r|^2) r^{2\alpha-1} (r^{2\alpha-1} + r^{2\lambda_l}) dr, \\ S_2^3 &:= C y^{2\alpha-1} e^{(s-s_1)/2} \int_{e^{\sigma s}}^{\infty} r^{2\lambda_l+2\alpha-1} \exp(-C |y e^{-(s-s_1)/2} - r|^2) dr. \end{aligned}$$

First, for S_2^1 , since $y \geq e^{(s-s_1)/2}$ and $K \in (e^{\beta s_1/8}, e^{\beta s_1/2})$, we have

$$\begin{aligned} S_2^1 &\leq C e^{-\beta s} y^{2\lambda_l} \{K^{2\alpha} e^{-2\alpha\beta s_1} y^{-2\lambda_l+1/2} e^{\beta s} e^{(4\alpha-1)(s-s_1)/4} (K e^{-\beta s_1})^{3/2}\}, \\ &\leq \nu e^{-\beta s} y^{2\lambda_l} \quad \text{if } s_1 \gg 1. \end{aligned}$$

Next, we consider S_2^2 . Define

$$D_1 := \{|y e^{-(s-s_1)/2} - r| \leq r/2\}, \quad D_2 := \{|y e^{-(s-s_1)/2} - r| \geq r/2\},$$

and write $S_2^2 = S_2^{2,1} + S_2^{2,2}$, where

$$\begin{aligned} S_2^{2,1} &:= C \nu e^{-\beta s_1} y^{2\alpha-1} e^{(s-s_1)/2} \int_{D_1} (1 + C e^{-(s-s_1)/2} r y)^{-2\alpha+1} \\ &\quad \cdot \exp(-C |y e^{-(s-s_1)/2} - r|^2) r^{2\alpha-1} (r^{2\alpha-1} + r^{2\lambda_l}) dr, \\ S_2^{2,2} &:= C \nu e^{-\beta s_1} y^{2\alpha-1} e^{(s-s_1)/2} \int_{D_2} (1 + C e^{-(s-s_1)/2} r y)^{-2\alpha+1} \\ &\quad \cdot \exp(-C |y e^{-(s-s_1)/2} - r|^2) r^{2\alpha-1} (r^{2\alpha-1} + r^{2\lambda_l}) dr. \end{aligned}$$

Since $e^{-(s-s_1)/2}ry \geq r^2/2$, if $|ye^{-(s-s_1)/2} - r| \leq r/2$. We get

$$\begin{aligned} S_2^{2,1} &\leq Cve^{-\beta s_1}y^{2\alpha-1}e^{(s-s_1)/2} \int_{D_1} \exp(-C|ye^{-(s-s_1)/2} - r|^2)(1 + r^{2\lambda_l-2\alpha+1})dr \\ &\leq Cve^{-\beta s_1}y^{2\alpha-1}e^{(s-s_1)/2}[1 + (ye^{-(s-s_1)/2})^{2\lambda_l-2\alpha+1}] \\ &\leq Cve^{-\beta s}y^{2\lambda_l}\{e^{(\beta+1/2)(s-s_1)}[y^{-2\lambda_l+2\alpha-1} + e^{-(2\lambda_l-2\alpha+1)(s-s_1)/2}]\} \\ &\leq Cve^{-\beta s}y^{2\lambda_l}. \end{aligned}$$

Moreover, we get

$$\begin{aligned} S_2^{2,2} &\leq Cve^{-\beta s_1}y^{2\alpha-1}e^{(s-s_1)/2} \int_{D_2} \exp(-Cr^2)r^{2\alpha-1}(r^{2\alpha-1} + r^{2\lambda_l})dr \\ &\leq Cve^{-\beta s}y^{2\lambda_l}[e^{(\beta+1/2)(s-s_1)}y^{-2\lambda_l+2\alpha-1}] \\ &\leq Cve^{-\beta s}y^{2\lambda_l}. \end{aligned}$$

Finally, we consider S_2^3 . Since $r - ye^{-(s-s_1)/2} \geq Cr$ if $r \geq e^{\sigma s_1}$ and $y \leq e^{\sigma s}$, we have

$$\begin{aligned} S_2^3 &\leq Cy^{2\alpha-1}e^{(s-s_1)/2} \int_{e^{\sigma s_1}}^{\infty} r^{2\lambda_l+2\alpha-1} \exp(-Cr^2)dr, \\ &\leq Cy^{2\alpha-1}e^{(s-s_1)/2}e^{-Ce^{2\sigma s_1}}, \\ &\leq ve^{-\beta s}y^{2\lambda_l} \quad \text{if } s_1 \gg 1. \end{aligned}$$

This completes the proof of the lemma.

LEMMA 7.10. *For any $v \in (0, 1)$, there exists $s_1 \gg 1$ such that*

$$|I_3(y, s)| \leq ve^{-\beta s}y^{2\lambda_l}$$

for $y \in \{Ke^{-\beta s} \leq y \leq e^{\sigma s}\} \cap \{y \geq e^{(s-s_1)/2}\}$ and $s_1 + 1 \leq s \leq s_2$.

PROOF. By Lemma 7.5, we only need to prove that

$$|B(y, s)| \leq ve^{-\beta s}y^{2\lambda_l},$$

where

$$B(y, s) = \int_{s_1}^{s-1} e^{-A(s-\tau)} f(v(y, \tau))d\tau.$$

Following the same argument as the proof of Lemma 7.5, we have

$$\begin{aligned} |B(y, s)| &\leq Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{(s-\tau)/2} \int_0^{\infty} (1 + Ce^{-(s-\tau)/2}ry)^{-2\alpha+1} \\ &\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2)r^{2\alpha-1}|f(v(r, \tau))|drd\tau \\ &= I_3^1 + I_3^2 + I_3^3, \end{aligned}$$

where

$$\begin{aligned}
I_3^1 &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{(s-\tau)/2} \int_0^{Ke^{-\beta\tau}} (1 + Ce^{-(s-\tau)/2}ry)^{-2\alpha+1} \\
&\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2)r^{2\alpha-1}|f(v)|drd\tau, \\
I_3^2 &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{(s-\tau)/2} \int_{Ke^{-\beta\tau}}^{e^{\sigma\tau}} (1 + Ce^{-(s-\tau)/2}ry)^{-2\alpha+1} \\
&\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2)r^{2\alpha-1}|f(v)|drd\tau, \\
I_3^3 &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{(s-\tau)/2} \int_{e^{\sigma\tau}}^{\infty} (1 + Ce^{-(s-\tau)/2}ry)^{-2\alpha+1} \\
&\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2)r^{2\alpha-1}|f(v)|drd\tau.
\end{aligned}$$

Consider I_3^1 . From (7.3) we have

$$\begin{aligned}
I_3^1 &\leq CK^{2\alpha}e^{-2\alpha\beta s_1}y^{2\alpha-1} \int_{s_1}^{s-1} e^{(s-\tau)/2} \int_0^{Ke^{-\beta\tau}} r^{2\alpha-3}drd\tau \\
&= CK^{2\alpha}e^{-2\alpha\beta s_1}y^{2\alpha-1} \int_{s_1}^{s-1} e^{(s-\tau)/2}(Ke^{-\beta\tau})^{2\alpha-2}d\tau \\
&\leq CK^{4\alpha-2}e^{-2\alpha\beta s_1}y^{2\alpha-1}e^{s/2} \int_{s_1}^{s-1} e^{-(1/2+\beta(2\alpha-2))\tau}d\tau \\
&\leq Ce^{-\beta s}y^{2\lambda_l}\{K^{4\alpha-2}y^{-2\lambda_l+2\alpha-1}e^{(\beta+1/2)s}e^{-(2\alpha\beta+\beta(2\alpha-2)+1/2)s_1}\} \\
&\leq ve^{-\beta s}y^{2\lambda_l} \quad \text{if } s_1 \gg 1 \text{ and } y \geq e^{(s-s_1)/2}.
\end{aligned}$$

Consider I_3^2 . From (7.4) and (7.5) we get $I_3^2 \leq I_3^{2,1} + I_3^{2,2}$, where

$$\begin{aligned}
I_3^{2,1} &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau}e^{(s-\tau)/2} \int_{Ke^{-\beta\tau}}^{e^{\sigma\tau}} (1 + Ce^{-(s-\tau)/2}ry)^{-2\alpha+1} \\
&\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2)r^{4\alpha-5}drd\tau, \\
I_3^{2,2} &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau}e^{(s-\tau)/2} \int_{Ke^{-\beta\tau}}^{e^{\sigma\tau}} (1 + Ce^{-(s-\tau)/2}ry)^{-2\alpha+1} \\
&\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2)r^{4\lambda_l-3}drd\tau.
\end{aligned}$$

Firstly, we consider $I_3^{2,1}$. Define

$$D_1 := \{|ye^{-(s-\tau)/2} - r| \leq r/2\}, \quad D_2 := \{|ye^{-(s-\tau)/2} - r| \geq r/2\},$$

and write $I_3^{2,1} = I_3^{2,1,1} + I_3^{2,1,2}$, where

$$\begin{aligned} I_3^{2,1,1} &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} \int_{D_1} (1 + Ce^{-(s-\tau)/2} ry)^{-2\alpha+1} \\ &\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2) r^{4\alpha-5} dr d\tau, \\ I_3^{2,1,2} &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} \int_{D_2} (1 + Ce^{-(s-\tau)/2} ry)^{-2\alpha+1} \\ &\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2) r^{4\alpha-5} dr d\tau. \end{aligned}$$

Consider $I_3^{2,1,1}$. If $r \in D_1$, we have

$$r/2 \leq ye^{-(s-\tau)/2} \leq 3r/2.$$

Hence

$$\begin{aligned} I_3^{2,1,1} &\leq Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} \int_{D_1} r^{-3} dr d\tau \\ &\leq Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} (ye^{-(s-\tau)/2})^{-2} d\tau \\ &\leq Cy^{2\alpha-3} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{3(s-\tau)/2} d\tau \\ &\leq Ce^{-\beta s} y^{2\lambda_l} [y^{-2\lambda_l+2\alpha-3} e^{(\beta+3/2)s} e^{-(2\beta+3/2)s_1}] \\ &\leq ve^{-\beta s} y^{2\lambda_l} \quad \text{if } s_1 \gg 1 \text{ and } y \geq e^{(s-s_1)/2}. \end{aligned}$$

For $I_3^{2,1,2}$, we have

$$\begin{aligned} I_3^{2,1,2} &\leq Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} \int_{D_2} e^{-Cr^2} r^{4\alpha-5} dr d\tau \\ &\leq Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} d\tau \\ &\leq Ce^{-\beta s} y^{2\lambda_l} (y^{-2\lambda_l+2\alpha-1} e^{(\beta+1/2)s} e^{-(2\beta+1/2)s_1}) \\ &\leq ve^{-\beta s} y^{2\lambda_l} \quad \text{if } s_1 \gg 1 \text{ and } y \geq e^{(s-s_1)/2}. \end{aligned}$$

Secondly, we consider $I_3^{2,2}$ and as before we write $I_3^{2,2} = I_3^{2,2,1} + I_3^{2,2,2}$, where

$$\begin{aligned} I_3^{2,2,1} &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} \int_{D_1} (1 + Ce^{-(s-\tau)/2} ry)^{-2\alpha+1} \\ &\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2) r^{4\lambda_l-3} dr d\tau, \\ I_3^{2,2,2} &:= Cy^{2\alpha-1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{(s-\tau)/2} \int_{D_2} (1 + Ce^{-(s-\tau)/2} ry)^{-2\alpha+1} \\ &\quad \cdot \exp(-C|ye^{-(s-\tau)/2} - r|^2) r^{4\lambda_l-3} dr d\tau. \end{aligned}$$

For $I_3^{2,2,1}$, by the same reasoning as the case of $I_3^{2,2,1}$, we obtain

$$\begin{aligned} I_3^{2,2,1} &\leq C y^{4\lambda_l - 2\alpha - 1} \int_{s_1}^{s-1} e^{-2\beta\tau} e^{-(2\lambda_l - 2\alpha - 1/2)(s-\tau)} d\tau \\ &\leq C e^{-\beta s} y^{2\lambda_l} (y^{2\lambda_l - 2\alpha - 1} e^{(-\lambda_l + \alpha + 1/2)s} e^{-s_1/2}) \\ &\leq v e^{-\beta s} y^{2\lambda_l} \quad \text{if } s_1 \gg 1 \text{ and } y \leq e^{\sigma s}. \end{aligned}$$

For $I_3^{2,2,2}$, by the same reasoning as the case of $I_3^{2,1,2}$, we also get

$$I_3^{2,2,2} \leq v e^{-\beta s} y^{2\lambda_l} \quad \text{if } s_1 \gg 1 \text{ and } y \geq e^{(s-s_1)/2}.$$

Now, we consider I_3^3 . Since $r - y e^{-(s-\tau)/2} \geq Cr$ if $r \geq e^{\sigma\tau}$, $y \leq e^{\sigma s}$ and $\tau \leq s - 1$, it follows from (7.6) that

$$\begin{aligned} I_3^3 &\leq C y^{2\alpha - 1} \int_{s_1}^{s-1} e^{(s-\tau)/2} \int_{e^{\sigma\tau}}^{\infty} e^{-Cr^2} r^{4\alpha - 3} dr d\tau \\ &\leq C e^{-C e^{2\sigma s_1}} y^{2\alpha - 1} \int_{s_1}^{s-1} e^{(s-\tau)/2} d\tau \\ &\leq C e^{-\beta s} y^{2\lambda_l} e^{-C e^{2\sigma s_1}} (y^{-2\lambda_l + 2\alpha - 1} e^{(\beta + 1/2)s} e^{-s_1/2}) \\ &\leq v e^{-\beta s} y^{2\lambda_l} \quad \text{if } s_1 \gg 1 \text{ and } y \geq e^{(s-s_1)/2}. \end{aligned}$$

This proves the lemma.

PROOF OF PROPOSITION 6.3. By Lemmas 7.4 and 7.5, it suffices to consider the case when $s_2 > s_1 + 1$. Fix \tilde{s}_1 such that $s_1 + 1 < \tilde{s}_1 < s_2$. Let $R = e^{(\tilde{s}_1 - s_1)/2}$. Set

$$v_0(y, s) = v(y, s) - e^{-\beta s} \langle \tilde{\phi}_l, \phi_l \rangle \phi_l.$$

Note that $v_0(y, s) = S_2(y, s) + S_3(y, s) = I_2(y, s) + I_3(y, s) + I_4(y, s)$. We easily see that (7.29) and (7.30) are true if s_1 is replaced by s for any $s \in [s_1, s_2]$. Moreover, from Lemmas 7.6, 7.7 and 7.8, we have

$$(7.31) \quad |v_0(y, s)| \leq v e^{-\beta s} (y^{2\alpha - 1} + y^{2\lambda_l})$$

for $K e^{-\beta s} \leq y \leq R$, $s \in [s_1 + 1, s_2]$; and from Lemmas 7.6, 7.9 and 7.10, we have

$$(7.32) \quad |v_0(y, s)| \leq v e^{-\beta s} (y^{2\alpha - 1} + y^{2\lambda_l})$$

for $R \leq y \leq e^{\sigma s}$, $s \in [s_1 + 1, \tilde{s}_1]$. Hence (7.28) is true if s_1 is replaced by s for any $s \in [s_1, \tilde{s}_1]$.

Let $s \in [\tilde{s}_1, \tilde{s}_1 + 2 \log R]$. If $R \leq y \leq e^{\sigma s}$, then we have $y \geq e^{(s - \tilde{s}_1)/2}$. Hence, by the same arguments as Lemmas 7.6, 7.9 and 7.10, we have

$$(7.33) \quad |I_2(y, s)| + |I_3(y, s)| + |I_4(y, s)| \leq v e^{-\beta s} (y^{2\alpha - 1} + y^{2\lambda_l})$$

for $R \leq y \leq e^{\sigma s}$ and $\tilde{s}_1 + 1 \leq s \leq \tilde{s}_1 + 2 \log R$. Moreover, by the same arguments as those of Lemmas 7.4 and 7.5, we obtain

$$(7.34) \quad |S_2(y, s)| + |S_3(y, s)| \leq v e^{-\beta s} (y^{2\alpha - 1} + y^{2\lambda_l})$$

for $Ke^{-\beta s} \leq y \leq e^{\sigma s}$ and $\tilde{s}_1 \leq s \leq \tilde{s}_1 + 1$. From (7.33) and (7.34), we have

$$(7.35) \quad |I_2(y, s)| + |I_3(y, s)| + |I_4(y, s)| \leq ve^{-\beta s}(y^{2\alpha-1} + y^{2\lambda_l})$$

for $R \leq y \leq e^{\sigma s}$ and $\tilde{s}_1 \leq s \leq \tilde{s}_1 + 2 \log R$. Continuing to set $\tilde{s}_2 = \tilde{s}_1 + 2 \log R$, $\tilde{s}_3 = \tilde{s}_2 + 2 \log R, \dots$, and following the same argument as above, we have

$$(7.36) \quad |I_2(y, s)| + |I_3(y, s)| + |I_4(y, s)| \leq ve^{-\beta s}(y^{2\alpha-1} + y^{2\lambda_l})$$

for $R \leq y \leq e^{\sigma s}$ and $\tilde{s}_1 \leq s \leq s_2$.

Combining (7.31), (7.32) and (7.36), with the help of Lemmas 7.4 and 7.5, we have

$$(7.37) \quad |v(y, s) - e^{-\beta s} \langle \tilde{\phi}_l, \phi_l \rangle \phi_l| \leq ve^{-\beta s}(y^{2\alpha-1} + y^{2\lambda_l})$$

for $Ke^{-\beta s} \leq y \leq e^{\sigma s}$ and $s_1 \leq s \leq s_2$. Since $\langle \tilde{\phi}_l, \phi_l \rangle \rightarrow 1$ as $s_1 \rightarrow \infty$, we get

$$|v(y, s) - e^{-\beta s} \phi_l| \leq 2ve^{-\beta s}(y^{2\alpha-1} + y^{2\lambda_l})$$

for $Ke^{-\beta s} \leq y \leq e^{\sigma s}$ and $s_1 \leq s \leq s_2$. Hence we have proved Proposition 6.3.

REFERENCES

- [1] M. ABRAMOWITZ AND I. A. STEGUN, Handbook of mathematical functions with formulas, graphs, and mathematical tables, National Bureau of Standards, Applied Mathematics Series 55, Washington, D.C., 1964.
- [2] C. BANDLE AND I. STAKGOLD, The formation of the dead core in parabolic reaction-diffusion problems, Trans. Amer. Math. Soc. 286 (1984), 275–293.
- [3] J. W. DOLD, V. A. GALAKTIONOV, A. A. LACEY AND J. L. VAZQUEZ, Rate of approach to a singular steady state in quasilinear reaction-diffusion equations, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 26 (1998), 663–687.
- [4] J.-S. GUO AND PH. SOUPLLET, Fast rate of formation of dead-core for the heat equation with strong absorption and applications to fast blow-up, Math. Ann. 331 (2005), 651–667.
- [5] D. HENRY, Geometric theory of semilinear parabolic equations, Lecture Notes in Math. 840, Springer, Berlin-New York, 1981.
- [6] M. A. HERRERO AND J. J. L. VELAZQUEZ, A blow up result for semilinear heat equations in the supercritical case, preprint.
- [7] PH. ROSENAU AND S. KAMIN, Thermal waves in an absorbing and convecting medium, Phys. D 8 (1983), 273–283.
- [8] L. K. MARTINSON, The finite velocity of propagation of thermal perturbations in media with constant thermal conductivity, Zh. Vychisl. Mat. Mat. Fiz. 16 (1976), 1233–1243.
- [9] N. MIZOGUCHI, Type-II blowup for a semilinear heat equation, Adv. Differential Equations 9 (2004), 1279–1316.
- [10] PH. SOUPLLET AND J. L. VAZQUEZ, Stabilization towards a singular steady state with gradient blow-up for a diffusion-convection problem, Discrete Contin. Dyn. Syst. 14 (2006), 221–234.
- [11] I. STAKGOLD, Reaction-diffusion problems in chemical engineering, Nonlinear diffusion problems (Montecatini Terme, 1985), 119–152, Lecture Notes in Math. 1224, Springer, Berlin, 1986.

DEPARTMENT OF MATHEMATICS
NATIONAL TAIWAN NORMAL UNIVERSITY
88, S-4, TING CHOU ROAD, TAIPEI 116
TAIWAN

E-mail addresses: jsguo@math.ntnu.edu.tw
: chin@hp715.math.ntnu.edu.tw