

The reduction exponent of socle ideals associated to parameter ideals in a Buchsbaum local ring of multiplicity two

By Shiro GOTO and Hideto SAKURAI

(Received Oct. 23, 2002)

(Revised Jun. 23, 2003)

Abstract. Let A be a Buchsbaum local ring with the maximal ideal \mathfrak{m} and let $e(A)$ denote the multiplicity of A . Let Q be a parameter ideal in A and put $I = Q : \mathfrak{m}$. Then the equality $I^2 = QI$ holds true, if $e(A) = 2$ and $\text{depth } A > 0$. The assertion is no longer true, unless $e(A) = 2$. Counterexamples are given.

1. Introduction.

Let A be a Noetherian local ring with the maximal ideal \mathfrak{m} and $d = \dim A$. Let Q be a parameter ideal in A and let $I = Q : \mathfrak{m}$. In this paper we are interested in the problem of when the equality $I^2 = QI$ holds true. This problem was completely solved by A. Corso, C. Huneke, C. Polini, and W. Vasconcelos [CHV], [CP], [CPV] in the case where A is a Cohen-Macaulay ring. When A is a Buchsbaum ring, partial answers only recently appeared in the authors' paper [GSa], supplying [Y1], [Y2] and [GN] with ample examples of ideals I , for which the Rees algebras $R(I) = \bigoplus_{n \geq 0} I^n$, the associated graded rings $G(I) = R(I)/IR(I)$, and the fiber cones $F(I) = R(I)/\mathfrak{m}R(I)$ are all Buchsbaum rings with certain specific graded local cohomology modules.

This research is a succession of [GSa] and the present purpose is to prove the following, in which $e(A) = e_{\mathfrak{m}}^0(A)$ denotes the multiplicity of A with respect to the maximal ideal \mathfrak{m} .

THEOREM 1.1. *Let A be a Buchsbaum local ring. Assume that $e(A) = 2$ and $\text{depth } A > 0$. Then the equality $I^2 = QI$ holds true for all parameter ideals Q in A , where $I = Q : \mathfrak{m}$.*

The readers may consult [G2] for the structure of Buchsbaum local rings A with $e(A) = 2$. There is given in [G2, Sections 3, 4] a complete list of equi-characteristic non-Cohen-Macaulay Buchsbaum complete local rings A with $e(A) = 2$, $\text{depth } A > 0$, and infinite residue class fields.

In their remarkable papers [CP], [CPV] A. Corso, C. Polini, and W. Vasconcelos proved that the equality $I^2 = QI$ holds true for every parameter ideal Q in A , if A is a Cohen-Macaulay local ring with $e(A) \geq 2$. This is no more true in the Buchsbaum case. As is shown in [GSa, Theorem 4.8], for every integer $e \geq 3$, there exists a

2000 *Mathematics Subject Classification.* Primary 13B22; Secondary 13H10.

Key Words and Phrases. Buchsbaum ring, generalized Cohen-Macaulay ring, Cohen-Macaulay ring, Gorenstein ring, Cohen-Macaulay type, local cohomology, multiplicity.

The first author is supported by the Grant-in-Aid for Scientific Researches (C(2), No. 13640044) Japan Society for the Promotion of Science.

Buchsbaum local ring A with $\dim A = 1$ and $e(A) = e$ which contains a parameter ideal Q such that $I^e = QI^{e-1}$ but $I^{e-1} \neq QI^{e-2}$. Accordingly, without additional assumptions on Buchsbaum local rings A , no hope is left for the equality $I^2 = QI$, at least in the case where $\dim A = 1$ and $e(A) \geq 3$. Our theorem 1.1 settles the case where $e(A) = 2$ and $\text{depth } A > 0$, providing a drastic break-through against the counterexamples of [GSa].

Before entering details, let us briefly note how this paper is organized. We shall prove Theorem 1.1 in Section 3. For the purpose we need some preliminaries, that we will summarize in Section 2. The counterexamples given by [GSa] are all of dimension 1, and according to Theorem 1.1, it might be natural to suspect that the equality $I^2 = QI$ holds true in higher dimensional cases of higher depth. The answer is, nevertheless, still negative. We shall construct examples, showing that for given integers $1 \leq d < m$, there exists a Buchsbaum local ring A with $\dim A = d$, $\text{depth } A = d - 1$, and $e(A) = 2m$, which contains a parameter ideal Q such that $I^3 = QI^2$ but $I^2 \neq QI$ (Theorem 4.5 and Proposition 4.7).

2. Preliminaries.

The purpose of this section is to summarize some preliminary steps, which we need to prove Theorem 1.1. The result might have its own significance. In such a case we shall include a closer proof.

Here let us fix our standard notation. Otherwise specified, let A be a Noetherian local ring with the maximal ideal \mathfrak{m} and $d = \dim A$. For an ideal \mathfrak{a} let $\mathfrak{a}^\#$ be the integral closure of \mathfrak{a} . Let $\ell_A(*)$ and $\mu_A(*)$ respectively denote the length and the number of generators. When A is a Cohen-Macaulay local ring, we denote by $r(A)$ the Cohen-Macaulay type of A , that is

$$r(A) = \ell_A(\text{Ext}_A^d(A/\mathfrak{m}, A)).$$

Let $H_{\mathfrak{m}}^i(*)$ ($i \in \mathbb{Z}$) be the local cohomology functors of A with respect to \mathfrak{m} . We denote by $e(A) = e_{\mathfrak{m}}^0(A)$ the multiplicity of A .

Let us begin with the following.

LEMMA 2.1. *Assume that $e(A) \geq 2$. Then $\mu_A(I) = \ell_A(I/Q) + d$ for every parameter ideal Q in A , where $I = Q : \mathfrak{m}$.*

PROOF. By [GSa, Proposition 2.3] Q is a minimal reduction of I , whence $\mathfrak{m}I = \mathfrak{m}Q$. Thus

$$\mu_A(I) = \ell_A(I/\mathfrak{m}I) = \ell_A(I/Q) + \ell_A(Q/\mathfrak{m}Q) = \ell_A(I/Q) + d$$

as is claimed. □

PROPOSITION 2.2. *Suppose that A is a Cohen-Macaulay local ring with $d = \dim A \geq 1$ and let Q be a parameter ideal in A . Then*

$$\ell_A((0) :_{\mathfrak{m}/Q\mathfrak{m}} \mathfrak{m}) = \begin{cases} r(A) + d & \text{if } Q \neq Q^\#, \\ d & \text{if } Q = Q^\#. \end{cases}$$

PROOF. Since $\dim A \geq 1$, we have $Q\mathfrak{m} : \mathfrak{m} \subseteq \mathfrak{m}$, whence

$$\ell_A((0) :_{\mathfrak{m}/Q\mathfrak{m}} \mathfrak{m}) = \ell_A((Q\mathfrak{m} : \mathfrak{m})/Q\mathfrak{m}).$$

Let $I = Q : \mathfrak{m}$. Firstly, assume that $Q \neq Q^\#$. Then Q is a minimal reduction of I , because $I^2 = QI$ (cf. e.g., [GH, Proposition 3.4]). Hence $\mathfrak{m}I = \mathfrak{m}Q$, so that we have $I \subseteq Q\mathfrak{m} : \mathfrak{m}$; thus $I = Q\mathfrak{m} : \mathfrak{m}$. Consequently,

$$\ell_A((0) :_{\mathfrak{m}/Q\mathfrak{m}} \mathfrak{m}) = \ell_A(I/\mathfrak{m}Q) = \ell_A(I/Q) + \ell_A(Q/\mathfrak{m}Q) = r(A) + d.$$

Suppose that $Q = Q^\#$. Then A is a regular local ring which contains a regular system a_1, a_2, \dots, a_d of parameters such that $Q = (a_1, \dots, a_{d-1}, a_d^q)$ for some $q \geq 1$ ([G3, Theorem 3.1]). Hence $Q\mathfrak{m} : \mathfrak{m} = Q$, because $\mathfrak{m} = (a_1, a_2, \dots, a_d)$ and $Q\mathfrak{m} : \mathfrak{m} \subseteq (a_1, \dots, a_{d-1}, a_d^{q+1}) : a_d = Q$. Thus

$$\ell_A((0) :_{\mathfrak{m}/Q\mathfrak{m}} \mathfrak{m}) = \ell_A(Q/\mathfrak{m}Q) = d$$

as is claimed. \square

Let A be a Buchsbaum local ring with the Buchsbaum invariant $I(A)$. Then all the local cohomology modules $H_{\mathfrak{m}}^i(A)$ ($i \neq d$) are killed by the maximal ideal \mathfrak{m} and one has the equality

$$I(A) = \sum_{i=0}^{d-1} \binom{d-1}{i} h^i(A),$$

where $h^i(A) = \ell_A(H_{\mathfrak{m}}^i(A))$ ([SV, Chapter I, Proposition 2.6]). Let

$$r(A) = \sup_Q \ell_A((Q : \mathfrak{m})/Q)$$

where Q runs over parameter ideals in A and call it the Cohen-Macaulay type of A . We then have

$$(2.3) \quad r(A) = \sum_{i=0}^{d-1} \binom{d}{i} h^i(A) + \mu_{\hat{A}}(K_{\hat{A}})$$

([GSu, Theorem 2.5]), where $K_{\hat{A}}$ denotes the canonical module of the \mathfrak{m} -adic completion \hat{A} of A . Consequently $r(A) < \infty$.

THEOREM 2.4. *Let B be a Gorenstein local ring with $d = \dim B \geq 2$. Let A be a subring of B . Assume that B is a module-finite extension of A and $\ell_A(B/A) = 1$. Then*

- (1) *A is a Buchsbaum local ring with $\dim A = d$ and $I(A) = d - 1$.*
- (2) *The equality $I^2 = QI$ holds true for all parameter ideals Q in A , where $I = Q : \mathfrak{m}$.*

PROOF. Since B is a module-finite extension of A , by Eakin-Nagata's theorem our ring A is a Noetherian local ring with $d = \dim A$. Let \mathfrak{m} and \mathfrak{n} be the maximal ideals in A and B , respectively. We look at the exact sequence

$$(2.5) \quad 0 \rightarrow A \xrightarrow{l} B \rightarrow A/\mathfrak{m} \rightarrow 0$$

of A -modules, where ι denotes the inclusion map. Then applying functors $H_m^i(*)$ to (2.5), we get that

$$H_m^i(A) = (0) \quad (i \neq 1, d) \quad \text{and} \quad H_m^1(A) \cong A/\mathfrak{m},$$

because $\text{depth}_A B = d$. Hence A is a Buchsbaum ring with $I(A) = d - 1$ (cf. [SV, Chapter I, Proposition 2.12]). Notice that $\mathfrak{m}B = \mathfrak{m}$, because $\mathfrak{m} \cdot (B/A) = (0)$. Hence \mathfrak{m} is an ideal in B . On the other hand, we naturally have by (2.5) the exact sequence

$$0 \rightarrow A/\mathfrak{m} \rightarrow B/\mathfrak{m}B \rightarrow A/\mathfrak{m} \rightarrow 0,$$

whence $\mu_A(B) = 2$. Let us write $B = A + At$ with $t \in B$. Then $t \notin A$. We have $r(A) = d + 2$ by (2.3), since $K_A = B$. Notice that our local ring A is unmixed, because A is a Buchsbaum ring with $\text{depth}_A A > 0$ (cf. [SV, Chapter I, Lemma 1.13, Lemma 2.2]). Hence $e(A) \geq 2$. In fact, assume that $e(A) = 1$. Then A is a regular local ring so that we have $A = B$, because B is a module-finite birational extension of A . This is impossible, since $A \subsetneq B$ by our assumption.

Now let $Q = (a_1, a_2, \dots, a_d)$ be a parameter ideal in A and put $I = Q : \mathfrak{m}$. Then I is an ideal in B , since so is \mathfrak{m} . Thus $QB \subseteq I$. We need the following.

CLAIM 2.6. $\ell_A(QB/Q) = d$.

PROOF OF CLAIM 2.6. Since $B = A + At$, we have $QB = Q + \sum_{i=1}^d A \cdot a_i t$. Let $\overline{a_i t}$ denote the reduction of $a_i t \bmod Q$. Then $QB/Q = \sum_{i=1}^d k \cdot \overline{a_i t}$ ($k = A/\mathfrak{m}$). Let $\alpha_i \in A$ ($1 \leq i \leq d$) and assume that $\sum_{i=1}^d \alpha_i(a_i t) \in Q$. We write $\sum_{i=1}^d \alpha_i(a_i t) = \sum_{i=1}^d \beta_i a_i$ with $\beta_i \in A$. Then $\sum_{i=1}^d a_i(\alpha_i t - \beta_i) = 0$. Because a_1, a_2, \dots, a_d forms a B -regular sequence, $\alpha_i t - \beta_i \in (a_j \mid j \neq i)B \subseteq A$, so that $\alpha_i t \in A$. Hence $\alpha_i \in \mathfrak{m}$, because $t \notin A$. Thus the classes $\{\overline{a_i t}\}_{1 \leq i \leq d}$ form a k -basis of QB/Q . Hence $\ell_A(QB/Q) = d$. \square

If $\ell_A(I/Q) = r(A)$, then $I^2 = QI$ by [GSa, Theorem 3.9]. Therefore to prove $I^2 = QI$, we may assume that $\ell_A(I/Q) \leq d + 1$. Hence, either $\ell_A(I/Q) = d$, or $\ell_A(I/Q) = d + 1$ (cf. Claim 2.6). If $\ell_A(I/Q) = d$, then $I = QB$, so that $I^2 = QB \cdot IB = QI$. Assume that $\ell_A(I/Q) = d + 1$. Then $\ell_A(I/QB) = 1$. Therefore $\ell_B(I/QB) = 1$ and $\mathfrak{n}I \subseteq QB$. Hence $I = QB : \mathfrak{n}$, because $QB \subsetneq I \subseteq QB : \mathfrak{n}$ and B/QB is an Artinian Gorenstein local ring. Accordingly, $I^2 = QB \cdot IB = QI$, if $QB \neq (QB)^\#$ in B (cf., e.g., [GH, Proposition 3.4]). Suppose that $QB = (QB)^\#$ in B . Then, since $e(A) \geq 2$, by [GSa, Proposition 2.3] we have $I \subseteq Q^\#$. Hence $I \subseteq (QB)^\# = QB$ so that $I = QB$, which is impossible, because $\ell_A(I/QB) = 1$. Thus $QB \neq (QB)^\#$ in B and $I^2 = QI$, which completes the proof of Theorem 2.4. \square

The proof of the following result Proposition 2.7 is essentially the same as that of Theorem 2.4. Let us indicate a sketch only.

PROPOSITION 2.7. *Let B be a Gorenstein local ring with the maximal ideal \mathfrak{n} and $d = \dim B \geq 2$. Let A be a subring of B such that B is a finitely generated A -module. Assume that $A \subsetneq B$ and $\mathfrak{n} \subseteq A$. Then*

- (1) *A is a Buchsbaum local ring and $I(A) = (d - 1) \cdot \ell_A(B/A)$.*
- (2) *Let \mathfrak{m} be the maximal ideal in A . Then $\mathfrak{m} = \mathfrak{n}$, and the equality $I^2 = QI$ holds true for all parameter ideals Q in A , where $I = Q : \mathfrak{m}$.*

PROOF. Similarly as in the proof of Theorem 2.4, A is a Buchsbaum local ring and $\mathfrak{m} = \mathfrak{n}$. We also have $H_{\mathfrak{m}}^i(A) = (0)$ ($i \neq 1, d$) and $H_{\mathfrak{m}}^1(A) \cong B/A$. Let Q be a parameter ideal in A and put $I = Q : \mathfrak{m}$. Then $QB \subseteq I \subseteq QB : \mathfrak{n}$, since I is an ideal of B and $\mathfrak{m} = \mathfrak{n}$ in our case. Therefore, either $QB = I$, or $I = QB : \mathfrak{n}$, since B/QB is an Artinian Gorenstein local ring. We certainly have $I^2 = QI$ if $QB = I$. Assume that $I \neq QB$. Then $I = QB : \mathfrak{n}$ and $I^2 = QB \cdot IB = QI$, because $QB \neq (QB)^{\#}$ in B for the same reason as in the proof of Theorem 2.4. \square

Before closing this section let us note one example satisfying the hypothesis of Proposition 2.7.

EXAMPLE 2.8. Let K/k be a finite extension of fields and assume that $\delta = [K : k] \geq 2$. Let $n = \delta - 1$ and choose a k -basis $\{\theta_0 = 1, \theta_1, \dots, \theta_n\}$ of K . Let $d \geq 2$ be an integer and let $B = K[[X_1, X_2, \dots, X_d]]$ be the formal power series ring over K . Let

$$A = k[[\theta_i X_j \mid 0 \leq i \leq n, 1 \leq j \leq d]].$$

Then B is a module-finite extension of A such that the maximal ideal \mathfrak{n} of B coincides with that of A , $\ell_A(B/A) = n$, and $e(A) = \delta$. Hence by Proposition 2.7 A is a Buchsbaum local ring, in which the equality $I^2 = QI$ holds true for all parameter ideals Q , where $I = Q : \mathfrak{m}$.

PROOF. Let \mathfrak{m} be the maximal ideal in A , that is $\mathfrak{m} = (\theta_i X_j \mid 0 \leq i \leq n, 1 \leq j \leq d)A$. Then $\mathfrak{m}B = \mathfrak{n}$ whence $B = \sum_{i=0}^n A\theta_i$. Let $b \in B$ and write $b = \sum_{i=0}^n a_i \theta_i$ with $a_i \in A$. Then, since $bX_j = \sum_{i=0}^n a_i(\theta_i X_j) \in \mathfrak{m}$ for all $1 \leq j \leq d$, we get $\mathfrak{n} \subseteq \mathfrak{m}$. Hence $\mathfrak{m} = \mathfrak{n}$. We have

$$\mu_A(B) = \ell_A(B/\mathfrak{m}B) = \ell_A(B/\mathfrak{n}) = [K : k] = \delta \geq 2.$$

Because $\ell_A(B/A) = \ell_A(B/\mathfrak{n}) - \ell_A(A/\mathfrak{m})$, we get $\ell_A(B/A) = \delta - 1 = n$. Let $\mathfrak{q} = (X_1, X_2, \dots, X_d)A$. Then since $\mathfrak{m}^2 = \mathfrak{q}\mathfrak{m}$, the ideal \mathfrak{q} is a minimal reduction of \mathfrak{m} , so that we have

$$e(A) = e_{\mathfrak{q}}^0(A) = e_{\mathfrak{q}}^0(B) = \ell_A(B/\mathfrak{q}B) = \ell_A(B/\mathfrak{n}) = \delta,$$

as is claimed. \square

3. Proof of Theorem 1.1.

Let A be a Noetherian local ring with $d = \dim A \geq 2$. Suppose that A is a reduced ring with $\#\text{Ass } A = 2$, say $\text{Ass } A = \{\mathfrak{p}_1, \mathfrak{p}_2\}$. We furthermore assume that A/\mathfrak{p}_i is a regular local ring with $\dim A/\mathfrak{p}_i = d$ ($i = 1, 2$) and $\mathfrak{p}_1 + \mathfrak{p}_2 = \mathfrak{m}$. Hence $\mathfrak{m} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$, because $\mathfrak{p}_1 \cap \mathfrak{p}_2 = (0)$. With this notation and assumption we have the following.

THEOREM 3.1. $I^2 = QI$ for all parameter ideals Q in A , where $I = Q : \mathfrak{m}$.

PROOF. We look at the exact sequences

$$(3.2) \quad 0 \rightarrow A \xrightarrow{I} A/\mathfrak{p}_1 \oplus A/\mathfrak{p}_2 \rightarrow A/\mathfrak{m} \rightarrow 0 \quad \text{and}$$

$$(3.3) \quad 0 \rightarrow \mathfrak{p}_i \rightarrow A \rightarrow A/\mathfrak{p}_i \rightarrow 0$$

of A -modules ($i = 1, 2$), where $\iota(a) = (a \bmod \mathfrak{p}_1, a \bmod \mathfrak{p}_2)$ for all $a \in A$. Then, applying functors $H_m^i(*)$ to (3.2), we get $H_m^i(A) = (0)$ ($i \neq 1, d$) and $H_m^1(A) \cong A/\mathfrak{m}$. Hence A is a Buchsbaum local ring with $I(A) = d - 1$ and $e(A) = 2$. We have $r(A) = d + 2$ by (2.3), because $K_A = A/\mathfrak{p}_1 \oplus A/\mathfrak{p}_2$. Let $Q = (a_1, a_2, \dots, a_d)$ be a parameter ideal in A and put $I = Q : \mathfrak{m}$. Then $\ell_A(I/Q) \leq r(A) = d + 2$. We may assume that $\ell_A(I/Q) \leq d + 1$ (cf. [GSa, Theorem 3.9]). Let $A_i = A/\mathfrak{p}_i$ and $\mathfrak{m}_i = \mathfrak{m}/\mathfrak{p}_i$ ($i = 1, 2$). We write $a_j = \ell_j + m_j$ ($1 \leq j \leq d$) with $\ell_j \in \mathfrak{p}_1$ and $m_j \in \mathfrak{p}_2$.

Firstly we consider the case $QA_2 \neq (QA_2)^\#$ in A_2 . Let $\varepsilon : A \rightarrow A_2$ be the canonical epimorphism. Then $\varepsilon(\mathfrak{p}_1) = \mathfrak{m}_2$ and $\mathfrak{p}_1 \cong \mathfrak{m}_2$ via ε , because $\mathfrak{m} = \mathfrak{p}_1 \oplus \mathfrak{p}_2$. Hence

$$(3.4) \quad \ell_A((0) :_{\mathfrak{p}_1/Q\mathfrak{p}_1} \mathfrak{m}) = \ell_A((0) :_{\mathfrak{m}_2/Q\mathfrak{m}_2} \mathfrak{m}) = \ell_{A_2}((0) :_{\mathfrak{m}_2/Q\mathfrak{m}_2} \mathfrak{m}_2) = d + 1$$

by Proposition 2.2. Let $\bar{*}$ denote the reduction mod \mathfrak{p}_2 . Then since A_2/QA_2 is an Artinian Gorenstein local ring and since $Q\mathfrak{m}_2 : \mathfrak{m}_2 = QA_2 : \mathfrak{m}_2 \subseteq \mathfrak{m}_2$ (cf. Proof of Proposition 2.2), the ideal $Q\mathfrak{m}_2 : \mathfrak{m}_2$ of A_2 is generated by $\{\bar{\ell}_j\}_{1 \leq j \leq d}$ together with one more element, say $\bar{\eta}$ ($\eta \in \mathfrak{p}_1$). Hence the A/\mathfrak{m} -space $(0) :_{\mathfrak{p}_1/Q\mathfrak{p}_1} \mathfrak{m}$ is spanned by $\{\ell_j \bmod Q\mathfrak{p}_1\}_{1 \leq j \leq d}$ and $\eta \bmod Q\mathfrak{p}_1$, because $\mathfrak{p}_1 \cong \mathfrak{m}_2$ via ε . Now look at the exact sequence

$$(3.5) \quad 0 \rightarrow \mathfrak{p}_i/Q\mathfrak{p}_i \rightarrow A/Q \xrightarrow{\varphi_i} A_i/QA_i \rightarrow 0 \quad (i = 1, 2)$$

of A -modules induced from (3.3) (notice that a_1, a_2, \dots, a_d is an A_i -regular sequence). Then, considering the socles of the terms in (3.5) with $i = 1$, we get

$$I = Q + (\ell_1, \ell_2, \dots, \ell_d) + (\eta),$$

since $\ell_A((0) :_{\mathfrak{p}_1/Q\mathfrak{p}_1} \mathfrak{m}) = d + 1$ by (3.4) and since $\ell_A(I/Q) \leq d + 1$, as is assumed at the beginning of this proof. Hence $I^2 = QI + (\eta^2)$, because

$$\ell_i \ell_j = (\ell_i + m_i) \ell_j = a_i \ell_j \quad \text{and} \quad \ell_i \eta = (\ell_i + m_i) \eta = a_i \eta$$

for all $1 \leq i, j \leq d$ (recall that $\mathfrak{p}_1 \cap \mathfrak{p}_2 = (0)$). On the other hand, since $QA_2 \neq (QA_2)^\#$, we get $(QA_2 : \mathfrak{m}_2)^2 = QA_2 \cdot (QA_2 : \mathfrak{m}_2)$ (cf., e.g., [GH, Proposition 3.4]). Hence $\bar{\eta}^2 \in (\bar{\ell}_1, \bar{\ell}_2, \dots, \bar{\ell}_d) \cdot [(\bar{\ell}_1, \bar{\ell}_2, \dots, \bar{\ell}_d) + (\bar{\eta})]$ so that

$$(3.6) \quad \eta^2 \in (\ell_1, \ell_2, \dots, \ell_d) \cdot [(\ell_1, \ell_2, \dots, \ell_d) + (\eta)] + \mathfrak{p}_2.$$

Because $\mathfrak{p}_1 \cap \mathfrak{p}_2 = (0)$ and $(\ell_1, \ell_2, \dots, \ell_d) + (\eta) \subseteq \mathfrak{p}_1$, by (3.6) we readily get that

$$\eta^2 \in (\ell_1, \ell_2, \dots, \ell_d) \cdot [(\ell_1, \ell_2, \dots, \ell_d) + (\eta)] \subseteq QI.$$

Hence $I^2 = QI$, since $I^2 = QI + (\eta^2)$. We get, by the symmetry, that $I^2 = QI$ also in the case where $QA_1 \neq (QA_1)^\#$.

We now consider the case $QA_i = (QA_i)^\#$ for $i = 1, 2$. Then thanks to the exact sequence (3.5) with $i = 1$, we have $\ell_A(I/Q) \geq d$, because

$$\ell_A((0) :_{\mathfrak{p}_1/Q\mathfrak{p}_1} \mathfrak{m}) = \ell_{A_2}((0) :_{\mathfrak{m}_2/Q\mathfrak{m}_2} \mathfrak{m}_2) = d$$

by Proposition 2.2. We actually have the following.

CLAIM 3.7. $\ell_A(I/Q) = d$.

PROOF OF CLAIM 3.7. Suppose that $\ell_A(I/Q) \neq d$. Then $\ell_A(I/Q) = d + 1$. Choose

a regular system $\overline{c_1}, \overline{c_2}, \dots, \overline{c_d}$ ($c_i \in \mathfrak{p}_1$) of parameters for A_2 so that $QA_2 = (\overline{c_1}, \dots, \overline{c_{d-1}}, \overline{c_d^q})$ for some $q > 0$ (cf. [G3, Theorem 3.1]). Hence

$$QA_2 : \mathfrak{m}_2 = (\overline{c_1}, \dots, \overline{c_{d-1}}, \overline{c_d^{q-1}}) = QA_2 + (\overline{c_d^{q-1}}).$$

We have

$$(3.8) \quad (\ell_1, \ell_2, \dots, \ell_d) = (c_1, \dots, c_{d-1}, c_d^q),$$

because $QA_2 = (\overline{\ell_1}, \overline{\ell_2}, \dots, \overline{\ell_d}) = (\overline{c_1}, \dots, \overline{c_{d-1}}, \overline{c_d^q})$ and $\mathfrak{p}_1 \cong \mathfrak{m}_2$ via ε . Notice that

$$\ell_A((0) :_{\mathfrak{p}_2/Q\mathfrak{p}_2} \mathfrak{m}) = \ell_{A_1}((0) :_{\mathfrak{m}_1/Q\mathfrak{m}_1} \mathfrak{m}_1) = d$$

and $(0) :_{\mathfrak{m}_1/Q\mathfrak{m}_1} \mathfrak{m}_1 = QA_1/Q\mathfrak{m}_1$ (cf. Proposition 2.2 and its proof). Hence the A/\mathfrak{m} -space $(0) :_{\mathfrak{p}_2/Q\mathfrak{p}_2} \mathfrak{m}$ is spanned by $\{m_j \bmod Q\mathfrak{p}_2\}_{1 \leq j \leq d}$. We now look at the exact sequence (3.5) with $i = 2$. Then since $\ell_A((0) :_{\mathfrak{p}_2/Q\mathfrak{p}_2} \mathfrak{m}) = d$ and $\ell_A(I/Q) = d + 1$, the canonical epimorphism $\varphi_2 : A/Q \rightarrow A_2/QA_2$ cannot be zero on the socles, so that we have $IA_2 = QA_2 : \mathfrak{m}_2$. Hence $IA_2 = QA_2 + (\overline{c_d^{q-1}})$. Choose $\eta \in I$ so that $\overline{\eta} = \overline{c_d^{q-1}}$ and write $\eta = c_d^{q-1} + \delta + \rho$ with $\delta \in Q$ and $\rho \in \mathfrak{p}_2$. Then thanks to the exact sequence (3.5) with $i = 2$, we get

$$\begin{aligned} I &= Q + (m_1, m_2, \dots, m_d) + (\eta) \\ &= Q + (m_1, m_2, \dots, m_d) + (c_d^{q-1} + \delta + \rho), \end{aligned}$$

because $(0) :_{\mathfrak{p}_2/Q\mathfrak{p}_2} \mathfrak{m}$ is spanned by $\{m_j \bmod Q\mathfrak{p}_2\}_{1 \leq j \leq d}$. Consequently we have

$$\begin{aligned} I &= Q + (m_1, m_2, \dots, m_d) + (c_d^{q-1} + \rho) && (\text{since } \delta \in Q) \\ &= (\ell_1, \ell_2, \dots, \ell_d) + (m_1, m_2, \dots, m_d) + (c_d^{q-1} + \rho) \\ &= (c_1, \dots, c_{d-1}, c_d^q) + (m_1, m_2, \dots, m_d) + (c_d^{q-1} + \rho) && (\text{by (3.8)}) \\ &= (c_1, \dots, c_{d-1}) + (m_1, m_2, \dots, m_d) + (c_d^{q-1} + \rho) && (\text{since } c_d^q = c_d(c_d^{q-1} + \rho)), \end{aligned}$$

which is impossible, because $\mu_A(I) = \ell_A(I/Q) + d = 2d + 1$ by Lemma 2.1. Thus $\ell_A(I/Q) = d$. \square

Therefore, in the exact sequence (3.5) with $i = 2$, the socle I/Q of A/Q coincides with the image of $(0) :_{\mathfrak{p}_2/Q\mathfrak{p}_2} \mathfrak{m}$, because $\ell_A((0) :_{\mathfrak{p}_2/Q\mathfrak{p}_2} \mathfrak{m}) = d$; that is

$$I = Q + (m_1, m_2, \dots, m_d).$$

Thus $I^2 = QI$, because $m_i m_j = (\ell_i + m_i)m_j \in QI$ for all $1 \leq i, j \leq d$. This completes the proof of Theorem 3.1. \square

We are now ready to prove Theorem 1.1. Let A be a Buchsbaum local ring with $e(A) = 2$ and $\text{depth } A > 0$.

PROOF OF THEOREM 1.1. To prove Theorem 1.1, passing to the ring $A[X]_{\mathfrak{m}_A[X]}$ where X is an indeterminate over A and then passing to the completion, we may assume that A is a complete local ring with the infinite residue class field. Thanks to [CP,

Theorem 2.2], we may assume that A is not a Cohen-Macaulay ring. Hence $d \geq 2$ and so by [G2, Theorem 1.1]

$$H_m^i(A) = (0) \quad (i \neq 1, d) \quad \text{and} \quad H_m^1(A) \cong A/\mathfrak{m}.$$

Let B be the Cohen-Macaulayfication of A , that is the intermediate ring $A \subseteq B \subseteq Q(A)$ where $Q(A)$ denotes the total quotient ring of A , such that B is a module-finite extension of A , $\text{depth}_A B = d$, and $\mathfrak{m}B = \mathfrak{m}$ (cf. [G1, Theorem 1.1]). Then $H_m^1(A) \cong B/A$ and hence $\ell_A(B/A) = 1$. Now there are two cases. One is the case where B is a local ring. The other one is the case where B is not a local ring. Firstly, suppose that B is a local ring and choose a minimal reduction \mathfrak{q} of \mathfrak{m} . Then $\ell_A(B/\mathfrak{q}B) = 2$, because

$$\ell_A(B/\mathfrak{q}B) = e_{\mathfrak{q}}^0(B) = e_{\mathfrak{q}}^0(A) = e(A).$$

Consequently $[B/\mathfrak{n} : A/\mathfrak{m}] \cdot \ell_B(B/\mathfrak{q}B) = 2$. If $A/\mathfrak{m} \neq B/\mathfrak{n}$, then $\ell_B(B/\mathfrak{q}B) = 1$ so that $\mathfrak{n} = \mathfrak{q}B$. Hence B is a regular local ring with $\mathfrak{n} = \mathfrak{m}$ and the assertion follows from Proposition 2.7. If $A/\mathfrak{m} = B/\mathfrak{n}$, then $\ell_B(B/\mathfrak{q}B) = 2$. Hence $\ell_B(\mathfrak{n}/\mathfrak{q}B) = 1$ and so B is a Gorenstein ring, because the ring $B/\mathfrak{q}B$ is Artinian and Gorenstein. The assertion now follows from Theorem 2.3, since $\ell_A(B/A) = 1$.

Assume that B is not a local ring. Then the proof of [G2, Proposition 4.2] still works in our context to show that A is a reduced ring with $\#\text{Ass } A = 2$, say $\text{Ass } A = \{\mathfrak{p}_1, \mathfrak{p}_2\}$, such that $\mathfrak{p}_1 + \mathfrak{p}_2 = \mathfrak{m}$, and A/\mathfrak{p}_i is a regular local ring with $d = \dim A/\mathfrak{p}_i$ for $i = 1, 2$. Hence the equality $I^2 = QI$ follows from Theorem 3.1. This completes the proof of Theorem 1.1. \square

4. Examples.

In this section we shall construct examples, showing that the equality $I^2 = QI$ fails in general to hold, even though A is a Buchsbaum local ring with sufficiently large depth and multiplicity.

Let k be a field and let $1 \leq d < m$ be integers. Let $S = k[X_1, \dots, X_m, V, A_1, \dots, A_d]$ be the polynomial ring with $m + d + 1$ variables over k and put

$$\mathfrak{a} = (X_i \mid 1 \leq i \leq m-1)^2 + (X_m^2) + (X_i V \mid 1 \leq i \leq m) + \left(V^2 - \sum_{i=1}^d A_i X_i \right).$$

We regard S as a \mathbf{Z} -graded ring such that $S_0 = k$ and $X_i, V, A_j \in S_1$ for all $1 \leq i \leq m$ and $1 \leq j \leq d$. We put

$$R = S/\mathfrak{a}, M = R_+, B = R_M, \quad \text{and} \quad \mathfrak{m} = MB.$$

Then $\dim R = d$, because $\sqrt{\mathfrak{a}} = (X_i \mid 1 \leq i \leq m) + (V)$. Let x_i, v , and a_j denote respectively the reduction of X_i, V , and $A_j \bmod \mathfrak{a}$. We put $Q = (a_1, a_2, \dots, a_d)$ and $\mathfrak{p} = (x_1, x_2, \dots, x_m) + (v)$ in R . Hence $M = Q + \mathfrak{p}$ and $M^3 = QM^2$, since $\mathfrak{p}^3 = (0)$. Consequently, $\mathfrak{q} = QB$ is a minimal reduction of the maximal ideal \mathfrak{m} in B .

Let us begin with the following.

LEMMA 4.1. $\ell_B(B/\mathfrak{q}) = 2m + 1$, $e_{\mathfrak{m}}^0(B) = 2m$, and B is not a Cohen-Macaulay ring.

PROOF. Since $R/Q \cong k[X_1, X_2, \dots, X_m, V]/\mathfrak{b}$ where

$$\mathfrak{b} = (X_i \mid 1 \leq i \leq m-1)^2 + (X_m^2, V^2) + (X_i V \mid 1 \leq i \leq m),$$

we have $\dim_k R/Q = 2m+1$ and

$$Q : M = Q + (x_i x_m \mid 1 \leq i \leq m-1) + (v).$$

Hence $\ell_B(B/\mathfrak{q}) = 2m+1$. Let $P = (X_i \mid 1 \leq i \leq m) + (V)$ in the ring S ; hence $\mathfrak{p} = P/\mathfrak{a}$. Then $\text{Min } B = \{\mathfrak{p}B\}$ and $B/\mathfrak{p}B$ is a regular local ring, so that we have $e(B) = \ell_{B/\mathfrak{p}B}(B/\mathfrak{p}B)$. Let $\tilde{S} = S[1/A_1]$ and $\tilde{k} = k[A_1, 1/A_1]$. Then

$$\tilde{S} = \tilde{k}[X'_1, \dots, X'_m, V', A'_2, \dots, A'_d],$$

where $X'_i = X_i/A_1$, $V' = V/A_1$, and $A'_j = A_j/A_1$ ($1 \leq i \leq m, 1 \leq j \leq d$). The elements $\{X'_i\}_{1 \leq i \leq m}$, V' , and $\{A'_j\}_{2 \leq j \leq d}$ are algebraically independent over \tilde{k} . We have

$$\mathfrak{a}\tilde{S} = (X'_i \mid 1 \leq i \leq m-1)^2 + (X_m'^2) + (X'_i V' \mid 1 \leq i \leq m) + \left(V'^2 - \sum_{i=1}^d A'_i X'_i \right)$$

and $P\tilde{S} = (X'_i \mid 1 \leq i \leq m) + (V')$. Hence $X'_1 - (V'^2 - \sum_{i=2}^d A'_i X'_i) \in \mathfrak{a}\tilde{S}$. Let $T = \tilde{k}[X'_2, \dots, X'_m, V', A'_2, \dots, A'_d]$ and we identify $T = \tilde{S}/(X'_1 - (V'^2 - \sum_{i=2}^d A'_i X'_i))$. Then

$$\begin{aligned} \mathfrak{a}T &= \left(V'^2 - \sum_{i=2}^d A'_i X'_i, \{X'_i\}_{2 \leq i \leq m-1} \right)^2 + (X_m'^2) + \left(\left(V'^2 - \sum_{i=2}^d A'_i X'_i \right) \cdot V' \right) \\ &\quad + (X'_i V' \mid 2 \leq i \leq m) \\ &= (V'^2, \{X'_i\}_{2 \leq i \leq m-1})^2 + (X_m'^2) + (V'^3) + (X'_i V' \mid 2 \leq i \leq m) \quad (\text{since } d < m) \\ &= (X'_i \mid 2 \leq i \leq m-1)^2 + (X_m'^2) + (V'^3) + (X'_i V' \mid 2 \leq i \leq m), \end{aligned}$$

and $PT = (X'_i \mid 2 \leq i \leq m) + (V')$. Therefore $\ell_{B/\mathfrak{p}B}(B/\mathfrak{p}B) = \ell_{S_P}(S_P/\mathfrak{a}S_P) = \ell_U(U)$, where

$$U = k(A'_1, A'_2, \dots, A'_d)[X'_2, \dots, X'_m, V']/\mathfrak{b}, \quad \text{and}$$

$$\mathfrak{b} = (X'_i \mid 2 \leq i \leq m-1)^2 + (X_m'^2) + (V'^3) + (X'_i V' \mid 2 \leq i \leq m).$$

Consequently, $e(B) = \ell_U(U) = 2m < \ell_B(B/\mathfrak{q}) = 2m+1$, whence B is not a Cohen-Macaulay ring. \square

Let \underline{a}^2 denote the sequence $a_1^2, a_2^2, \dots, a_d^2$ and let $e_{(\underline{a}^2)B}^0(B)$ denote the multiplicity of B with respect to the parameter ideal $(\underline{a}^2)B = (a_1^2, a_2^2, \dots, a_d^2)B$. We then have the following.

PROPOSITION 4.2. $\ell_B(B/(\underline{a}^2)B) - e_{(\underline{a}^2)B}^0(B) = 1$.

PROOF. Since B is not a Cohen-Macaulay ring, $\ell_B(B/(\underline{a}^2)B) - e_{(\underline{a}^2)B}^0(B) > 0$. It suffices to show $\ell_B(B/(\underline{a}^2)B) - e_{(\underline{a}^2)B}^0(B) \leq 1$. Let

$$\mathfrak{c} = (A_i^2 \mid 1 \leq i \leq d) + (X_i^2 \mid 1 \leq i \leq m) + \left(V^2 - \sum_{i=1}^d A_i X_i \right)$$

and put $C = S/\mathfrak{c}$ and $D = S/(\mathfrak{a} + (A_i^2 \mid 1 \leq i \leq d))$. Then D is a homomorphic image

of C . The ring C is a complete intersection with $\dim_k C = 2^{d+m+1}$. Let x_i, v , and a_j denote, for the moment, the reduction of X_i, V , and $A_j \bmod \mathfrak{a} + (A_i^2 \mid 1 \leq i \leq d)$. Let $\Lambda = \{1, 2, \dots, d\}$ and $\Gamma = \{1, 2, \dots, m\}$. For given subsets $I \subseteq \Lambda$ and $J \subseteq \Gamma$ we put

$$a_I = \prod_{i \in I} a_i \quad \text{and} \quad x_J = \prod_{j \in J} x_j.$$

Then the elements $\{a_I x_J\}_{I \subseteq \Lambda, J \subseteq \Gamma}$ and $\{a_I x_J v\}_{I \subseteq \Lambda, J \subseteq \Gamma}$ span the k -space D , because their preimages in C form a k -basis of C . Notice that $a_I x_J v = 0$ if $J \neq \emptyset$ and that $J \subseteq \{i, m\}$ for some $1 \leq i \leq m-1$ if $x_J \neq 0$. Hence the k -space D is actually spanned by the following $2^d(2m+1)$ elements

$$(4.3) \quad a_I, x_i a_I, x_m a_I, x_i x_m a_I, \text{ and } a_I v \quad \text{with } I \subseteq \Lambda, 1 \leq i \leq m-1.$$

Let $1 \leq i \leq d$ and $K \subseteq \Lambda$. Assume that $i \notin K$ but $\{1, \dots, i-1\} \subseteq K$. Then since $(\sum_{i=1}^d a_i x_i)(x_m a_K) = v^2 x_m a_K = 0$, we have

$$(4.4) \quad \sum_{1 \leq j < i} (a_j x_j)(x_m a_K) + (a_i x_i)(x_m a_K) + \sum_{i < j \leq d} (a_j x_j)(x_m a_K) = 0.$$

Notice that $\sum_{1 \leq j < i} (a_j x_j)(x_m a_K) = 0$, since $(a_\ell x_\ell)(x_m a_K) = (a_\ell a_K)(x_\ell x_m) = 0$ for all $\ell \in K$. If $i < j \leq d$ and $j \notin K$, then $(a_j x_j)(x_m a_K) = (x_j x_m) a_{K \cup \{j\}}$. Consequently by (4.4) we have the following expression

$$x_i x_m a_{K \cup \{i\}} = (a_i x_i)(x_m a_K) = - \sum_{i < j \leq d, j \notin K} (x_j x_m) a_{K \cup \{j\}}$$

of $x_i x_m a_{K \cup \{i\}}$. Hence, letting $K = I \setminus \{i\}$, it follows from this expression that for all $1 \leq i \leq d$, the set $\{x_i x_m a_I\}_{\{1, \dots, i\} \subseteq I \subseteq \Lambda}$ is contained in the k -subspace of D spanned by $\{x_j x_m a_J\}_{i < j \leq d, J \subseteq \Lambda}$. Therefore, in order to span the whole k -space D , for each $1 \leq i \leq d$ the elements $\{x_i x_m a_I \mid \{1, 2, \dots, i\} \subseteq I \subseteq \Lambda\}$ can be deleted from the system of generators given by (4.3), so that we have

$$\begin{aligned} \ell_B(B/(\underline{a}^2)B) &= \dim_k D \leq 2^d(2m+1) - \sum_{i=1}^d 2^{d-i} \\ &= 2^d(2m+1) - (2^d - 1) \\ &= 2^{d+1}m + 1 \\ &= 2^d e_{(\underline{a})B}^0(B) + 1 \\ &= e_{(\underline{a}^2)B}^0(B) + 1 \end{aligned}$$

as is claimed. □

By Proposition 4.2 we get

$$\ell_B(B/(\underline{a}^2)B) - e_{(\underline{a}^2)B}^0(B) = \ell_B(B/\mathfrak{q}) - e_{\mathfrak{q}}^0(B) = 1.$$

Hence the local cohomology modules $H_m^i(B)$ ($i \neq d$) are finitely generated B -modules

and $\sum_{i=0}^{d-1} \binom{d-1}{i} h^i(B) = 1$ (cf. [SV, Appendix, Theorem and Definition 17]). Accordingly, either $\text{depth } B = 0$, or $\text{depth } B = d - 1$. We have that $h^0(B) = 1$ and $h^i(B) = 0$ ($1 \leq i \leq d - 1$) if $\text{depth } B = 0$, and that $h^{d-1}(B) = 1$ if $\text{depth } B = d - 1$. In any case, $H_m^i(B) = (0)$ for all $i \neq t, d$, and $H_m^t(B) \cong B/\mathfrak{m}$, where $t = \text{depth } B$. Thus B is a Buchsbaum ring (cf. [SV, Chapter I, Proposition 2.6]). We actually have the following.

THEOREM 4.5. $H_M^{d-1}(R) \cong (R/M)(d - 3)$.

PROOF. (1) ($d = 1$). Use the fact that $0 \neq x_1 x_m \in (0) : M$.

(2) ($d = 2$). Assume that $\text{depth } B = 0$. Then applying functors $H_M^i(*)$ to the exact sequence

$$(4.6) \quad 0 \rightarrow H_M^0(R)(-1) \rightarrow R(-1) \xrightarrow{a_2} R \rightarrow R/a_2 R \rightarrow 0,$$

we get a natural isomorphism $H_M^0(R) \cong H_M^0(R/a_2 R)$. We apply the result of the case where $d = 1$ to the ring $R/a_2 R$ and choose $0 \neq \varphi \in R_2$ such that $M\varphi = (0)$ and $\varphi \equiv x_1 x_m \pmod{a_2 R}$. Let $\varphi = x_1 x_m + a_2 \psi$ with $\psi \in R_1$. Then $x_1 x_m a_2 + a_2^2 \psi = 0$, that is $X_1 X_m A_2 + A_2^2 \xi \in \mathfrak{a}$ for some $\xi \in S_1$, which is impossible. Hence $\text{depth } B = 1$ and by (4.6) we get an isomorphism $H_M^0(R/a_2 R) \cong H_M^1(R)(-1)$. Thus $H_M^1(R) \cong (R/M)(-1)$, because $H_M^0(R/a_2 R) \cong (R/M)(-2)$.

(3) ($d \geq 3$). We may assume that our assertion holds true for $d - 1$. Then $\text{depth } B = d - 1$, because

$$h^0(B/a_d B) = h^0(B) + h^1(B)$$

and $h^0(B/a_d B) = 0$ by the hypothesis on d . Hence a_d is a non-zerodivisor of R , so that we have

$$H_M^{d-2}(R/a_d R) \cong H_M^{d-1}(R)(-1).$$

Thus $H_M^{d-1}(R) \cong (R/M)(d - 3)$, because $H_M^{d-2}(R/a_d R) \cong (R/M)(d - 4)$. \square

Let $J = Q : M$ and $I = \mathfrak{q} : \mathfrak{m} (= JB)$. We then have the following.

PROPOSITION 4.7. $I^2 \neq \mathfrak{q}I$ but $I^3 = \mathfrak{q}I^2$.

PROOF. We have $J = Q + (x_i x_m \mid 1 \leq i \leq m - 1) + (v)$ (cf. Proof of Lemma 4.1). Assume that $J^2 = QJ$. Then $v^2 \in QJ$ whence

$$(4.8) \quad V^2 \in (A_i \mid 1 \leq i \leq d) \cdot [(A_i \mid 1 \leq i \leq d) + (X_i X_m \mid 1 \leq i \leq m - 1) + (V)] + \mathfrak{a}.$$

We now substitute $X_i = 0$ and $A_j = 0$ for all $2 \leq i \leq m$ and $2 \leq j \leq d$. Then by (4.8) we get

$$V^2 \in (A_1^2, A_1 V, X_1^2, X_1 V, V^2 - A_1 X_1)$$

in the polynomial ring $k[X_1, V, A_1]$, which is impossible. Hence $v^2 \notin QJ$ so that we have $J^2 \neq QJ$. We get $J^3 = QJ^2$, because $J^2 = QJ + (v^2)$ and $v^3 = 0$. \square

Therefore, for given integers $1 \leq d < m$, there exists a Buchsbaum local ring A with $\dim A = d$, $\text{depth } A = d - 1$, and $e(A) = 2m$, such that A contains a parameter ideal Q

which is a minimal reduction of \mathfrak{m} and $I^2 \neq QI$ but $I^3 = QI^2$, where $I = Q : \mathfrak{m}$. Thus the equality $I^2 = QI$ fails in general to hold, even though A is a Buchsbaum local ring with sufficiently large depth and multiplicity.

References

- [CHV] A. Corso, C. Huneke and W. V. Vasconcelos, On the integral closure of ideals, *Manuscripta Math.*, **95** (1998), 331–347.
- [CP] A. Corso and C. Polini, Links of prime ideals and their Rees algebras, *J. Algebra*, **178** (1995), 224–238.
- [CPV] A. Corso, C. Polini and W. V. Vasconcelos, Links of prime ideals, *Math. Proc. Cambridge Philos. Soc.*, **115** (1994), 431–436.
- [G1] S. Goto, On the Cohen-Macaulayfication of certain Buchsbaum rings, *Nagoya Math. J.*, **80** (1980), 107–116.
- [G2] S. Goto, Buchsbaum rings with multiplicity 2, *J. Algebra*, **74** (1982), 494–508.
- [G3] S. Goto, Integral closedness of complete-intersection ideals, *J. Algebra*, **108** (1987), 151–160.
- [GH] S. Goto and F. Hayasaka, Finite homological dimension and primes associated to integrally closed ideals II, *J. Math. Kyoto Univ.*, **42** (2002), No. 4, 631–639.
- [GN] S. Goto and K. Nishida, Hilbert coefficients and Buchsbaumness of associated graded rings, *J. Pure Appl. Algebra*, **181** (2003), No. 1, 61–74.
- [GSa] S. Goto and H. Sakurai, The equality $I^2 = QI$ in Buchsbaum rings, *Rendiconti del Seminario Matematico dell'Universit di Padova*, **110** (2003), 25–56.
- [GSu] S. Goto and N. Suzuki, Index of reducibility of parameter ideals in a local ring, *J. Algebra*, **87** (1984), 53–88.
- [SV] J. Stückrad and W. Vogel, *Buchsbaum rings and applications*, Springer-Verlag, Berlin-New York-Tokyo, 1986.
- [Y1] K. Yamagishi, The associated graded modules of Buchsbaum modules with respect to \mathfrak{m} -primary ideals in the equi- I -invariant case, *J. Algebra*, **225** (2000), 1–27.
- [Y2] K. Yamagishi, Buchsbaumness in Rees modules associated to ideals of minimal multiplicity in the equi- I -invariant case, *J. Algebra*, **251** (2002), 213–255.

Shiro GOTO

Department of Mathematics
School of Science and Technology
Meiji University
214-8571 Japan
E-mail: goto@math.meiji.ac.jp

Hideto SAKURAI

Department of Mathematics
School of Science and Technology
Meiji University
214-8571 Japan
E-mail: ee78052@math.meiji.ac.jp