

A CHARACTERIZATION OF THE MAXIMAL IDEAL

IN A FACTOR OF THE CASE (II_∞)

By Haruo SUNOUCHI

(Comm. by T. Kawata)

It is well-known ([5; Theorem VIII] and [3; Theorem 2]) that any factor on a Hilbert space can be classified into the cases (I_n) , (I_∞) , (II_1) , (II_∞) and (III) . Von Neumann [6] proved that a factor is simple, that is, contains no (two sided) ideal, if and only if it belongs to one of the cases (I_n) , (II_1) and (III) . Moreover, Calkin [1] showed that the factor of case (I_∞) contains only one maximal ideal, and which consists of all completely continuous operators. Recently, Misonou [4; Theorem 3] proved that any rings of operators is weakly central, that is, there is a one-to-one correspondence between the maximal ideals in the ring and the maximal ideals in its center. As a corollary of his theorem, we know that there exists only one maximal ideal in a factor of case (II_∞) . In this note we shall characterize this maximal ideal. For simplicity, we consider only in a separable Hilbert space, but the proof remains true in the non-separable case by the use of the corresponding results due to Dixmier [2].

Theorem. Each factor M of case (II_∞) has a unique maximal ideal, and this is characterized by all those operators, $A \in M$ of which $[R(A)]$ are finite. Here, by $[R(A)]$ we shall denote the closure of the range of A .

Proof proceeds by the Calkin's principle and the use of the results on the rings of operators ([5] and [2]).

Let J be the set of all $A \in M$ such that $[R(A)]$ is finite, then it is well known that J is an ideal in M by [6, p. 25]. But for completeness, we shall trace his proof. Let A, B be in J , then $A+B \in J$ follows from the facts that $[R(A+B)] \subseteq [R(A)] \vee [R(B)]$ and the right hand side is also finite ([5; Lemma 7.3.5]). Since $[R(A)] \supseteq [R(AX)]$, $A \in J$ implies $AX \in J$ for any $X \in M$. Finally, by $[R(A)] \sim [R(A^*)]$ ([5; Lemma 6.2.1]), we see that $A \in J$

implies $A^* \in J$. Hence $XA = (A^*X^*)^* \in J$, if $A \in J$. Now $[R(I)] = H$ is infinite, whence $I \notin J$, and there exists a non-zero projection $E \in M$ such that $[R(E)]$ is finite, so that $E \in J$. Consequently, we see that J is a non-trivial ideal in M . That J is unique, maximal and uniformly closed, will follow from the next paragraph.

Let K be any non-trivial ideal in M . Suppose that there exists a partially isometric operator $X \in M$ which gives the equivalency $H \sim [R(A)]$ ([5; Lemma 7.2.2]). It is well-known that $H \ominus [f \in H; Af=0] = [R(A^*)]$ so that we obtain $[R(A^*)] \sim [R(A)] \sim H$, and this implies the existence of a partially isometric operator $Y \in M$ which gives the equivalency $H \sim [R(A^*)]$. Now, put $B = XAY^*$, then $B \in K$. Moreover, it is easily proved that B is a one-to-one transformation on the whole space H , by the definition of X and Y . Hence there exists the inverse $B^{-1} \in M$, so that $I = BB^{-1} \in K$, and this implies $K = M$. Thus we obtain $K \subseteq J$. This completes the proof.

Bibliography.

- 1 J. W. Calkin, Two-sided ideals and congruences in the rings of bounded operators in Hilbert space. Ann. of Math. 42 (1941), 839-873.
- 2 J. Dixmier, Les anneaux d'opérateurs de classe finie, Ann. École Norm. Sup., 66 (1949), 209-261.
- 3 J. Dixmier, Sur la réduction des anneaux d'opérateurs, Ann. École Norm. Sup., 68 (1951), 185-202.
- 4 Y. Misonou, On a weakly central operator algebra, Tôhoku Math. Journ., 4 (1952), 194-202.
- 5 F. J. Murray and J. von Neumann, On rings of operators, Ann. of Math., 37 (1936), 116-229.
- 6 J. von Neumann, Continuous Geometry, Part II. Princeton.

Department of Mathematics,
Waseda University. Tokio.

(*) Received Oct. 5, 1953.