

BSΔEs and BSDEs with non-Lipschitz drivers: Comparison, convergence and robustness

PATRICK CHERIDITO¹ and MITJA STADJE²

¹Princeton University, Princeton, NJ 08544, USA. E-mail: dito@princeton.edu

²Department of Econometrics and Operations Research, Tilburg University and CentER, 5000 LE Tilburg, The Netherlands. E-mail: m.a.stadje@uvt.nl

We provide existence results and comparison principles for solutions of backward stochastic difference equations (BSΔEs) and then prove convergence of these to solutions of backward stochastic differential equations (BSDEs) when the mesh size of the time-discretization goes to zero. The BSΔEs and BSDEs are governed by drivers $f^N(t, \omega, y, z)$ and $f(t, \omega, y, z)$, respectively. The new feature of this paper is that they may be non-Lipschitz in z . For the convergence results it is assumed that the BSΔEs are based on d -dimensional random walks W^N approximating the d -dimensional Brownian motion W underlying the BSDE and that f^N converges to f . Conditions are given under which for any bounded terminal condition ξ for the BSDE, there exist bounded terminal conditions ξ^N for the sequence of BSΔEs converging to ξ , such that the corresponding solutions converge to the solution of the limiting BSDE. An important special case is when f^N and f are convex in z . We show that in this situation, the solutions of the BSΔEs converge to the solution of the BSDE for every uniformly bounded sequence ξ^N converging to ξ . As a consequence, one obtains that the BSDE is robust in the sense that if (W^N, ξ^N) is close to (W, ξ) in distribution, then the solution of the N th BSΔE is close to the solution of the BSDE in distribution too.

Keywords: backward stochastic difference equations; backward stochastic differential equations; comparison principle; convergence; robustness

1. Introduction

The aim of this paper is to obtain general convergence results of solutions of stochastic backward equations in discrete time (BSΔEs) to solutions of stochastic backward equations in continuous time (BSDEs). The discrete equations are governed by drivers $f^N(t, \omega, y, z)$, $N \in \mathbb{N}$, and the continuous one by $f(t, \omega, y, z)$. The new feature of this paper is that f^N and f may be non-Lipschitz in z . We assume that the BSΔEs are based on d -dimensional random walks W^N converging to the d -dimensional Brownian motion W underlying the BSDE and that f^N tends to f . Convergence results for Lipschitz drivers have been obtained by Briand *et al.* [4,5] as well as Toldo [28,29]. In these papers, existence and uniqueness of solutions follow from a Picard iteration argument. Using results on convergence of filtrations from Coquet *et al.* [12], it can be shown that the Picard sequences approach each other asymptotically, which yields general convergence results. In the case of non-Lipschitz drivers this approach does not work, and neither the existence of solutions of BSΔEs nor their convergence to their counterparts in continuous time are clear.

In this paper, we start with a careful analysis of BSΔEs. Central to our approach is Theorem 4.2 which provides a comparison principle for BSΔEs. It requires drivers that can grow faster than linearly but strictly less than quadratically in z . Kobylanski [21] showed existence, comparison and uniqueness of solutions to BSDEs with general bounded terminal conditions and drivers of quadratic growth in z . However, in discrete time the situation is different. Example 4.1 shows that neither a general comparison principle nor convergence of solutions for diminishing step sizes can hold for BSΔEs if the drivers grow quadratically in z . Our main convergence results are Theorems 5.9 and 6.2. Theorem 5.9 shows that if f^N and f grow less than quadratically in z , then for any bounded terminal condition ξ for the BSDE, there exist bounded terminal conditions ξ^N for the BSΔEs such that the corresponding solutions Y^N converge to the solution Y of the BSDE in the following sense:

$$\sup_{0 \leq t \leq T} |Y_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2 \text{ when } N \rightarrow \infty. \quad (1.1)$$

Furthermore, if ξ is of the form $\xi = \varphi(W_{s_1}, \dots, W_{s_n})$ for a bounded, uniformly continuous function φ , then the ξ^N can be chosen as $\xi^N = \varphi(W_{s_1}^N, \dots, W_{s_n}^N)$. In Theorem 6.2, we prove that if the drivers f^N are convex in z , then (1.1) holds for every sequence of uniformly bounded ξ^N converging to ξ in L^2 . As a corollary one obtains that if (W^N, ξ^N) is close to (W, ξ) in distribution, then Y^N is close to Y in distribution too.

Discrete schemes for the approximation of solutions of BSDEs have been studied by a number of authors; see for instance, Ma *et al.* [23], Douglas *et al.* [15], Bally [1], Chevance [9], Coquet *et al.* [11], Ma *et al.* [22], Zhang and Zheng [31], Zhang [30], Bouchard and Touzi [3], Gobet *et al.* [18] and Otmani [17]. However, in all these papers the drivers are assumed to be Lipschitz. Recently, Imkeller and Reis [20] as well as Richou [27] have obtained results on the convergence of solutions of discretized BSDEs with drivers of quadratic growth under regularity assumptions on the terminal conditions and for specially chosen discrete-time drivers f^N . In Cheridito and Stadje [8] convergence results are shown for convex drivers and terminal conditions that are Lipschitz continuous in the underlying Brownian motion. Our results hold for general terminal conditions and general drivers f^N converging to f . But they need subquadratic growth of f^N in z . Comparison results for BSΔEs have also been studied in Cohen and Elliott [10] but under different assumptions than here.

The structure of the paper is as follows: In Section 2, we introduce the notation and provide some background material. Then we give an example showing that BSΔEs with non-Lipschitz drivers need not converge if the terminal conditions are not uniformly bounded. In Section 3, we show that BSΔEs admit solutions under very mild assumptions if the time-discretization is fine enough. Section 4 starts with an example showing two facts about BSΔEs with drivers of quadratic growth: (a) a general comparison principle cannot hold and (b) solutions of BSΔEs can explode if the step-size goes to zero even if the terminal conditions are uniformly bounded and converge to zero in L^2 . We then prove a general comparison principle for subquadratic BSΔEs. Section 5 gives convergence results of solutions of general BSΔEs to solutions of BSDEs, and in Section 6 we prove convergence results for drivers that are convex in z .

2. Notation and setup

We fix a finite time horizon $T \in \mathbb{R}_+$. As underlying process for the BSDE, we take a d -dimensional Brownian motion $(W_t)_{t \in [0, T]}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and denote by $(\mathcal{F}_t)_{t \in [0, T]}$ the augmented filtration generated by $(W_t)_{t \in [0, T]}$. Equalities and inequalities between random variables will, as usual, be understood in the \mathbb{P} -almost sure sense. As approximating processes we consider a sequence $(W_t^N)_{t \in [0, T]}$, $N \in \mathbb{N}$, of d -dimensional square-integrable martingales on $(\Omega, \mathcal{F}, \mathbb{P})$ starting at 0 with independent increments satisfying the following three conditions:

(C1) For every N there exists a finite sequence $0 = t_0^N < t_1^N < \dots < t_{i_N}^N = T$ such that

$$\lim_{N \rightarrow \infty} \sup_i |t_{i+1}^N - t_i^N| = 0$$

and W_t^N is constant on each of the intervals $[t_i^N, t_{i+1}^N)$.

(C2)

$$\Delta \langle W^{N,k} \rangle_{t_i^N} = \Delta \langle W^{N,l} \rangle_{t_i^N} > 0 \quad \text{for all } i \text{ and } 1 \leq k, l \leq d.$$

(C3)

$$\lim_{N \rightarrow \infty} \mathbb{E} \left[\sup_{0 \leq t \leq T} |W_t^N - W_t|^2 \right] = 0,$$

where $|\cdot|$ denotes the standard Euclidean norm on \mathbb{R}^d : $|x| := (\sum_{i=1}^d x_i^2)^{1/2}$.

Let (\mathcal{F}_t^N) be the filtration generated by (W_t^N) and define $\langle W^N \rangle_t := \langle W^{N,k} \rangle_t$. Since W^N has independent increments, $\langle W^N \rangle_t = \langle W^{N,k} \rangle_t$ is equal to $\mathbb{E}[(W_t^{N,k})^2]$, and it follows from (C3) that

$$\sup_{0 \leq t \leq T} |\langle W^N \rangle_t - t| = \sup_{0 \leq t \leq T} |\mathbb{E}[(W_t^{N,k})^2] - (W_t^k)^2| \rightarrow 0 \quad \text{for } N \rightarrow \infty. \quad (2.1)$$

In particular,

$$\lim_{N \rightarrow \infty} \max_i |\Delta \langle W^N \rangle_{t_i^N}| = 0.$$

Our standard example for the approximating processes W^N will be d -dimensional Bernoulli random walks.

Example 2.1. Let

$$t_i^N = i \frac{T}{N} \quad \text{and} \quad \tilde{W}_{t_i^N}^{N,k} = \sqrt{\frac{T}{N}} \sum_{j=1}^i X_j^{N,k}$$

for i.i.d. random variables $X_j^{N,k}$ on a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ with distribution $\tilde{\mathbb{P}}[X_j^{N,k} = \pm 1] = 1/2$. Extend (\tilde{W}_t^N) to $[0, T]$ such that it is constant on the intervals $[t_i^N, t_{i+1}^N)$. Then conditions (C1) and (C2) are satisfied. To fulfill (C3), one must transfer the random walks to another probability space. Since they converge to d -dimensional Brownian motion in distribution, there exists a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with a d -dimensional Brownian motion (W_t) and random walks (W_t^N) having the same distributions as (\tilde{W}_t^N) such that

$$\sup_{0 \leq t \leq T} |W_t^N - W_t| \rightarrow 0 \quad \text{almost surely as } N \rightarrow \infty; \tag{2.2}$$

see, for instance, Theorem I.2.7 in Ikeda and Watanabe [19]. It can be shown that the sequence $\sup_{0 \leq t \leq T} |W_t^N - W_t|^2$ is uniformly integrable. Therefore, the convergence (2.2) also holds in L^2 , and condition (C3) is satisfied.

The *driver* of the BSDE is a $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R}^d)$ -measurable function

$$f : [0, T] \times \Omega \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R},$$

where \mathcal{P} denotes the predictable σ -algebra on $[0, T] \times \Omega$ with respect to (\mathcal{F}_t) and $\mathcal{B}(\mathbb{R})$ and $\mathcal{B}(\mathbb{R}^d)$ are the Borel σ -algebras on \mathbb{R} and \mathbb{R}^d , respectively. We will assume throughout the paper that for fixed (t, ω) , $f(t, \omega, y, z)$ is continuous in (y, z) . Then $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}) \otimes \mathcal{B}(\mathbb{R}^d)$ -measurability of f is equivalent to $(t, \omega) \mapsto f(t, \omega, y, z)$ being predictable for all fixed (y, z) .

The approximating BSDEs have *drivers*

$$f^N : [0, T] \times \Omega \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$$

that are continuous in (y, z) , constant on the intervals $(t_i^N, t_{i+1}^N]$ and such that $\omega \mapsto f^N(t_i^N, \omega, y, z)$ is $\mathcal{F}_{t_i^N}^N$ -measurable. As usual, we henceforth suppress the dependence of f and f^N on ω in the notation.

The *terminal conditions* for the BSDE and BSDEs are given by random variables ξ, ξ^N that are measurable with respect to \mathcal{F}_T and \mathcal{F}_T^N , respectively.

A *solution* of the BSDE consists of a pair of predictable processes (Y_t, Z_t) with values in $\mathbb{R} \times \mathbb{R}^d$ such that

$$\mathbb{E} \left[\sup_{0 \leq t \leq T} Y_t^2 \right] < \infty, \quad \mathbb{E} \left[\left(\int_0^T |Z_s|^2 ds \right)^{1/2} \right] < \infty,$$

and

$$Y_t = \xi + \int_t^T f(s, Y_s, Z_s) ds - \int_t^T Z_s dW_s, \quad 0 \leq t \leq T. \tag{2.3}$$

In contrast to (W_t) , the approximating processes (W_t^N) do in general not have the predictable representation property. Therefore, a *solution* of the N th BSDE is a triple of (\mathcal{F}_t^N) -adapted processes (Y_t^N, Z_t^N, M_t^N) taking values in $\mathbb{R} \times \mathbb{R}^d \times \mathbb{R}$ such that (Y_t^N) is constant on the intervals

$[t_i^N, t_{i+1}^N)$, (Z_t^N) is constant on the intervals $(t_i^N, t_{i+1}^N]$, (M_t^N) is a square-integrable martingale starting at 0 and orthogonal to (W_t^N) that is constant on the intervals $[t_i^N, t_{i+1}^N)$ and

$$Y_t^N = \xi^N + \int_{(t, T]} f^N(s, Y_{s-}^N, Z_s^N) d\langle W^N \rangle_s - \int_{(t, T]} Z_s^N dW_s^N - (M_T^N - M_t^N). \quad (2.4)$$

Due to the particular form of (Y_t^N, Z_t^N, M_t^N) , (2.4) is equivalent to

$$Y_{t_i^N}^N = Y_{t_{i+1}^N}^N + f^N(t_{i+1}^N, Y_{t_i^N}^N, Z_{t_i^N}^N) \Delta \langle W^N \rangle_{t_i^N} - Z_{t_i^N}^N \Delta W_{t_i^N}^N - \Delta M_{t_i^N}^N, \quad (2.5)$$

$$Y_T^N = \xi^N. \quad (2.6)$$

Note that if (W_t^N) is a one-dimensional Bernoulli random walk, it has the predictable representation property and the orthogonal martingale terms in (2.4) and (2.5) disappear.

It is well known that if the driver f is Lipschitz-continuous in (y, z) and the terminal condition ξ is in L^2 , the BSDE (2.3) admits a unique solution (Y, Z) ; see, for instance, Pardoux and Peng [25] or the survey paper by El Karoui *et al.* [16]. Concerning the approximation of BSDEs with Lipschitz drivers, we recall the following result from Briand *et al.* [5]. Their assumptions are slightly different. But the result also holds in our setup.

Theorem 2.2 (Briand *et al.* [5]). *Assume $\xi^N \rightarrow \xi$ in L^2 and there exists a constant $K \in \mathbb{R}_+$ such that for all $N \in \mathbb{N}$, $y, y' \in \mathbb{R}$ and $z, z' \in \mathbb{R}^d$ the following four conditions hold:*

- (i) $\mathbb{E}[\sup_t f(t, 0, 0)^2] < \infty$;
- (ii) $|f(t, y, z) - f(t, y', z')| \leq K(|y - y'| + |z - z'|)$;
- (iii) $|f^N(t, y, z) - f^N(t, y', z')| \leq K(|y - y'| + |z - z'|)$;
- (iv) $\sup_t |f^N(t, y, z) - f(t, y, z)| \rightarrow 0$ in L^2 as $N \rightarrow \infty$.

Then, for N large enough, the N th BSDE has a unique solution (Y^N, Z^N, M^N) , and

$$\sup_t \left(|Y_t^N - Y_t| + \left| \int_0^t Z_s^N dW_s^N - \int_0^t Z_s dW_s \right| + |M_t^N| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^2$$

as well as

$$\sup_t \left(\sum_{k=1}^d \left| \int_0^t Z_s^{N,k} d\langle W^N \rangle_s - \int_0^t Z_s^k ds \right|^2 + \left| \int_0^t |Z_s^N|^2 d\langle W^N \rangle_s - \int_0^t |Z_s|^2 ds \right| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^1,$$

where (Y, Z) is the unique solution of the BSDE (2.3).

Remark 2.3. Two special cases of terminal conditions satisfying $\xi^N \rightarrow \xi$ in L^2 are:

- (a) $\xi = \varphi(W_T)$ and $\xi^N = \varphi(W_T^N)$ for a continuous function $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\varphi^2(W_T^N)$, $N \in \mathbb{N}$, is uniformly integrable.

(b) $\xi \in L^2(\mathcal{F}_T)$ general and $\xi^N = \mathbb{E}[\xi | \mathcal{F}_T^N]$.

The aim of this paper is to obtain similar convergence results for non-Lipschitz drivers. However, the following example shows that we cannot hope for general results under the sole assumption $\xi^N \rightarrow \xi$ in L^2 .

Example 2.4. Consider a one-dimensional Bernoulli random walk with $T = 1$, $t_i^N = i/N$ and $\mathbb{P}[\Delta W_{t_i^N}^N = \pm\sqrt{1/N}] = 1/2$. Then

$$\Delta \langle W^N \rangle_{t_i^N} = \mathbb{E}[(\Delta W_{t_i^N}^N)^2] = 1/N.$$

Fix $q \in (1, 2)$ and a sequence of constants $a^N \geq 2N^{(1-q/2)/(q-1)}$. Consider the BSDEs

$$Y_{t_i^N}^N = Y_{t_{i+1}^N}^N + |Z_{t_{i+1}^N}^N|^q \Delta \langle W^N \rangle_{t_{i+1}^N} - Z_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^N, \quad Y_T^N = a^N \mathbf{1}_{\{W_{t_T^N}^N = \sqrt{N}\}}.$$

It can easily be checked that

$$Z_{t_N^N}^N = \frac{\sqrt{N}}{2} a^N \mathbf{1}_{\{W_{t_{N-1}^N}^N = (N-1)/\sqrt{N}\}} \quad \text{and} \quad Y_{t_{N-1}^N}^N = a_{t_{N-1}^N}^N \mathbf{1}_{\{W_{t_{N-1}^N}^N = (N-1)/\sqrt{N}\}}$$

for

$$a_{t_{N-1}^N}^N = \frac{a^N}{2} + 2^{-q} N^{q/2-1} (a^N)^q \geq a^N.$$

Continuing this way one gets

$$Z_{t_{N-1}^N}^N = \frac{\sqrt{N}}{2} a_{t_{N-1}^N}^N \mathbf{1}_{\{W_{t_{N-2}^N}^N = (N-2)/\sqrt{N}\}} \quad \text{and} \quad Y_{t_{N-2}^N}^N = a_{t_{N-2}^N}^N \mathbf{1}_{\{W_{t_{N-2}^N}^N = (N-2)/\sqrt{N}\}}$$

with

$$a_{t_{N-2}^N}^N = \frac{a_{t_{N-1}^N}^N}{2} + 2^{-q} N^{q/2-1} (a_{t_{N-1}^N}^N)^q \geq a_{t_{N-1}^N}^N,$$

and so on. In particular,

$$Y_0^N \geq a^N \geq 2N^{(1-(q/2))/(q-1)} \rightarrow \infty \quad \text{for } N \rightarrow \infty.$$

Note that for $a^N = 2N^{(1-q/2)/(q-1)}$, one has $\xi^N \rightarrow 0$ in L^p for all $p \in (0, \infty)$ but not in L^∞ .

The example shows that in the case of super-linear growth of f^N in z one cannot expect convergence of the discrete-time solutions if the terminal conditions are uniformly L^p -bounded and converge in L^p for $p < \infty$. This is not unexpected since in the literature on BSDEs with non-Lipschitz drivers it is usually required that the terminal condition be in L^∞ or sufficiently well exponentially integrable (see Kobylanski [21], or Briand and Hu [6]). Consequently, in this paper, we will always assume:

(C4)

$$\sup_N \|\xi^N\|_\infty < \infty \quad \text{and} \quad \|\xi\|_\infty < \infty.$$

We shortly summarize the notation and assumptions that have been introduced in this section:

- W^N , $N \in \mathbb{N}$, is a sequence of discrete-time martingales approximating the d -dimensional Brownian motion W .
- f and ξ are the driver and terminal condition of the BSDE (2.3). A solution to (2.3) will be denoted by (Y, Z) .
- f^N and ξ^N are the drivers and terminal conditions of the BSΔEs (2.4). Solutions will be denoted by (Y^N, Z^N, M^N) .
- We always assume (C1)–(C4).

3. Solutions of BSΔEs

In this section, we present two results on solutions of BSΔEs that will be needed later in the paper. Their proofs are straightforward and therefore, given in the [Appendix](#).

Lemma 3.1. *If a solution (Y^N, Z^N, M^N) of the N th BSΔE exists, one has*

$$Y_{t_i^N}^N - f^N(t_{i+1}^N, Y_{t_i^N}^N, Z_{t_{i+1}^N}^N) \Delta \langle W^N \rangle_{t_{i+1}^N} = \mathbb{E}[Y_{t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N], \tag{3.1}$$

and the pair (Z^N, M^N) is uniquely determined by Y^N through

$$Z_{t_{i+1}^N}^{N,k} = \frac{\mathbb{E}[Y_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^{N,k} | \mathcal{F}_{t_i^N}^N]}{\Delta \langle W^N \rangle_{t_{i+1}^N}}, \tag{3.2}$$

$$\Delta M_{t_{i+1}^N}^N = Y_{t_{i+1}^N}^N - \mathbb{E}[Y_{t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N] - Z_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^N. \tag{3.3}$$

Concerning the existence of solutions to BSΔEs, one has the following result. For the special case where W^N is a one-dimensional Bernoulli random walk, see Peng [26].

Proposition 3.2. *Assume there exists a constant $K \in \mathbb{R}_+$ and a locally bounded function $g : \mathbb{R}^d \rightarrow \mathbb{R}_+$ such that*

$$|f^N(t, y, z)| \leq K(1 + |y| + g(z)) \quad \text{and} \quad \max_i \Delta \langle W^N \rangle_{t_i^N} < 1/K.$$

Then the N th BSΔE has a solution (Y^N, Z^N, M^N) such that Y^N and Z^N are bounded. If W^N is bounded, then so is M^N .

Remark 3.3. For $\max_i \Delta \langle W^N \rangle_{t_i^N} \geq 1/K$ a solution of the N th BSΔE might not exist. For example, let W^1 be a one-dimensional Bernoulli random walk with $t_0^1 = 0$, $t_1^1 = 1 = T$,

$\mathbb{P}[\Delta W_1^1 = \pm 1] = 1/2$, $\xi^1 = 1$ and $f^1(t, y, z) = y$. Since the terminal condition is deterministic, one must choose $Z_1^1 = 0$, and (A.2) becomes

$$Y_0^1 - Y_1^1 = 1,$$

an equation without solution.

4. Comparison principle for BSΔEs

Our main tool to derive convergence results will be a comparison principle for BSΔEs of the following form: Let f_1^N, f_2^N be drivers and ξ_1^N, ξ_2^N terminal conditions such that $f_1^N(t, y, z) \geq f_2^N(t, y, z)$ for all t, y, z and $\xi_1^N \geq \xi_2^N$. Then the corresponding solutions satisfy $Y_{1,t}^N \geq Y_{2,t}^N$ for all t .

The next example shows that if the drivers grow quadratically in z , a general comparison principle for BSΔEs cannot hold.

Example 4.1. As in Example 2.4, let W^N be a one-dimensional Bernoulli random walk with $T = 1$, $t_i^N = i/N$ and $\mathbb{P}[\Delta W_{t_i^N}^N = \pm\sqrt{1/N}] = 1/2$. Consider the BSΔEs

$$Y_{t_i^N}^N = Y_{t_{i+1}^N}^N + (Z_{t_{i+1}^N}^N)^2 \Delta \langle W^N \rangle_{t_{i+1}^N} - Z_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^N, \tag{4.1}$$

$$Y_T^N = a 1_{\{W_{t_T^N}^N = \sqrt{N}\}} \tag{4.2}$$

for a constant $a > 2$ and define $\varepsilon > 0$ by $a = 2(1 + 2\varepsilon)$. Then

$$Z_{t_{i_N}^N}^N = \frac{\sqrt{N}}{2} a 1_{\{W_{t_{i_N-1}^N}^N = (N-1)/\sqrt{N}\}}, \quad Y_{t_{i_N-1}^N}^N = a_{t_{i_N-1}^N}^N 1_{\{W_{t_{i_N-1}^N}^N = (N-1)/\sqrt{N}\}},$$

where

$$a_{t_{i_N-1}^N}^N = \frac{a}{2} + \left(\frac{a}{2}\right)^2 = a(1 + \varepsilon)$$

and

$$Z_{t_{i_N-1}^N}^N = \frac{\sqrt{N}}{2} a_{t_{i_N-1}^N}^N 1_{\{W_{t_{i_N-2}^N}^N = (N-2)/\sqrt{N}\}}, \quad Y_{t_{i_N-2}^N}^N = a_{t_{i_N-2}^N}^N 1_{\{W_{t_{i_N-2}^N}^N = (N-2)/\sqrt{N}\}}$$

for

$$a_{t_{i_N-2}^N}^N = \frac{a_{t_{i_N-1}^N}^N}{2} + \left(\frac{a_{t_{i_N-1}^N}^N}{2}\right)^2 \geq \frac{a_{t_{i_N-1}^N}^N}{2} \left(1 + \frac{a}{2}\right) = a_{t_{i_N-1}^N}^N (1 + \varepsilon).$$

Continuing this computation, one obtains

$$Y_0^N \geq a(1 + \varepsilon)^N \rightarrow \infty \quad \text{as } N \rightarrow \infty.$$

Note that the terminal conditions Y_T^N are uniformly L^∞ -bounded in N and $Y_T^N \rightarrow 0$ in L^p for all $p < \infty$. But the solutions Y_t^N explode as $N \rightarrow \infty$. We also point out that for fixed N , the solutions to equation (4.1) are not monotone in the terminal condition. Indeed, $(\tilde{Y}_t^N, \tilde{Z}_t^N) \equiv (a, 0)$ is a solution of equation (4.1) with terminal condition $\tilde{Y}_T^N = a \geq Y_T^N$. However, $\tilde{Y}_0^N < Y_0^N$. In particular, the comparison principle is violated.

In view of Example 4.1, we restrict ourselves in the next theorem to drivers that grow less than quadratically in z . We need the following assumption on the increments of W^N :

$$(W1) \text{ There exists a constant } q \in [1, 2) \text{ such that } \lim_{N \rightarrow \infty} \max_{i,k} \frac{\|\Delta W_{t_i^N}^{N,k}\|_\infty}{\Delta(W^N)_{t_i^N}^{q/4}} = 0.$$

Note that the standard Bernoulli random walks of Example 2.1 satisfy (W1) for all $q \in [1, 2)$. The subsequent theorem establishes a comparison result for BSΔEs governed by non-Lipschitz drivers.

Theorem 4.2. *Let $C, K, L \in \mathbb{R}_+$ and assume (W1) holds for some $q \in [1, 2)$. Then there exists $N_0 \in \mathbb{N}$ such that for every $N \geq N_0$, all drivers $f_1^N \geq f_2^N$ and terminal conditions $\xi_1^N \geq \xi_2^N$ satisfying*

- (i) $\|\xi_m^N\|_\infty \leq C$,
- (ii) $|f_m^N(t, y, z)| \leq K(1 + |y| + |z|^q)$ for all $(t, y, z) \in [0, T] \times \mathbb{R}^{d+1}$,
- (iii) $|f_m^N(t, y_1, z) - f_m^N(t, y_2, z)| \leq L(1 + |z|^q)|y_1 - y_2|$ for all $(t, y_1, y_2, z) \in [0, T] \times \mathbb{R}^{d+2}$ such that $|y_1|, |y_2| \leq (C + 1) \exp(KT)$,
- (iv) $|f_m^N(t, y, z_1) - f_m^N(t, y, z_2)| \leq L(1 + (|z_1| \vee |z_2|)^{q/2})|z_1 - z_2|$ for all $(t, y, z_1, z_2) \in [0, T] \times \mathbb{R}^{2d+1}$ such that $|y| \leq (C + 1) \exp(KT)$,

the BSΔEs with parameters (f_m^N, ξ_m^N) have unique solutions (Y_m^N, Z_m^N, M_m^N) , $m = 1, 2$, and

$$(C + 1) \exp(K(T - t)) \geq Y_{1,t}^N \geq Y_{2,t}^N \geq -(C + 1) \exp(K(T - t)) \quad \text{for all } t \in [0, T].$$

To prove Theorem 4.2, we need the following two lemmas, whose proofs can be found in the Appendix. The first one provides a comparison principle under stronger assumptions than Theorem 4.2. The second one gives conditions under which the Y^N are uniformly bounded in N .

Lemma 4.3. *Let $C, K \in \mathbb{R}_+$ and assume (W1) holds for some $q \in [1, 2)$. Then there exists $N_0 \in \mathbb{N}$ such that for every $N \geq N_0$, all drivers $f_1^N \geq f_2^N$ and terminal conditions $\xi_1^N \geq \xi_2^N$ satisfying conditions (i) and (ii) of Theorem 4.2 as well as*

- (iii) $|f_m^N(t, y_1, z) - f_m^N(t, y_2, z)| \leq K(1 + |z|^q)|y_1 - y_2|$ for all $(t, y_1, y_2, z) \in [0, T] \times \mathbb{R}^{d+2}$,
- (iv) $|f_m^N(t, y, z_1) - f_m^N(t, y, z_2)| \leq qK(1 + (|z_1| \vee |z_2|)^{q/2})|z_1 - z_2|$ for all $(t, y, z_1, z_2) \in [0, T] \times \mathbb{R}^{2d+1}$,

the BSΔEs with parameters (f_m^N, ξ_m^N) have unique solutions (Y_m^N, Z_m^N, M_m^N) , $m = 1, 2$, and

$$(C + 1) \exp(K(T - t)) \geq Y_{1,t}^N \geq Y_{2,t}^N \geq -(C + 1) \exp(K(T - t)) \quad \text{for all } t \in [0, T]. \quad (4.3)$$

Lemma 4.4. *Let $C, K \in \mathbb{R}_+$ and assume (W1) holds for some $q \in [1, 2)$. Then there exists $N_0 \in \mathbb{N}$ such that for every $N \geq N_0$, all drivers f^N and terminal conditions ξ^N satisfying*

- (i) $\|\xi^N\|_\infty \leq C,$
- (ii) $|f^N(t, y, z)| \leq K(1 + |y| + |z|^q)$ for all $t \in [0, T], y \in \mathbb{R}$ and $z \in \mathbb{R}^d,$

every solution (Y^N, Z^N, M^N) of the N th BSΔE satisfies

$$|Y_t^N| \leq (C + 1) \exp(K(T - t)) \quad \text{for all } t \in [0, T]. \tag{4.4}$$

We now are ready for the proof.

Proof of Theorem 4.2. It follows from Proposition 3.2 and Lemma 4.4 that there exists an N_1 such that for all $N \geq N_1$, the N th BSΔE has a solution (Y^N, Z^N, M^N) for all f^N and ξ^N satisfying conditions (i) and (ii) of Theorem 4.2, and every such solution satisfies $|Y_t^N| \leq (C + 1) \exp(K(T - t)), 0 \leq t \leq T$. Now choose $N_0 \geq N_1$ such that Lemma 4.3 holds for $\tilde{K} = K \vee L$ instead of K and fix $N \geq N_0$. If $f_1^N \geq f_2^N$ and $\xi_1^N \geq \xi_2^N$ are drivers and terminal conditions satisfying conditions (i)–(iv) of Theorem 4.2, then there exist corresponding solutions $(Y_m^N, Z_m^N, M_m^N), m = 1, 2$, both of which satisfy $|Y_{m,t}^N| \leq (C + 1) \exp(K(T - t))$. So one can change the drivers f_m^N for $|y| > (C + 1) \exp(KT)$ such that they satisfy the conditions of Lemma 4.3, and it follows that $Y_{1,t}^N \geq Y_{2,t}^N$. In particular, both solutions are unique. \square

5. Convergence results for drivers with subquadratic growth

With a slight abuse of notation, the discrete-time drivers can be written as $f^N(t, W^N, y, z)$. By predictability, $f^N(t_{i+1}^N, W^N, y, z)$ only depends on $W_{t_1^N}^N, \dots, W_{t_i^N}^N$. Let $q \in [1, 2)$ and consider the following conditions on the drivers f^N : There exists a constant $K > 0$ such that

- (f1) For all $N \in \mathbb{N}, w \in \mathbb{R}^{d \times i_N}$ and $(t, y, z) \in [0, T] \times \mathbb{R}^{d+1},$

$$|f^N(t, w, y, z)| \leq K(1 + |y| + |z|^q).$$

- (f2) For all $N \in \mathbb{N}, w \in \mathbb{R}^{d \times i_N}$ and $(t, y_1, y_2, z) \in [0, T] \times \mathbb{R}^{d+2},$

$$|f^N(t, w, y_1, z) - f^N(t, w, y_2, z)| \leq K|y_1 - y_2|.$$

- (f3) For every $a \in \mathbb{R}_+$ there exists $b \in \mathbb{R}_+$ such that for all $N \in \mathbb{N}, t \in [0, T], y \in [-a, a], w \in \mathbb{R}^{d \times i_N}$ and $z_1, z_2 \in \mathbb{R}^d,$

$$|f^N(t, w, y, z_1) - f^N(t, w, y, z_2)| \leq b(1 + (|z_1| \vee |z_2|)^{q/2})|z_1 - z_2|.$$

- (f4) For all $N \in \mathbb{N}, i = 0, \dots, i_N - 1, w_1, w_2 \in \mathbb{R}^{d \times i_N}$ and $(y, z) \in \mathbb{R}^{d+1},$

$$|f^N(t_{i+1}^N, w_1, y, z) - f^N(t_{i+1}^N, w_2, y, z)| \leq K \sup_{0 \leq t \leq t_i^N} |w_1(t) - w_2(t)|.$$

(f5) For all $(y, z) \in \mathbb{R}^{d+1}$,

$$\sup_{0 \leq t \leq T} |f^N(t, y, z) - f(t, y, z)| \rightarrow 0 \quad \text{in } L^2 \text{ as } N \rightarrow \infty.$$

For a measurable function $g : [0, T] \times \Omega \times \mathbb{R}^{d+1} \rightarrow \mathbb{R}$, denote

$$\|g\|_\infty = \text{ess sup}_\omega \sup_{t, y, z} |g(t, \omega, y, z)|.$$

The following lemma shows that the solutions of the BSΔEs are stable in the terminal condition and the driver function. The proof relies on Theorem 4.2 and can be found in the Appendix.

Lemma 5.1. *Let $C, K \in \mathbb{R}_+$ and assume condition (W1) holds for some $q \in [1, 2)$. Then there exists $N_0 \in \mathbb{N}$ and a constant $D \in \mathbb{R}_+$ such that for all $N \geq N_0$, all terminal conditions ξ_1^N, ξ_2^N bounded by C and drivers f_1^N, f_2^N satisfying (f1)–(f3) as well as $\|f_1^N - f_2^N\|_\infty \leq K$, the BSΔEs with parameters (f_m^N, ξ_m^N) have unique solutions (Y_m^N, Z_m^N, M_m^N) , $m = 1, 2$, and*

$$\sup_{0 \leq t \leq T} |Y_{1,t}^N - Y_{2,t}^N| \leq D(\|f_1^N - f_2^N\|_\infty + \|\xi_1^N - \xi_2^N\|_\infty).$$

The next lemma shows that for Lipschitz-continuous terminal conditions, the Z^N are uniformly bounded. This will be a key ingredient in the proofs of our convergence results. The proof is given in the Appendix.

Lemma 5.2. *Assume (W1) and (f1)–(f4) hold for some $q \in [1, 2)$ and the ξ^N are of the form $\xi^N = \varphi(W_{s_1}^N, \dots, W_{s_n}^N)$ for fixed $n \in \mathbb{N}$, $0 \leq s_1 < \dots < s_n \leq T$, and a bounded Lipschitz-continuous function $\varphi : \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$. Then there exists an $N_0 \in \mathbb{N}$ such for all $N \geq N_0$, the N th BSΔE has a unique solution (Y^N, Z^N) and $\sup_{N \geq N_0} \|\sup_t |Z_t^N|\|_\infty < \infty$.*

Remark 5.3. In general $\sup_{N \geq N_0} \|\sup_t |Z_t^N|\|_\infty < \infty$ does not hold if φ is not Lipschitz-continuous. For example, consider one-dimensional Bernoulli random walks W^N with $T = 1$, $t_i^N = i/N$ and $\mathbb{P}[\Delta W_{t_i^N}^N = \pm\sqrt{1/N}] = 1/2$. Let the terminal conditions be of the form

$$\xi^N = \begin{cases} \sqrt{W_1^N} \wedge 1, & \text{if } W_1^N \geq 0, \\ -\sqrt{-W_1^N} \vee -1, & \text{if } W_1^N < 0. \end{cases}$$

On the set $\{W_{(N-1)/N}^N = 0\}$ one has $\xi^N = \text{sign}(\Delta W_1^N) \sqrt{|\Delta W_1^N|}$, and hence, by Lemma 3.1,

$$Z_1^N = \frac{\mathbb{E}[\xi^N \Delta W_1^N | W_{(N-1)/N}^N = 0]}{\Delta \langle W^N \rangle_1} = N^{1/4}.$$

In particular, $Z_1^N \rightarrow \infty$ as $N \rightarrow \infty$ on the set $\{W_{(N-1)/N}^N = 0\}$.

Before we prove convergence of solutions of BSΔEs to solutions of BSDEs, we recall the following result on quadratic BSDEs, which follows from Theorems 2.5–2.7 of Morlais [24].

Theorem 5.4 (Morlais [24]). *Let $K \in \mathbb{R}_+$ such that*

$$|f(t, y, z)| \leq K(1 + |y| + |z|^2), \tag{5.1}$$

$$|f(t, y_1, z) - f(t, y_2, z)| \leq K|y_1 - y_2| \quad \text{for all } y_1, y_2 \in \mathbb{R}, \tag{5.2}$$

and for every $a \in \mathbb{R}_+$ there exists $b \in \mathbb{R}_+$ such that

$$|f(t, y, z_1) - f(t, y, z_2)| \leq b(1 + (|z_1| \vee |z_2|))|z_1 - z_2| \tag{5.3}$$

for all $t \in [0, T]$, $y \in [-a, a]$ and $z_1, z_2 \in \mathbb{R}^d$. Then the BSDE (2.3) has a unique solution (Y, Z) such that Y is bounded. Furthermore, for bounded terminal conditions $\xi_1 \geq \xi_2$ and drivers $f_1 \geq f_2$ fulfilling (5.1)–(5.3), the corresponding solutions satisfy $Y_{1,t} \geq Y_{2,t}$ for all t .

Remark 5.5. Actually, Morlais [24] makes slightly different assumptions. In her paper, the underlying noise process is continuous but does not have to be a Brownian motion, and condition (5.3) is assumed to hold for a constant b independent of a . However, existence of a solution (Y, Z) with bounded Y already follows from (5.1), and if Y is bounded by a constant $a \in \mathbb{R}_+$, the driver $f(t, y, z)$ only matters for $y \in [-a, a]$ and can be modified so that it satisfies (5.3) for a constant b independent of a . Hence, assumptions (5.1)–(5.3) are sufficient for Theorem 5.4.

Proposition 5.6. *Assume there exists a $q \in [1, 2)$ such that (W1) and (f1)–(f5) hold. If ξ and ξ^N are of the form $\xi = \varphi(W_{s_1}, \dots, W_{s_n})$ and $\xi^N = \varphi(W_{s_1^N}, \dots, W_{s_n^N})$ for fixed $n \in \mathbb{N}$, $0 \leq s_1 < \dots < s_n \leq T$, and a bounded Lipschitz-continuous function $\varphi : \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$, then there exists an $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$, the N th BSΔE has a unique solution (Y^N, Z^N, M^N) satisfying $\sup_{N \geq N_0} \|\sup_t |Z_t^N|\|_\infty < \infty$, the BSDE (2.3) has a unique solution (Y, Z) with bounded Y , and*

$$\sup_t \left(|Y_t^N - Y_t| + \left| \int_0^t Z_s^N dW_s^N - \int_0^t Z_s dW_s \right| + |M_t^N| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^2$$

as well as

$$\sup_t \left(\sum_{k=1}^d \left| \int_0^t Z_s^{N,k} d\langle W^N \rangle_s - \int_0^t Z_s^k ds \right|^2 + \left| \int_0^t |Z_s^N|^2 d\langle W^N \rangle_s - \int_0^t |Z_s|^2 ds \right| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^1.$$

In particular, there exists a constant $R \in \mathbb{R}_+$ such that $|Z| \leq R \nu \otimes \mathbb{P}$ -almost everywhere, where ν denotes Lebesgue measure on $[0, T]$.

Proof. It follows from (f1)–(f5) that the driver f satisfies (5.1)–(5.3). So one obtains from Theorem 5.4 that the BSDE (2.3) has a unique solution (Y, Z) such that Y is bounded. By Lemma 5.2, there exists $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$, the N th BSΔE has a unique solution (Y^N, Z^N, M^N) and $\sup_{N \geq N_0} \|\sup_t |Z_t^N|\|_\infty \leq R$ for some constant $R \in \mathbb{R}_+$. Define

$$\tilde{f}^N(t, y, z) = \begin{cases} f^N(t, y, z), & \text{for } |z| \leq R, \\ f^N(t, y, Rz/|z|), & \text{for } |z| > R \end{cases}$$

and

$$\tilde{f}(t, y, z) = \begin{cases} f(t, y, z), & \text{for } |z| \leq R, \\ f(t, y, Rz/|z|), & \text{for } |z| > R. \end{cases}$$

Then the \tilde{f}^N are uniformly Lipschitz in (y, z) and

$$\sup_{0 \leq t \leq T} |\tilde{f}^N(t, y, z) - \tilde{f}(t, y, z)| \rightarrow 0 \quad \text{in } L^2 \text{ as } N \rightarrow \infty.$$

So it follows that \tilde{f}^N and \tilde{f} fulfill the conditions of Theorem 2.2. Denote by $(\tilde{Y}^N, \tilde{Z}^N, \tilde{M}^N)$ the solution to the N th BSΔE with parameters (\tilde{f}^N, ξ^N) and by (\tilde{Y}, \tilde{Z}) the solution of the BSDE corresponding to (\tilde{f}, ξ) . Since the Z^N are bounded by R , (Y^N, Z^N, M^N) is also a solution of the BSΔE corresponding to (\tilde{f}^N, ξ^N) . So it follows from Theorem 4.2 that for N large enough, $(Y^N, Z^N, M^N) = (\tilde{Y}^N, \tilde{Z}^N, \tilde{M}^N)$, and we may apply Theorem 2.2 to conclude that

$$\sup_t \left(|Y_t^N - \tilde{Y}_t| + \left| \int_0^t Z_s^N dW_s^N - \int_0^t \tilde{Z}_s dW_s \right| + |M_t^N| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^2, \quad (5.4)$$

and

$$\sup_t \left(\sum_{k=1}^d \left| \int_0^t Z_s^{N,k} d\langle W^N \rangle_s - \int_0^t \tilde{Z}_s^k ds \right|^2 + \left| \int_0^t |Z_s^N|^2 d\langle W^N \rangle_s - \int_0^t |\tilde{Z}_s|^2 ds \right| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad (5.5)$$

in L^1 . It follows from (5.5) that $|\tilde{Z}| \leq R \vee \otimes \mathbb{P}$ -almost everywhere. So (\tilde{Y}, \tilde{Z}) is also a solution of the original BSDE corresponding to (f, ξ) , and it follows from Theorem 5.4 that it is equal to (Y, Z) . This completes the proof. \square

Another result that we need below is the following proposition.

Proposition 5.7 (Briand and Hu [6]). *Let $(\xi_m)_{m \in \mathbb{N}}$ be a sequence of \mathcal{F}_T -measurable random variables such that $\sup_m \|\xi_m\|_\infty < \infty$ and $\xi_m \rightarrow \xi$ almost surely. Furthermore assume that f satisfies (5.1). Let (Y_m, Z_m) and (Y, Z) be solutions of the BSDEs corresponding to (f, ξ_m) and (f, ξ) , respectively, such that Y_m and Y are bounded. If Y_m is increasing (or decreasing) in m , then*

$$\sup_t |Y_{m,t} - Y_t| \rightarrow 0 \quad \text{a.s.} \quad \text{and} \quad \mathbb{E} \left[\int_0^T |Z_{m,s} - Z_s|^2 ds \right] \rightarrow 0 \quad \text{for } m \rightarrow \infty.$$

Remark 5.8. Note that if f satisfies (5.1)–(5.3), then Proposition 5.7 holds without the assumption that Y_m is increasing or decreasing in m . Indeed, by Theorem 5.4 one has $Y(\xi_1) \geq Y(\xi_2)$ for $\xi_1 \geq \xi_2$ (where $Y(\xi)$ denotes the solution of the BSDE with driver f and terminal condition ξ). Define $\hat{\xi}_m = \sup_{n \geq m} \xi_n$ and $\tilde{\xi}_m = \inf_{n \geq m} \xi_n$. Then one obtains from Proposition 5.7 that $\sup_t |Y_t(\hat{\xi}_m) - Y_t(\xi)| \rightarrow 0$ and $\sup_t |Y_t(\tilde{\xi}_m) - Y_t(\xi)| \rightarrow 0$ a.s., and therefore also $\sup_t |Y_t(\xi_m) - Y_t(\xi)| \rightarrow 0$ a.s. The convergence of $Z(\xi_m)$ to $Z(\xi)$ now follows exactly as in the proof of Proposition 2.4 in Kobylanski [21].

The next theorem shows that for any continuous-time terminal condition there exists a sequence of discrete-time terminal conditions such that the corresponding solutions of the BSΔEs converge to their counterparts in continuous time.

Theorem 5.9. Assume there exists a $q \in [1, 2)$ such that (W1) and (f1)–(f5) are satisfied. Then for every $\xi \in L^\infty(\mathcal{F}_T)$, there exist \mathcal{F}_T^N -measurable $\tilde{\xi}^N$ bounded by $\|\xi\|_\infty$ such that for N large enough, the N th BSΔE with terminal condition $\tilde{\xi}^N$ has a unique solution $(\tilde{Y}^N, \tilde{Z}^N, \tilde{M}^N)$ and

$$\sup_t \left(|\tilde{Y}_t^N - Y_t| + \left| \int_0^t \tilde{Z}_s^N dW_s^N - \int_0^t Z_s dW_s \right| + |\tilde{M}_t^N| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^2 \tag{5.6}$$

as well as

$$\sup_t \left(\left| \sum_{k=1}^d \int_0^t \tilde{Z}_s^{N,k} d(W^N)_s - \int_0^t Z_s^k ds \right|^2 + \left| \int_0^t |\tilde{Z}_s^N|^2 d(W^N)_s - \int_0^t |Z_s|^2 ds \right| \right) \xrightarrow{(N \rightarrow \infty)} 0 \quad \text{in } L^1, \tag{5.7}$$

where (Y, Z) is the unique solution of the BSDE (2.3) with bounded Y . Moreover, if $\xi = \varphi(W_{s_1}, \dots, W_{s_n})$ and $\xi^N = \varphi(W_{s_1}^N, \dots, W_{s_n}^N)$ for a bounded, uniformly continuous function $\varphi : \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$, then

$$\sup_t |Y_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2 \text{ as } N \rightarrow \infty,$$

where (Y^N, Z^N, M^N) solves the N th BSΔE with terminal condition ξ^N .

Proof. Given a random variable $\xi \in L^\infty(\mathcal{F}_T)$, there exists a sequence $n_m, m \in \mathbb{N}$, of positive integers together with times $0 \leq s_1^m < \dots < s_{n_m}^m \leq T$ and Lipschitz-continuous functions $\varphi_m : \mathbb{R}^{d \times n_m} \rightarrow \mathbb{R}$ bounded by $\|\xi\|_\infty$ such that the random variables $\xi_m := \varphi_m(W_{s_1}, \dots, W_{s_{n_m}})$ converge to ξ almost surely. It follows from (f1)–(f5) that the driver f satisfies (5.1)–(5.3). So one obtains from Theorem 5.4 that there exist unique solutions (Y, Z) and (Y_m, Z_m) to the BSDEs corresponding to (f, ξ) and (f, ξ_m) , respectively, such that Y and Y_m are bounded. Since for fixed m , φ_m is bounded and Lipschitz-continuous, one can apply Proposition 5.6 and choose $N_m \in \mathbb{N}$

increasing in m such that for all $N \geq N_m$, one has

$$\begin{aligned} & \mathbb{E} \left[\sup_t \left(|Y_{m,t}^N - Y_{m,t}|^2 + \left| \int_0^t Z_{m,s}^N dW_s^N - \int_0^t Z_{m,s} dW_s \right|^2 + |M_{m,t}^N|^2 \right. \right. \\ & \quad \left. \left. + \sum_{k=1}^d \left| \int_0^t Z_{m,s}^{N,k} d\langle W^N \rangle_s - \int_0^t Z_{m,s}^k ds \right|^2 \right. \right. \\ & \quad \left. \left. + \left| \int_0^t |Z_{m,s}^N|^2 d\langle W^N \rangle_s - \int_0^t |Z_{m,s}|^2 ds \right| \right) \right] \leq \frac{1}{m}, \end{aligned}$$

where (Y_m^N, Z_m^N, M_m^N) is the unique solution to the N th BSΔE with driver f^N and terminal condition $\xi_m^N := \varphi_m(W_{s_1}^N, \dots, W_{s_{m_N}}^N)$. Now set $\tilde{\xi}^N := \xi_{m_N}^N$ and $(\tilde{Y}^N, \tilde{Z}^N, \tilde{M}^N) := (Y_{m_N}^N, Z_{m_N}^N, M_{m_N}^N)$, where for given N , m_N is the largest m satisfying $N_m \leq N$. Then $\lim_{N \rightarrow \infty} m_N = \infty$, and therefore,

$$\begin{aligned} & \mathbb{E} \left[\sup_t \left(|\tilde{Y}_t^N - Y_{m_N,t}|^2 + \left| \int_0^t \tilde{Z}_s^N dW_s^N - \int_0^t Z_{m_N,s} dW_s \right|^2 + |\tilde{M}_t^N|^2 \right. \right. \\ & \quad \left. \left. + \sum_{k=1}^d \left| \int_0^t \tilde{Z}_s^{N,k} d\langle W^N \rangle_s - \int_0^t Z_{m_N,s}^k ds \right|^2 \right. \right. \\ & \quad \left. \left. + \left| \int_0^t |\tilde{Z}_s^N|^2 d\langle W^N \rangle_s - \int_0^t |Z_{m_N,s}|^2 ds \right| \right) \right] \stackrel{(N \rightarrow \infty)}{\rightarrow} 0. \end{aligned}$$

In particular,

$$\sup_{t \in [0, T]} |\tilde{M}_t^N| \stackrel{(N \rightarrow \infty)}{\rightarrow} 0 \quad \text{in } L^2.$$

Moreover, it follows from Proposition 5.7 and Remark 5.8 that

$$\sup_t |Y_{m_N,t} - Y_t| \rightarrow 0 \quad \text{a.s. and} \quad \mathbb{E} \left[\int_0^T |Z_{m_N,s} - Z_s|^2 ds \right] \rightarrow 0.$$

This implies (5.6)–(5.7).

If

$$\xi = \varphi(W_{s_1}, \dots, W_{s_n}) \quad \text{and} \quad \xi^N = \varphi(W_{s_1}^N, \dots, W_{s_n}^N)$$

for a bounded, uniformly continuous function $\varphi: \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$, there exist Lipschitz-continuous functions $\varphi_m: \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$ bounded by $\|\varphi\|_\infty$ such that $\sup_{x \in \mathbb{R}^{d \times n}} |\varphi_m(x) - \varphi(x)| \leq 1/m$. Choose m_N as in the first part of the proof and set

$$\tilde{\xi}^N := \varphi_{m_N}(W_{s_1}^N, \dots, W_{s_n}^N).$$

One then obtains as above that

$$\sup_t |\tilde{Y}_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2 \text{ as } N \rightarrow \infty.$$

By Lemma 5.1, there exists an $N_0 \in \mathbb{N}$ and a constant $D \in \mathbb{R}_+$ such that for $N \geq N_0$,

$$\sup_t |Y_t^N - \tilde{Y}_t^N| \leq D \|\xi^N - \tilde{\xi}^N\|_\infty.$$

Hence,

$$\sup_t |Y_t^N - \tilde{Y}_t^N| \rightarrow 0 \quad \text{in } L^2 \text{ for } N \rightarrow \infty,$$

and one can conclude that

$$\sup_t |Y_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2 \text{ for } N \rightarrow \infty. \quad \square$$

In the following corollary, we denote by $C^d[0, T]$ the set of all continuous functions from $[0, T]$ to \mathbb{R}^d and assume that the driver f is of the form

$$f(t, y, z) = \tilde{f}(t, W, y, z) \tag{5.8}$$

for a measurable function $\tilde{f}: [0, T] \times C^d[0, T] \times \mathbb{R} \times \mathbb{R}^d \rightarrow \mathbb{R}$ that is left-continuous in t and for which there exists a $q \in [1, 2)$ such that conditions (5.9)–(5.12) are satisfied:

$$|\tilde{f}(t, w, y, z)| \leq K(1 + |y| + |z|^q) \quad \text{for all } t, w, y, z, \tag{5.9}$$

$$|\tilde{f}(t, w, y_1, z) - \tilde{f}(t, w, y_2, z)| \leq K|y_1 - y_2| \quad \text{for all } t, w, y_1, y_2, z. \tag{5.10}$$

For every $a \in \mathbb{R}_+$ there exists $b \in \mathbb{R}_+$ such that

$$|\tilde{f}(t, w, y, z_1) - \tilde{f}(t, w, y, z_2)| \leq b(1 + (|z_1| \vee |z_2|)^{q/2})|z_1 - z_2| \tag{5.11}$$

for all $t \in [0, T]$, $y \in [-a, a]$ and $z_1, z_2 \in \mathbb{R}^d$.

There exists a constant $L \in \mathbb{R}_+$ such that

$$|\tilde{f}(t, w_1, y, z) - \tilde{f}(t, w_2, y, z)| \leq L \sup_{s \leq t} |w_1(s) - w_2(s)| \quad \text{for all } t, w_1, w_2, y, z. \tag{5.12}$$

We also assume that the discrete-time drivers f^N are of the form

$$f^N(t, W^N, y, z) = \tilde{f}(t_{i+1}^N, W^{N,c}, y, z) \quad \text{for } t_i^N < t \leq t_{i+1}^N, \tag{5.13}$$

where $W^{N,c}$ is the following continuous approximation of W^N : Set $h^N = \sup_t |t_i^N - t_{i-1}^N|$ and

$$W_t^{N,c} = \begin{cases} 0, & \text{for } t \leq h^N, \\ W_{t_{i-1}^N}^N + \frac{t - (t_{i-1}^N + h^N)}{t_i^N - t_{i-1}^N} (W_{t_i^N}^N - W_{t_{i-1}^N}^N), & \text{for } t_{i-1}^N + h^N \leq t \leq t_i^N + h^N. \end{cases}$$

Note that $W^{N,c}$ is adapted to the filtration (\mathcal{F}_t^N) and $f^N(t_{i+1}^N, W^N, y, z)$ only depends on $W_{t_1^N}^N, \dots, W_{t_i^N}^N$.

Corollary 5.10. *Assume the W^N fulfill (C1), (C2) and (W1) for some $q \in [1, 2)$, but instead of (C3) they converge to W in distribution and satisfy $\sup_N \mathbb{E}[\sup_t |W_t^N|^{2+\varepsilon}] < \infty$ for some $\varepsilon > 0$. Furthermore, suppose f and f^N are of the form (5.8) and (5.13), respectively. Then for every $\xi \in L^\infty(\mathcal{F}_T)$, there exists a sequence of \mathcal{F}_T^N -measurable random variables $\tilde{\xi}^N$ bounded by $\|\xi\|_\infty$ such that for N large enough, the N th BSΔE with terminal condition $\tilde{\xi}^N$ has a unique solution $(\tilde{Y}^N, \tilde{Z}^N, \tilde{M}^N)$ and*

$$\sup_t |\tilde{Y}_t^N - Y_t| \rightarrow 0 \quad \text{in distribution for } N \rightarrow \infty,$$

where (Y, Z) is the unique solution of the BSDE (2.3) with bounded Y . In the special case, where $\xi = \varphi(W_{s_1}, \dots, W_{s_n})$ for a uniformly continuous function $\varphi: \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$, one can choose $\tilde{\xi}^N = \varphi(W_{s_1}^N, \dots, W_{s_n}^N)$.

Proof. It can be shown as in Example 2.1 that there exists a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ supporting a d -dimensional Brownian motion \tilde{W} and random walks \tilde{W}^N with the same distributions as W^N such that $\mathbb{E}[\sup_t |\tilde{W}_t^N - \tilde{W}_t|^2] \rightarrow 0$ for $N \rightarrow \infty$. Then

$$\sup_t |\tilde{f}(t, \tilde{W}^{N,c}, y, z) - \tilde{f}(t, \tilde{W}, y, z)| \rightarrow 0 \quad \text{in } L^2 \text{ for } N \rightarrow \infty$$

and it follows from Theorem 5.9 that for every $\tilde{\xi} \in L^\infty(\tilde{\mathcal{F}}_T)$ one can choose \mathcal{F}_T^N -measurable terminal conditions $\tilde{\xi}^N$ bounded by $\|\tilde{\xi}\|_\infty$ such that the corresponding solutions satisfy $\sup_t |\tilde{Y}_t^N - \tilde{Y}_t| \rightarrow 0$ in L^2 as $N \rightarrow \infty$. Furthermore, if $\tilde{\xi}$ is of the form $\tilde{\xi} = \varphi(\tilde{W}_{s_1}, \dots, \tilde{W}_{s_n})$ for a uniformly continuous function $\varphi: \mathbb{R}^{d \times n} \rightarrow \mathbb{R}$, one can choose $\tilde{\xi}^N = \varphi(\tilde{W}_{s_1}^N, \dots, \tilde{W}_{s_n}^N)$. This proves the corollary. \square

Example 5.11. In the setting of Corollary 5.10, let $\xi = \varphi(W_T)$ and $\xi^N = \varphi(W_T^N)$ for

$$\varphi(x) = \begin{cases} \sqrt{x} \wedge 1, & \text{if } x \geq 0, \\ -\sqrt{-x} \vee -1, & \text{if } x < 0. \end{cases}$$

Then for every function \tilde{f} satisfying (5.9)–(5.12) the corresponding solutions Y^N converge to Y in distribution. Let us illustrate this result for the example

$$\tilde{f}(t, w, y, z) = K_1 y + K_2 |z|^{3/2}.$$

Let $T = 1$ and W^N be the Bernoulli random walks from Example 2.1. Then the discrete equations can numerically be solved using formulas (3.1)–(3.2).

Figure 1(a) and (b) show the convergence of Y_0^N for different values of K_1 and K_2 . It can be seen that for $(K_1, K_2) = (1, 1)$, Y_0^N converges rather fast. Already for $N = 20$, it is close to the limit value. On the other hand, for $(K_1, K_2) = (1, 5)$, the convergence is much slower.




Figure 1. (a) Y_0^N corresponding to $K_1 = 1$ and $K_2 = 1$. (b) Y_0^N corresponding to $K_1 = 1$ and $K_2 = 5$.

6. Convergence results for convex drivers

In this section, we consider BSΔEs with drivers that are convex in z and use convex duality to derive stronger convergence results than in Section 5. For the case where f does not depend on y it has been shown in Barrieu and El Karoui [2], Delbaen *et al.* [14] and Delbaen *et al.* [13] that BSDEs with convex drivers admit a convex dual representation. Here, we establish convex dual representations for solutions of BSΔEs and use them to show convergence. We need the following stronger version of condition (W1) on the approximating processes W^N :

$$(W2) \quad \mathbb{E}[\Delta W_{t_i^N}^{N,k} \Delta W_{t_i^N}^{N,l}] = 0 \text{ for all } N \in \mathbb{N}, i = 1, \dots, i_N, k \neq l \text{ and}$$

$$\sup_{N,i,k} \frac{\|\Delta W_{t_i^N}^{N,k}\|_\infty}{\sqrt{\Delta \langle W^N \rangle_{t_i^N}}} < \infty.$$

Note that this implies (W1) for all $q \in [1, 2)$. In the following, we assume that the drivers f^N are convex in z and define

$$g^N(t, y, \mu) := \operatorname{ess\,sup}_z \{ \mu z - f^N(t, y, z) \}, \quad \mu \in \mathbb{R}^d.$$

Let μ^N be an \mathbb{R}^d -valued (\mathcal{F}_i^N) -adapted process that is constant on the intervals $(t_{i-1}^N, t_i^N]$ and satisfies

$$\mu_{t_i^N}^N \Delta W_{t_i^N}^N > -1 \quad \text{for all } i. \tag{6.1}$$

Then

$$\frac{d\mathbb{P}^{\mu^N}}{d\mathbb{P}} = \prod_{i=1}^{i_N} (1 + \mu_{t_i^N}^N \Delta W_{t_i^N}^N) \tag{6.2}$$

defines a probability measure \mathbb{P}^{μ^N} equivalent to \mathbb{P} under which the processes

$$W_{t_i^N}^{N,\mu^N,k} = W_{t_i^N}^{N,k} - \sum_{j=1}^i \mu_{t_j^N}^k \Delta \langle W^N \rangle_{t_j^N}, \quad k = 1, \dots, d,$$

are martingales. The following proposition gives an implicit dual representation of solutions of BSΔEs. Its proof can be found in the [Appendix](#).

Proposition 6.1. *Assume (W2) and let $C, K, L \in \mathbb{R}_+, q \in [1, 2)$ be constants such that all terminal conditions ξ^N and drivers f^N fulfill the following conditions:*

- (i) $\|\xi^N\|_\infty \leq C$;
- (ii) f^N is convex in z ;
- (iii) $|f^N(t, y, z)| \leq K(1 + |y| + |z|^q)$ for all $(t, y, z) \in [0, T] \times \mathbb{R}^{d+1}$;
- (iv) $|f^N(t, y_1, z) - f^N(t, y_2, z)| \leq L|y_1 - y_2|$ for all $(t, y, z) \in [0, T] \times \mathbb{R}^{d+1}$;
- (v) $|f^N(t, y, z_1) - f^N(t, y, z_2)| \leq L(1 + (|z_1| \vee |z_2|)^{q/2})|z_1 - z_2|$ for all $(t, y, z_1, z_2) \in [0, T] \times \mathbb{R}^{2d+1}$ such that $|y| \leq (C + 1)\exp(KT)$.

Then there exists $N_0 \in \mathbb{N}$ such that for every $N \geq N_0$, the N th BSΔE has a unique solution (Y^N, Z^N, M^N) and Y^N can be represented as

$$Y_{t_i^N}^N = \operatorname{ess\,sup}_{\mu^N} \mathbb{E}^{\mu^N} \left[\xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \mu_{t_j^N}^N) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right], \tag{6.3}$$

where the essential supremum is taken over all \mathbb{R}^d -valued (\mathcal{F}_t^N) -adapted processes μ^N that are constant on the intervals $(t_{i-1}^N, t_i^N]$ and satisfy (6.1). Moreover, there exists a constant $R \in \mathbb{R}_+$ such that for each $N \geq N_0$, (6.3) admits a maximizer $\hat{\mu}^N$ satisfying

$$\mathbb{E}^{\hat{\mu}^N} \left[\sum_{j=i+1}^{i_N} |\hat{\mu}_{t_j^N}^N|^2 \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right] \leq R \quad \text{for all } i \leq i_N - 1. \tag{6.4}$$

We are now ready to prove our convergence result for convex drivers. It states that for any sequence of bounded discrete-time terminal conditions converging to ξ and every sequence of discrete-time drivers converging to f the discrete-time solutions Y^N converge to the continuous-time solution Y .

Theorem 6.2. *Assume (W2), the $f^N(t, y, z)$ are convex in z and one has $\sup_N \|\xi^N\|_\infty < \infty$ as well as $\xi^N \rightarrow \xi$ in L^2 . Moreover, suppose the f^N satisfy (f1)–(f5). Then for N large enough, the*

N th BS ΔE has a unique solution (Y^N, Z^N, M^N) and

$$\sup_t |Y_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2 \text{ for } N \rightarrow \infty,$$

where (Y, Z) is the unique solution of the BSDE (2.3) with bounded Y .

Proof. By Theorem 5.9, there exist \mathcal{F}_T^N -measurable terminal conditions $\tilde{\xi}^N$ bounded by $C := \sup_N \|\tilde{\xi}^N\|_\infty$ such that the corresponding solutions satisfy

$$\sup_t |\tilde{Y}_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2.$$

Choose $b \in \mathbb{R}_+$ such that condition (f3) holds for $a = (C + 1) \exp(KT)$. Then the conditions of Theorem 4.2 and Proposition 6.1 are satisfied with $L = K \vee b$. Hence, there exists $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$, $\sup_t |Y_t^N|$ and $\sup_t |\tilde{Y}_t^N|$ are bounded by $(C + 1) \exp(KT)$ and

$$\begin{aligned} Y_{t_i^N}^N &= \operatorname{ess\,sup}_\mu \mathbb{E}^\mu \left[\xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \mu_{t_j^N}) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right] \\ &= \mathbb{E}^{\hat{\mu}^N} \left[\xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \hat{\mu}_{t_j^N}^N) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right] \end{aligned}$$

as well as

$$\begin{aligned} \tilde{Y}_{t_i^N}^N &= \operatorname{ess\,sup}_\mu \mathbb{E}^\mu \left[\tilde{\xi}^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, \tilde{Y}_{t_{j-1}^N}^N, \mu_{t_j^N}) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right] \\ &= \mathbb{E}^{\tilde{\mu}^N} \left[\tilde{\xi}^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, \tilde{Y}_{t_{j-1}^N}^N, \tilde{\mu}_{t_j^N}^N) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right]. \end{aligned}$$

If we can show

$$\sup_t |\tilde{Y}_t^N - Y_t^N| \rightarrow 0 \quad \text{in } L^2,$$

we get

$$\sup_t |Y_t^N - Y_t| \rightarrow 0 \quad \text{in } L^2,$$

and the theorem is proved. As the supremum of K -Lipschitz functions, g^N is again K -Lipschitz in y . Hence, since $|\max\{a_1, a_2\} - \max\{b_1, b_2\}| \leq \max\{|a_1 - b_1|, |a_2 - b_2|\}$ for $a_1, a_2, b_1, b_2 \in \mathbb{R}$, and

$$Y_{t_i^N}^N = \max_{\mu \in \{\hat{\mu}^N, \tilde{\mu}^N\}} \mathbb{E}^\mu \left[\xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \mu_{t_j^N}) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right],$$

$$\tilde{Y}_i^N = \max_{\mu \in \{\hat{\mu}^N, \tilde{\mu}^N\}} \mathbb{E}^\mu \left[\tilde{\xi}^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, \tilde{Y}_{t_{j-1}^N}, \mu_{t_j^N}) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right],$$

one obtains

$$\begin{aligned} |\tilde{Y}_{t_i^N}^N - Y_{t_i^N}^N| &\leq \max_{\mu \in \{\hat{\mu}^N, \tilde{\mu}^N\}} \mathbb{E}^\mu \left[|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right] \\ &\leq \mathbb{E}^{\hat{\mu}^N} \left[|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right] \\ &\quad + \mathbb{E}^{\tilde{\mu}^N} \left[|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right]. \end{aligned}$$

From Proposition 6.1, we know that there exists a constant $R \in \mathbb{R}_+$ such that

$$\mathbb{E}^{\hat{\mu}^N} \left[\sum_{j=i+1}^{i_N} |\hat{\mu}_{t_j^N}^N|^2 |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right] \leq R \quad \text{for all } N \geq N_0 \text{ and } i = 0, \dots, i_N - 1.$$

Consequently, we obtain from Lemma A.3 in the Appendix that there exists a constant \tilde{R} such that

$$\mathbb{E} \left[\varphi \left(\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \right) \middle| \mathcal{F}_{t_i^N}^N \right] \leq \tilde{R} \quad \text{for all } N \geq N_0 \text{ and } i = 0, \dots, i_N - 1,$$

where $\varphi(x) = x \log(x) \vee 1$. Fix $\varepsilon > 0$ and set $D = 2[C + (C + 1) \exp(KT)K \sup_{N \geq N_0} \langle W^N \rangle_T]$. Since $\varphi(x)/x \uparrow \infty$, there exists $B \in \mathbb{R}_+$ such that for all $x > B$,

$$\frac{x}{\varphi(x)} \leq \frac{\varepsilon}{\tilde{R}D}.$$

Introduce the sets $E_{i+1}^N = \{\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) > B\}$. Then

$$\begin{aligned} &\sup_{N \geq N_0, 0 \leq i \leq i_N - 1} \mathbb{E} \left[1_{E_{i+1}^N} \prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \middle| \mathcal{F}_{t_i^N}^N \right] \\ &= \sup_{N \geq N_0, 0 \leq i \leq i_N - 1} \mathbb{E} \left[1_{E_{i+1}^N} \frac{\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N)}{\varphi(\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N))} \right] \end{aligned} \tag{6.5}$$

$$\begin{aligned} & \times \varphi \left(\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \middle| \mathcal{F}_{t_i^N}^N \right) \\ & \leq \frac{\varepsilon}{\tilde{R}D} \sup_{N \geq N_0, 0 \leq i \leq i_N - 1} \mathbb{E} \left[\varphi \left(\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \middle| \mathcal{F}_{t_i^N}^N \right) \right] \leq \frac{\varepsilon}{D}. \end{aligned}$$

This yields for all $N \geq N_0$,

$$\begin{aligned} & \mathbb{E}^{\hat{\mu}^N} \left[|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right] \\ & = \mathbb{E} \left[\prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \left(|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right) \right] \\ & = \mathbb{E} \left[1_{E_{i+1}^N} \prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \left(|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right) \right] \\ & \quad + \mathbb{E} \left[1_{E_{i+1}^{N,c}} \prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \left(|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right) \right] \\ & \leq D \mathbb{E} \left[1_{E_{i+1}^N} \prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \middle| \mathcal{F}_{t_i^N}^N \right] \\ & \quad + \mathbb{E} \left[1_{E_{i+1}^{N,c}} \prod_{j=i+1}^{i_N} (1 + \hat{\mu}_{t_j^N}^N \Delta W_{t_j^N}^N) \right. \\ & \quad \left. \times \left(|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right) \right] \\ & \leq \varepsilon + B \mathbb{E} \left[|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right]. \end{aligned}$$

In the first inequality, we used that the random variables $|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}|$ are uniformly bounded by D . In the second inequality, we used (6.5) and the definition of the sets E_{i+1}^N . Using the same estimate for $\tilde{\mu}^N$ instead of $\hat{\mu}^N$ gives

$$|\tilde{Y}_{t_i^N}^N - Y_{t_i^N}^N| \leq 2\varepsilon + 2B \mathbb{E} \left[|\tilde{\xi}^N - \xi^N| + K \sum_{j=i+1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| |\Delta \langle W^N \rangle_{t_j^N}| \middle| \mathcal{F}_{t_i^N}^N \right]. \tag{6.6}$$

Taking expectations, one gets

$$\mathbb{E}[|\tilde{Y}_{t_i^N}^N - Y_{t_i^N}^N|] \leq 2\varepsilon + 2B\mathbb{E}[|\tilde{\xi}^N - \xi^N|] + K \sum_{j=i+1}^{i_N} \mathbb{E}[|\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N|] \Delta\langle W^N \rangle_{t_j^N}.$$

Since $\varepsilon > 0$ was arbitrary, one obtains from a discrete version of Gronwall’s lemma (see Lemma A.4 in the Appendix) that

$$\sup_t \mathbb{E}[|\tilde{Y}_t^N - Y_t^N|] \rightarrow 0 \quad \text{as } N \rightarrow \infty,$$

and since Y^N and \tilde{Y}^N are both bounded by $(C + 1) \exp(KT)$, also

$$\sup_t \mathbb{E}[|\tilde{Y}_t^N - Y_t^N|^2] \rightarrow 0 \quad \text{as } N \rightarrow \infty. \tag{6.7}$$

It remains to show that \sup_t can be taken inside of the expectation in (6.7). To do this, note that (6.6) gives

$$\sup_i |\tilde{Y}_{t_i^N}^N - Y_{t_i^N}^N| \leq 2\varepsilon + 2B \left(\sup_i \mathbb{E}[|\tilde{\xi}^N - \xi^N| | \mathcal{F}_{t_i^N}^N] + K \sup_i A_{t_i^N}^N \right)$$

for the nonnegative martingale

$$A_{t_i^N}^N = \mathbb{E} \left[\sum_{j=1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| \Delta\langle W^N \rangle_{t_j^N} \mid \mathcal{F}_{t_i^N}^N \right], \quad i = 0, \dots, i_N.$$

Since ε was arbitrary, and $\sup_i \mathbb{E}[|\tilde{\xi}^N - \xi^N| | \mathcal{F}_{t_i^N}^N] \xrightarrow{(N \rightarrow \infty)} 0$ in L^2 by Doob’s maximal inequality, the only thing left to show is $\sup_i A_{t_i^N}^N \xrightarrow{(N \rightarrow \infty)} 0$ in L^2 . Applying Doob’s maximal inequality to A^N yields

$$\begin{aligned} \mathbb{E} \left[\sup_i |A_{t_i^N}^N|^2 \right] &\leq 2\mathbb{E} \left[\left(\sum_{j=1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N| \Delta\langle W^N \rangle_{t_j^N} \right)^2 \right] \\ &\leq 2\langle W^N \rangle_T \mathbb{E} \left[\sum_{j=1}^{i_N} |\tilde{Y}_{t_{j-1}^N}^N - Y_{t_{j-1}^N}^N|^2 \Delta\langle W^N \rangle_{t_j^N} \right] \\ &\leq 2(\langle W^N \rangle_T)^2 \sup_t \mathbb{E}[|\tilde{Y}_t^N - Y_t^N|^2] \rightarrow 0 \quad \text{as } N \rightarrow \infty, \end{aligned}$$

where we used Jensen’s inequality for the second inequality and (6.7) for the convergence in the last line. This proves the theorem. □

If one has convergence of (W^N, ξ^N) to (W, ξ) in distribution instead of L^2 together with

$$\sup_N \mathbb{E} \left[\sup_t |W_t^N|^{2+\varepsilon} \right] < \infty \quad \text{and} \quad \sup_N \|\xi^N\|_\infty < \infty,$$

one can show as in Example 2.1 that there exists a probability space $(\tilde{\Omega}, \tilde{\mathcal{F}}, \tilde{\mathbb{P}})$ carrying $(\tilde{W}^N, \tilde{\xi}^N)$ distributed as (W^N, ξ^N) and $(\tilde{W}, \tilde{\xi})$ distributed as (W, ξ) such that

$$\mathbb{E} \left[\sup_t |\tilde{W}_t^N - \tilde{W}_t|^2 \right] \rightarrow 0 \quad \text{and} \quad \mathbb{E} [|\tilde{\xi}^N - \tilde{\xi}|^2] \rightarrow 0 \quad \text{for } N \rightarrow \infty.$$

In the case where the drivers f and f^N are given as in (5.8) and (5.13), the following holds.

Corollary 6.3. *Assume the W^N fulfill (C1), (C2) and (W2), but instead of (C3), (W^N, ξ^N) converges in distribution to (W, ξ) and one has $\sup_N \mathbb{E}[\sup_t |W_t^N|^{2+\varepsilon}] < \infty$ for some $\varepsilon > 0$ and $\sup_N \|\xi^N\|_\infty < \infty$. Furthermore, suppose f and f^N are of the form (5.8) and (5.13), respectively. Then for N large enough, the N th BSΔE has a unique solution (Y^N, Z^N, M^N) and*

$$\sup_t |Y_t^N - Y_t| \rightarrow 0 \quad \text{in distribution for } N \rightarrow \infty,$$

where (Y, Z) is the unique solution of the BSDE (2.3) with bounded Y .

Appendix

A.1. Proofs of Section 3

Proof of Lemma 3.1. If (Y^N, Z^N, M^N) is a solution of the N th BSΔE, then

$$Y_{t_i^N}^N - f^N(t_{i+1}^N, Y_{t_i^N}^N, Z_{t_{i+1}^N}^N) \Delta \langle W^N \rangle_{t_{i+1}^N} + Z_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^N + \Delta M_{t_{i+1}^N}^N = Y_{t_{i+1}^N}^N. \tag{A.1}$$

Taking conditional expectations on both sides with respect to $\mathcal{F}_{t_i^N}^N$ gives (3.1). Multiplying both sides of (A.1) with $\Delta W_{t_{i+1}^N}^{N,k}$ and taking conditional expectations with respect to $\mathcal{F}_{t_i^N}^N$ yields (3.2). Finally, (3.3) is a consequence of (3.1) and (A.1). □

Proof of Proposition 3.2. We prove the proposition by backwards induction. Set $Y_T^N = \xi^N$, which by assumption (C4) is bounded. Now assume that there exist i and (Y_i^N, Z_i^N, M_i^N) solving the BSΔE (2.4) for $t \in [t_{i+1}^N, T]$ such that (Y_i^N) and (Z_i^N) are bounded. By Lemma 3.1, $Z_{t_{i+1}^N}^{N,k}$ must be of the form

$$Z_{t_{i+1}^N}^{N,k} = \frac{\mathbb{E}[Y_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^{N,k} | \mathcal{F}_{t_i^N}^N]}{\Delta \langle W^N \rangle_{t_{i+1}^N}}.$$

Since by induction hypothesis, $Y_{t_{i+1}}^N$ is bounded, $Z_{t_{i+1}}^{N,k}$ is well-defined and bounded. Next, we try to find $Y_{t_i}^N \in L^\infty(\mathcal{F}_{t_i}^N)$ such that

$$Y_{t_i}^N - f^N(t_{i+1}, Y_{t_i}^N, Z_{t_{i+1}}^N) \Delta \langle W^N \rangle_{t_{i+1}} = \mathbb{E}[Y_{t_{i+1}}^N | \mathcal{F}_{t_i}^N]. \quad (\text{A.2})$$

To do that, we introduce the mapping $A(\omega, y) := y - f(t_{i+1}, y, Z_{t_{i+1}}^N) \Delta \langle W^N \rangle_{t_{i+1}}$. It is $\mathcal{F}_{t_i}^N$ -measurable in ω and continuous in y . Moreover, it satisfies

$$y - \kappa(1 + |y| + g(Z_{t_{i+1}}^N)) \leq A(\omega, y) \leq y + \kappa(1 + |y| + g(Z_{t_{i+1}}^N)) \quad (\text{A.3})$$

for $\kappa = K \max_i \Delta \langle W^N \rangle_{t_i} < 1$. So it follows from Lemma A.1 below that there exists an $\mathcal{F}_{t_i}^N \otimes \mathcal{B}(\mathbb{R})$ -measurable function $B : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ such that $A(\omega, B(\omega, y)) = y$ for all $(\omega, y) \in \Omega \times \mathbb{R}$. Thus,

$$Y_{t_i}^N = B(\omega, \mathbb{E}[Y_{t_{i+1}}^N | \mathcal{F}_{t_i}^N]) \in L^0(\mathcal{F}_{t_i}^N)$$

solves (A.2), and since $Y_{t_{i+1}}^N$ and $Z_{t_{i+1}}^N$ are bounded, it follows from the estimate (A.3) that the same is true for $Y_{t_i}^N$. Finally, $M_0^N = 0$ and

$$\begin{aligned} \Delta M_{t_{i+1}}^N &= \Delta Y_{t_{i+1}}^N + f^N(t_{i+1}, Y_{t_i}^N, Z_{t_{i+1}}^N) \Delta \langle W^N \rangle_{t_{i+1}} - Z_{t_{i+1}}^N \Delta W_{t_{i+1}}^N \\ &= Y_{t_{i+1}}^N - \mathbb{E}[Y_{t_{i+1}}^N | \mathcal{F}_{t_i}^N] - Z_{t_{i+1}}^N \Delta W_{t_{i+1}}^N \end{aligned}$$

defines a square-integrable martingale M^N orthogonal to W^N which is bounded if W^N is so. This completes the proof. \square

Lemma A.1. *Let \mathcal{G} be a sub- σ -algebra of \mathcal{F} and $A : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ a function that is \mathcal{G} -measurable in $\omega \in \Omega$ and continuous in $y \in \mathbb{R}$. Assume that for every $\omega \in \Omega$, the set $\{y \in \mathbb{R} : A(\omega, y) \in C\}$ is nonempty and bounded for each nonempty bounded subset C of \mathbb{R} . Then there exists a $\mathcal{G} \otimes \mathcal{B}(\mathbb{R})$ -measurable function $B : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ such that $A(\omega, B(\omega, x)) = x$ for all $x \in \mathbb{R}$.*

Proof. For all $k, l \in \mathbb{N}$,

$$b_{kl}(\omega) = \inf\{y \in \mathbb{R} : A(\omega, y) \in ((k-1)2^{-l}, k2^{-l}]\}$$

is a \mathcal{G} -measurable mapping from Ω to \mathbb{R} and

$$B_l(\omega, x) = \sum_{k \in \mathbb{Z}} b_{kl}(\omega) 1_{\{(k-1)2^{-l} < x \leq k2^{-l}\}}$$

a $\mathcal{G} \otimes \mathcal{B}(\mathbb{R})$ -measurable map from $\Omega \times \mathbb{R}$ to \mathbb{R} such that

$$B_l(\omega, x) \rightarrow B(\omega, x) \quad \text{as } l \rightarrow \infty$$

for a $\mathcal{G} \otimes \mathcal{B}(\mathbb{R})$ -measurable function $B : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$. Since $y \mapsto A(\omega, y)$ is continuous for all $\omega \in \Omega$, one obtains

$$A(\omega, B(\omega, x)) = A\left(\omega, \lim_{l \rightarrow \infty} B_l(\omega, x)\right) = \lim_{l \rightarrow \infty} A(\omega, B_l(\omega, x)) = x$$

for all $x \in \mathbb{R}$. □

A.2. Proofs of Lemmas 4.3 and 4.4

To prove Lemmas 4.3 and 4.4, we need the following lemma.

Lemma A.2. *Assume the N th driver and terminal condition are of the special form*

$$f^N(t, y, z) = K(1 + |y| + g(z)) \quad \text{and} \quad \xi^N = C$$

for constants $C, K \in \mathbb{R}_+$ and a measurable function $g : \mathbb{R}^d \rightarrow \mathbb{R}$ with $g(0) = 0$. Then for all $N \in \mathbb{N}$ such that $\max_i \Delta \langle W^N \rangle_{t_i^N} < 1/K$, the N th BSΔE has a unique solution (Y^N, Z^N, M^N) given by

$$Y_T^N = C, \quad Y_{t_i^N}^N = \frac{Y_{t_{i+1}^N}^N + K \Delta \langle W^N \rangle_{t_{i+1}^N}}{1 - K \Delta \langle W^N \rangle_{t_{i+1}^N}}, \quad Z_{t_i^N}^N = 0, \quad M_{t_i^N}^N = 0. \quad (\text{A.4})$$

In particular, Y^N is deterministic and for $N \rightarrow \infty$, converges uniformly to the function

$$(C + 1) \exp(K(T - t)) - 1.$$

Proof. Since the terminal condition and the increments $\Delta \langle W^N \rangle_{t_i^N}$ are deterministic, Z^N and M^N are both zero and Y^N solves

$$Y_{t_i^N}^N = Y_{t_{i+1}^N}^N + K(1 + |Y_{t_i^N}^N|) \Delta \langle W^N \rangle_{t_{i+1}^N}, \quad Y_T^N = C. \quad (\text{A.5})$$

This shows (A.4). Moreover, since (A.5) are deterministic difference equations with Lipschitz coefficients, one obtains from Theorem 2.2 that their solutions converge uniformly to the solution of the ordinary differential equation

$$y'(t) = -K(1 + |y(t)|), \quad y(T) = C,$$

given by

$$y(t) = (C + 1) \exp(K(T - t)) - 1. \quad \square$$

Proof of Lemma 4.3. Since $\max_i \Delta \langle W^N \rangle_{t_i^N} < 1/K$ for N large enough, it follows from Lemma A.2 that there exists an $N_1 \geq 1$ such that for all $N \geq N_1$, the BSΔE with driver

$\hat{f}^N(t, y, z) = K(1 + |y| + |z|^q)$ and terminal condition $\hat{\xi}^N = C$ has a deterministic solution \hat{Y}^N that is bounded by $(C + 1)\exp(K(T - t))$. Set $D = 2(C + 1)\exp(KT)$. Since $q < 2$, one obtains from condition (C5) that there exists $N_0 \geq N_1$ such that

$$\sup_{N \geq N_0} \max_i K(1 + d^{q/2} D^q [\Delta \langle W^N \rangle_{t_i^N}]^{-q/2}) \Delta \langle W^N \rangle_{t_i^N} < 1 \quad (\text{A.6})$$

and

$$\sup_{N \geq N_0} \max_{i,k} dq K(1 + d^{q/4} D^{q/2} [\Delta \langle W^N \rangle_{t_i^N}]^{-q/4}) \|\Delta W_{t_i^N}^{N,k}\|_\infty \leq 1. \quad (\text{A.7})$$

Fix $N \geq N_0$ and let $f_1^N \geq f_2^N$ be drivers and $\xi_1^N \geq \xi_2^N$ terminal conditions satisfying assumptions (i)–(iv) of Lemma 4.3. By Proposition 3.2, both BSΔEs have a solution (Y_m^N, Z_m^N, M_m^N) , $m = 1, 2$, and (4.3) clearly holds at the final time T . We now go backwards in time and assume (4.3) is true on $[t_{i+1}, T]$. Then

$$(C + 1)\exp(K(T - t_{i+1}^N)) \geq Y_{1,t_{i+1}^N}^N \geq Y_{2,t_{i+1}^N}^N \geq -(C + 1)\exp(K(T - t_{i+1}^N)). \quad (\text{A.8})$$

By Lemma 3.1, one has

$$Z_{m,t_{i+1}^N}^{N,k} = \frac{\mathbb{E}[Y_{m,t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^{N,k} | \mathcal{F}_{t_i^N}^N]}{\Delta \langle W^N \rangle_{t_{i+1}^N}}, \quad (\text{A.9})$$

and

$$Y_{m,t_i^N}^N = \mathbb{E}[Y_{m,t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N] + f_m^N(t_{i+1}^N, Y_{m,t_{i+1}^N}^N, Z_{m,t_{i+1}^N}^N) \Delta \langle W^N \rangle_{t_{i+1}^N}.$$

Set

$$Y_t^N := Y_{1,t}^N - Y_{2,t}^N, \quad Z_t^N := Z_{1,t}^N - Z_{2,t}^N.$$

By (A.8), $Y_{1,t_{i+1}^N}^N$, $Y_{2,t_{i+1}^N}^N$ and $Y_{t_{i+1}^N}^N$ are bounded by D and

$$Y_{t_i^N}^N = \mathbb{E}[Y_{t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N] + (\alpha + Y_{t_i^N}^N \beta + Z_{t_i^N}^N \gamma) \Delta \langle W^N \rangle_{t_{i+1}^N}$$

for

$$\alpha = f_1^N(t_{i+1}^N, Y_{2,t_{i+1}^N}^N, Z_{2,t_{i+1}^N}^N) - f_2^N(t_{i+1}^N, Y_{2,t_{i+1}^N}^N, Z_{2,t_{i+1}^N}^N),$$

$$\beta = \frac{1}{Y_{t_i^N}^N} (f_1^N(t_{i+1}^N, Y_{1,t_{i+1}^N}^N, Z_{2,t_{i+1}^N}^N) - f_1^N(t_{i+1}^N, Y_{2,t_{i+1}^N}^N, Z_{2,t_{i+1}^N}^N)),$$

$$\begin{aligned} \gamma^k &= \frac{1}{Z_{t_{i+1}^N}^{N,k}} (f_1^N(t_{i+1}^N, Y_{1,t_{i+1}^N}^N, Z_{1,t_{i+1}^N}^{N,1}, \dots, Z_{1,t_{i+1}^N}^{N,k}, Z_{2,t_{i+1}^N}^{N,k+1}, \dots, Z_{2,t_{i+1}^N}^{N,d}) \\ &\quad - f_1^N(t_{i+1}^N, Y_{1,t_{i+1}^N}^N, Z_{1,t_{i+1}^N}^{N,1}, \dots, Z_{1,t_{i+1}^N}^{N,k-1}, Z_{2,t_{i+1}^N}^{N,k}, \dots, Z_{2,t_{i+1}^N}^{N,d})). \end{aligned}$$

It can be seen from (A.9) that for $m = 1, 2$,

$$|Z_{m,t_{i+1}^N}^N|^2 = \sum_{k=1}^d (Z_{m,t_{i+1}^N}^{N,k})^2 \leq \sum_{k=1}^d \frac{\mathbb{E}[(Y_{m,t_{i+1}^N}^N)^2 | \mathcal{F}_{t_i^N}^N] \mathbb{E}[(\Delta W_{t_{i+1}^N}^{N,k})^2]}{(\Delta \langle W^N \rangle_{t_{i+1}^N})^2} \leq \frac{dD^2}{\Delta \langle W^N \rangle_{t_{i+1}^N}}. \tag{A.10}$$

So by assumption (iii) and (A.6),

$$\begin{aligned} |\beta \Delta \langle W^N \rangle_{t_{i+1}^N}| &\leq K(1 + |Z_{2,t_{i+1}^N}^N|^q) \Delta \langle W^N \rangle_{t_{i+1}^N} \\ &\leq K(1 + d^{q/2} D^q [\Delta \langle W^N \rangle_{t_{i+1}^N}]^{-q/2}) \Delta \langle W^N \rangle_{t_{i+1}^N} < 1. \end{aligned}$$

Hence,

$$Y_{i+1}^N = \frac{\mathbb{E}[Y_{t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N] + (\alpha + Z_{t_{i+1}^N}^N \gamma) \Delta \langle W^N \rangle_{t_{i+1}^N}}{1 - \beta \Delta \langle W^N \rangle_{t_{i+1}^N}}. \tag{A.11}$$

From assumption (iv) and (A.10) one obtains

$$|\gamma| \leq d^{1/2} q K (1 + (|Z_{1,t_{i+1}^N}^N| \vee |Z_{2,t_{i+1}^N}^N|)^{q/2}) \leq d^{1/2} q K (1 + d^{q/4} D^{q/2} (\Delta \langle W^N \rangle_{t_{i+1}^N})^{-q/4})$$

and from (A.9),

$$|Z_{t_{i+1}^N}^N| \leq d^{1/2} \max_k \frac{\|\Delta W_{t_{i+1}^N}^{N,k}\|_\infty}{\Delta \langle W^N \rangle_{t_{i+1}^N}} \mathbb{E}[|Y_{t_{i+1}^N}^N| | \mathcal{F}_{t_i^N}^N].$$

By (A.7), this yields

$$\begin{aligned} |Z_{t_{i+1}^N}^N \gamma| \Delta \langle W^N \rangle_{t_{i+1}^N} &\leq |Z_{t_{i+1}^N}^N| |\gamma| \Delta \langle W^N \rangle_{t_{i+1}^N} \\ &\leq dqK (1 + d^{q/4} D^{q/2} (\Delta \langle W^N \rangle_{t_{i+1}^N})^{-q/4}) \max_k \|\Delta W_{t_{i+1}^N}^{N,k}\|_\infty \mathbb{E}[|Y_{t_{i+1}^N}^N| | \mathcal{F}_{t_i^N}^N] \\ &\leq \mathbb{E}[|Y_{t_{i+1}^N}^N| | \mathcal{F}_{t_i^N}^N]. \end{aligned}$$

Since $Y_{t_{i+1}^N}^N \geq 0$ and $\alpha \geq 0$, it follows from (A.11) that $Y_{1,t_i^N}^N - Y_{2,t_i^N}^N = Y_{t_i^N}^N \geq 0$. Now observe that \hat{f}^N satisfies assumptions (ii)–(iv). So the same argument applied to the equations corresponding to (\hat{f}^N, C) and (f_1^N, ξ^N) gives

$$(C + 1) \exp(K(T - t_i^N)) \geq \hat{Y}_{t_i^N}^N \geq Y_{1,t_i^N}^N.$$

Analogously, one deduces

$$Y_{2,t_i^N}^N \geq (C + 1) \exp(K(T - t_i^N)),$$

and the induction step is complete. □

Proof of Lemma 4.4. For N large enough, one has

$$\max_i \Delta \langle W^N \rangle_{t_i^N} < 1/K. \quad (\text{A.12})$$

So it follows from Lemma A.2 that there exists $N_1 \in \mathbb{N}$ such that for all $N \geq N_1$, the BSΔE with driver $\hat{f}^N(t, y, z) = K(1 + |y| + |z|^q)$ and terminal condition $\hat{\xi}^N = C$ has a deterministic solution \hat{Y}^N dominated by $(C + 1)\exp(K(T - t))$. Choose $N_0 \geq N_1$ such that for all $N \geq N_0$, the statement of Lemma 4.3 holds for all terminal conditions bounded by $(C + 1)\exp(KT)$ and drivers satisfying conditions (ii)–(iv) of Lemma 4.3. Now fix $N \geq N_0$ and assume (Y^N, Z^N, M^N) is a solution corresponding to ξ^N and f^N satisfying conditions (i) and (ii) of Lemma 4.4. Since $\hat{Y}_t^N \leq (C + 1)\exp(K(T - t))$, it is enough to show that

$$\hat{Y}_{t_i^N}^N \geq Y_{t_i^N}^N \geq -\hat{Y}_{t_i^N}^N \quad \text{for all } i. \quad (\text{A.13})$$

By condition (i), (A.13) holds for $t = T$. For $t < T$ we argue by backwards induction. So let us assume that (A.13) holds for $t = t_{i+1}^N$. We will only show $\hat{Y}_{t_i^N}^N \geq Y_{t_i^N}^N$. The second inequality in (A.13) follows analogously. From Lemma 3.1, we know that

$$Z_{t_{i+1}^N}^{N,k} = \frac{\mathbb{E}[Y_{t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^{N,k} | \mathcal{F}_{t_i^N}^N]}{\Delta \langle W^N \rangle_{t_{i+1}^N}}$$

and

$$A(\omega, Y_{t_i^N}^N) = \mathbb{E}[Y_{t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N],$$

where $A(\omega, y) = y - f(t_{i+1}^N, y, Z_{t_{i+1}^N}^{N,k}) \Delta \langle W^N \rangle_{t_{i+1}^N}$. Consider the BSΔE with driver

$$\tilde{f}^N(t_j^N, y, z) = \begin{cases} K(1 + |y| + |z|^q), & \text{for } j = i + 1, \\ 0, & \text{for } j \neq i + 1 \end{cases}$$

and terminal condition $Y_{t_{i+1}^N}^N$. By Lemma 4.3, it has a unique solution $(\tilde{Y}^N, \tilde{Z}^N, \tilde{M}^N)$, and it is easy to see that $\tilde{Y}_{t_{i+1}^N}^N = Y_{t_{i+1}^N}^N$. Due to (A.12), the mapping $\tilde{A}(\omega, y) = y - \tilde{f}(t_{i+1}^N, y, Z_{t_{i+1}^N}^{N,k}) \times \Delta \langle W^N \rangle_{t_{i+1}^N}$ is strictly increasing in y and since $\tilde{f}^N(t_{i+1}^N, \cdot, \cdot) \geq f^N(t_{i+1}^N, \cdot, \cdot)$, one has

$$\tilde{A}(\omega, \tilde{Y}_{t_i^N}^N) = \mathbb{E}[Y_{t_{i+1}^N}^N | \mathcal{F}_{t_i^N}^N] = A(\omega, Y_{t_i^N}^N) \geq \tilde{A}(\omega, Y_{t_i^N}^N).$$

This shows $\tilde{Y}_{t_i^N}^N \geq Y_{t_i^N}^N$. To conclude the proof, consider the solution \bar{Y}^N of the BSΔE with driver \tilde{f}^N and terminal condition $\hat{Y}_{t_{i+1}^N}^N$. Then $\bar{Y}_{t_i^N}^N = \hat{Y}_{t_i^N}^N$ and Lemma 4.3 yields $\bar{Y}_{t_i^N}^N \geq \tilde{Y}_{t_i^N}^N$. Consequently,

$$\hat{Y}_{t_i^N}^N = \bar{Y}_{t_i^N}^N \geq \tilde{Y}_{t_i^N}^N \geq Y_{t_i^N}^N,$$

which completes the induction step. \square

A.3. Remaining proofs of Section 5

Proof of Lemma 5.1. Set $\tilde{C} = 3C$ and $\tilde{K} = 2K(2C + K + 1)(\exp(KT) + 1)(T + 1)$. Choose $b \in \mathbb{R}_+$ such that condition (f3) holds for $a = (\tilde{C} + 1)\exp(\tilde{K}T)$. It follows from (2.1) that $\prod_{i=1}^{i_N} (1 - K\Delta\langle W^N \rangle_{t_i^N}) \rightarrow \exp(-KT)$ for $N \rightarrow \infty$. So there exists $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$,

$$\prod_{i=1}^{i_N} (1 - K\Delta\langle W^N \rangle_{t_i^N})^{-1} \leq \exp(KT) + 1, \quad \langle W^N \rangle_T \leq T + 1$$

and the statement of Theorem 4.2 holds for \tilde{C} instead of C , \tilde{K} instead of K and $L = K \vee b$. Set $D = (\exp(KT) + 1)(T + 1)$ and fix $N \geq N_0$ as well as terminal conditions ξ_1^N, ξ_2^N bounded by C and drivers f_1^N, f_2^N satisfying (f1)–(f3) such that $\|f_1^N - f_2^N\|_\infty \leq K$. Then the parameter pairs (f_m^N, ξ_m^N) , $m = 1, 2$, and $(\tilde{f}^N, \tilde{\xi}^N)$, where $\tilde{f}^N = f_2^N + \|f_1^N - f_2^N\|_\infty$ and $\tilde{\xi}^N = \xi_2^N + \|\xi_1^N - \xi_2^N\|_\infty$, satisfy the conditions of Theorem 4.2 for \tilde{C} instead of C , \tilde{K} instead of K and $L = K \vee b$. Therefore, the corresponding BSΔEs have unique solutions, which, since $\tilde{f}^N \geq f_1^N$ and $\tilde{\xi}^N \geq \xi_1^N$, satisfy $\tilde{Y}_t^N \geq Y_{1,t}^N$ for all t . Note that the solution of the deterministic BSΔE

$$\begin{aligned} \hat{Y}_{t_i^N}^N &= \hat{Y}_{t_{i+1}^N}^N + (\|f_1^N - f_2^N\|_\infty + K\hat{Y}_{t_i^N}^N)\Delta\langle W^N \rangle_{t_{i+1}^N}, \\ \hat{Y}_T^N &= \|\xi_1^N - \xi_2^N\|_\infty, \end{aligned} \tag{A.14}$$

is given by

$$\hat{Y}_{t_i^N}^N = \frac{\|\xi_1^N - \xi_2^N\|_\infty}{\prod_{j=i+1}^{i_N} (1 - K\Delta\langle W^N \rangle_{t_j^N})} + \|f_1^N - f_2^N\|_\infty \sum_{j=i+1}^{i_N} \frac{\Delta\langle W^N \rangle_{t_j^N}}{\prod_{l=i+1}^j (1 - K\Delta\langle W^N \rangle_{t_l^N})}.$$

In particular, \hat{Y}_t^N is positive and decreasing in t , and it satisfies

$$\hat{Y}_{t_i^N}^N \leq \frac{\|\xi_1^N - \xi_2^N\|_\infty + \|f_1^N - f_2^N\|_\infty \sum_{j=i+1}^{i_N} \Delta\langle W^N \rangle_{t_j^N}}{\prod_{j=i+1}^{i_N} (1 - K\Delta\langle W^N \rangle_{t_j^N})}.$$

Hence, by the choice of the constant D , one obtains the estimate

$$\sup_t \hat{Y}_t^N = \hat{Y}_0^N \leq D(\|\xi_1^N - \xi_2^N\|_\infty + \|f_1^N - f_2^N\|_\infty). \tag{A.15}$$

In particular, since $\|\xi_1^N - \xi_2^N\|_\infty \leq 2C$ and $\|f_1^N - f_2^N\|_\infty \leq K$, it follows from (A.15) that

$$\sup_t \hat{Y}_t^N \leq (2C + K)(\exp(KT) + 1)(T + 1). \tag{A.16}$$

Next, notice that the process

$$\tilde{Y}_t^N := Y_{2,t}^N + \hat{Y}_t^N$$

satisfies

$$\begin{aligned}\bar{Y}_{t_i^N}^N &= \bar{Y}_{t_{i+1}^N}^N + \{f_2^N(t_{i+1}^N, W^N, Y_{2,t_i^N}^N, Z_{2,t_{i+1}^N}^N) + \|f_1^N - f_2^N\|_\infty + K\hat{Y}_{t_i^N}^N\} \Delta \langle W^N \rangle_{t_{i+1}^N} \\ &\quad - Z_{2,t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^N - \Delta M_{2,t_{i+1}^N}^N, \\ \bar{Y}_T^N &= \xi_2^N + \|\xi_1^N - \xi_2^N\|_\infty,\end{aligned}$$

and since f_2^N is K -Lipschitz in y , one has

$$f_2^N(t_{i+1}^N, W^N, \bar{Y}_{t_i^N}^N, Z_{2,t_{i+1}^N}^N) \leq f_2^N(t_{i+1}^N, W^N, Y_{2,t_i^N}^N, Z_{2,t_{i+1}^N}^N) + K\hat{Y}_{t_i^N}^N.$$

Hence,

$$\alpha_{t_i^N} = f_2^N(t_{i+1}^N, W^N, Y_{2,t_i^N}^N, Z_{2,t_{i+1}^N}^N) - f_2^N(t_{i+1}^N, W^N, \bar{Y}_{t_i^N}^N, Z_{2,t_{i+1}^N}^N) + K\hat{Y}_{t_i^N}^N \geq 0$$

and \bar{Y}^N satisfies the BSΔE

$$\begin{aligned}\bar{Y}_{t_i^N}^N &= \bar{Y}_{t_{i+1}^N}^N + \{f_2^N(t_{i+1}^N, W^N, \bar{Y}_{t_i^N}^N, Z_{2,t_{i+1}^N}^N) + \|f_1^N - f_2^N\|_\infty + \alpha_{t_i^N}\} \Delta \langle W^N \rangle_{t_{i+1}^N} \\ &\quad - Z_{2,t_{i+1}^N}^N \Delta W_{t_{i+1}^N}^N - \Delta M_{2,t_{i+1}^N}^N, \\ \bar{Y}_T^N &= \xi_2^N + \|\xi_1^N - \xi_2^N\|_\infty.\end{aligned}\tag{A.17}$$

Since f_2^N is K -Lipschitz in y , one obtains from the estimate (A.16) that

$$\begin{aligned}\|\alpha_{t_i^N}\|_\infty &\leq \|f_2^N(t_{i+1}^N, W^N, Y_{2,t_i^N}^N, Z_{2,t_{i+1}^N}^N) - f_2^N(t_{i+1}^N, W^N, \bar{Y}_{t_i^N}^N, Z_{2,t_{i+1}^N}^N)\|_\infty + K\|\hat{Y}_{t_i^N}^N\|_\infty \\ &\leq 2K\|\hat{Y}_{t_i^N}^N\|_\infty \leq 2K(2C + K)(\exp(KT) + 1)(T + 1),\end{aligned}$$

which shows that the BSΔE (A.17) satisfies the assumptions of Theorem 4.2 for \tilde{C} , \tilde{K} and $L = K \vee b$. Hence, a comparison of \tilde{Y}^N to \bar{Y}^N yields

$$Y_{1,t}^N \leq \tilde{Y}_t^N \leq \bar{Y}_t^N = Y_{2,t}^N + \hat{Y}_t^N \leq Y_{2,t}^N + D(\|\xi_1^N - \xi_2^N\|_\infty + \|f_1^N - f_2^N\|_\infty)$$

for all t . By symmetry, one also has

$$Y_{2,t}^N \leq Y_{1,t}^N + D(\|f_1^N - f_2^N\|_\infty + \|\xi_1^N - \xi_2^N\|_\infty)$$

for all t , and the proof is complete. \square

Proof of Lemma 5.2. Let $C \in \mathbb{R}_+$ such that φ is bounded by C and $|\varphi(w_1) - \varphi(w_2)| \leq C \sup_{1 \leq i \leq n} |w_1(s_i) - w_2(s_i)|$ for all $w_1, w_2 \in \mathbb{R}^{d \times n}$. Choose $N_0 \in \mathbb{N}$ and $D \in \mathbb{R}_+$ such that

for all $N \geq N_0$, $\sup_i |\Delta W_{t_i^N}^N| \leq 1$ and the statement of Lemma 5.1 holds. From Lemma 3.1, we know that

$$Z_{t_i^N}^{N,k} = \frac{\mathbb{E}[Y_{t_i^N}^N \Delta W_{t_i^N}^{N,k} | \mathcal{F}_{t_{i-1}^N}^N]}{\Delta \langle W^N \rangle_{t_i^N}},$$

and since $Y_{t_i^N}^N$ is $\mathcal{F}_{t_i^N}^N$ -measurable, it can be written as

$$Y_{t_i^N}^N = y_i^N(W_{t_1^N}^N, \dots, W_{t_i^N}^N)$$

for a Borel measurable function $y_i^N : \mathbb{R}^{d \times i} \rightarrow \mathbb{R}$. We want to show that y_i^N can be chosen uniformly Lipschitz-continuous in the last argument. To do that, let us condition on $W_{t_j^N}^N = w(t_j^N)$, $j = 1, \dots, i - 1$ and $W_{t_i^N}^N = x$. Denote $\tilde{W}_t^N = W_t^N - W_{t_i^N}^N$, $t \in [t_i^N, T]$, and define $r = \max\{m : s_m \leq t_i^N\}$. Then for $t_j^N \geq t_i^N$, the conditioned BSΔE with solution $(Y^{N,x}, Z^{N,x}, M^{N,x})$ can be written as

$$\begin{aligned} Y_{t_j^N}^{N,x} &= Y_{t_{j+1}^N}^{N,x} + f^N(t_{j+1}^N, w(t_1^N), \dots, w(t_{i-1}^N), x + \tilde{W}^N, Y_{t_j^N}^{N,x}, Z_{t_{j+1}^N}^{N,x}) \Delta \langle \tilde{W}^N \rangle_{t_{j+1}^N} \\ &\quad - Z_{t_{j+1}^N}^{N,x} \Delta \tilde{W}_{t_{j+1}^N}^N - \Delta M_{t_{j+1}^N}^{N,x}, \end{aligned} \tag{A.18}$$

$$Y_T^{N,x} = \varphi(w(s_1), \dots, w(s_r), x + \tilde{W}_{s_{r+1}}^N, \dots, x + \tilde{W}_{s_n}^N).$$

Thus, for $t \geq t_i^N$ we have $Y_t^{N,x} = \bar{Y}_t^{N,x}$, where $\bar{Y}^{N,x}$ solves the BSΔE driven by the processes W^N with terminal conditions $\xi^{N,x} = \varphi(w(s_1), \dots, w(s_r), x + W_{s_{r+1}}^N - W_{t_i^N}^N, \dots, x + W_{s_n}^N - W_{t_i^N}^N)$ and drivers

$$\begin{aligned} &\bar{f}^{N,x}(t, w(t_1^N), \dots, w(t_{i-1}^N), W^N, y, z) \\ &= \begin{cases} f^N(t, w(t_1^N), \dots, w(t_{i-1}^N), x + W^N - W_{t_i^N}^N, y, z), & \text{for } t > t_i^N, \\ 0, & \text{for } t \leq t_i^N. \end{cases} \end{aligned}$$

Clearly, all $\bar{f}^{N,x}$ are adapted, left-continuous and satisfy (f1)–(f3). By our Lipschitz assumption on φ and f^N , one has,

$$\|\xi^{N,x_1} - \xi^{N,x_2}\|_\infty \leq C|x_1 - x_2|$$

and

$$\|\bar{f}^{N,x_1} - \bar{f}^{N,x_2}\|_\infty \leq K|x_1 - x_2|$$

for all $x_1, x_2 \in \mathbb{R}^d$. In particular,

$$\|\bar{f}^{N,x_1} - \bar{f}^{N,x_2}\|_\infty \leq K$$

if $|x_1 - x_2| \leq 1$. So one obtains from Lemma 5.1 that for all $x_1, x_2 \in \mathbb{R}^d$ satisfying $|x_1 - x_2| \leq 1$,

$$\begin{aligned} |Y_{t_i^N}^{N,x_1} - Y_{t_i^N}^{N,x_2}| &\leq \sup_{0 \leq t \leq T} |\bar{Y}_t^{N,x_1} - \bar{Y}_t^{N,x_2}| \\ &\leq D(\|\xi^{N,x_1} - \xi^{N,x_2}\|_\infty + \|\bar{f}^{N,x_1} - \bar{f}^{N,x_2}\|_\infty) \\ &\leq D(C + K)|x_1 - x_2|. \end{aligned}$$

Note that

$$\mathbb{E}[y_{t_i^N}^{N,N}(W_{t_1^N}^N, \dots, W_{t_{i-1}^N}^N, W_{t_{i-1}^N}^N) \Delta \langle W^N \rangle_{t_i^N} | \mathcal{F}_{t_{i-1}^N}^N] = 0,$$

and therefore,

$$\begin{aligned} |Z_{t_i^N}^{N,k}| &= \Delta \langle W^N \rangle_{t_i^N}^{-1} |\mathbb{E}[Y_{t_i^N}^N \Delta W_{t_i^N}^{N,k} | \mathcal{F}_{t_{i-1}^N}^N]| \\ &= |\mathbb{E}[(y_{t_i^N}^{N,N}(W_{t_1^N}^N, \dots, W_{t_{i-1}^N}^N, W_{t_{i-1}^N}^N + \Delta W_{t_i^N}^N) \\ &\quad - y_{t_i^N}^{N,N}(W_{t_1^N}^N, \dots, W_{t_{i-1}^N}^N, W_{t_{i-1}^N}^N)) \Delta W_{t_i^N}^{N,k} | \mathcal{F}_{t_{i-1}^N}^N]| / (\Delta \langle W^N \rangle_{t_i^N}) \\ &\leq \mathbb{E}[|y_{t_i^N}^{N,N}(W_{t_1^N}^N, \dots, W_{t_{i-1}^N}^N, W_{t_{i-1}^N}^N + \Delta W_{t_i^N}^N) \\ &\quad - y_{t_i^N}^{N,N}(W_{t_1^N}^N, \dots, W_{t_{i-1}^N}^N, W_{t_{i-1}^N}^N)| |\Delta W_{t_i^N}^{N,k}| | \mathcal{F}_{t_{i-1}^N}^N] / (\Delta \langle W^N \rangle_{t_i^N}) \\ &\leq D(C + K) \frac{\mathbb{E}[|\Delta W_{t_i^N}^N| |\Delta W_{t_i^N}^{N,k}| | \mathcal{F}_{t_{i-1}^N}^N]}{\Delta \langle W^N \rangle_{t_i^N}} \\ &\leq D(C + K) \frac{\mathbb{E}[|\Delta W_{t_i^N}^N| |\Delta W_{t_i^N}^{N,k}| | \mathcal{F}_{t_{i-1}^N}^N]}{\Delta \langle W^N \rangle_{t_i^N}} = D(C + K)d. \end{aligned}$$

□

A.4. Remaining proofs of Section 6

Proof of Proposition 6.1. Set $\bar{C} = (C + 1) \exp(KT)$ and denote

$$a = \sup_{N,i} \frac{\|\Delta W_{t_i^N}^N\|_\infty}{\sqrt{\Delta \langle W^N \rangle_{t_i^N}}} < \infty.$$

Choose $N_0 \in \mathbb{N}$ such that for all $N \geq N_0$ the conclusion of Theorem 4.2 holds and

$$\sqrt{d}La(\Delta \langle W^N \rangle_{t_i^N})^{1/2} + d^{(2+q)/4}L\bar{C}^{q/2}a(\Delta \langle W^N \rangle_{t_i^N})^{(2-q)/4} < 1. \quad (\text{A.19})$$

Then it follows from Theorem 4.2 that for fixed $N \geq N_0$, the N th BSΔE has a unique solution (Y^N, Z^N, M^N) with $|Y_t^N| \leq \bar{C}$ for all $t \in [0, T]$. Now choose an \mathbb{R}^d -valued (\mathcal{F}_t^N) -adapted

process μ^N that is constant on the intervals $(t_{i-1}^N, t_i^N]$ and satisfies (6.1). It follows from the definition of g^N that

$$\begin{aligned} Y_{t_i^N}^N &= \xi^N + \sum_{j=i+1}^{i_N} f^N(t_j^N, Y_{t_{j-1}^N}^N, Z_{t_j^N}^N) \Delta \langle W^N \rangle_{t_j^N} - \sum_{j=i+1}^{i_N} Z_{t_j^N}^N \Delta W_{t_j^N}^N - (M_T^N - M_{t_i^N}^N) \\ &\geq \xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \mu_{t_j^N}^N) \Delta \langle W^N \rangle_{t_j^N} - \sum_{j=i+1}^{i_N} Z_{t_j^N}^N \Delta W_{t_j^N}^{N, \mu^N} - (M_T^N - M_{t_i^N}^N). \end{aligned}$$

Since M^N is orthogonal to W^N , its components are still martingales under P^{μ^N} , and one obtains

$$Y_{t_i^N}^N \geq \mathbb{E}^{\mu^N} \left[\xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \mu_{t_j^N}^N) \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right]. \tag{A.20}$$

On the other hand, it can be shown (see, e.g., Cheridito *et al.* [7]) that for each i there exists a $\hat{\mu}_{t_i^N}^N \in L^0(\mathcal{F}_{t_{i-1}^N}^N)^d$ such that

$$f^N(t_i^N, Y_{t_{i-1}^N}^N, Z_{t_i^N}^N + z) - f^N(t_i^N, Y_{t_{i-1}^N}^N, Z_{t_i^N}^N) \geq z \hat{\mu}_{t_i^N}^N \quad \text{for all } z \in \mathbb{R}^d.$$

Set $\hat{\mu}_t^N = \hat{\mu}_{t_i^N}^N$ for $t \in (t_{i-1}^N, t_i^N]$. Then $\hat{\mu}^N$ is a left-continuous \mathbb{R}^d -valued (\mathcal{F}_t^N) -adapted process satisfying

$$f^N(t_i^N, Y_{t_{i-1}^N}^N, Z_{t_i^N}^N) + g^N(t_i^N, Y_{t_{i-1}^N}^N, \hat{\mu}_{t_i^N}^N) = \hat{\mu}_{t_i^N}^N Z_{t_i^N}^N \quad \text{for all } i. \tag{A.21}$$

So if we can show that $\hat{\mu}^N$ satisfies (6.1) and (6.4), the equality in (A.20) becomes an equality and the proposition is proved. To see that $\hat{\mu}^N$ satisfies (6.1), note that it follows from the Cauchy–Schwarz inequality that

$$\begin{aligned} |Z_{t_i^N}^{N,k}| &= |(\Delta \langle W^N \rangle_{t_i^N})^{-1} \mathbb{E}[Y_{t_i^N}^N \Delta W_{t_i^N}^{N,k} | \mathcal{F}_{t_{i-1}^N}^N]| \\ &\leq |(\Delta \langle W^N \rangle_{t_i^N})^{-1} \sqrt{\mathbb{E}[|Y_{t_i^N}^N|^2 | \mathcal{F}_{t_{i-1}^N}^N]} \sqrt{\mathbb{E}[|\Delta W_{t_i^N}^{N,k}|^2 | \mathcal{F}_{t_{i-1}^N}^N]} \\ &\leq \bar{C} (\Delta \langle W^N \rangle_{t_i^N})^{-1/2}, \end{aligned}$$

and therefore,

$$|Z_{t_i^N}^N| \leq \sqrt{d} \bar{C} (\Delta \langle W^N \rangle_{t_i^N})^{-1/2}. \tag{A.22}$$

From condition (v) one obtains

$$|\hat{\mu}_{t_i^N}^{N,k}| \leq L(1 + |Z_{t_i^N}^N|^{q/2}) \quad \text{for all } k.$$

Hence, it follows from estimate (A.22) that

$$|\hat{\mu}_{t_i^N}^N| \leq \sqrt{d}L(1 + |Z_{t_i^N}^N|^{q/2}) \leq \sqrt{d}L + d^{(2+q)/4}L\bar{C}^{q/2}(\Delta\langle W^N \rangle_{t_i^N})^{-q/4}.$$

This gives

$$|\hat{\mu}_{t_i^N}^N \Delta W_{t_i^N}^N| \leq |\hat{\mu}_{t_i^N}^N| |\Delta W_{t_i^N}^N| \leq \sqrt{d}La(\Delta\langle W^N \rangle_{t_i^N})^{1/2} + d^{(2+q)/4}L\bar{C}^{q/2}a(\Delta\langle W^N \rangle_{t_i^N})^{(2-q)/4} < 1$$

and shows that $\hat{\mu}^N$ satisfies condition (6.1).

To show (6.4), we first assume $q = 1$. Then one has

$$\begin{aligned} g^N(t_j^N, Y_{t_{j-1}^N}^N, \hat{\mu}_{t_j^N}^N) &= \operatorname{ess\,sup}_z \{ \hat{\mu}_{t_j^N}^N z - f^N(t_j^N, Y_{t_{j-1}^N}^N, z) \} \\ &\geq \operatorname{ess\,sup}_z \{ \hat{\mu}_{t_j^N}^N z - K(1 + |Y_{t_{j-1}^N}^N| + |z|) \}. \end{aligned}$$

It follows that

$$|\hat{\mu}_{t_j^N}^{N,k}| \leq K \quad \text{for all } k = 1, \dots, d,$$

and it is clear that $\hat{\mu}^N$ satisfies condition (6.4). If $q \in (1, 2)$, denote $|x|_q = (\sum_{i=1}^d |x_i|^q)^{1/q}$, and observe that there exist constants $C_1, C_2, C_3 > 0$ such that

$$\begin{aligned} g^N(t_j^N, Y_{t_{j-1}^N}^N, \hat{\mu}_{t_j^N}^N) &= \operatorname{ess\,sup}_z \{ \hat{\mu}_{t_j^N}^N z - f^N(t_{j+1}^N, Y_{t_j^N}^N, z) \} \\ &\geq \operatorname{ess\,sup}_z \{ \hat{\mu}_{t_j^N}^N z - K(1 + |Y_{t_j^N}^N| + |z|^q) \} \\ &\geq -K(1 + |Y_{t_j^N}^N|) + \operatorname{ess\,sup}_z \{ \hat{\mu}_{t_j^N}^N z - C_1|z|_q^q \} \\ &= -K(1 + |Y_{t_j^N}^N|) + C_2|\hat{\mu}_{t_j^N}^N|_{q/(q-1)}^{q/(q-1)} \geq -K(1 + |Y_{t_j^N}^N|) + C_3(|\hat{\mu}_{t_j^N}^N|^2 + 1). \end{aligned} \tag{A.23}$$

Since

$$Y_{t_i^N}^N = \mathbb{E}^{\hat{\mu}^N} \left[\xi^N - \sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \hat{\mu}_{t_j^N}^N) \Delta\langle W^N \rangle_{t_j^N} | \mathcal{F}_{t_i^N}^N \right]$$

and ξ^N and $Y_{t_i^N}^N$ are bounded by C and \bar{C} , respectively, one obtains

$$\mathbb{E}^{\hat{\mu}^N} \left[\sum_{j=i+1}^{i_N} g^N(t_j^N, Y_{t_{j-1}^N}^N, \hat{\mu}_{t_j^N}^N) \Delta\langle W^N \rangle_{t_j^N} | \mathcal{F}_{t_i^N}^N \right] \leq C + \bar{C}.$$

This together with (A.23) and the uniform boundedness of Y^N shows that $\hat{\mu}^N$ fulfills (6.4). \square

Lemma A.3. *Let μ be an (\mathcal{F}_t^N) -adapted process that is constant on the intervals $(t_{i-1}^N, t_i^N]$ and satisfies (6.1). Then one has*

$$\begin{aligned} & \mathbb{E} \left[\prod_{j=i+1}^{i_N} (1 + \mu_{t_j^N} \Delta W_{t_j^N}^N) \log \left(\prod_{j=i+1}^{i_N} (1 + \mu_{t_j^N} \Delta W_{t_j^N}^N) \right) \middle| \mathcal{F}_{t_i^N}^N \right] \\ & \leq \mathbb{E}^\mu \left[\sum_{j=i+1}^{i_N} |\mu_{t_j^N}|^2 \Delta \langle W^N \rangle_{t_j^N} \middle| \mathcal{F}_{t_i^N}^N \right]. \end{aligned}$$

Proof. One can write

$$\begin{aligned} & \mathbb{E} \left[\prod_{j=i+1}^{i_N} (1 + \mu_{t_j^N} \Delta W_{t_j^N}^N) \log \left(\prod_{j=i+1}^{i_N} (1 + \mu_{t_j^N} \Delta W_{t_j^N}^N) \right) \middle| \mathcal{F}_{t_i^N}^N \right] \\ & = \sum_{j=i+1}^{i_N} \mathbb{E}^\mu [\log(1 + \mu_{t_j^N} \Delta W_{t_j^N}^N) | \mathcal{F}_{t_i^N}^N] \leq \sum_{j=i+1}^{i_N} \log(\mathbb{E}^\mu [(1 + \mu_{t_j^N} \Delta W_{t_j^N}^N) | \mathcal{F}_{t_i^N}^N]), \end{aligned}$$

where the inequality follows from Jensen’s inequality. The right-hand side can be estimated as follows:

$$\begin{aligned} & \sum_{j=i+1}^{i_N} \log \left\{ 1 + \sum_{k=1}^d \mathbb{E}^\mu [\mu_{t_j^N}^k \mathbb{E}^\mu [\Delta W_{t_j^N}^{N,k} | \mathcal{F}_{t_{j-1}^N}^N] | \mathcal{F}_{t_i^N}^N] \right\} \\ & = \sum_{j=i+1}^{i_N} \log \left\{ 1 + \sum_{k=1}^d \mathbb{E}^\mu [(\mu_{t_j^N}^k)^2 \Delta \langle W^N \rangle_{t_j^N} | \mathcal{F}_{t_i^N}^N] \right\} \leq \sum_{j=i+1}^{i_N} \mathbb{E}^\mu [|\mu_{t_j^N}|^2 \Delta \langle W^N \rangle_{t_j^N} | \mathcal{F}_{t_i^N}^N]. \end{aligned}$$

The equality holds because

$$\mathbb{E}^\mu [\Delta W_{t_j^N}^{N,k} | \mathcal{F}_{t_{j-1}^N}^N] = \mu_{t_j^N}^k \Delta \langle W_{t_j^N}^N \rangle.$$

For the inequality we used $\log(1 + x) \leq x$. □

Lemma A.4. *For all $N \in \mathbb{N}$, let $h^N : [0, T] \rightarrow \mathbb{R}$ be a function that is constant on the intervals $[t_i^N, t_{i+1}^N)$. If there exist constants $a, b \in \mathbb{R}_+$ such that*

$$|h^N(T)| \leq a \quad \text{and} \quad |h^N(t_i^N)| \leq a + b \sum_{j=i+1}^{i_N} |h^N(t_{j-1}^N)| \Delta \langle W^N \rangle_{t_j^N} \quad \text{for all } N \text{ and } i \leq i_N - 1,$$

there exists an $N_0 \in \mathbb{N}$ such that

$$|h^N(t_i^N)| \leq 2a \exp(b(T - t_i^N)) \quad \text{for all } N \geq N_0 \text{ and } i = 0, \dots, i_N.$$

Proof. For N so large that $\sup_i \Delta \langle W^N \rangle_{t_i^N} < 1/b$, the function given by

$$H^N(T) := a \quad \text{and} \quad H^N(t) := a \prod_{j:t_j^N > t} (1 - b \Delta \langle W^N \rangle_{t_j^N})^{-1}, \quad t < T$$

solves

$$H^N(t_i^N) = a + b \sum_{j=i+1}^{i_N} H^N(t_{j-1}^N) \Delta \langle W^N \rangle_{t_j^N} \quad \text{for all } i \leq i_N - 1,$$

and converges uniformly to $a \exp(b(T - t))$. In particular, there exists an $N_0 \in \mathbb{N}$ such that

$$H^N(t) \leq 2a \exp(b(T - t)) \quad \text{for all } t \text{ and } N \geq N_0.$$

So the lemma follows if we can show that $|h^N(t_i^N)| \leq H^N(t_i^N)$ for all $N \geq N_0$ and $i = 0, \dots, i_N$. For $i = i_N$ this is obvious, and if it holds for $j \geq i + 1$, then

$$\begin{aligned} |h^N(t_i^N)| &\leq \frac{a + b \sum_{j=i+2}^{i_N} |h^N(t_{j-1}^N)| \Delta \langle W^N \rangle_{t_j^N}}{1 - b \Delta \langle W^N \rangle_{t_{i+1}^N}} \\ &\leq \frac{a + b \sum_{j=i+2}^{i_N} |h^N(t_{j-1}^N)| \Delta \langle W^N \rangle_{t_j^N}}{1 - b \Delta \langle W^N \rangle_{t_{i+1}^N}} = H^N(t_i^N). \end{aligned} \quad \square$$

Acknowledgments

Financial support from NSF Grant DMS-06-42361 is gratefully acknowledged.

References

- [1] Bally, V. (1997). Approximation scheme for solutions of BSDE. In *Backward Stochastic Differential Equations (Paris, 1995–1996)*. Pitman Res. Notes Math. Ser. **364** 177–191. Harlow: Longman. [MR1752682](#)
- [2] Barrieu, P. and El Karoui, N. (2009). Pricing, hedging and optimally designing derivatives via minimization of risk measures. In *Indifference Pricing* (R. Carmona, ed.). Princeton: Princeton Univ. Press. [MR2547456](#)
- [3] Bouchard, B. and Touzi, N. (2004). Discrete-time approximation and Monte-Carlo simulation of backward stochastic differential equations. *Stochastic Process. Appl.* **111** 175–206. [MR2056536](#)
- [4] Briand, P., Delyon, B. and Mémin, J. (2001). Donsker-type theorem for BSDEs. *Electron. Commun. Probab.* **6** 1–14 (electronic). [MR1817885](#)
- [5] Briand, P., Delyon, B. and Mémin, J. (2002). On the robustness of backward stochastic differential equations. *Stochastic Process. Appl.* **97** 229–253. [MR1875334](#)
- [6] Briand, P. and Hu, Y. (2006). BSDE with quadratic growth and unbounded terminal value. *Probab. Theory Related Fields* **136** 604–618. [MR2257138](#)

- [7] Cheridito, P., Kupper, M. and Vogelpoth, N. (2011). Conditional analysis on \mathbb{R}^d . Preprint.
- [8] Cheridito, P. and Stadje, M. (2012). Existence, minimality and approximation of solutions to BSDEs with convex drivers. *Stochastic Process. Appl.* **122** 1540–1565. [MR2914762](#)
- [9] Chevance, D. (1997). Numerical methods for backward stochastic differential equations. In *Numerical Methods in Finance. Publ. Newton Inst.* 232–244. Cambridge: Cambridge Univ. Press. [MR1470517](#)
- [10] Cohen, S.N. and Elliott, R.J. (2010). A general theory of finite state backward stochastic difference equations. *Stochastic Process. Appl.* **120** 442–466. [MR2594366](#)
- [11] Coquet, F., Mackevičius, V. and Mémin, J. (1999). Corrigendum to: “Stability in D of martingales and backward equations under discretization of filtration” [*Stochastic Processes Appl.* **75** (1998) 235–248; [MR1632205](#) (99f:60109)]. *Stochastic Process. Appl.* **82** 335–338. [MR1700013](#)
- [12] Coquet, F., Mémin, J. and Słominski, L. (2001). On weak convergence of filtrations. In *Séminaire de Probabilités, XXXV. Lecture Notes in Math.* **1755** 306–328. Berlin: Springer. [MR1837294](#)
- [13] Delbaen, F., Hu, Y. and Bao, X. (2011). Backward SDEs with superquadratic growth. *Probab. Theory Related Fields* **150** 145–192. [MR2800907](#)
- [14] Delbaen, F., Peng, S. and Rosazza Gianin, E. (2010). Representation of the penalty term of dynamic concave utilities. *Finance Stoch.* **14** 449–472. [MR2670421](#)
- [15] Douglas, J. Jr., Ma, J. and Protter, P. (1996). Numerical methods for forward–backward stochastic differential equations. *Ann. Appl. Probab.* **6** 940–968. [MR1410123](#)
- [16] El Karoui, N., Peng, S. and Quenez, M.C. (1997). Backward stochastic differential equations in finance. *Math. Finance* **7** 1–71. [MR1434407](#)
- [17] El Otmani, M. (2006). Approximation scheme for solutions of backward stochastic differential equations via the representation theorem. *Ann. Math. Blaise Pascal* **13** 17–29. [MR2233010](#)
- [18] Gobet, E., Lemor, J.P. and Warin, X. (2005). A regression-based Monte Carlo method to solve backward stochastic differential equations. *Ann. Appl. Probab.* **15** 2172–2202. [MR2152657](#)
- [19] Ikeda, N. and Watanabe, S. (1989). *Stochastic Differential Equations and Diffusion Processes*, 2nd ed. *North-Holland Mathematical Library* **24**. Amsterdam: North-Holland. [MR1011252](#)
- [20] Imkeller, P. and Reis, G. (2009). Path regularity and explicit truncation order for BSDE with drivers of quadratic growth. *Stochastic Process. Appl.* **120** 348–379.
- [21] Kobylanski, M. (2000). Backward stochastic differential equations and partial differential equations with quadratic growth. *Ann. Probab.* **28** 558–602. [MR1782267](#)
- [22] Ma, J., Protter, P., San Martín, J. and Torres, S. (2002). Numerical method for backward stochastic differential equations. *Ann. Appl. Probab.* **12** 302–316. [MR1890066](#)
- [23] Ma, J., Protter, P. and Yong, J.M. (1994). Solving forward–backward stochastic differential equations explicitly—a four step scheme. *Probab. Theory Related Fields* **98** 339–359. [MR1262970](#)
- [24] Morlais, M.A. (2009). Quadratic BSDEs driven by a continuous martingale and applications to the utility maximization problem. *Finance Stoch.* **13** 121–150. [MR2465489](#)
- [25] Pardoux, É. and Peng, S.G. (1990). Adapted solution of a backward stochastic differential equation. *Systems Control Lett.* **14** 55–61. [MR1037747](#)
- [26] Peng, S. (2004). Nonlinear expectations, nonlinear evaluations and risk measures. In *Stochastic Methods in Finance. Lecture Notes in Math.* **1856** 165–253. Berlin: Springer. [MR2113723](#)
- [27] Richou, A. (2011). Numerical simulation of BSDEs with drivers of quadratic growth. *Ann. Appl. Probab.* **21** 1933–1964. [MR2884055](#)
- [28] Toldo, S. (2006). Stability of solutions of BSDEs with random terminal time. *ESAIM Probab. Stat.* **10** 141–163 (electronic). [MR2218406](#)
- [29] Toldo, S. (2007). Corrigendum to: “Stability of solutions of BSDEs with random terminal time”. *ESAIM Probab. Stat.* **11** 381–384 (electronic). [MR2339299](#)
- [30] Zhang, J. (2004). A numerical scheme for BSDEs. *Ann. Appl. Probab.* **14** 459–488. [MR2023027](#)

- [31] Zhang, Y. and Zheng, W. (2002). Discretizing a backward stochastic differential equation. *Int. J. Math. Math. Sci.* **32** 103–116. [MR1937828](#)

Received November 2010 and revised September 2011